1 Chronic intermittent alcohol yields sex-specific disruptions in cortical-striatal-limbic oscillations

3 Kelly A. Hewitt[1], Skylar E. Nicholson[1], Madilyn J. Peterson[1], Lucas L. Dwiel[2], Angela M.

4 Henricks[1]

6 [1]Department of Psychology, Washington State University, Pullman, WA, USA

7 [2]Department of Psychiatry, Geisel School of Medicine at Dartmouth, Hanover, NH, USA

9 Corresponding author: Angela M. Henricks

10 Assistant Professor

11 Department of Psychology

12 Washington State University

13 Pullman, WA 99163

14 Email: angela.henricks@wsu.edu

ABSTRACT

Background: While the neurobiology of alcohol use disorder (AUD) has been extensively

researched, the vast majority of these studies included only male organisms. However, there

are significant sex differences in both the causes and consequences of alcohol misuse and

dependence, suggesting sex-specific neurobiological mechanisms. The current study used a

rodent model to determine whether chronic alcohol exposure impacts sex-specific neural

circuits, and whether these changes contribute to the development of alcohol misuse.

Methods: Male and female Sprague-Dawley rats were trained to self-administer 10% alcohol

before implanting bilateral electrodes into the infralimbic medial prefrontal cortex (IL), nucleus

accumbens shell (NAcSh), and central nucleus of the amygdala (CeA). Half of the rats were

then exposed to four weeks of chronic intermittent alcohol (CIA) vapor (14 hours on/10 hours

off). During acute withdrawal (6-8 after the vapor turns off), local field potentials (LFPs) were

recorded from the IL, NAcSh, and CeA during 30-minute self-administration sessions. Using an

unbiased machine learning approach, we built predictive models to determine whether/which

LFP features could distinguish CIA-exposed from control rats in each sex, as well as if any of

these LFP features correlated with rates of alcohol self-administration.

Results: Female rats self-administered more alcohol in general compared to males, but only

males exposed to CIA showed increased alcohol intake during acute withdrawal. LFPs

predicted CIA exposure in both sexes better than chance estimates, but models built on IL and

NAcSh oscillations performed the best in males, while models built on IL and CeA LFPs

performed best in females. High γ LFPs recorded in the NAcSh correlated with rates of alcohol

self-administration in males exposed to CIA, while only left-right NAcSh β coherence correlated

with drinking in control females.

Conclusions: These data provide support for the hypothesis that the neural circuits driving

alcohol dependence development are sex-specific, and that high frequency oscillations in the

NAcSh may be related to the increased drinking observed in males exposed to CIA. Overall,

these data add to our understanding of the neurobiological underpinnings behind the sex

differences observed in AUD and offer promising biomarkers for future therapeutic research.

Keywords: Alcohol, local field potentials, self-administration, neural circuits, sex differences

BACKGROUND

Alcohol use disorder (AUD) is the most prevalent substance use disorder in the United

States, causing serious social, behavioral, and health-related problems (1,2). Despite several

FDA-approved treatments for AUD (3), relapse is still extremely common with estimated long-

term relapse rates as high as 85% (4).

There are significant sex differences in the reasons for drinking and the physical,

behavioral, and neurobiological consequences of chronic alcohol misuse. While men consume

more alcohol and are more likely to be diagnosed with AUD compared to women, this gender

gap has begun to shrink over the last decade, with more women engaging in risky alcohol

drinking (5–7). Commonly used medications for AUD, which were developed and tested

primarily in male animals, are less effective in women compared to men (8,9). In fact, the vast

majority of preclinical neuroscience research aimed at understanding AUD has not included

female animals, which may contribute to poor translatability from preclinical animal research to

clinical treatment development (10). Therefore, there is a vital need for a more inclusive and

thorough understanding of the neurobiological mechanisms driving alcohol misuse in both

sexes.

We have previously demonstrated that local field potentials (LFPs) collected in the

nucleus accumbens shell (NAcSh) and medial prefrontal cortex (mPFC) predict the volume of

alcohol consumed in non-dependent male, but not female, rats (11). LFPs reflect the dynamic

flow of information across neural circuits, capturing electrical information within (power) and

between (coherence) brain regions that provides insight into how synchronized/asynchronized

activity is across areas of interest (12,13). LFPs can be collected in awake, freely behaving

animals over long periods of time, and therefore have the potential to serve as biomarkers for

different behavioral states (14). Our previous results suggest that cortical-striatal oscillations

may be particularly relevant to male, but not female, drinking behavior. While the mPFC and

NAcSh play vital roles in voluntary alcohol consumption, contributing to the rewarding,

motivational, and cognitive components of AUD (15), other circuits play important roles as well

and may be more relevant to females.

Women are more likely to use alcohol to alleviate stress (negative reinforcement), while

men are more likely to use alcohol as a positive reinforcer (16). Further, in both humans and

rodents, females are more susceptible to stress-induced relapse (17). We therefore hypothesize

that oscillations recorded in stress-related brain regions, like the amygdala, may contain more

information regarding drinking behavior in females.

In order to collect data relevant to AUD, the current study employed a commonly used

model of alcohol dependence in rodents: chronic intermittent alcohol vapor exposure (CIA). The

main objective was to determine whether CIA alters LFPs recorded in the infralimbic (IL) mPFC,

NAcSh, and central nucleus of the amygdala (CeA) in a sex-specific manner. Using an unbiased

machine-learning approach, we hypothesized that IL, NAcSh, and CeA LFPs would predict

whether animals were exposed to CIA and correlate with rates of alcohol self-administration,

with cortical-striatal and cortical-limbic LFPs containing the most information for males and

females, respectively.

METHODS

Animals

Adult (approximately 70 days old) male and female Sprague-Dawley rats were used for

these experiments (n = 6-7/group/sex; 27 total). Animals were obtained from Inotiv (West

Lafayette, IN) and were housed in standard polycarbonate cages with *ad libitum* access to food

and water. Rats were kept under a 12-hour reverse light-dark cycle (lights off at 08:00), allowing

for all experimental procedures to be conducted during the rat's active phase and to avoid

potential circadian rhythm disruptions. Animal facilities were accredited by the American

Association of Laboratory Animal Care (AALAC), and all procedures were approved by the

Institutional Animal Care and Use Committee (IACUC) at Washington State University. Animals

were kept in same-sex pairs prior to electrode implantation and single housed after surgery to

prevent damage to the electrodes. All animals were handled daily for a minimum of 5 days the

week prior to the start of testing.

Alcohol self-administration training

Self-administration took place in standard, sound attenuating operant boxes (10" L x

12.5" W x 8.5" H) programmed using MED-Associates IV software. Boxes are equipped with two

retractable levers on the same wall of the chamber with a center receptacle for liquid

distribution. Presses to the active lever resulted in the delivery of 0.1mL of liquid. Responses on

the inactive lever resulted in no reinforcer. A house light was illuminated for the duration that

levers were available for each session.

Rats were trained to self-administer 10% ethanol in water under a fixed ratio-1 (FR-1)

reinforcement schedule using a sucrose fade technique, as described in our previous work

(18,19). Rats were trained with the following sequence of solutions: 10% sucrose, 10% sucrose

+ 10% ethanol, 5% sucrose + 10% ethanol, and finally 10% ethanol. Each session in the

operant chambers lasted 30 minutes, and sessions were conducted 5 days per week, Monday-

Friday. Training was considered complete after 15 sessions of 10% ethanol self-administration.

The average g/kg of alcohol consumed over the last four sessions was calculated for each

animal as the baseline drinking level.

Electrode implants

Following self-administration training, rats were anesthetized with isoflurane gas before

electrodes were implanted via stereotaxic surgery. Electrodes were constructed in-house, and

designed to collect LFPs from the NAcSh (from bregma: DV -8.0mm; AP +1.2 mm; ML +/-

1.0mm) and IL mPFC (from bregma: DV –4.5mm, AP +3.4mm, ML +/-0.75mm), as outlined

previously (11,20), as well as the CeA (from bregma: DV –8.5mm , AP –2.0mm, ML +/-3.5mm).

Four stainless steel screws were attached to the skull around the electrode site and dental

acrylic (Lang Dental, Wheeling, IL, USA) was used to secure the electrode in place. No rats had

to be excluded based on electrode placement (see *Histology*). Animals were allowed to recover

for at least seven days before the next phase of the experiment.

Chronic intermittent alcohol exposure

After recovery, animals were either exposed to CIA vapor (n = 6-7/sex) or normal room

air (control; n = 7/sex) for four weeks, as described previously (18,19). Our custom-built, 4-

chamber apparatus (La Jolla Alcohol Research Inc., La Jolla, CA, USA) allows blood alcohol

levels (BALs) to be titrated by the experimenter by adjusting the rate of 95% ethanol that is

vaporized and integrated into the air flow supplying the sealed Plexiglas chambers in which the

rats were housed (25.5" L x 21" W x 13" H). Food and water were available *ad libitum*. CIA

animals were exposed to vaporized ethanol for 14 hours per day from approximately 18:00-

08:00; for 10 hours a day they were just exposed to room air that was filtered into the chamber.

Animals in the CIA condition remained in the vapor chambers for the entirety of the experiment.

Blood was collected at least two times per week to ensure BALs remained at an appropriate

level (between 150-250mg%), as described previously (18,19). BALs were determined by

centrifuging approximately 0.1mL of blood collected from the tail vein. Plasma samples were

then analyzed for ethanol content using the Analox AM1 (Analox Instruments Ltd., Lunenburg,

MA, USA).

Alcohol self-administration after CIA

Following the four-week period of initial CIA exposure, all rats completed four baseline

alcohol self-administration sessions (with a minimum of 1 day between sessions) during acute

alcohol withdrawal (~7 hours after the vapor turned off).

Collecting local field potentials

Next, three separate alcohol self-administration sessions were completed while

collecting LFP data with an OmniPlex Neural Recording Data Acquisition System (Plexon,

Dallas, TX, USA). Time-locked behavior was also recorded via a CinePlex camera system

(Plexon, Dallas, TX, USA). Twelve rats completed an additional session due to technical issues

during one of the original recording sessions. Signals passed through a digital head stage and

custom-made cable connected to a commutator, which allowed free movement and behavior

while tethered to the data acquisition system in the operant chamber.

Histology

At the end of the experiment, rats were euthanized using CO_2 gas and brains were snap

frozen in 2-methylbutane on dry ice. Tissue was stored at -20℃ prior to sectioning. Tissue

sections were then stained with thionine and electrode placement was verified using a Leica

CTR 6500 microscope (Wetzler, Germany) (Figure 1).

Statistical Analyses

Alcohol self-administration

To determine how alcohol self-administration was impacted by CIA, a 2 x 5 mixed

factorial ANOVA was conducted with dependence state (controls, CIA) as the between-subjects

factor, and session as the within-subjects factor (pre-CIA baseline, plus 4 post-CIA sessions),

and g/kg of alcohol consumed as the dependent variable. To see how animals behaved during

LFP collection, a 2 x 3 mixed factorial ANOVA was conducted with dependence state (controls,

CIA) as the between-subjects factor, three LFP self-administration sessions as the within-

subject factors, and g/kg of alcohol consumed as the dependent variable. Because we had an *a*

priori hypothesis that there would be sex differences in behavior, based on our previous data

(18), we performed analyses for each sex separately.

Unbiased prediction modeling

Custom code written in MATLAB R2023a allowed for the characterization of the LFP

power spectral densities (PSDs) within and coherence between brain regions for each rat and

each session, as previously reported (11,20,21). Data from each recording was analyzed using

frequency ranges from the rodent literature (δ =1-4 Hz, θ = 5-10 Hz, α = 11-14 Hz, β = 15-30

Hz, low γ = 45-65 Hz, and high γ = 70-90 Hz) (22,23). After applying a 60Hz noise-filter, PSDs

and coherence for each 5-second segment were averaged together across each 30-minute

recording session. We used a penalized regression model (lasso) to predict binary (CIA vs.

controls) and continuous (g/kg of alcohol consumed) outcomes using IL, NAcSh, and CeA

LFPs, as outlined previously (20).

Each recording session produced 126 LFP features: 36 measures of power (6 frequency

bands x 6 channels) and 90 measures of coherence (6 frequency bands x 15 channel

combinations). The MATLAB package Glmnet was used to implement the lasso using a 5-fold

cross-validation. Models were trained on 80% of the data and tested on a left-out 20%, which

was then repeated 100 times with different, randomly selected sets of training and testing data.

The real models were then compared to 100 random permutations of the data (permuted

models), as described previously (21). Real and permuted models were generated for each sex

separately. Additionally, we performed logistic regressions on each LFP feature individually to

determine the relative predictive accuracy of said feature, as we have previously described in

detail (21). We further used simple linear regressions to determine whether each of the top five

LFP features with the highest prediction accuracy correlated with g/kg of alcohol consumed in

self-administration sessions for each group (male control, male CIA, female control, female

CIA). Lastly, we used moderation analyses to determine if CIA exposure influenced the strength

and directions of any correlations.

RESULTS

Alcohol self-administration

For males, a mixed factorial ANOVA revealed a significant effect of session [$F(4,44) =$

4.01, $p<0.01$, $\eta_p^2 = 0.27$], a significant effect of group [$F(1,11) = 7.68$, $p<0.05$, $\eta_p^2 = 0.41$], and a

significant session*group interaction [$F(4,44) = 3.27$, $p<0.05$, $\eta_p^2 = 0.23$] for g/kg of alcohol

consumed. Post-hoc tests showed that CIA-exposed males drank more alcohol in sessions 3

and 4 ($p<0.05$; Figure 3A) compared to control males.

For females, a mixed factorial ANOVA revealed a significant effect of session [$F(4,48) =$

3.75, $p<0.05$, $\eta_p^2 = 0.24$], but no effect of group [$F(1,12) = 0.65$, $p=0.44$, $\eta_p^2 = 0.05$] or a

group*session interaction [$F(4,48) = 0.30$, $p=0.88$, $\eta_p^2 = 0.02$; Figure 3B] for g/kg of alcohol

consumed. These data are consistent with our and others' previous findings (18,26). It is also

important to note that, in both sexes, inactive lever presses did not differ by group

(Supplemental Figure 1).

During LFP collection, tethering the animals to the commutator eliminated any session or

group effects on g/kg of alcohol consumed, and drinking was suppressed below baseline levels

(Supplementary Figure 2).

Unbiased prediction models

Using LFPs from all regions (IL, NAcSh, and CeA), Figure 4 shows the average area

under the receiver operating characteristic curve (AUC) for the models predicting CIA vs. control

exposure in males (real AUC = 0.63 ± 0.04, permuted AUC = 0.52 ± 0.04; 4A) and females (real

AUC = 0.81 ± 0.03, permuted AUC = 0.50 ± 0.03; 4B). We also determined the prediction

accuracy of models built on just cortical-striatal vs. just cortical-limbic LFPs. In males, models

built on IL and NAcSh LFPs (real AUC = 0.78 ± 0.04, permuted AUC = 0.49 ± 0.04) performed

better than models built on IL and CeA LFPs (real AUC = 0.65 ± 0.04, permuted AUC = 0.48 ±

0.03; Figure 4A). The opposite was true in females; models built on IL and CeA LFPs (real AUC

= 0.83 ± 0.03, permuted AUC = 0.49 ± 0.03) performed better than models built on IL and

NAcSh LFPs (real AUC = 0.57 ± 0.04, permuted AUC = 0.49 ± 0.03; Figure 4B).

Table 1 indicates the top five LFP features with the highest prediction accuracies

distinguishing CIA vs. control exposure for each sex, including the AUCs for each feature. IL

and NAcSh LFPs best predicted CIA exposure in males, while primarily CeA LFPs best

predicted CIA exposure in females.

We also attempted to predict g/kg of alcohol consumed within each tethered session

from the corresponding LFPs. These prediction models did not outperform chance estimates,

likely due to the very low drinking levels exhibited by both sexes when tethered (Supplemental

Figure 3).

As an alternative, we measured whether the average value of each of the top neural

features correlated with the average g/kg of alcohol consumed during acute withdrawal but prior

to tethering (Figure 5). Table 2 shows the correlation coefficients for each of the top neural

features in males and females. In males, moderation analyses revealed a significant interaction

between right NAcSh high γ power and amount of alcohol consumed ($p<0.05$). Further probing

of the interaction revealed that right NAcSh high γ power significantly correlated with g/kg of

alcohol consumed in males exposed to CIA [t(9) = -2.89, $p<0.05$], but not in control males [t(9)

= 1.05, $p=0.32$] (Figure 5A).

In females, moderation analyses revealed a significant interaction between left-right

NAcSh β coherence and amount of alcohol consumed in females ($p<0.01$). Further probing of

the interaction revealed that left-right NAcSh β coherence significantly correlated with g/kg of

alcohol consumed in room air control females [$t(9)$ = -3.50, $p<0.01$], but not in females exposed

to CIA [$t(9)$ = 1.29, $p=0.23$] (Figure 5B). There was also a trend for right CeA δ power to

correlate with g/kg consumed in CIA females ($p=0.08$).

DISCUSSION

Using an unbiased approach, the current study demonstrates that the neural circuits

most impacted by CIA are sex-specific. CIA caused disruptions primarily in cortical-striatal

regions (IL mPFC and NAcSh) in males and in cortical-limbic regions (IL mPFC and CeA) in

females. Further, high γ oscillations recorded in the NAcSh significantly correlated with drinking

during acute withdrawal in males exposed to CIA, but not in control males. In females, we

identified that β coherence between the right and left NAcSh correlated with drinking in control

females, but not in females exposed to CIA. However, there was a trend for a correlation

between low frequency (δ) oscillations recorded in the CeA and drinking in females exposed to

CIA. It is important to note that though many of our single feature correlations were not

significant, our prediction models performed much better than chance estimates, highlighting the

power of combining multiple small effects that would usually be undetectable without much

larger sample sizes. Overall, these data support the notion that chronic alcohol exposure

significantly impacts different neural circuits in males and females, and that high-frequency

striatal oscillations may be related to increased alcohol drinking seen in males, which could

greatly inform our understanding of the sex differences observed in AUD.

284   Consistent with our hypothesis, the neural features that contained the most information

regarding CIA exposure in males primarily came from LFPs recorded in cortical-striatal regions.

This is particularly interesting given the fact that deep brain stimulation (DBS) of the accumbens

decreases alcohol consumption in both male rats and men with AUD (20,32,33). Chronic

alcohol exposure also leads to dysregulated connectivity between the PFC and accumbens,

which may further drive alcohol misuse (34). In line with these findings, CIA exposed males in

this study showed lower IL – NAcSh high γ coherence compared to same-sex controls (Figure

5A), suggesting reduced connectivity between those regions. We also showed that high γ power

in the right NAcSh significantly correlated with g/kg of alcohol consumed in males exposed to

CIA, which is interesting considering that alcohol craving is associated with increased BOLD

activity in the accumbens in men, but not in women (35). Further, high frequency oscillations

have specifically been tied to substance use; cortical, striatal, and limbic γ power changes

correlate with drug delivery or administration of dopamine agonists, suggesting that these

oscillations may play a role in motivated behavior and processing reward (22,36–38). Thus,

CIA-induced changes to high-frequency cortical-striatal oscillations may be especially relevant

to the development of alcohol misuse in males.

300   On the other hand, the current data suggest that oscillations recorded in the CeA, a

region vital to moderating responses to stress and negative affect (39), are significantly

disrupted by CIA in females. Increased negative affect plays a critical role in the initiation and

maintenance of alcohol misuse (39,40,41). Women also more often report using substances to

relieve negative affect compared to men, as well as experience higher levels of subjective

stress in general (16,17). In light of the current data, combined with the fact that others have

shown that repeated cycles of alcohol withdrawal leads to significant neuroadaptations in the

CeA (39,42), we propose that this region is a promising research target for potentially unlocking

more information regarding alcohol dependence development in females.

However, although there was a trend for right CeA δ power to correlate with alcohol

consumption in CIA-exposed females, none of the CeA-based neural features significantly

correlated with drinking. We hypothesize that these findings may be related to inadequate

power due to a small sample size in the present experiment, but this requires further

investigation.

A novel finding of the current study is that left-right NAcSh β coherence correlated with

alcohol consumption in control females. In humans, frontal cortex β oscillations have been

associated with genetic variants that predispose an individual to AUD (43). In the NAc, β

oscillations are associated with action preparation (44), are inversely related to dopamine

availability, and predict relapse to alcohol drinking in rodents (45). While the functional meaning

behind NAcSh β coherence is currently unknown, coherence measures are thought to reflect

communication between brain regions. We therefore hypothesize that disruptions in inter-

accumbens communication may be related to motivation to drink in non-dependent females, but

this theory requires significant further investigation.

We also observed significant sex differences in self-administration behavior, which

aligns with previous work (18,24–27). In a non-dependent state (pre-CIA), female rats drank

more alcohol than their male counterparts. However, CIA males drank more than same-sex

controls during acute withdrawal, which is a hallmark of dependence-like behavior (29,30).

Although dependence-induced escalation was not observed in females, which is consistent with

other studies (18,26), it is important to note that CIA does elicit other signs of dependence in

female rats, including increased signs of physical withdrawal and negative affective behavior

(18,26,31). We therefore reason that the current data is still relevant to alcohol dependence in

females.

There are a few limitations of the current work that require consideration. First, as

mentioned above, females exposed to CIA did not show escalated alcohol consumption during

acute withdrawal, which is a key component of alcohol dependence (29,46). There are several

potential explanations, including that females appear less sensitive than males to the aversive

effects of drug withdrawal (47), or that the exposure time of CIA may not have been adequate to

induce escalated alcohol self-administration for females. Although as little as ten days of CIA

exposure can lead to escalated alcohol intake in male rats (26), other studies have shown

escalations in female self-administration only after ten weeks of CIA exposure (27).

This limitation highlights a long-standing issue in preclinical research, namely that there

is a dearth of information regarding female behavior and neurobiology compared to males, and

that most animal models, including CIA, have been built using only male subjects. The

outcomes are often different when these models are retroactively applied to females. This

doesn't necessarily mean that female rats don't experience alcohol withdrawal or dependence,

but that the phenotype may be different in females or that the way we measure behavior is not

adequate to capture such phenotypes in female rodents, as has been observed with other

behaviors (48,49). Our ongoing and future work aims to tackle some of these issues by

determining whether females show escalation after longer periods of CIA exposure, require a

different withdrawal period, and/or require more or longer operant sessions.

A second limitation is that we were unable to predict current drinking behavior using

LFPs from the tethered self-administration sessions. Tethering animals significantly reduced

drinking, to 0 in many cases, despite attempts to habituate animals to the tether. Alcohol

drinking behavior, especially during withdrawal, is highly sensitive to stressors that often

suppress rather than enhance drinking (50). Our future work aims to rectify this problem by

increasing the number of habituation sessions or using wireless recording equipment that would

eliminate the need for a tether.

Overall, the current study has allowed us to determine whether and how cortical, striatal,

and limbic oscillations are impacted by CIA in males and females, as well as which specific LFP

features correlate with drinking. Based on these findings, our future work will determine whether

the CIA-induced changes in LFPs observed here are functionally relevant. While LFPs are

thought to reflect neuronal activity, the precise nature of the relationship between what is

happening at a cellular level and LFPs is poorly understood (51). Our future work will use

chemogenetic and optogenetic techniques to selectively target cortical-striatal vs. cortical-limbic

circuits in CIA-exposed animals to determine whether manipulation of these circuits will reduce

drinking and other measures of dependence in a sex-specific manner, while also ameliorating

the effect of CIA on LFPs.

CONCLUSIONS

Although it is very probable that CIA-induced changes to cortical-striatal and cortical-

limbic circuits play a role in both male and female alcohol misuse, the current experiment

supports the notion that the neural circuits *most* impacted by chronic alcohol exposure are sex-

specific. This study also highlights the vital need to incorporate both sexes into preclinical work,

enhancing our ability to translate preclinical findings to clinical populations. For example, DBS

targeting the NAc is currently going through clinical trials for treatment-resistant AUD, but

previous preclinical and clinical DBS studies were performed almost exclusively in males (52),

and targeting different circuits might better benefit women. Understanding that limbic regions

might play a stronger role in driving alcohol misuse in females could provide an important

biological treatment target for future studies aimed at developing more efficacious therapies for

AUD in both sexes.

DECLARATIONS

Ethics approval and consent to participate

Consent for publication

Not applicable.

Availability of data and material

Data sets generated during the current study are available from the corresponding author on reasonable request. Matlab codes to generate the predictive models are available on GitHub: https://github.com/lucasdwi/code/tree/greenlab.

Competing interests

The authors declare that they have no competing interests.

Funding

Authors' contributions

KH assisted in developing the research design, collected and analyzed the data, and drafted the manuscript. SN and MP assisted in collecting the data and drafting the manuscript. LD assisted in data analysis and drafting the manuscript. AH developed the research question and research design, assisted in analyzing the data, and drafted and edited the manuscript.

Acknowledgements

Not applicable.

FIGURE CAPTIONS

Figure 1: Electrode schematic (A) and example placements in the IL (B), NAcSh (C) and CeA

(D).

Figure 2: Experimental timeline.

Figure 3: Average g/kg of alcohol consumed in male (A) and female (B) rats exposed to CIA or

room air (control) during 30-minute self-administration sessions before (pre-CIA baseline) and

after (acute withdrawal) four weeks of CIA exposure. CIA led to increased drinking in the last

two self-administration sessions in males (*p<0.05; n=6-7/group), but no changes in female

drinking (p>.05; n=7/group).

Figure 4: Average area under the receiver operating characteristic curve (AUC) +/- SD for real

and permuted binary prediction models (n = 6-7/sex × 3 recordings/rat) in males (A) and

females (B). Models built from IL and NAcSh LFPs best predicted whether males were exposed

to CIA. Models built from IL and CeA LFPs best predicted whether females were exposed to

CIA.

Figure 5: Correlations between average g/kg of alcohol consumed during acute withdrawal in

CIA and control rats and average power or coherence across sessions. In males (A), CIA

significantly moderated the correlation between alcohol consumption and right NAcSh hγ power,

with power and g/kg consumed negatively correlated in CIA males. In females (B), CIA

significantly moderated the correlation between alcohol consumption and L-R NAcSh β

coherence, with coherence and g/kg consumed negatively correlated in control females.

[+]p<0.08; *p<0.05, **p<0.01.

Table 1. Neural features that best predict CIA exposure

Males		Females	
Feature	Mean AUC	Feature	Mean AUC
IL_R - $NAcSh_L$ lγ	0.75	CeA_L α	0.87
$NAcSh_R$ hγ	0.75	CeA_R α	0.81
IL_R δ	0.74	CeA_L hγ	0.80
IL_L - CeA_L θ	0.73	$NAcSh_L$ – $NAcSh_R$ β	0.79
IL_L δ	0.72	CeA_R δ	0.76

The top 5 LFP features used in binary models predicting CIA exposure in males and females. Frequency bands [delta (δ), theta (θ), alpha (α), beta (β), low gamma (lγ), and high gamma (hγ)] are described for power features within and coherence features between neural sites. Subscript indicates left (L) or right (R) hemisphere.

Table 2. Correlations between average neural feature and g/kg consumed

Males			Females		
Feature	Control	CIA	Feature	Control	CIA
IL_R - $NAcSh_L$ lγ	0.42	0.48	CeA_L α	0.01	0.21
$NAcSh_R$ hγ	0.41	0.83*	CeA_R α	0.62	0.20
IL_R δ	0.24	0.36	CeA_L hγ	0.03	0.52
IL_L - CeA_L θ	0.29	0.49	$NAcSh_L$ − $NAcSh_R$ β	0.87**	0.48
IL_L δ	0.37	0.14	CeA_R δ	0.07	0.71[+]

Note: [+]p<0.08; *p<0.05, **p<0.01

Correlation coefficients for average g/kg of alcohol consumed in CIA and control rats and average power or coherence across sessions. Frequency bands [delta (δ), theta (θ), alpha (α), beta (β), low gamma (lγ), and high gamma (hγ)] are described for power features within and coherence features between neural sites. Subscript indicates left (L) or right (R) hemisphere.

References

1.   Griswold MG, Fullman N, Hawley C, Arian N, Zimsen SRM, Tymeson HD, et al. Alcohol use
and burden for 195 countries and territories, 1990–2016: a systematic analysis for the
Global Burden of Disease Study 2016. The Lancet. 2018 Sep;392(10152):1015–35.

2.   National Survey on Drug Use and Health 2021 [Internet]. U.S. Department of Health and
Human Services, Substance Abuse and Mental Health Services Administration, Center for
Behavioral Health Statistics and Quality; 2022. Available from:
https://www.samhsa.gov/data/sites/default/files/reports/rpt39441/NSDUHDetailedTabs2021/
NSDUHDetailedTabs2021/NSDUHDetTabsSect5pe2021.htm

3.   Witkiewitz K, Litten RZ, Leggio L. Advances in the science and treatment of alcohol use
disorder. Sci Adv. 2019 Sep 6;5(9):eaax4043.

4.   Batra A, Müller CA, Mann K, Heinz A. Alcohol Dependence and Harmful Use of Alcohol:
Diagnosis and Treatment Options. Dtsch Ärztebl Int [Internet]. 2016 Apr 29 [cited 2024 Mar
18]; Available from: https://www.aerzteblatt.de/10.3238/arztebl.2016.0301

5.   Erol A, Karpyak VM. Sex and gender-related differences in alcohol use and its
consequences: Contemporary knowledge and future research considerations. Drug Alcohol
Depend. 2015 Nov;156:1–13.

6.   Grant BF, Goldstein RB, Saha TD, Chou SP, Jung J, Zhang H, et al. Epidemiology of *DSM-
5* Alcohol Use Disorder: Results From the National Epidemiologic Survey on Alcohol and
Related Conditions III. JAMA Psychiatry. 2015 Aug 1;72(8):757.

7.   White A. Gender Differences in the Epidemiology of Alcohol Use and Related Harms in the
United States. Alcohol Res Curr Rev. 2020;40(2):01.

8.   Canidate SS, Carnaby GD, Cook CL, Cook RL. A Systematic Review of Naltrexone for
Attenuating Alcohol Consumption in Women with Alcohol Use Disorders. Alcohol Clin Exp
Res. 2017 Mar;41(3):466–72.

9.   Garbutt JC, Kranzler HR, O'Malley SS, Gastfriend DR, Pettinati HM, Silverman BL, et al.
Efficacy and Tolerability of Long-Acting Injectable Naltrexone for Alcohol Dependence: A
Randomized Controlled Trial. JAMA. 2005 Apr 6;293(13):1617.

10. Karp NA, Reavey N. Sex bias in preclinical research and an exploration of how to change
the status quo. Br J Pharmacol. 2019 Nov;176(21):4107–18.

11. Henricks AM, Sullivan EDK, Dwiel LL, Keus KM, Adner ED, Green AI, et al. Sex differences
in the ability of corticostriatal oscillations to predict rodent alcohol consumption. Biol Sex
Differ. 2019 Dec;10(1):61.

12. Howe MW, Atallah HE, McCool A, Gibson DJ, Graybiel AM. Habit learning is associated
with major shifts in frequencies of oscillatory activity and synchronized spike firing in
striatum. Proc Natl Acad Sci. 2011 Oct 4;108(40):16801–6.

13. Pesaran B, Vinck M, Einevoll GT, Sirota A, Fries P, Siegel M, et al. Investigating large-scale
brain dynamics using field potential recordings: analysis and interpretation. Nat Neurosci.
2018 Jul;21(7):903–19.

14. Tort ABL, Kramer MA, Thorn C, Gibson DJ, Kubota Y, Graybiel AM, et al. Dynamic cross-
frequency couplings of local field potential oscillations in rat striatum and hippocampus
during performance of a T-maze task. Proc Natl Acad Sci. 2008 Dec 23;105(51):20517–22.

15. Haber SN. Corticostriatal circuitry. Dialogues Clin Neurosci. 2016 Mar 31;18(1):7–21.

16. Peltier MR, Verplaetse TL, Mineur YS, Petrakis IL, Cosgrove KP, Picciotto MR, et al. Sex
differences in stress-related alcohol use. Neurobiol Stress. 2019 Feb;10:100149.

17. Logrip ML, Milivojevic V, Bertholomey ML, Torregrossa MM. Sexual dimorphism in the
neural impact of stress and alcohol. Alcohol. 2018 Nov;72:49–59.

18. Henricks AM, Berger AL, Lugo JM, Baxter-Potter LN, Bieniasz KV, Craft RM, et al. Sex
differences in alcohol consumption and alterations in nucleus accumbens endocannabinoid
mRNA in alcohol-dependent rats. Neuroscience. 2016 Oct;335:195–206.

19. Williams AM, Reis DJ, Powell AS, Neira LJ, Nealey KA, Ziegler CE, et al. The effect of
intermittent alcohol vapor or pulsatile heroin on somatic and negative affective indices
during spontaneous withdrawal in Wistar rats. Psychopharmacology (Berl). 2012
Sep;223(1):75–88.

20. Henricks AM, Dwiel LL, Deveau NH, Simon AA, Ruiz-Jaquez MJ, Green AI, et al.
Corticostriatal Oscillations Predict High vs. Low Drinkers in a Rat Model of Limited Access
Alcohol Consumption. Front Syst Neurosci. 2019 Aug 13;13:35.

21. Dwiel LL, Khokhar JY, Connerney MA, Green AI, Doucette WT. Finding the balance
between model complexity and performance: Using ventral striatal oscillations to classify
feeding behavior in rats. Cohen KB, editor. PLOS Comput Biol. 2019 Apr
22;15(4):e1006838.

22. Catanese J, Carmichael JE, Van Der Meer MAA. Low- and high-gamma oscillations deviate
in opposite directions from zero-phase synchrony in the limbic corticostriatal loop. J
Neurophysiol. 2016 Jul 1;116(1):5–17.

23. McCracken CB, Grace AA. Nucleus Accumbens Deep Brain Stimulation Produces Region-
Specific Alterations in Local Field Potential Oscillations and Evoked Responses *In Vivo*. J
Neurosci. 2009 Apr 22;29(16):5354–63.

24. Almeida OF, Shoaib M, Deicke J, Fischer D, Darwish MH, Patchev VK. Gender differences
in ethanol preference and ingestion in rats. The role of the gonadal steroid environment. J
Clin Invest. 1998 Jun 15;101(12):2677–85.

25. Flores-Bonilla A. Sex Differences in the Neurobiology of Alcohol Use Disorder. Alcohol Res
Curr Rev. 2020;40(2):03.

26. Morales M, McGinnis MM, McCool BA. Chronic ethanol exposure increases voluntary home
cage intake in adult male, but not female, Long–Evans rats. Pharmacol Biochem Behav.
2015 Dec;139:67–76.

27. Priddy BM, Carmack SA, Thomas LC, Vendruscolo JCM, Koob GF, Vendruscolo LF. Sex,
strain, and estrous cycle influences on alcohol drinking in rats. Pharmacol Biochem Behav.
2017 Jan;152:61–7.

28. Torres OV, Walker EM, Beas BS, O'Dell LE. Female Rats Display Enhanced Rewarding
Effects of Ethanol That Are Hormone Dependent. Alcohol Clin Exp Res. 2014
Jan;38(1):108–15.

29. Roberts AJ, Heyser C, Cole M, Griffin P, Koob G. Excessive Ethanol Drinking Following a
History of Dependence Animal Model of Allostasis. 2000 Jun;22(6):581–94.

30. Roberts AJ, Cole M, Koob GF. Intra-amygdala Muscimol Decreases Operant Ethanol
Self-administration in Dependent Rats. Alcohol Clin Exp Res. 1996 Oct;20(7):1289–98.

31. Quijano Cardé NA, De Biasi M. Behavioral characterization of withdrawal following chronic
voluntary ethanol consumption via intermittent two-bottle choice points to different
susceptibility categories. Alcohol Clin Exp Res. 2022 Apr;46(4):614–27.

32. Bach P, Luderer M, Müller UJ, Jakobs M, Baldermann JC, Voges J, et al. Deep brain
stimulation of the nucleus accumbens in treatment-resistant alcohol use disorder: a double-
blind randomized controlled multi-center trial. Transl Psychiatry. 2023 Feb 8;13(1):49.

33. Davidson B, Giacobbe P, George TP, Nestor SM, Rabin JS, Goubran M, et al. Deep brain
stimulation of the nucleus accumbens in the treatment of severe alcohol use disorder: a
phase I pilot trial. Mol Psychiatry. 2022 Oct;27(10):3992–4000.

34. Volkow ND, Wang GJ, Fowler JS, Tomasi D, Telang F. Addiction: Beyond dopamine reward
circuitry. Proc Natl Acad Sci. 2011 Sep 13;108(37):15037–42.

35. Seo D, Jia Z, Lacadie CM, Tsou KA, Bergquist K, Sinha R. Sex differences in neural
responses to stress and alcohol context cues. Hum Brain Mapp. 2011 Nov;32(11):1998–
2013.

36. Morra JT, Glick SD, Cheer JF. Cannabinoid receptors mediate methamphetamine induction
of high frequency gamma oscillations in the nucleus accumbens. Neuropharmacology. 2012
Sep;63(4):565–74.

37. Tsurugizawa T, Abe Y, Le Bihan D. Water apparent diffusion coefficient correlates with
gamma oscillation of local field potentials in the rat brain nucleus accumbens following
alcohol injection. J Cereb Blood Flow Metab. 2017 Sep;37(9):3193–202.

38. Van Der Meer M. Integrating early results on ventral striatal gamma oscillations in the rat.
Front Neurosci [Internet]. 2010 [cited 2024 Apr 1]; Available from:
http://journal.frontiersin.org/article/10.3389/fnins.2010.00300/abstract

39. Gilpin NW, Herman MA, Roberto M. The Central Amygdala as an Integrative Hub for
Anxiety and Alcohol Use Disorders. Biol Psychiatry. 2015 May;77(10):859–69.

40. Koob GF, Volkow ND. Neurocircuitry of Addiction. Neuropsychopharmacology. 2010
Jan;35(1):217–38.

41. Roberto M, Kirson D, Khom S. The Role of the Central Amygdala in Alcohol Dependence.
Cold Spring Harb Perspect Med. 2021 Feb;11(2):a039339.

42. Breese GR, Overstreet DH, Knapp DJ. Conceptual framework for the etiology of alcoholism:
a□?kindling?/stress hypothesis. Psychopharmacology (Berl). 2005 Apr;178(4):367–80.

43. Meyers JL, Zhang J, Manz N, Rangaswamy M, Kamarajan C, Wetherill L, et al. A genome
wide association study of fast beta EEG in families of European ancestry. Int J
Psychophysiol. 2017 May;115:74–85.

44. Stenner MP, Dürschmid S, Rutledge RB, Zaehle T, Schmitt FC, Kaufmann J, et al.
Perimovement decrease of alpha/beta oscillations in the human nucleus accumbens. J
Neurophysiol. 2016 Oct 1;116(4):1663–72.

45. Hadar R, Voget M, Vengeliene V, Haumesser JK, Van Riesen C, Avchalumov Y, et al.
Altered neural oscillations and elevated dopamine levels in the reward pathway during
alcohol relapse. Behav Brain Res. 2017 Jan;316:131–5.

46. Walker BM. Conceptualizing withdrawal-induced escalation of alcohol self-administration as
a learned, plasticity-dependent process. Alcohol. 2012 Jun;46(4):339–48.

47. Carroll ME, Anker JJ. Sex differences and ovarian hormones in animal models of drug
dependence. Horm Behav. 2010 Jun;58(1):44–56.

48. Shansky RM. Sex differences in behavioral strategies: avoiding interpretational pitfalls. Curr
Opin Neurobiol. 2018 Apr;49:95–8.

49. Shansky RM, Murphy AZ. Considering sex as a biological variable will require a global shift
in science culture. Nat Neurosci. 2021 Apr;24(4):457–64.

50. Wille-Bille A, Ferreyra A, Sciangula M, Chiner F, Nizhnikov ME, Pautassi RM. Restraint
stress enhances alcohol intake in adolescent female rats but reduces alcohol intake in
adolescent male and adult female rats. Behav Brain Res. 2017 Aug;332:269–79.

51. Herreras O. Local Field Potentials: Myths and Misunderstandings. Front Neural Circuits
[Internet]. 2016 Dec 15 [cited 2023 Jun 12];10. Available from:
http://journal.frontiersin.org/article/10.3389/fncir.2016.00101/full

52. Chang R, Peng J, Chen Y, Liao H, Zhao S, Zou J, et al. Deep Brain Stimulation in Drug
Addiction Treatment: Research Progress and Perspective. Front Psychiatry. 2022 Apr
7;13:858638.

A **Males**

B **Females**

A **Males**

IL, NAcSh, and CeA Oscillations

Real AUC: 0.63 ± 0.04
Perm AUC: 0.52 ± 0.04

IL and NAcSh Oscillations

Real AUC: 0.78 ± 0.04
Perm AUC: 0.49 ± 0.04

IL and CeA Oscillations

Real AUC: 0.65 ± 0.04
Perm AUC: 0.48 ± 0.03

B **Females**

IL, NAcSh, and CeA Oscillations

Real AUC: 0.81 ± 0.03
Perm AUC: 0.50 ± 0.03

IL and NAcSh Oscillations

Real AUC: 0.57 ± 0.04
Perm AUC: 0.49 ± 0.03

IL and CeA Oscillations

Real AUC: 0.83 ± 0.03
Perm AUC: 0.49 ± 0.03