Temperature and Pressure Shaped the Evolution of Antifreeze Proteins in Polar and Deep Sea Zoarcoid Fishes

Authors: Samuel N. Bogan[1], Nathan Surendran[1], Scott Hotaling[2], Thomas Desvignes[3], Iliana Bista[4,5,6], Luana S.F. Lins[7], Mari H. Eilertsen[8], Nathalie R. Le François[9], Tait Algayer[10], Scott L. Hamilton[11], Paul B. Frandsen[12], Federico G. Hoffmann[13,14], and Joanna L. Kelley[1]

Affiliations:
[1] Department of Ecology and Evolutionary Biology, University of California Santa Cruz, Santa Cruz, CA USA
[2] Department of Watershed Sciences, Utah State University, Logan, UT, USA
[3] Institute of Neuroscience, University of Oregon, Eugene, OR, USA
[4] LOEWE Centre for Translational Biodiversity Genomics, Frankfurt, 60325, Germany
[5] Senckenberg Research Institute and Natural History Museum, Frankfurt, 60325, Germany
[6] Wellcome Sanger Institute, Tree of Life, Wellcome Genome Campus, Hinxton CB10 1SA, United Kingdom
[7] CSIRO, Health and Biosecurity, Canberra, Australia
[8] Department of Biological Sciences, Centre for Deep Sea Research, University of Bergen, Norway
[9] Laboratoire de Physiologie et Aquaculture de la Conservation (LPAC), Division des collections vivantes, de la conservation et de la recherche, Biodôme de Montréal/ Espace pour la vie, 4777, av. Pierre-De Coubertin, Montréal, Québec H1V 1B3
[10] School of Biological Sciences, Washington State University, Pullman, WA, USA
[11] Moss Landing Marine Laboratories, Moss Landing, CA, USA
[12] Department of Plant and Wildlife Sciences, Brigham Young University, Provo, UT, USA
[13] Department of Biochemistry, Molecular Biology, Entomology, and Plant Pathology, Mississippi State University, Mississippi State, MS, USA
[14] Institute for Genomics, Biocomputing and Biotechnology, Mississippi State University, Mississippi State, MS, USA

Keywords: Long-read sequencing; antifreeze protein; adaptation; genomics; temperature; gene copy number

Running head: Long reads reveal antifreeze protein evolution

Abstract

Antifreeze proteins (AFPs) have enabled teleost fishes to repeatedly colonize polar seas.

Four AFP types have convergently evolved in several fish lineages. AFPs inhibit ice crystal growth

and lower cellular freezing point. In lineages with AFPs, species inhabiting colder environments

may possess more AFP copies. Elucidating how differences in AFP copy number evolve is

challenging due to the genes' tandem array structure and consequently poor resolution of these

repetitive regions. Here we explore the evolution of type III AFPs (AFP III) in the globally

distributed suborder Zoarcoidei, leveraging six new long-read genome assemblies. Zoarcoidei has

fewer genomic resources relative to other polar fish clades while it is one of the few groups of

fishes adapted to both the Arctic and Southern Oceans. Combining these new assemblies with

additional long-read genomes available for Zoarcoidei, we conducted a comprehensive

phylogenetic test of AFP III evolution and modeled the effects of thermal habitat and depth on

AFP III gene family evolution. We confirm a single origin of AFP III via neofunctionalization of

the enzyme sialic acid synthase B and show that AFP gains and losses were not only correlated

with temperature, but also with depth, which reduces freezing point via pressure. Associations

between the environment and AFP III copy number were driven by duplications of paralogs that

were translocated out of the ancestral locus at which Zoarcoidei AFP arose. Our results reveal

novel environmental effects on AFP evolution and demonstrate the value of high-quality genomic

resources for studying how structural genomic variation shapes convergent adaptation.

1. Introduction

The evolution of gene family copy number variation can contribute to environmental adaptation (Schmidt et al. 2010) and be stimulated by environmental change (Hull et al. 2017). Studying the causes and consequences of copy number's evolution is difficult when duplications produce large tandem repeats or affect synteny, but the advent of long-read sequencing has helped resolve repetitive variants clouding the origins of copy number evolution (Marx 2023). The evolution of antifreeze proteins (AFPs) and antifreeze glycoproteins (AFGPs) are remarkable examples of convergent environmental adaptation in ectotherms living at low temperatures (A. L. DeVries 1971; Baskaran et al. 2021). Increased AFP 'dosage' driven by gene duplication can facilitate adaptation to chronic cold (Desjardins et al. 2012). In theory, temperate species derived from cold-adapted ancestors should undergo relaxed selection on AFP copy number and subsequent contractions (Hayes, Davies, and Fletcher 1991; Hew et al. 1988). Studying contractions and expansions of AFPs across thermally-distributed clades is critical for understanding thermal adaptation by ectotherms but has been hampered by poor resolution of their copy number due to low quality reference genome assemblies.

Antifreeze proteins were first discovered in Antarctic notothenioid fish in 1971, which possess antifreeze glycoproteins (AFGPs) and lower both the freezing and boiling point of cellular fluid (A. L. DeVries 1971; Raymond and DeVries 1977; A. L. DeVries and Wohlschlag 1969). Since this discovery, AFPs that induce thermal hysteresis, an increase in degrees of difference between a fluid's freezing and boiling points (Barrett 2001), have been discovered in microbes (Gilbert, Davies, and Laybourn-Parry 2005), plants (Gupta and Deswal 2014), invertebrates (Duman 2015), and vertebrates (Arthur L. DeVries and Cheng 2005). AFPs generally function by binding to the leading edges of newly-crystalized nuclei of ice, attenuating their expansion and

reducing cellular freezing point (Jia et al. 1996). In fishes, AFPs have independently evolved at

least three times (C. Cheng and Devries 2002). These convergently evolved AFPs make up four

known categories termed AFGPs, and AFP types I - III. The evolution of AFPs by ectotherms is a

classical example of environmental adaptation, but we still know little about the environmental

processes that shaped their evolution and the diversity and variation they exhibit across species

and their genomes.

It has been repeatedly suggested that, among lineages possessing AFPs, species or

populations inhabiting colder environments should exhibit more AFP copies (Hew et al. 1988;

Hayes, Davies, and Fletcher 1991; Kelley et al. 2010). Comparative studies of AFP copy number

in population- and species-pairs of fishes from nominally warmer versus colder habitats have

confirmed this expectation (Hayes, Davies, and Fletcher 1991; Desjardins et al. 2012). Studying

AFGPs in Arctic cod and Antarctic notothenioids, Baalsrud et al. (2018) observed losses of AFGPs

in secondarily sub-polar species of both groups. Long-read sequencing of Notothenioids has

revealed pseudogenization and contraction of AFGP copies in individual species inhabiting milder

latitudes suggestive of relaxed selection on antifreeze function (Bista et al. 2023; Rivera-Colón et

al. 2023). However, an analysis of AFP copy number across continuous thermal clines has not

been reported. Whether and how AFP copy number and gene dosage scales with freezing risk

remains unknown. One environmental variable that has been proposed to affect freezing risk and

AFP evolution is pressure. Because increased pressure in seawater affects the stability of ice

crystals and decreases phase change temperatures, lowering freezing point (Fujino, Lewis, and

Perkin 1974), it has been proposed that polar fishes inhabiting different depths and exhibiting

divergence in AFP copy number have experienced differential positive selection for AFP copy

number (Desjardins et al. 2012).

AFPs of bony fishes have evolved from neofunctionalization, and AFGPs by both

neofunctionalization and *de novo* gene birth. These events were driven in part by repetitive

genomic regions that provided substrate for *de novo* gene birth of an AFGP in cods (Baalsrud et

al. 2018; Zhuang et al. 2019) and translocations of ancestral genes that underwent

neofunctionalization and expanded into tandem arrays of AFPs (Deng et al. 2010; Hotaling et al.

2023). In addition to influencing AFP origins, structural variation such as translocations,

inversions, and repetitive element content have the potential to influence expansions and

contractions of AFPs across environmental gradients (Lauer and Gresham 2019). For example,

cytogenetically-balanced and unbalanced translocations can both facilitate variation in gene copy

number (Harewood et al. 2010; S. K. Watson et al. 2007). The accurate detection of translocations

and measurement of gene copy number in repetitive regions require long-read sequencing

approaches, which enable identification of structural variants such as translocations and better

resolve complex genomic regions relative to short read sequencing (Marx 2023). Long-read

sequencing has shown success in assembling repetitive AFGP arrays in notothenioid fishes (Bista

et al. 2023).

The bony fish suborder Zoarcoidei is an ideal system in which to study the joint effects of

environmental and structural genomic variation on AFP evolution. Zoarcoidei likely originated in

north temperate seas, followed by stepwise invasions of (i) the Arctic Ocean, (ii) south temperate

seas and (iii) the Southern Ocean (Hotaling et al. 2021) where minimum temperatures can fall

below -2.0 °C (Auger et al. 2021; Fetterer 2017). Sanger sequencing of BAC clones in the zoarcoid

Lycodichthys dearborni suggested that type III AFP arose within Zoarcoidei via

neofunctionalization of the enzyme sialic acid synthase B (*sasb*). *sasb* likely duplicated before

undergoing deletions of all exons except 1 and 6, which are homologous to AFP III exons 1 and 2

(Deng et al. 2010). Zoarcoidei repeatedly invaded and adapted to the poles (Hotaling et al. 2021)

while Zoarcoidei AFP III likely evolved once (Deng et al. 2010; Hobbs et al. 2020). AFP III copies

have likely expanded and contracted multiple times in Zoarcoidei (Hotaling et al. 2023), but the

extent of gene family evolution of AFP III in zoarcoids, of AFPs across fishes, and how genomic

and environmental variation interactively affected AFP evolution remain unclear. Zoarcoidei

diversified across depth faster than any group of fishes (Friedman and Muñoz 2023), potentiating

a complex landscape of selection for AFP function and dosage across the suborder's distributions

through latitude and depth. Previously identified tandem arrays of zoarcoid AFP III exhibited

conserved synteny with several flanking genes (Hotaling et al. 2023), enabling identification of

translocations out of this region.

Here we report high-quality, long-read genome assemblies from six species of Zoarcoidei

from diverse thermal habitats and depths (Fig. 1; Table 1). We integrated annotations of AFP III

genes with oceanographic measures of thermal minima, pressure, and depth within species'

habitats to test whether temperature and pressure are associated with AFP III copy number

evolution. Using analyses of synteny, we traced AFP III copies to ancestral and translocated arrays

and determined whether translocation drove AFP expansions under high freezing risk. These new

reference genome assemblies provide a valuable resource for studying how abiotic environments

and structural variation jointly influence convergent adaptation.

2. Results and Discussion

2.1. Genome assembly - Reference genomes were assembled with Pacific Biosciences

(PacBio) long-read sequence data for six zoarcoid fishes spanning a range of temperatures and

depths: the ronquil *Bathymaster signatus*, the wolffishes *Anarhichas lupus* and *Anarhichas minor*,

and the eelpouts *Lycenchelys platyrhina*, *Lycodes diapterus*, and *Ophthalmolycus amberensis*.

These species inhabit polar seas with latitudes as northerly as 73.0 °N and southerly as -65.8 °S.

Assembled genomes included temperate species with absolute latitudes as low as 37.8 ° (Fig. 1).

Species' maximum depths spanned 24 - 2,229 meters. These mark the first long-read genome

assemblies for the families Bathymasteridae (ronquils) and Anarhichadidae (wolffishes), and

increase the number of long-read genome assemblies among Zoarcidae (eelpouts) by 2.5x.

Genome sizes, sequencing coverage, N50 values, BUSCO complete percentages, and

percent repetitive elements are reported in Table 1 for the six new assemblies, five existing

zoarcoid long-read assemblies (Bista, Collins, and Wellcome Sanger Institute Tree of Life

Management, Samples and Laboratory team 2024), and one scorpaeniform outgroup. Genome

sizes of all Zoarcoidei examined equaled a mean of 647.80 ± 74.3 Mb. Genome assemblies

reported here averaged 11.7x greater coverage compared to publicly available, long-read zoarcoid

assemblies. Contig N50 values were an average of 6.5 Mb longer than available assemblies (Table

1). As expected, repetitive element content linearly and positively correlated with genome size

across all species (Kidwell 2002) as shown in Fig. S1 (r^2 = 0.93).

Zoarcoidei families are among the few fish groups inhabiting both the Arctic and Southern

Oceans (Møller, Nielsen, and Anderson 2005), positioning this suborder to be a valuable system

in which to study convergent evolution and adaptation to extreme cold. However, a paucity of

genomic resources for zoarcoid fishes stands as an obstacle to their value in research. The

improved reference genome quality of these new assemblies among other members of Zoarcoidei

and the extension of thermal habitats and depths represented among them significantly strengthen

Zoarcoidei as a tractable, valuable system in which to study convergent adaptation.

Species	Genome size (Mb)	Sequencing coverage	Contig N50 (Mb)	Scaffold N50 (Mb)	% Repetitive elements	BUSCO % complete (*n* orthologs)
Anarhichas lupus	627.7	80x	23.8	*na*	27.0%	98.5% (3587)
Anarhichas minor	643.9	74x	13.2	*na*	28.0%	98.5% (3587)
Bathymaster signatus	714.3	42x	3.1	*na*	32.2%	98.3% (3587)
Lycodes diapterus	711.9	43x	6.0	*na*	34.1%	98.5% (3585)
Lycenchelys platyrhina	718.5	34x	2.8	*na*	31.0%	98.3% (3579)
Ophthalmolycus amberensis	712.6	95x	5.8	*na*	30.6%	98.5% (3586)
Cebidichthys violaceus	575.7	31x	1.0	16.4	25.4%	93.3% (3396)
Gasterosteus aculeatus*	471.9	75x	0.5	20.5	18.6%	97.0% (3534)
Leptoclinus maculatus	699.9	65x	2.4	23.5	32.8%	98.0% (3569)
Lycodes pacificus	646.4	68x	4.7	27.7	29.0%	98.1% (3573)
Melanostigma gelatinosum	662.0	39x	1.9	28.0	29.3%	98.1% (3569)
Pholis gunnellus	588.7	45x	3.5	25.4	23.3%	98.6% (3591)

Table 1 | *Genome assembly metrics.* Assemblies reported by the authors are shaded in gray. Publicly available zoarcoid genomes leveraged in this study are unhighlighted. A non-zoarcoid outgroup is designated by an asterisk. BUSCO analysis was performed on universal single-copy orthologs of Actinopterygii.

2.2. Variation in type III antifreeze protein copy number - The number of AFP III copies

in the 11 zoarcoids studied ranged from 0 to 42 putatively functional copies. Between 3 – 16

pseudogenized AFP III copies were detected across all zoarcoids (Fig. 1A). The copy number of

putatively-functional and pseudogenized AFP genes correlated non-linearly such that pseudogenes

increased in species with more AFP III toward a plateau (Fig. S2; 2nd order polynomial $r^2 = 0.31$).

This association was explained by phylogeny and became insignificant after controlling for species

relatedness (95% posterior interval = -1.54 – 0.67). The presence of pseudogenized AFP III copies

in all zoarcoids indicates that the gene's origin predates the most-recent common ancestor of

zoarcoids included in this study (Fig. 1A).

We evaluated whether AFP III may have arisen in a common ancestor predating Zoarcoidei

by screening for the gene in long-read assemblies of 12 species of order Scorpaeniformes

representing 7 families and found no evidence of complete or partial AFP III paralogs (Table S1).

Past phylogenetic analyses of zoarcoid AFP III also detected a partial, pseudogenized AFP III

sequence in Bathymasteridae, the earliest-branching family of Zoarcoidei (Hobbs et al. 2020).

However, Hobbs et al. (2020) and others have not ruled out an emergence of AFP III prior to the

branching of Zoarcoidei. Our results therefore strengthen evidence that Zoarcoidei AFP III likely

arose in an early branch of Bathymasteridae as opposed to a common ancestor within

Scorpaeniformes predating Zoarcoidei. Long-read sequencing of Zoarcoidei offers the potential to

trace the origin of AFP III within Zoarcoidei by mapping structural variants contributing to the

gene's evolution. These events include duplication-translocation of *sasb* and deletions of *sasb*

exons 2 – 5.

212
213

Figure 1 | *Species' phylogeny, AFP copy numbers, and distributions.* (A) A species tree of Zoarcoidei included in this study including outgroup *Gasterosteus aculeatus* is plotted adjacent to AFP copy number variation colored by pseudogenization or location of homologs in the ancestral syntenic region or translocation(s). The red diamond signifies the origin of type III AFP. Species highlighted in blue denote genome assemblies reported by the authors. The scale bar denotes substitutions per amino acid. *Ophthalmolycus amberensis* and *Lycodes diapterus* branch tip lengths equal 0.000069 and 0.000074 substitutions per amino acid, respectively. (B) Species' distributions are plotted as the point of maximum likelihood of occurrence from AquaMaps standardized distributions. The size of white circles around occurrences are scaled according to species abundance. Mean sea surface temperature (SST, °C) recorded in August, 2020, is visualized according to color. Illustrations by SNB.

2.3. Environmental effects on type III antifreeze protein copy number - Habitat temperature

and pressure jointly influence freezing risk (Fujino, Lewis, and Perkin 1974) and were additively

correlated with AFP copy number. A phylogenetically corrected Bayesian model revealed that

AFP copy number significantly increased with decreases in (i) thermal minima of species' habitats

and (ii) mean pressure of species' depth ranges (Fig. 2). Thermal minima equaled the lowest

recorded temperature at a species' coordinate of maximum likelihood of occurrence within their

depth range. Pressure was derived from habitat depth and latitude. Thermal minima of species

habitats spanned -3.00 – 7.21 °C (Fig. S2). AFP copy number significantly increased in species

occupying lower thermal minima (Fig. 2A). This effect was significant after controlling for

phylogeny (scaled effect = -1.55 ± 0.97).

AFP III gene copy number declined in species inhabiting greater pressure (Fig. 2B).

Pressure had a stronger effect on AFP copy number than temperature and was non-linear,

occupying a negative exponential curve. Pressure affected AFP copy number after controlling for

habitat depth as well as phylogeny (scaled linear effect = -4.72 ± 2.58). As expected, habitat

temperatures correlated with depth and pressure (Fig. S4 – S5). To account for this autocorrelation,

predicted effects of thermal minima and pressure on AFP copy number were modeled in a

statistically independent manner.

The finding that AFP III copy number is reduced in deep sea species experiencing high

pressure is pertinent to the biogeography of Zoarcoidei. Zoarcoidei includes the eelpout family

Zoarcidae, which has exhibited faster rates of evolution across depth than any group of fishes

(Friedman and Muñoz 2023). Associations between depth and AFP copy number were previously

proposed by Desjardins et al., who observed that the wolffishes *A. lupus* and *A. minor* exhibit

higher AFP copy number in the shallow water species *A. lupus* rather than the species occupying

a colder thermal environment (Desjardins et al. 2012). Our results mark the first quantitative

evidence that selection on antifreeze proteins across thermal environments was shaped by depth

and pressure.

Figure 2 | *Associations between temperature, pressure, and type III antifreeze protein copy number.*
Relationships between AFP III copy number and (A) thermal minima of species habitats and (B) mean
pressure of species' habitats are plotted. Fitted lines represent fit from a phylogenetically controlled model
of type III antifreeze protein copy number.
258

A direct explanation of pressure's association with AFP copy number is that increased

energy imposed by pressure lowers the freezing point of intracellular fluids (Fujino, Lewis, and

Perkin 1974). An indirect explanation also exists that is not mutually exclusive to this direct

hypothesis. Deep sea fishes generally regulate the hydrostatic pressure of cells through

osmoregulation such that increased pressure is countered by greater intracellular osmolality

(Yancey, Blake, and Conley 2002; Gillett et al. 1997; Yancey et al. 2014). Because increased

osmolality also reduces the freezing point of fluids (Bodnar 1993; Potter, Clynne, and Brown

1978), it is possible that pressure and osmolality additively affect cellular freezing point.

267
Figure 3 | *Synteny and origin of type III antifreeze proteins.* (A) Synteny of AFP III tandem arrays. Full
lengths of scaffolds and contigs are shown. Colored bars represent regions conserved in at least two species.
Regions sharing the same color are homologous. Gray lines connect these homologous regions. Narrow
gray bars depict regions lacking homology in another species. Positions of ancestral, syntenic AFP arrays
are marked by blue circles and translocated arrays by orange circles. Curved lines connect single AFP arrays
split between contigs. All contigs containing syntenic AFP are homologous to *P. gunnelus* Chr 21. *L.*
maculatus contigs containing translocated AFP fall within different regions of Chr 12. *A. minor* and *A.*
lupus contigs with translocated AFP are homologous to Chr 2. (B) A rooted gene tree of *sasa*, *sasb* and
AFP III paralogs is shown. Branches are colored by whether they denote syntenic versus translocated AFPs,
sasa, or *sasb* genes. The scale bar represents nucleotide substitutions per base pair. The node most closely
related to *sasb* branches into gene glades of translocated and syntenic AFPs, prohibiting resolution of
whether ancestral Zoarcoidei AFP evolved in the syntenically conserved region. Tests of whether Zoaroidei
AFP arose in the syntenic region are described under Supplemental Material.
281

2.4. Gene family evolution and convergence of antifreeze protein copy number - Restricted

maximum likelihood ancestral state reconstruction of AFP III copy number demonstrated that at

least 8 expansions and 12 contractions of AFP III occurred across the studied phylogeny (Fig. S6).

To evaluate whether AFP contractions and expansions convergently evolved during independent

invasions of polar or deep seas, the phylogenetically-controlled model of environmental

associations with AFP III copy number was expanded as a structural equation model predicting

both habitat thermal minima and mean pressure as a function of phylogenetic relatedness. This

enabled regression of AFP III copy number against phylogenetically-independent invasions of

colder or higher-pressure environments, which is necessary for quantifying convergence.

Independent invasions of lower thermal minima were significantly associated with increased AFP

III copy number (95% posterior interval = -1.15 – -0.03 AFP copies per °C). Despite having a

stronger association with pressure before testing for convergence, independent invasions of higher

pressure habitats were not associated with decreased AFP III copy number.

While the association between AFP III copy number and habitat pressure among extant

branches was stronger than with thermal minimum, there is little evidence that this association

resulted from convergent evolution between independent invasions of the deep ocean. By contrast,

weaker associations between temperature and AFP III are more likely to have evolved as a result

of convergent gains and losses of AFP III during invasions of Zoarcoidei into colder and warmer

environments, respectively. The pole-to-pole distribution of Zoarcoidei is marked by a likely

origin in the Arctic followed by invasion of the deep ocean, trans-polar migration across deep-

ocean regions, and subsequent reinvasion of shallow waters in the Southern Ocean (Hobbs et al.

2020; Hotaling et al. 2023). The ancestral state reconstruction of thermal minima shows repeated

invasions of sub-zero oceans across Zoarcoidei (Fig. S7). As shown in Figure S9, the

reconstruction of habitat pressure shows one primary lineage of deep ocean fishes belonging to the

eelpout family Zoarcidae. Zoarcidae is the deepest dwelling family of zoarcoid fishes. Thus, the

strong effect of pressure on AFP III copies across Zoarcoidei was more likely attributed to a

singular invasion of the deep sea by Zoarcidae rather than multiple convergent events. Decreases

in habitat pressure in Zoarcidae that corresponded with increased AFP III copy number represent

instances of parallel rather than convergent evolution.

Using maximum likelihood gene tree inference, we confirmed the origin of AFP III

evolution in Zoarcoidei. Our tree of all *sasa*, *sasb*, and AFP III homologs across study species is

consistent with a single origin of AFP III from *sasb (Deng et al. 2010)* as shown in Figure 3B.

Maximum likelihood ancestral state reconstructions were used to predict binary characters of

whether species possessed AFP III in the syntenic locus or in one of the six translocated loci. These

reconstructions demonstrated that ancestral Zoarcoidei AFP III most likely arose in the syntenic

region rather than one of the translocated positions (relative likelihood = 727.89 – 1440.39).

2.5. Role of translocation in AFP III expansions - The ancestral AFP III array falls within

a conserved, syntenic region flanked by genes including synuclein gamma (*sncgb*) and LIM-

binding domain protein (*ldb3b*) as previously shown by Deng et al. 2010. We identified at least

six independent translocations of AFP III out of the ancestral syntenic region (Fig. 3A). All

zoarcoids with AFP III retained this narrow scale synteny such that at least some AFP III copies

were flanked upstream by *sncgb* and downstream by *ldb3b*. Whole-contig alignment revealed

conservation of synteny across larger genomic scales. The ancestral AFP III tandem repeat region

flanked by *sncgb* and *ldb3b* fell in the same homologous region of chromosome 21 across

Zoarcoidei (Fig. 3A). In *L. maculatus*, however, the ancestral AFP III array was positioned in a ~3

Mb chromosomal inversion (Fig. 3A). Translocated AFPs clustered into three gene clades (Fig.

3B).

AFP III copies outside of the ancestral array mapped to two translocated regions in each of

three species, totalling six translocated arrays across all study species. The wolffishes *A. lupus* and

A. minor each contained two translocations on contigs homologous to *P. gunnelus* chromosome 2.

However, the two translocated arrays in *A. lupus* and the two in *A. minor* possessed differing

synteny, falling into four distinctly different regions of chromosome 2 (Fig. 3A). Thus, these arrays

likely represent four discrete translocation events. The two translocated arrays in *L. maculatus*

were distributed across two contigs mapping to different regions of *P. gunnelus* chromosome 12

(Fig. 3A). Translocated copies contributed significantly to AFP III expansions across Zoarcoidei,

making up 13 of 18 AFP copies in *A. lupus*, 2 of 5 copies in *A. minor*, and 38 of 42 copies in *L.*

maculatus (Fig. 1A). Considering the short lengths of contigs in which translocated AFP III were

located *L. maculatus* (Fig. 3A), there is potential for misassembly to have inflated the total number

of translocations, the AFP copy number within translocated arrays, and the total AFP copy number

in this species. Conversely, translocated arrays that border the ends of these contigs may also cause

underreporting of total AFP copy number at these locations (Fig. 3A).

A multivariate phylogenetic model predicting syntenic and translocated AFP III copy

number as a function of thermal minima and pressure revealed that environmental gradients in

AFP copy number were driven by expansions of translocated homologs. After controlling for

phylogeny, habitat thermal minima was associated with syntenic AFP copies at a scaled effect

strength of -0.62 ± 0.72 and translocated AFP copies at -1.36 ± 0.70 (Fig. 4A). Mean habitat

pressure yielded an effect on syntenic copies of -0.88 ± 0.54 and an effect on translocated copies

of -9.84 ± 3.08 (Fig. 4B).

Figure 4 | *Effects of thermal minimum and pressure on copy number of syntenicically-conserved and translocated type III antifreeze protein genes.* Densities represent the posterior probabilities for fixed effects (β) of (A) thermal minimum and (B) mean pressure on antifreeze protein copy in ancestral, syntenically-conserved versus translocated sites.

Molecular repair mechanisms causing variation in copy number such as break-induced

replication can directly facilitate translocation of gene copies to regions sharing microhomology

with break sites (Hastings et al. 2009; Hastings, Ira, and Lupski 2009; Kramara, Osia, and Malkova

2018). It is possible that expansions of AFP III via translocation are simply a consequence of

selection for high copy number. AFP III translocations were not more likely among species with

a large number of AFPs, however. Rather, copy number in translocated arrays correlated with

habitat temperature and pressure. This is suggestive of directional selection acting more strongly

on AFP copy number outside of the syntenic, ancestral AFP III locus. Differences in directional

selection on copies may arise as a consequence of subfunctionalization stemming from

transcriptional changes that occur following translocation into new regulatory landscapes (Y.

Wang, Wang, and Paterson 2012; Harewood et al. 2010). Alternatively, copy number at the

ancestral AFP array may be under stronger stabilizing selection and evolutionary constraint

relative to translocated arrays.

Studies observing environmental effects on gene copy number are abundant (Kondrashov

2012; Lauer and Gresham 2019; Mérot et al. 2020), but the role of translocation in mediating

environmental gradients of copy number is less understood. In populations of salmon, a

chromosomal translocation shows strong genotype-environment association with temperature (K.

B. Watson et al. 2022) while in soybean, a reciprocal chromosomal translocation may have

facilitated climatic adaptation (W. Wang et al. 2021). Furthermore, translocation duplications of

AFGP have been observed in notothenioid icefishes (Rivera-Colón et al. 2023), indicating

potential for translocation to affect environmental gradients of antifreeze gene copy number in

other systems. Our results expand on these studies by demonstrating that not only translocation,

but its association with copy number, can contribute to molecular evolution across environments.

Population genomic studies of AFP III structural variation within Zoarcoidei species are

necessary for further understanding environmental gradients in AFP copy number and the potential

role of translocation in shaping these gradients. Intraspecific AFP copy number variation has been

observed in populations of winter flounder inhabiting different depths and extents of sea ice cover

(Hayes, Davies, and Fletcher 1991). Zoarcoids included in our study inhabit ranges that, in some

cases, span both non-freezing and freezing conditions. Neutral and adaptive variation shaping AFP

III copy number across the Zoarcoidei suborder can be more finely resolved by resequencing of

conspecifics and populations distributed across gradients of temperature, pressure, and freezing

risk. Detection and modeling of adaptive and neutral variation in AFP copy number is necessary

for evaluating (i) additional evidence of whether copy number-environment associations arose via

drift and (ii) whether non-neutral copy number variation is greater among translocated AFP III

arrays.

3. Conclusion

Structural variation observable via long-read sequencing appears to have been critical for

AFP III gene family evolution during the convergent adaptation of Zoarcoidei to both poles. Our

findings highlight the role of multiple environmental stressors (temperature and pressure) in

shaping the genetic architecture of adaptations like antifreeze function. Furthermore, structural

variants such as translocations may act as gatekeepers of adaptation by constraining or facilitating

molecular evolution under the selective pressures of those stressors. Future research should address

whether translocation-associated gene family expansions are driven by mechanisms associated

with transposable element architecture at translocation endpoints or break-induced replication. The

reported, long-read Zoarcoidei reference genomes offer further potential to study the role of

structural variation and chromosomal translocation in convergent environmental adaptation, as

well as the complex evolution of antifreeze proteins.

4. Materials and Methods

4.1. DNA extraction, sequencing, and genome assembly - Lycenchelys platyrhina was collected from the Norwegian Sea, Ægir vent field, Norway (72.3286 °N, 1.5178 °E) on July 28th, 2021. The specimen for *Lycenchelys platyrhina* (originally described as *Lycodes platyrhinus* Jensen, 1902) received a putative identification during collection which required genomic identification and statistical consideration. *L. platyrhina* is a rarely observed species associated with only 11 occurrences in the Global Biodiversity Information Facility. The sequenced specimen was collected inside the range of *L. platyrhina* observations (67.1 - 73.8 °N, 10.3 °W - 8.2 °E). To this end, mitochondrial scaffolds of the *L. platyrhina* assembly were aligned to the NCBI nucleotide database using blastn (Altschul et al. 1990). No sequence of *Lycenchelys platyrhina* was available in the database. The top 10 greatest percent similarity blast hits were with species of the family Lycodinae to which *L. platyrhina* belongs; top 5 = *O. amberensis* (95.88%) *Lycenchelys kolthoffi* (95.57%), *Lycenchelys raridens* (95.05%), *Lycenchelys muraena* (94.60%), and *Lycenchelys sarsii* (94.60%). Skeletal muscle from a single individual was used for DNA isolation using the Circulomics Nanobind, BIG DNA prep, Tissue protocol. Isolated genomic DNA was quantified with a Fragment Analyzer (Agilent). Ampure-PB beads were used to purify and remove short fragments from the sample prior to library prep according to manufacturer protocol. A second size selection for >10 kb fragments was performed on the sample using a BluePippin system (Sage Science Inc., Beverly, MA, USA) with a 10kb cutoff. A library was prepared without shearing using SMRTbell® ExpressTemplate Prep Kit 3.0. Final library was size selected using BluePippin with an 8 kb cutoff. The library was sequenced on one 8M SMRT cells on Sequel IIe instrument using Sequel II Binding kit 2.2 and Sequencing Chemistry v2.0. Loading was performed by adaptive loading, movie time: 30 hours, pre-extension 2 hours.

Sampling of *Anarhichas lupus* was performed in July, 2022, on a captive-reared individual

derived from eggs collected in Conception Bay, NewFoundland, Canada (47.697222 °N, -

52.921944 °N) via diving. Sampling of *Anarhichas minor* was performed in July, 2022, on a

captive-reared individual bred from broodstock collected in St-Lawrence Gulf, Banc Beaugé,

Canada (49.750000 °N, -60.166667 °W). Tissue was ground using dry ice. Cells were gently added

to pre-warmed 0.8% agarose and mixture was placed into the wells of a pre-chilled plug mold.

Following solidification, plugs were dislodged and treated with proteinase K and EDTA at 50°C.

HMW DNA was extracted by dissolving the plugs at 70°C to remove the agarose followed by a

phenol-chloroform extraction. Purification was performed on 15.0 μg of gDNA with AMPure PB

beads and a bead-to-template ratio of 1. The DNA libraries were prepared following the Pacific

Biosciences Preparing HiFi SMRTbell® Libraries using the SMRTbell Express Template Prep Kit

2.0 protocol. 10.0 μg of high molecular weight purified genomic DNA (final volume of 120 μl)

was sheared using the Diagenode Megaruptor 3 instrument (Diagenode Inc., Denville, NJ, USA)

and a two-cycle shear method at speed 33 followed by speed 34. The DNA Damage repair, End

repair and SMRT bell ligation steps were performed as described in the template preparation

protocol with the SMRTbell Template Prep Kit 2.0 reagents (Pacific Biosciences, Menlo

Park, CA, USA). The DNA libraries were size selected on a BluePippin system using a 10 kb

cutoff. The libraries underwent an AMPure PB bead cleanup following the SMRTlink v10

calculator procedure before being sequenced on a PacBio Sequel II instrument using adaptive

loading, Sequel II Sequencing Kit 2.0, SMRT Cell 8M and 30 hours movies with a pre-extension

time of 2h. Sequence data was processed using the SMRT Link Analysis

v. 10.1.0.119588 pipeline.

Bathymaster signatus was collected in Alaska, USA (52.1861 °N, -173.596 °W), by the

Alaska Trawl Survey on July 28th, 2012, via trawl net. DNA was extracted from gill filament via

the Circulomics Nanobind Tissue Kit. HiFi Library prep was done with SMRTbell prep kit 3.0

(version 102-166-600 Apr 2022). As part of the library prep, the genomic DNA was sheared to

25kb on a Megaruptor 2.0. The library was size selected on the Blue Pippin according to PacBio's

recommended settings (10kb-50kb collection). The sample was sequenced on one SMRT cell on

the Sequel II with v5 primer, v2.2 polymerase, 1 hr binding time, a 30 hr movie, 2 hr pre-extension,

and adaptive loading.

Lycodes diapterus was collected in Monterey Bay, California, USA (36.8009 °N, -

121.903867 °W), on October, 20th, 2021. High molecular weight DNA was extracted from

dissected muscle tissue using a Qiagen Genomic-tip DNA extraction kit. DNA was then sheared

to ~20 kbp using a Diagenode Megaruptor 3 and size selected for DNA for fragments greater than

10 kb using a BluePippin system. A PacBio HiFi library was prepared with the SMRTbell Express

Template Prep Kit 2.0. The library was then sequenced on a PacBio Sequel II instrument on an

8M SMRT cell in CCS mode with a 30-hour movie time. DNA extraction, library preparation, and

long-read sequencing of *Ophthalmolycus amberensis* are described in detail by Hotaling et al.

(2023).

Haplotype-phased genomes of all species except for *O. amberensis* were assembled from

long-read libraries using hifiasm v0.19.5-r592 (H. Cheng et al. 2021) set to default parameters.

Assemblies of primary contigs are reported here. Read processing and genome assembly of *O.*

amberensis was achieved using Canu (Koren et al. 2017) v1.8 and is described in detail in

Hotaling, et al. (2023). Assembly completeness was evaluated using BUSCO v5.4.7 (Simão et al.

2015) input with universal, single-copy Actinopterygii orthologs.

4.2. AFP III and sasa/b annotation - Homologs of AFP III were annotated in each genome

assembly included in the study using Exonerate v2.78.4 input with a query sequence of the

translated *Macrozoarces americanus* AFP III (J03924.1) set to the model parameter

'protein2genome' (Slater and Birney 2005). *sasa*, *sasb*, and genes flanking the 5' and 3' ends of

the ancestral syntenic region of Zoarcoidei AFP III (*sncgb* and *ldb3b*) were annotated using

Exonerate (Slater and Birney 2005) set to the same parameters input with protein sequences of *sas*

paralogs from *Lycodichtys dearborni* and *Gasterosteus aculeatus* protein sequences of *sncgb*

(XP_040031911.1) and *ldb3b* (XP_040031907.1). Exonerate BED file annotations of AFP III

were filtered for whether hits contained start and stop codons and complete sequences of exons 1

and 2. Translocated AFP III copies outside of the ancestral syntenic AFP III region were manually

reviewed for evidence of misassembly by aligning translocated AFP III regions to (i) known,

characterized AFP translocation in *A. lupus* (Desjardins et al. 2012) and (ii) additional regions of

translocation within single species if they possessed multiple translocation events to identify

artificial duplications. This ancestral region is defined as the locus of conserved synteny at which

AFP tandem arrays are flanked by the genes *sncgb* and *lbd3b*. Evidence of artificial AFP III

translocations driven by misassembly was not apparent. Pseudogenized copies of AFP III were

identified as regions mapping to AFP III query sequences that were complete but lacked a start

codon, lacked one of the two AFP III exons, exhibited truncations of the majority of an exon, or

conserved 5'/3' untranslated regions that lacked a proximal AFP coding sequence. Publicly-

available long-read genomes used in this study included *Cebidichthys violaceus*

(GCA_023349555.1), *Gasterosteus aculeatus* (GCA_016920845.1), *Leptoclinus maculatus*

(GCA_032191485.1), *Lycodes pacificus* (GCA_028022725.1), *Melanostigma gelatinosum*

(GCA_949748355.1), *Pholis gunnellus* (GCA_910591455.2).

4.3. Environmental data - Depth-informed temperatures and salinities were mined from the

Copernicus Global Ocean Reanalysis (GLORYS) database (Jean-Michel et al. 2021) within ±1°

latitude/longitude of species' coordinate of maximum likelihood of occurrence and the expected

depth range of each species spanning a 10 year period between June 1st 2011 - June 1st 2021.

Coordinates of maximum likelihood of occurrence were extracted from AquaMaps standardized

range predictions (Kaschner, Rius-Barile, and Kesner-Reyes 2013), which are probabilistic

predictions of abundance and occurrence derived from Global Biodiversity Information Facility

occurrence data ("GBIF.org," n.d.). AquaMaps ranges were constrained within coordinates of

known species occurrences. Species minimum and maximum habitat depths were extracted from

FishBase (Froese 1996). Thermal minima were calculated as the minimum temperature

experienced by each species within their representative latitude-longitude-depth range. Pressure

was calculated as a function of sea water depth and latitude using conversions defined by the

UNESCO Joint Panel on Oceanographic Tables and Standards (Fofonoff 1983) and expressed as

a mean.

2.4. Species tree construction and phylogenetic modeling - A maximum likelihood species

tree was constructed from MUSCLE (Edgar 2004) alignments of concatenated protein sequences

of universal 3,603 Actinopterygii single copy orthologs using IQ-Tree2 v1.6.12 parameterized

with a WAG substitution model and fixed rate heterogeneity (Minh et al. 2020; Nguyen et al.

2015). A Bayesian phylogenetically-corrected model was then fitted to predict AFP III copy

number variation as a function of species' habitat temperature and pressure. The model assumed

negative binomial variance for the AFP III count data and uninformative, uniform priors.

Phylogenetic relatedness was fitted as a random effect of the genetic distance matrix derived from

the species tree. The thermal minima and mean pressure of species' habitats were included as fixed

effects, with pressure fitted as a second order polynomial function. An iteration of this model was

fitted that predicted AFP III copy number as a function of both pressure and depth, revealing that

pressure exhibited a significant and equivalent effect after controlling for depth. Model selection

performed using Bayesian marginal likelihoods (Berger and Pericchi 1996) revealed that pressure

was a better predictor of AFP III copy number than depth.

A multivariate iteration of the phylogenetic model was also fitted to predict two separate

outcome variables for copy numbers of (i) AFP III within its ancestral, syntenic region and (ii)

AFP III genes translocated out of the syntenic region. This multivariate model did not permit

residual correlations. A third iteration of the phylogenetic model fit structural equations predicting

variation in AFP III copy number as a function of phylogenetically independent variation in habitat

thermal minimum and pressure. Phylogenetically independent variance in abiotic environments

were estimated by predicting environmental variables as a function of phylogeny, and AFP III

copy number as a function of environment and phylogeny. Thus, any effect of thermal minimum

or pressure on AFP III copy number is quantified after conditioning on phylogenetic variation in

abiotic environment.

Posterior sampling across all models was performed using four MCMC chains and 40,000

iterations including 10,000 warmups. All models were fitted using brms v2.20.4 (Bürkner 2017),

an R interpreter of the Stan Bayesian programming language (Carpenter et al. 2017). The

significance of fixed effects was assessed using a probability of direction test such that effects

described as 'significant' exhibited posterior probabilities with 95% confidence intervals above or

below 0 (Makowski et al. 2019). Because of potential misidentification of *L. platyrhina*, we

evaluated the effect of the identification on model predictions. The leverage (the effect of a single

observation on model predictions) that *L. platyrhina* imposed on parameter estimation in

phylogenetic models of AFP gene family evolution across environments was quantified using a

leave-one-out analysis. Removal of *L. platyrhina* from the univariate model imposed a 6.45%

reduction in the effect of thermal minima on AFP III copy number and 4.24% reduction in the

linear effect of pressure.

4.5. Analyses of AFP III gene family evolution and synteny - FASTA sequences of AFP III

exons 1 and 2 and *sasa/b* exons 1 and 6 were extracted from annotation bed files using bed2fasta

from bedtools v2.31.1 (Quinlan and Hall 2010). The two exons were concatenated per gene and

aligned using MAFFT v7.520 (Katoh et al. 2002). A maximum likelihood gene tree was

constructed from the alignment using IQ-TREE2 v1.6.12 set up fixed rate heterogeneity and

substitution model optimization, which selected a HKY+F+G4 model (Minh et al. 2020; Nguyen

et al. 2015). Restricted maximum likelihood ancestral state reconstructions of AFP III copy

number, habitat thermal minimum, and mean habitat pressure were performed using the 'ace'

function of the R package ape v5.7-1 (Paradis, Claude, and Strimmer 2004; Paradis and Schliep

2019) in order to model contractions and expansions of AFP III across the studied phylogeny.

Ancestral state reconstructions of AFP III positions at the ancestral Zoarcoidei node were

performed using ace(). Individual, discrete reconstructions were performed on 7 binary characters

associated with whether species possessed putatively functional copies of AFP III at the

syntenically-conserved locus or the six translocation loci. The relative likelihoods of the

Zoarcoidei common ancestor possessing AFP III at these regions was measured as a ratio of

maximum likelihoods for AFP III presence at the syntentically-conserved region divided by

maximum likelihood of presence at each of the six translocation loci.

A multiple alignment of scaffolds or contigs containing translocated AFP III copies was

performed using progressiveMAUVE v2.4.0 (Darling, Mau, and Perna 2010), which identified

conservation and synteny of genomic regions conserved in at least two species of Zoarcoidei that

exhibited translocated AFPs. This multiple alignment and the positions of translocated AFP arrays

relative to conserved, syntenic regions are plotted in Fig. 3A in order to visualize conserved

synteny of the ancestral AFP III locus, as well as the locations and number of AFP III

translocations.

Acknowledgements

We acknowledge funding from National Science Foundation awards OPP-1543383 and

OPP-1947040 (supporting TD), and OPP-2312253 (to JLK) as well as funding from the Antarctic

Science Bursary. We acknowledge computing resources from the Center for Institutional Research

Computing at Washington State University, the National Science Foundation Extreme Science

and Engineering Discovery Environment (XSEDE), and University of California, Santa Cruz

Research Computing. We thank the veterinary clinic staff of the Biodôme de Montréal, including

Karine Béland, DMV. We also thank the captain and crew of the ARSV *Laurence M. Gould*, the

personnel of the U.S. Antarctic Program Support Contractors for assistance in Chile, at sea and at

Palmer Station, as well as the logistics in Denver, CO, for their support related to the Antarctic

fieldwork required for this study. We acknowledge funding from the Trond Mohn Science

Foundation through the Centre for Deep Sea Research (grant number TMS2020TMT13) and the

Norwegian Biodiversity Information Centre (the Taxonomy Initiative, project nr. 3-20-70184243)

and thank the scientists and students from University of Bergen (Norway), the crew of the RV GO

Sars and the pilots of the ROV ÆGIR6000 for their assistance with sampling in the Arctic. Lastly,

we would like to thank the Alaskan Trawl Survey, and the Ocean Genome Legacy Center,

Northeastern University, for assisting with specimen collections and archiving.

Author Contributions

Designed research (JLK, SNB)

Performed research (FH, LSFL, IB, JLK, MHE, NRLF, NS, PBF, SH, SNB, TA, TD)

Analyzed data (JLK, NS, SNB)

Wrote the paper (JLK, NS, SNB)

Edited the paper (FH, LSFL, IB, JLK, MHE, NRLF, NS, PBF, SH, SNB, TA, TD)

Data Availability

Genome assemblies, raw sequencing reads, and associated files are available on NCBI under the

following Bioprojects: PRJNA980960 (*Anarhichas lupus*), PRJNA982125 (*Anarhichas minor*),

PRJNA982427 (*Bathymaster signatus*), PRJNA985917 (*Lycenchelys platyrhina*), PRJNA982746

(*Lycodes diapterus*) and PRJNA701078 (*Omphalmolycus amberensis*). All code used for

bioinformatic processing and statistical analysis are available under the GitHub repository

https://github.com/snbogan/Long_AFP [available upon acceptance and publication] in addition to

intermediate data files and specimen metadata. An archived copy of this repository is available on

Zenodo under the DOI [to be generated upon acceptance].

References

Altschul, S. F., W. Gish, W. Miller, E. W. Myers, and D. J. Lipman. 1990. "Basic Local Alignment Search Tool." *Journal of Molecular Biology* 215 (3): 403–10.

Auger, Matthis, Rosemary Morrow, Elodie Kestenare, Jean-Baptiste Sallée, and Rebecca Cowley. 2021. "Southern Ocean in-Situ Temperature Trends over 25 Years Emerge from Interannual Variability." *Nature Communications* 12 (1): 514.

Baalsrud, Helle Tessand, Ole Kristian Tørresen, Monica Hongrø Solbakken, Walter Salzburger, Reinhold Hanel, Kjetill S. Jakobsen, and Sissel Jentoft. 2018. "De Novo Gene Evolution of Antifreeze Glycoproteins in Codfishes Revealed by Whole Genome Sequence Data." *Molecular Biology and Evolution* 35 (3): 593–606.

Barrett, J. 2001. "Thermal Hysteresis Proteins." *The International Journal of Biochemistry & Cell Biology* 33 (2): 105–17.

Baskaran, Abirami, Manigundan Kaari, Gopikrishnan Venugopal, Radhakrishnan Manikkam, Jerrine Joseph, and Parli V. Bhaskar. 2021. "Anti Freeze Proteins (Afp): Properties, Sources and Applications - A Review." *International Journal of Biological Macromolecules* 189 (October):292–305.

Berger, James O., and Luis R. Pericchi. 1996. "The Intrinsic Bayes Factor for Model Selection and Prediction." *Journal of the American Statistical Association* 91 (433): 109–22.

Bista, Iliana, Martin Collins, and Wellcome Sanger Institute Tree of Life Management, Samples and Laboratory team. 2024. "The Genome Sequence of the Limp Eelpout, Melanostigma Gelatinosum Günther, 1881." *Wellcome Open Research* 9:192.

Bista, Iliana, Jonathan M. D. Wood, Thomas Desvignes, Shane A. McCarthy, Michael Matschiner, Zemin Ning, Alan Tracey, et al. 2023. "Genomics of Cold Adaptations in the Antarctic Notothenioid Fish Radiation." *Nature Communications* 14 (1): 3412.

Bodnar, R. J. 1993. "Revised Equation and Table for Determining the Freezing Point Depression of H2O-Nacl Solutions." *Geochimica et Cosmochimica Acta* 57 (3): 683–84.

Bürkner, Paul-Christian. 2017. "Brms: An R Package for Bayesian Multilevel Models Using Stan." *Journal of Statistical Software* 80 (August):1–28.

Carpenter, Bob, Andrew Gelman, Matthew D. Hoffman, Daniel Lee, Ben Goodrich, Michael Betancourt, Marcus A. Brubaker, Jiqiang Guo, Peter Li, and Allen Riddell. 2017. "Stan: A Probabilistic Programming Language." *Journal of Statistical Software* 76 (January). https://doi.org/10.18637/jss.v076.i01.

Cheng, C., and A. Devries. 2002. "Origins and Evolution of Fish Antifreeze Proteins." *Fish Antifreeze Proteins*, October, 83–107.

Cheng, Haoyu, Gregory T. Concepcion, Xiaowen Feng, Haowen Zhang, and Heng Li. 2021. "Haplotype-Resolved de Novo Assembly Using Phased Assembly Graphs with Hifiasm." *Nature Methods* 18 (2): 170–75.

Darling, Aaron E., Bob Mau, and Nicole T. Perna. 2010. "progressiveMauve: Multiple Genome Alignment with Gene Gain, Loss and Rearrangement." *PloS One* 5 (6): e11147.

Deng, Cheng, C-H Christina Cheng, Hua Ye, Ximiao He, and Liangbiao Chen. 2010. "Evolution of an Antifreeze Protein by Neofunctionalization under Escape from Adaptive Conflict." *Proceedings of the National Academy of Sciences of the United States of America* 107 (50): 21593–98.

Desjardins, Mariève, Laurie A. Graham, Peter L. Davies, and Garth L. Fletcher. 2012. "Antifreeze Protein Gene Amplification Facilitated Niche Exploitation and Speciation in Wolffish." *The FEBS Journal* 279 (12): 2215–30.

DeVries, A. L. 1971. "Glycoproteins as Biological Antifreeze Agents in Antarctic Fishes." *Science* 172 (3988): 1152–55.

DeVries, A. L., and D. E. Wohlschlag. 1969. "Freezing Resistance in Some Antarctic Fishes." *Science* 163 (3871): 1073–75.

DeVries, Arthur L., and C. -H Christina Cheng. 2005. "Antifreeze Proteins and Organismal Freezing Avoidance in Polar Fishes." In *Fish Physiology*, 22:155–201. Academic Press.

Duman, John G. 2015. "Animal Ice-Binding (antifreeze) Proteins and Glycolipids: An Overview with Emphasis on Physiological Function." *The Journal of Experimental Biology* 218 (Pt 12): 1846–55.

Edgar, Robert C. 2004. "MUSCLE: Multiple Sequence Alignment with High Accuracy and High Throughput." *Nucleic Acids Research* 32 (5): 1792–97.

Fetterer, F. ;. Knowles K. ;. Meier W. ;. Savoie M. ;. Windnagel A. 2017. "Sea Ice Index, Version 3." NSID. https://doi.org/10.7265/N5K072F8.

Fofonoff, N. P. 1983. "Algorithms for the Computation of Fundamental Properties of Seawater." https://repository.oceanbestpractices.org/handle/11329/109.

Friedman, Sarah T., and Martha M. Muñoz. 2023. "A Latitudinal Gradient of Deep-Sea Invasions for Marine Fishes." *Nature Communications* 14 (1): 773.

Froese, R. 1996. "FishBase." *Oceanographic Literature Review* 3 (43): 321.

Fujino, K., E. L. Lewis, and R. G. Perkin. 1974. "The Freezing Point of Seawater at Pressures up to 100 Bars." *Journal of Geophysical Research* 79 (12): 1792–97.

"GBIF.org." n.d. Global Biodiversity Information Facility. Accessed 2024. https://www.gbif.org.

Gilbert, Jack A., Peter L. Davies, and Johanna Laybourn-Parry. 2005. "A Hyperactive, Ca2+-Dependent Antifreeze Protein in an Antarctic Bacterium." *FEMS Microbiology Letters* 245 (1): 67–72.

Gillett, Matthew B., Jolyn R. Suko, Freddy O. Santoso, and Paul H. Yancey. 1997. "Elevated Levels of Trimethylamine Oxide in Muscles of Deep-Sea Gadiform Teleosts: A High-Pressure Adaptation?" *The Journal of Experimental Zoology* 279 (4): 386–91.

Gupta, Ravi, and Renu Deswal. 2014. "Antifreeze Proteins Enable Plants to Survive in Freezing Conditions." *Journal of Biosciences* 39 (5): 931–44.

Harewood, Louise, Frédéric Schütz, Shelagh Boyle, Paul Perry, Mauro Delorenzi, Wendy A. Bickmore, and Alexandre Reymond. 2010. "The Effect of Translocation-Induced Nuclear Reorganization on Gene Expression." *Genome Research* 20 (5): 554–64.

Hastings, P. J., Grzegorz Ira, and James R. Lupski. 2009. "A Microhomology-Mediated Break-Induced Replication Model for the Origin of Human Copy Number Variation." *PLoS Genetics* 5 (1): e1000327.

Hastings, P. J., James R. Lupski, Susan M. Rosenberg, and Grzegorz Ira. 2009. "Mechanisms of Change in Gene Copy Number." *Nature Reviews. Genetics* 10 (8): 551–64.

Hayes, Pliny H., Peter L. Davies, and Garth L. Fletcher. 1991. "Population Differences in Antifreeze Protein Gene Copy Number and Arrangement in Winter Flounder." *Genome / National Research Council Canada = Genome / Conseil National de Recherches Canada* 34 (1): 174–77.

Hew, C. L., N. C. Wang, S. Joshi, G. L. Fletcher, G. K. Scott, P. H. Hayes, B. Buettner, and P. L. Davies. 1988. "Multiple Genes Provide the Basis for Antifreeze Protein Diversity and Dosage in the Ocean Pout, Macrozoarces Americanus." *The Journal of Biological Chemistry* 263 (24): 12049–55.

Hobbs, Rod S., Jennifer R. Hall, Laurie A. Graham, Peter L. Davies, and Garth L. Fletcher.

2020. "Antifreeze Protein Dispersion in Eelpouts and Related Fishes Reveals Migration and Climate Alteration within the Last 20 Ma." *PloS One* 15 (12): e0243273.

Hotaling, Scott, Marek L. Borowiec, Luana S. F. Lins, Thomas Desvignes, and Joanna L. Kelley. 2021. "The Biogeographic History of Eelpouts and Related Fishes: Linking Phylogeny, Environmental Change, and Patterns of Dispersal in a Globally Distributed Fish Group." *Molecular Phylogenetics and Evolution* 162 (September):107211.

Hotaling, Scott, Thomas Desvignes, John S. Sproul, Luana S. F. Lins, and Joanna L. Kelley. 2023. "Pathways to Polar Adaptation in Fishes Revealed by Long-read Sequencing." *Molecular Ecology* 32:1381–97.

Hull, Ryan M., Cristina Cruz, Carmen V. Jack, and Jonathan Houseley. 2017. "Environmental Change Drives Accelerated Adaptation through Stimulated Copy Number Variation." *PLoS Biology* 15 (6): e2001333.

Jean-Michel, Lellouche, Greiner Eric, Bourdallé-Badie Romain, Garric Gilles, Melet Angélique, Drévillon Marie, Bricaud Clément, et al. 2021. "The Copernicus Global 1/12° Oceanic and Sea Ice GLORYS12 Reanalysis." *Frontiers of Earth Science in China* 9. https://doi.org/10.3389/feart.2021.698876.

Jia, Z., C. I. DeLuca, H. Chao, and P. L. Davies. 1996. "Structural Basis for the Binding of a Globular Antifreeze Protein to Ice." *Nature* 384 (6606): 285–88.

Kaschner, K., J. Rius-Barile, and K. Kesner-Reyes. 2013. "About AquaMaps." aquamaps.org. 2013. https://www.aquamaps.org/AboutAquaMaps.htm.

Katoh, Kazutaka, Kazuharu Misawa, Kei-Ichi Kuma, and Takashi Miyata. 2002. "MAFFT: A Novel Method for Rapid Multiple Sequence Alignment Based on Fast Fourier Transform." *Nucleic Acids Research* 30 (14): 3059–66.

Kelley, Joanna L., Jan E. Aagaard, Michael J. MacCoss, and Willie J. Swanson. 2010. "Functional Diversification and Evolution of Antifreeze Proteins in the Antarctic Fish Lycodichthys Dearborni." *Journal of Molecular Evolution* 71 (2): 111–18.

Kidwell, Margaret G. 2002. "Transposable Elements and the Evolution of Genome Size in Eukaryotes." *Genetica* 115 (1): 49–63.

Kondrashov, Fyodor A. 2012. "Gene Duplication as a Mechanism of Genomic Adaptation to a Changing Environment." *Proceedings. Biological Sciences / The Royal Society* 279 (1749): 5048–57.

Koren, Sergey, Brian P. Walenz, Konstantin Berlin, Jason R. Miller, Nicholas H. Bergman, and Adam M. Phillippy. 2017. "Canu: Scalable and Accurate Long-Read Assembly via Adaptive K-Mer Weighting and Repeat Separation." *Genome Research* 27 (5): 722–36.

Kramara, J., B. Osia, and A. Malkova. 2018. "Break-Induced Replication: The Where, The Why, and The How." *Trends in Genetics: TIG* 34 (7): 518–31.

Lauer, Stephanie, and David Gresham. 2019. "An Evolving View of Copy Number Variants." *Current Genetics* 65 (6): 1287–95.

Makowski, Dominique, Mattan S. Ben-Shachar, S. H. Annabel Chen, and Daniel Lüdecke. 2019. "Indices of Effect Existence and Significance in the Bayesian Framework." *Frontiers in Psychology* 10 (December):2767.

Marx, Vivien. 2023. "Method of the Year: Long-Read Sequencing." *Nature Methods* 20 (1): 6–11.

Mérot, Claire, Rebekah A. Oomen, Anna Tigano, and Maren Wellenreuther. 2020. "A Roadmap for Understanding the Evolutionary Significance of Structural Genomic Variation." *Trends in Ecology & Evolution* 35 (7): 561–72.

Minh, Bui Quang, Heiko A. Schmidt, Olga Chernomor, Dominik Schrempf, Michael D.
Woodhams, Arndt von Haeseler, and Robert Lanfear. 2020. "IQ-TREE 2: New Models and
Efficient Methods for Phylogenetic Inference in the Genomic Era." *Molecular Biology and*
Evolution 37 (5): 1530–34.
Møller, Peter Rask, Jørgen G. Nielsen, and M. Eric Anderson. 2005. "Systematics of Polar
Fishes." In *Fish Physiology*, 22:25–78. Academic Press.
Nguyen, Lam-Tung, Heiko A. Schmidt, Arndt von Haeseler, and Bui Quang Minh. 2015. "IQ-
TREE: A Fast and Effective Stochastic Algorithm for Estimating Maximum-Likelihood
Phylogenies." *Molecular Biology and Evolution* 32 (1): 268–74.
Paradis, Emmanuel, Julien Claude, and Korbinian Strimmer. 2004. "APE: Analyses of
Phylogenetics and Evolution in R Language." *Bioinformatics* 20 (2): 289–90.
Paradis, Emmanuel, and Klaus Schliep. 2019. "Ape 5.0: An Environment for Modern
Phylogenetics and Evolutionary Analyses in R." *Bioinformatics* 35 (3): 526–28.
Potter, R. W., M. A. Clynne, and D. L. Brown. 1978. "Freezing Point Depression of Aqueous
Sodium Chloride Solutions." *Economic Geology and the Bulletin of the Society of*
Economic Geologists 73 (2): 284–85.
Quinlan, Aaron R., and Ira M. Hall. 2010. "BEDTools: A Flexible Suite of Utilities for
Comparing Genomic Features." *Bioinformatics* 26 (6): 841–42.
Raymond, J. A., and A. L. DeVries. 1977. "Adsorption Inhibition as a Mechanism of Freezing
Resistance in Polar Fishes." *Proceedings of the National Academy of Sciences of the United*
States of America 74 (6): 2589–93.
Rivera-Colón, Angel G., Niraj Rayamajhi, Bushra Fazal Minhas, Giovanni Madrigal, Kevin T.
Bilyk, Veronica Yoon, Mathias Hüne, Susan Gregory, C. H. Christina Cheng, and Julian M.
Catchen. 2023. "Genomics of Secondarily Temperate Adaptation in the Only Non-Antarctic
Icefish." *Molecular Biology and Evolution* 40 (3): msad029.
Schmidt, Joshua M., Robert T. Good, Belinda Appleton, Jayne Sherrard, Greta C. Raymant,
Michael R. Bogwitz, Jon Martin, et al. 2010. "Copy Number Variation and Transposable
Elements Feature in Recent, Ongoing Adaptation at the Cyp6g1 Locus." *PLoS Genetics* 6
(6): e1000998.
Simão, Felipe A., Robert M. Waterhouse, Panagiotis Ioannidis, Evgenia V. Kriventseva, and
Evgeny M. Zdobnov. 2015. "BUSCO: Assessing Genome Assembly and Annotation
Completeness with Single-Copy Orthologs." *Bioinformatics* 31 (19): 3210–12.
Slater, Guy St C., and Ewan Birney. 2005. "Automated Generation of Heuristics for Biological
Sequence Comparison." *BMC Bioinformatics* 6 (February):31.
Wang, Weidong, Liyang Chen, Xutong Wang, Jingbo Duan, Rachel D. Flynn, Ying Wang,
Chancelor B. Clark, et al. 2021. "A Transposon-Mediated Reciprocal Translocation
Promotes Environmental Adaptation but Compromises Domesticability of Wild Soybeans."
The New Phytologist 232 (4): 1765–77.
Wang, Yupeng, Xiyin Wang, and Andrew H. Paterson. 2012. "Genome and Gene Duplications
and Gene Expression Divergence: A View from Plants." *Annals of the New York Academy*
of Sciences 1256 (May):1–14.
Watson, K. Beth, Sarah J. Lehnert, Paul Bentzen, Tony Kess, Antony Einfeldt, Steven Duffy,
Ben Perriman, Sigbjørn Lien, Matthew Kent, and Ian R. Bradbury. 2022. "Environmentally
Associated Chromosomal Structural Variation Influences Fine-Scale Population Structure
of Atlantic Salmon (Salmo Salar)." *Molecular Ecology* 31 (4): 1057–75.
Watson, Spencer K., Ronald J. deLeeuw, Doug E. Horsman, Jeremy A. Squire, and Wan L. Lam.

2007. "Cytogenetically Balanced Translocations Are Associated with Focal Copy Number
798   Alterations." *Human Genetics* 120 (6): 795–805.
799 Yancey, Paul H., Wendy R. Blake, and James Conley. 2002. "Unusual Organic Osmolytes in
800   Deep-Sea Animals: Adaptations to Hydrostatic Pressure and Other Perturbants."
801   *Comparative Biochemistry and Physiology. Part A, Molecular & Integrative Physiology*
802   133 (3): 667–76.
803 Yancey, Paul H., Mackenzie E. Gerringer, Jeffrey C. Drazen, Ashley A. Rowden, and Alan
804   Jamieson. 2014. "Marine Fish May Be Biochemically Constrained from Inhabiting the
805   Deepest Ocean Depths." *Proceedings of the National Academy of Sciences of the United*
806   *States of America* 111 (12): 4461–65.
807 Zhuang, Xuan, Chun Yang, Katherine R. Murphy, and C-H Christina Cheng. 2019. "Molecular
808   Mechanism and History of Non-Sense to Sense Evolution of Antifreeze Glycoprotein Gene
809   in Northern Gadids." *Proceedings of the National Academy of Sciences of the United States*
810   *of America* 116 (10): 4400–4405.

A

B

A

P. gunnelus	
O. amberensis	
L. platyrhina	
L. maculatus	
A. minor	
A. lupus	

0   10   20   **Mb**

0   10   20   **Mb**

B

Gene family
- AFP syntenic
- AFP translocated
- *sasa*
- *sasb*

0.1