Adaptation to yeast restriction through the evolution of resource acquisition,

macronutrient reserve, and ovarian function

Purbasha Dasgupta[1,2], purbasha_dasgupta@brown.edu

Anish Koner[1,3], anish.koner@students.iiserpune.ac.in

Rabi Sankar Pal[1], rabisp@iiserbpr.ac.in

Pranav Nandan Pradhan[1], pranav19@iiserbpr.ac.in

Kingkini Roychoudhury[1], kingkini19@iiserbpr.ac.in

Bodhisatta Nandy*[1]

*Corresponding author: nandy@iiserbpr.ac.in, contact number: +91-680-222 7755

Affiliation:

[1]Indian Institute of Science Education and Research Berhampur, Transit Campus, Govt. ITI

Building, NH 59, Engineering School Junction, Ganjam, Berhampur 760 010, Odisha, India.

Present addresses:

[2]Brown University, 225 Dyer Street, Providence, RI 02903

[3]Indian Institute of Science Education and Research Pune, Dr. Homi Bhabha Road, Pashan,

Maharashtra, Pune 411008, India.

Abstract

Optimization of reproduction under dietary adversity is an important aspect of diet-dependent

adaptation. Yet, little is known about the mechanism of such adaptive evolution. Here, we

investigated a set of experimentally evolved populations of *Drosophila melanogaster* where

early-life fecundity evolved as an adaptation to chronic protein restriction. We tested the role of

resource acquisition and macronutrient storage, and changes in ovarian function that could have

allowed such reproductive adaptation. We show that higher early-life fecundity was associated

with the increased larval feeding rate, aiding in accumulation of higher protein content at the

time of eclosion. Further evidence also suggests increase in reproductively allocated lipid

content. Evolved females were found to have larger ovaries that also had a higher number of

mature, post-vitellogenic oocytes that seem to readily allow the attainment of higher fecundity.

Our results further show the evolution of plasticity in ovariole count (depending on mating

status) and total egg-chamber count in females. These results are important in understanding the

putative role of genetic variance and covariances that aid or limit the evolution of reproductive

optimization, especially under nutritional adversity.

Key words: Dietary restriction, ovariole count, oocyte count, lipid content, protein content

Introduction

Natural selection should lead to the optimization of reproductive investment since resource availability is never unlimited (Kirkwood, 1977; Roff, 1992; Stearns, 1989, 1992, 2000). Allocation in reproductive traits often results in reduced survival and/or reduced future fertility due to constrained resource availability to growth, immune function, and tissue repair (Stearns, 1992). Reproductive optimization is thus pivotal to successful life history strategies. Pianka (1976) conceptualized organisms as "input-output systems" where foraging tactics represented the input of resources, and progeny number represented the output. Thus, it is important to integrate foraging and resource allocation with life history to better understand adaptive reproductive optimization (Pianka, 1976; Boggs 1992; Gatto, Matessi & Slobodkin, 1989). Putatively important role of dietary ecology in reproductive physiology and strategy have not been fully explored beyond the literature on dietary restriction leading to temporary shutting down of reproduction (Mair et al., 2004; Carey et al., 2008; Inness & Metcalfe, 2008; Adler & Bonduriansky, 2014; Moatt et al., 2016; Zajitschek et al., 2016; Zajitschek et al., 2019). Since metabolic energy serves as the currency that links foraging to optimal reproduction, investment is often physiologically regulated (Boggs, 1992; Ellison, 2003; Jervis et al., 2008). Further, the quantity and timing of reproductive allocation have profound implications for lifetime reproductive success (Long et al., 2010; Desprez et al., 2017; Churchill et al., 2019). However, genetic variance and covariances that aid or limit the evolution of reproductive allocation and its timing are poorly understood.

Dietary restriction (DR), a well-known intervention, can be a key experimental paradigm to probe resource allocation and reallocation. DR is known to limit reproductive allocation, often

resulting in an increased investment in stress resistance and survival (Mair et al., 2005; Partridge

et al., 2005; Nakagawa et al., 2012; Fontana & Partridge, 2015). Theoretically, such resource

reallocation may enable organisms overcome transient (nutritionally) challenging periods by

prioritizing essential functions for immediate survival (Shanley & Kirkwood, 2000; Adler &

Bonduriansky, 2014; Zajitschek et al., 2019; Moatt et al., 2020). Interestingly, long-term (multi-

generational) nutritional restriction can result in the evolution of strategies aimed at optimizing

reproduction (Williams, 1957; Bonduriansky et al., 2008; Zajitschek et al., 2019). Such

evolutionary responses can be used to infer on the genetic variance and covariances pertaining to

resource acquisition, utilization and storage traits underlying adaptive reproductive optimization

(Chippindale et al., 1996; Schwasinger-Schmidt et al., 2012; Ahmad et al., 2018).

Experimental evolution under a variety of dietary conditions has been known to result in

adaptations resulting in improved survival and/or reproduction in fruit flies (Kolss et al., 2009;

Zajitschek et al., 2016; Cavigliasso et al., 2020). Similar to several other organisms, phenotypic

response to short-term nutritional deprivation in *Drosophila* involves a shift from reproduction to

survival mode (Rion & Kawecki, 2007; Ballard, Melvin & Simpson, 2008; Lee & Jang, 2014;

Shu et al., 2022). For instance, protein restriction caused by yeast limitation has been shown to

increase longevity and starvation resistance at the cost of fecundity (Chippindale, 1993;

Zimmerman et al., 2003; Mair et al., 2005; Piper et al., 2005), indicating physiological trade-off

between egg production and somatic storage, especially lipid reserve (Simmons & Bradley,

1997). Further, in response to a low protein: carbohydrate diet, adult *Drosophila melanogaster*

showed an increase in lipid reserve accompanied by greater resistance to starvation stress (Lee &

Jang, 2014). Importantly, in laboratory and wild populations, *Drosophila* females largely depend

on yeast as the primary protein source, which has a profound impact on reproductive output.

Restricted access to yeast, especially live-yeast supplement, appears to have a more targeted

effect on decreasing female fecundity, and extending lifespan (Chippindale et al., 1993; Nandy et

al., 2012; Magwere et al., 2004). However, such reallocation of resources from reproduction to

survival, is unlikely to be adaptive under long-term nutritional restriction. Generations under

yeast-deprivation have been shown to lead to the evolution of higher early-life fecundity in

experimental populations of *D. melanogaster* (Dasgupta et al., 2022). Interestingly, such

reproductive evolution seems to be a readjustment of reproductive schedule, wherein increased

early-life fecundity evolved with faster reproductive senescence such that cumulative life-time

fecundity and lifespan were unaffected (Dasgupta et al., 2022). Thus, it is amply clear that

selection can act on variation affecting early-life fecundity, driving such reproductive adaptation.

However, it is not yet clear whether the underlying physiological mechanism of such adaptation

involves evolution of resource acquisition (larval or adult) or macronutrient metabolism (for

example, storage) or both.

Here, we addressed this unsolved problem by investigating the populations used by Dasgupta et

al. (2022). Theoretically, there are two non-mutually exclusive ways of achieving higher early-

life reproductive output in females, without an associated survival cost - (a) increased larval

resource acquisition (juvenile resource acquisition theory), and (b) increased macronutrient

reserve that can cater to the increased metabolic energy demand of higher reproduction

(macronutrient content theory). Once the metabolic resources are made available to the female

reproductive system, the system can either ramp up the production of eggs and/or increase the

number of egg-producing units. Since in holometabolous insects such as fruit flies, early-life

fitness heavily depends on larval resource acquisition, we compared early third-instar larval

feeding rates across evolved and control regimes to test juvenile resource acquisition theory.

Considering the role of protein and lipid in egg production (Drummond-Barbosa & Spradling,

2001) as well as in survival and stress resistance (Chippindale, Chu & Rose, 1996; Rehman &

Varghese, 2021), we compared the whole-body protein content and lipid content in females of

evolved vs. control lines to test the macronutrient content theory. Further, we also investigated

the putative role of changes in ovarian function by measuring ovarian size (ovariole count),

which is known to be a strong determinant of reproductive potential (Wayne et al., 2006; Lobell

et al., 2017), and the abundance of mature stages of oocytes in each ovariole. We hypothesized

that the increased early-life fecundity in evolved females should be ascribed to either an increase

in ovariole count and/or a surge in the production of the number of oocytes per ovariole.

Materials and methods

<u>Study populations and selection regime</u>

All the experiments described here were conducted using the four replicate experimental

populations (YLB_{1-4}) and their corresponding control populations (BL_{1-4}). The details of these

populations are discussed in Dasgupta et al. 2022. In short, experimental populations (YLB_{1-4})

were derived from four replicate baseline populations (BL_{1-4}) of *D. melanogaster*, which are

outbred (N ~2800), wild-type populations. BL populations were maintained under a 14-day

discrete generation cycle at 25°C on a banana-barley-sugarcane jaggery-yeast diet. Adults are

given a supplement of *ad-lib* yeast-paste smeared on food plates. Every generation, eggs are

collected during an 18-hour window of the day14 of the generation cycle to start the next

generation. This 18-hour window constitutes the (Darwinian) fitness window as in this system,

progeny produced only during this window can be passed on to the next generation. Hereafter,

"fitness window" refers to this time period of the maintenance regime.

To create YLB populations, ∼2800 eggs were collected from each BL population, and these

populations were maintained identically to BL populations, except the yeast supplementation

step for adults. Thus, YLB populations are maintained in absence of yeast-supplement at the

adult. Prior to assays, all populations were passed through one generation of common-garden

rearing conditions to equalize the putative role of parental diet. During this common-garden

rearing phase, both BL and YLB populations were provided with *ad-lib* yeast-paste, matching

the ecology of BL regime exactly. Wherever needed virgin flies were collected within 6 hours of

eclosion under light CO_2-anaesthesia. A total of six experiments were performed. Experiments 1-

3 were conducted within 67-74 generations; experiment 4 on the 79[th] generation; experiment 5

within the 94-95 generations and finally, experiment 6 on the 120[th] generation of the

experimental evolution.

Experiment 1: Whole-body protein content in females

The general plan was to measure protein content at two time points – on eclosion, and during the

fitness window (i.e., day14 of the generation cycle) in reproducing condition. In this assay, one

set of flies was used to measure whole body protein content using the Bradford method (see

Supplementary information S1). The other set was used to measure whole body mass, which is

used as a scaling factor at the time of statistical analysis.

For protein quantification on eclosion, females were collected as virgins in two sets and

immediately frozen at -20□. In one set, groups of three females were collected in an empty vial,

which were processed for protein content measurement within 24 hours. A second set of females

were collected in groups of five, which were processed for body weight measurement within 24

hours. Twelve such vials were measured per population. For body weight measurement, flies

were dried in a hot air oven for 24 hours at 30□ (following Colinet & Renault, 2014) and

weighed using Shimadzu AUW220D to the nearest 0.01 mg in the pre-formed groups. Each

group of five females was weighed thrice and the average of the three readings was noted. A

population mean was calculated using values from all replicate vials. These population means

were used to normalize the protein content value obtained from the Bradford assay (see

Supplementary information S1). For protein quantification in the fitness window, adult flies were

collected as virgins and held in single-sex food vials in groups of ten. 10 such vials were set up

for each sex per population. On day12 post-egg collection, 2-3 days old virgin males and females

were transferred into fresh food vials and were allowed to interact for 48 hours during which

they completed mating. Afterwards, the vials containing flies were frozen at -20□. Frozen

females were sorted in groups of three in empty glass vials. These females were later used for

protein quantification using the Bradford assay (see Supplementary information S1). Here, each

group of three females served as the unit of replication. Additionally, 12 vials with five females

in each were kept for dry weight measurement. The latter was done within 24 hours of freezing,

following the same protocol described above.

Experiment 2: Whole-body dry weight and lipid content in females

The general plan was to measure whole body dry weight and lipid content in freshly eclosed

females and in reproducing females at the fitness window. For the first time point, i.e., on

eclosion, virgin females were collected in groups of 10 and held in a glass vial. 12 such vials

were collected for a population and were immediately frozen at -20□. Within 24 hours of

freezing, all the flies were transferred to a hot air oven and allowed to dry for 48 hours at 60□

before being processed and weighed.

For the second time point, adult flies were collected as virgins and allowed to pass through the

following steps before being subjected to the body weight and lipid content measurement.

Freshly eclosed flies were sexed and held in single-sex vials in groups of 10 until day12. For a

population, 12 such vials were set up per sex. On day12 post egg collection, males and females

were transferred into fresh food vials, and were allowed to interact for 48 hours, following which

males were discarded and females were frozen at -20□. The frozen dead flies were sorted into

groups of 10 females and 12 such vials were set. Within 24 hours of freezing, all the flies were

transferred to a hot air oven and allowed to dry for 48 hours at 60□ before being processed and

weighed.

Dried females were weighed in groups of 10 using Shimadzu AUW220D to the nearest 0.01 mg.

Each set of 10 females is weighed thrice and the average across these three measurements were

used as Dry Mass (DM) value. After that, each set of 10 individuals was transferred to 1.5 ml

centrifuge tubes with 1.5ml Diethyl ether (Marron et al. 2003) and incubated in an incubator

shaker at 200 RPM for 24 hours at room temperature (~25-30□). After 24 hours, the centrifuge

tubes were kept in a hot air oven, with the lids open, at 60□ for another 24 hours to allow the

Diethyl ether to evaporate. Following this, each set of 10 individuals was weighed thrice again

and the average across these three measurements was obtained. By subtracting this value from

the DM value previously obtained, an Absolute Lipid Mass (ALM) value was obtained for each

set of ten females. Thereafter, relative lipid content per unit dry mass was calculated by using the

formula mentioned below:

$$Relative\ lipid\ content = \frac{ALM}{DM} \times 100$$

This was done for all twelve samples from a population. This lipid extraction method using

Diethyl ether integrates all lipid classes and does not distinguish between lipid sequestered in

eggs and that found in somatic tissues (Ballard et al. 2008). From one round of lipid estimation

experiments, three sets of data were obtained - dry mass, absolute lipid mass and relative lipid

content (i.e., lipid content per unit dry mass). For the final analysis, after the exclusion of

outliers, based on 1.5 IQR assessment, values from 10-12 samples were considered for each

population.

<u>Experiment 3: Starvation resistance</u>

Starvation resistance was assayed by quantifying longevity of the females of both regimes under

starved condition. Like other assays mentioned above, the populations were passed through one

generation of common rearing following which eggs were collected under standard density. On

eclosion, flies were collected and held in single-sex vials in groups of 12. On day 3 post-

eclosion, the 'interaction vials' were set up by combining 12 males and 12 females in a food vial.

12 such vials were set-up for each population. The vials were left undisturbed for 2 days,

following which the females from each vial were separated under light CO_2 anaesthesia and

transferred to 'starvation vials'. These starvation vials were standard 8-dram vials provisioned

with non-nutritive agar gel (1% Agar-agar solution + 1% p-hydroxy benzoic acid solution). The

Agar-gel provided moisture for the flies and ensured that flies are only starved but not

desiccated. Mortality was recorded in each starvation vial at 6-hours intervals. The mean

longevity of 12 individuals per vial under starved condition provided the measure of starvation

resistance and was used as the unit of analysis. We removed a few vials from the analysis due to

a variety of handling errors, including escaped flies and broken vials, during the assay. The final

number of vials used for the analysis ranged from 11 to 12 vials for a population.

Experiment 4: Ovariole count

After one generation of common garden rearing, eggs were collected from commonly reared

cages and cultured following the standard conditions. On eclosion, adult flies were collected as

virgins every 6 hours. Virgin males and females were collected in groups of 10 and held in

single-sex vials. These vials were left undisturbed for two days. On day 3 post eclosion, flies

were transferred into fresh food vials, and these vials were randomly divided into two (mating

status) sets – (a) Mated: 10 males and 10 females placed in a vial and (b) Virgin:10 virgin

females placed in a vial. These vials were left undisturbed for two days.

After two days of conditioning, the flies were frozen at -20□. The number of ovarioles were

counted in both the ovaries. This was done on 15 randomly picked females per population for

both virgin and mated. Ovaries were dissected out on a drop of PBS and stained with a saturated

solution of potassium dichromate for 10-20 minutes, which facilitated the easy identification and

counting of ovarioles (Coyne et al., 1991). The ovariole number of each female was defined as

the mean of the number of ovarioles in the right and left ovaries. The mean ovariole count of

each female was used as the unit of analysis.

Before dissection, thorax length of the females was measured as a proxy for body size to look for

possible correlations between body size and ovariole count. Lateral view images of the thorax

were captured using *ZEISS AxiocamERc 5s* mounted on a stereozoom microscope. Thorax length

was measured as the distance between the highest point of the anterior tip of the thorax and the

posterior-most point of the scutellum. The image analysis was done on *Zeiss ZEN 2 Blue edition*

software. Each individual was measured thrice, and the mean was used for the analysis.

Experiment 5: Larval feeding rate

After common garden rearing, two small food plates were introduced in each population cage

and the females were allowed 1 hour for oviposition to ensure developmental synchrony of the

experimental larvae. Following the protocol by Joshi and Mueller (1996), about a hundred eggs

were then randomly picked from the two oviposition plates and gently transferred to two Petri

dishes containing a thin layer of solidified agar gel (1% agar-agar solution) using a fine brush. 24

hours later, 25 newly hatched larvae were gently transferred to a Petri dish containing a thin

layer of 1% non-nutritive agar overlaid with 1.5 ml of 37.5% yeast suspension. Four such Petri

dishes were set up for each population. They were left undisturbed till the larvae reached an early

third-instar stage. 25 larvae from each population were used to quantify the feeding rate by

placing them individually on Petri dishes containing a thin layer of agar overlaid with 10% yeast

suspension. Each larva was allowed to acclimatize for 15 seconds following which the feeding

rate was quantified by counting the number of cephalo-pharyngeal sclerite retractions per minute

under a stereo zoom microscope (*ZEISS Stemi 508 Stereo Microscope*). Blind-fold observation

was adopted to avoid observer bias. Feeding rate of each larva was taken as the unit of analysis.

Here and in all subsequent analyses, a few outliers were removed from the final analysis using

'±1.5×Interquartile range' rule. Hence, the final number of larvae used for the analysis ranged

between 24-25 for each population.

Experiment 6: Stage-composition of the ovarioles

In *D. melanogaster*, each ovariole comprises a string of progressively developing egg chambers

or follicles. The anterior end of an ovariole is the germarium containing somatic and germline

stem cells. Egg chambers bud off from the germaria and mature through 14 different stages as

they move towards the posterior end near the oviduct. Stage 1-7 are in pre-vitellogenic, stage 8-

10 are in vitellogenic and stage 11-14 are in post-vitellogenic stages of development

(Drummond-Barbosa et al. 2001). Protocols for generation of flies and dissection were identical

to those followed in Experiment 4. During the assay, the ovarioles were carefully teased apart

using stainless steel needles. Stage-specific quantification of the egg chambers was performed on

either the left or right ovary, randomly selected from 15 randomly chosen females per population

for both virgin and mated groups. The ovaries were dissected in 1X PBS and fixed in 4%

paraformaldehyde for 10-30 minutes. Following fixation, the ovaries were washed four times in

1X PBS, with each wash lasting 15 minutes. To increase tissue permeability, 0.1-0.2% Triton-X

was added for 10 minutes. The tissues were then washed in PBS for 15 minutes. Subsequently,

the ovaries were stained with 1 µg/ml DAPI for 5-10 minutes and washed again in PBS for 5-10

minutes. Next, the ovarioles were carefully teased apart using a fine tungsten needle and

mounted with Vectashield antifade mounting media. The total number of egg chambers at each

stage was counted for every female using a fluorescence microscope (*Zeiss Axio Scope A1*). The

proportion of pre-vitellogenic, vitellogenic, and post-vitellogenic egg chambers was calculated

for each female and used as the unit of analysis. The different developmental stages of the egg

chambers were identified based on the characteristics described in Jia et al. (2016).

<u>Data analysis</u>

All analyses were done on R version 4.0.3 (R Core Team 2020). Linear and generalized linear

mixed-effects models (LMM and GLMM respectively) were performed using the package `lme4`

(Bates et al. 2014). Further, to obtain p-values based on the Satterthwaite approximation for

denominator degrees of freedom, we used the R package `lmerTest` (Kuznetsova et al. 2017).

Selection regime was modeled as a fixed factor. Block and its interaction with other fixed-effects

were fitted as random effects.

Larval feeding rate was analysed as a GLMM with a Poisson distribution. An OLRE

(Observation-level random effect) was included in the model to correct for overdispersion.

Female whole-body protein content, absolute lipid mass, relative lipid content and longevity

under starvation were analysed using LMM. The macronutrient contents (protein and lipid) were

quantified at two different adult life-stages/time points - on eclosion and in fitness window.

Initially, time point was fitted as a fixed effect, however, a significant time point $\times$ block

interaction was observed. Hence, separate analyses were performed to assess the effect of

selection regime on the macronutrient content at each of these time points. On the other hand,

female dry body mass was analysed using LMM with selection regime, time point and their

interaction as fixed factors.

Ovariole count was analysed using an LMM. Selection regime, mating status and their

interaction were fitted as fixed effects, whereas thorax length was included as a covariate. The

effect of selection on thorax length was analysed using a GLMM with a gamma distribution and

a log link function. To analyse the proportion of pre-vitellogenic, vitellogenic, and post-

vitellogenic egg chambers per ovary per female, separate models were used. The proportion of

pre-vitellogenic egg chambers was analyzed using a LMM, whereas the proportions of

vitellogenic and post-vitellogenic egg chambers were analyzed using GLMMs with a Poisson

distribution and OLRE to account for overdispersion. In these models, the number of egg

chambers from each stage was the dependent variable, scaled by the total number of egg

chambers across all stages, included as an offset.

Wherever applicable, post hoc analyses were done using Tukey's HSD with `emmeans` package

(Lenth et al. 2018). The final models used for the analyses are described in Table S5. Descriptive

statistics of all the traits quantified here are mentioned in Table S4.

Results

<u>Experiment 1:</u>

Analysis of the whole-body protein content results suggested a significant effect of the selection

regime in freshly eclosed females. Compared to BL females, YLB females eclosed with about

15% higher protein content per unit dry mass (Figure 1a, Table 1, Table S4). At fitness window,

the effect of the selection regime on protein content was not significant (Figure 1b, Table 1).

<u>Experiment 2:</u>

Analysis of the whole-body dry weight suggested a significant effect of selection regime, time

point and the interaction between them (Table 1). On an average, dry mass of the YLB females

was significantly higher compared to that of the BL females. Pairwise comparisons suggested

that, on eclosion the dry weight difference was not significant, however, at the fitness window,

dry weight of the YLB females were found to be significantly higher compared to that of the BL

females (Table S1). While females of both the regimes showed a significant increase in dry mass

from eclosion to their fitness window, the degree to which it increased appeared to be different.

YLB females showed a 16% increase in dry weight whereas, a 12% increase was observed for

the BL females between these two time points (Figure 1c,d; Table 1, Table S4).

Analysis of the lipid content measurement indicated that the effect of the selection regime was

not significant at the first time point, i.e., on-eclosion (Table 1, Figure 1e). This was true for both

measures of lipid content, i.e., absolute lipid content as well as relative lipid content, at this time

point. At the second point, i.e., during the fitness window, the effect of the selection regime on

absolute lipid content was significant but the same was not true for relative lipid content (Table

1). YLB females were found to have ~16% higher mean absolute lipid mass compared to the BL

females at the second time point (Figure 1f, Table S4). However, this difference turns out to be

non-significant after factoring in the dry mass difference. This implied that the observed

difference in dry mass between the two types of females at the second time point was primarily

due to the difference in lipid content.

Experiment 3:

We did not find a significant effect of the selection regime on females' mean longevity under

starved conditions (Table 1, Figure 2b). This indicated that there was no difference in starvation

resistance between the YLB and BL females.

Experiment 4:

Analysis of the ovariole count results indicated that the fixed effect of selection was not

significant (Table 1). However, the effects of mating status, and selection regime × mating status

interaction were significant (Table 1). Tukey's adjusted P-value obtained by pairwise

comparisons suggested that while there was no significant difference in ovariole numbers

between BL and YLB females under virgin condition. However, under mated condition, a

significantly higher (~7%) number of ovarioles was observed in YLB females (Table S2, Figure

3). The effect of thorax length, as a covariate, was not significant (Table 1). We also tested the

effect of selection regime on thorax length and found the effect to be non-significant, indicating

that body size did not respond to experimental evolution (Table 2).

Experiment 5:

The analysis of the feeding rate of the early third-instar larvae suggested a significant effect of

the selection regime (Table 2). The third instar larvae of YLBs were found to have a significantly

higher feeding rate compared to those of the BLs (Figure 2a).

Experiment 6:

The analysis of the total number of oocytes per unit ovariole, regardless of developmental stage,

revealed significant effects of selection regime and selection regime × mating status interaction

(Table 1). BL females had significantly more total egg chambers per ovariole compared to YLB

females (Table S3). Pairwise comparisons with Tukey's adjustment indicated that this difference

was particularly pronounced in mated conditions (9.01% more in mated BL females, Table S4).

The proportion of pre-vitellogenic egg chambers showed no significant effect from selection

regime, mating status, or their interaction (Table 1, Figure 4c). For vitellogenic egg chambers,

the main effects of selection regime and the effect of selection regime × mating status interaction

were not significant (Table 2). However, mating status had a significant effect (Table 2, Figure

4c), wherein regardless of selection regime, mated females were found to have 55.02 % more

vitellogenic egg chambers than virgin females (Table S4, Figure 4c). Analysis of the proportion

of post-vitellogenic egg chambers indicated significant effects of both selection regime and

mating status (Table 2, Figure 4c). YLB females were found to possess 19.87% more post-

vitellogenic stages than that in the BL females (Table S4). Virgin females were found to have a

higher number of mature oocytes regardless of selection regime (Table S4). The effect of

interaction between selection regime and mating status on the proportion of post-vitellogenic egg

chambers was marginally non-significant (Table 2).

Discussion

Reproduction is a protein-intensive process and hence, most animals need persistent access to

proteins to reproduce (Skorupa et al., 2008). We found that as a result of adaptation to chronic

protein deprivation experimental populations (YLB) evolved significantly higher larval feeding

rate, and an associated increase in whole-body weight and protein content at the time of eclosion

(emergence of an adult from puparium). Thus, the results indicate the possibility of evolution of

nutrient specific resource acquisition and/or assimilation at the larval stage. Interestingly, YLB

females were also found to have a higher body mass and absolute lipid content in the fitness

window – indicating the evolution of adult stage resource acquisition and/or differential

utilization and storage as well. Our results further suggest changes in ovarian size (ovariole

count) and function (production of mature oocytes) aiding the attainment of a higher rate of egg

production. The YLB females maintained a higher number of ovarioles and a greater number of

mature (post-vitellogenic) oocytes per ovariole. These results shed light on the proximate

mechanisms underlying the attainment of a higher early-life fecundity, and indicate the putative

role of genetic variance and covariance between juvenile and adult fitness traits pertinent to such

evolution.

<u>Resource acquisition and storage:</u>

In holometabolous insects, early-life reproduction is reliant on larval resource acquisition and

storage (Telang & Wells, 2004; Aguila et al., 2013). In *D. melanogaster*, ovarian development

begins during metamorphosis (Bodenstein, 1950) and hence, early-life ovarian output could rely

primarily on larval nutrition. Unsurprisingly, YLBs were found to have an increased early-third

instar feeding rate, the most prolific feeding stage (Li et al., 2008; Wang et al., 2023). Evolution

of feeding rate under a variety of life history selection is fairly well-known in the literature (Joshi

& Mueller, 1988; Rajamani et al., 2006; Mueller & Barter, 2015; Bitner et al., 2021). However,

such change in feeding behaviour is expected to result in an increase in overall body mass at

eclosion (Bierbaum et al., 1989; Klepsatel et al., 2018). This trend was not observed in our YLB

females. Instead, only (whole-body) protein content seemed to have increased, at least at the time

of eclosion. To the best of our knowledge, this is the first report of a macronutrient-nutrient

specific effect of increased feeding rate. Nonetheless, since protein is known to be a limiting

macronutrient for egg production, such an observation is consistent with the *macronutrient*

content theory mentioned earlier. The observed increase in protein content on eclosion is

expected to have allowed the attainment of elevated early-life fecundity, especially under a

protein restricted ecology, assuming that at least part of this protein reserve is eventually

allocated to egg production. Curiously, increased whole body protein content was not observed

at the fitness window (i.e., 4-5 days post-eclosion). Given that Bradford assay is less sensitive to

conjugated proteins in the body (Giraudi et al., 1997), the observed results could indicate

reproductive allocation of stored protein, as much of the protein molecules in the developing

oocyte, especially yolk protein such as, vitellogenin, is usually found in conjugated forms such

as glycolipid and phosphoprotein (Raikhel & Dhadialla, 1992). Thus, the observed results on

protein content could simply indicate reproductive allocation of stored protein.

Apart from juvenile phase resource acquisition, adult foraging can significantly contribute to

biomass accumulation that can boost reproductive output (Partridge et al., 1999). Our results on

whole body dry weight corroborated this theory and showed that body mass in females increased

from eclosion to the fitness window, i.e., during 4-5 days of adult life. Interestingly, the evolved

YLB females showed a greater increase compared to the control females, indicating that the

evolved females were putting on more biomass during the adult phase until the reproductive

(fitness) window. Such biomass accumulation can either be specific to the reproductive system,

i.e., in reproductive structures – especially increased ovary size (Simmons & Bradley, 1997), or

be more generic long-term storage, especially lipid. The latter would naturally make individuals

more resistant to starvation (Chippindale, Chu & Rose, 1996; Rion & Kawecki, 2007; Rehman &

Varghese, 2021), while the former would allow females to be more fecund.

<u>Allocation of acquired resource to ovaries:</u>

Two lines of evidence uphold the theory of reproductive biomass accumulation. First, YLB

females were found to have significantly higher total body lipid content at the time of their

reproductive window, without a correspondingly higher resistance to starvation. This implied

that stored lipid is not available to general somatic metabolism, and perhaps was locked in

reproductive tissues. Importantly, *Drosophila* ovaries are known to contain ~15% of the whole-

body lipid content (Lease & Wolf, 2011; Colinet & Renault, 2014). It is important to note that

the observed increase in lipid content among the evolved females corresponds with their increase

in whole-body biomass. This suggests that changes in resource acquisition and/or utilization

during the adult phase, which are reflected in biomass accumulation, primarily influenced lipid

storage. Secondly, we found ovariole count, a proxy for ovary size (Mendes & Mirth, 2016;

Sarikaya et al., 2019), to be higher in YLB females. Since a substantial part of ovarian biomass is

known to be lipid, it is not unreasonable to suggest that the increase in lipid content discussed

earlier corresponds to increase in ovary size, at least partly. Thus, our results appear to suggest

an evolution of larval and adult macronutrient specific resource assimilation, but a preferential

allocation of it to the reproductive system. The evolution of such preferential allocation of

metabolic resources is only beginning to emerge (also see Vijendravarma, 2018). It is important

to understand how this can translate into changes in egg production.

<u>Ovarian function:</u>

Notwithstanding the limited information in the literature (Hodin, 2009; Sarikaya et al., 2012; Mendes & Mirth, 2016), our results suggest plasticity in ovariole count in females, such that the ovariole count in mated females was significantly less than that in the virgin females (but see Carlson et al., 1998). Putatively, programmed cell death, which is commonly observed in insect ovaries, including *Drosophila* sp., can potentially modulate ovariole number (Pritchett et al., 2009; Almeida & Suesdek, 2017; Ronai et al., 2016, 2017). In adult fruit flies, protein restriction leads to increased cell death in the germaria and the mid-stages of oogenesis (Drummond-Barbosa et al., 2001; McCall, 2004). In face of nutritional stress, cell deaths are induced in these regions which act as checkpoints determining whether egg chambers should proceed to the vitellogenesis process (Drummond-Barbosa et al., 2001). Mating is known to promote increased oogenesis (Soller et al., 1997; Avila et al., 2011; McDonough-Goldstein et al., 2021). It is not unreasonable to suggest that rapid increase in oogenesis quickly becomes unsustainable, especially under sub-optimal diets such as, protein restriction similar to our assay condition - triggering programmed cell death of developing egg chambers and/or ovarioles. This could indeed act as an effective checkpoint, and regulation on reproductive investment (Moore & Attisano, 2011; Adler & Bonduriansky, 2014).

Interestingly, we did not detect this difference in ovariole count across the two mating statuses in evolved (YLB) females. This could have been achieved either by evolving a higher threshold for the apoptotic control to set in, and/or by making more metabolic resources (e.g., protein, lipid

etc.) available to the reproductive tissues. While the latter seems likely, the former cannot be

ruled out at this point.

Interestingly, our results on oocyte developmental stages suggested that although the ovaries of

the females of the control population had significantly more total egg chambers, the evolved

females produced a higher number of post-vitellogenic egg chambers. There are at least three

ways in which this could have been achieved by the evolved females – (a) increased rate of

oocyte development and oviposition, (b) reduced oocyte arrest at the pre-vitellogenic stages, or

(c) reduced apoptosis at the checkpoints (Drummond-Barbosa, 2001). These three possibilities

are not mutually exclusive. Further investigations are needed to test these theories. Irrespective

of the regime, ovaries of mated females were found to possess fewer mature egg chambers

probably because mating stimulated oviposition (Carlson et al., 1998).

Evolutionary genetics of nutrition-dependent adaptation:

Response in experimental evolution usually depends on standing genetic variation (Teotónio et

al. 2009, Kawecki et al. 2012). Hence, our results directly indicate the presence of a substantial

standing genetic variation in resource acquisition, storage, and allocation traits, in line with the

existing literature (Joshi & Mueller, 1996; Hoffmann & Harshman, 1999; Rion & Kawecki,

2007). Further, though biomass at eclosion seems to be genetically correlated with macronutrient

content (Chippindale et al. 1996, Kristensen et al. 2011, Enriquez et al. 2022), it is evident that

protein and lipid content at eclosion are, at least, partly genetically uncorrelated. Whether these

results were due to changes in protein uptake during larval stage, or metabolic conversion of

other macronutrients to protein, should be a topic of future investigations. Dasgupta et al. (2022)

posed several plausible scenarios in which females can maximize early-life egg output without a

corresponding trade-off with lifespan. We now know that, not only reduction in late-life

fecundity (Dasgupta et al., 2022), increased resource availability to reproduction, which

culminated in increased ovarian function was the mechanistic basis of the observed nutrient

dependent adaptation.

Interestingly, we showed mating-dependent plasticity in ovariole number and rate of production

of mature stages of oocyte, and the evolution of such plasticity. To the best of our knowledge

this is the first report of such evolution of female reproductive physiology. Evolution of

phenotypic plasticity itself is a fairly underexplored area (but see Maggu et al., 2020). Whether

such evolution is fundamentally reliant on the evolution of control elements of ovarian function

remains to be explored. Nonetheless, our results are indicative of plausible involvement of some

signaling pathway that connects nutrient sensing and ovarian function (Durdevic & Ephrussi,

2019), an area yet to be fully explored.

Figure 1: Female whole body protein content (a & b), dry mass (c & d), and lipid content (e & f). Whole body

protein content in females from Experiment 1, quantified at two different adult life-stages – on eclosion and in

fitness window. Dry body mass from Experiment 2, quantified at two different adult life-stages/time-points – on

eclosion and in fitness window. Absolute lipid mass in females from Experiment 3. Whole body absolute lipid mass

of females, from Experiment 2, was quantified at two different adult life-stages – on eclosion and in fitness window.

The cyan box represents data from BL (control regime) females, while the orange box represents that from YLB

(experimental regime). The box thickness represents the Inter-Quartile Range. The black horizontal line in the

middle is the median. Wherever the BL vs. YLB difference is statistically significant, it is indicated with an asterisk

(*).

Figure 2: Important results from Experiment 3 and 5. (a) Larval feeding rate (Experiment 5) was quantified as the

number of sclerite retractions per min in an early third-instar larva. (b): Starvation resistance (Experiment 3) was

quantified as the females' mean longevity under starvation. The cyan box represents data from BL (control regime),

while the orange box represents that from YLB (experimental regime). The box thickness represents the Inter-

Quartile Range. Selection regime had a significant effect on larval feeding rate. Wherever the BL vs. YLB

difference is statistically significant, it is indicated with an asterisk (*).

560 **Figure 3:** Mean ovariole count from Experiment 4. Ovariole number was quantified for virgin and mated females.

561 The cyan box represents data from BL (control regime) females, while the orange box represents that from YLB

562 (experimental regime). The box thickness represents the Inter-Quartile Range. The black horizontal line in the

563 middle is the median and the notch denotes 95% CI around the median. Statistically significant differences, as

564 indicated by multiple comparisons, are indicated with an asterisk (*).

Figure 4: Important results from Experiment 6. (a) a typical DAPI stained image of an ovariole from a BL female, (b) a typical DAPI stained image of an ovariole from a YLB female. The arrowheads points to the different stages of egg chambers. A: Pre-vitellogenic, B: vitellogenic, and C: post-vitellogenic. (c): Percentage of egg chambers at different developmental stages from Experiment 6. The percentage of pre-vitellogenic, vitellogenic, and post-vitellogenic egg chambers was quantified for virgin and mated females. The cyan bar represents data from BL (control regime) females, while the orange bar represents data from YLB (experimental regime) females. The error bar represents standard error of means. Statistically significant differences, as indicated by multiple comparisons, are indicated with an asterisk (*).

Traits	Stage	Effects	SS	MS	Den. DF	F	p
Protein content	On eclosion	Selection regime	16332	16332	85.01	20.68	**<0.01**

Traits		Effect	Estimate	Std. Error	z value	p
	In fitness window	Selection regime	737.1	737.1	2.94	0.35	0.4
Larval feeding rate		Selection regime (SR)	4808.3	4808.3	5.78	25.55	<0.01
		Selection regime (SR)	4.97		0.03	138.99	<0.01
Dry body mass		Time point	20764	20764	2.99	110.33	**<0.01**
		SR × Time point	1383.4	1383.4	5.78	7.35	**0.03**
Absolute lipid mass	On eclosion	Selection regime	571.21	571.21	2.99	4.93	0.11
	In fitness window	Selection regime	2332.5	2332.5	86	17.43	**<0.01**
Relative lipid content	On eclosion	Selection regime	59.9	59.9	2.94	8.33	0.06
	In fitness window	Selection regime	41.84	41.84	2.96	6.32	0.09
Starvation resistance		Selection regime	117.21	117.21	3.07	3.02	0.18
Ovariole count		Selection regime (SR)	2.15	2.15	222.12	0.73	0.39
		Mating status (MSt)	25.57	25.57	222.17	8.62	**<0.01**
		Thorax length	8.22	8.22	189.93	2.78	0.09
		SR × MSt	29.47	29.47	221.89	9.94	**<0.01**
Pre-vitellogenic egg chambers		Selection regime (SR)	965.91	965.91	5.98	4.59	0.07
		Mating status (MSt)	320.10	320.10	3.01	1.52	0.30
		SR × MSt	190.08	190.08	5.98	0.90	0.38

585 **Table 1:** Results of linear mixed effect model (LMM) analyses of various traits. All tests were performed

586 considering α= 0.05 and significant p-values are mentioned in bold.

Total egg chamber count	Selection regime (SR)	-0.09	0.02	-4.42	**<0.01**
	Mating status (MSt)	-0.01	0.02	-0.78	0.44
	SR × MSt	0.07	0.03	2.59	**<0.01**
Vitellogenic egg chamber count	Selection regime (SR)	0.04	0.07	0.63	0.53
	Mating status (MSt)	-0.54	0.07	-7.84	**<0.01**
	SR × MSt	0.10	0.09	1.09	0.28
Post-vitellogenic egg chamber count (egg chamber staging experiment)	Selection regime (SR)	0.37	0.10	3.66	**<0.01**
	Mating status (MSt)	0.99	0.12	8.32	**<0.01**
	SR × MSt	-0.21	0.11	-1.85	0.06

Table 2: Results of generalized linear mixed effect model (GLMM) analyses of various traits. All tests were

performed considering α= 0.05 and significant p-values are mentioned in bold.

References

Adler, M. I., & Bonduriansky, R. (2014). Why do the well☐fed appear to die young? A new

evolutionary hypothesis for the effect of dietary restriction on lifespan. *Bioessays*, *36*, 439-450.

Aguila, J. R., Hoshizaki, D. K., & Gibbs, A. G. (2013). Contribution of larval nutrition to adult

reproduction in *Drosophila melanogaster*. *Journal of Experimental Biology*, *216*(3), 399-406.

Ahmad, M., Keebaugh, E. S., Tariq, M., & Ja, W. W. (2018). Evolutionary responses of

Drosophila melanogaster under chronic malnutrition. *Frontiers in ecology and evolution*, *6*, 47.

Almeida, F., & Suesdek, L. (2017). Effects of Wolbachia on ovarian apoptosis in *Culex*

quinquefasciatus (Say, 1823) during the previtellogenic and vitellogenic periods. *Parasites and*

Vectors, *10*(1), 1–8.

Avila, F. W., Sirot, L. K., Laflamme, B. A., Rubinstein, C. D., & Wolfner, M. F. (2011). Insect

seminal fluid proteins: Identification and function. *Annual Review of Entomology*, *56*, 21–40.

Ballard, J. W. O., Melvin, R. G., & Simpson, S. J. (2008). Starvation resistance is positively

correlated with body lipid proportion in five wild caught *Drosophila simulans* populations.

Journal of insect physiology, *54*(9), 1371-1376.

Bates, D., Mächler, M., Bolker, B., & Walker, S. (2014). Fitting linear mixed-effects models

using lme4. *arXiv preprint arXiv:1406.5823*.

Bierbaum, T. J., Mueller, L. D., & Ayala, F. J. (1989). Density-Dependent Evolution of Life-

History Traits in *Drosophila melanogaster*. *Evolution*, *43*(2), 382.

Bitner, K., Rutledge, G. A., Kezos, J. N., & Mueller, L. D. (2021). The effects of adaptation to

urea on feeding rates and growth in Drosophila larvae. *Ecology and evolution*, *11*(14), 9516-

9529.

Bodenstein, D. (1950). The postembryonic development of Drosophila. *Biology of Drosophila*,

275-367.

Boggs, C. L. (1992). Resource allocation: exploring connections between foraging and life

history. *Functional Ecology*, *6*(5), 508-518.

Bonduriansky, R., Maklakov, A., Zajitschek, F., & Brooks, R. (2008). Sexual selection, sexual

conflict and the evolution of ageing and life span. *Functional Ecology*, *22*(3), 443–453.

Bradford, M. M. (1976). A rapid and sensitive method for the quantitation of microgram

quantities of protein utilizing the principle of protein-dye binding. *Analytical biochemistry*, *72*(1-

2), 248-254.

Carey, J. R., Harshman, L. G., Liedo, P., Müller, H. G., Wang, J. L., & Zhang, Z. (2008).

Longevity–fertility trade☐offs in the tephritid fruit fly, Anastrepha ludens, across

dietary☐restriction gradients. *Aging cell*, *7*(4), 470-477.

Carlson, K. A., Nusbaum, T. J., Rose, M. R., & Harshman, L. G. (1998). Oocyte maturation and

ovariole number in lines of *Drosophila melanogaster* selected for postponed senescence.

Functional Ecology, *12*(4), 514–520.

Cavigliasso, F., Dupuis, C., Savary, L., Spangenberg, J. E., & Kawecki, T. J. (2020).

Experimental evolution of post-ingestive nutritional compensation in response to a nutrient-poor

diet. *Proceedings of the Royal Society B*, *287*(1940), 20202684.

Chippindale, A. K., Chu, T. J. F., & Rose, M. R. (1996). Complex trade-offs and the evolution of

starvation resistance in *Drosophila melanogaster*. *Evolution*, *50*(2), 753–766.

Chippindale, A. K., Leroi, A. M., Kim, S. B., & Rose, M. R. (1993). Phenotypic plasticity and

selection in *Drosophila* life☐history evolution. I. Nutrition and the cost of reproduction. *Journal*

of Evolutionary Biology, *6*(2), 171–193.

Churchill, E. R., Dytham, C., & Thom, M. D. (2019). Differing effects of age and starvation on

reproductive performance in Drosophila melanogaster. *Scientific reports*, *9*(1), 2167.

Colinet, H., & Renault, D. (2014). Dietary live yeast alters metabolic profiles, protein

biosynthesis and thermal stress tolerance of *Drosophila melanogaster*. *Comparative*

Biochemistry and Physiology - A Molecular and Integrative Physiology, *170*, 6–14.

Coyne, J. A., Rux, J., & David, J. R. (1991). Genetics of morphological differences and hybrid

sterility between *Drosophila sechellia* and its relatives. *Genetics Research*, *57*(2), 113-122.

Dasgupta, P., Halder, S., Dari, D., Nabeel, P., Vajja, S. S., & Nandy, B. (2022). Evolution of a

novel female reproductive strategy in Drosophila melanogaster populations subjected to long-

term protein restriction. *Evolution*, *76*(8), 1836-1848.

Desprez, M., Gimenez, O., McMahon, C. R., Hindell, M. A., & Harcourt, R. G. (2018).

Optimizing lifetime reproductive output: Intermittent breeding as a tactic for females in a

long□lived, multiparous mammal. *Journal of Animal Ecology*, *87*(1), 199-211.

Drummond-Barbosa, D., & Spradling, A. C. (2001). Stem cells and their progeny respond to

nutritional changes during *Drosophila* oogenesis. *Developmental biology*, *231*(1), 265-278.

Durdevic, Z., & Ephrussi, A. (2019). Germ cell lineage homeostasis in *Drosophila* requires the

Vasa RNA helicase. *Genetics*, *213*(3), 911-922.

Ellison, P. T. (2003). Energetics and reproductive effort. *American Journal of Human Biology*,

15(3), 342-351.

Enriquez, T., Lievens, V., Nieberding, C. M., & Visser, B. (2022). Pupal size as a proxy for fat

content in laboratory-reared and field-collected Drosophila species. *Scientific Reports*, *12*(1),

12855.

Fontana, L., & Partridge, L. (2015). Promoting health and longevity through diet: From model

organisms to humans. *Cell*, 161(1), 106–118.

Foray, V., Pelisson, P. F., Bel-Venner, M. C., Desouhant, E., Venner, S., Menu, F., ... & Rey, B.

(2012). A handbook for uncovering the complete energetic budget in insects: the van Handel's

method (1985) revisited. Physiological Entomology, 37(3), 295-302.

Gatto, M., Matessi, C., & Slobodkin, L. B. (1989). Physiological profiles and demographic rates

in relation to food quantity and predictability: an optimization approach. *Evolutionary Ecology*,

3, 1-30.

Giraudi, G., Baggiani, C., & Giovannoli, C. (1997). Inaccuracy of the Bradford method for the

determination of protein concentration in steroid-horseradish peroxidase conjugates. *Analytica*

chimica acta, *337*(1), 93-97.

Hodin, J. (2009). She shapes events as they come: Plasticity in female insect reproduction.

Phenotypic Plasticity of Insects□: Mechanisms and Consequences, 99.

Hoffmann, A. A., & Harshman, L. G. (1999). Desiccation and starvation resistance in

Drosophila: Patterns of variation at the species, population and intrapopulation levels. *Heredity*,

83(6), 637–643.

Inness, C. L., & Metcalfe, N. B. (2008). The impact of dietary restriction, intermittent feeding

and compensatory growth on reproductive investment and lifespan in a short-lived fish.

Proceedings of the Royal Society B: Biological Sciences, *275*(1644), 1703-1708.

Jervis, M. A., Ellers, J., & Harvey, J. A. (2008). Resource acquisition, allocation, and utilization

in parasitoid reproductive strategies. *Annu. Rev. Entomol.*, *53*(1), 361-385.

Jia, D., Xu, Q., Xie, Q., Mio, W., & Deng, W. M. (2016). Automatic stage identification of

Drosophila egg chamber based on DAPI images. Scientific reports, 6(1), 18850.

Joshi, A., & Mueller, L. D. (1988). Evolution of higher feeding rate in *Drosophila* due to

density-dependent natural selection. *Evolution*, 42, 1090– 1093.

Joshi, A., & Mueller, L. D. (1996). Density-dependent natural selection in *Drosophila*: Trade-

offs between larval food acquisition and utilization. *Evolutionary Ecology*, *10*(5), 463–474.

Kawecki, T. J., Lenski, R. E., Ebert, D., Hollis, B., Olivieri, I., & Whitlock, M. C. (2012).

Experimental evolution. *Trends in Ecology and Evolution*, *27*(10), 547–560.

Kirkwood, T. B. (1977). Evolution of ageing. *Nature*, *270*(5635), 301-304.

Klepsatel, P., Procházka, E., & Gáliková, M. (2018). Crowding of Drosophila larvae affects

lifespan and other life-history traits via reduced availability of dietary yeast. *Experimental*

Gerontology, *110*, 298-308.

Kolss, M., Vijendravarma, R. K., Schwaller, G., & Kawecki, T. J. (2009). Life-history

consequences of adaptation to larval nutritional stress in *Drosophila*. *Evolution*, *63*(9), 2389–

2401.

Kristensen, T. N., Overgaard, J., Loeschcke, V., & Mayntz, D. (2011). Dietary protein content

affects evolution for body size, body fat and viability in Drosophila melanogaster. *Biology*

letters, *7*(2), 269-272.

Kristensen, T. N., Ketola, T., & Kronholm, I. (2020). Adaptation to environmental stress at

different timescales. *Annals of the New York Academy of Sciences*, 1476(1), 5–12.

Kuznetsova, A., Brockhoff, P. B., & Christensen, R. H. (2017). lmerTest package: tests in linear

mixed effects models. *Journal of statistical software*, 82, 1-26.

Lease, H. M., & Wolf, B. O. (2011). Lipid content of terrestrial arthropods in relation to body

size, phylogeny, ontogeny and sex. *Physiological Entomology*, *36*(1), 29–38.

Lee, K. P., & Jang, T. (2014). Exploring the nutritional basis of starvation resistance in

Drosophila melanogaster. *Functional Ecology*, *28*(5), 1144–1155.

Lenth, R., Singmann, H., Love, J., Buerkner, P., &Herve, M. (2018). Emmeans: Estimated

marginal means, aka least-squares means. *R package version*, 1, 3.

Li, H. M., Buczkowski, G., Mittapalli, O., Xie, J., Wu, J., Westerman, R., ... & Pittendrigh, B. R.

(2008). Transcriptomic profiles of Drosophila melanogaster third instar larval midgut and

responses to oxidative stress. *Insect molecular biology*, *17*(4), 325-339.

Lobell, A. S., Kaspari, R. R., Serrano Negron, Y. L., & Harbison, S. T. (2017). The genetic

architecture of Ovariole number in *Drosophila melanogaster*: Genes with major, quantitative,

and pleiotropic effects. *G3: Genes, Genomes, Genetics*, *7*(7), 2391–2403.

Long, T. A. F., Pischedda, A., Nichols, R. V., & Rice, W. R. (2010). The timing of mating

influences reproductive success in Drosophila melanogaster: implications for sexual conflict.

Journal of Evolutionary Biology, *23*(5), 1024-1032.

Maggu, K., Ahlawat, N., Geeta Arun, M., Meena, A., & Prasad, N. G. (2021). Divergence of

responses to variable socio-sexual environments in laboratory populations of Drosophila

melanogaster evolving under altered operational sex ratios. *Evolution*, *75*(2), 414-426.

Magwere, T., Chapman, T., & Partridge, L. (2004). Sex differences in the effect of dietary

restriction on life span and mortality rates in female and male *Drosophila melanogaster. The*

Journals of Gerontology Series A: Biological Sciences and Medical Sciences, *59*(1), B3-B9.

Mair, W., Sgrò, C. M., Johnson, A. P., Chapman, T., & Partridge, L. (2004). Lifespan extension

by dietary restriction in female Drosophila melanogaster is not caused by a reduction in

vitellogenesis or ovarian activity. *Experimental gerontology*, *39*(7), 1011-1019.

Mair, W., Piper, M. D. W., & Partridge, L. (2005). Calories do not explain extension of life span

by dietary restriction in *Drosophila. PLoS Biology*, 3(7), 1305–1311.

Marron, M. T., Markow, T. A., Kain, K. J., & Gibbs, A. G. (2003). Effects of starvation and

desiccation on energy metabolism in desert and mesic *Drosophila. Journal of Insect Physiology*,

49(3), 261-270.

McCall, K. (2004). Eggs over easy: Cell death in the *Drosophila* ovary. *Developmental Biology*,

274(1), 3–14.

McDonough-Goldstein, C. E., Pitnick, S., & Dorus, S. (2021). Drosophila oocyte proteome

composition covaries with female mating status. *Scientific reports*, *11*(1), 3142.

Mendes, C. C., & Mirth, C. K. (2016). Stage-specific plasticity in ovary size is regulated by

insulin/insulin-like growth factor and ecdysone signaling in *Drosophila*. *Genetics*, *202*(2), 703–

719.

Moatt, J. P., Nakagawa, S., Lagisz, M., & Walling, C. A. (2016). The effect of dietary restriction

on reproduction: a meta-analytic perspective. *BMC Evolutionary Biology*, *16*, 1-9.

Moatt, J. P., Savola, E., Regan, J. C., Nussey, D. H., & Walling, C. A. (2020). Lifespan

extension via dietary restriction: time to reconsider the evolutionary mechanisms?. *BioEssays,*

42, 1900241.

Moore, P. J., & Attisano, A. (2011). Oosorption in response to poor food: Complexity in the

trade-off between reproduction and survival. *Ecology and Evolution*, *1*(1), 37–45.

Mueller, L. D., & Barter, T. T. (2015). A model of the evolution of larval feeding rate in

Drosophila driven by conflicting energy demands. *Genetica*, *143*, 93-100.

Nakagawa, S., Lagisz, M., Hector, K. L., & Spencer, H. G. (2012). Comparative and meta-

analytic insights into life extension via dietary restriction. *Aging Cell*, 11(3), 401–409.

Nandy, B., Joshi, A., Ali, Z. S., Sen, S., & Prasad, N. G. (2012). Degree of adaptive male mate

choice is positively correlated with female quality variance. *Scientific Reports*, 2, 1-8.

Partridge, L., Langelan, R., Fowler, K., Zwaan, B., & French, V. (1999). Correlated responses to

selection on body size in *Drosophila melanogaster*. *Genetics Research*, *74*(1), 43-54.

Partridge, L., Piper, M. D. W., & Mair, W. (2005). Dietary restriction in *Drosophila*.

Mechanisms of Ageing and Development, 126(9 SPEC. ISS.), 938–950.

Pianka, E. R. (1976). Natural selection of optimal reproductive tactics. *American Zoologist*,

16(4), 775-784.

Piper, M. D., Mair, W., & Partridge, L. (2005). Counting the calories: the role of specific

nutrients in extension of life span by food restriction. *The Journals of Gerontology Series A:*

Biological Sciences and Medical Sciences, 60, 549-555.

Pritchett, T. L., Tanner, E. A., & McCall, K. (2009). Cracking open cell death in the *Drosophila*

ovary. *Apoptosis*, *14*(8), 969–979.

Raikhel, A. S., & Dhadialla, T. S. (1992). Accumulation of yolk proteins in insect oocytes.

Rajamani, M., Raghavendra, N., Prasad, N. G., Archana, N., Joshi, A., & Shakarad, M. (2006).

Reduced larval feeding rate is a strong evolutionary correlate of rapid development in Drosophila

melanogaster. *Journal of Genetics*, *85*(3), 209-212.

Rehman, N., & Varghese, J. (2021). Larval nutrition influences adult fat stores and starvation

resistance in *Drosophila*. *PLoS One*, *16*(2), e0247175.

Rion, S., & Kawecki, T. J. (2007). Evolutionary biology of starvation resistance: What we have

learned from *Drosophila*. *Journal of Evolutionary Biology*, 20(5), 1655–1664.

Roff, D. (Ed.). (1993). *Evolution of life histories: theory and analysis*. Springer Science &

Business Media.

Ronai, I., Allsopp, M. H., Tan, K., Dong, S., Liu, X., Vergoz, V., & Oldroyd, B. P. (2017). The

dynamic association between ovariole loss and sterility in adult honeybee workers. *Proceedings*

of the Royal Society B: Biological Sciences, *284*(1851).

Ronai, I., Oldroyd, B. P., & Vergoz, V. (2016). Queen pheromone regulates programmed cell

death in the honey bee worker ovary. *Insect Molecular Biology*, *25*(5), 646–652.

Sarikaya, D. P., Belay, A. A., Ahuja, A., Dorta, A., Green, D. A., & Extavour, C. G. (2012). The

roles of cell size and cell number in determining ovariole number in *Drosophila*. *Developmental*

Biology, *363*(1), 279–289.

Sarikaya, D. P., Church, S. H., Lagomarsino, L. P., Magnacca, K. N., Montgomery, S. L., Price,

D. K., … Extavour, C. G. (2019). Reproductive Capacity Evolves in Response to Ecology

through Common Changes in Cell Number in Hawaiian Drosophila. *Current Biology*, *29*(11),

1877-1884.

Schwasinger☐schmidt, T. E., Kachman, S. D., & Harshman, L. G. (2012). Evolution of

starvation resistance in *Drosophila melanogaster*: measurement of direct and correlated

responses to artificial selection. *Journal of evolutionary biology*, *25*(2), 378-387.

Shanley, D. P., & Kirkwood, T. B. (2000). Calorie restriction and aging: A life☐history analysis.

Evolution, *54*(3), 740-750.

Shu, R., Uy, L., & Wong, A. C. N. (2022). Nutritional phenotype underlines the performance

trade-offs of *Drosophila suzukii* on different fruit diets. *Current Research in Insect Science*, *2*,

100026.

Simmons, F. H., & Bradley, T. J. (1997). An analysis of resource allocation in response to

dietary yeast in *Drosophila melanogaster*. *Journal of insect physiology*, *43*(8), 779-788.

Skorupa, D. A., Dervisefendic, A., Zwiener, J., & Pletcher, S. D. (2008). Dietary composition

specifies consumption, obesity, and lifespan in *Drosophila melanogaster*. *Aging cell*, *7*(4), 478–

490.

Soller, M., Bownes, M., & Kubli, E. (1997). Mating and sex peptide stimulate the accumulation

of yolk in oocytes of *Droosophila melanogaster*. *European Journal of Biochemistry*, *243*(3),

732–738.

Stearns, S. C. (1989). Trade-offs in life-history evolution. *Functional ecology*, *3*(3), 259-268.

Stearns, S. C. (1992). *The evolution of life histories* (Vol. 249, p. xii). Oxford: Oxford university

press.

Stearns, S. C. (2000). Life history evolution: successes, limitations, and prospects.

Naturwissenschaften, *87*(11), 476-486.

Telang, A., & Wells, M. A. (2004). The effect of larval and adult nutrition on successful

autogenous egg production by a mosquito. *Journal of Insect Physiology*, *50*(7), 677-685.

Teleman, A. A. (2010). Molecular mechanisms of metabolic regulation by insulin in *Drosophila*.

Biochemical Journal, *425*(1), 13–26.

Teotónio, H., Chelo, I. M., Bradić, M., Rose, M. R., & Long, A. D. (2009). Experimental

evolution reveals natural selection on standing genetic variation. *Nature genetics*, *41*(2), 251-

257.

Vijendravarma, R. K. (2018). Experimental evolution demonstrates evolvability of preferential

nutrient allocation to competing traits in response to chronic malnutrition. *Journal of*

Evolutionary Biology, *31*(11), 1743–1749.

Wang, L., Wei, D. D., Wang, G. Q., Huang, H. Q., & Wang, J. J. (2023). High-sucrose diet

exposure on larvae contributes to adult fecundity and insecticide tolerance in the oriental fruit

fly, Bactrocera dorsalis (Hendel). *Insects*, *14*(5), 407.

Wayne, M. L., Soundararajan, U., & Harshman, L. G. (2006). Environmental stress and

reproduction in *Drosophila melanogaster*: Starvation resistance, ovariole numbers and early age

egg production. *BMC Evolutionary Biology*, *6*, 1–10.

Williams, G. C. (1957). Pleiotropy, natural selection, and the evolution of senescence. *Evolution*,

11, 398-411.

Zajitschek, F., Georgolopoulos, G., Vourlou, A., Ericsson, M., Zajitschek, S. R. K., Friberg, U.,

& Maklakov, A. A. (2019). Evolution under dietary restriction decouples survival from fecundity

in *Drosophila melanogaster* females. *Journals of Gerontology - Series A Biological Sciences and*

Medical Sciences, 74(10), 1542–1548.

Zajitschek, F., Zajitschek, S. R., Canton, C., Georgolopoulos, G., Friberg, U., & Maklakov, A.

A. (2016). Evolution under dietary restriction increases male reproductive performance without

survival cost. *Proceedings of the Royal Society B: Biological Sciences*, *283*(1825), 20152726.

Zimmerman, J. A., Malloy, V., Krajcik, R., & Orentreich, N. (2003). Nutritional control of

aging. *Experimental gerontology*, 38, 47-52.