Linking Tissue Morphology and Tissue Healing in a Cell-Fate Model

Somya Mani [1,2,*] **and Tsvi Tlusty** [1,3]

[1] *Institute for Basic Science-Center for Soft and Living Matter, South Korea*
[2] *Konrad Lorenz Institute for Evolution and Cognition Research, Austria*
[3] *Department of Physics, Ulsan National Institute of Science and Technology, South Korea*

Correspondence*:
Somya Mani
somyamn@gmail.com

ABSTRACT

Multicellular tissues are immensely diverse, and yet are notably similar at the microanatomical level – most tissues are organized into distinct domains with well-defined cellular compositions and precise adjacencies among themselves. Concurrently, tissues across organisms are also similar in an important functional property: the ability to heal from injury, even in organisms with poor regenerative capacities, such as mammals. Both the cellular organization within tissues and the ability of tissues to heal are outcomes of developmental processes, suggesting that the two may be intrinsically linked. In this work, we explore this connection using an agent-based model of developmental cell-fate decisions. Our model produces a rich diversity of tissue morphologies: By tuning only four parameters, which control the density of intercellular interactions and the propensity of cellular differentiation, we generate tissue structures ranging from disordered and sparse to tissues organized into dense, contiguous domains. Only tissues with large contiguous domains had the ability to heal from injuries, thus demonstrating that tissue healing is strongly coupled to tissue morphology. Moreover, model-generated tissues predominantly heal through the replacement of injured cells by cells dividing in the neighborhood, which recapitulates natural mechanisms in animals as well as plants. More generally, our modeling framework allows a systematic sampling of diverse developmental programs and reproduces healing mechanisms in organisms as phylogenetically distant as plants and animals. Our work thus points to general patterns which often are difficult to directly compare in empirical research. Despite the multiple evolutionary origins of multicellularity, the demonstrated link between tissue domain-level organization and healing capacity may represent a unifying feature of multicellular life.

Keywords: Cell-fate, cell differentiation, cell migration, cellular neighborhood, tissue domains, tissue healing

INTRODUCTION

Tissues are systems of cells of one or more cell-types that together perform specific functions beyond what any single cell is capable of (Adler et al., 2023). Tissue level organization is posited to be essential for the evolution of functionally specialized cells in multicellular organisms (Pavlicev et al., 2024). Correspondingly, tissue level organization is widely distributed across the different multicellular lineages

of animals (Technau and Steele, 2011), land plants (Oliva and Lister, 2023), brown algae (Theodorou and
Charrier, 2021) and fungi (Nagy et al., 2023). In keeping with the diversity of these organisms, their tissue
compositions are also diverse, ranging from the Cnidarian *Hydra* which contains two tissue layers that
have 3-4 cell-types (Holstein, 2023), to complex animals like humans which contain at least sixty tissues
that are composed of tens of cell-types (Hatton et al., 2023).

Irrespective of cellular complexity, cells within any tissue interact closely with one another using molecular
signals and mechanical cues. These inter-cellular interactions shape cellular decisions to divide, die,
differentiate or migrate (O'Connor et al., 2010). These cellular decisions in turn dictate tissue form and
function (Ramos et al., 2024), and tissues across organisms tend to organize into distinct microanatomical
domains that are characterized by their cellular compositions and form niches for inter-cellular interactions:
For example, there are three distinct domains in the mammalian intestinal epithelium: the crypt base
which contains the stem cells, the crypt walls which contain transit amplifying cells and the villi which
contain differentiated cells (Clevers, 2013). Similarly, in plants, the shoot apical meristem is composed of
three domains: the central zone which contains the stem cells, the organizing center whose cells provide
hormonal cues to the central zone, and the peripheral zone which contains differentiated cells (Gaillochet
et al., 2015). At the same time, cellular interactions are also essential for healing of tissues from injuries:
For example, in mammalian muscles (Koike et al., 2022) and spinal cord (Brennan et al., 2022), and in
plants (Sugimoto et al., 2019).

In this work, we developed a spatial agent-based model to explore how the control of cell-fate decisions by
inter-cellular interactions regulates cellular organization within tissues, and how tissue healing is related
to tissue cellular organization. Our model produces a rich variety of tissue-level organizations which,
similarly to real tissues, are arranged into distinct domains. The model's tissues vary widely in the relative
arrangements of these domains, ranging from those with sparse cellular distribution to full and densely
occupied grids, and tissues with small, disperse domains to large, contiguous domains. Crucially, we
found tissue healing to be strongly linked with tissue form: almost all regenerative tissues were densely
occupied with cells which formed large and contiguous domains. Moreover, the model's tissues healed
predominantly via a localized response where cells lost upon injury were replaced through cell-division
and differentiation of neighboring cells. This is the most probable mechanism that emerges robustly across
a wide sampling of cellular interactions, though specific organisms may employ alternative strategies
shaped by their evolutionary histories. Our model also produces experimentally testable predictions
since our parameters represent elementary developmental processes which are amenable to experimental
manipulation. Specifically, we predict that increasing the level of inter-cellular interactions should cause
tissues with contiguous domains to become disordered and reduce their ability to heal.

MODEL OF CELLULAR INTERACTIONS AND CELL-FATES

A tissue in the model is composed of N distinct cell types. These cells are distributed across the tissue's
body: In the model, the tissue is housed in a 2-dimensional grid with G^2 grid-squares and periodic
boundaries (Fig.1(B)). Multiple cells are allowed to reside within any grid-square: $C^{n,g}$ gives the number
of cells of cell-type n in the grid-square g, for $n \in \{1, 2, 3, ...N\}$ and $g \in \{1, 2, 3, ...G^2\}$.

The geometry of the body is given by a distance matrix D: $D(g_a, g_b) = d \in \mathbb{W}$ implies the grid-squares
g_a and g_b are a Manhattan distance d apart. Specifically, g_a and g_b are *adjacent* if $D(g_a, g_a) \leq 1$. In a
2-dimensional grid, each grid-square is adjacent to five grid-squares including itself (Fig.1(B)).

Figure 1. An example of cellular interactions and cell-fate rules for a tissue with $N = 3$ cell-types in a $G = 3$ grid. (A) Cellular interactions: Circles represent distinct cell-types. Sharp arrows indicate stabilizing interactions and blunt arrows indicate destabilizing interactions. Cell-types *1* and *3* are intrinsically stable whereas cell-type *2* is intrinsically unstable. **(B)** Adjacent grid-squares: The grid has periodic boundaries and the dark blue squares are all adjacent to the light blue square. **(C)** An example of Cell-fate rules for cell-types that interact according to (A): Cell-type-1 is intrinsically stable and has only stabilizing interactions, therefore, it neither differentiates not migrates in any cellular neighborhood. We show here cell-fate rules for cell-type-2 (top) and cell-type-3 (bottom). Light blue squares in the first row represent the grid-square occupied by the cell of interest: The cell-type and the location of the grid-square are indicated within the square. The numbers in bold written above the square represent the cellular neighborhood: black numbers indicate the presence of the corresponding cell-type and grey numbers indicate the absence of the corresponding cell-types. Arrows point to the cell-fate of the cell of interest in the corresponding cellular neighborhood: Cellular decisions of unstable cells to differentiate or migrate are are highlighted in blue font.**(D)** A tissue represented by the grid on the left updates to the grid on the right following Cell-Fate Rules given in (C). **(E)** An example of a stable $N = 5$, $G = 15$ tissue. Colors of grid-squares indicate their cellular compositions. We call contiguous regions of the tissue that have the same composition a *domain*: Cells in the center of the domain have a cellular neighborhood identical to the domain composition, while cells at domain edges have richer cellular neighborhoods. Domains have variable sizes: we highlight with dashed borders the three large domains in this tissue. The cellular compositions of these three domains are given alongside the image: 1 indicates the presence of at least one cell of the corresponding cell-type while 0 indicates absence of the corresponding cell-type. Black grid-squares are empty.

Cells in adjacent grid-squares are called *neighbors*. The *neighborhood* of any cell in grid-square g is given

by a N-length binary vector H_g: $H_g(n) = 1$ if a cell of cell-type n is in the neighborhood of square g;

$H_g(n) = 0$ otherwise. There are 2^N distinct possible neighborhoods, which can be arranged in the standard

binary ordering: $H^0 = [0, 0, 0..., 0]$, $H^1 = [0, 0, 0..., 1]$, ..., $H^{2^N-1} = [1, 1, 1..., 1]$.

By allowing multiple cells to reside within the same grid-square, our model presents a coarser picture

of cellular arrangements within tissues compared to standard models where the location, geometry and

neighborhood of each cell (such as in cellular automaton (Lee et al., 1995) and vertex models (Farhadifar

et al., 2007)) or even subparts of cells (such as in the cellular Potts model (Marée et al., 2007)) are explicitly
described. Thus our formalism offers a more general representation of tissues since we do not impose
constraints on the geometries or adjacencies of individual cells (Osborne et al., 2017). At the same time,
the simplicity of our model as compared to the cellular Potts model and lattice-free models (such as in
(Bock et al., 2010; Sandersius and Newman, 2008)) makes it computationally feasible to sample a broad
space of tissue patterns.

Tissue developmental rules

Cellular interactions are prescribed by the interaction matrix I (Fig.1(A), Methods: Generating tissue
developmental rules): A cell of type n_i in square g_a interacts with a cell of type n_j in square g_b if
$I(n_i, n_j) \neq 0$ and g_a and g_b are adjacent. That is, cellular interactions in the model are short to medium-
range and a cell can interact with another cell some tens of cells away. We do not impose any constraints on
the form of cellular interaction and do not distinguish between molecular and mechanical signals (Pavlicev
and Wagner, 2023). The model's cellular interactions can represent short-range molecular signaling,
which are known to be important during tissue development (Armingol et al., 2024), for example, during
embryonic development across animals (Perrimon et al., 2012) and during the growth and development
of tissues in the plant shoot (Gaillochet et al., 2015). The model's cellular interactions can also represent
mechanical signals (Chan et al., 2017) such as via the local modification of the extracellular matrix by one
of the cells which impacts the behavior of nearby cells. For example, during skull development in mice,
differentiated osteoblasts secrete collagen into the extracellular matrix and induce differentiation of nearby
mesenchymal cells (Dang et al., 2023).

We call a cellular interaction *stabilizing* if $I(n_i, n_j) = 1$ and *destabilizing* if $I(n_i, n_j) = -1$. Particularly,
a cell-type n_i is *intrinsically stable* if $I(n_i, n_i) = 0$ or 1 and *intrinsically unstable* if $I(n_i, n_i) = -1$.
Intrinsically stable cells are destabilized in the presence of any destabilizing interactions and intrinsically
unstable cells are stabilized in the presence of any stabilizing interactions. For example, the germline stem
cells (GSC) in the worm *Caenorhabditis elegans* are intrinsically unstable and tend to differentiate into
germ cells. Interaction of GSCs with distal tip cells stabilizes them by repressing their differentiation (Byrd
and Kimble, 2009).

A stable cell maintains its cell-type and its position within the body, whereas an unstable cell either
differentiates into a new cell-type or migrates to a new grid-square. In the model, we allow a migrating cell
to move a randomly chosen distance $m \leq G$ away. In order to maintain tissue connectedness, cells in the
model can only migrate through grid-squares that are already occupied, though they are allowed to migrate
into an empty grid-square which is adjacent to an occupied square.

Additionally, unstable cells of the same cell-type behave differently in different cellular neighborhoods
(Fig.1(C)): For example, let a cell of cell-type n_i be unstable in two distinct neighborhoods H^a and
H^b. In neighborhood H^a, the cell differentiates into cell-type n_j; i.e. $n_i | H^a \rightarrow [n_j, m = 0]$, while in
neighborhood H^b, the cell migrates; i.e. $n_i | H^b \rightarrow [n_i, m = m_1]$. In nature, the effect of such positional
inputs on cell-fate are especially apparent in migratory multipotent stem cells, such as neural crest cells in
zebrafish (Subkhankulova et al., 2023).

While for the most part, plasticity of cell-fate is restricted to stem cells, under special circumstances,
such as stem cell ablation, differentiated cells can revert to a stem-like state capable of proliferation
and differentiation (Yao and Wang, 2020). Hence in the model, we do not impose any restrictions on
cell-fate potential, and instead allow hierarchical structures in cell-fate potential to emerge from the
interplay of cellular interactions. Cell Fate Rules in the model comprise the set of all cell-fate transitions

$n_i|H^a \to [n_j, m]$, such that $n_i, n_j \in \{1, 2, ..., N\}$, $m \leq G$ and $a \in \{0, 1, ..., 2^N - 1\}$ (see Methods:
Generating tissue developmental rules).

In addition to cellular interactions, differentiation and migration, cells in the model can also divide and die
(Methods:Updating tissue states). In nature, cell proliferation is often regulated by cell density (Fan and
Meyer, 2021; Pavel et al., 2018). To capture this in the model, cell division and cell death are regulated by
the number of cells in grid-squares: Let TC^g be the total number of cells in some grid-square g. Any cell in
grid-square g divides into an identical cell-type with probability $P_{div} = max(0, (10 - TC^g)/9)$ and a cell
dies with probability $P_{death} = min(0.8, 0.02(TC^g - 10))$. The fixed cell-division and death parameters
we use lead to grid-squares having on average 10 cells; therefore the neighborhood of a single cell, which
comprises five adjacent grid-squares (Fig.1(B)), is composed of 50 cells. The fixed parameters in these
formulae can be tuned to suit different cellular densities.

The daughter cell of a dividing cell in grid-square g can be born either in the same square or in some
adjacent square. In nature, many cells divide asymmetrically during embryonic development, and in adults,
tissue stem cells divide asymmetrically to produce daughter cells with distinct fates (Sunchu and Cabernard,
2020). In the model, daughter cells are identical to each other immediately after cell division, but cells
born in adjacent grid-squares can end up with distinct cell-fates, which allows us to capture the effects of
asymmetric cell-division.

Model parameters

We draw the interaction matrix I using parameters $0 \leq P_{den} \leq 1$ and $0 \leq P_{stable} \leq 1$: P_{den} controls the
level of inter-cellular signaling – $I(n_i, n_j) \neq 0$ with probability P_{den}; P_{stable} controls cell-fate potential –
$I(n_i, n_j)$ is positive with probability P_{stable}.

We draw Cell Fate Rules using parameter $0 \leq P_{diff} \leq 1$: for some cell of cell-type n_i that is unstable in
neighborhood H^a, the cell differentiates with probability P_{diff} and migrates otherwise. In neighborhood
H^a, if cell-type n_i differentiates, the final cell-type n_j is chosen uniform randomly from among the
N distinct cell-types. Instead if cell-type n_i migrates, the number of grid-squares it migrates is drawn
uniformly from within a distance $m \leq G$. Together, the interaction matrix I and the Cell Fate Rules
comprise *tissue development rules* (Methods:Generating tissue developmental rules). *Tissue development
rules*, once drawn, remain fixed throughout a simulation.

Finally, the parameter $0 \leq F_{adj} \leq 1$ controls the location of daughter cells after cell division: A fraction
$1 - F_{adj}$ of daughter cells are born within the same grid-square as the mother cell, and a fraction F_{adj}
are born in a different adjacent grid-square. F_{adj} is a complex parameter that captures aspects of cellular
organization within grid-squares, including cell shape (McKinley et al., 2018) and the directions of the
axes of cell-division (Gillies and Cabernard, 2011).

At any time-step t, the state of the tissue is completely characterized by the cellular compositions of
its grid-squares $C_t^{n,g}$ for all $n \in \{1, 2, ...N\}$ and $g \in \{1, 2, ...G^2\}$. At $t = 0$, we initialize the tissue by
randomly placing 20 cells of random cell-types across the connected set of grid-squares $[0, 1, 2, 3, 4]$. We
update the tissue by synchronously updating the state of each cell in the tissue according to the *tissue
development rules* (Fig.1(D), Methods:Updating tissue states). We simulate tissue development in the
model for $T = 500$ time-steps. We explore the spatial features and cellular compositions of tissues in the
final time-step (Fig.1(E)), and analyze how tissues grow and heal from injuries.

RESULTS

159 **Model-generated tissues are rich in cell-types**

Figure 2. Number of tissue cell-types and their cell-fate potential. **(A)** Histogram of the number of cell-types in model-generated tissues across all parameters. **(B)** Histogram of cell-fate potential averaged across all cell-types in a tissue. **(C)** Heat map of the fraction of tissues with number of cell-types given by the rows and average cell-fate potential given by the columns.

We first analyse how Cell Fate Rules regulate the cellular compositions of model-generated tissues. A
majority of model-generated tissues across all parameters contain all 5 cell-types in the final time-step
(Fig.2(A), Supplementary Information:FigS2). We find that the number of tissue cell-types increases with
the probability of stabilizing interactions, P_{stable} (Pearson's $r = 0.15$) and decreases with the propensity
of cells to differentiate, P_{diff} ($r = -0.41$). Intuitively, we expect that increasing the number of cellular
interaction partners should increase the probability of destabilizing interactions. In keeping with this, we
find that the number of tissue cell-types decreases with P_{den} which controls the probability of inter-cellular
interactions ($r = -0.14$).

To understand the nature of tissue cell-types, we calculate their cell-fate potentials as the set of all distinct
cell-types which it differentiates into across all possible cellular neighborhoods (Fig.2(B), Supplementary
Information:FigS2). Note that here cell-fate potential refers to the set of cell-types which any given cell-
type *directly* differentiates into. Cell-fate potentials are expected to be wider if one additionally considers
indirect differentiation involving multiple transitions. Average cell-fate potential in the model decreases
with P_{stable} ($r = -0.31$), and increases with P_{diff} ($r = 0.62$). Average cell-fate potential also increases with
P_{den} ($r = 0.28$): Increasing cellular interaction partners in turn increases the probability that a cell has
different fates in different cellular neighborhoods.

Overall, model-generated tissues with more cell-types tend to have lower average cell-fate potential
(Fig.2(C), Pearson's r between number of tissue cell-types and average cell-fate potential $= -0.35$). That
is, most model-generated tissues tend to be rich in cell-types and the constituent cell-types tend to
have low plasticity. Nevertheless, 31% (57%) of model-generated tissues contain at least one cell-type
which can directly (through multiple transitions) differentiate into all 5 tissue cell-types (Supplementary
Information:FigS3). These highly plastic cell-types in the model's tissues are reminiscent of stem cells in
regenerative tissues in real organisms (Fuchs and Blau, 2020).

Figure 3. Sectors define qualitatively distinct tissues. (A) Spread of model-generated tissues across PC1 and PC2. Each point represents the state of a model-generated tissue at the final time-point. Tissues belonging to the 4 different sectors are indicated by different colors (yellow = stunted, green = sparse-contiguous, orange = full-contiguous, purple = random-disperse). Lines indicate borders between sectors. Red points represent examples of tissues from each sector shown in (E-H). (B) Values of *dispersity* across model-generated tissues. (C) Tissue *fullness*, which we calculate as the number of occupied grid-squares, across model-generated tissues. (D) Histograms for model parameters (rows) encoding tissues belonging to different sectors (columns). (E-H) Representative model-generated tissues belonging to different sectors: (E)*Stunted* tissues show very little growth; (F) the *random-disperse* sector contains tissues in which the grid is densely populated by small contiguous domains. (G) the *full-contiguous* sector contains tissues in which the grid is densely populated by large contiguous domains; (H) the *sparse-contiguous* sector contains diverse forms of tissues. Depending on the distance of the tissue from the sector boundaries, it can resemble tissues in the stunted sector (top-left image), or in the full-contiguous sector (bottom-right image), or be a sparse and disperse tissue (bottom-left image). Importantly, this sector contains tissues where the grid is sparsely populated, yet the domains are contiguous.

Tissues vary widely in the fullness and contiguity of cellular arrangements

Tissues are organized into domains

In nature, cells of different types are not randomly scattered in a tissue; rather, they are arranged into
distinct, but functionally and developmentally interconnected domains. Each tissue domain has a defined
cellular composition and sits in a specific position with respect to other tissue domains.

Model-generated tissues resemble natural tissues and are organized into domains. 77% of tissues reach
steady states where each cell is part of a stable cellular neighborhood (Supplementary Information: FigS1).
Since cellular neighborhoods also encompass cells in adjacent grid-squares, at steady state, contiguous grid-
squares tend to be arranged into domains defined by their cellular compositions (Fig.1(E)). We characterize
tissues using five features that describe the spatial arrangement of cells and domains (Methods:Analysing
final tissues, Supplementary Information:FigS4):

1. *Fullness*, or the number of grid-squares in the final tissue that are occupied by cells, describes tissue size
and sparsity (Fig.3(C)).
2. The *number of distinct tissue domains* describes how cells of different types are distributed into distinct
stable cellular neighborhoods.
3. *Coverage equality* describes whether or not the tissue is dominated by a domain of one particular cellular
composition, as opposed to being more uniformly covered by different domains.
4. *Dispersity* describes whether domains of identical cellular composition are present as a single connected
region, or are scattered across the tissue in smaller disjoint regions (Fig.3(B)).
5. *Mixedness* describes the diversity of cellular compositions of grid-squares adjacent to any given grid-
square. For example, grid-squares at the edges of domains are more *mixed* than those in the center; in this
sense, all disperse tissues are expected to have a high overall mixedness, but some non-disperse tissues
with large but intertwined domains (resembling noodles) also have high *mixedness*.

Of these five features, *dispersity* and *mixedness* are related to clinically relevant quantities that are calculated
from spatial omics studies (Feng et al., 2023).

Model-generated tissues fall into four sectors

We perform a Principal Components Analysis of all model-generated tissues against these five features
and find that the first two Principal Components (PC1 and PC2) capture 86% of the variance in the data
(Methods:Analysing final tissues, Supplementary Information:FigS5, FigS6). Correlations between the
PCs and the five tissue features indicate that tissues mostly vary in *dispersity* and *fullness* (Fig.3(B,C)).

We also find that model parameters are significantly predictive of tissue spatial properties: we can divide
the plane described by PC1 and PC2 into four sectors such that for 126 out of the 135 parameter sets tested,
at least 50 out of 100 tissues independently generated using a single parameter set lie within a single sector
(Fig.3(A), black lines). Note that the Principal Components themselves do not separate these tissues into
distinct clusters, and tissue properties vary more or less continuously (Supplementary Information:FigS7).
Our separation of tissues into sectors is for instructive purposes and for ease of discussion.

Sectors capture qualitative features of tissue morphology and spatial distributions of tissue domains;
appropriately, we name these four sectors *stunted* (17% of tissues, Fig.3(E)), *random-disperse* (19%
of tissues, Fig.3(F)), *full-contiguous* (48% of tissues, Fig.3(G)) and *sparse-contiguous* (16% of tissues,
Fig.3(H)).

Our model parameters are based on local cellular interactions, nonetheless they provide clues about global
cellular spatial arrangements: for example, the *stunted* sector is populated by tissues whose cells tend
to be non-interactive (low P_{den}) and intrinsically stable (high P_{stable}). Additionally, growth of these
tissues is hampered because daughter cells tend to be born within the same grid-square as the mother cell
rather than an adjacent square (low F_{adj}) and the levels of cell migration are low (high P_{diff}) (Fig.3(D),
first column). In contrast, the *random-disperse* sector is populated by tissues with highly interactive
and unstable cells that are highly migratory and tend to produce daughter cells in adjacent grid-squares

(Fig.3(D), fourth column). Model parameters that generate *sparse-contiguous* and *full-contiguous* tissues
are more moderately distributed than in the *stunted* and *random-disperse* sectors (Fig.3(D), second and
third columns).

Figure 4. Tissue growth trajectories and sector crossings. Representative tissue growth trajectories projected onto PC1 and PC2 for tissues whose final states fall into (A) *stunted*, (B) *random-disperse*, (C) *sparse-contiguous*, (D) *full-contiguous* sectors. Blue circles represent the initial state of tissues; red circles represent final states of tissues; colored lines indicate growth trajectories. Insets: Graphs indicating probabilities of sector crossings. Nodes (colored circles) represent different sectors (yellow = stunted, green = sparse-contiguous, orange = full-contiguous, purple = random-disperse). Numbers on edges (arrows) indicate the fraction of updates in a trajectory that take a tissue from the source to the target sector averaged across all trajectories for (A) *stunted*, (B) *random-disperse*, (C) *sparse-contiguous*, (D) *full-contiguous* tissues. Numbers within circles represent the average fraction of updates that stay within the corresponding sector.

Tissue development resembles a sequence of sector crossings

In real organisms, tissue development is expressed as a sequence of intertwined morphogenetic and cell-fate
changes (Thowfeequ and Srinivas, 2022; Armenta-Medina et al., 2021). In our model, the four sectors
capture tissue morphology as well as cellular compositions and spatial arrangements of tissue domains.
Therefore, we project the states of growing tissues onto the two Principal Components described earlier
(Fig.3(A)) and express tissue development as sequences of sector crossings (Fig.4, Methods:Analysing
tissue developmental trajectories, Supplementary Information:FigS8).

We summarize tissue developmental trajectories of different tissue types (stunted, full-contiguous, sparse-
contiguous and random-disperse) as graphs whose nodes represent sectors and edges represent sector
crossings (Fig.4, insets). The edges of the graph are weighted to reflect the fraction of developmental
update steps where a particular sector crossing occurs averaged across all tissues of a given tissue-type.
For example, we see from the inset in Fig.4(D) that all *full-contiguous* tissues have a *sparse-contiguous*
intermediate stage. A majority of these tissue developmental trajectories involve only two sector crossings:
stunted → sparse-contiguous → full-contiguous, with multiple update steps within each of the intermediate
sectors (e.g. Fig.4(D), red and blue trajectories). A small number of trajectories transition via the *random-*
disperse sector and some of these trajectories involve oscillations between the *random-disperse* sector and
the final *full-contiguous* sector (e.g. Fig.4(D), black trajectory).

Full and contiguous tissues heal well

Simulating tissue injury and recovery

We test the healing ability of model-generated tissues by simulating injuries to final time-step tissues and
assessing their ability to return to their original states (Methods:Analysing tissue healing). We perform
three types of injuries:

type A: *Single square-all cells* – we ablate all cells of any cell-type from a single randomly chosen grid-square
(Fig.5(A), first row).
type B: *Adjacent squares-single cell-type* – we ablate all cells of a single randomly chosen cell-type from
all grid-squares adjacent to a single randomly chosen grid-square. This amounts to a single cell-type
being ablated from 5 contiguous grid-squares (Fig.5(A), second row).
type C: *5 neighbors-single cell-type* – we ablate all cells of a single randomly chosen cell-type from all
grid-squares that can be reached in ≤ 5 steps from a randomly chosen grid-square. This amounts to a
single cell-type being ablated from 61 contiguous grid-squares, which is 27% of the tissue's body
(Fig.5(A), third row).

After injuring a tissue, we allow it to heal by updating the injured tissue for 100 time-steps using the same
Cell Fate Rules originally used during tissue development (see Methods:Analysing tissue healing). We
assess the ability of a tissue to heal by comparing the Euclidean distance between the state of the tissue
before injury and immediately after injury ($IM_{t=0}$) against the distance between the state of the tissue
before injury and after 100 time-steps of healing ($IM_{t=100}$). We say that a tissue heals if $IM_{t=100} < IM_{t=0}$;
degrades if $IM_{t=100} > IM_{t=0}$; does not recover if $IM_{t=100} = IM_{t=0}$. Notably, cell migration and cell
division are probabilistic decisions in our model, therefore tissues are not expected to heal perfectly
and do not achieve the exact configuration before injury (Fig.5(B)). We perform 25 independent injuries
of type A and type B for each tissue and 10 independent injuries of type C for each tissue and report
the fraction of injuries for which the tissue heals (*recovery-fraction*: Fig.5(A), first column), does not
recover (*no-recovery-fraction*: Fig.5(A), second column), or degrades (*degradation-fraction*: Fig.5(A),
third column).

In order to gain some intuition for the process of tissue healing, consider how cell-fate in the model depends
on the cellular neighborhood: For cells present in the middle of a domain, ablation of one or more cells in
a single adjacent grid-square does not change its neighborhood, and thus does not influence its behavior.
Therefore, injuries affecting cells in a single grid-square, such as type A injuries, are typically not sufficient
to evoke a healing response; these injuries only perturb the tissue if the affected grid-square is at the edge
of a domain. In keeping with this intuition, the *degradation-fraction* for type A injuries is highest in tissues

Figure 5. Tissue responses to injury. (A) Tissue healing ability. Points in the scatter plots indicate distinct tissues. Colors of points indicate the fraction of injuries for which the tissue showed recovery (left column), no recovery (middle column) or degradation (right column). Rows indicate injury type: first row – all cells are ablated from a single random grid-square, second row – all cells of a single random cell-type are ablated from a randomly chosen set of adjacent grid-squares, third row – all cells of a single random cell-type are ablated from a randomly chosen set of adjacent grid-squares. (B) Tissue recovery trajectories. We demonstrate three distinct modes of tissue recovery prevalent in the model for type C injuries: first row – the ablated cell-type is replenished through cell division in adjacent grid-squares, second row – the injury occurs at a boundary between two domains (white box), thus promoting cell division in both domains, ultimately leading to a shifted domain boundary in the tissue after healing, third row – the injury affects cell fates throughout the tissue, ultimately destabilizing a domain (red domain in white box) leading to its loss.

with high *mixedness*, in which a majority of grid-squares lie on the edges of domains (Pearson's *r* between
degradation-fraction and *mixedness* for type A injuries = 0.51 (p-value = 0.0)). Overall, for type A injuries,
61.5% of all tissues tend to neither heal nor get worse, and the *no-recovery-fraction* is greater than 0.5
(green dots in Fig.5(A) second column, row 1).

In contrast, type B and type C ablations are bound to change cellular neighborhoods for cells occupying
grid-squares in the middle of the ablation. We find that the ability to heal from type B and type C injuries was
enriched in the *full-contiguous* sector: For type B injuries, only 16.8% of all tissues have a recovery-fraction
≥ 0.5 , and 99.4% of these tissues are in the *full-contiguous* sector (Supplementary Information:FigS9).
For type C injuries, 24.7% of all tissues have a recovery-fraction ≥ 0.5 , and 97.1% of these tissues are in
the *full-contiguous* sector. These results reveal a strong connection between tissue morphology and healing,
and highlight evolutionary constraints on tissue architecture stemming from the requirement of tissues to
heal. We expect that these constraints apply universally to all multicellular groups despite their independent
origins (Birnbaum and Alvarado, 2008).

Mechanism of tissue recovery in the model

In order to understand the general mechanism of tissue healing in the model, we tested how model
parameters are correlated with *recovery-fractions*: for both type B and type C injuries, the ability to heal
shows the strongest correlation with F_{adj} (*r* = 0.44 for type B and *r* = 0.30 for type C, see Supplementary
Information:FigS10). This suggests that the mechanism of tissue healing in the model depends on cell
division in adjacent grid-squares that replenishes lost cells in the injury sites (for example Fig.5(B),rows
1,2). This mechanism is notably similar to that seen in mammalian tissues (Clevers, 2013; Wei et al., 2021),
as well as during tissue healing in plants, where cells adjacent to injuries switch their cell division axes in
order to replace lost cells (Marhava et al., 2019).

DISCUSSION

We present here an agent-based spatial model of cell-fate decisions in tissues. Notably, our model trades the
ability to explicitly encode the exact cellular neighbors and cell shapes that are captured in more traditional
models (Farhadifar et al., 2007; Marée et al., 2007; Bock et al., 2010) for simplicity, generality, and the
ability to survey vast spaces of tissue developmental rules. We use our model to explore the diversity of
tissue architectures produced by simple developmental rules and the association between tissue architecture
and tissue healing. We categorize model-generated tissues into four sectors – *stunted*, *sparse-contiguous*,
full-contiguous and *random-disperse* – which represent tissue morphology as well as cellular compositions.
These sectors capture a wide variety of tissue architectures: For example, most natural tissues tend to be
contiguous and ordered, such as animal epithelia (Buckley and St Johnston, 2022), mammalian striated
muscle bundles (Mukund and Subramaniam, 2020), and plant leaf epidermis (Zuch et al., 2022), which are
examples of tissues resembling *full-contiguous* tissues. In contrast, the leaf spongy parenchyma forms a
sparse but contiguous tissue (Borsuk et al., 2022) and resembles tissues in the *sparse-contiguous* sector,
while animal blood resembles model-generated *random-disperse* tissues (Nagahata et al., 2022).

We characterize tissue growth in the model in terms of crossings between different sectors, which
recapitulates the manner in which tissue development in real organisms is described, where morphological
changes are coupled with cell-fate changes. For example, during human embryo development, compaction
of the ball of cells during Morula is coupled with cellular differentiation into the inner cell mass and the
trophectoderm, and subsequent cavitation of the ball of cells is coupled with differentiation of the inner
cell mass into the epiblast and the hypoblast (Thowfeequ and Srinivas, 2022). These cell-fate decisions are

not cell autonomous; instead, all through this process, different cell-types communicate with each other
and play an important role in regulating each other's cellular decisions (Okubo et al., 2023).

Importantly, we find a strong association between model-generated tissue structures with their ability
to regenerate; particularly, the ability to heal is enriched in *full-contiguous* tissues. Thus our results
indicate that tissue healing could act as a selective pressure that moulds tissue architecture. Moreover, the
mechanism of healing in the model is through replacement of damaged cells by cell-division in adjacent
regions; this resembles the process of healing in plant (Marhava et al., 2019) as well as animal tissues
(Clevers, 2013; Wei et al., 2021). However, our model is statistical in nature; therefore, it identifies the most
probable mechanisms across a wide range of modes of cellular interaction. Thus outputs of our model may
not match specific mechanisms in particular organisms: For example, regeneration in planarian flatworms
depends on widely circulating neoblast cells (Reddien, 2018), which is in contrast to the local mechanism
enriched in the model.

Overall, our spatial model of cell-fate captures essential features of tissue growth, form and function.
Our model also maps how varying local intercellular interactions can lead to changes in global tissue
architecture, which in turn is closely related to the ability of tissues to heal. These inter-related tissue
properties allow experimental tests of the model's predictions. We highlight here one such experimental
test: our model predicts that increasing the density of inter-cellular interactions should lead to more
disperse tissues (Fig.3(D), first row), and disperse tissues tend to have low healing capacity (Fig.5). To
experimentally test this prediction, one could tune the density of cellular interaction by manipulating the
types of receptors expressed on tissue cell-types and observe the ensuing effects on tissue architecture
and tissue healing. Additionally, our framework is flexible and can be modified to encode special features
of particular tissues; we highlight two natural extensions: First, while two- and three-dimensional grids
may be appropriate to study many solid tissues, tissues such as the plant spongy mesophyll (Borsuk et al.,
2022), or the branched animal lung epithelium (Schittny, 2017) can be more faithfully represented with
more irregular geometries. Second, the model can be expanded by including external signals associated
with different grid locations irrespective of their cellular compositions, which could represent morphogen
gradients that form an integral aspect of animal (Romanova-Michaelides et al., 2022) as well as plant
development (Hu et al., 2021).

METHODS

Sampling the space of tissue developmental parameters

We simulate tissues with $N = 5$ cell-types in a $G \times G$ 2-dimensional grid with periodic boundaries, where
$G = 15$. The geometry of the tissue is described by the grid's adjacency matrix A, where $A(g_i, g_j) = 1$
if grid-squares g_i and g_j are adjacent to each other. In a 2-dimensional grid, each square is adjacent to
4 other squares, as shown in Fig.1(B). We uniformly survey parameter values from the following sets:
$P_{den} \in \{0.3, 0.5, 0.7\}$, $P_{stable} \in \{0.3, 0.5, 0.7\}$, $P_{diff} \in \{0, 0.3, 0.5, 0.7, 1\}$ and $F_{adj} \in \{0.3, 0.5, 0.7\}$.
For each combination of parameter values, we draw 100 replicate *tissue development rules*, and overall,
we examine tissues resulting from 13,500 distinct rules. For 77% of these simulated tissues, cellular
compositions of all grid-squares reach steady state within 400 time-steps.

Generating tissue developmental rules

Interaction matrix

The interaction matrix I is a matrix of size $N \times N$ where

$I(n_i, n_j) = 0$ if $Ber(P_{den}) = 0$ and implies that cell n_j does not interact with n_i,

$I(n_i, n_j) = 1$ if $(Ber(P_{den}) = 1) \bigwedge (Ber(P_{stable}) = 1)$ and implies that n_i stabilizes n_j,

$I(n_i, n_j) = -1$ if $(Ber(P_{den}) = 1) \bigwedge (Ber(P_{stable}) = 0)$ and implies that n_i destabilizes n_j

Where $Ber()$ symbolizes a Bernoulli random variable. Note that the interaction matrix for each tissue is
randomly generated according to parameters P_{den} and P_{stable} and remains fixed for the duration of the
simulation.

Cell Fate Rules

Let $H^a = 0, 1^N$ be an N-length binary vector describing an arbitrary cellular neighborhood, such that
$H^a(n) = 1(0)$ indicates the presence (absence) of cell-type n in neighborhood H^a. An N-length vector
$Stable^a$ encodes the stability of cell-types present in the neighborhood H^a.

$Stable^a(n_i) = 1$ and cells of cell-type n_i are stable in neighborhood H^a if:

• it is intrinsically stable and receives no destabilizing interactions: that is, $I(n_i, n_i) = 0 \, or \, 1$, and
$\nexists n_j s.t.(H^a(n_j) = 1) \bigwedge (I(n_j, n_i) = -1)$
• it is intrinsically unstable and receives stabilizing interactions: that is, $I(n_i, n_i) = -1$, and
$\exists n_j s.t.(H^a(n_j) = 1) \bigwedge (I(n_j, n_i) = 1)$

$Stable^a(n_i) = -1$ and cells of cell-type n_i are unstable in neighborhood H^a if:

• it is intrinsically stable and receives destabilizing interactions: that is, $I(n_i, n_i) = 0 \, or \, 1$, and
$\exists n_j s.t.(H^a(n_j) = 1) \bigwedge (I(n_j, n_i) = -1)$
• it is intrinsically unstable and receives no stabilizing interactions: that is, $I(n_i, n_i) = -1$, and
$\nexists n_j s.t.(H^a(n_j) = 1) \bigwedge (I(n_j, n_i) = 1)$

$Stable^a(n_i) = 0$ for cell-types not present in neighborhood H^a. Unstable cells either differentiate or
migrate. For cells present in H^a, we encode Cell Fate Rules as a pair of look-up tables: A differentiation
table $Diff^a$ and a migration table Mig^a.

• $Diff^a(n_i) = n_j$ indicates that cell-type n_i differentiates into cell-type n_j in neighborhood H^a. If
$(Stable^a(n_i) = -1) \bigwedge (Ber(P_{diff}) = 1)$, n_j is randomly drawn from among the N cell-types. On the
other hand, if $(Ber(P_{diff}) = 1) \bigvee (Stable^a(n_i) = 1)$, then $n_j = n_i$.
• $Mig^a(n_i) = m$ such that $m \leq G$ indicates that cell-type n_i in neighborhood H^a migrates to a grid-
square a distance m away from its current grid-square. If $(Stable^a(n_i) = -1) \bigwedge (Ber(P_{diff}) = 0)$, m is
uniform randomly drawn from $0, 1, 2, ...G$. On the other hand, if $(Ber(P_{diff}) = 1) \bigvee (Stable^a(n_i) = 1)$,
then $m = 0$.

Note that although Cell Fate Rules for cellular differentiation and cell migration are randomly generated
according to the parameter P_{diff}, they remain fixed for the duration of the simulation.

Updating tissue states

At any time-step, the state of the tissue is completely characterized by the cellular compositions of its
grid-squares: $C_t^{n,g}$ gives the number of cells of cell-type n for $n \in 1, 2, 3, ...N$ in grid-square g for
$g \in 1, 2, 3, ...G^2$ at time-step t. At $t = 0$, we initialize the tissue by randomly placing 20 cells of random

cell-types across the connected set of grid-squares $0, 1, 2, 3, 4$. At each time-step of the simulation, the state
of the tissue is updated through sequential steps of cell death, cell division, and cell differentiation and
migration.

Cell death

403 In the tissue at time t, the total number of cells in a grid-square g is given by $TC_t^g = \sum_{n=1}^N C_t^{n,g}$. Then the
404 probability of cell-death in this grid-square is given by

$$P_{death}^g = min(0.8, 0.02(TC_t^g - 10)) \text{ if } TC_t^g > 10,$$

$$P_{death}^g = 0 \text{ otherwise.}$$

407 That is, there is no cell-death in grid-squares with 10 or fewer cells, and the probability that any cell dies
408 increases linearly as the number of cells in the grid-square increases beyond 10.

409 We draw the number of cells of cell-type n in grid-square g that die in this time-step, $\phi_t^{n,g}$ from the binomial
410 distribution $Bin(C_t^{n,g}, P_{death}^g)$. Then, $RC_t^{n,g} = C_t^{n,g} - \phi_t^{n,g}$ gives the cells of cell-type n that remain in
411 grid-square g.

Cell division

After cell death, the total number of cells remaining in some grid-square g_i is given by $RC_t^{g_i} =$
$\sum_{n=1}^N RC_t^{n,g_i}$. Then the probability of cell-division in this grid-square is given by $P_{div}^{g_i} = max(0, \frac{10-RC_t^{g_i}}{9})$.
That is, there is no cell-division in unoccupied grid-squares and in squares with 10 or more cells, and the
probability that any cell divides decreases linearly as the number of cells in the grid-square increases up to
10.

We draw the number of cells of cell-type n in grid-square g_i that divide in this time-step, θ_t^{n,g_i} from the
binomial distribution $Bin(RC_t^{n,g_i}, P_{div}^{g_i})$. θ_t^{n,g_i} is then the number of daughter cells of cell-type n produced
in grid-square g_i.

A fraction $(1 - P_{adj})$ of these daughter cells are born within grid-square g_i. The rest of the daughter cells
are assigned to randomly chosen grid-squares adjacent to g_i. Let $DC_{g_i}^{n,g_j}$ be the number of daughter cells
of cell-type n produced by division of cells in grid-square g_i that are born in grid-square g_j. For any g_j not
adjacent to g_i, $DC_{g_i}^{n,g_j} = 0$.

Then, $C_t^{n,g_j} = RC_t^{n,g_j} + \sum_{g_i=1}^{G^2} DC_{g_i}^{n,g_j}$ gives the total number of cells of type n in grid-square g_j after
cell-division.

Note that in the model, the probability of cell-division is the same for all cells of all cell-types in a given
grid-square. Additionally, two cells of the same cell-type in the same grid-square may divide differently:
Depending on P_{div}^g, it is possible that only one of these cells divides. In case both these cells divide,
depending on F_{adj}, it is also possible that the daughters are produced in different grid-squares.

Cell differentiation and migration

432 Let N-length binary vector H_{g_i} represent the cellular neighborhood for grid-square g_i in C_t.

$$H_{g_i}(n) = 1 \text{ if } \sum_{g \in [1,2,..,G^2]} (C_t^{n,g} > 0) \bigwedge (A(g_i, g) = 1).$$

Cell migration and differentiation rules for cells in grid-square g_i can then be calculated using the Cell Fate
Rule $Mig^{H_{g_i}}(n) = m$: cells of type n migrate from grid-square g_i if $m > 0$, else not.

In order to find the new locations of cells migrating from grid-square g_i, we use matrix powers of the
adjacency matrix, A^m: The set of all grid-squares m steps away from g_i is $J_{g_i}^m = [g_j \,|\mathrm{A}^m(g_i, g_j) > 0]$. For
each cell in g_i that migrates a distance m away, we randomly choose its new grid-square from $J_{g_i}^m$. We
encode updated locations of migrating cells in μ: number of cells of cell-type n originally in grid-square g_i
which have migrated to grid-square g_j is given by μ_{g_i,g_j}^n.

Note that μ only contains the new locations of cells that migrate, i.e., for all cells of any cell-type n
such that $Mig^{H_{g_i}}(n) > 0$. Non-migrating cells are either stable in the current cellular neighborhood
H_{g_i}, or they differentiate. Cell Fate Rules governing the fate of non-migrating cell-types are given by
$Diff^{H_{g_i}}$. We encode updated cell-types for non-migrating cells in $CellDiff$: $\nu^{g_i}(n) = \sum_{n_j=1}^{N}(C_t^{n_j,g_i} >$
$0)(\delta(Diff^{H_{g_i}}(n_j), n))(\delta(Mig^{H_{g_i}}(n_j), 0))$. That is, $\nu^{g_i}(n)$ is the number of non-migrating cells of any
cell-type in g_i which differentiate into cell-type n.

The state of the tissue in the next time-step is then given by:

$C_{t+1}^{n,g_i} = \nu^{g_i}(n) + \sum_{g_j=1}^{G^2} \mu_{g_j,g_i}^n, \, \forall n \in [1, 2, ..N]$.

We synchronously update each grid-square at each model time-step, and perform 500 updates for each
tissue.

Analysing final tissues

We characterize tissue morphology based on how tissue domains (regions of uniform cellular composition
within tissues) are arranged relative to each other. In order to do this, we measure five quantities:

<u>Fullness:</u> We calculate tissue fullness as the number of grid-squares that are occupied by cells. This is then
given as $fullness = G^2 - \sum_{g=1}^{G^2} \delta(TC_{500}^g, 0)$.

<u>Number of unique domains:</u> We express the cellular composition of any occupied grid-square as a N-length
binary vector κ^g, where $\kappa^g(n) = 1$ if $C_{500}^{n,g} > 0$. κ^g can take on one of 2^N values depending on the presence
or absence of the N cell-types. We arrange these 2^N distinct cellular compositions according to the standard
binary ordering. Tissues in the model tend to be composed of contiguous domains of uniform cellular
composition, such that the cellular compositions of grid-squares within a single domain are identical to
each other. Therefore, the number of unique cellular compositions (UD) across grid-squares gives us
the number unique tissue domains (NUD). In this work, the number of cell-types $N = 5$, therefore the
maximum value of $NUD = 2^5$, and $NUD = 1$ for completely uniform tissues with a single domain.

<u>Coverage equality:</u> Let DS be an NUD length vector where $DS(i)$ reflects the size of the i^{th} domain, and
is given by the number of grid-squares with cellular composition identical to the i^{th} unique domain. We
say that a tissue has high *coverage equality* if each unique domain cellular composition is represented
by roughly equal number of grid-squares; i.e., $Eq(i) = \frac{G^2}{NUD} \forall i \in [1, 2, .., NUD]$. At the other end, for a
tissue with extremely unequal distribution of cellular compositions across the grid, $Uneq(1) = G^2$ and
$Uneq(i) = 0 \forall i in [2, 3, .., nud]$. We calculate *coverage equality* of the tissue as:

$$CovEq = 1 - \|\mathrm{DS - Eq}\|/\|\mathrm{Uneq - Eq}\|.$$

Dispersity: Any set of grid-squares of identical composition, say $UD(i)$, is either completely contiguous,
or splits into smaller subsets of contiguous regions. Contiguity of grid-squares can be assessed using
the adjacency matrix A: A subset of grid-squares is contiguous if it forms a connected component, i.e.,
any grid-square in the subset can be reached from any other grid-square in the subset by traversing the
grid. We encode the number of contiguous components with cellular compositions identical to $UD(i)$ as
$DComp(i)$. We call a tissue *disperse* if grid-squares with identical cellular compositions are split into
multiple distinct connected components. We measure the dispersity of domains with a cellular composition
$UD(i)$ as $DomDisp(i) = \frac{DComp(i)-1}{DS(i)-1}$. We measure *dispersity* of the tissue as the average $DomDisp$;
i.e., $dispersity = \frac{\sum_{i=1}^{NUD} DomDisp(i)}{NUD}$.

<u>Mixness:</u> Finally, we examine the local neighborhood of each grid-square by counting $GridMix_i$: the
number of unique cellular compositions across the set of grid-squares adjacent to g_i, including itself.
For a 2-dimensional grid, $GridMix_i$ is at most 5. We describe a G^2 length vector $SqMix$, where
$SqMix(i) = \frac{GridMix_i-1}{5}$, such that for a grid-square in the middle of a contiguous domain, $SqMix(i) = 0$.
We also assign $SqMix(i) = 0$ if grid-square g_i is unoccupied. We define tissue *mixedness* by averaging
$SqMix$ over all grid-squares.

We characterize the morphology of any model-generated tissue in the final time-point by the set of five
numbers: $fullness$, NUD, $CovEq$, $dispersity$ and $mixness$.

Principal Components and sectors

These 5 properties are not independent of each other; particularly, $mixness$ and $dispersity$ are closely
related quantities (Pearson's correlation coefficient = 0.74). To find relevant dimensions across which
properties of model-generated tissues in the final time-point ($t = 500$) vary, we perform Principal
Components Analysis. We find that the first two Principal Components (PC1 and PC2) explain 85.6% of
the variance, and we now characterize tissues based on PC1 and PC2.

In order to assess how tissue properties (PC1 and PC2) change with model parameters, and for ease of
discussion, we further split the space of tissues into *sectors*, such that tissues generated using the same
set of model parameters ($P_{den}, P_{stable}, P_{diff}, F_{adj}$) are likely to all fall within the same *sector*. Based
on our raw assessment of model-generated tissues, we choose to divide the space of tissues into *four*
sectors. Sector boundaries are straight lines: boundaries between the *stunted* and *sparse-contiguous*, and
the *sparse-contiguous* and *full-contiguous* sectors are characterized by their slopes and y-intercepts. The
boundary separating the *random-disperse* sector is vertical, and is characterized by its x-intercept. We find
optimal values for the slopes and intercepts of sector boundaries which maximize the number of parameter
sets for which at least 50% of generated tissues lie within the same sector. For the sector boundaries we
use, 50% of generated tissues lie within the same sector for 93.3% of the 135 parameter sets sampled in
this study.

Analysing tissue developmental trajectories

For each tissue, for each intermediate time-point t during its development, we calculate the five quantities
that describe its morphology: $fullness_t$, NUD_t, $CovEq_t$, $dispersity_t$ and $mixness_t$. We scale each
of these properties: For example, $scaled - fullness_t = \frac{fullness_t - MFF}{SDF}$; where MFF is the average
$fullness$ across all tissues at $t = 500$, and SDF is the standard deviation of $fullness$ across all tissues at
$t = 500$. We then project scaled values of properties of tissues at intermediate time-points onto the Principal
Components (PC1 and PC2) that describe model-generated tissues in the final time-point. This allows us

to place developing tissues at intermediate time-points into the four *sectors*: *stunted, sparse-contiguous,*
full-contiguous and random-disperse.

For each tissue T_i such that $i \in [1, 2, ..., 13500]$, let the the sector assignment at time t be $sec_t^{T_i} = x$, for
$x \in$ [*stunted, sparse-contiguous, full-contiguous, random-disperse*]. We describe tissue developmental
trajectories in terms of sequences of sector crossings: $cross^{T_i}(x, y)$ is the number of update steps during
its development in which the tissue transitions from a sector x to a sector y, where $x, y \in$ [*stunted, sparse-*
contiguous, full-contiguous, random-disperse]. For example, $cross^{T_i}(stunted, sp - contig)$ is the number
of times T_i transitions from the *stunted* to the *sparse-contiguous* sector, and $cross^{T_i}(sp - contig, sp -$
$contig)$is the number of update steps in which T_i stays within the *sparse-contiguous* sector.

We then estimate *sector transition probabilities* (STP) for each sector: For all tissues which end up in
sector $z \in$ [*stunted, sparse-contiguous, full-contiguous, random-disperse*] at $t = 500$,

$$\text{Let } T^z = [T_i : sec_{500}^{T_i} = z].$$

$$\text{Then, } STP^z(x, y) = \frac{\sum_{T_i \in T^z} cross^{T_i}(x,y)}{\sum_{T_i \in T^z} \sum_{t=1}^{500} \delta(sec_t^{T_i}, x)}.$$

Analysing tissue healing

Producing injuries

For each tissue, we test its ability to heal by simulating three kinds of injuries:

type A: For each tissue, we perform 25 independent injuries of this type. For some tissue T_i among the
13500 model-generated tissues, we set the initial state of the injured tissue as the final state of tissue T_i,
that is:

$$InjA_{t=0}^{n,g}(i) = C_{t=500}^{n,g}, \forall \text{ cell-types } n \in [1, 2, ...N] \text{ and grid-squares } g \in [1, 2, ...G^2],$$

where $i \in [1, 2, ...25]$ represents the injury-id.

Then we ablate all cells of all cell-types from a single grid-square. That is, for some randomly chosen
grid-square g_r such that $InjA_{t=0}^{all,g_r}(i) > 0$, we set $InjA_1^{n,g_r}(i) = 0 \,\forall n \in [1, 2, ...N]$.

type B: For each tissue, we perform 25 independent injuries of this type. For some tissue T_i among the
13500 model-generated tissues, we set the initial state of the injured tissue as the final state of tissue T_i,
that is:

$$InjB_{t=0}^{n,g}(i) = C_{t=500}^{n,g}, \forall \text{ cell-types } n \in [1, 2, ...N] \text{ and grid-squares } g \in [1, 2, ...G^2],$$

where $i \in [1, 2, ...25]$ represents the injury-id.

In this type of injury, we ablate all cells of one particular cell-type from all grid-squares adjacent to a
given grid-square. To do this, for some randomly chosen grid-square g_r and a cell-type n_r such that
$InjB_0^{n_r,g_r}(i) > 0$, we set $InjB_1^{n_r,g_{adj}}(i) = 0 \forall g_{adj}$ such that $A(g_r, g_{adj}) <= 1$.

type C: For each tissue, we perform 10 independent injuries of this type. For some tissue T_i among the
13500 model-generated tissues, we set the initial state of the injured tissue as the final state of tissue T_i,
that is:

$$InjC_{t=0}^{n,g}(i) = C_{t=500}^{n,g}, \forall \text{ cell-types } n \in [1, 2, ...N] \text{ and grid-squares } g \in [1, 2, ...G^2],$$

where $i \in [1, 2, ...25]$ represents the injury-id.

In this type of injury, we ablate all cells of one particular cell-type from all grid-squares ≤ 5 steps away
from a given grid-square. To do this, for some randomly chosen grid-square g_r and a cell-type n_r such that
$InjC_0^{n_r, g_r}(i) > 0$, we set $InjC_1^{n_r, g_{adj}}(i) = 0 \forall g_{adj}$ such that $A(g_r, g_{adj}) <= 5$.

Injury Magnitude

552 Finally, we calculate the magnitude of an injury as the Euclidean distance between the uninjured tissue
553 and the tissue immediately after injury: $IMX_{t=1}(i) = \|(InjX_{t=0}(i) - InjX_{t=1}(i))\|$, for injury-ids
554 $i \in [1, 2, ..., 25]$.

Tissue recovery

556 In order to simulate tissue recovery, we update the state of injured tissues for 100 time-steps using the same
557 Cell Fate Rules that were used to simulate tissue development as described above in Updating tissue states.
558 For each tissue, we test how the magnitude of the injury changes during recovery for the 25 replicates of
559 type A and type B injuries, and for the 10 replicates of type C injuries): For the i^{th} replicate injury, the
560 magnitude of the injury after 100 time-steps of recovery is given by

$$IMX_{t=100}(i) = \|(InjX_{t=0}(i) - InjX_{t=100}(i))\|, \text{ where } X \in [A, B, C]$$

We say that a tissue is from injury i of type X:

• is healing if $IMX_{t=100}(i) < IMX_{t=1}(i)$.
• shows no recovery if $IMX_{t=100}(i) = IMX_{t=1}(i)$.
• is degrading if $IMX_{t=100}(i) > IMX_{t=1}(i)$.

For each tissue, for each injury type, we measure the fraction of replicate injuries out of 25 (10 for type C)
that result in healing ($RecFrac$), no recovery ($NoRecFrac$), or tissue degradation ($DegFrac$).

ACKNOWLEDGEMENTS

568 This work was supported by the taxpayers of South Korea through the Institute for Basic Science, Project
569 Code IBS-R020-D1. We thank all members of the Living Matter Theory group at IBS-CSLM for engaging
570 discussions during the conception of this work. We thank John McBride for constructive comments during
571 the writing of this manuscript.

SUPPLEMENTARY INFORMATION

The supplementary file contains the following figures:

• FigS1: Histogram for the number of time-steps to tissue steady-state.
• FigS2: Scatter-plots showing the dependence on model parameters of number of distinct tissue cell-types,
average cell-fate potential.
• FigS3: Scatter-plots showing the dependence on model parameters of maximum cell-fate potential of
tissue cell-types.
• FigS4: Heatmaps displaying dependence on model parameters of the 5 spatial properties using which we
characterize model-generated tissues.

• FigS5: Elbow plot showing the variance in the data explained by each PC
• FigS6: Scatter-plots showing the dependence on model parameters of Principal Components which
represent tissue spatial properties.
• FigS7: Hierarchical clustering of model-generated tissues.
• FigS8: Histograms for the number of sector crossings, number of time-steps to reach the final sector.
• FigS9: Scatterplots displaying the extent of tissue healing for tissues in different sectors in response to
type B injuries.
• FigS10: Heatmaps displaying dependence of tissue healing in response to type B injuries on model
parameters.

REFERENCES

Adler, M., Moriel, N., Goeva, A., Avraham-Davidi, I., Mages, S., Adams, T. S., et al. (2023). Emergence
of division of labor in tissues through cell interactions and spatial cues. *Cell Reports* 42

Armenta-Medina, A., Gillmor, C. S., Gao, P., Mora-Macias, J., Kochian, L. V., Xiang, D., et al. (2021).
Developmental and genomic architecture of plant embryogenesis: from model plant to crops. *Plant*
Communications 2

Armingol, E., Baghdassarian, H. M., and Lewis, N. E. (2024). The diversification of methods for studying
cell–cell interactions and communication. *Nature Reviews Genetics* , 1–20

Birnbaum, K. D. and Alvarado, A. S. (2008). Slicing across kingdoms: regeneration in plants and animals.
Cell 132, 697–710

Bock, M., Tyagi, A. K., Kreft, J.-U., and Alt, W. (2010). Generalized voronoi tessellation as a model of
two-dimensional cell tissue dynamics. *Bulletin of mathematical biology* 72, 1696–1731

Borsuk, A. M., Roddy, A. B., Théroux-Rancourt, G., and Brodersen, C. R. (2022). Structural organization
of the spongy mesophyll. *New Phytologist* 234, 946–960

Brennan, F. H., Li, Y., Wang, C., Ma, A., Guo, Q., Li, Y., et al. (2022). Microglia coordinate cellular
interactions during spinal cord repair in mice. *Nature communications* 13, 4096

Buckley, C. E. and St Johnston, D. (2022). Apical–basal polarity and the control of epithelial form and
function. *Nature reviews Molecular cell biology* 23, 559–577

Byrd, D. T. and Kimble, J. (2009). Scratching the niche that controls caenorhabditis elegans germline stem
cells. In *Seminars in cell & developmental biology* (Elsevier), vol. 20, 1107–1113

Chan, C. J., Heisenberg, C.-P., and Hiiragi, T. (2017). Coordination of morphogenesis and cell-fate
specification in development. *Current Biology* 27, R1024–R1035

Clevers, H. (2013). The intestinal crypt, a prototype stem cell compartment. *Cell* 154, 274–284

Dang, Y., Lattner, J., Lahola-Chomiak, A., Alves-Afonso, D., Ulbricht, E., Taubenberger, A. V., et al.
(2023). Self-propagating wave drives noncanonical antidurotaxis of skull bones in vivo. *bioRxiv* ,
2023–07

Fan, Y. and Meyer, T. (2021). Molecular control of cell density-mediated exit to quiescence. *Cell reports*
36

Farhadifar, R., Röper, J.-C., Aigouy, B., Eaton, S., and Jülicher, F. (2007). The influence of cell mechanics,
cell-cell interactions, and proliferation on epithelial packing. *Current biology* 17, 2095–2104

Feng, Y., Yang, T., Zhu, J., Li, M., Doyle, M., Ozcoban, V., et al. (2023). Spatial analysis with spiat and
spasim to characterize and simulate tissue microenvironments. *Nature Communications* 14, 2697

Fuchs, E. and Blau, H. M. (2020). Tissue stem cells: architects of their niches. *Cell stem cell* 27, 532–556

Gaillochet, C., Daum, G., and Lohmann, J. U. (2015). O cell, where art thou? the mechanisms of shoot
meristem patterning. *Current opinion in plant biology* 23, 91–97

Gillies, T. E. and Cabernard, C. (2011). Cell division orientation in animals. *Current biology* 21,
R599–R609

Hatton, I. A., Galbraith, E. D., Merleau, N. S., Miettinen, T. P., Smith, B. M., and Shander, J. A. (2023).
The human cell count and size distribution. *Proceedings of the National Academy of Sciences* 120,
e2303077120

Holstein, T. W. (2023). The hydra stem cell system–revisited. *Cells & Development* 174, 203846

Hu, Y., Omary, M., Hu, Y., Doron, O., Hoermayer, L., Chen, Q., et al. (2021). Cell kinetics of auxin
transport and activity in arabidopsis root growth and skewing. *Nature Communications* 12, 1657

Koike, H., Manabe, I., and Oishi, Y. (2022). Mechanisms of cooperative cell-cell interactions in skeletal
muscle regeneration. *Inflammation and Regeneration* 42, 48

Lee, Y., Kouvroukoglou, S., McIntire, L. V., and Zygourakis, K. (1995). A cellular automaton model for
the proliferation of migrating contact-inhibited cells. *Biophysical journal* 69, 1284–1298

Marée, A. F., Grieneisen, V. A., and Hogeweg, P. (2007). The cellular potts model and biophysical
properties of cells, tissues and morphogenesis. *Single-cell-based models in biology and medicine* ,
107–136

Marhava, P., Hoermayer, L., Yoshida, S., Marhavỳ, P., Benková, E., and Friml, J. (2019). Re-activation of
stem cell pathways for pattern restoration in plant wound healing. *Cell* 177, 957–969

McKinley, K. L., Stuurman, N., Royer, L. A., Schartner, C., Castillo-Azofeifa, D., Delling, M., et al.
(2018). Cellular aspect ratio and cell division mechanics underlie the patterning of cell progeny in
diverse mammalian epithelia. *Elife* 7, e36739

Mukund, K. and Subramaniam, S. (2020). Skeletal muscle: A review of molecular structure and function,
in health and disease. *Wiley Interdisciplinary Reviews: Systems Biology and Medicine* 12, e1462

Nagahata, Y., Masuda, K., Nishimura, Y., Ikawa, T., Kawaoka, S., Kitawaki, T., et al. (2022). Tracing
the evolutionary history of blood cells to the unicellular ancestor of animals. *Blood, The Journal of the*
American Society of Hematology 140, 2611–2625

Nagy, L. G., Vonk, P. J., Künzler, M., Földi, C., Virágh, M., Ohm, R. A., et al. (2023). Lessons on fruiting
body morphogenesis from genomes and transcriptomes of agaricomycetes. *Studies in Mycology* 104,
1–85

Okubo, T., Rivron, N., Kabata, M., Masaki, H., Kishimoto, K., Semi, K., et al. (2023). Hypoblast from
human pluripotent stem cells regulates epiblast development. *Nature* , 1–3

Oliva, M. and Lister, R. (2023). Exploring the identity of individual plant cells in space and time. *New*
Phytologist 240, 61–67

Osborne, J. M., Fletcher, A. G., Pitt-Francis, J. M., Maini, P. K., and Gavaghan, D. J. (2017).
Comparing individual-based approaches to modelling the self-organization of multicellular tissues.
PLoS computational biology 13, e1005387

O'Connor, C. M., Adams, J. U., and Fairman, J. (2010). Essentials of cell biology. *Cambridge, MA: NPG*
Education 1, 54

Pavel, M., Renna, M., Park, S. J., Menzies, F. M., Ricketts, T., Füllgrabe, J., et al. (2018). Contact inhibition
controls cell survival and proliferation via yap/taz-autophagy axis. *Nature communications* 9, 2961

Pavlicev, M., DiFrisco, J., Love, A. C., and Wagner, G. P. (2024). Metabolic complementation between
cells drives the evolution of tissues and organs. *Biology letters* 20

Pavlicev, M. and Wagner, G. P. (2023). Reading the palimpsest of cell interactions: what questions may we
ask of the data? *iScience*

Perrimon, N., Pitsouli, C., and Shilo, B.-Z. (2012). Signaling mechanisms controlling cell fate and
embryonic patterning. *Cold Spring Harbor perspectives in biology* 4, a005975

Ramos, R., Swedlund, B., Ganesan, A. K., Morsut, L., Maini, P. K., Monuki, E. S., et al. (2024). Parsing patterns: Emerging roles of tissue self-organization in health and disease. *Cell* 187, 3165–3186

Reddien, P. W. (2018). The cellular and molecular basis for planarian regeneration. *Cell* 175, 327–345

Romanova-Michaelides, M., Hadjivasiliou, Z., Aguilar-Hidalgo, D., Basagiannis, D., Seum, C., Dubois, M., et al. (2022). Morphogen gradient scaling by recycling of intracellular dpp. *Nature* 602, 287–293

Sandersius, S. A. and Newman, T. J. (2008). Modeling cell rheology with the subcellular element model. *Physical biology* 5, 015002

Schittny, J. C. (2017). Development of the lung. *Cell and tissue research* 367, 427–444

Subkhankulova, T., Camargo Sosa, K., Uroshlev, L. A., Nikaido, M., Shriever, N., Kasianov, A. S., et al. (2023). Zebrafish pigment cells develop directly from persistent highly multipotent progenitors. *Nature Communications* 14, 1258

Sugimoto, K., Temman, H., Kadokura, S., and Matsunaga, S. (2019). To regenerate or not to regenerate: factors that drive plant regeneration. *Current opinion in plant biology* 47, 138–150

Sunchu, B. and Cabernard, C. (2020). Principles and mechanisms of asymmetric cell division. *Development* 147, dev167650

Technau, U. and Steele, R. E. (2011). Evolutionary crossroads in developmental biology: Cnidaria. *Development* 138, 1447–1458

Theodorou, I. and Charrier, B. (2021). Brown algae: Ectocarpus and saccharina as experimental models for developmental biology. In *Handbook of Marine Model Organisms in Experimental Biology* (CRC Press). 27–47

Thowfeequ, S. and Srinivas, S. (2022). Embryonic and extraembryonic tissues during mammalian development: shifting boundaries in time and space. *Philosophical Transactions of the Royal Society B* 377, 20210255

Wei, Y., Wang, Y. G., Jia, Y., Li, L., Yoon, J., Zhang, S., et al. (2021). Liver homeostasis is maintained by midlobular zone 2 hepatocytes. *Science* 371, eabb1625

Yao, Y. and Wang, C. (2020). Dedifferentiation: inspiration for devising engineering strategies for regenerative medicine. *NPJ Regenerative medicine* 5, 14

Zuch, D. T., Doyle, S. M., Majda, M., Smith, R. S., Robert, S., and Torii, K. U. (2022). Cell biology of the leaf epidermis: Fate specification, morphogenesis, and coordination. *The Plant Cell* 34, 209–227