Functional maps of a genomic locus reveal confinement of an enhancer by its target gene

Mathias Eder[1,2,3]†, Christina J.I. Moene[1,2,3]†, Lise Dauban[1,2,3], Christ Leemans[1,3], Bas van Steensel[1,2,3]*

Divisions of [1]Gene Regulation and [2]Molecular Genetics, Netherlands Cancer Institute, Amsterdam, the Netherlands.
[3]Oncode Institute, the Netherlands

† These authors contributed equally to this work
* Corresponding author. Email: b.v.steensel@nki.nl

Abstract

Genes are often activated by enhancers located at large genomic distances. The importance of this positioning is poorly understood. By relocating promoter-reporter constructs into >1,000 alternative positions within a single locus, we dissected the positional relationship between the mouse *Sox2* gene and its distal enhancer. This revealed an intricate, sharply confined activation landscape, in which the native *Sox2* gene occupies an optimal position for its activation. Deletion of the gene relaxes this confinement and broadly increases reporter activity. Surprisingly, the confining effect of the *Sox2* gene is partially conferred by its ~1 kb coding region. Our local relocation approach provides high-resolution functional maps of a genomic locus and reveals that a gene can strongly constrain the realm of influence of its enhancer.

MAIN TEXT

Mammalian enhancers can activate genes across large genomic distances, up to hundreds of kilobases (*1, 2*), presumably by contacting their target promoter through folding of the chromatin fiber. Indeed, enhancers tend to have increased contact with the promoters they regulate (*3*), but it is unclear whether this is a cause or effect of their function. In addition, enhancer – promoter pairs often reside within the same topologically associated domain (TAD), a region of preferential self-interaction in the genome that is often demarcated by CTCF binding sites (*4, 5*). These TADs are thought to form functional domains (*6, 7*), but it is largely unclear whether enhancers can activate promoters anywhere within a TAD, or whether some positions within a TAD are more favorable than others. Recent studies in two loci with little three-dimensional (3D) structure indicated that promoter activity increases with decreasing genomic distance to its enhancer (*8, 9*). However, in more structured genomic loci the effect of position on enhancer – promoter communication has remained elusive. Here, we addressed these questions by constructing high-resolution maps of promoter activity as function of position in a native locus with pronounced 3D folding.

High throughput relocation by Sleeping Beauty transposition

To unravel the logic that underlies the linear arrangement of regulatory elements, it is necessary to alter the genomic position of such elements systematically and probe the functional consequences. Here we provide such a systematic transplantation of regulatory elements by adapting a methodology based on the cut-and-paste mechanism of transposable elements (*9-11*). When mobilized from the genome by its transposase, the Sleeping Beauty (SB) transposon excises and re-integrates, usually within ~1-2 Mb from the original location (**Fig. 1A**) (*10, 12*). Thus, any sequence inserted between the two SB inverted terminal repeats (ITRs) can be randomly relocated ("hopped") throughout a genomic locus. We developed a workflow to generate and map thousands of integrations in a single locus, and link the locations to the expression of a transcribed reporter, yielding high-resolution functional maps of the local regulatory landscape.

For this study we chose the *Sox2* locus in mouse embryonic stem cells (mESCs). In these cells, *Sox2* expression is primarily determined by the *Sox2* control region (SCR), which is a cluster of enhancers located 110 kb downstream of the gene in the same TAD (*13-17*). Within this TAD, the *Sox2* gene has been found to preferentially contact the SCR (*18*) (**Fig. 1B**). The 1.5 Mb region around *Sox2* contains no other protein-coding genes (*13*). To ease genetic engineering, we used F1-hybrid (*129/Sv:CAST/EiJ*) mESCs, in which the *CAST* and the *129S1* alleles can be distinguished by a high density of sequence polymorphisms. We tagged both alleles of *Sox2* with either *eGFP* (*CAST*) or *mCherry* (*129S1*) to be able to quantify allelic *Sox2* expression by flow cytometry (**Fig. S1A**). To confirm the regulation by the SCR, we transfected the

dual-tagged cell line with Cas9 and two gRNAs that cause deletion of the SCR. In line with the reported

key role of the SCR (*13, 14, 16*), a proportion of cells lost more than 90% of *eGFP* or *mCherry* expression,

most likely corresponding to a deletion of the SCR on the corresponding allele (**Fig. S1B**).

Next, we used Cas9 editing to create a "launch pad" by integrating a SB transposon cassette 6

kb upstream of the *Sox2::mCherry* allele (-116 kb relative to the SCR, **Fig. 1B**). The cassette contained a

double selection marker (mPGK-HyTK) flanked by a pair of heterotypic Flp-recombinase recognition sites

(FRT/F3), enabling us to easily change the DNA sequence within the transposon by recombination-

mediated cassette exchange (RMCE) (**Fig. S1C**).

In a proof-of-principle experiment, we induced hopping of the SB cassette containing an

arbitrary 282 bp sequence (see methods). After culturing the cells for one week to dilute the transposase-

containing plasmid and prevent further re-hopping events, we isolated and expanded 200,000 cells and

mapped the integrations using a Tn5-based sequencing approach (*19*). In some of the cells, the transposon

was never mobilized, which is reflected by 33% of the mapping reads originating from the launch site. Of

all the hopped integrations, 72% were located on the same chromosome as the *Sox2* locus (chr3) (**Fig.**

S1D). Allele-specific mapping indicated that most integrations happened in *cis* (**Fig. S1E**). Within -/+ 2 Mb

of the launch pad we mapped 2699 unique integrations (**Fig. 1C**), which were equally distributed between

the forward (1342) and reverse (1357) orientations (**Fig. S1F**). We note that this mapping is not exhaustive

and the actual pool likely contains more integrations, since more than 70% of the integrations were only

supported by one unique mapping read (**Fig S1G**). This experiment demonstrates the effectiveness of SB

hopping to generate thousands of genomic integrations.

Eder, Moene et al. Fig 1

Fig. 1 Sleeping Beauty mobilization creates integrations covering the *Sox2* locus.

(A) Cut-and-paste local relocation of a Sleeping Beauty (SB) transposon by transient expression of SB transposase. **(B)** Overview of the *Sox2* locus. Top to bottom: contact map in mESCs (region-capture micro-C data [(*18*)], 1000-bp resolution) and insulation scores based on the same data; positions and orientations of CTCF binding sites (triangles); and positions of SCR and eGFP/mCherry tagged *Sox2* alleles, the SCR, and a launch pad (129S1 allele) containing SB with a recombination cassette. **(C)** Distribution of hopped SB carrying a 282 bp inert DNA sequence, mobilized from the launch pad; n, total number of integrations mapped in the plotted region.

A highly detailed activation landscape of the *Sox2* locus

Next, we set out to generate a detailed map of the expression potential of the *Sox2* locus, *i.e.*, to determine quantitatively how the location throughout the locus affects the ability of the *Sox2* promoter to be activated. For this we engineered a reporter comprising 1.9 kb of the *Sox2* promoter (excluding a promoter-proximal CTCF binding site) driving the expression of mTurquoise2 (*mTurq*), a blue fluorescent protein (**Fig. 2A**). We integrated this reporter into the launch pad using RMCE. This resulted in low but detectable reporter expression, without affecting endogenous *Sox2::mCherry* expression (**Fig. S2A**, reporter -116 kb).

We then induced SB hopping and monitored reporter fluorescence by flow cytometry. Compared to the non-hopped control, the resulting cell pool showed a broadened distribution of reporter fluorescence, indicating that the relocation at least in part affected reporter expression (**Fig. 2B**). We then split the full range of reporter expression values into six gates (P1-P6) and sorted pools of cells from each gate (**Fig. 2B**), and subsequently mapped the locations of the reporters in each pool. As a control, we also mapped the locations in unsorted cells. Since the integration patterns were reproducible between biological replicates (**Fig. S2B-C**), we combined the replicates for further downstream analysis.

As we observed previously with the arbitrary insert sequence (**Fig. 1C**), reporter integrations in the control pool are mostly concentrated in a 2 Mb region surrounding the launch site (**Fig. 2C**, top track). In contrast, the cell pools with different expression levels yielded clearly distinct distributions (**Fig. 2C-D**, P1 – P6 tracks). Reporters with the highest activity (P1) were concentrated almost exclusively in the center of the SCR or close to the endogenous *Sox2* gene, while the second-most active reporters (P2) were more dispersed around these regions. Intermediate activity (P3 – P5) occurred predominantly near the launch site and between *Sox2* and SCR. Finally, silent (P6) reporters were almost exclusively located outside the *Sox2*-SCR TAD.

We converted these raw data into a reporter expression score, which is the average expression level estimated in a sliding window across the locus (**Fig. 2E**, see Methods). This graph highlights that the SCR and the region immediately surrounding the endogenous *Sox2* gene are the most optimal locations for transcriptional activity. Precise fluorescence measurements in a panel of clonal cell lines with 36 unique integrations confirms this peak in activity at the center of the SCR and validates the pattern of expression scores inferred from the sorted pools (**Fig. S2D**). Reporter integrations in the center of the SCR display a 10-fold upregulation in reporter expression compared to the -116 kb launch pad position. A remarkable lack of reporter expression occurs outside of the *Sox2*-SCR TAD, with a very sharp drop-off (within about 10 kb) on either side (**Fig. 2E**). A plateau of intermediate expression is found between *Sox2* and SCR. Together, this detailed landscape points to a strong confinement of the realm over which the SCR is able to activate transcription.

The relatively large confidence intervals of the expression scores limited interpretation of the fine-
scale pattern between *Sox2* and SCR (**Fig. 2E**). This is due to a relative paucity of reporter integrations in
this range. The distribution of reporter expression in cells where the reporter was not mobilized, overlaps
the medium-level expression gates (P4 – P5, **Fig. 2C**), leading to a low number of sorted integrations for
those gates. To overcome this, we repeated the hopping experiments starting from a launch pad 51 kb
upstream of *Sox2::mCherry* (-161 kb relative to the SCR) (**Fig. 2F**). At this position the reporter showed no
measurable expression (**Fig. S2A**), and hence from this launch pad it should be easier to detect hopping
events in the medium-level expression bins. Indeed, this was the case (**Fig. 2G-H**). The resulting expression
score map (**Fig. 2I**) has narrow confidence intervals within the *Sox2*-SCR TAD (at a resolution of 5 kb), at
the cost of a paucity of data outside this range.

This high-resolution map confirmed that the highest reporter expression occurs at the SCR and
around the endogenous *Sox2* gene (**Fig. 2I**). Reporter activity throughout the region is independent of the
orientation of the reporter (**Fig. S2E**). Within the *Sox2*-SCR TAD, the map reveals an intricate activity
landscape with several local peaks of reporter expression, two of which overlap CTCF binding sites (**Fig.
2I**, black arrows). Strikingly, this landscape closely matches a map of SCR contact frequencies that was
obtained in the absence of reporter integrations (*18*) (**Fig. 2J**, top panel; **Fig. 2K**). Maps of open chromatin
(ATAC-seq (*20*)) and the histone mark H3K27ac (*21*), both thought to be features of regulatory elements
(*22*), showed only partial overlap with the fine pattern of reporter activity (**Fig. 2J**, bottom panel). These
results indicate that reporter expression primarily follows contact frequency with the SCR, suggesting that
these contacts are a major determinant of quantitative expression levels.

Fig. 2 A highly detailed activation landscape of the Sox2 locus. (A) Left panel: design of mTurquoise2 reporter driven by *Sox2* promoter and 5'UTR, followed by SV40 polyadenylation signal (PAS). Right panel: this reporter was inserted in the -116 kb launch pad (LP), upstream of the mCherry-tagged endogenous *Sox2* allele. **(B)** Expression levels of reporter (mTurq) and *Sox2::eGFP* (control) in mock-transfected cells and in cells transfected with SB transposase, measured by FACS. Boxes mark sorting gates used to obtain cell populations P1-P6 in **C-D**. The percentage of cells in each gate is indicated. Representative result of three biological replicates. **(C)** Mapped integrations from a large pool of unsorted cells (ctrl) and the six sorted populations (P1-P6), mobilized from the -116 kb launch pad. Each dot indicates one mapped integration, number of plotted integrations in each population is indicated on the right. Combined data from three biological replicates. Grey shading highlights populations that are most different from the starting expression of the unhopped reporter. LP, launch pad position. **(D)** Zoom-in of C. Grey triangles indicate CTCF binding sites (CBS) and their orientation. **(E)** Expression score derived from data shown in **D**, smoothened using a 10 kb running window shifted in 1 kb steps and only plotted when the window contains 3 or more SB integrations. Shaded region indicates 95% confidence interval. **(F-I)** Same as **A-B, D-E**, except that the reporter was mobilized from a launch pad at position -161 kb. Data are from 2 biological replicates combined. Expression scores in **I** are calculated using a 5 kb running window shifted in 500 bp steps. Arrows mark two peaks near CBSs. **(J)** Annotation of the locus. Top: virtual 4C contact profile (from RCMC data, (*18*)) with the center of the SCR (indicated in lightgrey) as viewpoint, at 1 kb resolution. Dashed lines indicate CBSs and grey triangles indicate their orientation. Bottom: ATAC-seq signal (*20*), H3K27ac ChIP-seq p-value signal (*21*) and CTCF ChIP-seq coverage (own data), smoothened using a 500bp running window. **(K)** Correlation between expression score and virtual 4C contact with the core SCR, for the datasets from -161 kb launch pad, per 1 kb bin on the genomic region plotted in **H-J**. Only bins with 3 or more SB integrations are included. Rho is Spearman's correlation coefficient.

Random deletion of the endogenous *Sox2* gene boosts reporter expression

Close examination of the flow cytometry data of the transposase-transfected reporter cells (-116 kb launch pad) revealed a small but distinct population of cells with high reporter expression but no *Sox2::mCherry* expression (**Fig. 3A**). We isolated a panel of clones with these unusual characteristics and mapped the SB location. Surprisingly, in these clones the left ITRs were still located at the launch site (upstream of *Sox2::mCherry*), while the right ITRs were linked to sequences distributed throughout the *Sox2* locus (**Fig. 3B, S3A**). This suggests that the genomic sequence between the launch site and integration site has been lost. Sanger sequencing confirmed the loss of the 129S1 allele in this region (**Fig. S3B**). We hypothesize that in rare instances, errors in the SB excision-reinsertion process can cause such a deletion, which can lead to the loss of the *Sox2::mCherry* gene.

Strikingly, clones with such hopping-induced deletions exhibited very high reporter expression, up to 49-fold higher than the -116 kb reporter clone (**Fig. 3C**). Reporters close to the SCR (within 20 kb) were expressed about 5-fold higher when a deletion was present, compared to regular insertions in the same area. Even the smallest deletions of 8.5 kb, removing only the endogenous *Sox2::mCherry* gene, resulted in dramatically increased reporter expression. These data suggest that the endogenous *Sox2* gene

strongly limits reporter expression when present. Other sequences within the *Sox2*-SCR range may further

modulate this effect, indicated by the variable increases in expression across the deletions, which show

no correlation with distance to the SCR (Spearman's rho = 0.19, p = 0.19). In summary, these fortuitous

rare deletions suggested that the endogenous *Sox2* gene strongly reduces activity of the reporter gene

throughout the *Sox2*-SCR range.

***Fig. 3* Hopping-induced deletions of the endogenous Sox2 gene increase reporter expression. (A)** Expression of

Sox2::mCherry and *Sox2* reporter (mTurq) after mobilization of SB from the -116 kb launch pad, first biological

replicate (also shown in **Fig. 2B**). Box indicates rare *Sox2::mCherry*-negative cells with high reporter expression. **(B)**

Unique deletions inferred from mapping in *Sox2::mCherry*-negative sorted clones (**Fig. S3A**), sorted by deletion size.

Horizontal dashed lines indicate individual deletions. Dots indicate the locations of the re-insertions. Almost all

deletions start at the launch pad position. **(C)** Relative reporter expression of clones with a hopping-induced deletion

(blue dots, location corresponds to the end of the deletion) or regular insertion (grey, same data as in **Fig. S2D**,

bottom).

***Sox2* gene strongly controls the regulatory landscape of the enhancer**

To verify this interpretation, we used CRISPR/Cas9 to delete the entire *Sox2* gene, including its

promoter. Furthermore, to test whether the CTCF binding site (CBS) just upstream of the *Sox2* promoter

(which was also lost in the hopping-induced deletions) might play a role, we deleted either the *Sox2*

promoter and gene body (Δ*Sox2*), or the same region including this CBS (ΔCBS-*Sox2*). The guide RNAs used

for these deletions were not allele-specific, and thus could independently delete either the *Sox2::eGFP* or

Sox2::mCherry allele (homozygous *Sox2* deletion causes mESC differentiation (*23*)). By FACS analysis we

identified cells with either heterozygous deletion and measured the respective impacts on mTurq reporter

activity (**Fig. 4A)**. We performed these experiments in three cell lines, with the reporter located at

positions -161 kb (no detectable activity), -116 kb (low activity), or -39 kb (between *Sox2* and SCR, medium

activity) (**Fig. S4A**) relative to the SCR.

Compared to cells that retained both *Sox2* alleles, cells which only lost *Sox2::mCherry* showed
strong upregulation of the reporter (on the same allele) in all three locations (**Fig. 4B**). This upregulation
did not occur in cells lacking the *Sox2::eGFP* allele, indicating that the effect is strictly *in cis*. The increase
in reporter expression was similar between the Δ*Sox2* and ΔCBS-*Sox2*. The upregulation of the reporter
at position -161 kb is remarkable, as it indicates that the endogenous *Sox2* gene normally prevents the
SCR from reaching sequences upstream, outside of the TAD. For reporters in the other two positions only
a sub-population of cells showed pronounced upregulation (**Fig. 4B**). The basis of this bimodality is
unclear.

To gain better understanding of the relationship between reporter location and activation, we
sorted out pools and clones of *Sox2::mCherry* negative cells (*cis*-deletion) with high reporter activity. After
cell expansion, the reporters at -161 kb and -39 kb displayed stable and uniform reporter expression (**Fig.**
S4B-D). In contrast, the -116 kb reporters again exhibited a bimodal expression pattern: a fraction of cells
lost reporter expression after sorting the pools (**Fig. S4B**) and clones (**S4E-F**). Comparison of the highest
reporter expression peaks of the sorted clones shows that the highest expression was not reached by the
reporter closest to the SCR (-39 kb), but rather by the reporter closest to the native location of *Sox2* (-116
kb) (**Fig. 4C**). Overall, these results underscore that the region around the endogenous *Sox2* gene is
optimal for high expression, even when this gene including its promoter is deleted.

To further explore the impact of the native *Sox2* gene on the regulatory landscape throughout the
locus, we repeated the reporter hopping experiment from the -161 kb launch site in a ΔCBS-*Sox2::mCherry*
clone (**Fig. S4C**). Since the founder cell line has intermediate reporter expression (**Fig. S4G**), hopped
integrations were expected to be most enriched in the highest (P1) and lowest (P6) expression gates. In
the presence of the endogenous *Sox2::mCherry* (*Sox2*$^{+/+}$)*,* P1-level integrations were rare and only found
at the SCR and endogenous gene (**Fig. 4D**, bottom). In contrast, when the endogenous *Sox2* gene is
deleted, such highly active reporter integrations are found throughout the *Sox2*-SCR range and even
extend beyond the cluster of CTCF binding sites ~7 kb upstream of the deleted *Sox2* promoter. Most silent
(P6) reporters are only found at least 55 kb upstream of the deleted gene and 38 kb downstream of the
SCR, indicating that the region permissive for activity expanded by roughly 50%, extending beyond the
original TAD-boundaries (**Fig. 4D**, top). Together, these data demonstrate that the *Sox2* gene strongly
constrains both the level of activation of a competing promoter and the realm of influence of the SCR.

Fig. 4 Sox2 gene strongly confines the activation realm of the SCR. (A) Left panel: Cas9-mediated deletion of tagged *Sox2* alleles, with or without CBS (grey triangle); right panel: gating and analysis strategy. **(B)** Distribution of mTurq reporter expression levels in three cell lines with reporter inserted in positions -161 kb, -116 kb, and -39 kb, each treated and analyzed as in **A**. Numbers indicate the number of cells measured. **(C)** Maximum relative reporter expression of clonal cell lines after *Sox2::mCherry* or CBS_*Sox2::mCherry* deletion, at reporter locations as indicated. The maximum relative reporter expression is the maximum expression peak indicated in **Fig. S4C-F** (vertical lines), which is then autofluorescence subtracted, normalized to *Sox2::GFP*, and then normalized to the median of the standard -116 kb reporter cell line measured on the same day. Dashed line indicates 1 (standard cell line). Each dot represents one clone (CRISPR edited clones) or one measurement day (untreated clones). **(D)** Top: hopping of reporter (Sox2P) from the -161 kb launch pad in a ΔCBS_*Sox2::mCherry* cell line. Mapped integrations from a large pool of unsorted cells (ctrl) and the six sorted populations P1-P6 (same as in **Fig. 2B**). Data are from 2 biological replicates combined. Grey shading indicates the populations with reporter expression levels that are most different from the starting expression of the unhopped reporter. Bottom: the P1 and P6 sorted integrations combined from hopping in the original -161 kb (2 replicates) and -116 kb (3 replicates) reporter cell lines with *Sox2::mCherry* intact (same data as in **Fig 2D** and **2H**). Grey triangles indicate CBSs and their orientation.

234 **Unbalanced competition is partially driven by the *Sox2* gene body**

235   While the endogenous *Sox2* gene significantly restricts reporter expression, we found that the
236 reciprocal impact of the reporter on the *Sox2* gene is very minor. First, the reporter inserted in the launch
237 site at -116 kb did not detectably affect *Sox2::mCherry* expression (**Fig. S2A**). Second, among all clonal
238 lines with a hopped reporter, *Sox2::mCherry* expression is remarkably robust. Even in clones with a
239 reporter in the center of the SCR (with ~10-fold higher expression than at -116 kb) *Sox2::mCherry*
240 expression is reduced by only ~25% (**Fig. 5A**). This is extremely modest compared to the 3-30 fold
241 upregulation of the reporter upon loss of *Sox2::mCherry* (**Fig. 4C**). Hence, the endogenous gene affects
242 the reporter considerably more than *vice versa*.

243   We wondered what might give the endogenous *Sox2* gene this competitive edge. The reporter
244 contained the exact promoter and 5' UTR sequence from the *Sox2* gene, but it lacked the 1.0 kb coding
245 sequence (CDS) and the 1.1 kb 3' UTR and polyadenylation signal (PAS) (*Sox2* has no introns). Re-analysis
246 of high-resolution region capture micro-C data (*18*) indicates that the SCR not only contacts the promoter
247 of the endogenous *Sox2* gene, but also the gene body (**Fig. S5A**), indicating that the latter sequence might
248 play a role. We therefore generated two new reporters in which we added the *Sox2* CDS alone or in
249 combination with the 3' UTR/PAS (**Fig. 5B**). We kept the *mTurquoise2* sequence separated from the CDS
250 by a P2A cleavage signal (*24*), so the stability of the fluorescent protein would not be affected by a fusion
251 with the SOX2 protein.

252   We first tested these new reporters in positions -161 kb and -116 kb after deletion of
253 *Sox2::mCherry*. Strikingly, across multiple independent clonal cell lines, the CDS and CDS/3'UTR additions
254 boosted reporter activity in the -161 kb location by 6-8 fold (**Fig. 5C, Fig. S5B**). Reporter insertion in the -
255 116 kb location, just 6 kb upstream of the deleted *Sox2* gene, is a close approximation of replacing the
256 gene with a reporter. Here, the promoter-only reporters showed very high but bimodal expression (**Fig**
257 **5C, S5B, S5C**), as observed before (**Fig S4B, E, F**). Surprisingly, the CDS-containing reporters had lower but
258 stable expression (**Fig. 5C, S5C**). We speculate that this lower expression is the result of a negative
259 feedback mechanism on high SOX2 protein levels, which can be triggered by the CDS-containing reporter
260 but not the original *Sox2P* reporter (*25, 26*). Overall, these results suggest that the regulation of the CDS-
261 containing reporter more closely resembles the native *Sox2* gene.

262   Next, we tested the effect of the CDS when the competing *Sox2::mCherry* gene is also present in
263 the locus. In both the -116 kb and -39 kb position, clonal cell lines with the CDS-containing reporters had
264 3-7 fold higher expression than those with the original reporter (**Fig. 5D** top, **Fig. S5D**). We then asked
265 whether the presence of the CDS also shifts the competitive balance with the endogenous *Sox2* gene.
266 Indeed, in these cells the addition of the CDS to the reporter caused a 13-18% reduction in *Sox2::mCherry*
267 expression, while additional inclusion of the 3' UTR in the reporter even caused a 21-31% reduction. (**Fig.**
268 **5D** bottom, **Fig. S5D**).

Fig. 5 Unbalanced competition is partially driven by Sox2 gene body. (A) Relative reporter (mTurq) and *Sox2::mCherry* expression in clones around *Sox2* and the SCR (median expression, estimated autofluorescence subtracted, normalized to GFP, and then normalized to the unhopped reporters at -116 kb). Loess curve shows the trend in expression. Same clones and mTurq data as in **Fig. S2D**, bottom. **(B)** Design of original *Sox2* promoter reporter (Sox2P) and two coding sequence (CDS)-containing reporters. PAS, poly-adenylation signal. **(C)** Relative expression levels of the three reporters at -161 kb and -116 kb, in a ΔCBS_*Sox2::mCherry* cell line. Each dot represents an individual clone. Relative expression was calculated as in **A**. Dashed line indicates 1 (control cell line). **Fig. S5B** shows the same expression values, not normalized to *eGFP*. P-values according to Welch's unequal variances *t*-test.

(D) As C, but for the relative reporter (top) and endogenous *Sox2*::*mCherry* (bottom) expression of the three reporters and a minimal insert (not containing any promoter or mTurq) inserted at -116 kb and -39 kb launch pads with *Sox2*::mCherry intact. **Fig. S5D** shows the same expression values, not normalized to *eGFP*. **(E)** Left panels: schematic of hopping (from launch pad -161 kb) and sorting Sox2P and Sox2P_CDS reporters based on reporter expression (P1-P6). Right panel: correlation between relative *Sox2*::*mCherry* and reporter (mTurq) expression from the Sox2P and Sox2P_CDS reporter, from cells with reporter expression in the gates P2-P4 (see **Fig. 2G and S5H**). Relative expression was calculated as in **A**, but per individual cell. Rho, Spearman's correlation coefficient. Data are from two biological replicates combined. For visualization, the y-axis is cut at 1.5, excluding 8 outlier points (they are included in the correlation calculation); **Fig. S5F** shows the same data with uncut axes. **(F)** Mapping of the integrations in sorted populations P1-P6 (**Fig. S5E**) and unsorted control cells (ctrl). Visualization as in **Fig. 2D**; data from 2 biological replicates combined. **(G)** Expression score of the Sox2P and Sox2P_CDS reporter across the *Sox2* locus (-161 kb launch pad); 5 kb running window shifted in 500 bp steps. Expression score of the Sox2P reporter is same as in **Fig. 2I**. Shaded region indicates the 95% confidence interval. Expression score is only plotted when the window contains 3 or more SB integrations.

To test whether this competitive characteristic of the CDS reporters is a general feature throughout the locus, we mobilized the CDS-containing reporter from the -161 kb location – where it is not detectably expressed (**Fig. S5E**) – in cells carrying the intact *Sox2*::*mCherry* (**Fig. 5E**). We compared the relative *Sox2*::*mCherry* and reporter expression of individual cells with active reporters, as measured by flow cytometry. If the reporter competes with *Sox2*::*mCherry*, this should be reflected in reduced expression of the latter when the reporter is more active. With the reporter lacking the CDS, this effect is detectable but weak (**Fig. 5E, S5F**). However, when the CDS is present, this effect is about three times stronger (p = 0.008), as indicated by the respective slopes of a fitted regression model (**Fig. 5E, S5G**). Thus, across the random active locations sampled in this hopping experiment, the CDS strengthens the reporter in its competition with the *Sox2* gene.

To further assess if the presence of the CDS in the reporter can change its activation landscape throughout the locus, we mapped the integrations in cells sorted for different reporter expression levels from the CDS hopping experiment (**Fig. S5H**). The sorted integrations of CDS-containing reporter follow a similar pattern as the original reporter (**Fig. 5F**), indicating that the two reporters are stimulated by the SCR in a similar way. However, we did not detect any cells with the highest reporter expression (P1), causing flattening of the activity landscape near the SCR and the endogenous *Sox2* location. This is possibly explained by the same negative feedback that we proposed above. Despite this, the CDS boosted the reporter activity throughout the *Sox2*-SCR range (excluding *Sox2*::*mCherry* and SCR regions) (**Fig. 5G**). Together, these data indicate that the CDS has two effects on promoter activity: it can boost expression independent of competition; and it increases the competitive edge of a promoter.

Discussion

Here we developed a novel high-throughput "hopping" approach to create highly detailed quantitative
maps of the activation potential of a native genomic locus, based on thousands of reporter integrations.
We expect that this approach may also be used to systematically relocate other regulatory elements, such
as enhancers, CBSs and other structural elements, and a wide range of synthetic elements.

Earlier probing at low resolution in mouse embryos suggested that promoters respond to an
enhancer whenever they are in the same TAD (*11, 27*). Systematic insertion of enhancers and promoters
in heterologous genomic loci supported this notion, and pointed to a monotonous decay of promoter
activity with increasing genomic distance to the enhancer (*8, 9, 28*). Our high-resolution functional maps
of the native *Sox2* locus confirm that the influence realm of the SCR coincides with the TAD, but also reveal
an intricate landscape in which pre-existing SCR contact frequency rather than genomic distance is a key
determinant of promoter activity. Moreover, the region around the endogenous *Sox2* gene is a "sweet
spot" for activation, even in the absence of the gene itself. Possibly, synergistic interactions between the
SCR and enhancer-like elements surrounding the gene (*16*) contribute to this.

Another striking result is that the *Sox2* gene is not only passively controlled by the SCR, but also
actively constrains the SCR, as indicated by its strong dominance over the reporter. Various studies of
native and artificial loci, both in *Drosophila* (*29, 30*) and mammals (*31-33*), showed competition between
two promoters for activation by one enhancer. In each case, the enhancer preferentially activated the
closest promoter. In contrast, our results show that the SCR strongly favors the native *Sox2* gene
regardless of the location of the reporter. Surprisingly, this preference is in part mediated by the *Sox2*
coding sequence. Possibly the CDS DNA sequence binds one or more transcription factors that boost SCR
interactions; or the transcribed CDS RNA facilitates SCR interactions, as has been proposed for other
transcripts (*34*). Our functional maps of SCR – gene interplay lay the foundation for further mechanistic
dissection.

References

1. J. H. Yang, A. S. Hansen, Enhancer selectivity in space and time: from enhancer-promoter interactions to promoter activation. *Nat Rev Mol Cell Biol* **25**, 574-591 (2024).

2. C. Paliou *et al.*, Preformed chromatin topology assists transcriptional robustness of Shh during limb development. *Proc Natl Acad Sci U S A* **116**, 12390-12399 (2019).

3. C. P. Fulco *et al.*, Activity-by-contact model of enhancer-promoter regulation from thousands of CRISPR perturbations. *Nat Genet* **51**, 1664-1669 (2019).

4. J. R. Dixon *et al.*, Topological domains in mammalian genomes identified by analysis of chromatin interactions. *Nature* **485**, 376-380 (2012).

5. E. P. Nora *et al.*, Spatial partitioning of the regulatory landscape of the X-inactivation centre. *Nature* **485**, 381-385 (2012).

6. G. R. Cavalheiro, T. Pollex, E. E. Furlong, To loop or not to loop: what is the role of TADs in enhancer function and gene regulation? *Curr Opin Genet Dev* **67**, 119-129 (2021).

7. J. Dekker, E. Heard, Structural and functional diversity of Topologically Associating Domains. *FEBS Lett* **589**, 2877-2884 (2015).

8. N. J. Rinzema *et al.*, Building regulatory landscapes reveals that an enhancer can recruit cohesin to create contact domains, engage CTCF sites and activate distant genes. *Nat Struct Mol Biol* **29**, 563-574 (2022).

9. J. Zuin *et al.*, Nonlinear control of transcription through enhancer-promoter interactions. *Nature* **604**, 571-577 (2022).

10. V. W. Keng *et al.*, Region-specific saturation germline mutagenesis in mice using the Sleeping Beauty transposon system. *Nat Methods* **2**, 763-769 (2005).

11. O. Symmons *et al.*, Functional and topological characteristics of mammalian regulatory domains. *Genome Res* **24**, 390-400 (2014).

12. L. Dauban *et al.*, Genome - nuclear lamina interactions are multivalent and cooperative. *bioRxiv*, 2024.2001.2010.574825 (2024).

13. Y. Li *et al.*, CRISPR reveals a distal super-enhancer required for Sox2 expression in mouse embryonic stem cells. *PLoS One* **9**, e114485 (2014).

14. H. Y. Zhou *et al.*, A Sox2 distal enhancer cluster regulates embryonic stem cell differentiation potential. *Genes Dev* **28**, 2699-2711 (2014).

15. T. Taylor *et al.*, Transcriptional regulation and chromatin architecture maintenance are decoupled functions at the Sox2 locus. *Genes Dev* **36**, 699-717 (2022).

16. R. Brosh *et al.*, Synthetic regulatory genomics uncovers enhancer context dependence at the Sox2 locus. *Mol Cell* **83**, 1140-1152 e1147 (2023).

17. B. Bonev *et al.*, Multiscale 3D Genome Rewiring during Mouse Neural Development. *Cell* **171**, 557-572 e524 (2017).

18. V. Y. Goel, M. K. Huseyin, A. S. Hansen, Region Capture Micro-C reveals coalescence of
enhancers and promoters into nested microcompartments. *Nat Genet* **55**, 1048-1056 (2023).

19. D. L. Stern, Tagmentation-Based Mapping (TagMap) of Mobile DNA Genomic Insertion Sites.
bioRxiv, 037762 (2017).

20. H. W. King, N. A. Fursova, N. P. Blackledge, R. J. Klose, Polycomb repressive complex 1 shapes
the nucleosome landscape but not accessibility at target genes. *Genome Res* **28**, 1494-1507
(2018).

21. A. Sethi *et al.*, Supervised enhancer prediction with epigenetic pattern recognition and targeted
validation. *Nat Methods* **17**, 807-814 (2020).

22. M. Gasperini, J. M. Tome, J. Shendure, Towards a comprehensive catalogue of validated and
target-linked human enhancers. *Nat Rev Genet* **21**, 292-310 (2020).

23. S. Masui *et al.*, Pluripotency governed by Sox2 via regulation of Oct3/4 expression in mouse
embryonic stem cells. *Nat Cell Biol* **9**, 625-635 (2007).

24. A. L. Szymczak, D. A. Vignali, Development of 2A peptide-based strategies in the design of
multicistronic vectors. *Expert Opin Biol Ther* **5**, 627-638 (2005).

25. J. L. Kopp, B. D. Ormsbee, M. Desler, A. Rizzino, Small increases in the level of Sox2 trigger the
differentiation of mouse embryonic stem cells. *Stem Cells* **26**, 903-911 (2008).

26. B. D. Ormsbee Golden, E. L. Wuebben, A. Rizzino, Sox2 expression is regulated by a negative
feedback loop in embryonic stem cells that involves AKT signaling and FoxO1. *PLoS One* **8**,
e76345 (2013).

27. O. Symmons *et al.*, The Shh Topological Domain Facilitates the Action of Remote Enhancers by
Reducing the Effects of Genomic Distances. *Dev Cell* **39**, 529-543 (2016).

28. H. Thomas *et al.*, Enhancer cooperativity can compensate for loss of activity over large genomic
distances. *bioRxiv*, 2023.2012.2006.570399 (2023).

29. T. Fukaya, B. Lim, M. Levine, Enhancer Control of Transcriptional Bursting. *Cell* **166**, 358-368
(2016).

30. J. R. Bateman, J. E. Johnson, Altering enhancer-promoter linear distance impacts promoter
competition in cis and in trans. *Genetics* **222**, (2022).

31. N. Dillon, T. Trimborn, J. Strouboulis, P. Fraser, F. Grosveld, The effect of distance on long-range
chromatin interactions. *Mol Cell* **1**, 131-139 (1997).

32. S. Oh *et al.*, Enhancer release and retargeting activates disease-susceptibility genes. *Nature* **595**,
735-740 (2021).

33. S. K. Topfer *et al.*, Disrupting the adult globin promoter alleviates promoter competition and
reactivates fetal globin gene expression. *Blood* **139**, 2107-2118 (2022).

34. L. Liang *et al.*, Complementary Alu sequences mediate enhancer-promoter selectivity. *Nature* **619**,
868-875 (2023).

35. J. C. Rivera-Mulia *et al.*, Allele-specific control of replication timing and genome organization during
development. *Genome Res* **28**, 800-811 (2018).

36.   C. E. Grant, T. L. Bailey, W. S. Noble, FIMO: scanning for occurrences of a given motif.
Bioinformatics **27**, 1017-1018 (2011).

37.   R. Schep *et al.*, Impact of chromatin context on Cas9-induced DNA double-strand break repair
pathway balance. *Mol Cell* **81**, 2216-2230 e2210 (2021).

38.   L. Cong *et al.*, Multiplex genome engineering using CRISPR/Cas systems. *Science* **339**, 819-823
(2013).

39.   F. Hahne *et al.*, flowCore: a Bioconductor package for high throughput flow cytometry. *BMC*
Bioinformatics **10**, 106 (2009).

40.   A. G. Doran *et al.*, Deep genome sequencing and variation analysis of 13 inbred mouse strains
defines candidate phenotypic alleles, private variation and homozygous truncating mutations.
Genome Biol **17**, 167 (2016).

41.   T. M. Keane *et al.*, Mouse genomic variation and its effect on phenotypes and gene regulation.
Nature **477**, 289-294 (2011).

42.   G. Finak, W. Jiang, R. Gottardo, CytoML for cross-platform cytometry data sharing. *Cytometry A*
93, 1189-1196 (2018).

43.   M. Malek *et al.*, flowDensity: reproducing manual gating of flow cytometry data by automated
density-based cell population identification. *Bioinformatics* **31**, 606-607 (2015).

44.   S. Chen, Ultrafast one-pass FASTQ data preprocessing, quality control, and deduplication using
fastp. *Imeta* **2**, e107 (2023).

45.   R. H. van der Weide *et al.*, Hi-C analyses with GENOVA: a case study with cohesin variants. *NAR*
Genom Bioinform **3**, lqab040 (2021).

Acknowledgments

We thank the Giorgetti lab for sharing the *Sox2* fosmid; the NKI Genomics, Flow Cytometry, Protein Production and Research High Performance Computing core facilities for excellent support; Marantha Kaagman for valuable experimental contributions; and Elphege Nora, Luca Giorgetti, Elzo de Wit, Tineke Lenstra and members from our lab and the divisions of Gene Regulation and Molecular Genetics for inspiring and helpful discussions.

Funding

This project was funded by the Boehringer Ingelheim Fonds (C.J.I.M.); the European Research Council: GoCADiSC, 694466 (B.v.S.); RE_LOCATE, 101054449 (B.v.S.) and the Oncode Institute. Views and opinions expressed are however those of the authors only and do not necessarily reflect those of the European Union or the European Research Council. Neither the European Union nor the granting authority can be held responsible for them. Research at the Netherlands Cancer Institute is supported by an institutional grant of the Dutch Cancer Society and of the Dutch Ministry of Health, Welfare and Sport. The Oncode Institute is partially funded by the Dutch Cancer Society.

Author Contribution

Conceptualization: M.E., C.J.I.M., B.v.S.

Wet-lab experiments: M.E., C.J.I.M., L.D.

Data analysis and visualization: C.J.I.M., M.E.

Method development: M.E., L.D.

Code development: C.L., C.J.I.M.

Writing: M.E., C.J.I.M., B.v.S.

Funding Acquisition: B.v.S., C.J.I.M.

Supervision: B.v.S.

Competing interests:

The authors declare no conflict of interest

Data and materials availability:

Laboratory notebooks and supplementary data (including primer sequences, gRNAs, plasmids, etc.) are available on Zenodo (10.5281/zenodo.13350269). Raw sequencing data is available on GEO under accession numbers GSE275427.

Code and supplementary files used as input for the scripts are available on Zenodo (10.5281/zenodo.13362255).

All plasmids used will be made available to Addgene.

Fig. S1. Cell line generation and proof of principle experiment. (A) Left and middle: PCR and Sanger sequencing with mCherry and eGFP specific primers indicates on which allele each fluorophore is inserted. Right: expression level of Sox2::mCherry and Sox2::eGFP measured by flow cytometry. **(B)** Allele-specific Sox2 expression in the dual tagged cell line, in cells treated with CRISPR to delete the SCR and in untreated control cells. **(C)** Recombination-mediated cassette exchange: The hygromycin phosphotransferase-thymidine kinase fusion gene (HyTK) is flanked by heterotypic Flp recombinase recognition sites (FRT & F3) and embedded into a Sleeping Beauty transposon (SB-ITRs), integrated via CRISPR-Cas9-mediated knock-in 6 kb upstream of the Sox2::mCherry fusion gene. Co-transfection of the Flp recombinase expression plasmid (pFlp) and donor plasmid (pInsert) with subsequent ganciclovir selection for 7 days results in cells with the intended insert. (D-G) Statistics from the proof-of-principle hopping experiment, where a SB transposon carrying a 282 bp arbitrary DNA sequence was relocated. Number of integrations is indicated (above the bars or n). **(D)** Fraction of integrations mapped per chromosome. The launch pad is on chr 3. **(E)** Fraction of all integrations on chr 3 mapped to each allele, or not assigned to either allele. The launch pad is on the 129S1 allele. **(F)** Number of integrations per 10 kb bin, split by SB orientation. **(G)** Distribution of the number of unique mapping reads supporting each integration. **(B, D-G)** Data from one (proof-of-principle) biological replicate.

Fig. S2. Reproducibility of reporter hopping and validation of the expression score. (A) Distribution of expression of *Sox2::eGFP* (green), *Sox2::mCherry* (red) and the reporter (mTurq, blue), in the -161 kb and -116 kb reporter cell lines and the parental control cell lines measured by flow cytometry. **(B)** Reproducibility of the fraction of sorted integrations across the *Sox2* locus, from the -116 kb launch pad. Bottom left panels: each dot represents the fraction of all integrations of that population found in a 5 kb bin in the *Sox2* locus, compared between two biological replicates. Colors indicate the 6 sorted populations. Panels along diagonal: distribution of observed fractions in each replicate. Top right panels: Spearman correlation between two replicates. **(C)** As B, but for the -161 kb launch pad, with two biological replicates. **(D)** Top: the computed expression scores around *Sox2* gene and SCR based on the sorted integrations from the -116 kb and -161 kb launch pad (see Fig. 2D, H), smoothened using a 5 kb running window shifted in 500 bp steps. Shaded region indicates 95% ¬confidence interval. Grey triangles indicate CTCF binding sites (CBS) and their orientation. LP, launch pad position. Bottom: relative reporter expression (median mTurq, estimated autofluorescence subtracted, normalized to eGFP, and then normalized to the unhopped reporters at -116 kb) of clones around the *Sox2* gene and the SCR. Loess curve shows the trend in expression. **(E)** Expression score of the positive and negative strand integrations (forward and reverse oriented promoter, respectively), across the *Sox2* locus, based on the integrations from the -116 kb and -161 kb launch pad. Smoothened using a 10 kb running window, shifted in 1 kb steps. Shaded region indicates 95% confidence interval. Locus annotation as in D.

Eder, Moene et al. Fig S3

Fig. S3. Mapping of hopping induced deletions and insertions. (A) Schematic of Tn5-based mapping in cells with a regular hopping event or a hopping-induced deletion. In case of a hopping-induced deletion, one SB ITR will be mapped to the launch site while the other is mapped to the insertion location. **(B)** Confirmation of the loss of the 129S1 allele in the deleted region of deletion clones. Top: genomic location and expression of two hopping-induced deletion clones (1, 2, blue) and one regular insertion clone (3, grey). Horizontal dashed grey lines indicate expected deletions, vertical dashed green lines indicate the locations of three known SNPs between the 129S1 and CAST alleles.
Bottom: Sanger sequencing across the three SNPs in the three cell lines. Black arrowheads indicate double peaks at the SNPs, indicating that both alleles are present. When only the CAST allele is present, the SNP is indicated in bold in the sequence above the Sanger trace.

Eder, Moene et al. Figure S4

Fig. S4. Sox2P-reporter expression in *Sox2::mCherry* deleted pools and clones. (A) Reporter expression in untreated control cell lines (-39 kb, -116 kb, -161 kb). **(B)** Distribution of reporter expression in *Sox2::mCherry*-negative, reporter-high sorted pools. **(C)** Distribution of reporter expression in CBS_*Sox2*::mCherry deleted, reporter-high sorted clones from the -161 kb reporter cell line. Clone D4 (in bold) was used for hopping (see Fig. 4D). **(D)** As in C, but for the -39 kb reporter cell line. Untreated reporter cell line is indicated in black. **(E)** As D, but for *Sox2*::mCherry deletion in the -116 kb reporter cell line. **(F)** As D, but for CBS_*Sox2*::mCherry deletion in the -116 kb reporter cell line. **(C-F)** Vertical lines indicate the (at most) two highest peaks in the density distribution, the peak with the highest mTurq level is used in Fig. 4B. **(G)** Expression of *Sox2*::eGFP (control) and the *Sox2* reporter (mTurq) in the untreated cells and cells transfected with the SB transposase plasmid, for the -161 kb ΔCBS_*Sox2*::mCherry reporter cell line. The percentage of cells in each sorted gate is indicated. Cells are gated for *Sox2*::mCherry-negative single cells. Representative image of 2 biological replicates.

Fig. S5. Impact of the Sox2 coding sequence on reporter and Sox2 expression. (A) Contact between *Sox2* and the SCR (region-capture micro-C data [(18)], 200-bp resolution). Positions of the components of the *Sox2* gene (blue, orange, red; x-axis) and SCR (yellow; y-axis) are indicated. Grey triangles indicate CTCF binding sites and their orientation. Arrows indicates the genomic direction. **(B)** Relative expression of the reporter (mTurq), *Sox2*::mCherry and *Sox2*::eGFP, not normalized to *Sox2*::eGFP, for the three reporters inserted at -161 kb and -116 kb. Each dot represents one clone. Expression values are autofluorescence subtracted, and then normalized to the control reporter cell line (*Sox2* reporter at -116 kb) measured on the same day. Dashed line indicates 1 (level of the control cell line). P-values according to Welch's unequal variances t-test. **(C)** Distribution of reporter expression of reporters integrated at -116 kb in a ΔCBS_*Sox2*::mCherry cell line. Each density plot is one clone, vertical lines indicate the (at most) two highest peaks in the density distribution. **(D)** As B, for the three reporters and a minimal insert (not containing any promoter or mTurq) inserted at -116 kb and -39 kb launch pads with *Sox2*::mCherry intact. **(E)** Reporter expression in a panel of clones with the Sox2P_CDS reporter integrated at the -161 kb launch pad with *Sox2*::mCherry intact. The arrow indicates the clone used for hopping (see Fig. 5F). **(F)** Correlation between relative *Sox2*::mCherry and reporter (mTurq) expression from the Sox2P and Sox2P_CDS reporter, from cells with reporter expression in the gates P2-P4 after induction of hopping. As in Fig. 5E, but also showing the outlier points. **(G)** Linear model of the data in F, describing how the relative *Sox2*::mCherry expression (rel_mCherry) depends on the identity of the reporter (CDS_reporter, is 0 for the Sox2P reporter and 1 for the Sox2P_CDS reporter), and the relative reporter expression (rel_mTurq). Right side indicates the p-values of the coefficients of the model, testing whether they are significantly different from 0. **(H)** Expression of *Sox2*::eGFP (control) and the *Sox2* reporter (mTurq) in the untreated cells and cells transfected with the SB transposase plasmid, for the -161 kb Sox2P_CDS reporter cell line. The percentage of cells in each sorted gate is indicated. Cells are gated for *Sox2*::mCherry-positive single cells. Representative image of 2 biological replicates.