Dissecting heterogeneity of tumor microenvironment in colorectal cancer using high-resolution single-cell atlas

Valentin Marteau[1, 11], Niloofar Nemati[1, 11], Kristina Handler[2, 11], Deeksha Raju[2], Erika Kvalem Soto[1], Georgios Fotakis[1], Sandro Carollo[1], Nina Böck[1], Alexander Kirchmair[1], Alexandra Scheiber[3], Arno Ammann[3], Andreas Seeber[3], Elisabeth Gasser[4], Lena Horvath[3], Hassan Fazilaty[5], Tomas Valenta[5, 8], Gregor Sturm[1,7], Dietmar Rieder[1], Sieghart Sopper[3], Andreas Pircher[3], Paul Ziegler[9], Markus Tschurtschenthaler[10], Dominik Wolf[3], Isabelle Arnold[2, 6, 12*], Stefan Salcher[3, 12*], Zlatko Trajanoski[1, 12*]

[1]Biocenter, Institute of Bioinformatics, Medical University of Innsbruck, Austria

[2]Institute of Experimental Immunology, University of Zurich, Switzerland

[3]Department of Internal Medicine V, Haematology & Oncology, Comprehensive Cancer Center Innsbruck (CCCI) and Tyrolean Cancer Research Institute (TKFI), Medical University of Innsbruck (MUI), Innsbruck, Austria

[4]Department of Visceral, Transplant and Thoracic Surgery, Medical University Innsbruck, Anichstrasse 35, 6020, Innsbruck, Austria

[5]Department of Molecular Life Sciences, University of Zürich, Zürich, Switzerland

[6]Comprehensive Cancer Center Zürich, Zürich, Switzerland

[7]Boehringer Ingelheim International Pharma GmbH & Co KG, Biberach, Germany

[8]Institute of Molecular Genetics of the Czech Academy of Sciences, Prague, Czech Republic

[9]Senckenberg. Institute of Pathology, Goethe University Frankfurt, Frankfurt, Germany

[10]Translational Cancer Research and Institute of Experimental Cancer Therapy, Klinikum rechts der Isar, School of Medicine, Technical University of Munich, Munich, Germany

[11]These authors contributed equally

[12]These authors contributed equally

*Correspondence: stefan.salcher@i-med.ac.at

*Correspondence: zlatko.trajanoski@i-med.ac.at

SUMMARY

Colorectal cancer (CRC) features molecular heterogeneity and differing immune cell infiltration
patterns, affecting how patients respond to therapy. However, a comprehensive understanding of
immune cell phenotypes across patient subgroups is missing. Here, we dissect the CRC tumor
microenvironment by integrating 4,27 million single cells from 1670 samples and 650 patients across
48 studies (76 datasets) representing 7 billion expression values. These samples encompass the full
spectrum of disease progression, from normal colon to polyps, primary tumors, and metastases, covering
both early and advanced stages of CRC. We present a high-resolution view of CRC with 62 major cell
types or states, revealing different cell-type composition patterns in CRC subtypes. The atlas allows for
refined tumor classification and patient stratification into four immune phenotypes: immune desert,
myeloid, B cell, and T cell subtypes. Using bulk samples with genomic and clinical information, we
identify cellular components associated with tumor histology and genotypes. These findings could have
important implications for developing improved cancer immunotherapy approaches in CRC.

KEYWORDS

Single-cell sequencing, cell-cell communication, patient stratification, therapy response

INTRODUCTION

Colorectal cancer (CRC) remains the most common gastrointestinal cancer and the second leading cause of cancer-related death according to the GLOBOCAN data[1]. The growing incidence in some countries reflects a modification in lifestyle and its consequences related with "Westernization" such as diet, obesity, physical inactivity, and alcohol consumption. Similarly, "Westernization" of diets and lifestyle have been attributed to the alarming growing incidence of early-onset CRCs (individuals <50 years of age)[2]. Despite the improvement of early detection screening policies and the development of many novel treatments, the median overall survival (OS) for patients with metastatic CRC (mCRC) is ~30 months[3]. Progress in targeted therapy for mCRC has been limited, relative to other solid cancers like lung cancer or melanoma[4]. The genetic heterogeneity[5] as well as paucity of drugs targeting common driver genes, pose considerable challenges for developing precision oncology approaches for patients with CRC. Moreover, CRC seems to be refractory to therapy with immune checkpoint inhibitors (ICI) with the notable exception of 4% of patients with tumors characterized by microsatellite instability (MSI) or POLE proofreading mutations[6]. This is somehow paradoxical since CRCs are known to be under immunological control, as we have shown in the past[7]. Thus, there is an urgent need to explore new immunomodulatory strategies for the vast majority of CRC patients with microsatellite stable (MSS) tumors. Strategies that convert the "immune cold" tumor microenvironment (TME) to "immune hot" could sensitize cancers to ICI therapy. A number of clinical trials with drug combinations have been carried out to overcome resistance to immunotherapy in MSS CRC[8,9]. The benefits of using a dual combination of ICIs (CTLA-4 and PD-1 blockade) or adding ICIs to chemotherapy appear to be very limited[8,10]. Combination strategies that exploit synergistic potential of ICI and targeted drugs showed increase of the anti-tumor activity, however, the objective response rates were disappointingly low[8]. Thus, to address this unmet clinical need a better understanding of the immunosuppressive TME and its heterogeneity is of utmost importance.

The intrinsic complexity of the interaction of the two interwoven systems, the tumor and the immune system, poses considerable challenges and requires comprehensive approaches to interrogate cancer immunity during tumor initiation and progression, and following therapeutic modulation thereof. A number of established and novel high-throughput technologies enable the generation of the necessary data and thereby provide the basis for mechanistic understanding. Specifically, next-generation sequencing (NGS) of cancer samples provide large data sets that can be mined for immunologically relevant parameters (see our reviews[11,12]). Using these tools we were able to chart the TME in CRC[13] and developed The Cancer Immunome Atlas[14]. While these attempts uncovered the enormous heterogeneity of MSS CRC and revealed unique cancer cell and TME subtypes with prognostic and therapeutic relevance, the advent of single-cell RNA sequencing (scRNA-seq) technologies opened fundamental new avenues to dissect cancer heterogeneity. We recently developed a novel computational

approach to dissect the cellular diversity in the TME at high resolution by integrating expression profiles

of single cells from multiple studies using cutting-edge artificial intelligence (AI) methods[15].

Here we used our approach to build a high-resolution single-cell transcriptome atlas for CRC with 4.27

million cells from 650 patients and 1,670 samples. This high-resolution atlas offers superior coverage

of histological and clinical variables and provides a unique resource for dissecting the cellular diversity

in the TME. We then used bulk RNA-seq data from The Cancer Genome Atlas (TCGA)[5] to evaluate the

association of atlas-derived single-cell derived transcriptomic signatures with survival data. The

exploitation of this high-resolution single-cell transcriptome atlas for CRC will provide novel biological

insights that will be the basis to improve therapy in MSS CRC patients.

RESULTS

Generation of a core large-scale CRC single-cell atlas

We developed a core CRC atlas by compiling scRNA-seq data from 48 studies and 76 datasets comprising 4.27 million cells from 650 patients and 1,670 samples (Figure 1A). This comprehensive CRC single-cell atlas integrates expert-curated, quality-assured and pre-analyzed transcriptomic data from publicly available studies encompassing the full spectrum of disease progression, from normal colon to polyps, primary tumors, and metastases, covering both early and advanced stages of CRC (see Methods). Important study characteristics are summarized in Table S1. In total, the core atlas includes transcriptomic data from 487 patients with CRC, 99 patients with polyps, and 64 control individuals. Of the 487 CRC patients, 254 were histopathologically annotated as colon (COAD), 63 as rectal cancer (READ), and 170 were not otherwise specified. CRC samples include tissue of the primary tumor (n=674) or metastasis (n=175) that were obtained by surgical resection. Among the control samples, 284 were derived from distant non-malignant tissue of colon tumor patients, of which 220 have a patient-matched tumor sample. 186 non-tumor control samples as well as 116 polyp and 90 matched normal samples were obtained through patients undergoing colonoscopy. Further, 103 blood samples of CRC patients as well as 24 samples derived from tumor affected lymph nodes, with another 18 non-tumor affected lymph node samples were included.

We clustered the CRC patients' disease stages as early (UICC stage I – II) *versus* advanced (UICC III-IV) disease, as not all studies provided sufficient information on tumor stages. Of the control patients none had a history of IBD. Overall, the core atlas integrates 4,264,946 single cells, which are annotated to 11 coarse cell type identities and 62 major cell subtypes or cell states (e.g. dividing cells) based on previously established canonical single-cell signatures including 1,146,196 epithelial cells, 1,552,108 T/NK cells, 804,042 B/Plasma cells, 385,589 myeloid cells, and 33,9767 stromal and endothelial cells (Figure 1B).

Cancer cells (in total 600,740 cells, from primary tumor and metastatic tissue) in the atlas showed high heterogeneity of their transcriptomic profiles (Figure 1B). Due to the large patient cohort, we were able to apply high-resolution cancer cell classification based on their specific marker gene expression signatures (Figure S1D). We divided the following main clusters Colonocyte-like (*FABP1, CEACAM7, ANPEP, SI*), BEST4 (BEST4, *OTOP2, SPIB*), Goblet-like (*MUC2, FCGBP, ZG16, ATOH1*), Crypt-like (*LGR5, SMOC2*), and TA-like (*TOP2A, UBE2C, PCLAF, HELLS, TK1*) (Figure 1B, Figure S1D). There are highly mitotic/proliferative clusters (*LGR5, TOP2A, MKI67*) of both COAD and READ that may resemble highly aggressive and invasive cancer cells. In addition, *EPCAM* expressing cells in the blood of CRC patients were annotated as circulating cancer cells.

Single-cell composition of the TME reveals distinct CRC tumor immunotypes

Next, we stratified CRC patients based on infiltration patterns of their TME using the high-resolution atlas. Unsupervised clustering on batch-corrected cell-type fractions revealed four distinct tumor immune phenotypes (Figure 2A): 1) immune deserted tumors (ID; i.e. no significant immune cell infiltration but high cancer cell fraction); 2) the subtype of tumors with B cell dominance (B; B cell, plasma IgA and IgG cell); 3) the subtype of tumors with myeloid dominance (M; macrophage/monocyte); and 4) subtype of tumors with T cell dominance (T; CD8+, CD4+, T regulatory cells). Using logistic regression, we did not find any association of the different patient strata to tumor stages (early *vs.* advanced) or sex.

We identified gene signatures based on the patient immune infiltration groups by selecting the top 15 differentially expressed genes. These gene signatures were then used to conduct a survival analysis utilizing data from the TCGA COAD-READ cohorts (n = 629). The Kaplan-Meier curves in Figure 2C demonstrate that higher expression of the B (p-value = 0.0015), M (p-value = 0.0096), and T (p-value = 0.00078) signatures is significantly correlated with improved overall survival. The analysis showed that higher expression of B, M, and T gene signatures is significantly linked to better overall survival.

Analysis of cell-cell communication uncovers interactions between different cell types in the tumor microenvironment (TME)

Differences in cellular crosstalk were analyzed using LIANA[29] framework, which leverages a consensus of expert-curated ligand-receptor databases, as detailed in the methods section. In this study, our primary focus is on the interactions between cancer cells and the main immune cell types. Specifically, we examined the top 10 differentially expressed ligands in cancer cells (Figure 3). *CD69,* which is known to be involved in the early activation and proliferation of T lymphocytes[31], was found to be upregulated in CD8+ T cells, regulatory T cells, and CD4+ T cells. We observed increased signaling of several chemokine receptors. This finding may indicate that cancer cell-secreted chemokine gradients may be contributing to immune cell infiltration into the tumor microenvironment.

Integration of bulk RNA-seq data reveals genotype-immune phenotype associations

scRNA-seq provides an unprecedented view on the cellular heterogeneity in the TME. However, the majority of the scRNA-seq studies lack both comprehensive cancer genotype information and survival data. In contrast, the TCGA reference dataset includes both of these, alongside bulk RNA-sequencing data. Using the computational method SCISSOR[36], we integrated bulk RNA-seq data from TCGA with single-cell transcriptomic signatures to assess the relationship between specific genotypes and survival outcomes in CRC. We evaluated the association of cell type transcriptomic signatures derived from the atlas with genotype and survival data from the TCGA reference dataset, which includes 629 patients (UICC I-IV)[5]. Building on our previous work that identified genomic features influencing immune

phenotypes in a pan-cancer study using bulk RNA-seq data[37], we leveraged the high resolution of the

single-cell CRC atlas for a more detailed analysis of these determinants.

We next analyzed the cell type transcriptomic signatures and their association with survival in 629

patients from the TCGA cohort. Overall, the presence of high levels of Plasma IgA, $CD4^+$ T cells $CD8^+$

T cells and B cells showed a prominent association with improved survival in CRC patients. Conversely,

CRC patients with tumors enriched in Schwann cells were associated with worst survival outcomes, as

also reported previously[38]. Notably, high infiltration of various endothelial cells and fibroblasts also

emerged as strong negative predictors of survival (Figure 4A). Mutations in *BRAF, KRAS* were

associated with distinct immune infiltrates (Figure 4B-C). For example, cDC1 and cDC2 showed high

infiltration patterns in CRC patients with BRAF mutations and same for Plasma IgM where the patterns

were present in both BRAF and KRAS mutations. In contrast, mutations in BRAF were associated with

low Plasma IgA infiltration, whereas KRAS mutations correlated with high Plasma IgA infiltration. MSI

and MSS status were associated with opposite cell type patterns (Figure 4D-E).

DISCUSSION

We built a large-scale atlas of single-cell transcriptomes of CRC through integration of 4.27 million cells from 650 patients and 1,670 samples representing 7 billion expression values. Reduction of dataset-specific batch effects due to variation in experimental design and used platforms while retaining biological information resulted in a high-quality reference atlas for CRC, offering superior coverage of histological and clinical variables and thereby providing a unique resource for dissecting the cellular diversity in the TME and generating hypotheses.

We provide a high-resolution view of the TME in CRC with 44 major cell types/states and show different cell type composition patterns including more precise functional transcriptomic classification of malignant epithelial cells. The single-cell composition of the TME in CRC enabled refined tumor classification and patient stratification into four immune phenotypes: immune deserted, myeloid, B cell, and T cell subtypes. These findings may have important implications for enabling cancer immunotherapy in CRC. For example, combination therapies that target both, myeloid cells and lymphoid cells could represent an immunotherapeutic strategy to treat M subtypes as shown recently in a melanoma mouse model[41]. Similarly, given the heterogeneity of the intratumoral B cells and the importance of tertiary lymphoid structures[42], further analysis of the B subtypes might open promising therapeutic avenues by additional refined B cell targeting.

Furthermore, integration of bulk RNA-seq data from the TCGA CRC cohort uncovered cell subsets associated with alterations in major driver genes, such as *BRAF*, *KRAS*, as well as with the CRC subtypes MSI and MSS. Hence, we provide further evidence that genetic aberrations in cancer cells dictate the immune contexture in tumors. This knowledge could be exploited to derive rationale for personalized therapeutic combination strategies based on the underlying genetic profile of the tumor. The biological insights we present here and future discoveries arising from the exploitation of the high-resolution CRC atlas could provide the basis for developing combination therapies for CRC patients who are not sufficiently responding to immune checkpoint blockers.

ACKNOWLEDGMENTS

This work was supported by the European Research Council (grant agreement No 786295 to ZT), by the Austrian Science Fund (FWF) (project I3978 to ZT). ZT is a member of the German Research Foundation (DFG) project TRR 241(INF). The results published here are in part based upon data generated by the TCGA Research Network: https://www.cancer.gov/tcga. VM was funded by 10.55776/DOC82. For open access purposes, the author has applied a CC BY public copyright license to any author accepted manuscript version arising from this submission.

FIGURE LEGENDS

Figure 1. Schematic outline of the overall concept used in this study.

(A) Summary of the data integration and analysis workflow.

(B) Overview of the CRC atlas and the epithelial, immune, and stromal/endothelial components depicted as UMAP plots.

(C) Fractions of depicted cell types per scRNA-sequencing platform.

(D) Tissue samples and the corresponding cell types.

Figure 2. Tumor immune phenotypes in CRC.

(A) Patient characteristics and stratification of the tumor immune phenotypes. *Tumor type (histopathological)* refers to the histological subtypes as provided by the original datasets based on pathological assessment; *tumor type (transcriptomic)* is based on the most abundant transcriptomically annotated cancer cell-subtype in the scRNA-seq atlas.

(B) Differential activation of (B) PROGENy cancer pathways- (C) CytoSig cytokine signaling signatures in cancer cells between the four tumor immune phenotypes. Heatmap colors indicate the deviation from the overall mean, independent of tumor histology and stage. White dots indicate significant interactions at different false-discovery-rate (FDR) thresholds. Only cytokine signatures with an FDR < 0.1 in at least one patient group are shown.

See also Figure S2.

(C) Kaplan-Meier curves of the survival analysis according to the patient immune infiltration groups (B, desert, M, T), performed using the molecular data of the TCGA COAD-READ cohorts (n = 461).

Figure 3. Cellular crosstalk analysis

(A) Circos plot illustrating the interaction network between cancer cells and major immune cells in paired tumor and normal samples. The plot highlights the top 10 differentially expressed ligands from cancer cells. Red lines indicate interactions upregulated in tumors, while blue lines represent those upregulated in normal.

Figure 4. Association of cellular composition and distinct genotypes and survival in the TCGA data.

(A-E) *SCISSOR* analysis relating phenotypic information from bulk RNA-seq data from TCGA with single cells. UMAP plots indicate the position of cells positively (blue) or negatively (red) associated with mutation or better survival. A log2 ratio > 0 indicates a positive association with mutation, or better survival, respectively. Shown are cell-types with a log2 ratio significantly different from 0 at an FDR < 0.01 (paired Wilcoxon test).

255   (A) Association of cellular composition with overall survival.

256   (B) Association of cellular composition with BRAF mutation in CRC patients (n = 136).

257   (C) Association of cellular composition with KRAS mutation in CRC patients (n = 489).

258   (D) Association of cellular composition with MSI status in CRC patients (n = 177).

259   (E) Association of cellular composition with MSS status in CRC patients (n = 412).

METHODS

Generation of the atlas

The data for the core atlas was from previously published studies[46-56], our own data, and 12 studies for control purpose[50,57-62]. The selected studies were published between May 2017 and July 2024 and the incorporated datasets were generated with six different sequencing platforms, including the most commonly applied 10x Chromium (10x Genomics) as well as Smart-seq2[63], SMARTer (C1), scTrio-seq2, GEXSCOPE (Singleron), TruDrop[64] and DNBelab C4[65]. We further integrated our own data generated with the microwell-based BD Rhapsody scRNA-seq platform (BD Biosciences). We specifically selected studies using comparable protocols for sample processing and data generation, such as sequencing of whole cells. We did not exclude studies that applied flow cytometry-based cell-sorting prior to sequencing, as these incorporate relevant information on rare cell types[48,49,66]. From non-CRC studies we exclusively included those parts of the published data that were relevant for our atlas: from the Garrido Trigo dataset [57] we only included colon samples, from Burclaff[60], Kong[58], Mazzurana[61], Parikh[59], Smillie, Thomas, and Scheid[46] datasets we only used the control samples.

Integration of scRNA-seq datasets

Individual datasets were merged into a single AnnData object. Since genome annotations partly differed between the datasets, we mapped gene symbols to the ensembl ids of the used annotation version and updated the symbols to the latest version of HGNC. If gene ids were missing from a dataset, the values were filled with zeros. Genes that were missing in more than 19 datasets (25%) were excluded altogether.

We integrated the datasets using the scANVI algorithm[68,69], as it has been demonstrated to be one of the top-performing methods for atlas-level integration and to scale to >1M cells[70]. Since scANVI requires cell-type annotations for at least one of the input datasets, we manually annotated two "seed" datasets based on unsupervised clustering as described below. Raw counts were used as input for scANVI. Counts from plate-based assays such as Smart-seq2 were scaled by the gene length as recommended on the scvi-tools website. The scANVI model was initialized with a pre-trained scVI model[71], as recommended in the scvi-tools tutorial. The scVI model was trained on the 6000 most highly variable genes as determined with scanpy's[72] *pp.highly_variable_genes* with parameters *flavor="seurat_v3"* and *batch_key="dataset"*. Each sample was considered as an individual batch for both scVI and scANVI. Other than that the algorithms were run with default parameters.

Unsupervised clustering and cell-type annotation

We computed UMAP embeddings[73] and unsupervised Leiden-clustering[74] with scanpy[72], based on a cell-cell neighborhood graph derived from scANVI latent space. Coarse, lineage-specific clusters were

297   iteratively sub-clustered to identify cell-types at a more fine-grained resolution. Cell type clusters were

298   annotated based on previously reported marker genes[57,75,76]

Comparing cell-type abundances

Since comparing cell-type fractions between groups is challenging due to different characteristics of the

datasets and the inherent compositional nature of cell-type fractions, we applied the scCODA [77] model,

which addresses this issue. We were interested in the differences between conditions (COAD *vs.*

READ). To this end, we ran the scCODA model with the formula ~ *condition + tumor_stage + dataset*

with 500,000 iterations using "cancer cells" as the reference cell-type, where *tumor_stage* is a binary

vector classifying datasets into early (stages I-II) and advanced (stages III-IV), and dataset is a

categorical vector encoding the different datasets. The final result shows *credible effects* with a false-

discovery-rate (FDR) of 0.1.

Patient stratification

We stratified patients into immune phenotypes based on immune cell-type fractions. We selected all

patients with primary tumor samples and excluded enriched datasets. Neutrophil fractions were

excluded, since they are not appropriately captured in the majority of datasets. Cell-type fractions of

primary tumor samples were loaded into a patient × cell-type AnnData container. Dataset-specific batch-

effects were removed using a linear model as implemented in *scanpy.pp.regress_out*. Patients were

clustered using graph-based Leiden clustering with the "correlation" distance metric for computing the

neighborhood graph. Patient clusters were labeled according to their predominant cell-types. In addition

to the histological subtypes based on the annotation of the original datasets, we annotated tumor types

based on the transcriptomics data according to the most abundant cancer cell cluster.

Differential gene expression testing

We used DESeq2[78] on pseudo-bulk samples for differential expression testing which has been

demonstrated to perform well and properly correct for false discoveries[79]. For each cell-type and patient,

we summed up transcript counts for each gene. Pseudo-bulk samples consisting of fewer than 10 cells

were discarded. We compared primary tumor samples from tumor *vs.* normal (*sample type*), including

the dataset as a covariate. For comparisons between multiple groups, we used contrasts with sum-to-

zero coding. P-values were adjusted for multiple hypothesis testing with independent hypothesis

weighting (IHW)[80].

Pathway, TF and cytokine signaling signatures

We performed pathway, transcription factor (TF), and cytokine signaling analysis on primary tumor

samples with PROGENy[22,25], DoROthEA[24,25] and CytoSig[27], respectively. Scores were computed using

the *dorothea-py* and *progeny-py* packages. The top 1,000 target genes of the progeny model were used,

as recommended for single-cell data. For dorothea, only regulons of the highest confidence levels "A"

and "B" were used. The cytosig signature matrix was obtained from the *data2intelligence/CytoSig*

GitHub repository and used with the scoring function implemented in the *progeny-py* package. The

methods were run with the options *num_perm=0, center=True, norm=True scale=True,* and

min_size=5. No permutations were used, as we perform statistics in a separate step at the level of

biological replicates. Pathway-, transcription factor-, and cytosig scores were then compared between

condition and patient *group* (T *vs*. B *vs.* M *vs.* deserted) using an ordinary least-squares (OLS) linear

model, as implemented in the statsmodels package [81]. Scores were aggregated into pseudobulk samples

by computing the mean of each variable for each patient and cell-type. Samples consisting of less than

10 cells were discarded. For each variable, we fitted a model with the formulas *~ condition + dataset +*

tumor_stage or *~ group + dataset + condition + tumor_stage*, respectively. Coefficients were obtained

from the linear model and p-values calculated with the f-test. P-values were adjusted for multiple testing

with the Benjamini-Hochberg procedure.

LIANA analysis

The ligand-receptor inference was computed using LIANA[29] framework to explore differences in cell-

to-cell communication in primary tumor samples. The default *consensus* resource was used, which

combines multiple expert-curated ligand-receptor databases, such as CellPhoneDB[82], CellChat[83],

ICELLNET[84], connectomeDB2020, and CellTalkDB[85]. The *rank_aggregate* method was chosen

combining ligand-receptor scores from multiple methods. Each method infers relevant interactions

based on different assumptions, providing scores that reflect the interaction's strength and its specificity

to particular cell identities. Our study focused on differences between conditions (tumor vs normal),

treating patients as biological replicates. To identify differentially expressed in CCC events, we

combined differential expression results for each gene into statistics of potentially downregulated

ligand-receptor interactions. This analysis used adjusted p-values and the Wald test statistic. Interactions

with low expression were filtered out when either a gene was expressed in less than 0.1 of cells, which

is the default threshold. The fraction of cells expressing a signaling molecule was calculated as the mean

of fractions per patient.

SCISSOR analysis

We used SCISSOR[86] to associate phenotypic data from bulk RNA-seq experiments with our single-cell

data. TCGA mutation and gene expression data was obtained from the GDC portal, survival data from

[86]. SCISSOR was run on primary tumor cells of each patient individually according to the SCISSOR

tutorial using mutation data (logistic regression) and overall survival (cox-regression) as dependent

variables. A grid search for the alpha-parameter was performed in $2^{-i/2}$ with $i \in [24, 23, \ldots, 2]$ and a

cutoff parameter of 0.3. 62 of 674 samples with low overall cell count failed during SCISSOR's Seurat-

preprocessing step and were excluded from the subsequent analysis. For each patient and cell-type, we

computed the fraction of *scissor+ cells* (i.e. positively associated with a mutation or worse survival),

scissor- cells (i.e. negatively associated), and *neutral cells* and added a pseudo-count of 0.01. A sample

was excluded from a cell-type if it contributed ≤10 cells. For each cell-type, we computed the log2-

ratio of scissor+ and scissor- cells as the mean fraction of scissor+ cells *vs.* the mean fraction of scissor-

cells. Significant differences were determined by comparing the fractions of scissor+ and scissor- cells

with a paired wilcoxon test with *zero_method="zsplit"* as implemented in the scipy package. P-values

were Benjamini-Hochberg-adjusted and considered significant at an FDR < 0.01.

Survival analysis

Initially, gene lists were extracted from the patient groups according to their immune infiltration profiles

(B, desert, M and T). The gene lists consisted of differentially expressed genes, as returned from DESeq2

for the comparisons of each immune infiltration group with the rest, that were filtered using as thresholds

the adjusted p-values (< 0.05) and the logFold Change (greater than 0.5 or lower than -0.5).

Consequently, the gene lists were condensed into gene signatures expression values using the GSVA

(v1.50.5) R package. Once the gene signature expression values were calculated the TCGA COAD-

READ cohort (n = 629) was stratified into "high" and "low" expression, using as threshold the optimal

cutpoint as computed by the surv_cutpoint() function of the survminer (v0.4.9) R package. The survival

analysis was carried out using the survival (v3.7-0) R package and the Kaplan-Meier curves were plotted

using the ggsurvplot function of the survminer (v0.4.9) R package.

Quantification and statistical analysis

Statistical analysis was performed using the statsmodels library in Python (scRNA-seq data) or

GraphPad Prism (flow cytometry and imaging data) using a linear model, t-test or wilcoxon test as

appropriate. Single cell-data were aggregated into pseudobulk samples by biological replicates.

Compositional analysis of cell-type fractions was performed using scCODA; survival analysis using the

Kaplan-Meier formula in R. P-values for untargeted analyses (DE genes, TFs, or pathways) were FDR-

adjusted. Significance levels and more details on the statistical tests are indicated in the figure captions.

REFERENCES

1. Bray, F., Laversanne, M., Sung, H., Ferlay, J., Siegel, R.L., Soerjomataram, I., and Jemal, A. (2024). Global cancer statistics 2022: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries. CA Cancer J Clin *74*, 229-263. 10.3322/caac.21834.

2. Akimoto, N., Ugai, T., Zhong, R., Hamada, T., Fujiyoshi, K., Giannakis, M., Wu, K., Cao, Y., Ng, K., and Ogino, S. (2021). Rising incidence of early-onset colorectal cancer - a call to action. Nat Rev Clin Oncol *18*, 230-243. 10.1038/s41571-020-00445-1.

3. Cremolini, C., Loupakis, F., Antoniotti, C., Lupi, C., Sensi, E., Lonardi, S., Mezi, S., Tomasello, G., Ronzoni, M., Zaniboni, A., et al. (2015). FOLFOXIRI plus bevacizumab versus FOLFIRI plus bevacizumab as first-line treatment of patients with metastatic colorectal cancer: updated overall survival and molecular subgroup analyses of the open-label, phase 3 TRIBE study. Lancet Oncol *16*, 1306-1315. 10.1016/S1470-2045(15)00122-9.

4. Di Nicolantonio, F., Vitiello, P.P., Marsoni, S., Siena, S., Tabernero, J., Trusolino, L., Bernards, R., and Bardelli, A. (2021). Precision oncology in metastatic colorectal cancer - from biology to medicine. Nat Rev Clin Oncol *18*, 506-525. 10.1038/s41571-021-00495-z.

5. Cancer Genome Atlas, N. (2012). Comprehensive molecular characterization of human colon and rectal cancer. Nature *487*, 330-337. 10.1038/nature11252.

6. Le, D.T., Durham, J.N., Smith, K.N., Wang, H., Bartlett, B.R., Aulakh, L.K., Lu, S., Kemberling, H., Wilt, C., Luber, B.S., et al. (2017). Mismatch-repair deficiency predicts response of solid tumors to PD-1 blockade. Science. 10.1126/science.aan6733.

7. Galon, J., Costes, A., Sanchez-Cabo, F., Kirilovsky, A., Mlecnik, B., Lagorce-Pages, C., Tosolini, M., Camus, M., Berger, A., Wind, P., et al. (2006). Type, density, and location of immune cells within human colorectal tumors predict clinical outcome. Science *313*, 1960-1964. 10.1126/science.1129139.

8. Sahin, I.H., Ciombor, K.K., Diaz, L.A., Yu, J., and Kim, R. (2022). Immunotherapy for Microsatellite Stable Colorectal Cancers: Challenges and Novel Therapeutic Avenues. Am Soc Clin Oncol Educ Book *42*, 1-12. 10.1200/EDBK_349811.

9. Ros, J., Balconi, F., Baraibar, I., Saoudi Gonzalez, N., Salva, F., Tabernero, J., and Elez, E. (2023). Advances in immune checkpoint inhibitor combination strategies for microsatellite stable colorectal cancer. Front Oncol *13*, 1112276. 10.3389/fonc.2023.1112276.

10. Antoniotti, C., Rossini, D., Pietrantonio, F., Catteau, A., Salvatore, L., Lonardi, S., Boquet, I., Tamberi, S., Marmorino, F., Moretto, R., et al. (2022). Upfront FOLFOXIRI plus

bevacizumab with or without atezolizumab in the treatment of patients with metastatic
colorectal cancer (AtezoTRIBE): a multicentre, open-label, randomised, controlled, phase 2
trial. Lancet Oncol *23*, 876-887. 10.1016/S1470-2045(22)00274-1.

11. Hackl, H., Charoentong, P., Finotello, F., and Trajanoski, Z. (2016). Computational genomics
tools for dissecting tumour-immune cell interactions. Nat Rev Genet *17*, 441-458.
10.1038/nrg.2016.67.

12. Finotello, F., Rieder, D., Hackl, H., and Trajanoski, Z. (2019). Next-generation computational
tools for interrogating cancer immunity. Nat Rev Genet *20*, 724-746. 10.1038/s41576-019-
0166-7.

13. Angelova, M., Charoentong, P., Hackl, H., Fischer, M.L., Snajder, R., Krogsdam, A.M.,
Waldner, M.J., Bindea, G., Mlecnik, B., Galon, J., and Trajanoski, Z. (2015). Characterization
of the immunophenotypes and antigenomes of colorectal cancers reveals distinct tumor escape
mechanisms and novel targets for immunotherapy. Genome biology *16*, 64. 10.1186/s13059-
015-0620-6.

14. Charoentong, P., Finotello, F., Angelova, M., Mayer, C., Efremova, M., Rieder, D., Hackl, H.,
and Trajanoski, Z. (2017). Pan-cancer immunogenomic analyses reveals genotype-
immunophenotype relationships and predictors of response to checkpoint blockade. Cell Rep
18, 248-262. 10.1101/056101.

15. Salcher, S., Sturm, G., Horvath, L., Untergasser, G., Kuempers, C., Fotakis, G., Panizzolo, E.,
Martowicz, A., Trebo, M., Pall, G., et al. (2022). High-resolution single-cell atlas reveals
diversity and plasticity of tissue-resident neutrophils in non-small cell lung cancer. Cancer cell
40, 1503-1520 e1508. 10.1016/j.ccell.2022.10.008.

16. Oliveira, G., Stromhaug, K., Klaeger, S., Kula, T., Frederick, D.T., Le, P.M., Forman, J.,
Huang, T., Li, S., Zhang, W., et al. (2021). Phenotype, specificity and avidity of antitumour
CD8(+) T cells in melanoma. Nature *596*, 119-125. 10.1038/s41586-021-03704-y.

17. He, X., Wang, L., Riedel, H., Wang, K., Yang, Y., Dinu, C.Z., and Rojanasakul, Y. (2017).
Mesothelin promotes epithelial-to-mesenchymal transition and tumorigenicity of human lung
cancer and mesothelioma cells. Molecular cancer *16*, 63. 10.1186/s12943-017-0633-8.

18. Francart, M.E., Vanwynsberghe, A.M., Lambert, J., Bourcy, M., Genna, A., Ancel, J., Perez-
Boza, J., Noel, A., Birembaut, P., Struman, I., et al. (2020). Vimentin prevents a miR-
dependent negative regulation of tissue factor mRNA during epithelial-mesenchymal
transitions and facilitates early metastasis. Oncogene *39*, 3680-3692. 10.1038/s41388-020-
1244-1.

19.   Kubala, M.H., Punj, V., Placencio-Hickok, V.R., Fang, H., Fernandez, G.E., Sposto, R., and
467   DeClerck, Y.A. (2018). Plasminogen Activator Inhibitor-1 Promotes the Recruitment and
468   Polarization of Macrophages in Cancer. Cell Rep *25*, 2177-2191 e2177.
469   10.1016/j.celrep.2018.10.082.

470   20.   Ji, P., Diederichs, S., Wang, W., Boing, S., Metzger, R., Schneider, P.M., Tidow, N., Brandt,
471   B., Buerger, H., Bulk, E., et al. (2003). MALAT-1, a novel noncoding RNA, and thymosin
472   beta4 predict metastasis and survival in early-stage non-small cell lung cancer. Oncogene *22*,
473   8031-8041. 10.1038/sj.onc.1206928.

474   21.   Qi, L., Liu, F., Zhang, F., Zhang, S., Lv, L., Bi, Y., and Yu, Y. (2018). lncRNA NEAT1
475   competes against let-7a to contribute to non-small cell lung cancer proliferation and
476   metastasis. Biomed Pharmacother *103*, 1507-1515. 10.1016/j.biopha.2018.04.053.

477   22.   Schubert, M., Klinger, B., Klunemann, M., Sieber, A., Uhlitz, F., Sauer, S., Garnett, M.J.,
478   Bluthgen, N., and Saez-Rodriguez, J. (2018). Perturbation-response genes reveal signaling
479   footprints in cancer gene expression. Nature communications *9*, 20. 10.1038/s41467-017-
480   02391-6.

481   23.   Jiang, G., Shi, L., Zheng, X., Zhang, X., Wu, K., Liu, B., Yan, P., Liang, X., Yu, T., Wang,
482   Y., and Cai, X. (2020). Androgen receptor affects the response to immune checkpoint therapy
483   by suppressing PD-L1 in hepatocellular carcinoma. Aging *12*, 11466-11484.
484   10.18632/aging.103231.

485   24.   Garcia-Alonso, L., Holland, C.H., Ibrahim, M.M., Turei, D., and Saez-Rodriguez, J. (2019).
486   Benchmark and integration of resources for the estimation of human transcription factor
487   activities. Genome research *29*, 1363-1375. 10.1101/gr.240663.118.

488   25.   Holland, C.H., Tanevski, J., Perales-Paton, J., Gleixner, J., Kumar, M.P., Mereu, E., Joughin,
489   B.A., Stegle, O., Lauffenburger, D.A., Heyn, H., et al. (2020). Robustness and applicability of
490   transcription factor and pathway analysis tools on single-cell RNA-seq data. Genome Biol *21*,
491   36. 10.1186/s13059-020-1949-z.

492   26.   Peng, S.L. (2008). Foxo in the immune system. Oncogene *27*, 2337-2344.
493   10.1038/onc.2008.26.

494   27.   Jiang, P., Zhang, Y., Ru, B., Yang, Y., Vu, T., Paul, R., Mirza, A., Altan-Bonnet, G., Liu, L.,
495   Ruppin, E., et al. (2021). Systematic investigation of cytokine signaling activity at the tissue
496   and single-cell levels. Nat Methods *18*, 1181-1191. 10.1038/s41592-021-01274-5.

497   28.   Yang, W.C., Hwang, Y.S., Chen, Y.Y., Liu, C.L., Shen, C.N., Hong, W.H., Lo, S.M., and
498   Shen, C.R. (2017). Interleukin-4 Supports the Suppressive Immune Responses Elicited by
499   Regulatory T Cells. Front Immunol *8*, 1508. 10.3389/fimmu.2017.01508.

29. Dimitrov, D., Türei, D., Garrido-Rodriguez, M., Burmedi, P.L., Nagai, J.S., Boys, C., Ramirez Flores, R.O., Kim, H., Szalai, B., Costa, I.G., et al. (2022). Comparison of methods and resources for cell-cell communication inference from single-cell RNA-seq data. Nat. Commun. 13, 3224. 10.1038/s41467-022-30755-0.

30. Pylaeva, E., and Jablonska, J. (2023). Sellhi neutrophils: a key to triumph in cancer therapy. *Signal Transduction and Targeted Therapy* **8**, 421 10.1038/s41392-023-01671-6.

31. Marzio, R., Mauël, J., and Betz-Corradin, S. (1999). CD69 and regulation of the immune function. Immunopharmacology and Immunotoxicology 21, 565–582. 10.3109/08923979909007126.

32. Del Prete, A., Martínez-Muñoz, L., Mazzon, C., Toffali, L., Sozio, F., Za, L., Bosisio, D., Gazzurelli, L., Salvi, V., Tiberio, L., et al. (2017). The atypical receptor CCRL2 is required for CXCR2-dependent neutrophil recruitment and tissue damage. Blood 130, 1223–1234. 10.1182/blood-2017-04-777680.

29. Motz, G.T., and Coukos, G. (2011). The parallel lives of angiogenesis and immunosuppression: cancer and other tales. Nat Rev Immunol *11*, 702-711. 10.1038/nri3064.

30. La Fleur, L., Botling, J., He, F., Pelicano, C., Zhou, C., He, C., Palano, G., Mezheyeuski, A., Micke, P., Ravetch, J.V., et al. (2021). Targeting MARCO and IL37R on Immunosuppressive Macrophages in Lung Cancer Blocks Regulatory T Cells and Supports Cytotoxic Lymphocyte Function. Cancer research *81*, 956-967. 10.1158/0008-5472.CAN-20-1885.

31. Yin, Y., Xu, X., Tang, J., Zhang, W., Zhangyuan, G., Ji, J., Deng, L., Lu, S., Zhuo, H., and Sun, B. (2018). CD97 Promotes Tumor Aggressiveness Through the Traditional G Protein-Coupled Receptor-Mediated Signaling in Hepatocellular Carcinoma. Hepatology *68*, 1865-1878. 10.1002/hep.30068.

32. Shojaei, F., Scott, N., Kang, X., Lappin, P.B., Fitzgerald, A.A., Karlicek, S., Simmons, B.H., Wu, A., Lee, J.H., Bergqvist, S., and Kraynov, E. (2012). Osteopontin induces growth of metastatic tumors in a preclinical model of non-small lung cancer. J Exp Clin Cancer Res *31*, 26. 10.1186/1756-9966-31-26.

33. Zhang, J., Takahashi, K., Takahashi, F., Shimizu, K., Ohshita, F., Kameda, Y., Maeda, K., Nishio, K., and Fukuchi, Y. (2001). Differential osteopontin expression in lung cancer. Cancer Lett *171*, 215-222. 10.1016/s0304-3835(01)00607-3.

34. Meng, J., Jiang, Y.Z., Zhao, S., Tao, Y., Zhang, T., Wang, X., Zhang, Y., Sun, K., Yuan, M., Chen, J., et al. (2022). Tumor-derived Jagged1 promotes cancer progression through immune evasion. Cell Rep *38*, 110492. 10.1016/j.celrep.2022.110492.

533   35.   Mikucki, M.E., Fisher, D.T., Matsuzaki, J., Skitzki, J.J., Gaulin, N.B., Muhitch, J.B., Ku,
534   A.W., Frelinger, J.G., Odunsi, K., Gajewski, T.F., et al. (2015). Non-redundant requirement
535   for CXCR3 signalling during tumoricidal T-cell trafficking across tumour vascular
536   checkpoints. Nature communications *6*, 7458. 10.1038/ncomms8458.

537   36.   Choi, H.Y., Jo, H., Zhao, X., Hoadley, K.A., Newman, S., Holt, J., Hayward, M.C., Love,
538   M.I., Marron, J.S., and Hayes, D.N. (2021). SCISSOR: a framework for identifying structural
539   changes in RNA transcripts. Nature communications *12*, 286. 10.1038/s41467-020-20593-3.

540   37.   Charoentong, P., Finotello, F., Angelova, M., Mayer, C., Efremova, M., Rieder, D., Hackl, H.,
541   and Trajanoski, Z. (2017). Pan-cancer Immunogenomic Analyses Reveal Genotype-
542   Immunophenotype Relationships and Predictors of Response to Checkpoint Blockade. Cell
543   Rep *18*, 248-262. 10.1016/j.celrep.2016.12.019.

544   38.   Chen S, Chen M. Schwann cells promote the migration and invasion of colorectal cancer cells
545   via the activated NF-κB/IL-8 axis in the tumor microenvironment. Front Oncol. 2022 Nov
546   17;12:1026670. doi: 10.3389/fonc.2022.1026670.

547   39.   Wellenstein, M.D., and de Visser, K.E. (2018). Cancer-Cell-Intrinsic Mechanisms Shaping the
548   Tumor Immune Landscape. Immunity *48*, 399-416. 10.1016/j.immuni.2018.03.004.

549   40.   Patel, A.J., Richter, A., Drayson, M.T., and Middleton, G.W. (2020). The role of B
550   lymphocytes in the immuno-biology of non-small-cell lung cancer. Cancer immunology,
551   immunotherapy : CII *69*, 325-342. 10.1007/s00262-019-02461-2.

552   41.   Li, T., Li, X., Zamani, A., Wang, W., Lee, C.N., Li, M., Luo, G., Eiler, E., Sun, H., Ghosh, S.,
553   et al. (2020). c-Rel Is a Myeloid Checkpoint for Cancer Immunotherapy. Nat Cancer *1*, 507-
554   517. 10.1038/s43018-020-0061-3.

555   42.   Fridman, W.H., Meylan, M., Petitprez, F., Sun, C.M., Italiano, A., and Sautès-Fridman, C.
556   (2022). B cells and tertiary lymphoid structures as determinants of tumour immune contexture
557   and clinical outcome. Nat Rev Clin Oncol *19*, 441-457. 10.1038/s41571-022-00619-z.

558   43.   Dong, Z.Y., Zhang, J.T., Liu, S.Y., Su, J., Zhang, C., Xie, Z., Zhou, Q., Tu, H.Y., Xu, C.R.,
559   Yan, L.X., et al. (2017). EGFR mutation correlates with uninflamed phenotype and weak
560   immunogenicity, causing impaired response to PD-1 blockade in non-small cell lung cancer.
561   Oncoimmunology *6*, e1356145. 10.1080/2162402x.2017.1356145.

562   44.   Pinto, R., Petriella, D., Lacalamita, R., Montrone, M., Catino, A., Pizzutilo, P., Botticella,
563   M.A., Zito, F.A., Del Bene, G., Zonno, A., et al. (2019). KRAS-Driven Lung Adenocarcinoma
564   and B Cell Infiltration: Novel Insights for Immunotherapy. Cancers (Basel) *11*.
565   10.3390/cancers11081145.

566   45.   Leader, A.M., Grout, J.A., Maier, B.B., Nabet, B.Y., Park, M.D., Tabachnikova, A., Chang,
567   C., Walker, L., Lansky, A., Le Berichel, J., et al. (2021). Single-cell analysis of human non-
568   small cell lung cancer lesions refines tumor classification and patient stratification. Cancer
569   Cell *39*, 1594-1609 e1512. 10.1016/j.ccell.2021.10.009.

570   57.   Madissoon, E., Wilbrey-Clark, A., Miragaia, R.J., Saeb-Parsy, K., Mahbubani, K.T.,
571   Georgakopoulos, N., Harding, P., Polanski, K., Huang, N., Nowicki-Osuch, K., et al. (2019).
572   scRNA-seq assessment of the human lung, spleen, and esophagus tissue stability after cold
573   preservation. Genome Biol *21*, 1. 10.1186/s13059-019-1906-x.

574   58.   Reyfman, P.A., Walter, J.M., Joshi, N., Anekalla, K.R., McQuattie-Pimentel, A.C., Chiu, S.,
575   Fernandez, R., Akbarpour, M., Chen, C.I., Ren, Z., et al. (2019). Single-Cell Transcriptomic
576   Analysis of Human Lung Provides Insights into the Pathobiology of Pulmonary Fibrosis. Am
577   J Respir Crit Care Med *199*, 1517-1536. 10.1164/rccm.201712-2410OC.

578   59.   Vieira Braga, F.A., Kar, G., Berg, M., Carpaij, O.A., Polanski, K., Simon, L.M., Brouwer, S.,
579   Gomes, T., Hesse, L., Jiang, J., et al. (2019). A cellular census of human lungs identifies novel
580   cell states in health and in asthma. Nature medicine *25*, 1153-1163. 10.1038/s41591-019-
581   0468-5.

582   60.   Adams, T.S., Schupp, J.C., Poli, S., Ayaub, E.A., Neumark, N., Ahangari, F., Chu, S.G.,
583   Raby, B.A., DeIuliis, G., Januszyk, M., et al. (2020). Single-cell RNA-seq reveals ectopic and
584   aberrant lung-resident cell populations in idiopathic pulmonary fibrosis. Sci Adv *6*, eaba1983.
585   10.1126/sciadv.aba1983.

586   61.   Habermann, A.C., Gutierrez, A.J., Bui, L.T., Yahn, S.L., Winters, N.I., Calvi, C.L., Peter, L.,
587   Chung, M.I., Taylor, C.J., Jetter, C., et al. (2020). Single-cell RNA sequencing reveals
588   profibrotic roles of distinct epithelial and mesenchymal lineages in pulmonary fibrosis. Sci
589   Adv *6*, eaba1972. 10.1126/sciadv.aba1972.

590   62.   Travaglini, K.J., Nabhan, A.N., Penland, L., Sinha, R., Gillich, A., Sit, R.V., Chang, S.,
591   Conley, S.D., Mori, Y., Seita, J., et al. (2020). A molecular cell atlas of the human lung from
592   single-cell RNA sequencing. Nature *587*, 619-625. 10.1038/s41586-020-2922-4.

593   63.   Picelli, S., Björklund, Å., Faridani, O.R., Sagasser, S., Winberg, G., and Sandberg, R. (2013).
594   Smart-seq2 for sensitive full-length transcriptome profiling in single cells. Nat Methods *10*,
595   1096-1098. 10.1038/nmeth.2639.

596   64.   Klein, A.M., Mazutis, L., Akartuna, I., Tallapragada, N., Veres, A., Li, V., Peshkin, L., Weitz,
597   D.A., and Kirschner, M.W. (2015). Droplet barcoding for single-cell transcriptomics applied
598   to embryonic stem cells. Cell *161*, 1187-1201. 10.1016/j.cell.2015.04.044.

599 65. Macosko, E.Z., Basu, A., Satija, R., Nemesh, J., Shekhar, K., Goldman, M., Tirosh, I., Bialas,
600   A.R., Kamitaki, N., Martersteck, E.M., et al. (2015). Highly Parallel Genome-wide Expression
601   Profiling of Individual Cells Using Nanoliter Droplets. Cell *161*, 1202-1214.
602   10.1016/j.cell.2015.05.002.

603 66. Maier, B., Leader, A.M., Chen, S.T., Tung, N., Chang, C., LeBerichel, J., Chudnovskiy, A.,
604   Maskey, S., Walker, L., Finnigan, J.P., et al. (2020). A conserved dendritic-cell regulatory
605   program limits antitumour immunity. Nature *580*, 257-262. 10.1038/s41586-020-2134-y.

606 67. Xin, J., Mark, A., Afrasiabi, C., Tsueng, G., Juchler, M., Gopal, N., Stupp, G.S., Putman, T.E.,
607   Ainscough, B.J., Griffith, O.L., et al. (2016). High-performance web services for querying
608   gene and variant annotation. Genome Biol *17*, 91. 10.1186/s13059-016-0953-9.

609 68. Xu, C., Lopez, R., Mehlman, E., Regier, J., Jordan, M.I., and Yosef, N. (2021). Probabilistic
610   harmonization and annotation of single-cell transcriptomics data with deep generative models.
611   Mol Syst Biol *17*, e9620. 10.15252/msb.20209620.

612 69. Gayoso, A., Lopez, R., Xing, G., Boyeau, P., Valiollah Pour Amiri, V., Hong, J., Wu, K.,
613   Jayasuriya, M., Mehlman, E., Langevin, M., et al. (2022). A Python library for probabilistic
614   analysis of single-cell omics data. Nat Biotechnol *40*, 163-166. 10.1038/s41587-021-01206-w.

615 70. Luecken, M.D., Buttner, M., Chaichoompu, K., Danese, A., Interlandi, M., Mueller, M.F.,
616   Strobl, D.C., Zappia, L., Dugas, M., Colome-Tatche, M., and Theis, F.J. (2022).
617   Benchmarking atlas-level data integration in single-cell genomics. Nat Methods *19*, 41-50.
618   10.1038/s41592-021-01336-8.

619 71. Lopez, R., Regier, J., Cole, M.B., Jordan, M.I., and Yosef, N. (2018). Deep generative
620   modeling for single-cell transcriptomics. Nat Methods *15*, 1053-1058. 10.1038/s41592-018-
621   0229-2.

622 72. Wolf, F.A., Angerer, P., and Theis, F.J. (2018). SCANPY: large-scale single-cell gene
623   expression data analysis. Genome Biol *19*, 15. 10.1186/s13059-017-1382-0.

624 73. Becht, E., McInnes, L., Healy, J., Dutertre, C.A., Kwok, I.W.H., Ng, L.G., Ginhoux, F., and
625   Newell, E.W. (2018). Dimensionality reduction for visualizing single-cell data using UMAP.
626   Nat Biotechnol. 10.1038/nbt.4314.

627 74. Traag, V.A., Waltman, L., and van Eck, N.J. (2019). From Louvain to Leiden: guaranteeing
628   well-connected communities. Sci Rep *9*, 5233. 10.1038/s41598-019-41695-z.

629 75. Schupp, J.C., Adams, T.S., Cosme, C., Jr., Raredon, M.S.B., Yuan, Y., Omote, N., Poli, S.,
630   Chioccioli, M., Rose, K.A., Manning, E.P., et al. (2021). Integrated Single-Cell Atlas of
631   Endothelial Cells of the Human Lung. Circulation *144*, 286-302.
632   10.1161/CIRCULATIONAHA.120.052318.

76.   Sikkema, L., Strobl, D., Zappia, L., Madissoon, E., Markov, N.S., Zaragosi, L., Ansari, M.,
Arguel, M., Apperloo, L., Bécavin, C., et al. (2022). An integrated cell atlas of the human lung
in health and disease. bioRxiv. 10.1101/2022.03.10.483747.

77.   Buttner, M., Ostner, J., Muller, C.L., Theis, F.J., and Schubert, B. (2021). scCODA is a
Bayesian model for compositional single-cell data analysis. Nat Commun *12*, 6876.
10.1038/s41467-021-27150-6.

78.   Love, M.I., Huber, W., and Anders, S. (2014). Moderated estimation of fold change and
dispersion for RNA-seq data with DESeq2. Genome Biol *15*, 550. 10.1186/s13059-014-0550-
8.

79.   Squair, J.W., Gautier, M., Kathe, C., Anderson, M.A., James, N.D., Hutson, T.H., Hudelle, R.,
Qaiser, T., Matson, K.J.E., Barraud, Q., et al. (2021). Confronting false discoveries in single-
cell differential expression. Nature communications *12*, 5692. 10.1038/s41467-021-25960-2.

80.   Ignatiadis, N., Klaus, B., Zaugg, J.B., and Huber, W. (2016). Data-driven hypothesis
weighting increases detection power in genome-scale multiple testing. Nat Methods *13*, 577-
580. 10.1038/nmeth.3885.

81.   Seabold, S., and Perktold, J. (2010). Statsmodels: Econometric and Statistical Modeling with
Python. Proceedings of the Python in Science Conference. 10.25080/majora-92bf1922-011.

82.   Efremova, M., Vento-Tormo, M., Teichmann, S. A., & others. (2020). CellPhoneDB:
Inferring cell–cell communication from combined expression of multi-subunit ligand–receptor
complexes. Nat. Protoc. 15, 1484–1506. 10.1038/s41596-020-0292-x.

83.   Jin, S., Guerrero-Juarez, C. F., Zhang, L., & others. (2021). Inference and analysis of cell-cell
communication using CellChat. Nat. Commun. 12, 1088. 10.1038/s41467-021-21246-9.

84.   Noël, F., Massenet-Regad, L., Carmi-Levy, I., & others. (2021). Dissection of intercellular
communication using the transcriptome-based framework ICELLNET. Nat. Commun. 12,
1089.10.1038/s41467-021-21244-x.

85.   Shao, X., Liao, J., Li, C., Lu, X., Cheng, J., & Fan, X. (2021). CellTalkDB: A manually
curated database of ligand-receptor interactions in humans and mice. Brief. Bioinform. 22,
bbaa269.10.1093/bib/bbaa269.

86.   Sun, D., Guan, X., Moran, A.E., Wu, L.Y., Qian, D.Z., Schedin, P., Dai, M.S., Danilov, A.V.,
Alumkal, J.J., Adey, A.C., et al. (2022). Identifying phenotype-associated subpopulations by
integrating bulk and single-cell sequencing data. Nat Biotechnol *40*, 527-538.
10.1038/s41587-021-01091-3.

Figure 1:

Figure 2

A

Figure 3

Figure 4