Single-cell integration and multi-modal profiling reveals phenotypes and spatial organization of neutrophils in colorectal cancer

Valentin Marteau[1,20], Niloofar Nemati[1,20], Kristina Handler[2,20], Deeksha Raju[2], Erika Kvalem Soto[1], Georgios Fotakis[1], Sandro Carollo[1], Nina Boeck[1], Alessia Rossi[1], Alexander Kirchmair[1], Alexandra Scheiber[3], Arno Amann[3], Andreas Seeber[3], Elisabeth Gasser[4], Steffen Ormanns[5], Michael Günther[5], Agnieszka Martowicz[3,6], Zuzana Loncova[1,19], Giorgia Lamberti[1], Marie Theres Eling[14], Lena Horvath[3], Hassan Fazilaty[7], Tomas Valenta[7,10], Gregor Sturm[1,9], Dietmar Rieder[1], Sieghart Sopper[3], Andreas Pircher[3], Patrizia Stoitzner[11], Paul Ziegler[12], Markus Tschurtschenthaler[13], Florian Huemer[16], Daniel Neureiter[15], Richard Greil[16], Lukas Weiss[16,17], Marieke Ijsselsteijn[18], Noel F.C.C. de Miranda[18], Dominik Wolf [3], Isabelle C. Arnold[2,8,21]*, Stefan Salcher[3,21]*, Zlatko Trajanoski[1,21,22]*

[1]Biocenter, Institute of Bioinformatics, Medical University of Innsbruck, Austria

[2]Institute of Experimental Immunology, University of Zurich, Switzerland

[3]Department of Internal Medicine V, Haematology & Oncology, Comprehensive Cancer Center Innsbruck and Tyrolean Cancer Research Institute, Medical University of Innsbruck, Austria

[4]Department of Visceral, Transplant and Thoracic Surgery, Medical University Innsbruck, Austria

[5]Innpath, Tirol Kliniken, Medical University Innsbruck, Austria

[6]Tyrolpath Obrist Brunhuber GmbH, Zams, Austria

[7]Department of Molecular Life Sciences, University of Zürich, Switzerland

[8]Comprehensive Cancer Center Zürich, Zürich, Switzerland

[9]Boehringer Ingelheim International Pharma GmbH & Co KG, Biberach, Germany

[10]Institute of Molecular Genetics of the Czech Academy of Sciences, Prague, Czech Republic

[11]Department of Dermatology, Venereology and Allergology, Medical University of Innsbruck, Austria

[12]Senckenberg Institute of Pathology, Goethe University Frankfurt, Germany

[13]Translational Cancer Research and Institute of Experimental Cancer Therapy, Klinikum rechts der Isar, School of Medicine & Health, Technical University of Munich, Germany

[14]Department of Therapeutic Radiology and Oncology, Medical University Innsbruck, Austria

[15]Institute of Pathology, Paracelsus Medical University, Salzburg, Austria

[16]Department of Internal Medicine III, Paracelsus Medical University, Salzburg, Austria

[17]Austrian Breast & Colorectal Cancer Study Group (ABCSG), Vienna, Austria

[18]Department of Pathology, Leiden University Medical Center, The Netherlands

[19]Present address: Institute of Pathology, Neuropathology and Molecular Pathology, Medical University of Innsbruck, Austria

[20]These authors contributed equally

[21]These authors contributed equally

37   [22]Lead contact

38   *Correspondence: stefan.salcher@i-med.ac.at, zlatko.trajanoski@i-med.ac.at

SUMMARY

Precision oncology including immunotherapy with checkpoint blockers for patients with colorectal cancer (CRC) continues to lag behind other cancer types. The immune composition of the tumor microenvironment (TME) has a major impact on the therapeutic response and clinical outcome. Here, we comprehensively characterize the TME at the single-cell level by first building a large-scale atlas that integrates 4.27 million single cells from 1,670 samples and then complementing the atlas by profiling single cells with low mRNA content. The analysis of the atlas allows refined tumor classification into four immune phenotypes: immune desert, B cell enriched, T cell enriched, and myeloid enriched subtypes. Within the myeloid compartment we uncover distinct subpopulations of neutrophils that acquire new functional properties in blood and in the TME, including antigen-presenting capabilities. Further, spatial multimodal single-cell profiling reveals that neutrophils are organized in clusters within distinct multi-cellular niches. Finally, using an orthotopic mouse model we show that cancer-derived systemic signals modify neutrophil production in the bone marrow, providing evidence for tumor-induced granulopoiesis. Our study provides a high-resolution resource for CRC TME and suggests novel therapeutic strategies targeting neutrophils.

KEYWORDS

Single-cell sequencing, spatial single-cell profiling, single-cell atlas, cellular niches

INTRODUCTION

Colorectal cancer (CRC) remains the most common gastrointestinal cancer and the second leading cause of cancer-related death according to the GLOBOCAN data[1]. The growing incidence in some countries reflects a modification in lifestyle and its consequences related with "Westernization" such as diet, obesity, physical inactivity, and alcohol consumption. Similarly, "Westernization" of diets and lifestyle have been attributed to the alarming growing incidence of early-onset CRCs (individuals <50 years of age)[2]. Despite the improvement of early detection screening policies and the development of many novel treatments, the median overall survival (OS) for patients with metastatic CRC (mCRC) is ~30 months[3]. Progress in targeted therapy for mCRC has been limited, relative to other solid cancers like lung cancer or melanoma[4]. The genetic heterogeneity[5] as well as paucity of drugs targeting common driver genes, pose considerable challenges for developing precision oncology approaches for patients with CRC. Moreover, CRC seems to be refractory to therapy with immune checkpoint inhibitors (ICI) with the notable exception of 4% of metastatic patients with tumors characterized by microsatellite instability (MSI) or *POLE* proofreading mutations[6]. This is somehow paradoxical since CRCs are known to be under immunological control, as we have shown in the past[7]. Thus, there is an urgent need to explore new immunomodulatory strategies for the vast majority of CRC patients with microsatellite stable (MSS) tumors. Strategies that convert the "immune cold" tumor microenvironment (TME) to "immune hot" could sensitize cancers to ICI therapy. A number of clinical trials with drug combinations have been carried out to overcome resistance to immunotherapy in MSS CRC[8,9]. The benefits of using a dual combination of ICIs (CTLA-4 and PD-1 blockade) or adding ICIs to chemotherapy appear to be very limited[8,10]. Combination strategies that exploit synergistic potential of ICI and targeted drugs showed increase of the anti-tumor activity, however, the objective response rates were disappointingly low[8]. Thus, to address this unmet clinical need a better understanding of the immunosuppressive TME and its heterogeneity is of utmost importance.

The intrinsic complexity of the interaction of the two interwoven systems, the tumor and the immune system, poses considerable challenges and requires comprehensive approaches to interrogate cancer immunity during tumor initiation and progression, and following therapeutic modulation thereof. A number of established and novel high-throughput technologies enable the generation of the necessary data and thereby provide the basis for mechanistic understanding. Specifically, next-generation sequencing (NGS) of cancer samples provide large data sets that can be mined for immunologically relevant parameters (see our reviews[11,12]). Using these tools we were able to chart the TME in CRC[13] and developed The Cancer Immunome Atlas[14]. While these attempts uncovered the enormous heterogeneity of MSS CRC and revealed unique cancer cell and TME subtypes with prognostic and therapeutic relevance, the advent of single-cell RNA sequencing (scRNA-seq) technologies opened fundamental new avenues to dissect cancer heterogeneity. We recently developed a novel computational approach to dissect the cellular diversity in the TME at high resolution by integrating expression profiles

of single cells from multiple studies using cutting-edge artificial intelligence (AI) methods[15]. Similar

effort was recently carried out in the context of colorectal cancer[16], enabling stratification of patients.

However, limited by the sampling bias of the commonly used 10x Genomics platform, this atlas is

lacking neutrophils. Neutrophils are the most abundant cells circulating in the human body and are

emerging as important regulators of cancer[17–20]. Neutrophils are highly abundant in intratumoral as well

as stromal regions in CRC[18] and co-localize with CD8$^+$ T cells[19]. Neutrophils exhibit remarkable

plasticity, show different pro-tumor and anti-tumor functions[20], and can acquire anti-tumor phenotype

in response to immunotherapy[21]. The importance of neutrophil-T cell interactions is supported by the

observation that CD4$^+$ PD-1$^+$ T cells in granulocyte-dominated cell neighborhoods are associated with

favorable prognosis in CRC[22]. Hence, deep characterization of neutrophils in the TME, in normal tissue,

and blood from CRC patients would provide novel insights into the myeloid compartment and the

myeloid-lymphoid cell-cell interactions.

Here, we first integrated CRC single-cell RNA sequencing datasets spanning 4.27 million cells from

650 patients and 1,670 samples. We then carried out single-cell profiling of samples from CRC patients

using optimized protocol and platform that efficiently recovers low mRNA-content immune cells. The

analyses refined tumor classification into four immune phenotypes. Within the myeloid compartment

we uncovered distinct subpopulations of neutrophils that acquire new functional properties in the TME

including antigen-presenting capabilities (HLA-DR$^+$ neutrophils). Furthermore, spatial multimodal

single-cell profiling revealed that neutrophils are organized in clusters within distinct multi-cellular

niches. Finally, using an orthotopic mouse model we show that cancer-derived systemic signals modify

neutrophil production in the bone marrow, providing evidence for tumor-induced granulopoiesis.

RESULTS

Generation of a large-scale CRC single-cell atlas

Based on our previous approach[15] we compiled CRC related scRNA-seq data from 48 studies and 76 datasets comprising 4.27 million cells from 650 patients and 1,670 samples (Figure 1A). This comprehensive CRC single-cell atlas integrates curated and quality-checked transcriptomic data from publicly available studies, complemented with extensive metadata curation, and encompasses the full spectrum of disease progression, from normal colon to polyps, primary tumors, and metastases, covering both early and advanced stages of CRC (see STAR Methods, Figures 1B-C and Supplementary Figure S1A-E). Study characteristics are summarized in Table S1. In total, the atlas includes transcriptomic data from 487 CRC patients (median: 58 years [22-91]), 99 patients with polyps (median: 60 years [40-75]), and 64 control individuals (median: 57 years [23-79]). Of the 487 CRC patients, 254 had colon cancer (COAD), 63 rectal cancer (READ), and 170 were not otherwise specified. Of these, 89 were classified as microsatellite instable (MSI), 268 as microsatellite stable (MSS), and 130 had an unknown microsatellite status. For disease stage 41 patients were classified as TNM stage I, 121 as stage II, 171 as stage III, and 110 as stage IV, while the stage was not reported for 44 patients. CRC samples include tissue of the primary tumor (n=674) or CRC liver metastasis (n=175) that were obtained by surgical resection. Among the control samples, 284 were derived from distant non-malignant tissue of colon tumor patients, of which 220 have a patient-matched tumor sample. 186 non-tumor control samples as well as 116 polyp and 90 matched normal samples were obtained from patients undergoing colonoscopy. Further, 103 blood samples of CRC patients as well as 24 samples derived from tumor affected lymph nodes, with another 18 non-tumor affected lymph node samples were included. Of the control patients none had a history of inflammatory bowel diseases.

Although neutrophils play a prominent role in cancer biology, they are underrepresented in most available scRNA-seq datasets generated with the widely used 10x Genomics technology[23] (Figure S1 F). To address this, we generated a scRNA-seq dataset from peripheral blood, adjacent normal tissue, and tumor tissue of 12 CRC patients using the BD Rhapsody platform and our optimized protocol[24] that allows capture of cells with very low mRNA content[23]. The data was then integrated into the final high-resolution single-cell atlas for CRC. Overall, the atlas integrates 4,264,946 single cells, which are annotated to 12 coarse cell type identities and 62 major cell subtypes or cell states (e.g. dividing cells) based on previously established canonical single-cell signatures including 1,146,196 epithelial cells, 1,552,108 T/NK cells, 804,042 B/Plasma cells, 385,589 myeloid cells, and 33,9767 stromal and endothelial cells (Figure 1D). Cancer cells (in total 600,740 cells, from primary tumor and metastatic tissue) in the atlas showed high heterogeneity in their transcriptomic profiles (Figure 1B). Due to the large patient cohort, we were able to apply high-resolution cancer cell classification based on their specific marker gene expression signatures (Figure S1D). We divided the following main clusters: 1) colonocyte-like (*FABP1, CEACAM7, ANPEP, SI*), 2) BEST4 (BEST4, *OTOP2, SPIB*), Goblet-like

(*MUC2, FCGBP, ZG16, ATOH1*), 3) Crypt-like (*LGR5, SMOC2*), and 4) TA-like (*TOP2A, UBE2C,*

PCLAF, HELLS, TK1) (Figure 1D, Figure S1C). There are highly mitotic/proliferative clusters (*LGR5,*

TOP2A, MKI67) of both COAD and READ that resemble highly aggressive and invasive cancer cells.

In addition, *EPCAM* expressing cells in the blood of CRC patients were annotated as circulating cancer

cells.

With a total of 2,741,739 immune cells, this single-cell atlas enables high-resolution classification of

immune cell types across both innate and adaptive subsets (Figure 1C, Figure S1A-C). These include T

cells - CD4 (*CD4*), CD8 (*CD8A*), and T regulatory cells (*FOXP3*) - as well as NK cells (*NKG7,*

TYROBP). B cells are represented as naïve (*IGHD, TCL1A*), memory (*CD27,CD44*), and activated

(*CD69, CD83*) subsets, while plasma cells (*JCHAIN*) express immunoglobulins such as IgA (*IGHA1*),

IgG (*IGHG1*), and IgM (*IGHM*). The myeloid lineage encompasses monocytes (*VCAN, CD14*),

macrophages (*C1QA, C1QB*), mast cells (*TPSB2, CPA3*), and dendritic cells, including classical

(*CLEC9A, CD1C*) and plasmacytoid subsets (*IL3RA, GZMB*), as well as eosinophils (*CLC, CCR3*) and

neutrophils (*CXCR2, FCGR3B*). Specialized populations such as gamma-delta T cells (*TRDV1, TRGC1*)

and innate lymphoid cells (ILCs; *IL4I1, KIT*) are also depicted, along with immune cell developmental

stages, such as progenitors (*CD24*, *AXL*) and stem-like cells (*LEF1*), and cycling states (*MKI67, CDK1*).

Single-cell composition of the TME highlights distinct immune infiltration patterns in primary

CRC tumors.

Utilizing the high-resolution atlas, we stratified 284 CRC untreated patients with primary tumor samples

based on the infiltration patterns of their tumor microenvironment (TME). The high number of patients

enabled us to perform unsupervised clustering of batch-corrected cell-type fractions, revealing four

major immune clusters with distinct subclusters (Figure 2A and Supplementary Figure 2A): 1) immune

deserted tumors (ID; i.e. no significant immune cell infiltration but high cancer cell fraction); 2) the

subtype of tumors with B cell dominance (B; B cell, plasma IgA and IgG cell); 3) the subtype of tumors

with myeloid dominance (M; macrophage/monocyte); and 4) subtype of tumors with T cell dominance

(T; CD8$^+$, CD4$^+$, T regulatory cells). Using logistic regression, we did not find any association of the

four different patient strata to tumor stages (early I/II *vs.* advanced III/IV), sex, or other features.

However, patients in the M (p=0.013) or B (p=0.089) subcluster were associated with a higher

likelihood of being in an early tumor stage (I/II). In addition, the CD8 (p=0.026) subcluster was

associated with a higher likelihood of the tumor being MSI compared to MSS.

In order to identify tumor-cell based TME imprinting characteristics, we next analyzed differentially

enriched pathways[25] in the cancer cells of each immune infiltration type. The immune B and M subtype

showed significant downregulation (FDR < 0.1) of the androgen pathway (Figure 2B).

Integration of bulk RNA-seq data reveals genotype-immune phenotype associations

The majority of the scRNA-seq studies lack both comprehensive cancer genotype information and survival data. In contrast, the TCGA reference dataset includes this information, alongside bulk RNA-sequencing data. Using the computational method SCISSOR[26], we integrated bulk RNA-seq data from TCGA with single-cell transcriptomic signatures to assess the relationship between specific genotypes and survival outcomes in CRC (Figure 3A). We evaluated the association of cell type transcriptomic signatures derived from the atlas with genotype and survival data from the TCGA reference dataset, which includes 629 patients (UICC I-IV)[5]. Building on our previous work that identified genomic features influencing immune phenotypes in a pan-cancer study using bulk RNA-seq data[14], we leveraged the high resolution of the single-cell CRC atlas for a more detailed analysis of these determinants. We therefore analyzed the cell type transcriptomic signatures and their association with survival in 629 patients from the TCGA cohort. Overall, the presence of high levels of IgA$^+$ plasma cells, CD4$^+$ T cells CD8$^+$ T cells and B cells showed a prominent association with improved survival in CRC patients. Conversely, CRC patients with tumors enriched in Schwann cells were associated with worst survival outcomes, as also reported previously[30]. Notably, high infiltration of various endothelial cells and fibroblasts also emerged as strong negative predictors of survival (Figure 3A). Mutations in *BRAF* and *KRAS* were associated with distinct immune infiltrate patterns (Figure 3B-C). For example, CRC patients with *BRAF* mutations exhibited high infiltration of cDC1 and cDC2, as well as increased levels of IgM$^+$ plasma cells. Similarly, IgM$^+$ plasma cell infiltration was also elevated in patients with *KRAS* mutations. Conversely, *BRAF* mutations were linked to low levels of Plasma IgA infiltration, while *KRAS* mutations were associated with high Plasma IgA infiltration. Additionally, MSI and MSS status displayed opposing cell type infiltration patterns (Figure S3A and S3B).

Finally, we identified gene signatures based on the patient immune infiltration groups by selecting the top 15 differentially expressed genes. These gene signatures were then used to conduct a survival analysis utilizing data from the TCGA COAD-READ cohorts (n = 629). The Kaplan-Meier curves in Figure 2D demonstrate that higher expression of the gene signatures of the B (p-value = 0.0015), M (p-value = 0.0096), and T (p-value = 0.00078) subgroups is significantly correlated with improved overall survival.

Comprehensive characterization of neutrophils reveals highly diverse and plastic phenotypes

We then carried out comprehensive characterization of the data we generated using the BD Rhapsody platform. The obtained scRNA-seq dataset comprises 126,991 single cells categorized into 23 main cell types based on specific marker genes, including 55,245 epithelial/tumor cells, 4,428 stromal/endothelial cells, and 65,782 immune cells, including 30,924 neutrophils (Figure 4A). Compositional analysis using

scRNA-seq and flow cytometry showed high concordance when comparing blood, normal, and tumor

tissue samples (Figure 4B). As expected, neutrophils were the most abundant leukocyte subtype in blood

samples (60–70%), followed by T lymphocytes and monocytes (Figure 4C). Normal-adjacent tissue was

dominated by epithelial cells, while tumor tissue contained a markedly higher proportion of immune

cells. Notably, neutrophils comprised ~20% of all cells and ~40% of leukocytes in tumor tissue but were

present only in small numbers in normal-adjacent tissue (Figure 4C). Also within the granulocytes

lineage, neutrophils were the most abundant cells (26.2% of all cells) followed by eosinophils (1.2%)

and mast cells (0.4%).

We observed distinct transcriptomic differences between blood *versus* tumor-associated neutrophils

(TAN) (Figure 4D) with 2,214 genes differentially expressed (padj<0.05; Figure 4E). We evaluated

previously described marker genes distinguishing TAN from normal tissue neutrophils (NAN) across

cancers, including *OLR1*, *VEGFA*, *CXCR2*, and *PLAU*[27]. TAN markers (*OLR1*, *PLAU*, *VEGFA*) were

highly upregulated in TAN, whereas *CXCR2*, a marker of mature neutrophils, was highly expressed in

blood neutrophils but downregulated in TAN (Figure 4F). Spatial analysis of Lectin-type oxidized LDL

receptor 1 (LOX-1; encoded by *OLR1*) expression using multiplex immunofluorescence (IF) staining

demonstrated the presence of segregated LOX-1+ neutrophil cell clusters exclusively in CRC tumor

tissue, but not in normal-adjacent colonic tissue (Figure 4G; Supplementary Figure S4A) LOX-1+ cells

co-express the neutrophil marker CD15). Of note, LOX-1 has previously been identified as a marker for

TAN in lung cancer[15], indicating LOX-1 as a putative tumor-agnostic TAN marker.

Sub-clustering of neutrophils revealed three distinct blood neutrophil clusters (BN1-BN3) and four TAN

clusters (TAN1-TAN4) (Figure 4H). Trajectory analysis indicated a progressive transition from BN1 to

BN3 and subsequently to TAN1 (Figure 4I), demonstrating that neutrophil migration from the

bloodstream into tissue is accompanied by distinct transcriptomic changes. From TAN1, neutrophils

diverged into two trajectories: one leading to a putative pro-tumor TAN phenotype (TAN4)

characterized by high *IL1A* expression, a key driver of emergency myelopoiesis that fuels inflammation

and immunosuppression in the TME[28], and another toward a phenotype defined by high expression of

antigen-presentation-associated markers, such as *HLA-DRA* (TAN2/TAN3) (Figure 4J). Neutrophils

with antigen-presenting capabilities, referred to as hybrid neutrophils[27], have previously been identified

in early-stage lung cancer and are associated with an anti-tumor phenotype[15,29]. Interestingly, the

expression of *HLA-DRA* in blood or tissue-resident neutrophils was significantly higher in early-stage

CRC compared to advanced-stage CRC (Figure 4K), suggesting that these antigen-presenting

neutrophils may be diminished as the tumor progresses to more advanced stages, potentially contributing

to immune evasion. Finally, we validated HLA-DR protein expression using flow cytometry and

confirmed that the expression was markedly higher in TAN compared to both blood neutrophils and

NAN (Figure 4L).

Spatial single-cell multi-modal profiling reveals neutrophil clusters within multicellular niches

Increasing evidence indicates that immune cells in the TME occupy spatially distinct areas or patches characterized by specific cellular compositions and discrete histological properties: so-called cellular neighborhoods[22] or cell niches. Evidence for this type of segregated myeloid populations in CRC was recently shown using a limited immunofluorescence (IF) panel[30] as well as in our data (Figure 4G). In order to quantify spatial niches we analyzed the TME of MSS CRC patients using a spatial transcriptomic technique with 380 genes and sub-cellular resolution (10x Genomics Xenium). The spatial single-cell transcriptomic profiling revealed distinct spatial architecture and cell type composition in both, center of the tumor and the invasive margin (Supplementary Figure 5A). Spatial co-localization of T cell and B cell clusters pinpointed tertiary lymphoid structures in these samples (Figure S5B-C). Neutrophils were the other immune cell type that was organized as clusters in the tumor and the invasive margin (Figure 5B). This spatial organization of neutrophils was confirmed using immunohistochemistry (IHC) stainings of the tissue from the same patients (Supplementary Figure 5D).

In order to quantify spatial multicellular interactions we used the graph deep-learning method NicheCompass that utilizes existing knowledge of inter- and intracellular interaction pathways to identify distinct cellular niches[31]. At the resolution level of two niches, we identified cancer and non-cancer niches (Supplementary Figure 5E), whereas at the resolution of three niches we identified cancer, boundary, and stroma niches (Supplementary Figure 5F). At the resolution level of ten we quantified five cancer-cell dominated cell niches (CN1-CN5), two fibroblast-dominated niches (CN6-7) and a plasma-cell-enriched niche (CN8) comprising the tumor-stroma boundary regions, a stromal niche (CN9) and an immune-cell dominated niche that identified tertiary lymphoid structures (CN10) (Figure 5A). The cancer-cell dominated niche CN5 was enriched with macrophages and neutrophils (Figure 5B). Analysis of cancer cells within this niche revealed elevated expression of the chemokines *CXCL1*, *CXCL3*, *CXCL10*, and *CXCL11*, along with the immunomodulatory enzyme *IDO1* (Figure 5C). Neutrophils may amplify *IDO1* expression in cancer cells[32], contributing to an immunosuppressive milieu. The identified chemokines, which recruit neutrophils through CXCR2 and CXCR3 signaling[33], likely play a central role in driving cancer-cell mediated localized neutrophil aggregation and cluster formation.

We then used an orthogonal method based on imaging mass-cytometry (IMC) and profiled FFPE samples from patients with stage III colon cancer treated with adjuvant chemotherapy (CAPOX) and for whom relapse and overall survival data were available (n=30). We previously developed and evaluated a panel of antibodies for IMC on FFPE samples for a comprehensive overview of the TME and cancer-immune cell interactions[34], including lineage and functional immune cell markers, surrogates of cancer cell states (proliferation, apoptosis) and structural markers (epithelium, stroma, vessels). Following data-driven identification of single-cell phenotypes segmentation, and image analysis, we quantified the

densities of major cell types (Figure 5D). The spatial protein profiling revealed similar cellular composition of the TME as identified using single-cell mRNA profiling. Given the limited number of features, quantification of cell niches based on protein expression was not possible. We therefore used a statistical approach to define niches and applied cell neighborhood analysis by defining cell nuclei and associating polygons (Voronoi diagram) to each nucleus[34] (see Methods), thereby allowing cells of different sizes and distances to be assessed as neighbors. We used a permutation approach[34] to identify pairwise interactions between cell phenotypes that occurred more or less frequently than expected by chance. This spatial information revealed neutrophil clusters similar to the clusters quantified using transcriptomic data (Figure 5E). It appears that the neutrophil clusters are more frequent in the tumor compared to the invasive margin, albeit the differences in this cohort were not significant (Figure 5H). Analyses of the survival data showed that the neutrophil clusters in both regions are associated with worse prognosis (Figure 5I).

Overall, the multimodal profiling revealed that neutrophils are not randomly distributed but are localized in spatial clusters. Moreover, neutrophil clusters are situated within distinct multicellular niches composed of different fractions of cancer cells, endothelial cells, and other immune cells.

Mouse neutrophils recapitulate human neutrophil phenotypes in various compartments

To carry out functional studies and define underlying mechanisms, we deployed an orthotopic mouse model based on tumor organoid transplantation (Figure 6A-B). AKPS organoids – harboring the four most common CRC-driving mutations (*Apc*-LOF (loss of function), *p53*-LOF, *Kras*-GOF (gain of function), *Smad4*-LOF) – were successfully engrafted following colonoscopy-guided injection into the colonic mucosa of C57BL/6JRj mice[35] (Supplementary Figure 6A). Tumors developed within six to eight weeks (Figure 6C) and progressively expanded through the colonic layers toward the peritoneal cavity (Supplementary Figure 6B), mirroring some of the invasive characteristics of late-stage human CRC. Notably, neutrophil infiltration was significantly higher in the tumor compared to the healthy colon (Figure 6D-F). Interestingly, there was already an increase of neutrophils within the blood of AKPS-tumor bearing compared to naive mice (Figure 6G) suggesting a systemic effect during tumor development. Analysis of neutrophil activation markers like Siglec-F, Cd11b, CD80 and PD-L1 further showed a progressive activation of neutrophils from the blood to the tumor (Figure 6G-J) while MHC-II proteins were upregulated in tumor and colon tissue compared to blood neutrophils (Figure 6L). Overall, the results confirm the validity of the orthothopic mouse model for human CRC.

Cross-species analyses reveal tumor-induced granulopoiesis in CRC

In order to carry out cross-species analyses of mouse and human CRC neutrophils, we performed scRNA-seq on CD45-enriched leukocytes isolated from tumors and blood of mice six weeks post-AKPS-injection. Colons and blood of naïve (control) mice were further included as controls.

Additionally CD45-enriched leukocytes were isolated from the bone-marrow of both tumor and non-
tumor bearing mice. Neutrophils from all tissues and phenotypes were extracted and integrated (Figures
7A andB). Clustering analysis identified 10 clusters with distinct transcriptional patterns (Figures 7C
and D) and a high concordance of BN and TAN marker genes derived from our human neutrophil dataset
(Supplementary Figure 7A), especially within the colon and tumor specific clusters. We then performed
pseudotime trajectory analysis, which showed a progressive transition of neutrophils from the bone-
marrow to the blood and colonic tissues (Figures 7E and F). Differential gene expression (DEG) analysis
between tumor and healthy bone-marrow and blood revealed that neutrophils already changed their
transcriptional phenotype prior to their recruitment into the tumor tissue (Figures 7G and H). Gene set
enrichment analysis (GSEA) of KEGG terms showed changes in the metabolic phenotypes
(Supplementary Figure 7C). Neutrophils from the tumor bone-marrow upregulated multiple genes
encoding for ribosomal proteins (*Rps2, Rpl32, Rpsa, Rps27l*) and proteins involved in cell proliferation[36]
(*Ccnb2, Mcm4, Pcna, Fen1*). Overall, the results from the single-cell profiling suggest tumor-induced
granulopoiesis which is likely driven by secreted factors.
350
To provide further evidence for tumor-induced granulopoiesis we carried out scRNA-seq profiling using
blood samples from healthy volunteers and compared the expression profiles of neutrophils with the
expression profiles of blood neutrophils from the CRC patients (Figure 7I). The results showed profound
changes in the expression with 2287 genes differentially expressed (padj<0.05; Figure 7J). GSEA on
KEGG terms indicated changes in the metabolic phenotypes. We then carried out functional experiments
with neutrophils isolated from blood of healthy donors and treated the neutrophils with supernatants
from a colorectal adenocarcinoma cell line (CaCO) and a non-small cell lung cancer cell line (SW900).
Seahorse analysis showed different O_2 consumption rates (Figure 7K), indicating context-dependent
metabolic rewiring of the neutrophils.
360
361 Together, these findings provide evidence of tumor-induced granulopoiesis and metabolic rewiring
362 which is likely driven by secreted factors from the TME.

DISCUSSION

We built a large-scale atlas of single-cell transcriptomes of CRC through integration of 4.27 million cells from 650 patients and 1,670 samples representing 7 billion expression values. We then complemented the atlas with single-cell data enriched with immune cells with low-mRNA content including neutrophils, and carried out multi-modal spatail profiling at the single-cell level of the TME in CRC. Our unique resource provides novel biological findings and has important translational relevance.

First, we provide a high-resolution view of the TME in CRC with 61 major cell types/states and show different cell type composition patterns including more precise functional transcriptomic classification of malignant epithelial cells. The single-cell composition of the TME in CRC enabled refined tumor classification and patient stratification into four immune phenotypes: immune deserted, myeloid, B cell, and T cell subtypes. These findings may have important implications for enabling cancer immunotherapy in CRC. For example, combination therapies that target both, myeloid cells and lymphoid cells could represent an immunotherapeutic strategy to treat M subtypes as shown recently in a melanoma mouse model[37]. Similarly, given the heterogeneity of the intratumoral B cells and the importance of tertiary lymphoid structures[38], further analysis of the B cell subtypes might open promising therapeutic avenues by additional refined B cell targeting.

Second, we characterized for the first time neutrophil subsets in various compartments in CRC patients and show remarkable diversity and plasticity. Of particular interest for cancer immunotherapy is the TAN phenotype with immunogenic antigen-presenting feature. This observation implies acquisition of antigen presenting-like properties by neutrophils at the tumor site, as previously reported[39]. Such conversion of neutrophils to antigen presenting cells may elicit anti-tumor immunity and has recently been shown in a murine model[40].

Third, multi-modal spatial profiling revealed that neutrophils build clusters of cells within distinct multicellular niches. Similar neutrophil clustering has been recently observed in the context of wound healing using a zebrafish model[41]. The driving force for this type of myeloid structures remains unknown and encourages further studies in the context of CRC and possibly other solid cancers.

Finally, using an orthotopic mouse model and cross-species analyses, we show that tumor-derived signals modulate the neutrophils in the bone-marrow, implicating tumor-induced granulopoiesis. Expression and functional data indicate metabolic rewiring in the bone marrow and imprinting the neutrophil cell fate via progenitors.

Beyond these biological insights, the results from this study have also important translational implications. As shown here, TANs can acquire antigen presenting properties and such conversion of

abundant neutrophils to antigen-presenting cells could overcome the limitations of the low abundance of cross-presenting DCs. Additionally, our study highlights CRC as a systemic disease that requires therapeutic strategies beyond the primary disease site. Such a strategy was recently shown for lung cancer[42]. However, as we show here, the disease context has profound effects on the neutrophil phenotypes and hence, requires further disease-specific investigations.

In conclusion, we provide a CRC atlas with single-cell resolution as well as a web portal that enables interactive exploration of the dataset through cell-x-gene (https://crc.icbi.at) that allows visualization of metadata and gene expression. The biological insights we present here and future discoveries arising from the exploitation of the high-resolution CRC atlas could provide the basis for developing combination therapies for CRC patients who are not sufficiently responding to immune checkpoint blockers.

ACKNOWLEDGMENTS

This work was supported by the European Research Council (grant agreement No 786295 to ZT), by the Austrian Science Fund (FWF) (project I3978 to ZT). ZT is a member of the German Research Foundation (DFG) project TRR 241(INF). The results published here are in part based upon data generated by the TCGA Research Network: https://www.cancer.gov/tcga. VM was funded by the Austrian Science Fund (FWF) 10.55776/DOC82. ICA was funded through an Eccellenza Professorial Fellowship from the Swiss National Science Foundation PCEFP3_187021 and through a Tandem Grant (2022 call) of the ISREC Foundation.

     FIGURE LEGENDS

Figure 1. Building a large-scale single-cell atlas for CRC. (**A**) Summary of the data integration and

analysis workflow. (**B**) Number of samples for each type. (**C**) Overview of the cell fractions in different

tissues. (**D**) Overview of the CRC atlas and the epithelial, and stromal/endothelial components depicted

as UMAP plots. (**E**) Overview of the CRC atlas and the immune components depicted as UMAP plots.

Figure 2. Tumor immune phenotypes in CRC. (**A**) Patient characteristics and stratification of the

tumor immune phenotypes. (**B**) Differential activation of PROGENy cancer pathways across different

cell types in the TME. Heatmap colors indicate the deviation from the overall mean, independent of

tumor histology and stage. White dots indicate significant interactions at different false-discovery-rate

(FDR) thresholds. Only cytokine signatures with an FDR < 0.1 in at least one patient group are shown.

(**C**) Kaplan-Meier curves of the survival analysis according to the patient immune infiltration groups

(B, desert, M, T), performed using the molecular data of the TCGA COAD-READ cohorts (n = 461).

Figure 3. Association of cellular composition and distinct genotypes and survival in the TCGA

data. (**A-E**) *SCISSOR* analysis relating phenotypic information from bulk RNA-seq data from TCGA

with single cells. UMAP plots indicate the position of cells positively (blue) or negatively (red)

associated with mutation or better survival. A log2 ratio > 0 indicates a positive association with

mutation, or better survival, respectively. Shown are cell-types with a log2 ratio significantly different

from 0 at an FDR < 0.01 (paired Wilcoxon test). (**A**) Association of cellular composition with overall

survival. (**B**) Association of cellular composition with BRAF mutation in CRC patients (n = 136). (**C**)

Association of cellular composition with KRAS mutation in CRC patients (n = 489).

Figure 4. Neutrophil characterization in CRC. (**A**) UMAP of the generated CRC scRNA-seq dataset

(n = 12 patients; matched blood, tumor tissue, and normal-adjacent tissue) colored by cell type. (**B**) Cell-

type fractions obtained by scRNA-seq and by flow cytometry. (**C**) Neutrophil fractions (as percentage

of leukocytes) obtained by flow cytometry (n = 12 patients; matched blood, tumor tissue, and normal-

adjacent tissue). (**D**) UMAP of neutrophils from blood and tumor tissue analyzed by scRNA-seq. (**E**)

Volcano plot of the gene expression from blood and tumor-associated neutrophils (TANs). (**F**)

Expression of *OLR1*, *PLAU*, *CXCR2*, and *VEGFA* in neutrophils from blood and tumor tissue, analyzed

by scRNA-seq. (**G**) Selected multiplex immunofluorescence (mpIF) staining of LOX-1 (pink), pan-

cytokeratin (turquoise), and DAPI (blue) in tumor tissue and matched normal-adjacent colorectal cancer

(CRC) tissue. Scale bar: 100 μm. See also Figure S4. (**H**) Top 10 markers for BN1-BN3 and TAN1-

TAN4 sub-clusters. The marker gene quality is reflected by the area under the receiver operator

characteristics curve (AUROC; 1 = marker gene perfectly distinguishes the respective cluster from other

clusters in all patients; AUROC 0.5 = no better than random). (**I**) Partition-based graph abstraction

(PAGA) based on RNA velocities, projected on the UMAP plot of neutrophils from blood and tumor

tissue. Edges represent cell-type transitions called by PAGA. (**J**) Expression of *HLA-DRA* in neutrophils

from blood and tumor tissue, analyzed by scRNA-seq. **K**) Expression of *HLA-DRA* in neutrophils from

blood, tumor tissue, and normal tissue.. (**L**) Quantification of HLA-DR expression on neutrophils by

flow cytometry across blood, normal-adjacent tissue, and tumor tissue. Each dot represents the mean

value for an individual patient (n = 12).

Figure 5. Spatial single-cell multi-modal profiling. (**A**) Cell niches identified from imaging-based

spatial transcriptomics profiling of five CRC tumor sections (CN1-4: Intratumoral niches, CN5:

Intratumoral neutrophil-enriched niche, CN6-8: Tumor-stroma boundary, CN9: Stroma, CN10: TLS).

(**B**) Cell type composition of each niche. (**C**) Top 5 genes upregulated in cancer cells of CN5. (**D**)

Cellular neighborhoods of neutrophil clusters. (**E**) Fraction of all neutrophils and of neutrophils in

clusters within all cell types per niche.(**F**) UMAP of the imaging mass-cytometry data using 40 markers

and samples from 30 CRC patients.(**G**) Neutrophil niches identified using the imaging mass-cytometry

data at the tumor core and invasive margin. (**H**) Violin plot of the neutrophil niches by location. (**I**)

Kaplan-Meier curves for the neutrophil niches identified at the tumor core and invasive margin.

Fig. 6. Neutrophils in a murine orthotopic organoid transplant model for CRC. (**A-B**) Mouse AKPS

tumor organoid cells harbouring the four most common CRC driver mutations (APC^{KO}, $KRAS^{G12D}$,

$P53^{KO}$, $SMAD4^{KO}$) were injected intra-rectally into 8- to 10-week-old C57BL/6J males. At the study

endpoint (4-6-weeks post-injection), blood, tumor and adjacent colonic tissue were harvested for flow

cytometric analysis. (**C**) Survival curve 4 weeks post-injection. (**D**) Neutrophil numbers per milligram

tissue and frequency among live CD45+ leukocytes (right). (**E**) Representative plots of neutrophil

frequencies among CD45+ cells, as determined by flow cytometry. (**F**) Neutrophil frequencies among

CD45+ cells in the blood of naïve versus AKPS-injected mice. (**G**) Frequencies and (**H**) representative

flow-cytometry plots of Siglec-F-expressing neutrophils among total neutrophils in blood, colon and

tumour of AKPS-bearing mice. Mean Fluorescence Intensity (MFI) of cell-surface CD11b integrin (**I**)

or CD80 and PDL1 (**J**) expression in neutrophils. (**K**) MFI (left) and frequencies of MHC II expression

among neutrophils. *, $P < 0.05$; **, $P < 0.01$; ***, $P < 0.001$; ****, $P < 0.0001$; as calculated by two-

tailed unpaired Student's t-test for comparison of two groups (**D, F**) or by one-way ANOVA for

statistical comparison of more than two groups (**G-K**). n=5, medians are displayed.

Figure 7. Neutromap of mouse CRC. Uniform manifold approximation and projection (UMAP) of

integrated neutrophils from tumor, colon, blood and bone marrow of AKPS tumor bearing and non-

tumor bearing mice. Cells are colored by (**A**) phenotype, (**B**) tissue type (colon is a combination of

colon, adjacent colon, tumor, and disseminated tumor) and (**C**) clusters generated via the Seurat

clustering algorithm. (**D**) Heatmap depicting the scaled average gene expression of the top 10

differentially expressed genes (DEGs) per cluster. Columns are clustered by tissue presence. (**E**)

Heatmap showing the scaled average gene expression by cluster of the ortholog genes representing

human neutrophil clusters. Columns are clustered by tissue presence. (**F**) UMAP representing RNA

velocity with cells colored by tissue type. The arrows indicate the velocity flow. (A-F) n = pool of 6

mice each condition and tissue. (**G**) Volcano plot of gene expression data from bone-marrow

neutrophils from tumor-bearing mice and control mice. (**H**) Volcano plot of gene expression data from

blood neutrophils from tumor-bearing mice and control mice. (**I**) UMAP of blood neutrophils from CRC

patients and controls. (**J**) Volcano plot of gene expression data from blood neutrophils from CRC

patients and controls. (**K**) OCR (oxygen consumption rate) calculated from AUC of XF neutrophil

activation assay kinetic traces of healthy human neutrophils activated with PMA, after 24h culture in

SW900 supernatant, or CaCo2 supernatant or RPMI control medium. The data shown are the mean and

SD, tested for differences using t test, n=3 biological replicates.

527   **Key resources table**

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies		
Antibodies list (human samples)		See Table S2
Antibodies list (mouse samples)		See Table S2

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Biological samples		
Resected colorectal tumors and adjacent healthy colorectal tissue (CRC) samples, naive/untreated	Tirol Kliniken Innsbruck	N/A
Whole blood samples from patients undergoing colorectal tumor resections	Tirol Kliniken Innsbruck	N/A
Human neutrophils from healthy donors	This study	N/A
Chemicals, peptides, and recombinant proteins		
StemPro hESC SFM	Gibco	Cat# A1000701
Penicillin-Streptomycin	Sigma-Aldrich	Cat# P4333
Primocin	InvivoGen	Cat# ant-pm-2
Fetal bovine serum (FBS), sterile-filtered	Sigma-Aldrich	Cat# F7524
Dulbecco's phosphate-buffered saline, without calcium and magnesium chloride, sterile-filtered, DPBS	Sigma-Aldrich	Cat# D8537
DMEM, high glucose, pyruvate, no glutamine	Gibco	Cat# 21969-035
L-glutamine, 200 mM, 100 mL	Sigma-Aldrich	Cat# G7513
Liberase DH, research grade	Roche	Cat# 5401054001
DNAse I, grade II, from bovine pancreas	Roche	Cat# 10104159001
OptiMEM	Gibco	Cat# 31985062
Matrigel	Corning	Cat# 354277
RPMI-1640 Medium	Gibco	Cat# 11875093
HEPES	Sigma-Aldrich	Cat# H0887
Collagenase type IV	Sigma-Aldrich	Cat# C5138
Collagenase type VIII	Sigma-Aldrich	Cat# C2139
Bovine Serum Albumin, BSA	Roche	Cat# 10738328103
Penicillin/streptomycin	Sigma-Aldrich	Cat# P0781
EDTA	SIgma-Aldrich	Cat# P0781
Percoll	Cytiva	Cat# 17089101
RPMI-1640 Medium	Sigma-Aldrich	Cat# R0883
Seahorse XF RPMI assay medium pack	Agilent Technologies	Cat# 103681
Seahorse XFp Cell Culture Miniplates	Agilent Technologies	Cat# 103022
Seahorse XFp FluxPak	Agilent Technologies	Cat# 103022
Poly-L-lysine	Sigma-Aldrich	Cat# P4832
Rotenone/Antimycin A	Agilent Technologies	Cat# 103010
Phorbol 12-myristate 13-acetate (PMA)	Fisher Scientific	Cat# J63916.MCR
Critical commercial assays		
Agencourt AMPure XP magnetic beads	Beckman Coulter	Cat# A63880
BD Human Single-Cell Multiplexing Kit	BD Biosciences	Cat# 633781
BD Rhapsody Cartridge Kit	BD Biosciences	Cat# 633733
BD Rhapsody cDNA Kit	BD Biosciences	Cat# 633773
BD Rhapsody Enhanced Cartridge Reagent Kit	BD Biosciences	Cat# 664887
BD Rhapsody WTA Amplification Kit	BD Biosciences	Cat# 633801
BD Stain Buffer (FBS)	BD Biosciences	Cat# 554656
High Sensitivity D1000 Reagents	Agilent	Cat# 5067-5585
High Sensitivity D1000 Screen Tape	Agilent	Cat# 5067-5584
Qubit dsDNA HS Assay Kit	Invitrogen	Cat# Q32854
BD Pharm Lyse Lysing Buffer	BD Biosciences	Cat# 555899
Calcein AM	Invitrogen	Cat# C1430
Draq7	BD Pharmingen	Cat# 564904
Anti-CD45 microbeads	Miltenyi Biotec	Cat# 130-052-301

REAGENT or RESOURCE	SOURCE	IDENTIFIER
BD Mouse Single-Cell Multiplexing Kit	BD Biosciences	Cat# 633793
BD Rhapsody Whole Transcriptome Analysis Amplification Kit	BD Biosciences	Cat# 633801
Qubit dsDNA HS Kit	Thermo Fisher Scientific	Cat# Q32851
TapeStation 4200 HS D5000 tapes and reagents	Agilent Technologies	Cat# 5067-5592
NovaSeq 6000 SP Reagent Kit v1.5 (100 cycles)	Illumina	Cat# 20028401
Opal 6-plex Detection Kits	Akoya Biosciences	Cat# NEL821001KT
BOND Epitope Retrival 1	Leica Biosystems	Cat# AR9961
BOND Epitope Retrival 2	Leica Biosystems	Cat# AR9640
BOND Dewax Solution	Leica Biosystems	Cat# AR9222
BOND Wash Solution 10x	Leica Biosystems	Cat# AR9590
Spectral DAPI	Akoya Biosciences	Cat# FP1490
Prolong Diamond Antifade	Thermo Fisher	Cat# P36961
BOND Research Detection System	Leica Biosystems	Cat# DS9455
BOND Titration Kit	Leica Biosystems	Cat# OPT9049
Xenium Human Immuno-Oncology Profiling Panel	10x Genomics	Cat# 1000654
Xenium Slides & Sample Prep Reagents	10x Genomics	Cat# 1000460
Xenium Cell Segmentation Add-on Kit	10x Genomics	Cat# 1000662
Xenium Decoding Reagents	10x Genomics	Cat# 1000461
Xenium Cassettes	10x Genomics	Cat# 1000566
Xenium Decoding Consumables	10x Genomics	Cat# 1000487
MACSxpress Whole Blood Neutrophil Isolation Kit	Miltenyi Biotec	Cat# 130-104-434
Deposited data		
MUI_Innsbruck	This study	Zenodo link
Processed input data	This study	Zenodo link
Core atlas as cell-x-gene instance and h5ad/Seurat file formats with standardized metadata according to the cell-x-gene schema	This study	https://cellxgene.cziscience.com/collections/3a844375-60ea-474b-adf1-98e76928baee
Bian_2018_Science (fastq)	SRA	PRJNA382695
Borras_2023_Cell_Discov (processed)	synapse	syn41687327
Burclaff_2022_Cell_Mol_Gastroenterol_Hepatol (processed)	SRA	PRJNA768275
Che_2021_Cell_Discov (fastq)	SRA	PRJNA725335
Chen_2021_Cell (processed)	synapse	syn26044148
Chen_2024_Cancer_Cell (processed)	GEO	PRJNA991601
Conde_2022_Science (fastq)	ENA	PRJEB51634
Elmentaite_2021_Nature (fastq)	ENA	PRJEB40273
GarridoTrigo_2023_Nat_Commun (fastq)	SRA	PRJNA886695
Giguelay_2022_Theranostics (processed)	SRA	PRJNA666199
Guo_2022_JCI_Insight (fastq)	SRA	PRJNA779978
Harmon_2023_Nat_Cancer (processed)	SRA	PRJNA863562
He_2020_Genome_Biol (fastq)	SRA	PRJNA670909
HTAPP_HTAN (processed)	synapse	syn24181445
Huang_2024_Nat_Cancer (processed)	NGDC	PRJCA007578
James_2020_Nat_Immunol (fastq)	ENA	PRJEB32912
Ji_2024_Cancer_Lett (fastq)	SRA	PRJNA915211
Ji_2024_PLoS_Genet (processed)	GEO	GSE217774
Joanito_2022_Nat_Genet (fastq/processed)	EGA	EGAD00001008555

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Khaliq_2022_Genome_Biol (processed)	GEO	PRJNA827831
Kong_2023_Immunity (processed)	Broad Single Cell Portal	SCP1884
Lee_2020_Nat_Genet (fastq/processed)	GEO	PRJNA604751
Li_2017_Nat_Genet (processed)	GEO	PRJNA323703
Li_2023_Cancer_Cell (processed)	GEO	PRJNA846169
Liu_2024_Cancer_Res (processed)	GEO	PRJNA1028966
Mazzurana_2021_Cell_Res (processed)	GEO	PRJNA630996
Parikh_2019_Nature (processed)	SRA	PRJNA477814
Pelka_2021_Cell (processed)	GEO	PRJNA738517
Qi_2022_Nat_Commun (processed)	NGDC	HRA000979
Qian_2020_Cell_Res (processed)		http://blueprint.lambrechtslab.org/
Qin_2023_Cell_Rep_Med (processed)	CNGBdb	CNP0004138
Sathe_2023_Clin_Cancer_Res (processed)		https://dna-discovery.stanford.edu/research/datasets/
Scheid_2023_J_EXP_Med (processed)	Broad Single Cell Portal	SCP1690
Smillie_2019_Cell (processed)	Broad Single Cell Portal	SCP259
Terekhanova_2023_Nature (processed)	synapse	syn53214678
Thomas_2024_Nat_Med (processed)	GEO	PRJNA849977
Tian_2023_Nat_Med (processed)	Broad Single Cell Portal	SCP2079
Uhlitz_2021_EMBO_Mol_Med (processed)	GEO	PRJNA701264
deVries_2023_Nature (processed)	GEO	PRJNA894171
Wang_2020_J_Exp_Med (fastq)	SRA	PRJNA518129
Wang_2021_Adv_Sci (fastq)	SRA	PRJNA937723
Wang_2023_Sci_Adv (processed)	GEO	PRJNA937723
Wu_2022_Cancer_Discov (processed)		http://www.cancerdiversity.asia/scCRLM
Wu_2024_Cell (processed)		http://www.pancancer.cn/neu/
Yang_2023_Front_Oncol (fastq)	SRA	PRJNA972665
Zhang_2018_Nature (processed)	GEO	PRJNA429424
Zhang_2020_Cell (processed)	figshare	https://figshare.com/articles/dataset/CRC_CD45_rda/14820318
Zheng_2022_Signal_Transduct_Target_Ther (processed)	GEO	PRJNA676119
TCGA data RNA-seq and mutation data	GDC	https://portal.gdc.cancer.gov/
Cytosig signatures	Jiang *et al.*, 2021 (PMID:34594031)	https://github.com/data2intelligence/CytoSig/
Experimental models: Cell lines		

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Vil-creER^{T2};APC^{KO};Trp53^{KO};Kras$^{G12D/WT}$Smad4^{KO} (AKPS) organoids	Dr. Tomas Valenta and Dr. Hassan Fazilaty (DMLS, University of Zürich, Basler Lab)[35]	N/A
SW900	ATCC	HTB-59
CaCo2	ATCC	HTB-37
Experimental models: Organisms/strains		
Mouse: C57BL/6J Strain #:000664	The Jackson Laboratory	RRID:IMSR_JAX:000664
Software and algorithms		
Seven Bridges - BD Rhapsody™ WTA Analysis Pipeline	BD Biosciences	v2.0
FlowJo™	BD Biosciences	v10.7
GraphPad Prism v9	GraphPad Software	www.graphpad.com
GraphPad Prism v10.4	GraphPad Software	www.graphpad .com
Cellranger	10x Genomics	v7.1.0
nf-core/scrnaseq pipeline	Philip et al., 2022	v2.4.1
nf-core/rnaseq pipeline	Philip et al., 2022	v3.12.0
Nextflow	Di Tommaso et al., 2017	v23.04.0
Singularity/Apptainer	Kurtzer et al., 2017	v1.3.1-1.el9
Nextflow workflow to reproduce this study	this study	https://github.com/icbi-lab/crc-atlas
Atlas web resource	this study	https://crc.icbi.at/
FlowJo Software	BD Life Sciences	v.10.8.0
Prism 10	GraphPad	v10
Bcl2fastq v2.20.0433	Illumina	
STAR v2.5.2b	(Dobin et al., 2023)[43]	
R v4.3	R Core Team (2021) R: A Language and Environment for Statistical Computing. R Foundation for Statistical Computing, Vienna.	https://www.R-project.org/
Seurat v5.1.0		
decontX v1.0.0	(Yang et al., 2020)[44]	
SingleR v2.4.1	(Aran et al., 2019)[45]	
Celldex v1.12.0	(Heng et al., 2008)[46]	
Pheatmap v1.0.12		
Orthogene v1.8.0		
Slingshot v2.10.0	(Street et al., 2018)[47]	
Sceasy v0.0.7		

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Fiji v1.54f	(Schindelin et al., 2012)[48]	https://imagej.net/software/fiji/downloads
QuPath v0.5.1	(Bankhead et al., 2017)[49]	https://qupath.github.io/
CellProfiler v4.2.8	(Stirling et al., 2021)[50]	https://cellprofiler.org/releases
R v4.4.2	R Core Team (2021) R: A Language and Environment for Statistical Computing. R Foundation for Statistical Computing, Vienna.	https://www.R-project.org/
Seahorse XFe Wave Software	Agilent Technolgies	www.agilent-com
Compute node CPU	Intel	Intel(R) Xeon(R) Gold 6342 CPU @ 2.80GHz (2x)
GPU node GPU	Nvidia	NVIDIA A100 80GB PCIe
Other		
50 mL centrifuge tube	Sarstedt	Cat# 62.547.254
Stainless steel surgical blades, sterile, Nr. 22	VWR International GmbH	Cat# 233-54840
pluriStrainer, 100 mm strainer, sterile	pluriSelect	Cat# 43-50100-51
CytoOne 6 cm Petri dish, uncoated	Starlab	Cat# CC7672-3359
Mediware, disposable scalpels, stainless steel, sterile (for tissue dissection)	medizinbedarf.at	Cat# I1 09
Mediware, disposable forceps, sterile (for tissue dissection)	medizinbedarf.at	Cat# H7 301
Mediware pre-powdered gloves	medizinbedarf.at	Cat# 9321
Mediware sterile drapes	medizinbedarf.at	Cat# H2 50
gentleMACS Octo Dissociator with heaters	Miltenyi Biotec	Cat# 130-096-427
gentleMACS Octo Dissociator with heaters, C tubes	Miltenyi Biotec	Cat# 130-093-237
Filter tips, 10 mL, sterile	Starlab	Cat# S1121-3810
Filter tips, 20 mL, sterile	Starlab	Cat# S1120-1810
Filter tips, 100 mL, sterile	Starlab	Cat# S1123-1840
Filter tips, 200 mL, sterile	Starlab	Cat# S1120-8810
Filter tips, 1000 mL, sterile	Starlab	Cat# S1126-7810
Serological pipette, 5 mL	Sarstedt	Cat# 86.1253.001
Serological pipette, 10 mL	Sarstedt	Cat# 86.1254.001
Serological pipette, 25 mL	Sarstedt	Cat# 86.1685.001
Serological pipette, 50 mL	Sarstedt	Cat# 86.1256.001
Water bath	N/A	N/A
Vacuum pump, Vacusafe aspiration system with integrated pump	INTEGRA Biosciences	Cat# 158320
Qubit Fluorometer	Thermo Fisher Scientific	N/A
TapeStation 4200	Agilent Technologies	N/A
NovaSeq6000	Illumina	N/A
FACSymphony A5	BD Biosciences	N/A
LSRII Fortessa	BD Biosciences	N/A
Xenium Analyzer	10x Genomics	N/A

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Seahorse XF Analyzer	Agilent Technologies	N/A

METHODS

Patient recruitment

Twelve naive tumor and matched adjacent normal tissues from CRC patients who underwent surgical resections at the Department of Surgery, Medical University of Innsbruck, Austria were obtained. Additionally, autologous whole blood samples were collected during surgery. Prior to study enrollment, all patients signed the informed consent for participation in the study. The ethical committee of the Medical University of Innsbruck approved the specimen collection and analysis of patients' data. Tumor samples subjected to imaging mass cytometry (IMC) were obtained from patients in the Salzburg cohort. These patients uniformly had R0-resected stage III colon cancer and received adjuvant CAPOX treatment. This analysis was approved by the Ethics Committee of the Provincial Government of Salzburg, Austria (415-E/2343/5-2018). Patients were otherwise unselected and treated at the Department of Internal Medicine III, Paracelsus Medical University, Salzburg, Austria.

Dissociation of normal and tumor colorectal tissue

Surgically resected human colorectal tumor and adjacent normal tissues were enzymatically and mechanically dissociated into single cells as described in detail in the following publication[51]. Briefly, after supervised sampling of tumor and adjacent normal CRC tissues from a pathologist, the specimens were transported to the lab on ice, washed, weighed and minced into small pieces (<1 mm). The tissue pieces were further enzymatically and mechanically digested by using DMEM medium (Gibco) with Liberase DH (Roche) and DNAse I (Roche/Sigma-Aldrich) on a gentleMACS dissociator with heaters (Miltenyi Biotec) at 37°C for ~45 min (selected program: 37C_h_TDK_1). The obtained single cell solutions were sieved through a 100 μm cell strainer (pluriSelect) and kept on ice.

BD Rhapsody library preparation and sequencing

Freshly isolated single-cells were immediately processed with the BD Rhapsody scRNA-seq platform (BD Biosciences). Red blood cells were removed from dissociated tissues and from whole blood samples using the BD Pharm Lyse lysing solution (BD Biosciences). Cell number and viability were assessed with the BD Rhapsody scRNA-seq platform using Calcein-AM (Invitrogen) and Draq7 (BD Biosciences). The BD Single-Cell Multiplexing Kit (BD Biosciences) was used to combine and load multiple samples (human colorectal tumor tissue, adjacent normal tissue, blood) onto a single BD Rhapsody™ cartridge (BD Biosciences). Sample-tag staining was performed according to the

manufacturer's protocol (sample-tag staining at room temperature for 20 min and washing by
centrifugation at 400 g for 5 min). Single-cell isolation in microwells (cell load: 20 min incubation at
room temperature, 20°C ± 2°C) with subsequent cell-lysis and capturing of poly-adenylated mRNA
molecules with barcoded, magnetic capture-beads was performed according to the manufacturer's
instructions. Beads were magnetically retrieved from the microwells, pooled into a single tube before
reverse transcription. Unique molecular identifiers (UMIs) were added to the cDNA molecules during
cDNA synthesis. Whole transcriptome amplification (WTA) and sample-tag sequencing libraries were
generated according to the BD Rhapsody single-cell whole-transcriptome amplification workflow. The
quantity and quality of the sequencing libraries was analyzed with the Qubit dsDNA HS (High
Sensitivity) assay kit (Invitrogen) and the 4200 TapeStation (Agilent) system. Libraries were sequenced
on the Novaseq 6000 system (Illumina) targeting a sequencing depth of 50.000 reads/cell.
573
Generation of the atlas
The data for the core atlas was from previously published studies[52–98]. The selected studies were
published between May 2017 and July 2024 and the incorporated datasets were generated with six
different sequencing platforms, including the most commonly applied 10x Chromium (10x Genomics)
as well as Smart-seq2[99], SMARTer (C1), scTrio-seq2, GEXSCOPE (Singleron), TruDrop[100] and
DNBelab C4[101]. We further integrated our own data generated with the microwell-based BD Rhapsody
scRNA-seq platform (BD Biosciences). We specifically selected studies using comparable protocols for
sample processing and data generation, such as sequencing of whole cells. We did not exclude studies
that applied flow cytometry-based cell-sorting prior to sequencing, as these incorporate relevant
information on rare cell types[102,103,104]. From non-CRC studies we exclusively included those parts of
the published data that were relevant for our atlas: from the Garrido Trigo dataset[105] we only included
colon samples, from Burclaff[106], Kong[107], Mazzurana[108], Parikh[109], Smillie, Thomas, and Scheid[110]
datasets we only used the control samples.
587
Integration of scRNA-seq datasets
Individual datasets were merged into a single AnnData object. Since genome annotations partly differed
between the datasets, we mapped gene symbols to the ensembl ids of the used annotation version and
updated the symbols to the latest version of HGNC. If gene ids were missing from a dataset, the values
were filled with zeros. Genes that were missing in more than 19 datasets (25%) were excluded
altogether.
We integrated the datasets using the scANVI algorithm[111,112], as it has been demonstrated to be one of
the top-performing methods for atlas-level integration and to scale to >1M cells[113]. Since scANVI
requires cell-type annotations for at least one of the input datasets, we manually annotated two "seed"
datasets based on unsupervised clustering as described below. Raw counts were used as input for
scANVI. Counts from plate-based assays such as Smart-seq2 were scaled by the gene length as

recommended on the scvi-tools website. The scANVI model was initialized with a pre-trained scVI

model[114], as recommended in the scvi-tools tutorial. The scVI model was trained on the 6000 most

highly variable genes as determined with scanpy's[115] *pp.highly_variable_genes* with parameters

flavor="seurat_v3" and *batch_key="dataset"*. Each sample was considered as an individual batch for

both scVI and scANVI. Other than that the algorithms were run with default parameters.

Unsupervised clustering and cell-type annotation

606   We computed UMAP embeddings[116] and unsupervised Leiden-clustering[117] with scanpy[115], based on a

607   cell-cell neighborhood graph derived from scANVI latent space. Coarse, lineage-specific clusters were

608   iteratively sub-clustered to identify cell-types at a more fine-grained resolution. Cell type clusters were

609   annotated based on previously reported marker genes[105,118,119]

Comparing cell-type abundances

Since comparing cell-type fractions between groups is challenging due to different characteristics of the

datasets and the inherent compositional nature of cell-type fractions, we applied the scCODA[120] model,

which addresses this issue. We were interested in the differences between conditions (COAD *vs.*

READ). To this end, we ran the scCODA model with the formula ~ *condition + tumor_stage + dataset*

with 500,000 iterations using "cancer cells" as the reference cell-type, where *tumor_stage* is a binary

vector classifying datasets into early (stages I-II) and advanced (stages III-IV), and dataset is a

categorical vector encoding the different datasets. The final result shows *credible effects* with a false-

discovery-rate (FDR) of 0.1.

Patient stratification

We stratified patients into immune phenotypes based on immune cell-type fractions. We selected all

patients with primary tumor samples and excluded enriched datasets. Neutrophil fractions were

excluded, since they are not appropriately captured in the majority of datasets. Cell-type fractions of

primary tumor samples were loaded into a patient × cell-type AnnData container. Dataset-specific batch-

effects were removed using a linear model as implemented in *scanpy.pp.regress_out*. Patients were

clustered using graph-based Leiden clustering with the "correlation" distance metric for computing the

neighborhood graph. Patient clusters were labeled according to their predominant cell-types. In addition

to the histological subtypes based on the annotation of the original datasets, we annotated tumor types

based on the transcriptomics data according to the most abundant cancer cell cluster.

Differential gene expression testing

We used DESeq2[121] on pseudo-bulk samples for differential expression testing which has been

demonstrated to perform well and properly correct for false discoveries[122]. For each cell-type and

patient, we summed up transcript counts for each gene. Pseudo-bulk samples consisting of fewer than

10 cells were discarded. We compared primary tumor samples from tumor *vs.* normal (*sample type*),

primary tumor samples from the patient subtypes M/B/T/desert (*group*), NANs *vs.* TANs

(*cell_type_tan_nan*), and Neutrophil clusters (BN1-3, TAN1-4), including the dataset as a covariate. For

comparisons between multiple groups, we used contrasts with sum-to-zero coding. P-values were

adjusted for multiple hypothesis testing with independent hypothesis weighting (IHW)[123].

Pathway, TF and cytokine signaling signatures

We performed pathway, transcription factor (TF), and cytokine signaling analysis on primary tumor

samples with PROGENy[25,124], and CytoSig[125], respectively. Scores were computed using the *decoupler-*

py[126] package.

Transcription factor activity was inferred from the Wald statistics of the differential expression analysis

output, using a univariate linear model and the CollecTRI[127] database. Activities were defined as

significantly activated or suppressed based on an FDR threshold of less than 0.1. The top 1,000 target

genes of the progeny model were used, as recommended for single-cell data. The cytosig signature

matrix was obtained from the *data2intelligence/CytoSig* GitHub repository and used with the scoring

function implemented in the *progeny-py* package. The methods were run with the options *num_perm=0,*

center=True, norm=True scale=True, and *min_size=5*. No permutations were used, as we perform

statistics in a separate step at the level of biological replicates. Pathway-, transcription factor-, and

cytosig scores were then compared between *condition* (tumor *vs.* normal) and patient *group* (T *vs.* B *vs.*

M *vs.* deserted) using an ordinary least-squares (OLS) linear model, as implemented in the statsmodels

package[115]. Sc[128]s were aggregated into pseudobulk samples by computing the mean of each variable

for each patient and cell-type. Samples consisting of less than 10 cells were discarded. For each variable,

we fitted a model with the formulas ~ *condition + dataset + tumor_stage* or ~ *group + dataset +*

condition + tumor_stage, respectively. Coefficients were obtained from the linear model and p-values

calculated with the f-test. P-values were adjusted for multiple testing with the Benjamini-Hochberg

procedure.

SCISSOR analysis

We used SCISSOR[129] to associate phenotypic data from bulk RNA-seq experiments with our single-cell

data. TCGA mutation and gene expression data was obtained from the GDC portal, survival data from

SCISSOR[129] was run on primary tumor cells of each patient individually according to the SCISSOR

tutorial using mutation data (logistic regression) and overall survival (cox-regression) as dependent

variables. A grid search for the alpha-parameter was performed in $2^{-i/2}$ with $i \in [24, 23, \ldots, 2]$ and a

cutoff parameter of 0.3. 62 of 674 samples with low overall cell count failed during SCISSOR's Seurat-

preprocessing step and were excluded from the subsequent analysis. For each patient and cell-type, we

computed the fraction of *scissor+ cells* (i.e. positively associated with a mutation or worse survival),

scissor- cells (i.e. negatively associated), and *neutral cells* and added a pseudo-count of 0.01. A sample

was excluded from a cell-type if it contributed ≤10 cells. For each cell-type, we computed the log2-ratio of scissor+ and scissor- cells as the mean fraction of scissor+ cells *vs.* the mean fraction of scissor-cells. Significant differences were determined by comparing the fractions of scissor+ and scissor- cells with a paired wilcoxon test with *zero_method="zsplit"* as implemented in the scipy package. P-values were Benjamini-Hochberg-adjusted and considered significant at an FDR < 0.01.

Survival analysis

Initially, gene lists were extracted from the patient groups according to their immune infiltration profiles (B, desert, M and T). The gene lists consisted of differentially expressed genes, as returned from DESeq2 for the comparisons of each immune infiltration group with the rest, that were filtered using as thresholds the adjusted p-values (< 0.05) and the logFold Change (greater than 0.5 or lower than -0.5). Consequently, the gene lists were condensed into gene signatures expression values using the GSVA (v1.50.5) R package. Once the gene signature expression values were calculated the TCGA COAD-READ cohort (n = 629) was stratified into "high" and "low" expression, using as threshold the optimal cutpoint as computed by the surv_cutpoint() function of the survminer (v0.4.9) R package. The survival analysis was carried out using the survival (v3.7-0) R package and the Kaplan-Meier curves were plotted using the ggsurvplot function of the survminer (v0.4.9) R package.

Quantification and statistical analysis

Statistical analysis was performed using the statsmodels library in Python (scRNA-seq data) or GraphPad Prism (flow cytometry and imaging data) using a linear model, t-test or wilcoxon test as appropriate. Single cell-data were aggregated into pseudobulk samples by biological replicates. Compositional analysis of cell-type fractions was performed using scCODA; survival analysis using the Kaplan-Meier formula in R. P-values for untargeted analyses (DE genes, TFs, or pathways) were FDR-adjusted. Significance levels and more details on the statistical tests are indicated in the figure captions.

Flow Cytometry staining of human samples

Cells isolated from surgically resected CRC tumor and adjacent normal tissues as well as whole blood obtained just prior surgery were stained with a cocktail of the following 20 antibodies (CD3, CD8, CD4, CD14, CD16, CD19, CD15, CD28, CD34, CD31, CD38, CD45, CD56, CD90, CD123, CD161, CD193, CD326, HLA-DR, TCRgd) to define all cell populations of interest. After washing and addition of 5 μL 7-AAD, cells were measured on a FACSymphony A5 flow cytometer (BD Biosciences). All source data of the aforementioned antibodies is provided in the key resources table, applied fluorochromes are listed in Table S2. Data were analyzed using FlowJo v10.7 software. For details of the gating strategy, see Figure S8.

Multiplex immunofluorescence of human samples

CRC tumor and tumor-adjacent tissue samples were fixed in 4% formalin and embedded in paraffin.

Three-micrometer sections were used for immunofluorescence staining. The immunofluorescence

staining on formalin-fixed paraffin-embedded (FFPE) tissue was conducted using the Opal 6-Plex

Detection Kit (Cat. No: NEL821001KT, Akoya Biosciences, Menlo Park, USA). A multiplex panel of

immune markers was established utilizing antibodies against CD15 (1:200, clone MMA, Abcam), LOX-

1/OLR1 (1:200, polyclonal, Sigma-Aldrich), and cytokeratin (1:1000, clone AE1/AE3, Dako; clone C-

11, Abcam). The staining procedure was performed using an automated staining system (BOND-RX,

Leica Biosystems). Markers were applied sequentially and paired with their respective Opal

fluorophores (CD15 – Opal650, LOX-1/OLR1 – Opal570, cytokeratin – Opal690). To visualize cell

nuclei, the tissue was stained with 4',6-diamidino-2-phenylindole (spectral DAPI, Akoya Biosciences).

The stained slides were scanned using the Mantra 2 Quantitative Pathology Workstation (Akoya

Biosciences), and representative images were acquired using the Mantra Snap software version 1.0.4.

Spectral unmixing, multispectral image analysis, and cell phenotyping were carried out using the inForm

Tissue Analysis Software version 2.4.10 (Akoya Biosciences). Pseudo-DAB staining projections were

achieved by converting fluorescent channel to a pseudo-brightfield images using InForm, where

fluorescent markers are displayed as pseudo-DAB staining and DAPI as hematoxylin.

Preparation of sections for Xenium spatial transcriptomics

Sections were prepared according to manufacturer's instructions (Tissue Preparation Guide

Demonstrated Protocol CG000578) at the Department of Anatomy, Histology and Embryology, Medical

University of Innsbruck. 5 µm sections were taken from the FFPE tissue blocks with a microtome

(Epredia, HM355S), floated in a 45°C water bath, and adhered to Xenium slides (10x Genomics).

Samples were baked at 42°C for 3 hours and stored in a desiccator at room temperature, overnight,

followed by deparaffinization and decrosslinking, according to manufacturer's instructions

(Deparaffinization & Decrosslinking Demonstrated Protocol CG000580).

10x Xenium workflow and data acquisition

Xenium samples were processed according to manufacturer's protocols (Xenium In Situ Gene

Expression with Morphology-based Cell Segmentation Staining User Guide CG0000749 and Decoding

& Imaging, User Guide CG000584). Samples were stained utilizing the predesigned Human Immuno-

Oncology panel from 10x genomics (PN 1000654), as they became available from the manufacturer.

Slides were stained with a Xenium Cell Segmentation kit according to manufacturer's instructions (User

Guide CG0000749). Briefly, probes of the immuno-oncology panel were hybridized overnight before

rolling circle amplification and native protein autofluorescence was reduced with a chemical

autofluorescence quencher. Slides were processed on a 10X Xenium Analyzer, with ROIs selected to

cover the selected regions. The Xenium Onboard Analysis pipeline v2.0.0 (10x Genomics) was run

directly on the instrument for imaging processing, cell segmentation, image registration, decoding,

deduplication, and secondary analysis, as previously described[130].

Post Xenium H&E staining

After processing tissue sections through the Xenium workflow, the same sections on the Xenium slide

were used for H&E staining, according to the manufacturer's instructions (Post-Xenium Analyzer H&E

Staining CG000613) and imaged using a PANNORAMIC slide scanners (Institute for Pathology,

Neuropathology & Molecular Pathology, Medical University of Innsbruck).

Xenium data analysis

Output files of the Xenium Onboard Analysis were further processed with scanpy[115], squidpy[131] and the

SpatialData framework[132] for region selection, quality control and calculation of spatial neighbor graphs.

We applied SpaNorm[133] to normalize transcript counts. Integration of samples was done with scvi on

the raw counts, followed by leiden clustering to define distinct cell types. We utilized tangram[134] in

cluster mode to annotate leiden clusters by mapping cell type annotations from a downsampled version

of the CRC atlas generated with scSampler[135]. NicheCompass[136] was run with OmniPath[137] and

NicheNet[138] as prior gene programs. Leiden clustering was then performed at different resolutions on

the niches latent space to identify distinct spatial niches. Neutrophil clusters were identified from the

spatial neighbor graph (neighbors=20, distance=25 µm) by first defining cluster seeds as cells of cell

type 'Neutrophil' with at least three homotypic neighbors. Cluster seeds were subsequently expanded

in five iterations to add all neutrophils adjacent to seed cells to the respective cluster. We further defined

cluster neighborhoods as all cells in the spatial neighbors graph (neighbors=20, distance=10 µm)

connected to a neutrophil cluster.

Imaging mass cytometry and data analysis

Imaging mass cytometry was performed on tissue microarrays from microsatellite-stable CRC patients

as previously described[34], using the antibody panel provided in Table S2. Images were obtained from a

total of 37 patient samples, including intratumoral regions from 27 patients and tumor margin regions

from 29 patients. Among these, both intratumoral and tumor margin regions were imaged for 19 patients.

Per sample, one or two 1000x1000µm regions of interest were ablated, depending on tissue availability.

Images were visually inspected and exported to ome.tiff using MCD viewer (standard biotools). Image

normalization, cell segmentation and phenotype identification were performed as described previously[34]

and steps were validated throughout on the original images.

Spatial analysis of granulocyte niches

To analyze the spatial organization in tumor and margin regions, we constructed Voronoi diagrams

based on the center coordinates of all detected nuclei. Each Voronoi region was then assigned the

corresponding cell type. The DBscan clustering algorithm was used to find niches consisting of at least 3 granulocytes in a maximum pairwise distance of 15 μm. For these niches all other cell types that share a Voronoi ridge (edge) with a granulocyte were searched and added as "niche members". To determine if the number of detected niches differed significantly from what would be expected by chance, a Monte Carlo simulation with 1,000 iterations was conducted. In this simulation, the cell type labels for each cell on the imaged slide was randomly assigned while maintaining the number of cells for each type constant and preserving the overall spatial arrangement. A z-score and p-value were then calculated.

Tissue Microarrays and Image Acquisition

Five tissue microarrays (TMAs) containing a total of 549 valid cores were processed and stained for CD15 and HLA-DR using immunofluorescence (IF) and six TMAs containing in total 649 valid cores were processed and stained with MPO and HLA-DR using IF. Images of the stained TMAs and whole slides were acquired in .czi format.

Image Preprocessing and core extraction

The .czi images were converted to a compatible format (.tif) for downstream analysis using Fiji (ImageJ, v1.54f). Images were carefully inspected during this step to ensure no loss of resolution in any of the RGB layers.

Individual TMA cores were extracted from the pre-processed images using QuPath (v0.5.1), an open-source software platform for digital pathology. The cores were manually annotated to exclude damaged or improperly stained regions, and only valid cores were included in subsequent analyses.

Cell Identification and Quantification

A customized image analysis pipeline was developed in CellProfiler (v4.2.8) to identify and quantify nuclei and marker-positive cells for CD15 and HLA-DR and in the second batch for MPO and HLA-DR. The pipeline consisted of the following steps:

1.	Nuclear Identification: Nuclei were segmented and identified based on fluorescence intensity.

2.	Marker Co-Localization: Positive staining for CD15, MPO and HLA-DR was identified and quantified, with nuclei serving as reference points for spatial localization and co-expression analysis.

Output: The pipeline produced quantitative data on nuclei counts, their spatial position and the number of nuclei positive for each marker, based on staining intensity to distinguish specific marker expression from background fluorescence.

Data Extraction and evaluation

Quantitative data, including total nuclei count, all CD15-positive cells, all HLA-DR-positive cells, all MPO-positive cells and co-expression counts, were exported from CellProfiler for statistical analysis. Data evaluation and statistical analyses were conducted in R (v4.4.2).

Mice

C57BL/6J (B6J, stock no. 000664) mice were obtained from The Jackson laboratory and used for injections of AKPS organoids, or kept uninjected as non-tumour bearing control mice.

All mouse experiments were performed under the approval of Cantonal Veterinary Office of Zurich, Switzerland and in accordance with Swiss guidelines. All mice were maintained on a 12h light and 12h dark schedule, and water and chow were provided ad libitum.

Colonoscopy submucosal injection of AKPS organoids

Apc-LOF, *p53*-LOF, *Kras*-GOF, *Smad4*-LOF (AKPS) tumor organoids were generated from AKP organoids harbouring a RNP-mediated Smad4 mutation introduced by CRISPR-Cas9[35]. Organoids were cultured in Matrigel and passaged every 2-3 days[139]. Colonoscopy-guided orthotopic cell transplantation was done as described previously with modifications[140]. Briefly, following mechanical dissociation, the equivalent of one Matrigel done was resuspended in 50 mL OptiMEM (Gibco) to equate to one injection in one mouse. 8-to 10-week-old males were anesthetized with isoflurane and placed facing up on a 37°C heating pad. Fecal content was flushed from the colon with warm DPBS and AKPS organoid solution was injected carefully below the mucosa using a custom needle, syringe and colonoscope. Observation of a visible injection bubble was noted as an indicator for successful injection in the correct location. Mice were then regularly monitored according to official guidelines until experimental endpoint.

Tissue dissociation of mouse samples

AKPS tumors

Tumors were collected at the study endpoint and cut into small pieces. Tissue was digested at 37 °C for 50 min in RPMI-1640 containing 15 mM HEPES (H0887 Sigma), 500 U/mL^{-1} type IV (C5138 Sigma) and type VIII collagenase, (C2139 Sigma) and 0.05 mg/mL DNase I (10104159001 Roche). Non-digested tissue was mechanically disrupted with a syringe and cell suspension was pushed through a 70 μm cell strainer. Flow-through was topped with DPBS buffer containing 2% BSA, 100 U mL/$^{-1}$ penicillin/streptomycin (P0781 Sigma) and 5 mM EDTA and cells were spun prior and resuspended prior analysis.

Blood

Blood was collected from mice post-mortem by cardiac puncture in 2% BSA 5 mM EDTA, followed by red blood cell lysis with distilled cold water for 30 sec.

Colon from naive mice and adjacent colon to AKPS tumor

Colonic tissue was collected, cleaned and opened longitudinally, prior to washing with PBS. Tissue was

cut into small pieces and washed twice while shaking for 25 min at 37 °C with HBSS 2% BSA, 100 U

mL/$^{-1}$ penicillin/streptomycin (P0781 Sigma) and 5 mM EDTA. Tissue was digested at 37 °C for 50 min

in RPMI-1640 containing 15 mM HEPES (H0887 Sigma), 500 U/mL^{-1} type IV (C5138 Sigma) and type

VIII collagenase, (C2139 Sigma) and 0.05 mg/mL DNase I (10104159001 Roche). Non-digested tissue

was mechanically disrupted with a syringe and cell suspension was pushed through a 70 µm cell strainer.

Cells were centrifuged for 8 min and layered onto a 40/80% Percoll (17089101 Cytiva) gradient (18

min, 2.100 g, 20 °C, no brake). The interphase was collected and washed in DPBS.

Bone marrow

Tibia and femur were collected and flushed with RPMI medium using a 23-gauge needle. The content

was filtered through a 40 µm strainer followed by red blood cell lysis via incubation in ice-cold distilled

water for 30 sec.

Flow cytometry staining of mouse samples

Staining

Cells were surface stained in DPBS at 4 °C for 30 minutes with fixable viability dye eFluor 780, and

combinations of the following listed antibodies (1:200, all from BioLegend; unless otherwise stated):

CD11b BV510 (M1/70, 101263), CD45 BV650 (30-F11,103151), CD80 BV605 (1:100, 16-10A1,

104729), Ly6G Percp-Cy5.5 (1A8, 127616), MHCII AF700 (M5/114.15.2,107622), PD-L1 PE-Cy7

(1:100, 10F.9G2, 124314), Siglec-F BV421 (E50-2440, 552681 BD Biosciences). Samples were

acquired on a LSRII Fortessa (BD Biosciences).

Data analysis

Analysis of flow cytometry data was performed using FlowJo Software. Relative cell frequencies, cell

counts, or Mean Fluorescent Intensities (MFIs) were subsequently plotted in graphical formats using

GraphPad Prism.

Statistical analysis

Statistical analyses were performed using GraphPad Prism. For statistical comparisons of two groups,

two-tailed unpaired t-tests were used. For comparisons of more than two groups, a one-way analysis of

variance (ANOVA) was used. Differences were considered statistically significant when $P < 0.05$.

Single-cell RNA sequencing of mouse neutrophils - experimental

CD45+ cell enrichment using magnetic beads

CD45 positive cells were enriched using anti-CD45 microbeads (130-052-301 Miltenyi Biotec)
according to the manufacturer's instructions.

Cell hashing

Samples were pooled for shared scRNAseq runs after labeling with sample tags (633793 BD Mouse
Single-Cell Multiplexing Kit). Thereby, 10^6 cells from each sample were incubated in a staining buffer
(DPBS, 1% BSA, 1% EDTA) and the respective sample tag for 20 min at room temperature (20°C ±
2°C). After incubation, cells were washed twice with staining buffer before counting. From each sample
10-20.000 cells were pooled to have a total of around 60-80.000 cells for single cell capture.

Single cell capture, library preparation and sequencing

Single cell capture was done using the BD Rhapsody Express Single-Cell Analysis System (BD
Biosciences) following the manufacturer's protocol. In short, cells were resuspended in 650 µL BD
Sample Buffer supplemented with Rnase inhibitors (1:1.000 SUPERase 20 U µL^{-1}, AM2694 Thermo
Fisher Scientific; NxGen Rnase Inhibitor 40 U µL^{-1}, 30281-2 Lucigen) and loaded on a cartridge. To
extract their RNA capture beads (v1 for tumor, disseminated and adjacent colon; enhanced beads for
colon, blood and bone marrow) were added, followed by cell lysis. After single cell capture, libraries
from the cDNA and the sample tags were prepared using the BD Rhapsody Whole Transcriptome
Analysis Amplification Kit (633801 BD Biosciences). Libraries were then quantified using a Qubit
Fluorometer (dsDNA HS kit: Q32851 Thermo Fisher Scientific) and a TapeStation 4200 (HS D500
types and reagents, 5067-5592 Agilent Technologies). Sequencing was performed on a NovaSeq6000
system (Illumina) with SP100 Reagents (100 cycles) and PhiX spike-in (20% for v1, 1% for enhanced
beads) with a depth of around 20.000 reads/cell for WTA and 500-1000 reads/cell for the sample tags.

Single-cell RNA sequencing of mouse neutrophils - data analysis

Pre-processing, normalization, clustering and data integration

Demultiplexing was done using Bcl2fastq (v2.20.0433, Illumina). FASTQ files were then further
processed using the BD Rhapsody[TM] WTA analysis pipeline (v1.12.1) on the Seven Bridge Genomics
platform with default settings. Gene mapping was done with a STAR index generated using STAR[43]
(v2.5.2b) with the gene code GRCm38 v25. R (v4.3) and Python (v3) were used for the downstream
analysis. If not specified, the Seurat[141] (v5.1.0) R package was used. After the first quality assessment,
cells with fewer than 200 gene features, more than 5000 gene features and more than 25 % of
mitochondrial genes were removed from the dataset. Additionally, the R package decontX[44] (v1.0.0)
was applied to adjust the dataset for cell free RNA. This approach uses a Bayesian model to estimate
and remove cell free RNA contaminations. Data was log normalized and scaled with the Seurat functions

'NormalizeData' and 'ScaleData'. Thereby, the number of features, UMI counts and percentage of
mitochondrial genes were regressed out. The top 2.000 variable feature genes were used for principal
component analysis (PCA) followed by clustering using the Louvain algorithm with multi level
refinements. The colon and tumor samples were processed separate from the blood and bone marrow
samples. To account for batch effects in the colon and tumor samples the fast mutual nearest neighbors
(FastMNN) algorithm (Seurat function 'FastMNNIntegration') was applied. The blood and bone
marrow samples were merged without batch effect correction.

Annotation, extraction and integration of neutrophils

Neutrophil clusters were identified using on the one hand neutrophil specific marker genes (*S100a8,*
S100a9) and an automatic annotation approach called SingleR[45] (v2.4.1) with the murine ImmGen[46]
(Immunological Genome Project) as reference which was loaded with the celldex[45] (v1.12.0) R package.
Neutrophil clusters were then extracted and integrated from all tissues using the FastMNN approach
mentioned before.

DEG analysis

To analyze DEGs between neutrophil clusters the Seurat function 'FindAllMarkers' with min.pct = 0.25
and logfc.threshold = 0.25 was used on log normalized data. This method applies a non-parametric
Wilcoxon rank sum test and corrects the p-values using the Bonferroni method. Genes of interest were
plotted in heatmap using pheatmap (v1.0.12).

938
939
940

Analysis of human neutrophil markers in mouse data
To analyze the expression of marker genes identified in human neutrophil clusters, the genes were first
converted to mouse gene symbols using the orthogene Bioconductor package (v1.8.0) and then plotted
in a heatmap using pheatmap (v1.0.12).

Pseudotime trajectory analysis with Slingshot
A pseudotime trajectory was inferred on the mnn.clusters of the combined mouse neutrophils dataset
(bone marrow, blood and colon) using the slingshot[47] (v2.10.0) R package. Therefore, the Seurat object
was first converted to a Single Cell Expression (SCE) object using sceasy R package (v0.0.7) , which
was then used for the slingshot analysis on the UMAP dimensional space. In brief, by minimum
spanning tree (MST) construction using the 'getLineages' function, a global lineage structure was

identified and pseudotime variables were inferred using the 'getCurves' function. To compare the
pseudotime between tissue, the cellular distribution was visualized in a density plot across pseudotime.
954
Neutrophil Activation Assay
Human neutrophils were isolated from freshly collected peripheral blood from healthy donors.
Neutrophils were purified using the MACSxpress Whole Blood Neutrophil Isolation Kit (Miltenyi
Biotec) according to the manufacturer's instructions. Neutrophils were washed with DPBS (Sigma-
Aldrich) and centrifuged at 115 g for 10 min. Then, 2 million neutrophils were seeded in a 12-well plate
in 1 mL of either RPMI 1640 (Sigma-Aldrich) supplemented with 2 mM L-glutamine (Sigma-Aldrich),
10% FBS (Sigma-Aldrich) and 10 mL/L Penicillin-Streptomycin (Sigma-Aldrich), or with conditioned
medium obtained as supernatant of SW900 or CaCo2 cell lines after 48 hours of growth in RPMI
(supplemented with L-glutamine, FBS and Penicillin-Streptomycin) to approximately 70% confluency.
Neutrophils were cultured for 24 hours at 37°C and 5% CO_2. Neutrophils were collected, counted, and
centrifuged at 450 g for 10 min at room temperature (20°C ± 2°C). Cells were resuspended at a
concentration of 2.5×10^5 cells/mL in Seahorse XF RPMI Medium pH 7.4 supplemented with 1 mM
pyruvate, 2 mM glutamine, 10 mM glucose (all from Agilent Technologies). Seahorse XFp Cell Culture
Miniplates (Agilent Technologies) were pre-coated with poly-L-lysine (Sigma-Aldrich) and 5×10^4 cells
were plated into each well. Miniplates were centrifuged at 300 g for 1 min with low-brake deceleration.
Seahorse XP Neutrophil Activation assay was performed on an XFp instrument with sequential injection
of 0.5 µM Rotenone/Antimycin A (Agilent Technologies) and 100 ng/mL phorbol 12-myristate 13-
acetate (PMA) (Fisher Scientific) according to the manufacturer's protocol. Data was processed by
Seahorse XFe Wave software (Agilent Technologies) and by GraphPad Prism software (v 10.4;
GraphPad Software). O2 consumption was calculated from the area under the curve (AUC) of the kinetic
range between activation and return to basal rates.

REFERENCES

1. Bray, F., Laversanne, M., Sung, H., Ferlay, J., Siegel, R.L., Soerjomataram, I., and Jemal, A. (2024). Global cancer statistics 2022: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries. CA. Cancer J. Clin. *74*, 229–263. https://doi.org/10.3322/caac.21834.

2. Akimoto, N., Ugai, T., Zhong, R., Hamada, T., Fujiyoshi, K., Giannakis, M., Wu, K., Cao, Y., Ng, K., and Ogino, S. (2021). Rising incidence of early-onset colorectal cancer - a call to action. Nat. Rev. Clin. Oncol. *18*, 230–243. https://doi.org/10.1038/s41571-020-00445-1.

3. Cremolini, C., Loupakis, F., Antoniotti, C., Lupi, C., Sensi, E., Lonardi, S., Mezi, S., Tomasello, G., Ronzoni, M., Zaniboni, A., et al. (2015). FOLFOXIRI plus bevacizumab versus FOLFIRI plus bevacizumab as first-line treatment of patients with metastatic colorectal cancer: updated overall survival and molecular subgroup analyses of the open-label, phase 3 TRIBE study. Lancet Oncol. *16*, 1306–1315. https://doi.org/10.1016/S1470-2045(15)00122-9.

4. Nicolantonio, F. (2021). Precision oncology in metastatic colorectal cancer - from biology to medicine. Nat. Rev. Clin. Oncol. *18*, 506-525,. https://doi.org/10.1038/s41571-021-00495-z.

5. Cancer Genome Atlas Network (2012). Comprehensive molecular characterization of human colon and rectal cancer. Nature *487*, 330–337. https://doi.org/10.1038/nature11252.

6. Le, D.T., Durham, J.N., Smith, K.N., Wang, H., Bartlett, B.R., Aulakh, L.K., Lu, S., Kemberling, H., Wilt, C., Luber, B.S., et al. (2017). Mismatch repair deficiency predicts response of solid tumors to PD-1 blockade. Science *357*, 409–413. https://doi.org/10.1126/science.aan6733.

7. Galon, J. (2006). Type, density, and location of immune cells within human colorectal tumors predict clinical outcome. Science *313*, 1960-1964,. https://doi.org/10.1126/science.1129139.

8. Sahin, I.H., Ciombor, K.K., Diaz, L.A., Yu, J., and Kim, R. (2022). Immunotherapy for Microsatellite Stable Colorectal Cancers: Challenges and Novel Therapeutic Avenues. Am. Soc. Clin. Oncol. Educ. Book Am. Soc. Clin. Oncol. Annu. Meet. *42*, 1–12. https://doi.org/10.1200/EDBK_349811.

9. Ros, J., Balconi, F., Baraibar, I., Saoudi Gonzalez, N., Salva, F., Tabernero, J., and Elez, E. (2023). Advances in immune checkpoint inhibitor combination strategies for microsatellite stable colorectal cancer. Front. Oncol. *13*, 1112276. https://doi.org/10.3389/fonc.2023.1112276.

10. Antoniotti, C., Rossini, D., Pietrantonio, F., Catteau, A., Salvatore, L., Lonardi, S., Boquet, I., Tamberi, S., Marmorino, F., Moretto, R., et al. (2022). Upfront FOLFOXIRI plus bevacizumab with or without atezolizumab in the treatment of patients with metastatic colorectal cancer (AtezoTRIBE): a multicentre, open-label, randomised, controlled, phase 2 trial. Lancet Oncol. *23*, 876–887. https://doi.org/10.1016/S1470-2045(22)00274-1.

11. Hackl, H., Charoentong, P., Finotello, F., and Trajanoski, Z. (2016). Computational genomics tools for dissecting tumour-immune cell interactions. Nat. Rev. Genet. *17*, 441–458. https://doi.org/10.1038/nrg.2016.67.

12. Finotello, F., Rieder, D., Hackl, H., and Trajanoski, Z. (2019). Next-generation computational tools for interrogating cancer immunity. Nat. Rev. Genet. *20*, 724–746. https://doi.org/10.1038/s41576-019-0166-7.

13. Angelova, M., Charoentong, P., Hackl, H., Fischer, M.L., Snajder, R., Krogsdam, A.M., Waldner, M.J., Bindea, G., Mlecnik, B., Galon, J., et al. (2015). Characterization of the

immunophenotypes and antigenomes of colorectal cancers reveals distinct tumor escape mechanisms and novel targets for immunotherapy. Genome Biol. *16*, 64. https://doi.org/10.1186/s13059-015-0620-6.

14. Charoentong, P., Finotello, F., Angelova, M., Mayer, C., Efremova, M., Rieder, D., Hackl, H., and Trajanoski, Z. (2017). Pan-cancer Immunogenomic Analyses Reveal Genotype-Immunophenotype Relationships and Predictors of Response to Checkpoint Blockade. Cell Rep. *18*, 248–262. https://doi.org/10.1016/j.celrep.2016.12.019.

15. Salcher, S., Sturm, G., Horvath, L., Untergasser, G., Kuempers, C., Fotakis, G., Panizzolo, E., Martowicz, A., Trebo, M., Pall, G., et al. (2022). High-resolution single-cell atlas reveals diversity and plasticity of tissue-resident neutrophils in non-small cell lung cancer. Cancer Cell *40*, 1503-1520.e8. https://doi.org/10.1016/j.ccell.2022.10.008.

16. Chu, X., Li, X., Zhang, Y., Dang, G., Miao, Y., Xu, W., Wang, J., Zhang, Z., and Cheng, S. (2024). Integrative single-cell analysis of human colorectal cancer reveals patient stratification with distinct immune evasion mechanisms. Nat. Cancer *5*, 1409–1426. https://doi.org/10.1038/s43018-024-00807-z.

17. Hirschhorn, D., Budhu, S., Kraehenbuehl, L., Gigoux, M., Schröder, D., Chow, A., Ricca, J.M., Gasmi, B., De Henau, O., Mangarin, L.M.B., et al. (2023). T cell immunotherapies engage neutrophils to eliminate tumor antigen escape variants. Cell *186*, 1432-1447.e17. https://doi.org/10.1016/j.cell.2023.03.007.

18. Väyrynen, J.P., Lau, M.C., Haruki, K., Väyrynen, S.A., Dias Costa, A., Borowsky, J., Zhao, M., Fujiyoshi, K., Arima, K., Twombly, T.S., et al. (2020). Prognostic Significance of Immune Cell Populations Identified by Machine Learning in Colorectal Cancer Using Routine Hematoxylin and Eosin-Stained Sections. Clin. Cancer Res. Off. J. Am. Assoc. Cancer Res. *26*, 4326–4338. https://doi.org/10.1158/1078-0432.CCR-20-0071.

19. Governa, V., Trella, E., Mele, V., Tornillo, L., Amicarella, F., Cremonesi, E., Muraro, M.G., Xu, H., Droeser, R., Däster, S.R., et al. (2017). The Interplay Between Neutrophils and CD8+ T Cells Improves Survival in Human Colorectal Cancer. Clin. Cancer Res. Off. J. Am. Assoc. Cancer Res. *23*, 3847–3858. https://doi.org/10.1158/1078-0432.CCR-16-2047.

20. Hedrick, C.C., and Malanchi, I. (2022). Neutrophils in cancer: heterogeneous and multifaceted. Nat. Rev. Immunol. *22*, 173–187. https://doi.org/10.1038/s41577-021-00571-6.

21. Gungabeesoon, J., Gort-Freitas, N.A., Kiss, M., Bolli, E., Messemaker, M., Siwicki, M., Hicham, M., Bill, R., Koch, P., Cianciaruso, C., et al. (2023). A neutrophil response linked to tumor control in immunotherapy. Cell *186*, 1448-1464.e20. https://doi.org/10.1016/j.cell.2023.02.032.

22. Schürch, C.M., Bhate, S.S., Barlow, G.L., Phillips, D.J., Noti, L., Zlobec, I., Chu, P., Black, S., Demeter, J., McIlwain, D.R., et al. (2020). Coordinated Cellular Neighborhoods Orchestrate Antitumoral Immunity at the Colorectal Cancer Invasive Front. Cell *183*, 838. https://doi.org/10.1016/j.cell.2020.10.021.

23. Salcher, S., Heidegger, I., Untergasser, G., Fotakis, G., Scheiber, A., Martowicz, A., Noureen, A., Krogsdam, A., Schatz, C., Schäfer, G., et al. (2024). Comparative analysis of 10X Chromium vs. BD Rhapsody whole transcriptome single-cell sequencing technologies in complex human tissues. Heliyon *10*, e28358. https://doi.org/10.1016/j.heliyon.2024.e28358.

24. Scheiber, A., Trebo, M., Pittl, A., Heidegger, I., Hautz, T., Oberhuber, R., Trajanoski, Z., Augustin, F., Sopper, S., Wolf, D., et al. (2024). Profiling low-mRNA content cells in complex human tissues using BD Rhapsody single-cell analysis. STAR Protoc. *5*, 103475. https://doi.org/10.1016/j.xpro.2024.103475.

25.   Schubert, M. (2018). Perturbation-response genes reveal signaling footprints in cancer
gene expression. Nat Commun *9, 20*. https://doi.org/10.1038/s41467-017-02391-6.
26.   Choi, H.Y., Jo, H., Zhao, X., Hoadley, K.A., Newman, S., Holt, J., Hayward, M.C.,
Love, M.I., Marron, J.S., and Hayes, D.N. (2021). SCISSOR: a framework for
identifying structural changes in RNA transcripts. Nat. Commun. *12*, 286.
https://doi.org/10.1038/s41467-020-20593-3.
27.   Horvath, L., Puschmann, C., Scheiber, A., Martowicz, A., Sturm, G., Trajanoski, Z.,
Wolf, D., Pircher, A., and Salcher, S. (2024). Beyond binary: bridging neutrophil
diversity to new therapeutic approaches in NSCLC. Trends Cancer *10*, 457–474.
https://doi.org/10.1016/j.trecan.2024.01.010.
28.   Park, M.D., Le Berichel, J., Hamon, P., Wilk, C.M., Belabed, M., Yatim, N., Saffon, A.,
Boumelha, J., Falcomatà, C., Tepper, A., et al. (2024). Hematopoietic aging promotes
cancer by fueling IL-1α-driven emergency myelopoiesis. Science *386*, eadn0327.
https://doi.org/10.1126/science.adn0327.
29.   Singhal, S., Bhojnagarwala, P.S., O'Brien, S., Moon, E.K., Garfall, A.L., Rao, A.S.,
Quatromoni, J.G., Stephen, T.L., Litzky, L., Deshpande, C., et al. (2016). Origin and
Role of a Subset of Tumor-Associated Neutrophils with Antigen-Presenting Cell
Features in Early-Stage Human Lung Cancer. Cancer Cell *30*, 120–135.
https://doi.org/10.1016/j.ccell.2016.06.001.
30.   Matusiak, M., Hickey, J.W., van IJzendoorn, D.G.P., Lu, G., Kidziński, L., Zhu, S.,
Colburg, D.R.C., Luca, B., Phillips, D.J., Brubaker, S.W., et al. (2024). Spatially
Segregated Macrophage Populations Predict Distinct Outcomes in Colon Cancer. Cancer
Discov. *14*, 1418–1439. https://doi.org/10.1158/2159-8290.CD-23-1300.
31.   Birk, S., Bonafonte-Pardàs, I., Feriz, A.M., Boxall, A., Agirre, E., Memi, F., Maguza,
A., Fan, R., Castelo-Branco, G., Bayraktar, O.A., et al. (2024). Large-scale
characterization of cell niches in spatial atlases using bio-inspired graph learning.
Preprint at bioRxiv, https://doi.org/10.1101/2024.02.21.581428
https://doi.org/10.1101/2024.02.21.581428.
32.   Ouyang, Y., Zhong, W., Xu, P., Wang, B., Zhang, L., Yang, M., Chen, J., Li, H., Li, S.,
Chen, X., et al. (2024). Tumor-associated neutrophils suppress CD8+ T cell immunity in
urothelial bladder carcinoma through the COX-2/PGE2/IDO1 Axis. Br. J. Cancer *130*,
880–891. https://doi.org/10.1038/s41416-023-02552-z.
33.   Metzemaekers, M., Gouwy, M., and Proost, P. (2020). Neutrophil chemoattractant
receptors in health and disease: double-edged swords. Cell. Mol. Immunol. *17*, 433–450.
https://doi.org/10.1038/s41423-020-0412-0.
34.   Plattner, C., Lamberti, G., Blattmann, P., Kirchmair, A., Rieder, D., Loncova, Z., Sturm,
G., Scheidl, S., Ijsselsteijn, M., Fotakis, G., et al. (2023). Functional and spatial
proteomics profiling reveals intra- and intercellular signaling crosstalk in colorectal
cancer. iScience *26*, 108399. https://doi.org/10.1016/j.isci.2023.108399.
35.   Dalessi, T., Valenta, T., Moro, G., Brügger, M.D., Meijs, A., Szczerba, B.M., Brunner,
T., Lampart, S., Nater, M., Platt, R.J., et al. (2024). An oncoembryology approach
uncovers SoxC-driven regulation of colon development and cancer. Preprint at bioRxiv,
https://doi.org/10.1101/2024.10.08.617191 https://doi.org/10.1101/2024.10.08.617191.
36.   Locard-Paulet, M., Palasca, O., and Jensen, L.J. (2022). Identifying the genes impacted
by cell proliferation in proteomics and transcriptomics studies. PLoS Comput. Biol. *18*,
e1010604. https://doi.org/10.1371/journal.pcbi.1010604.
37.   Li, T., Li, X., Zamani, A., Wang, W., Lee, C.-N., Li, M., Luo, G., Eiler, E., Sun, H.,
Ghosh, S., et al. (2020). c-Rel Is a Myeloid Checkpoint for Cancer Immunotherapy. Nat.
Cancer *1*, 507–517. https://doi.org/10.1038/s43018-020-0061-3.
38.   Fridman, W.H., Meylan, M., Petitprez, F., Sun, C.-M., Italiano, A., and Sautès-Fridman,

C. (2022). B cells and tertiary lymphoid structures as determinants of tumour immune
contexture and clinical outcome. Nat. Rev. Clin. Oncol. *19*, 441–457.
https://doi.org/10.1038/s41571-022-00619-z.

39. Moffat, A., and Gwyer Findlay, E. (2024). Evidence for antigen presentation by human
neutrophils. Blood *143*, 2455–2463. https://doi.org/10.1182/blood.2023023444.

40. Wu, Y., Ma, J., Yang, X., Nan, F., Zhang, T., Ji, S., Rao, D., Feng, H., Gao, K., Gu, X.,
et al. (2024). Neutrophil profiling illuminates anti-tumor antigen-presenting potency.
Cell *187*, 1422-1439.e24. https://doi.org/10.1016/j.cell.2024.02.005.

41. Poplimont, H., Georgantzoglou, A., Boulch, M., Walker, H.A., Coombs, C.,
Papaleonidopoulou, F., and Sarris, M. (2020). Neutrophil Swarming in Damaged Tissue
Is Orchestrated by Connexins and Cooperative Calcium Alarm Signals. Curr. Biol. CB
30, 2761-2776.e7. https://doi.org/10.1016/j.cub.2020.05.030.

42. LaMarche, N.M., Hegde, S., Park, M.D., Maier, B.B., Troncoso, L., Le Berichel, J.,
Hamon, P., Belabed, M., Mattiuz, R., Hennequin, C., et al. (2024). An IL-4 signalling
axis in bone marrow drives pro-tumorigenic myelopoiesis. Nature *625*, 166–174.
https://doi.org/10.1038/s41586-023-06797-9.

43. Dobin, A., Davis, C.A., Schlesinger, F., Drenkow, J., Zaleski, C., Jha, S., Batut, P.,
Chaisson, M., and Gingeras, T.R. (2013). STAR: ultrafast universal RNA-seq aligner.
Bioinforma. Oxf. Engl. *29*, 15–21. https://doi.org/10.1093/bioinformatics/bts635.

44. Yang, S., Corbett, S.E., Koga, Y., Wang, Z., Johnson, W.E., Yajima, M., and Campbell,
J.D. (2020). Decontamination of ambient RNA in single-cell RNA-seq with DecontX.
Genome Biol. *21*, 57. https://doi.org/10.1186/s13059-020-1950-6.

45. Aran, D., Looney, A.P., Liu, L., Wu, E., Fong, V., Hsu, A., Chak, S., Naikawadi, R.P.,
Wolters, P.J., Abate, A.R., et al. (2019). Reference-based analysis of lung single-cell
sequencing reveals a transitional profibrotic macrophage. Nat. Immunol. *20*, 163–172.
https://doi.org/10.1038/s41590-018-0276-y.

46. Heng, T.S.P., Painter, M.W., and Immunological Genome Project Consortium (2008).
The Immunological Genome Project: networks of gene expression in immune cells. Nat.
Immunol. *9*, 1091–1094. https://doi.org/10.1038/ni1008-1091.

47. Street, K., Risso, D., Fletcher, R.B., Das, D., Ngai, J., Yosef, N., Purdom, E., and
Dudoit, S. (2018). Slingshot: cell lineage and pseudotime inference for single-cell
transcriptomics. BMC Genomics *19*, 477. https://doi.org/10.1186/s12864-018-4772-0.

48. Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T.,
Preibisch, S., Rueden, C., Saalfeld, S., Schmid, B., et al. (2012). Fiji: an open-source
platform for biological-image analysis. Nat. Methods *9*, 676–682.
https://doi.org/10.1038/nmeth.2019.

49. Bankhead, P., Loughrey, M.B., Fernández, J.A., Dombrowski, Y., McArt, D.G., Dunne,
P.D., McQuaid, S., Gray, R.T., Murray, L.J., Coleman, H.G., et al. (2017). QuPath:
Open source software for digital pathology image analysis. Sci. Rep. *7*, 16878.
https://doi.org/10.1038/s41598-017-17204-5.

50. Stirling, D.R., Swain-Bowden, M.J., Lucas, A.M., Carpenter, A.E., Cimini, B.A., and
Goodman, A. (2021). CellProfiler 4: improvements in speed, utility and usability. BMC
Bioinformatics *22*, 433. https://doi.org/10.1186/s12859-021-04344-9.

51. Nemati, N., Boeck, N., Lamberti, G., Lisandrelli, R., and Trajanoski, Z. (2024). Protocol
for functional profiling of patient-derived organoids for precision oncology. STAR
Protoc. *5*, 102887. https://doi.org/10.1016/j.xpro.2024.102887.

52. Wu, Y., Ma, J., Yang, X., Nan, F., Zhang, T., Ji, S., Rao, D., Feng, H., Gao, K., Gu, X.,
et al. (2024). Neutrophil profiling illuminates anti-tumor antigen-presenting potency.
Cell *187*, 1422-1439.e24. https://doi.org/10.1016/j.cell.2024.02.005.

53. Bian, S., Hou, Y., Zhou, X., Li, X., Yong, J., Wang, Y., Wang, W., Yan, J., Hu, B., Guo,

H., et al. (2018). Single-cell multiomics sequencing and analyses of human colorectal
cancer. Science *362*, 1060–1063. https://doi.org/10.1126/science.aao3791.

54.   Borràs, D.M., Verbandt, S., Ausserhofer, M., Sturm, G., Lim, J., Verge, G.A.,
Vanmeerbeek, I., Laureano, R.S., Govaerts, J., Sprooten, J., et al. (2023). Single cell
dynamics of tumor specificity vs bystander activity in CD8+ T cells define the diverse
immune landscapes in colorectal cancer. Cell Discov. *9*, 114.
https://doi.org/10.1038/s41421-023-00605-4.

55.   Burclaff, J., Bliton, R.J., Breau, K.A., Ok, M.T., Gomez-Martinez, I., Ranek, J.S., Bhatt,
A.P., Purvis, J.E., Woosley, J.T., and Magness, S.T. (2022). A Proximal-to-Distal
Survey of Healthy Adult Human Small Intestine and Colon Epithelium by Single-Cell
Transcriptomics. Cell. Mol. Gastroenterol. Hepatol. *13*, 1554–1589.
https://doi.org/10.1016/j.jcmgh.2022.02.007.

56.   Che, L.-H., Liu, J.-W., Huo, J.-P., Luo, R., Xu, R.-M., He, C., Li, Y.-Q., Zhou, A.-J.,
Huang, P., Chen, Y.-Y., et al. (2021). A single-cell atlas of liver metastases of colorectal
cancer reveals reprogramming of the tumor microenvironment in response to
preoperative chemotherapy. Cell Discov. *7*, 80. https://doi.org/10.1038/s41421-021-
00312-y.

57.   Chen, Y., Wang, D., Li, Y., Qi, L., Si, W., Bo, Y., Chen, X., Ye, Z., Fan, H., Liu, B., et
al. (2024). Spatiotemporal single-cell analysis decodes cellular dynamics underlying
different responses to immunotherapy in colorectal cancer. Cancer Cell *42*, 1268-
1285.e7. https://doi.org/10.1016/j.ccell.2024.06.009.

58.   Domínguez Conde, C., Xu, C., Jarvis, L.B., Rainbow, D.B., Wells, S.B., Gomes, T.,
Howlett, S.K., Suchanek, O., Polanski, K., King, H.W., et al. (2022). Cross-tissue
immune cell analysis reveals tissue-specific features in humans. Science *376*, eabl5197.
https://doi.org/10.1126/science.abl5197.

59.   Elmentaite, R., Kumasaka, N., Roberts, K., Fleming, A., Dann, E., King, H.W.,
Kleshchevnikov, V., Dabrowska, M., Pritchard, S., Bolt, L., et al. (2021). Cells of the
human intestinal tract mapped across space and time. Nature *597*, 250–255.
https://doi.org/10.1038/s41586-021-03852-1.

60.   Garrido-Trigo, A., Corraliza, A.M., Veny, M., Dotti, I., Melón-Ardanaz, E., Rill, A.,
Crowell, H.L., Corbí, Á., Gudiño, V., Esteller, M., et al. (2023). Macrophage and
neutrophil heterogeneity at single-cell spatial resolution in human inflammatory bowel
disease. Nat. Commun. *14*, 4506. https://doi.org/10.1038/s41467-023-40156-6.

61.   Giguelay, A., Turtoi, E., Khelaf, L., Tosato, G., Dadi, I., Chastel, T., Poul, M.-A.,
Pratlong, M., Nicolescu, S., Severac, D., et al. (2022). The landscape of cancer-
associated fibroblasts in colorectal cancer liver metastases. Theranostics *12*, 7624–7639.
https://doi.org/10.7150/thno.72853.

62.   Guo, W., Zhang, C., Wang, X., Dou, D., Chen, D., and Li, J. (2022). Resolving the
difference between left-sided and right-sided colorectal cancer by single-cell sequencing.
JCI Insight *7*, e152616. https://doi.org/10.1172/jci.insight.152616.

63.   Harmon, C., Zaborowski, A., Moore, H., St Louis, P., Slattery, K., Duquette, D.,
Scanlan, J., Kane, H., Kunkemoeller, B., McIntyre, C.L., et al. (2023). γδ T cell
dichotomy with opposing cytotoxic and wound healing functions in human solid tumors.
Nat. Cancer *4*, 1122–1137. https://doi.org/10.1038/s43018-023-00589-w.

64.   He, S., Wang, L.-H., Liu, Y., Li, Y.-Q., Chen, H.-T., Xu, J.-H., Peng, W., Lin, G.-W.,
Wei, P.-P., Li, B., et al. (2020). Single-cell transcriptome profiling of an adult human
cell atlas of 15 major organs. Genome Biol. *21*, 294. https://doi.org/10.1186/s13059-
020-02210-0.

65.   Huang, C., Wang, X., Wang, Y., Feng, Y., Wang, X., Chen, S., Yan, P., Liao, J., Zhang,
Q., Mao, C., et al. (2024). Sirpα on tumor-associated myeloid cells restrains antitumor

immunity in colorectal cancer independent of its interaction with CD47. Nat. Cancer *5*,
500–516. https://doi.org/10.1038/s43018-023-00691-z.

66. James, K.R., Gomes, T., Elmentaite, R., Kumar, N., Gulliver, E.L., King, H.W., Stares,
M.D., Bareham, B.R., Ferdinand, J.R., Petrova, V.N., et al. (2020). Distinct microbial
and immune niches of the human colon. Nat. Immunol. *21*, 343–353.
https://doi.org/10.1038/s41590-020-0602-z.

67. Ji, T., Fu, H., Wang, L., Chen, J., Tian, S., Wang, G., Wang, L., and Wang, Z. (2024).
Single-cell RNA profiling reveals classification and characteristics of mononuclear
phagocytes in colorectal cancer. PLoS Genet. *20*, e1011176.
https://doi.org/10.1371/journal.pgen.1011176.

68. Ji, L., Fu, G., Huang, M., Kao, X., Zhu, J., Dai, Z., Chen, Y., Li, H., Zhou, J., Chu, X., et
al. (2024). scRNA-seq of colorectal cancer shows regional immune atlas with the
function of CD20+ B cells. Cancer Lett. *584*, 216664.
https://doi.org/10.1016/j.canlet.2024.216664.

69. Joanito, I., Wirapati, P., Zhao, N., Nawaz, Z., Yeo, G., Lee, F., Eng, C.L.P., Macalinao,
D.C., Kahraman, M., Srinivasan, H., et al. (2022). Single-cell and bulk transcriptome
sequencing identifies two epithelial tumor cell states and refines the consensus
molecular classification of colorectal cancer. Nat. Genet. *54*, 963–975.
https://doi.org/10.1038/s41588-022-01100-4.

70. Khaliq, A.M., Erdogan, C., Kurt, Z., Turgut, S.S., Grunvald, M.W., Rand, T., Khare, S.,
Borgia, J.A., Hayden, D.M., Pappas, S.G., et al. (2022). Refining colorectal cancer
classification and clinical stratification through a single-cell atlas. Genome Biol. *23*, 113.
https://doi.org/10.1186/s13059-022-02677-z.

71. Kong, L., Pokatayev, V., Lefkovith, A., Carter, G.T., Creasey, E.A., Krishna, C.,
Subramanian, S., Kochar, B., Ashenberg, O., Lau, H., et al. (2023). The landscape of
immune dysregulation in Crohn's disease revealed through single-cell transcriptomic
profiling in the ileum and colon. Immunity *56*, 444-458.e5.
https://doi.org/10.1016/j.immuni.2023.01.002.

72. Lee, H.-O., Hong, Y., Etlioglu, H.E., Cho, Y.B., Pomella, V., Van den Bosch, B.,
Vanhecke, J., Verbandt, S., Hong, H., Min, J.-W., et al. (2020). Lineage-dependent gene
expression programs influence the immune landscape of colorectal cancer. Nat. Genet.
52, 594–603. https://doi.org/10.1038/s41588-020-0636-z.

73. Li, H., Courtois, E.T., Sengupta, D., Tan, Y., Chen, K.H., Goh, J.J.L., Kong, S.L., Chua,
C., Hon, L.K., Tan, W.S., et al. (2017). Reference component analysis of single-cell
transcriptomes elucidates cellular heterogeneity in human colorectal tumors. Nat. Genet.
49, 708–718. https://doi.org/10.1038/ng.3818.

74. Li, J., Wu, C., Hu, H., Qin, G., Wu, X., Bai, F., Zhang, J., Cai, Y., Huang, Y., Wang, C.,
et al. (2023). Remodeling of the immune and stromal cell compartment by PD-1
blockade in mismatch repair-deficient colorectal cancer. Cancer Cell *41*, 1152-1169.e7.
https://doi.org/10.1016/j.ccell.2023.04.011.

75. Liu, X., Wang, X., Yang, Q., Luo, L., Liu, Z., Ren, X., Lei, K., Li, S., Xie, Z., Zheng,
G., et al. (2024). Th17 Cells Secrete TWEAK to Trigger Epithelial-Mesenchymal
Transition and Promote Colorectal Cancer Liver Metastasis. Cancer Res. *84*, 1352–1371.
https://doi.org/10.1158/0008-5472.CAN-23-2123.

76. Mazzurana, L., Czarnewski, P., Jonsson, V., Wigge, L., Ringnér, M., Williams, T.C.,
Ravindran, A., Björklund, Å.K., Säfholm, J., Nilsson, G., et al. (2021). Tissue-specific
transcriptional imprinting and heterogeneity in human innate lymphoid cells revealed by
full-length single-cell RNA-sequencing. Cell Res. *31*, 554–568.
https://doi.org/10.1038/s41422-020-00445-x.

77. Parikh, K., Antanaviciute, A., Fawkner-Corbett, D., Jagielowicz, M., Aulicino, A.,

1276  Lagerholm, C., Davis, S., Kinchen, J., Chen, H.H., Alham, N.K., et al. (2019). Colonic
1277  epithelial cell diversity in health and inflammatory bowel disease. Nature *567*, 49–55.
1278  https://doi.org/10.1038/s41586-019-0992-y.

1279 78. Pelka, K., Hofree, M., Chen, J.H., Sarkizova, S., Pirl, J.D., Jorgji, V., Bejnood, A.,
1280  Dionne, D., Ge, W.H., Xu, K.H., et al. (2021). Spatially organized multicellular immune
1281  hubs in human colorectal cancer. Cell *184*, 4734-4752.e20.
1282  https://doi.org/10.1016/j.cell.2021.08.003.

1283 79. Qi, J., Sun, H., Zhang, Y., Wang, Z., Xun, Z., Li, Z., Ding, X., Bao, R., Hong, L., Jia,
1284  W., et al. (2022). Single-cell and spatial analysis reveal interaction of FAP+ fibroblasts
1285  and SPP1+ macrophages in colorectal cancer. Nat. Commun. *13*, 1742.
1286  https://doi.org/10.1038/s41467-022-29366-6.

1287 80. Qian, J., Olbrecht, S., Boeckx, B., Vos, H., Laoui, D., Etlioglu, E., Wauters, E., Pomella,
1288  V., Verbandt, S., Busschaert, P., et al. (2020). A pan-cancer blueprint of the
1289  heterogeneous tumor microenvironment revealed by single-cell profiling. Cell Res. *30*,
1290  745–762. https://doi.org/10.1038/s41422-020-0355-0.

1291 81. Qin, P., Chen, H., Wang, Y., Huang, L., Huang, K., Xiao, G., Han, C., Hu, J., Lin, D.,
1292  Wan, X., et al. (2023). Cancer-associated fibroblasts undergoing neoadjuvant
1293  chemotherapy suppress rectal cancer revealed by single-cell and spatial transcriptomics.
1294  Cell Rep. Med. *4*, 101231. https://doi.org/10.1016/j.xcrm.2023.101231.

1295 82. Sathe, A., Mason, K., Grimes, S.M., Zhou, Z., Lau, B.T., Bai, X., Su, A., Tan, X., Lee,
1296  H., Suarez, C.J., et al. (2023). Colorectal Cancer Metastases in the Liver Establish
1297  Immunosuppressive Spatial Networking between Tumor-Associated SPP1+
1298  Macrophages and Fibroblasts. Clin. Cancer Res. Off. J. Am. Assoc. Cancer Res. *29*,
1299  244–260. https://doi.org/10.1158/1078-0432.CCR-22-2041.

1300 83. Scheid, J.F., Eraslan, B., Hudak, A., Brown, E.M., Sergio, D., Delorey, T.M., Phillips,
1301  D., Lefkovith, A., Jess, A.T., Duck, L.W., et al. (2023). Remodeling of colon plasma cell
1302  repertoire within ulcerative colitis patients. J. Exp. Med. *220*, e20220538.
1303  https://doi.org/10.1084/jem.20220538.

1304 84. Smillie, C.S., Biton, M., Ordovas-Montanes, J., Sullivan, K.M., Burgin, G., Graham,
1305  D.B., Herbst, R.H., Rogel, N., Slyper, M., Waldman, J., et al. (2019). Intra- and Inter-
1306  cellular Rewiring of the Human Colon during Ulcerative Colitis. Cell *178*, 714-730.e22.
1307  https://doi.org/10.1016/j.cell.2019.06.029.

1308 85. Terekhanova, N.V., Karpova, A., Liang, W.-W., Strzalkowski, A., Chen, S., Li, Y.,
1309  Southard-Smith, A.N., Iglesia, M.D., Wendl, M.C., Jayasinghe, R.G., et al. (2023).
1310  Epigenetic regulation during cancer transitions across 11 tumour types. Nature *623*,
1311  432–441. https://doi.org/10.1038/s41586-023-06682-5.

1312 86. Thomas, M.F., Slowikowski, K., Manakongtreecheep, K., Sen, P., Samanta, N., Tantivit,
1313  J., Nasrallah, M., Zubiri, L., Smith, N.P., Tirard, A., et al. (2024). Single-cell
1314  transcriptomic analyses reveal distinct immune cell contributions to epithelial barrier
1315  dysfunction in checkpoint inhibitor colitis. Nat. Med. *30*, 1349–1362.
1316  https://doi.org/10.1038/s41591-024-02895-x.

1317 87. Tian, J., Chen, J.H., Chao, S.X., Pelka, K., Giannakis, M., Hess, J., Burke, K., Jorgji, V.,
1318  Sindurakar, P., Braverman, J., et al. (2023). Combined PD-1, BRAF and MEK inhibition
1319  in BRAFV600E colorectal cancer: a phase 2 trial. Nat. Med. *29*, 458–466.
1320  https://doi.org/10.1038/s41591-022-02181-8.

1321 88. Uhlitz, F. (2021). Mitogen-activated protein kinase activity drives cell trajectories in
1322  colorectal cancer. EMBO Mol Med *13, e14123*.
1323  https://doi.org/10.15252/emmm.202114123.

1324 89. Chen, B., Scurrah, C.R., McKinley, E.T., Simmons, A.J., Ramirez-Solano, M.A., Zhu,
1325  X., Markham, N.O., Heiser, C.N., Vega, P.N., Rolong, A., et al. (2021). Differential pre-

malignant programs and microenvironment chart distinct paths to malignancy in human
colorectal polyps. Cell *184*, 6262-6280.e26. https://doi.org/10.1016/j.cell.2021.11.031.
90. Wang, Y., Song, W., Wang, J., Wang, T., Xiong, X., Qi, Z., Fu, W., Yang, X., and Chen,
Y.-G. (2020). Single-cell transcriptome analysis reveals differential nutrient absorption
functions in human intestine. J. Exp. Med. *217*, e20191130.
https://doi.org/10.1084/jem.20191130.
91. Wang, H., Gong, P., Chen, T., Gao, S., Wu, Z., Wang, X., Li, J., Marjani, S.L., Costa, J.,
Weissman, S.M., et al. (2021). Colorectal Cancer Stem Cell States Uncovered by
Simultaneous Single-Cell Analysis of Transcriptome and Telomeres. Adv. Sci. Weinh.
Baden-Wurtt. Ger. *8*, 2004320. https://doi.org/10.1002/advs.202004320.
92. Wang, F., Long, J., Li, L., Wu, Z.-X., Da, T.-T., Wang, X.-Q., Huang, C., Jiang, Y.-H.,
Yao, X.-Q., Ma, H.-Q., et al. (2023). Single-cell and spatial transcriptome analysis
reveals the cellular heterogeneity of liver metastatic colorectal cancer. Sci. Adv. *9*,
eadf5464. https://doi.org/10.1126/sciadv.adf5464.
93. Wu, Y., Yang, S., Ma, J., Chen, Z., Song, G., Rao, D., Cheng, Y., Huang, S., Liu, Y.,
Jiang, S., et al. (2022). Spatiotemporal Immune Landscape of Colorectal Cancer Liver
Metastasis at Single-Cell Level. Cancer Discov. *12*, 134–153.
https://doi.org/10.1158/2159-8290.CD-21-0316.
94. Yang, M., Yang, C., Ma, D., Li, Z., Zhao, W., and Yang, D. (2023). Single-cell analysis
reveals cellular reprogramming in advanced colon cancer following FOLFOX-
bevacizumab treatment. Front. Oncol. *13*, 1219642.
https://doi.org/10.3389/fonc.2023.1219642.
95. Zhang, L., Yu, X., Zheng, L., Zhang, Y., Li, Y., Fang, Q., Gao, R., Kang, B., Zhang, Q.,
Huang, J.Y., et al. (2018). Lineage tracking reveals dynamic relationships of T cells in
colorectal cancer. Nature *564*, 268–272. https://doi.org/10.1038/s41586-018-0694-x.
96. Zhang, L., Li, Z., Skrzypczynska, K.M., Fang, Q., Zhang, W., O'Brien, S.A., He, Y.,
Wang, L., Zhang, Q., Kim, A., et al. (2020). Single-Cell Analyses Inform Mechanisms
of Myeloid-Targeted Therapies in Colon Cancer. Cell *181*, 442-459.e29.
https://doi.org/10.1016/j.cell.2020.03.048.
97. Zheng, X., Song, J., Yu, C., Zhou, Z., Liu, X., Yu, J., Xu, G., Yang, J., He, X., Bai, X.,
et al. (2022). Single-cell transcriptomic profiling unravels the adenoma-initiation role of
protein tyrosine kinases during colorectal tumorigenesis. Signal Transduct. Target. Ther.
7, 60. https://doi.org/10.1038/s41392-022-00881-8.
98. de Vries, N.L., van de Haar, J., Veninga, V., Chalabi, M., Ijsselsteijn, M.E., van der
Ploeg, M., van den Bulk, J., Ruano, D., van den Berg, J.G., Haanen, J.B., et al. (2023).
γδ T cells are effectors of immunotherapy in cancers with HLA class I defects. Nature
613, 743–750. https://doi.org/10.1038/s41586-022-05593-1.
99. Picelli, S., Björklund, Å.K., Faridani, O.R., Sagasser, S., Winberg, G., and Sandberg, R.
(2013). Smart-seq2 for sensitive full-length transcriptome profiling in single cells. Nat.
Methods *10*, 1096–1098. https://doi.org/10.1038/nmeth.2639.
100. Klein, A.M., Mazutis, L., Akartuna, I., Tallapragada, N., Veres, A., Li, V., Peshkin, L.,
Weitz, D.A., and Kirschner, M.W. (2015). Droplet barcoding for single-cell
transcriptomics applied to embryonic stem cells. Cell *161*, 1187–1201.
https://doi.org/10.1016/j.cell.2015.04.044.
101. Macosko, E.Z., Basu, A., Satija, R., Nemesh, J., Shekhar, K., Goldman, M., Tirosh, I.,
Bialas, A.R., Kamitaki, N., Martersteck, E.M., et al. (2015). Highly Parallel Genome-
wide Expression Profiling of Individual Cells Using Nanoliter Droplets. Cell *161*, 1202–
1214. https://doi.org/10.1016/j.cell.2015.05.002.
102. Goveia, J., Rohlenova, K., Taverna, F., Treps, L., Conradi, L.-C., Pircher, A., Geldhof,
V., de Rooij, L.P.M.H., Kalucka, J., Sokol, L., et al. (2020). An Integrated Gene

Expression Landscape Profiling Approach to Identify Lung Tumor Endothelial Cell
Heterogeneity and Angiogenic Candidates. Cancer Cell *37*, 21-36.e13.
https://doi.org/10.1016/j.ccell.2019.12.001.

103. Guo, X., Zhang, Y., Zheng, L., Zheng, C., Song, J., Zhang, Q., Kang, B., Liu, Z., Jin, L.,
Xing, R., et al. (2018). Global characterization of T cells in non-small-cell lung cancer
by single-cell sequencing. Nat. Med. *24*, 978–985. https://doi.org/10.1038/s41591-018-
0045-3.

104. Maier, B., Leader, A.M., Chen, S.T., Tung, N., Chang, C., LeBerichel, J., Chudnovskiy,
A., Maskey, S., Walker, L., Finnigan, J.P., et al. (2020). A conserved dendritic-cell
regulatory program limits antitumour immunity. Nature *580*, 257–262.
https://doi.org/10.1038/s41586-020-2134-y.

105. Madissoon, E., Wilbrey-Clark, A., Miragaia, R.J., Saeb-Parsy, K., Mahbubani, K.T.,
Georgakopoulos, N., Harding, P., Polanski, K., Huang, N., Nowicki-Osuch, K., et al.
(2019). scRNA-seq assessment of the human lung, spleen, and esophagus tissue stability
after cold preservation. Genome Biol. *21*, 1. https://doi.org/10.1186/s13059-019-1906-x.

106. Adams, T.S., Schupp, J.C., Poli, S., Ayaub, E.A., Neumark, N., Ahangari, F., Chu, S.G.,
Raby, B.A., DeIuliis, G., Januszyk, M., et al. (2020). Single-cell RNA-seq reveals
ectopic and aberrant lung-resident cell populations in idiopathic pulmonary fibrosis. Sci.
Adv. *6*, eaba1983. https://doi.org/10.1126/sciadv.aba1983.

107. Reyfman, P.A., Walter, J.M., Joshi, N., Anekalla, K.R., McQuattie-Pimentel, A.C.,
Chiu, S., Fernandez, R., Akbarpour, M., Chen, C.-I., Ren, Z., et al. (2019). Single-Cell
Transcriptomic Analysis of Human Lung Provides Insights into the Pathobiology of
Pulmonary Fibrosis. Am. J. Respir. Crit. Care Med. *199*, 1517–1536.
https://doi.org/10.1164/rccm.201712-2410OC.

108. Habermann, A.C., Gutierrez, A.J., Bui, L.T., Yahn, S.L., Winters, N.I., Calvi, C.L.,
Peter, L., Chung, M.-I., Taylor, C.J., Jetter, C., et al. (2020). Single-cell RNA
sequencing reveals profibrotic roles of distinct epithelial and mesenchymal lineages in
pulmonary fibrosis. Sci. Adv. *6*, eaba1972. https://doi.org/10.1126/sciadv.aba1972.

109. Vieira Braga, F.A., Kar, G., Berg, M., Carpaij, O.A., Polanski, K., Simon, L.M.,
Brouwer, S., Gomes, T., Hesse, L., Jiang, J., et al. (2019). A cellular census of human
lungs identifies novel cell states in health and in asthma. Nat. Med. *25*, 1153–1163.
https://doi.org/10.1038/s41591-019-0468-5.

110. Mayr, C.H., Simon, L.M., Leuschner, G., Ansari, M., Schniering, J., Geyer, P.E.,
Angelidis, I., Strunz, M., Singh, P., Kneidinger, N., et al. (2021). Integrative analysis of
cell state changes in lung fibrosis with peripheral protein biomarkers. EMBO Mol. Med.
13, e12871. https://doi.org/10.15252/emmm.202012871.

111. Xu, C., Lopez, R., Mehlman, E., Regier, J., Jordan, M.I., and Yosef, N. (2021).
Probabilistic harmonization and annotation of single-cell transcriptomics data with deep
generative models. Mol. Syst. Biol. *17*, e9620. https://doi.org/10.15252/msb.20209620.

112. Gayoso, A., Lopez, R., Xing, G., Boyeau, P., Valiollah Pour Amiri, V., Hong, J., Wu,
K., Jayasuriya, M., Mehlman, E., Langevin, M., et al. (2022). A Python library for
probabilistic analysis of single-cell omics data. Nat. Biotechnol. *40*, 163–166.
https://doi.org/10.1038/s41587-021-01206-w.

113. Luecken, M.D., Büttner, M., Chaichoompu, K., Danese, A., Interlandi, M., Mueller,
M.F., Strobl, D.C., Zappia, L., Dugas, M., Colomé-Tatché, M., et al. (2022).
Benchmarking atlas-level data integration in single-cell genomics. Nat. Methods *19*, 41–
50. https://doi.org/10.1038/s41592-021-01336-8.

114. Lopez, R., Regier, J., Cole, M.B., Jordan, M.I., and Yosef, N. (2018). Deep generative
modeling for single-cell transcriptomics. Nat. Methods *15*, 1053–1058.
https://doi.org/10.1038/s41592-018-0229-2.

115. Wolf, F.A., Angerer, P., and Theis, F.J. (2018). SCANPY: large-scale single-cell gene
expression data analysis. Genome Biol. *19*, 15. https://doi.org/10.1186/s13059-017-
1382-0.
116. Becht, E., McInnes, L., Healy, J., Dutertre, C.-A., Kwok, I.W.H., Ng, L.G., Ginhoux, F.,
and Newell, E.W. (2018). Dimensionality reduction for visualizing single-cell data using
UMAP. Nat. Biotechnol. https://doi.org/10.1038/nbt.4314.
117. Traag, V.A., Waltman, L., and van Eck, N.J. (2019). From Louvain to Leiden:
guaranteeing well-connected communities. Sci. Rep. *9*, 5233.
https://doi.org/10.1038/s41598-019-41695-z.
118. Schupp, J.C., Adams, T.S., Cosme, C., Raredon, M.S.B., Yuan, Y., Omote, N., Poli, S.,
Chioccioli, M., Rose, K.-A., Manning, E.P., et al. (2021). Integrated Single-Cell Atlas of
Endothelial Cells of the Human Lung. Circulation *144*, 286–302.
https://doi.org/10.1161/CIRCULATIONAHA.120.052318.
119. Sikkema, L., Strobl, D., Zappia, L., Madissoon, E., Markov, N.S., Zaragosi, L., Ansari,
M., Arguel, M., Apperloo, L., Bécavin, C., et al. (2022). An integrated cell atlas of the
human lung in health and disease. Preprint at bioRxiv,
https://doi.org/10.1101/2022.03.10.483747 https://doi.org/10.1101/2022.03.10.483747.
120. Büttner, M., Ostner, J., Müller, C.L., Theis, F.J., and Schubert, B. (2021). scCODA is a
Bayesian model for compositional single-cell data analysis. Nat. Commun. *12*, 6876.
https://doi.org/10.1038/s41467-021-27150-6.
121. Love, M.I., Huber, W., and Anders, S. (2014). Moderated estimation of fold change and
dispersion for RNA-seq data with DESeq2. Genome Biol. *15*, 550.
https://doi.org/10.1186/s13059-014-0550-8.
122. Squair, J.W., Gautier, M., Kathe, C., Anderson, M.A., James, N.D., Hutson, T.H.,
Hudelle, R., Qaiser, T., Matson, K.J.E., Barraud, Q., et al. (2021). Confronting false
discoveries in single-cell differential expression. Nat. Commun. *12*, 5692.
https://doi.org/10.1038/s41467-021-25960-2.
123. Ignatiadis, N., Klaus, B., Zaugg, J.B., and Huber, W. (2016). Data-driven hypothesis
weighting increases detection power in genome-scale multiple testing. Nat. Methods *13*,
577–580. https://doi.org/10.1038/nmeth.3885.
124. Holland, C.H., Tanevski, J., Perales-Patón, J., Gleixner, J., Kumar, M.P., Mereu, E.,
Joughin, B.A., Stegle, O., Lauffenburger, D.A., Heyn, H., et al. (2020). Robustness and
applicability of transcription factor and pathway analysis tools on single-cell RNA-seq
data. Genome Biol. *21*, 36. https://doi.org/10.1186/s13059-020-1949-z.
125. Jiang, P., Zhang, Y., Ru, B., Yang, Y., Vu, T., Paul, R., Mirza, A., Altan-Bonnet, G.,
Liu, L., Ruppin, E., et al. (2021). Systematic investigation of cytokine signaling activity
at the tissue and single-cell levels. Nat. Methods *18*, 1181–1191.
https://doi.org/10.1038/s41592-021-01274-5.
126. Badia-i-Mompel, P., Vélez Santiago, J., Braunger, J., Geiss, C., Dimitrov, D., Müller-
Dott, S., Taus, P., Dugourd, A., Holland, C.H., Ramirez Flores, R.O., et al. (2022).
decoupleR: ensemble of computational methods to infer biological activities from omics
data. Bioinforma. Adv. *2*, vbac016. https://doi.org/10.1093/bioadv/vbac016.
127. Müller-Dott, S., Tsirvouli, E., Vazquez, M., Ramirez Flores, R.O., Badia-I-Mompel, P.,
Fallegger, R., Türei, D., Lægreid, A., and Saez-Rodriguez, J. (2023). Expanding the
coverage of regulons from high-confidence prior knowledge for accurate estimation of
transcription factor activities. Nucleic Acids Res. *51*, 10934–10949.
https://doi.org/10.1093/nar/gkad841.
128. Seabold, S., and Perktold, J. (2010). Statsmodels: Econometric and Statistical Modeling
with Python. scipy. https://doi.org/10.25080/Majora-92bf1922-011.
129. Sun, D., Guan, X., Moran, A.E., Wu, L.-Y., Qian, D.Z., Schedin, P., Dai, M.-S.,

Danilov, A.V., Alumkal, J.J., Adey, A.C., et al. (2022). Identifying phenotype-associated
subpopulations by integrating bulk and single-cell sequencing data. Nat. Biotechnol. *40*,
527–538. https://doi.org/10.1038/s41587-021-01091-3.

130. Oliveira, M.F., Romero, J.P., Chung, M., Williams, S., Gottscho, A.D., Gupta, A.,
Pilipauskas, S.E., Mohabbat, S., Raman, N., Sukovich, D., et al. (2024). Characterization
of immune cell populations in the tumor microenvironment of colorectal cancer using
high definition spatial profiling. Preprint at bioRxiv,
https://doi.org/10.1101/2024.06.04.597233 https://doi.org/10.1101/2024.06.04.597233.

131. Palla, G., Spitzer, H., Klein, M., Fischer, D., Schaar, A.C., Kuemmerle, L.B., Rybakov,
S., Ibarra, I.L., Holmberg, O., Virshup, I., et al. (2022). Squidpy: a scalable framework
for spatial omics analysis. Nat. Methods *19*, 171–178. https://doi.org/10.1038/s41592-
021-01358-2.

132. Marconato, L., Palla, G., Yamauchi, K.A., Virshup, I., Heidari, E., Treis, T., Vierdag,
W.-M., Toth, M., Stockhaus, S., Shrestha, R.B., et al. (2024). SpatialData: an open and
universal data framework for spatial omics. Nat. Methods.
https://doi.org/10.1038/s41592-024-02212-x.

133. Salim, A., Bhuva, D.D., Chen, C., Yang, P., Davis, M.J., and Yang, J.Y. (2024).
SpaNorm: spatially-aware normalisation for spatial transcriptomics data. Preprint at
bioRxiv, https://doi.org/10.1101/2024.05.31.596908
https://doi.org/10.1101/2024.05.31.596908.

134. Biancalani, T., Scalia, G., Buffoni, L., Avasthi, R., Lu, Z., Sanger, A., Tokcan, N.,
Vanderburg, C.R., Segerstolpe, Å., Zhang, M., et al. (2021). Deep learning and
alignment of spatially resolved single-cell transcriptomes with Tangram. Nat. Methods
18, 1352–1362. https://doi.org/10.1038/s41592-021-01264-7.

135. Song, D., Xi, N.M., Li, J.J., and Wang, L. (2022). scSampler: fast diversity-preserving
subsampling of large-scale single-cell transcriptomic data. Bioinforma. Oxf. Engl. *38*,
3126–3127. https://doi.org/10.1093/bioinformatics/btac271.

136. Birk, S., Bonafonte-Pardàs, I., Feriz, A.M., Boxall, A., Agirre, E., Memi, F., Maguza,
A., Fan, R., Castelo-Branco, G., Theis, F.J., et al. (2024). Quantitative characterization
of cell niches in spatial atlases. Preprint at bioRxiv,
https://doi.org/10.1101/2024.02.21.581428 https://doi.org/10.1101/2024.02.21.581428.

137. Türei, D., Valdeolivas, A., Gul, L., Palacio-Escat, N., Klein, M., Ivanova, O., Ölbei, M.,
Gábor, A., Theis, F., Módos, D., et al. (2021). Integrated intra- and intercellular
signaling knowledge for multicellular omics analysis. Mol. Syst. Biol. *17*, e9923.
https://doi.org/10.15252/msb.20209923.

138. Browaeys, R., Saelens, W., and Saeys, Y. (2020). NicheNet: modeling intercellular
communication by linking ligands to target genes. Nat. Methods *17*, 159–162.
https://doi.org/10.1038/s41592-019-0667-5.

139. Sato, T., and Clevers, H. (2013). Growing self-organizing mini-guts from a single
intestinal stem cell: mechanism and applications. Science *340*, 1190–1194.
https://doi.org/10.1126/science.1234852.

140. Fumagalli, A., Suijkerbuijk, S.J.E., Begthel, H., Beerling, E., Oost, K.C., Snippert, H.J.,
van Rheenen, J., and Drost, J. (2018). A surgical orthotopic organoid transplantation
approach in mice to visualize and study colorectal cancer progression. Nat. Protoc. *13*,
235–247. https://doi.org/10.1038/nprot.2017.137.

141. Hao, Y., Stuart, T., Kowalski, M.H., Choudhary, S., Hoffman, P., Hartman, A.,
Srivastava, A., Molla, G., Madad, S., Fernandez-Granda, C., et al. (2024). Dictionary
learning for integrative, multimodal and scalable single-cell analysis. Nat. Biotechnol.
42, 293–304. https://doi.org/10.1038/s41587-023-01767-y.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7

Figure S1: related to Figure 1. Composition of CRC atlas

D

cell_type
- B cell
- Dendritic cell
- Endothelial cell
- Epithelial cell
- Mast cell
- Monocyte/Macrophage
- NK cell
- Neutrophil
- Plasma cell
- Stromal cell
- T cell

sample_type
- tumor
- adjacent normal
- healthy normal
- blood
- metastasis
- polyp
- lymph node

crc_type
- COAD
- READ
- CRC type unknown
- non-cancer

E

cell_type
- Monocyte
- Macrophage
- Dendritic cell
- B cell
- Plasma IgA
- Plasma IgG
- Neutrophil
- Mast cell
- T cell CD4
- T cell regulatory
- T cell CD8
- NK cell
- gamma-delta

F

platform
- 10x 3p
- 10x 5p
- BD Rhapsody
- DNBelab C4
- GEXSCOPE Singleron
- SMARTer (C1)
- Smart-seq2
- TruDrop
- scTrio-seq2

Figure S1: related to Figure 1. Composition of CRC atlas

Figure S2: related to Figure 2. Tumor immune phenotypes in CRC. (**A**) Immune phenotypes and clinical, genetic features. (**B**) Results of the Progeny analysis. (**C**) Results of the cytosig analysis.

A

B

Figure S3: related to Figure 3. Association of cellular composition and distinct genotypes and survival in the TCGA data. (**A**) Association of cellular composition with MSI status in CRC patients (n = 177). (**B**) Association of cellular composition with MSS status in CRC patients (n = 412).

A

Figure S4: related to Figure 4. Characterization of neutrophils in CRC. (A) Selected multiplex immunofluorescence (mpIF) staining of LOX-1 (pink), CD15 (yellow), pan-cytokeratin (turquoise), and DAPI (blue) in colorectal cancer (CRC) tumor tissue. Scale bar: 100 μm.

Figure S5: related to Figure 5. Spatial multi-modal profiling of neutrophils. (**A**) Cell type annotation of the spatial transcriptomics data. (**B-C**) Localization of the T cells and B cells (representatively shown for one patient). (**D**) IHC of neutrophils (**F-E**) Cell niches at a resolution of two and 3, respectively.

G

H

Figure S5: related to Figure 5. Spatial multi-modal profiling of neutrophils. (**G**) Neutrophil niche composition in the tumor core. (**H**) Neutrophil niche composition at the invasive margin.

Figure S6: related to Figure 6. Orthotopic mouse model for CRC. (**A**) Orthotopic transplantation of AKPS organoids. (**B**) Tumor weights and tumor volumes of the mice. (**C**) Picture of the respective tumors. (**D**) Neutrophils in the bone marrow of the mice after 6 weeks. (**E**) Representative plots of the neutrophils in the bone marrow

Figure S7: related to Figure 7. Tumors are reprogramming neutrophil metabolism. (**A**) Human-mouse comparisons of the expression profiles. Gene Set Enrichment Analysis (GSEA) on KEGG terms for (**B**) the expression profiles of bone marrow neutrophils from AKPC mice and controls, and (**C**) the expression profiles of blood neutrophils from healthy volunteers and patients with CRC. (**D**) XF Neutrophil Activation Assay kinetic traces of oxygen consumption rate (OCR) of healthy human neutrophils after 24h culture in SW900 supernatant, or CaCo2 supernatant or RPMI control medium. Mitochondrial function is blocked by Rot/AA injection, oxidative burst is activated by PMA injection. The data shown are the mean and SD, n=3 biological replicates.

Figure S8: related to STAR Methods. Flow cytometry gating strategy to define cell populations from CRC tumor tissue, normal adjacent tissue and whole blood samples. In initial cleaning steps dead cells, debris and doublets were removed using 7-AAD staining and scatter characteristics. Leukocytes were defined by CD45 staining and sequentially gated into subtypes including neutrophils, monocytes, T cells and B cells. Non-CD45+ cells were gated into epithelial cells, endothelial cells and fibroblasts.