# What lurks in the dark? - An innovative framework for studying diverse wild insect microbiota

Karol H. Nowak[1,2*], Emily Hartop[3], Monika Prus-Frankowska[1], Mateusz Buczek[1], Michał Kolasa[1,4], Tomas Roslin[5,6], Otso T. Ovaskainen[7,8], Piotr Łukasik[1*]

1) Institute of Environmental Sciences, Faculty of Biology, Jagiellonian University, Kraków, Poland

2) Doctoral School. PhD Programme in Biology, Faculty of Biology, Jagiellonian University, Kraków, Poland

3) Department of Natural History, NTNU University Museum, Norwegian University of Science and Technology, Trondheim, NO-7491, Norway

4) Department of Farm Animal Biodiversity Conservation and Horse Breeding, National Research Institute of Animal Production, Balice, Poland

5) Department of Agricultural Sciences, University of Helsinki, Helsinki, Finland

6) Department of Ecology, Swedish University of Agricultural Sciences, Uppsala, Sweden

7) Department of Biological and Environmental Science, Faculty of Mathematics and Science, University of Jyväskylä, Jyväskylä, Finland

8) Organismal and Evolutionary Biology Research Programme, Faculty of Biological and Environmental Sciences, University of Helsinki, Helsinki, Finland

* To whom correspondence should be addressed.

E-mails: karol.h.nowak@doctoral.uj.edu.pl, p.lukasik@uj.edu.pl

All supporting data available at: https://github.com/KarolHub/Phorid-Microbiota.

## Abstract

Symbiotic microorganisms can profoundly impact insects, their life history traits, population dynamics, and evolutionary trajectories. However, microbiota remain poorly understood in natural insect communities, especially in 'dark taxa' - hyperdiverse, yet understudied clades.

Here, we implemented a novel multi-target amplicon sequencing approach to study microbiota in complex, species-rich communities. It combines four methodological innovations: (1) To establish a host taxonomic framework, we sequenced amplicons of host marker gene (COI) and reconstructed barcodes alongside microbiota characterisation. (2) To assess microbiota abundance, we incorporated synthetic spike-in-based quantification. (3) To improve the phylogenetic resolution for the dominant endosymbiont, *Wolbachia*, we analysed bycatch data from the COI amplicon sequencing. (4) To investigate the primary drivers of host-microbe associations in massive multi-dimensional dataset, we performed HMSC modelling.

Applying this approach to 1,842 wild-caught scuttle flies (Diptera: Phoridae) from northern Sweden, we organised them into 480 genotypes and 186 species, and gained unprecedented insights into their microbiota. We found orders-of-magnitude differences in bacterial abundance and massive intrapopulational variation in microbiota composition. Differences emerged in patterns and drivers among microbial functional categories: the distribution and abundance of facultative endosymbionts (*Wolbachia*, *Rickettsia*, *Spiroplasma*) were shaped by host species, genotype and sex. In contrast, many other bacterial taxa were broadly distributed across species and sites.

This study highlights facultative endosymbionts as key players in insect microbiota and reveals striking variation in distributional patterns of microbial clades. It also demonstrates the power of integrative sequencing approaches in uncovering the ecological complexity and significance of symbiotic microorganisms in multi-species natural communities.

**Keywords:** Arthropods, microbiome, endosymbiosis, next-generation sequencing, holobiont, Hierarchical Modelling of Species Communities, HMSC, joint species distribution models

## 1. Introduction

Microbial symbioses have played tremendously important roles in the biology and evolution of higher organisms [1]. The many levels at which different functional categories of symbionts have shaped hosts' biology are relatively well-understood in insects [2]. The range of microbial symbionts' effects on insect life-history traits - nutrition, reproduction, and defence [3,4] translate to population processes and patterns [5–7], host evolutionary trajectories [8,9], and effects at the level of multi-species communities [10]. There is little doubt that through such combined effects, microbiota have shaped the global biodiversity, and will continue to play key roles in ecosystems rapidly changing under anthropogenic pressures [11,12].

Despite their massive importance, insect microbiota are poorly known outside of model species. Some biological systems, including honey bees [13] or pea aphids [14] have been the focus of extensive research on symbiosis diversity, function, mechanisms, and various other aspects. However, for some 6 million of the remaining insect species on our planet [15,16], our understanding of microbiota is non-existent or, at best, very limited. Among the primary reasons is the sheer diversity of insects and microorganisms, which poses a significant challenge in generating and handling vast amounts of multidimensional ecological data. We lack reliable and cost-effective workflows for processing large numbers of wild-caught insects, encompassing host identity validation and microbiome characterisation, while including controls for different types of contamination [17–19]. Additionally, existing approaches rarely consider and address questions about microbial abundance, which is critical, as insects differ dramatically in microbial load [20]. Data analysis is another neglected area, as the value of conventional approaches, including diversity estimation and comparison, is questionable when microbial communities are dominated by one or few strains or when

69    the abundance is low and the contamination is not controlled. Finally, analytical solutions for the

70    reconstruction of microbiome patterns and drivers in highly multidimensional datasets are lacking.

71            As a result, we know very little about the microbial composition and abundance across the

arthropod diversity. Host biology, especially the diet, has been identified as an important driver [8],

alongside host taxonomy [21]. However, even within a species, there can be major differences among

populations, within a population over time, and among individuals sampled at the same time from a

single population [22]. Similarly, the differences in microbiota between host sexes have rarely been

considered outside of the context of reproductive manipulation [23,24]. Interactions among these and

other possible microbiota drivers can make analyses especially challenging. At the same time, we

would anticipate that microbial taxa and functional categories would differ in how the infections are

determined. A systematic assessment of these and other host-microbe interaction drivers would require

a combination of comprehensive laboratory methods applied at a large scale and robust data analysis

approaches, but also suitable study organisms.

82            Much of the global insect diversity is made up of so-called 'dark taxa': insect clades that are

hyperdiverse, abundant, and broadly distributed yet remain poorly studied [25]. The study of dark taxa

has greatly benefitted from recent advances in molecular workflows [26,27], particularly in

Megabarcoding - high-throughput barcoding of single specimens [28], which has enabled large-scale

analyses of hyperdiverse insects on a broader scale than previously possible [29]. Growing barcoding

data provide a promising opportunity to illuminate the hidden microbial diversity across these

enigmatic clades.

89            Scuttle flies (Diptera: Phoridae) are a quintessential example of a "dark taxon": they are

hyperdiverse, abundant, and globally distributed but remain poorly understood [25,30]. They exhibit

diverse life strategies, including parasitism, predation, herbivory, and saprophagy [31]. Such varied life

histories allow them to occupy numerous ecological niches, from bustling urban centres [32] and

subterranean caves [33] to intertidal zones [34] and the cavities of pitcher plants [35]. Despite their

ubiquity and diverse ecological roles, our understanding of their diversity, distribution, and biology

remains grossly incomplete - highlighted by recent reporting of more species captured over eight

weeks at a single Ugandan site than had been previously described from the entire Afrotropical

bioregion [26]. Given our poor state of knowledge about scuttle flies generally, it is not surprising that

little is known about their interactions with microbial symbionts, with the few studies that addressed

questions about these associations limited to single-species investigations with shallow sampling depth

[36,37].

In this study, we leverage a novel combination of molecular and analytical approaches to

comprehensively characterise microbiota abundance, composition, and distribution in diverse scuttle

flies as a model dark taxon across six sites in Northern Sweden. We employ high-resolution COI

amplicon sequencing for host identification, bacterial 16S-V4 rRNA sequencing to resolve microbiota

composition, using synthetic spike-ins to quantify microbial abundance, and state-of-the-art

Hierarchical Modelling of Species Communities (HMSC) to disentangle the ecological drivers of

microbial variation. By integrating these innovative techniques, we assess how microbial communities

are shaped by site, season, host species identity, and individual host traits such as sex and genotype.

This innovative methodological pipeline provides an unprecedented resolution of host-microbiota

interactions in a hyperdiverse insect group, advancing our understanding of the ecological and

evolutionary forces shaping microbial associations in wild communities.

## 2. Materials and methods

*2.1. Field collections*

Samples were collected using Malaise traps as a part of the Swedish Insect Inventory Project

(SIIP) [38]. The collection was carried out in northern Sweden at six sites, separated by distances

varying from less than 50 m to about 390 km (for exact locations, see Supplementary Table S1 and

Fig. 1). Selected samples come from late spring (June to July) and summer (July to August) of 2018

(Supplementary Table S1). After collection, samples in 95% ethanol were stored at -20 °C in Station

Linné, Sweden, where taxonomists manually sorted the scuttle flies from the samples. We randomly

selected 88 males and 88 females for processing from each of the six sites and two seasons (except for

three combinations where only 87, 86 and 84 individuals were available; see Supplementary Table S1

for details), totalling 2,105 insects.

*2.2 DNA extraction, amplicon library preparation and sequencing*

The insects were processed and prepared for sequencing in the Molecular Ecology Laboratory

at the Institute of Environmental Sciences of Jagiellonian University, following a protocol

comprehensively described recently [39]. First, each insect was placed in a 2 ml tube with 190 µl of

the lysis buffer (0.4 M NaCl, 10 mM Tris-HCl, 2 mM EDTA, 2% SDS), 10 µl proteinase K and

different-sized ceramic beads. After two rounds of 30-second homogenisation on a homogenizer (Bead

Ruptor Elite, Omni International), the samples were incubated for 2 hours at 55 °C. Subsequently, 40

µl aliquot of the homogenate was combined with 2 µl of the solution containing 10,000 or 20,000

copies (depending on the batch) of plasmids carrying artificial 16S rRNA amplification targets,

Ec5502 [40], placed in a 96-well PCR plate, and purified using SPRI beads [41]. DNA was eluted in

20 µl of TE buffer.

The amplicon libraries were prepared using a two-step PCR procedure (see method 4 [42]). In

the first step, two marker regions of interest were simultaneously amplified using QIAGEN Multiplex

Master Mix and template-specific primers with variable-length inserts and Illumina adapter stubs.

Specifically, for the V4 region of bacterial 16S rRNA, we used primers 515F:

GTGYCAGCMGCCGCGGTAA and 806R: GGACTACNVGGGTWTCTAAT [43], and for a 418 bp

portion of the mitochondrial cytochrome oxidase I (COI) gene, primers BF3:

CCHGAYATRGCHTTYCCHCG and BR2: TCDGGRTGNCCRAARAAYCA [44]. The PCR

reaction mix also contained 500 copies of another plasmid carrying an artificial 16S rRNA target,

Ec5001 [40], but it was not used in this study and subsequently filtered out from the data. SPRI bead-

purified first-step PCR products were used as a template for the second indexing PCR. The libraries

were verified on an agarose gel and based on band intensities pooled approximately equimolarly. The

purified pool was quality checked using BioAnalyzer 2100 (Agilent), combined with pools

representing other projects, and submitted to Swedish National Genomics Infrastructure for sequencing

on Illumina NovaSeq SPrime flow cells (2x250 bp read mode). To control for contamination from

different sources, for each of the 88 biological samples processed, we used eight different controls:

negative controls for the DNA extraction (3), first PCR (2), and indexing PCR stages (1), as well as

positive controls (2). Raw sequence data for all samples that passed filtering criteria and controls are

available under GenBank BioProject accession number PRJNA1099311.

## 152 *2.3. Bioinformatic analysis of amplicon data*

153     The amplicon data were processed using scripts and files available at the GitHub repository,

154     together with supplementary data: https://github.com/KarolHub/Phorid-Microbiota.

### 155 *2.3.1. Basic read sorting and processing*

Initially, we split the demultiplexed reads into batches corresponding to COI and 16S-V4

targets using a custom Python script (Multisplit.py), which divided the files based on primer sequences

and trimmed the primers. Further steps relied on the custom LSD.py script, a wrapper tool executing

several steps using PEAR [45], VSEARCH [46], and USEARCH [47] programs. Specifically, we

merged forward and reverse reads into contigs, discording those outside of the expected length range

(400-430 for COI and 250-260 bp for 16S-V4), or with lower quality regions (Phred Score $\leq$ 30).

Datasets representing both regions were then dereplicated to retain only unique sequences per sample

and their occurrence counts, with singleton genotypes discarded. We denoised each library separately

using the UNOISE algorithm [48]. Then, we clustered the zero-radius operational taxonomic units

(ZOTUs) into operational taxonomic units (OTUs) using a 97% identity cutoff. Finally, we assigned

taxonomy using the SINTAX algorithm [49], and taxonomic reference databases: MIDORI version GB

239 [50], with alphaproteobacterial sequences added, for the COI data and SILVA [51] version 138

SSU for 16S rRNA data. We retained identification to the lowest level that had at least an 80%

classification confidence score.

### 2.3.2. COI data analysis

The COI data contained a mix of sequences of different origins, including those representing

insect mitochondrial genes, nuclear pseudogenes, alphaproteobacterial symbionts, and others. To

obtain insect barcodes, we discarded sequences shorter or longer than the anticipated 418 nucleotides,

as well as bacterial, non-arthropod, unassigned, and chimeric sequences. Next, we assessed the number

of reads in the most abundant OTU classified as an arthropod and calculated its ratio relative to all

Arthropoda OTUs combined. We excluded libraries from further analyses if they had fewer than 300

reads of the dominant OTU or if the aforementioned read ratio was less than 95%. Then, we designated

the most abundant sequence within the dominant OTU of each sample as the individual's barcode,

dereplicated these barcodes, and clustered them with sequences from earlier barcoding studies [25] to

generate a dendrogram. Given the complex taxonomy of phorid flies, we manually reviewed the

proximity of barcodes and sequences of established phorid species within the dendrogram to determine

which groups of barcodes form distinct species-level clades. Once we had the sequences divided into

species-level clusters of barcodes, we compared them with sequences of named species from the same

study [25], to obtain the taxonomy labels. The species names were assigned if a sequence matched the

reference by at least 97%. Variants with no close matches were compared against the NCBI nt

database using the BLAST algorithm, retaining labels with a match of 97% or higher.

The COI data contained sequences assigned to the order Rickettsiales, which we separated for

further analyses. We filtered out non-*Wolbachia* sequences, chimeric sequences, and those of length

other than 418 bp. This resulted in tables on *Wolbachia* COI ZOTU and OTU read number distribution

among samples, analogous to standard bacterial 16S rRNA-based tables.

*2.3.3. 16S rRNA amplicon data filtering and abundance estimation*

We decontaminated the 16S-V4 data from 1,845 samples with reliable insect COI barcodes

using a custom QUACK.py script. This script classified as contaminants and removed ZOTUs unless

the percentage of reads representing that ZOTU in at least one experimental sample was at least five

times greater than the maximum percentage in any of the blank samples [52,53]. It also removed

sequences classified as Archaea, Eukaryota, chloroplasts, mitochondria and chimaeras. After

calculating the ratio between the spike-in and remaining symbiont reads in each sample, the spike-in

sequences were also removed. Afterwards, we calculated the absolute abundance of non-contaminant

bacterial 16S rRNA copies in each sample by multiplying the symbiont to extraction spike-in read

number ratio, the number of inserted extraction spike-in copies (10,000 or 20,000), and the inverse

proportion of insect homogenate used during the DNA extraction (5). The abundance was estimated

separately for each bacterial ZOTU, with the total bacterial abundance in the sample estimated as the

summed abundance of individual ZOTUs. Three samples without extraction spike-in reads were

discarded, along with control samples, resulting in the final total number of phorid samples of 1842.

For subsequent comparisons and modelling, we used tables on bacterial OTU and ZOTU estimated

16S rRNA copy number distributions across samples.

*2.3.4. Analyses*

To examine the drivers of variation in insect microbiomes we used Hierarchical Modelling of

Species Communities (HMSC; [54,55], a method of joint species distribution modelling (JSDM; [56]).

We modelled 75 species that were represented by at least 5 individuals for a total of 1634 samples. We

fitted separate HMSC models for each of three response variables: (1) estimated copy numbers for the

50 most abundant 16S-V4 OTUs, (2) estimated copy numbers for the 100 most dominant 16S-V4

ZOTUs, and (3) read counts of the 24 most dominant Wolbachia COI ZOTUs. For covariates, we used

effects of season (spring vs. summer), site (n = 6), sex of the host (male vs. female), host species (n =

75), host genotype (n = 328), and sampling unit (n = 1,634). This approach allowed us to quantify how

these ecological and host-related factors shape microbial community composition and abundance

across a hyperdiverse insect assemblage.

Because our data were zero-inflated—with many taxa being absent from individual samples,

resulting in numerous zero counts—we used a hurdle modelling approach to separately analyse

presence-absence and abundance. First, we modelled the presence-absence of each taxon using a

Bernoulli distribution with a probit link function. Then, for taxa that were present, we modelled

abundance conditional on presence by log-transforming and scaling the response, assuming a normal

distribution. In our models, season and sex were included as fixed effects, while site, host species, host

genotype, and individual host insect were treated as community-level random effects, structured using

latent variables to account for shared variation across taxa. This approach allowed us to better capture

the ecological and host-related factors influencing microbiota distributions.

We fitted the model using the Markov Chain Monte Carlo (MCMC) approach implemented in

the R-package HMSC [57] with default prior distributions [55]. The posterior distribution was sampled

with four chains, with 375 * thin MCMC iterations per chain. We ignored the first 125 * thin iterations

as burn-in and thinned the remaining iterations by thin to yield 250 samples per chain and hence 1000

samples in total. The thinning parameter was increased as thin = 1, 10, 100, … until satisfactory

MCMC convergence was reached. We considered MCMC convergence satisfactory if the third quartile

of the potential scale reduction factors over the parameters measuring the responses of the taxa to

season and sex was at most 1.05; this was reached with thin = 1000 for 16S-V4 ZOTU and *Wolbachia*

COI ZOTU presence-absence models, and with thin = 100 for the other four model components.

To evaluate model fit, we focused on explanatory and predictive powers – the former based on

fitting the models to all data and the latter through two-fold cross-validation. As metrics for evaluation,

we used AUC for presence-absence models and $R^2$ for models on abundance conditional on presence.

To examine the importance of the fixed and random effects in structuring microbiome variation, we

applied a variance partitioning approach to the explained variation. In other words, we examined what

proportion of the overall variation explained by the model could be attributed to the individual factors

and covariates. To examine the effects of sex and season, we counted the proportion of taxa that

showed either a positive or negative response with at least 95% posterior probability. To examine

patterns of co-variation not explained by sex or season, we examined which pairs of bacterial taxa

showed a positive or negative association for each random effect (site, species, genotype and

individual) with at least 90% posterior probability. The latter approach will reveal what microbial taxa

appear to covary in their presence and abundance beyond what can be expected based on the fixed

effects of sex and season.

## 3. Results

After stringent quality filtering of 2,105 amplicon sequencing libraries for COI and 16S-V4

marker regions of individual phorid flies, we retained on average 77 individuals for each of the 24 site-

season-sex combinations: 936 males and 906 females, or 1,842 individuals in total (Supplementary

Table S1). For individuals passing the filtering criteria, we obtained, on average, 27.1K (K = thousand)

COI reads (min 373, max 191.9K, median 20.4K) (Supplementary File S1). We then used these data to

reconstruct insect barcodes and classify specimens to species (Supplementary Table S2; see section

3.1), but also to reconstruct and compare COI sequences of *Wolbachia* (see section 3.3). We obtained,

on average, 29.7K 16S-V4 reads (range 682 - 295.7K, median 19.5K) per experimental library, but

after filtering and removal of putative laboratory and reagent-derived contaminants and removing

spike-in control reads, we retained, on average, 15.2K 16S-V4 non-contaminant reads (range 29 -

227K, median 5.4K) (Supplementary Table S3). We used these data to reconstruct microbial

abundance and diversity across individuals (section 3.2). For statistical modelling of the data (section

3.4) we used all data types listed above, focusing on the 75 abundant species (for specific cut-offs, see

Methods).

## *3.1. Abundance, diversity and distribution of phorid flies*

Of the COI amplicon data for individual phorid flies, around 99.5% of reads were classified as matching

insects. On average, the most abundant insect ZOTU in each library, designated as the individual's

barcode, accounted for 98.8% of the filtered insect reads (min 52.2%, max 100%, median 99.5%). The

COI barcodes of 1,842 phorid individuals represented 480 genotypes, which we grouped into 186

species (Supplementary Table S2) through a comparison with previously sequenced and

morphologically characterised material [25]. These species varied in genotype numbers and abundance

(Fig. 1a). For example, the five most abundant species were represented by between 10 and 33

genotypes each (85 genotypes in total), but 59 of these genotypes were represented by a single

individual only. Many of these low-abundance genotypes differed from the more abundant genotypes

by a single nucleotide. 142 of the COI genotypes, representing 438 individuals, were confidently

assigned to 63 recognised, named species, mostly within the genus *Megaselia*. A further 42 genotypes

were assigned to two species complexes: the *Megaselia pulicaria* complex (132 individuals) and the

*Megaselia giraudii* complex (82 individuals). Most of the samples (295 genotypes, 1176 individuals)

were assigned to the genus level, and one genotype consisting of 14 individuals was assigned to the

family level only. Given the low prevalence of individual genotypes, there was little indication of

geographic structuring within species (Fig. 1, Supplementary Table S2). To allow unequivocal

reference to each species (whether recognised or not), we assigned unique numbers to all species -

generally, numbering them consecutively after sorting them by decreasing abundance, although

subsequent changes to the sample set caused some discrepancies. We are referring to species by these

numbers throughout the text (e.g., "sp. 1" and "sp. 15" - rather than "*Megaselia* sp. 001" and

"*Megaselia hendersoni*") (Fig. 1a, Supplementary Table S2).

Species' representation in the dataset was highly skewed, with a few abundant species and a

long tail of rare ones (Supplementary Table S2). Within the same season, we found an overlapping set

of abundant species across all sites (Fig. 1b). For example, the five most abundant species were found

at all six sites. At the same time, there was little overlap in community composition among seasons,

with most species well-represented either in spring or in the summer (Fig. 1b). NMDS based on Bray-

Curtis distances clearly separated spring and summer community samples (Supplementary Fig. S1).

**(a) Host diversity;** Host genotypes clustered into species-level clades

**(b) Host distribution;** Geographical locations, sampling periods and species composition of the host communities

**Fig. 1:** (**a**) Haplotype network of the 25 most abundant scuttle fly species highlights the diversity among the sampled insects and showcases challenges with their delimitation. Individual pie charts represent fly COI genotypes, coloured by collection site and season. Selected species are represented

296   by photos of individually barcoded individuals from various sites, with all images scaled uniformly for

297   consistency. Genotypes of the remaining 161 species are not shown. The network was drawn using

298   PopArt [58]. (**b**) The distribution and diversity of the 25 most commonly sampled scuttle fly species

299   from Northern Sweden highlights community composition variation among the two seasons and

300   overlap among the six sampled sites.

## 3.2. Abundance and composition of phorid microbiota

Across 1,842 phorid samples, a total of 28M (M = million) reads scored as non-contaminants

and represented 58.1K ZOTUs that clustered into 9.6K 97% OTUs (Supplementary File S3). Using the

ratio of quantification spike-in to non-contaminant reads as an estimate of the abundance of specific

bacterial ZOTUs, OTUs, and total bacteria [39,40], we were able to assess and interpret microbial

abundance and diversity at the same time.

In terms of microbe abundance, we observed high variation in the symbiont to spike-in ratios

across the samples. The ratio ranged between 0.015 and 354.8 (median 1.38), and with 10K or 20K

artificial template spike-in copies added to one-fifth of insect homogenate before purification, these

ratios translate to bacterial 16S rRNA copy numbers per insect ranging between 766 and 24M (median

815.4K) (Supplementary File S3). Even among individuals representing the same sex and species, the

range in 16S rRNA copy numbers typically spanned between two to four orders of magnitude. At the

same time, microbial abundance was often higher in females than males of the same species, by about

an order of magnitude on average (Fig. 2a).

**Fig. 2:** Simultaneous inspection of bacterial absolute abundance and microbial diversity data indicates strong variation across 1,193 specimens of the 25 dominant scuttle fly species. (**a**) The distribution of bacterial 16S-V4 rRNA copy numbers across individual insects, estimated based on quantification spike-ins added during the DNA extraction step. Here, each individual fly is represented by a narrow vertical bar, with bars sorted first by species assignment, then by sex, and then by abundance. (**b**) The relative abundances of the six most abundant bacterial 97% OTUs across individual insects sorted as in panel **a**. The category 'other bacteria' shows all the remaining bacterial OTUs combined.

Although the number of detected 16S-V4 97% OTUs was high, scuttle fly communities were dominated by relatively few bacterial clades (Supplementary Table S4). The most common OTU1, taxonomically assigned to the genus *Wolbachia*, was represented by 30.3% of decontaminated bacterial reads in a library. The next-most prevalent symbiont OTUs, *Serratia*, *Rickettsia*, *Providencia*, *Carnobacterium,* and *Pseudomonas,* comprised an average of 5.5%, 5.1%, 3.0%, 2.9%, and 2.7% of reads in a library, respectively. We identified multiple additional, less abundant bacterial OTUs representing the same six genera plus a long list of other genera (Supplementary Table S4). These OTUs were broadly distributed across species, but their abundances showed non-random distribution

patterns among individuals within species (Table 1; Fig. 2b). For example, *Rickettsia* comprised a

particularly large share of the microbiota of females of species 1 with high overall bacterial abundance.

The analysis of abundance and distribution of dominant bacterial clades revealed large

variation among individuals, even of the same species, population, and sex - as well as the complex

interactions among likely drivers of microbial abundance and composition (Fig. 3). For example, when

present, *Rickettsia* can be a quantitatively dominant microbe in females of phorid sp. 1, but its

prevalence varies widely among populations. At a specific site (M38), *Providencia* is common and

abundant in both sexes of phorid sp. 1 and sp. 2, but it is much less common or abundant at other sites

or in co-occurring phorid sp. 3.

**Fig. 3:** The comparison of total bacterial abundance and relative abundance of dominant bacterial OTUs showcased the extent and patterns of variation among three most abundant phorid species, two sexes, and six sites. Each panel represents a particular species-sex-site combination, with the top bars indicating bacterial abundance and the lower stacked bars indicating the relative abundance of the most abundant bacterial 16S OTUs (as in Fig. 2). Individual columns of varying width represent individual flies.

Inspection of the data at higher phylogenetic resolution - 16S rRNA ZOTUs instead of 97%

OTUs - provided additional information about the host-microbe association patterns. Bacterial OTUs

typically comprise several relatively abundant genotypes (ZOTUs), and these OTUs differ in how their

component ZOTUs are distributed across host species (Fig. 4). Specifically, insect species were found

to preferentially associate with only one or two specific 16S rRNA genotypes of facultative

endosymbiont *Wolbachia* and *Rickettsia*. Different species associate with different facultative

endosymbiont genotypes, and conversely, different facultative endosymbiont genotypes colonise

different sets of species. In contrast, for most other abundant bacterial OTUs including *Serratia*,

*Providencia*, *Carnobacterium*, and *Pseudomonas*, associations appear much less specific. Most insect

species harbour several genotypes from these OTUs, and genotypes are broadly distributed across host

species (Supplementary File S3).

**Fig. 4**: The distributions of dominant bacterial genotypes (ZOTUs) within most abundant OTUs across

host species reveal differences in specificity among bacterial functional categories: facultative

endosymbionts (denoted with an asterisk) form relatively specific associations with hosts, unlike other

microbes. Insect individuals, represented by columns, are sorted by species, sex, and decreasing

bacterial abundance (as in Fig. 2). Rows represent bacterial genotypes (ZOTUs) sorted into 97%

OTUs, and black lines indicate symbiont genotypes represented by at least 100 copies of 16S rRNA in

specific insect individuals.

We noted that cytochrome oxidase (I) amplicon data for Alphaproteobacteria provide even

greater phylogenetic resolution. Specifically, *Wolbachia* comprised most of the 0.5% of non-insect

COI sequences in the dataset. Based on these COI data, *Wolbachia* was represented by 2 or more reads

and thus scored as present in 862 (47%) of samples, with an average of 269 reads per sample (max

4,875, median 100). These *Wolbachia* COI reads represented 258 unique ZOTUs grouped into 23 97%

OTUs. A comparison of the 16S-V4 and COI data revealed > 90% agreement among the two datasets

regarding the *Wolbachia* presence in the samples, after applying cutoffs likely to exclude much of

likely cross-contamination (3 COI reads vs 400 reads of 16S rRNA) (Supplementary Table S5).

The comparison of *Wolbachia* COI OTU distributions across phorid species (Fig. 5a)

resembles *Wolbachia* 16S rRNA genotype distribution patterns (Fig. 4). These data show that phorid

species typically associate with a single dominant *Wolbachia* COI OTU and differ considerably in

infection prevalence. COI genotype-level data for both hosts and symbionts (Fig. 5b) provide much

more detailed information: they show that genotypes of the same fly species frequently associate with

different genotypes of *Wolbachia*, with substantial variation among genotypes in infection prevalence.

Interestingly, we observe some of the same *Wolbachia* COI genotypes across genotypes of different

phorid species. For example, *Wolbachia* ZOTU206 infects most genotypes of phorid species 2, at high

prevalence, but also colonizes several genotypes of sp. 7, sp. 9, and some less abundant species.

Within species, we found even greater specificity in host-symbiont interactions with respect to sex and

site (Fig. 5c). For example, in sp. 5., *Wolbachia* COI ZOTU495 colonised most host genotypes, and

tends to achieve higher prevalence in females, only sometimes infecting males. In contrast, *Wolbachia*

ZOTU333 infected all individuals - males and females - of one of the abundant phorid COI genotypes,

and a single processed male of another genotype, while being absent from others. Interestingly, two

other *Wolbachia* genotypes (ZOTU206 & ZOTU486) infected sp. 5 individuals representing different

genotypes, all collected in summer rather than spring.

**Fig. 5:** *Wolbachia* COI amplicon data, analysed at different phylogenetic resolution scales, provides

much more precise information about the patterns of host-symbiont interactions than 16S rRNA

amplicon data. (**a**) The relative abundance of *Wolbachia* COI OTUs among all *Wolbachia* COI reads,

across 25 most abundant scuttle fly species. Individual flies are represented by columns, and sorted by

species, sex and decreasing abundance (as in Fig. 2 and 4); (**b**) The relative abundance of *Wolbachia*

COI genotypes (ZOTUs) across individual flies sorted into species and additionally COI genotypes; (**c**)

The relative abundance of *Wolbachia* genotypes (ZOTUs) among genotypes, sexes, and populations of

sp. 5 (*Megaselia* sp. 004), highlights strong specificity in interactions. In all panels, individuals with

fewer than 2 *Wolbachia* COI reads are considered uninfected and are represented by white columns.

## 3.3. Drivers of microbiota variation

Within our dataset, we observed substantial variation in the prevalence and abundance of microbiota. To quantify the relative contributions of host traits, environmental factors, and microbial resolution levels to this variation, we turned to HMSC analyses. We reconstructed the drivers of presence/absence and abundance conditional on presence at three levels of resolution: 16S rRNA OTUs, 16S rRNA ZOTUs, and for *Wolbachia* only, COI ZOTUs.

### 3.3.1 Results of the HMSC models

Our predictors (site, season, host species, host sex and host genotype) explained a substantial proportion of the variation in bacterial 16S rRNA OTU and ZOTU presence-absence (Supplementary Table S6), with high explanatory and predictive power (AUC $\geq$ 0.69). The effects of host individual and site contributed the most to variance, with microbial taxa generally more prevalent in the spring than summer and more abundant in female than male hosts. For abundance conditional on presence, the explanatory power remained high, but the predictive power was low due to the strong host individual effect (Table S6). Some individuals had a high abundance of most bacterial taxa, while in others, microbial abundance was consistently low for most bacterial taxa, amplifying trends in presence-absence. Unlike presence-absence, abundance was generally higher in summer than in spring, indicating that spring samples contained many taxa in low abundance, whereas summer samples had fewer taxa in high abundance.

For *Wolbachia* COI genotypes, presence-absence was strongly shaped by host species and genotype, with additional effects of season and site. Model performance was exceptionally high (Supplementary Table S6; AUC $\geq$ 0.90), and most strains were more prevalent in spring and in female hosts. *Wolbachia* abundance conditional on presence showed high explanatory power (AUC = 0.94),

but low predictive power, with individual variation driving 36% of the observed differences. Strain-

specific abundances of *Wolbachia* were consistently higher in females than males.

At the level of individual microbial taxa, explained variance (Tjur's $R^2$) varied significantly,

with more prevalent taxa generally showing higher values (Supplementary Fig. S2). The relative

contributions of specific drivers also differed among taxa (Fig. 6).

*3.3.2 Drivers of variation in bacterial clade occurrence and abundance*

The HMSC model explained up to 40% of the variance in the presence-absence of individual

16S-V4 OTUs, with much variation occurring among host individuals (Fig. 6a and Supplementary Fig.

S3a). However, for facultative endosymbionts (*Wolbachia*, *Rickettsia*, *Spiroplasma*), variance was

primarily driven by host species, sex, and genotypes, while *Lactobacillus* showed the strongest

seasonal variation (Fig. 6a and Supplementary Fig. S3a). For abundance conditional on presence, the

model explained up to 70% of variance, with lowest contribution of the host individual factor in

facultative endosymbionts, exhibiting distinct patterns: *Wolbachia* varied most with host species, sex

and genotype, *Rickettsia* showed a strong host genotype effect, and *Spiroplasma* exhibited minimal

sex-based differences.

Similar patterns emerged for the 16S-V4 ZOTUs, where *Wolbachia* ZOTUs showed the

highest Tjur $R^2$ values for both presence/absence and abundance conditional on presence, with

variation attributed to host species and genotype. Facultative endosymbionts differed from other taxa,

in which host individual effects explained most variation (Supplementary Fig. S3b). This was

particularly clear in Actinomycetes, which showed the strongest host individual effect (Supplementary

Fig. S3b).

For *Wolbachia* strains identified via COI (Supplementary Fig. S3c), variance in

presence/absence was primarily shaped by host genotype, species, sex, or site, dependent on the

ZOTU. However, on the overall, a relatively low share of raw variance was explained. Abundance

conditional on presence showed a greater influence of the host individual but remained significantly

affected by host species, genotype, and sex.

**Fig. 6**: Proportion of variance explained per microbial (a) 16S-V4 OTU and (b) 16S-V4 ZOTU by both

the presence-absence and the abundance conditional on presence models, showing that facultative

endosymbionts (denoted with asterisks) are highly influenced by the host sex, species and genotype

factors when compared with other bacteria. For 16S-V4 ZOTUs, 50 of the most abundant ZOTUs are

shown out of 100 included in the models.

*3.3.3 Residual associations among microbial taxa*

After accounting for systematic effects, we examined residual associations among microbial

taxa to identify influences beyond shared external drivers. For 16S rRNA OTU model, at the host

species level, numerous associations were evident, with taxa showing positive co-occurrence within

groups and negative co-occurrence between groups, particularly for Actinomycetes and various

Proteobacteria (Fig. 7). At the host genotype level, Actinomycetes taxa formed two distinct clusters,

with contrasting associations with other clades, though abundance associations were less consistent. At

the individual host level, most taxa, except Alphaproteobacteria, Mollicutes, and some Actinomycetes,

showed positive associations, indicating that host individuals either had high or low microbial

diversity. Facultative endosymbionts showed limited interactions with other taxa. Site-level patterns of

associations were mostly positive, with some exceptions, and mixed trends were observed for

abundance conditional on presence, where more taxa showed negative associations.

Patterns observed for the 16S OTU models were similar to the ones shown for the 16S ZOTU

models (Supplementary Fig. S4). In contrast, for the *Wolbachia* models, few significant associations

were detected, likely due to low statistical power from the limited occurrence of specific ZOTUs

across host species and individuals (Supplementary Fig. S4).

**Fig. 7:** Residual associations among bacterial taxa explain how OTUs may influence each others' presence and abundance (conditional on presence) once shared external drivers are accounted for. The columns of panels correspond to the four random levels included in the HMSC models, and the rows of panels represent two model components. Within each panel, the bacterial taxa were ordered in the same way as in the presence-absence part of the model at the level of the insect host species. Red colour indicates pairs of taxa with a positive residual association and blue colour pairs of taxa with a negative residual association, with a posterior probability of 0.90 used as a threshold.

## 4. Discussion

Insect biodiversity largely consists of poorly known 'dark taxa', such as scuttle flies [15,30,59]. In these groups, addressing even basic questions, such as about their interactions, pose considerable difficulties [29,60]. The situation is changing with the advent of new conceptual approaches combined with progress in sequencing, bioinformatics, and statistical modelling [11,17]. Our study demonstrates how these innovations, especially when applied in combination, can greatly enhance our understanding of interactions within the diverse wild insect communities.

## 4.1. What have we learnt about scuttle flies?

This study shows how diverse communities of a dark taxon can be accurately resolved *at the same time* as their associations with microbes or other organisms. This approach opens up entirely new prospects for community ecology. To date, most studies of ecological communities have focused on characterising species' Grinnellian niches – that is, how communities are shaped by the match between abiotic conditions and species requirements [61–63]. This has largely neglected how species occurrence is determined by their interactions with other species – the Eltonian niche [64,65]. This study characterises both of these niches in one go. By simultaneously identifying the host species (here: scuttle flies) and their associated taxa (here: bacteria) across sites, we may effectively characterise entire wild communities of multi-partite symbiotic entities, or holobionts [66]. The current study focuses on a relatively small set of sites, thus constraining analyses of the Grinnellian niche – but another study shows how this can be achieved for broader sample collection [29]. By adding the microbial dimension, we show the way forward for the integration of the two [67].

## 4.2. What have we learnt about scuttle fly microbiota?

Our study resolves massive variation in the microbiota of scuttle flies at several levels of organization. We show a large variation in bacterial abundance – a rarely assessed, although critical aspect of microbiota [19,20,40]. We present up to four orders of magnitude differences in the number of 16S rRNA copies, even among individuals representing the same population and sex. This is likely to correlate with large differences in bacterial cell density, although variations in rRNA operon copy numbers within bacterial genomes [68,69] and copies of the genome per cell [70] can also play a role. In turn, bacterial density should correlate with the functional significance of bacteria in the host biology, even if the density-dependence of bacterial effects on the host [71,72] makes the direction of these effects hard to predict.

This variation in bacterial abundance is linked to the community composition. In particular,

facultative endosymbionts *Wolbachia* and *Rickettsia* tend to be highly abundant when present, driving

the overall abundance among species and sexes. These widespread microbes show highly specific

patterns in their distribution, with host species, genotype, and sex explaining a large share of the

overall variance across statistical models. Pronounced variation in the association of facultative

endosymbiont vs host genotypes suggests that these symbioses are dynamic at relatively short

timescales - with symbiont strains moving among host lines, spreading, and likely affecting the host

population structure, but rarely persisting for a long time [3,73,74]. At the same time, the broad

distribution of some *Wolbachia* COI genotypes across different phorid species indicates that they are

relatively frequently transmitted horizontally among distantly related host individuals [75].

*Wolbachia* is thought to spread in host populations through the combination of reproductive

manipulation and beneficial effects, including protection against natural enemies and nutrient

provisioning. Our community-level data strongly suggest that reproductive manipulation is common in

the phorid system. Specifically, major differences in *Wolbachia* prevalence and abundance between

males and females (Supplementary Table S6) argue for a negative link between infection and male

functions. These patterns are consistent with expectations for some types of reproductive manipulation:

male killing, feminisation of genetic males, or parthenogenesis induction [3,76]. Unfortunately, we

lack information about sex ratios in the surveyed populations since our sampling and specimen

selection strategy were not designed to assess sex ratios accurately. Nonetheless, in eight out of 25

most abundant species, the sex ratio among processed specimens significantly deviated from an

expected 1:1 sex ratio towards females, and in six others towards males (chi-squared test, $p < 0.05$).

The current tentative evidence for reproductive manipulation needs to be systematically validated

through the combination of genomics and experiments.

Conversely, many of the remaining abundant bacteria represent generalist clades. For example,

the genus *Pseudomonas* is known from a wide range of environments and host organisms that it

influences in a multitude of ways. While hundreds of *Pseudomonas* species have been formally named,

their environmental variation is much greater [77]. The same applies to most other abundant bacterial

genera in our dataset, including *Serratia* [78], *Carnobacterium* [79], *Providencia* [80], and others. To

date, research on these bacterial genera has focused on a limited set of species of direct relevance to

human health and economy, and our understanding of their taxonomic and functional diversity in

natural ecosystems has remained limited. At the same time, our 16S rRNA genotype data highlight the

surprisingly broad distribution of bacterial genotypes across phorid species, suggesting their recent and

likely ongoing transmission among distantly related hosts. Here, the relatively slow rate of evolution in

the rRNA gene [81] constrains further resolution of phorid-bacteria relationships. Higher-resolution

tools, especially genomics, will be necessary to conclude how these microbes are distributed and move

among hosts and environments and what their effects may be.

Overall, our data suggest that adult scuttle flies may not typically form specific associations

with microbes other than heritable facultative endosymbionts. We suspect that at the time of flies'

emergence from pupas, their body surfaces and digestive tracts may typically be largely sterile - and

that only later do they become colonised by some of the versatile opportunistic bacteria present in the

environment.

*4.3. What do scuttle flies teach us about broader insect–microbe interaction*

*patterns?*

Perhaps the most universal finding of this study, and the key to interpreting most patterns, is the

striking contrast between microbial functional categories - facultative endosymbionts and other

bacteria - in their associations with hosts [82]. In future studies, it will be critical to distinguish among

functional categories when assessing insect-microbe associations. At the same time, when

characterising the microbiota of a poorly known host group, distribution patterns can provide insights

into microbial biology and its role in the host's biology.

Facultative endosymbioses, represented here by *Wolbachia*, *Rickettsia*, and *Spiroplasma*, are

often thought to form relatively dynamic associations with hosts. We have convincing examples of

relatively recent symbiont host switches among related species [74,83] and across distantly related

species that form natural communities [84], as well as selective sweeps that they may drive

[5,7,85,86]. These processes can dramatically influence insect population structure, performance in

changing environmental conditions, or interactions with other organisms [11]. Conversely, this can

affect species' responses to ecological and environmental challenges and opportunities, their longer-

term evolutionary trajectories, as well as broader community processes [4,87]. However, we do not

know how often these processes take place or what their primary drivers and likely consequences are

[73,75]. We believe that the patterns observed in phorids - including a relatively short age of most

host-symbiont associations, frequent transmission of some *Wolbachia* lineages across host species, and

common reproductive manipulation - will eventually prove to be general features of facultative

endosymbiont infections across diverse insects. Pioneering methodological approaches such as those

used in our study will enable the systematic testing of these hypotheses about facultative

endosymbiosis stability and infection dynamics.

In contrast, microbial clades other than facultative endosymbionts are likely to have much

broader distributions. Most genera listed in Supplementary Table 4 have been reported repeatedly from

many diverse insects from around the world - for example, from butterflies or colonies of army ants

from Neotropics [52,53,88]. It is tempting to think of these microbes as highly mobile across global

environments and host clades. However, comparative data from vertebrates show how host biology

may shape the specificity of bacteria, including strains from broadly distributed genera [89]. The

systematic assessment of microbial diversity, transmission patterns, and roles across taxa and

environments would require not only massive sampling and stringent contamination control [18] but

also markers that provide much greater phylogenetic resolution than a short rRNA gene region -

ideally, entire genomes.

At the same time, scuttle flies do not represent the full range of known insect-microbe

associations. Many other insects form highly stable associations with specific bacterial clades, often

maintained through nutritional dependencies, whether with obligate endosymbionts [9] or with host-

adapted gut microbiota [82,90]. We would normally expect bacteria representing these categories to be

always present and abundant within the host species - and in surveys spanning multiple species, to

show evidence of co-diversification with hosts [91,92]. However, it is possible that the near-universal

associations with *Wolbachia* or *Rickettsia* that some scuttle fly species form may be due to the

indispensable effects of these microbes, as reported from some other insects [93,94].

## *4.4. Advancing the study of wild-insect microbiota: a comprehensive approach*

This study introduces a novel methodological framework for understanding wild insect

communities and their microbial associations, offering insights far beyond the capabilities of standard

marker gene amplicon sequencing or other methodologies. Our approach incorporates four key

innovations: 1: host DNA-based identification done alongside bacterial community characterisation, 2:

quantification of bacterial abundance, 3: the use of COI amplicons for high-resolution characterisation

of *Wolbachia*, and 4: the application of advanced modelling to decipher complex community patterns

of hosts and microbes. By combining these techniques at scale, we provide a uniquely nuanced and

biologically relevant understanding of microbiota diversity and its ecological role in wild insect

communities.

DNA barcoding has already revolutionized the understanding of arthropod biodiversity [95]. In

this study, while characterising microbiota, we simultaneously barcoded their hosts, providing a robust

framework for reconstructing insect microbiota patterns with virtually unprecedented resolution. As

shown in other insect systems, COI data may also enable correcting misidentifications and

reconstructing patterns such as parasitoid infections - potentially important confounding variables in

microbiota studies [96].

We incorporated synthetic spike-ins to estimate microbial abundance, a step that adds a crucial

dimension to our understanding of microbiota. Such quantification has been repeatedly recommended

due to the known variation in within-host bacterial abundance [17,19,20], but our approach shows how

it can be implemented in large-scale studies. While qPCR has been the most popular quantification

approach, spike-ins [40] are an extremely cost-effective alternative of obtaining consistent abundance

estimates spanning some six orders of magnitude. The patterns reconstructed are biologically realistic,

as revealed by the non-random bacterial distributions observed within scuttle fly species.

Quantification helps to assess samples where bacterial abundances are high and community

composition reliable, but also identify samples with low bacterial abundances, where we should treat

the community composition information with caution. This is preferable to discarding such samples for

being below the detection limit. While this approach awaits more systematic validation, we highly

recommend incorporating spike-in-based quantification into microbiota studies across small

organisms.

COI amplicon data for *Wolbachia* provide much higher phylogenetic resolution than standard

16S rRNA amplicon data and thus yield more accurate insights into relationships among insects and

these key facultative endosymbionts. This was realised over two decades ago when COI was

incorporated into the multi-locus sequence typing (MLST) scheme [97]. Interestingly, the realisation

that *Wolbachia* co-amplifies with insect mitochondrial COI [98] has motivated dedicated efforts to

block its amplification [99]. Here, we have considered this co-amplification as an opportunity rather

than a problem. The > 90% overlap between 16S rRNA-based and COI-based *Wolbachia* detection

(depending on the cutoffs used - K. H. Nowak, unpublished) suggests that this symbiont can be

reliably detected based on COI data alone. Increased phylogenetic resolution greatly enhances our

ability to reconstruct key distribution and transmission patterns. This suggests exciting prospects for

extracting infection information from rapidly growing barcoding datasets.

Finally, our implementation of the Hierarchical Modelling of Species Communities proved

indispensable for systematically reconstructing host and microbiota patterns and drivers. HMSC is

ideally suited to deal with the complexities of these multi-dimensional systems, but its applications in

the reconstruction of insect-bacteria interaction networks have so far been limited [100]. On the other

hand, in other host-microbe systems, the method has proven its power to reconstruct complex

processes and patterns and their drivers [101]. In particular, it offers a way for disentangling species

associations caused by a joint preference for the same environmental conditions from interactions

among species. As we address broader questions about biological interactions in complex multi-

species communities from environmentally diverse habitats undergoing rapid anthropogenic changes,

HMSC will serve as an essential statistical framework for understanding such complexity.

The solutions outlined above - combined with other recommended approaches such as

contamination filtering [17,19,102] - help address the key limitations of amplicon sequencing and put

this powerful, if somewhat out-of-fashion technique back into the mainstream of microbiota research.

The amplicon sequencing approach can be applied in tandem with complementary techniques that vary

in breadth and depth - high-throughput (meta-) barcoding-based insect surveys on one end and high-

resolution metagenomics and metatranscriptomics on the other [11,103]. Such a combination of tools

and approaches can revolutionise our understanding of the significance of microbiota in natural

ecosystems during the era of major anthropogenic changes.

## 4.5. Conclusions

In this paper, we have shown how a combination of quantification-enabled marker gene

amplicon sequencing and statistical modelling, when applied to large collections of diverse specimens,

can reveal ecological patterns of fundamental importance in a model dark taxon. By sequencing

amplicons of the host marker gene (COI) and reconstructing barcodes alongside microbiota

characterisation, we establish the critical host taxonomic framework even when species are impossible

to tell apart or undescribed. By incorporating synthetic spike-in-based quantification, we cost-

effectively assess microbiota abundance, a critical, if largely neglected, aspect of host-microbe interactions. By analysing the bacterial bycatch data from the COI amplicon sequencing, we improve the phylogenetic resolution for the dominant insect endosymbiont, *Wolbachia*. By implementing HMSC modelling, we reconstruct the primary drivers of host-microbe associations in a massive multi-dimensional dataset. More generally, we show how methodological and statistical innovations allow us to address questions about microbe associations across diverse natural communities at a combination of breadth and resolution that has, to our knowledge, not been achieved in a non-vertebrate system. Further refinement and broad implementation of these approaches will aid the research community's endeavours to understand the significance of microbial associations in the function and resilience of natural communities to rapidly changing conditions.

## Acknowledgements

The project was supported by the Polish National Agency for Academic Exchange grant PPN/PPO/2018/1/00015 and Polish National Science Centre grants 2018/31/B/NZ8/01158 and 2021/41/B/NZ8/04526 (to PŁ). EH's work on Swedish Phoridae was funded by the Swedish Taxonomy Initiative grant 2016-203 4.3. OO and TR were funded by the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation programme (grant agreement No 856506: ERC-Synergy project LIFEPLAN). OO was additionally funded by the Research Council of Finland (grant no. 336212 and 345110). The authors also acknowledge support from the National Genomics Infrastructure in Stockholm. We thank Dave Karlsson, Esben Bøggild, trap owners, and sample sorters for the Artdatabanken-funded Swedish Insect Inventory Project that provided the samples for this project.

## Supplementary item legends

Supplementary figures, tables and files are available at:

https://github.com/KarolHub/Phorid-Microbiota/tree/main/Supplementary_materials

Supplementary Fig. S1: Ordination of host communities' composition among sites and seasons (NMDS

plot based on Bray-Curtis dissimilarity).

Supplementary Fig. S2: Relationship between the explained variance ($R^2$) of bacterial taxa presence

and their prevalence.

Supplementary Fig. S3: Proportion of variance explained per microbial taxon by the presence-absence

model and the abundance conditional of presence model at the levels of microbial 16S-V4 OTUs (a),

16S-V4 ZOTUs (b) and *Wolbachia* COI ZOTUs (c).

Supplementary Fig. S4: Residual associations in the presence-absence model and the abundance

conditional of presence model among bacterial taxa at the levels of microbial 16S-V4 OTUs (a), 16S-

V4 ZOTUs (b) and *Wolbachia* COI ZOTUs (c).

Supplementary Table S1: Details of the sampling sites and the numbers of processed insect specimens

passing the quality criteria.

Supplementary Table S2: List of sample barcodes, species and genotype assignments, along with their

sexes, sampling locations and accession numbers.

Supplementary Table S3: Numbers and proportions of 16S rRNA reads classified as contaminants,

spike-ins, and symbionts across the samples, identified by the QUACK script.

Supplementary Table S4: List of 16S rRNA OTUs, their prevalence and abundance across samples.

Supplementary Table S5: Comparison of *Wolbachia* presence between COI and 16S-V4 datasets.

Supplementary Table S6: Results of the HMSC models

Supplementary File 1: Compressed table of COI ZOTUs read numbers and taxonomic assignments

across the samples.

Supplementary File 2: Compressed, decontaminated table of 16S-V4 ZOTU read numbers and

taxonomic assignments across the samples.

Supplementary File 3: Compressed, decontaminated table of estimated 16S-V4 ZOTU copies and

taxonomic assignments across the samples.

# References

1.    McFall-Ngai M *et al.* 2013 Animals in a bacterial world, a new imperative for the life sciences.

*Proc. Natl. Acad. Sci.* **110**, 3229–3236. (doi:10.1073/pnas.1218525110)

2.    Douglas AE. 2015 Multiorganismal Insects: Diversity and Function of Resident Microorganisms.

*Annu. Rev. Entomol.* **60**, 17–34. (doi:10.1146/annurev-ento-010814-020822)

3.    Kaur R, Shropshire JD, Cross KL, Leigh B, Mansueto AJ, Stewart V, Bordenstein SR,

Bordenstein SR. 2021 Living in the endosymbiotic world of Wolbachia: A centennial review.

*Cell Host Microbe* **29**, 879–893. (doi:10.1016/j.chom.2021.03.006)

4.    Oliver KM, Degnan PH, Burke GR, Moran NA. 2010 Facultative Symbionts in Aphids and the

Horizontal Transfer of Ecologically Important Traits. *Annu. Rev. Entomol.* **55**, 247–266.

(doi:10.1146/annurev-ento-112408-085305)

5.    Jaenike J. 2012 Population genetics of beneficial heritable symbionts. *Trends Ecol. Evol.* **27**,

226–232. (doi:10.1016/j.tree.2011.10.005)

6.    Jaenike J, Unckless R, Cockburn SN, Boelio LM, Perlman SJ. 2010 Adaptation via Symbiosis:

Recent Spread of a *Drosophila* Defensive Symbiont. *Science* **329**, 212–215.

(doi:10.1126/science.1188235)

7.    Kriesner P, Hoffmann AA, Lee SF, Turelli M, Weeks AR. 2013 Rapid Sequential Spread of Two

Wolbachia Variants in Drosophila simulans. *PLoS Pathog.* **9**, e1003607.

(doi:10.1371/journal.ppat.1003607)

8.   Cornwallis CK, Van 'T Padje A, Ellers J, Klein M, Jackson R, Kiers ET, West SA, Henry LM.

2023 Symbioses shape feeding niches and diversification across insects. *Nat. Ecol. Evol.* **7**,

1022–1044. (doi:10.1038/s41559-023-02058-0)

9.   Moran NA, McCutcheon JP, Nakabachi A. 2008 Genomics and Evolution of Heritable Bacterial

Symbionts. *Annu. Rev. Genet.* **42**, 165–190. (doi:10.1146/annurev.genet.41.110306.130119)

10.  Frago E, Dicke M, Godfray HCJ. 2012 Insect symbionts as hidden players in insect–plant

interactions. *Trends Ecol. Evol.* **27**, 705–711. (doi:10.1016/j.tree.2012.08.013)

11.  Łukasik P, Kolasa MR. 2024 With a little help from my friends: the roles of microbial symbionts

in insect populations and communities. *Philos. Trans. R. Soc. B Biol. Sci.* **379**, 20230122.

(doi:10.1098/rstb.2023.0122)

12.  Sudakaran S, Kost C, Kaltenpoth M. 2017 Symbiont Acquisition and Replacement as a Source of

Ecological Innovation. *Trends Microbiol.* **25**, 375–390. (doi:10.1016/j.tim.2017.02.014)

13.  Zheng H, Steele MI, Leonard SP, Motta EVS, Moran NA. 2018 Honey bees as models for gut

microbiota research. *Lab Anim.* **47**, 317–325. (doi:10.1038/s41684-018-0173-x)

14.  Shigenobu S, Yorimoto S. 2022 Aphid hologenomics: current status and future challenges. *Curr.*

*Opin. Insect Sci.* **50**, 100882. (doi:10.1016/j.cois.2022.100882)

15.  Li X, Wiens JJ. 2023 Estimating Global Biodiversity: The Role of Cryptic Insect Species. *Syst.*

*Biol.* **72**, 391–403. (doi:10.1093/sysbio/syac069)

16.  Stork NE. 2018 How Many Species of Insects and Other Terrestrial Arthropods Are There on

Earth? *Annu. Rev. Entomol.* **63**, 31–45. (doi:10.1146/annurev-ento-020117-043348)

17.   Knight R *et al.* 2018 Best practices for analysing microbiomes. *Nat. Rev. Microbiol.* **16**, 410–

422. (doi:10.1038/s41579-018-0029-9)

18.   Salter SJ *et al.* 2014 Reagent and laboratory contamination can critically impact sequence-based

microbiome analyses. *BMC Biol.* **12**, 87. (doi:10.1186/s12915-014-0087-z)

19.   Williamson E, Hammer T, Hogendoorn K, Eisenhofer R. 2024 Blanking on Blanks: Few Insect

Microbiota Studies Control for Contaminants - A Systematic Review.

(doi:10.1101/2024.08.30.610589)

20.   Hammer TJ, Sanders JG, Fierer N. 2019 Not all animals need a microbiome. *FEMS Microbiol.*

*Lett.* **366**, fnz117. (doi:10.1093/femsle/fnz117)

21.   Baumann P. 2005 BIOLOGY OF BACTERIOCYTE-ASSOCIATED ENDOSYMBIONTS OF

PLANT SAP-SUCKING INSECTS. *Annu. Rev. Microbiol.* **59**, 155–189.

(doi:10.1146/annurev.micro.59.030804.121041)

22.   Lange C *et al.* 2023 Impact of intraspecific variation in insect microbiomes on host phenotype

and evolution. *ISME J.* **17**, 1798–1807. (doi:10.1038/s41396-023-01500-2)

23.   Hurst GDD, Frost CL. 2015 Reproductive Parasitism: Maternally Inherited Symbionts in a

Biparental World. *Cold Spring Harb. Perspect. Biol.* **7**, a017699.

(doi:10.1101/cshperspect.a017699)

24.   Perlmutter JI, Bordenstein SR. 2020 Microorganisms in the reproductive tissues of arthropods.

*Nat. Rev. Microbiol.* **18**, 97–111. (doi:10.1038/s41579-019-0309-z)

25.   Hartop E, Srivathsan A, Ronquist F, Meier R. 2022 Towards Large-Scale Integrative Taxonomy

(LIT): Resolving the Data Conundrum for Dark Taxa. *Syst. Biol.* **71**, 1404–1422.

(doi:10.1093/sysbio/syac033)

26. Srivathsan A, Hartop E, Puniamoorthy J, Lee WT, Kutty SN, Kurina O, Meier R. 2019 Rapid,

large-scale species discovery in hyperdiverse taxa using 1D MinION sequencing. *BMC Biol.* **17**,

96. (doi:10.1186/s12915-019-0706-9)

27. Srivathsan A, Lee L, Katoh K, Hartop E, Kutty SN, Wong J, Yeo D, Meier R. 2021

ONTbarcoder and MinION barcodes aid biodiversity discovery and identification by everyone,

for everyone. *BMC Biol.* **19**, 217. (doi:10.1186/s12915-021-01141-x)

28. Chua PYS, Bourlat SJ, Ferguson C, Korlevic P, Zhao L, Ekrem T, Meier R, Lawniczak MKN.

2023 Future of DNA-based insect monitoring. *Trends Genet.* **39**, 531–544.

(doi:10.1016/j.tig.2023.02.012)

29. Hartop E, Lee L, Srivathsan A, Jones M, Peña-Aguilera P, Ovaskainen O, Roslin T, Meier R.

2024 Resolving biology's dark matter: species richness, spatiotemporal distribution, and

community composition of a dark taxon. *BMC Biol.* **22**, 215. (doi:10.1186/s12915-024-02010-z)

30. Srivathsan A *et al.* 2023 Convergence of dominance and neglect in flying insect diversity. *Nat.*

*Ecol. Evol.* **7**, 1012–1021. (doi:10.1038/s41559-023-02066-0)

31. Disney H. 1994 *Scuttle Flies*. Dordrecht: Springer Netherlands.

32. Hartop EA, Brown BV, Disney RHL. 2015 Opportunity in our Ignorance: Urban Biodiversity

Study Reveals 30 New Species and One New Nearctic Record for Megaselia (Diptera: Phoridae)

in Los Angeles (California, USA). *Zootaxa* **3941**. (doi:10.11646/zootaxa.3941.4.1)

33. Disney RH, Taylor S, Slay ME, Krejca JK. 2011 New species of scuttle flies (Diptera: Phoridae)

recorded from caves in Nevada, USA. *Subterr. Biol.* **9**, 73–84. (doi:10.3897/subtbiol.9.2511)

34.    Cheng L, Hogue CL. 1974 New distribution and habitat records of biting midges and mangrove

797           flies from the coasts of southern Baja California, Mexico (Diptera: Ceratopogonidae, Culicidae,

798           Chironomidae, and Phoridae. *Entomol. News* **85**, 211–218.

799    35.    Disney RHL. 1991 The aquatic Phoridae (Diptera). *Insect Syst. Evol.* **22**, 171–191.

800           (doi:10.1163/187631291X00048)

801    36.    Pennington MJ, Rothman JA, Jones MB, McFrederick QS, Gan J, Trumble JT. 2017 Effects of

802           contaminants of emerging concern on Megaselia scalaris (Lowe, Diptera: Phoridae) and its

803           microbial community. *Sci. Rep.* **7**, 8165. (doi:10.1038/s41598-017-08683-7)

804    37.    Sakamoto JM, Shikano I, Rasgon JL. 2024 Microbiomes of Two Pest Fly Species of

805           Pennsylvania Mushroom Houses. *Insects* **15**, 525. (doi:10.3390/insects15070525)

806    38.    Karlsson D, Hartop E, Forshage M, Jaschhof M, Ronquist F. 2020 The Swedish Malaise Trap

807           Project: A 15 Year Retrospective on a Countrywide Insect Inventory. *Biodivers. Data J.* **8**,

808           e47255. (doi:10.3897/BDJ.8.e47255)

809    39.    Buczek M *et al.* 2024 A new tool in a toolbox: Addressing challenges in high-throughput

810           microbiota surveys across diverse wild insects. (doi:10.1101/2024.08.26.609764)

811    40.    Tourlousse DM, Yoshiike S, Ohashi A, Matsukura S, Noda N, Sekiguchi Y. 2016 Synthetic

812           spike-in standards for high-throughput 16S rRNA gene amplicon sequencing. *Nucleic Acids Res.*

813           , gkw984. (doi:10.1093/nar/gkw984)

814    41.    Rohland N, Reich D. 2012 Cost-effective, high-throughput DNA sequencing libraries for

815           multiplexed target capture. *Genome Res.* **22**, 939–946. (doi:10.1101/gr.128124.111)

816    42.    Glenn TC *et al.* 2019 Adapterama II: universal amplicon sequencing on Illumina platforms

817           (TaggiMatrix). *PeerJ* **7**, e7786. (doi:10.7717/peerj.7786)

43.   Parada AE, Needham DM, Fuhrman JA. 2016 Every base matters: assessing small subunit rRNA

primers for marine microbiomes with mock communities, time series and global field samples.

*Environ. Microbiol.* **18**, 1403–1414. (doi:10.1111/1462-2920.13023)

44.   Elbrecht V, Braukmann TWA, Ivanova NV, Prosser SWJ, Hajibabaei M, Wright M, Zakharov

EV, Hebert PDN, Steinke D. 2019 Validation of COI metabarcoding primers for terrestrial

arthropods. *PeerJ* **7**, e7745. (doi:10.7717/peerj.7745)

45.   Zhang J, Kobert K, Flouri T, Stamatakis A. 2014 PEAR: a fast and accurate Illumina Paired-End

reAd mergeR. *Bioinformatics* **30**, 614–620. (doi:10.1093/bioinformatics/btt593)

46.   Rognes T, Flouri T, Nichols B, Quince C, Mahé F. 2016 VSEARCH: a versatile open source tool

for metagenomics. *PeerJ* **4**, e2584. (doi:10.7717/peerj.2584)

47.   Edgar RC. 2010 Search and clustering orders of magnitude faster than BLAST. *Bioinformatics*

**26**, 2460–2461. (doi:10.1093/bioinformatics/btq461)

48.   Antich A, Palacin C, Wangensteen OS, Turon X. 2021 To denoise or to cluster, that is not the

question: optimizing pipelines for COI metabarcoding and metaphylogeography. *BMC*

*Bioinformatics* **22**, 177. (doi:10.1186/s12859-021-04115-6)

49.   Edgar RC. 2016 SINTAX: a simple non-Bayesian taxonomy classifier for 16S and ITS

sequences. (doi:10.1101/074161)

50.   Leray M, Ho S-L, Lin I-J, Machida RJ. 2018 MIDORI server: a webserver for taxonomic

assignment of unknown metazoan mitochondrial-encoded sequences using a curated database.

*Bioinformatics* **34**, 3753–3754. (doi:10.1093/bioinformatics/bty454)

51.   Quast C, Pruesse E, Yilmaz P, Gerken J, Schweer T, Yarza P, Peplies J, Glöckner FO. 2012 The

SILVA ribosomal RNA gene database project: improved data processing and web-based tools.

*Nucleic Acids Res.* **41**, D590–D596. (doi:10.1093/nar/gks1219)

52.    Łukasik P, Newton JA, Sanders JG, Hu Y, Moreau CS, Kronauer DJC, O'Donnell S, Koga R,

842           Russell JA. 2017 The structured diversity of specialized gut symbionts of the New World army

843           ants. *Mol. Ecol.* **26**, 3808–3825. (doi:10.1111/mec.14140)

844    53.    Valdivia C, Newton JA, Von Beeren C, O'Donnell S, Kronauer DJC, Russell JA, Łukasik P.

845           2023 Microbial symbionts are shared between ants and their associated beetles. *Environ.*

846           *Microbiol.* **25**, 3466–3483. (doi:10.1111/1462-2920.16544)

847    54.    Ovaskainen O, Tikhonov G, Norberg A, Guillaume Blanchet F, Duan L, Dunson D, Roslin T,

848           Abrego N. 2017 How to make more out of community data? A conceptual framework and its

849           implementation as models and software. *Ecol. Lett.* **20**, 561–576. (doi:10.1111/ele.12757)

850    55.    Ovaskainen O, Abrego N. 2020 *Joint Species Distribution Modelling: With Applications in R.* 1st

851           edn. Cambridge University Press. (doi:10.1017/9781108591720)

852    56.    Warton DI, Blanchet FG, O'Hara RB, Ovaskainen O, Taskinen S, Walker SC, Hui FKC. 2015 So

853           Many Variables: Joint Modeling in Community Ecology. *Trends Ecol. Evol.* **30**, 766–779.

854           (doi:10.1016/j.tree.2015.09.007)

855    57.    Tikhonov G, Opedal ØH, Abrego N, Lehikoinen A, De Jonge MMJ, Oksanen J, Ovaskainen O.

856           2020 Joint species distribution modelling with the r □package H msc. *Methods Ecol. Evol.* **11**,

857           442–447. (doi:10.1111/2041-210X.13345)

858    58.    Leigh JW, Bryant D. 2015 popart□: full□feature software for haplotype network construction.

859           *Methods Ecol. Evol.* **6**, 1110–1116. (doi:10.1111/2041-210X.12410)

860    59.    Caruso V, Hartop E, Chimeno C, Noori S, Srivathsan A, Haas M, Lee L, Meier R, Whitmore D.

861           2024 An integrative framework for dark taxa biodiversity assessment at scale: A case study using

*Megaselia* (Diptera, Phoridae). *Insect Conserv. Divers.* **17**, 968–987. (doi:10.1111/icad.12762)

60. Roslin T. 2024 Understanding insect movements through space and time is vital for safeguarding
global ecosystem services. *One Earth* **7**, 537–540. (doi:10.1016/j.oneear.2024.03.007)

61. Grinnell J. 1917 Field Tests of Theories Concerning Distributional Control. *Am. Nat.* **51**, 115–
128. (doi:10.1086/279591)

62. Holt RD. 2009 Bringing the Hutchinsonian niche into the 21st century: Ecological and
evolutionary perspectives. *Proc. Natl. Acad. Sci.* **106**, 19659–19665.
(doi:10.1073/pnas.0905137106)

63. Soberón J. 2019 A Grinnellian Niche Perspective on Species-Area Relationships. *Am. Nat.* **194**,
760–775. (doi:10.1086/705898)

64. Colwell RK, Rangel TF. 2009 Hutchinson's duality: The once and future niche. *Proc. Natl.*
*Acad. Sci.* **106**, 19651–19658. (doi:10.1073/pnas.0901650106)

65. Hirzel AH, Le Lay G. 2008 Habitat suitability modelling and niche theory. *J. Appl. Ecol.* **45**,
1372–1381. (doi:10.1111/j.1365-2664.2008.01524.x)

66. Bordenstein SR, Theis KR. 2015 Host Biology in Light of the Microbiome: Ten Principles of
Holobionts and Hologenomes. *PLOS Biol.* **13**, e1002226. (doi:10.1371/journal.pbio.1002226)

67. Gravel D *et al.* 2019 Bringing Elton and Grinnell together: a quantitative framework to represent
the biogeography of ecological interaction networks. *Ecography* **42**, 401–415.
(doi:10.1111/ecog.04006)

68. Klappenbach JA. 2001 rrndb: the Ribosomal RNA Operon Copy Number Database. *Nucleic*
*Acids Res.* **29**, 181–184. (doi:10.1093/nar/29.1.181)

69.  Louca S, Doebeli M, Parfrey LW. 2018 Correcting for 16S rRNA gene copy numbers in

microbiome surveys remains an unsolved problem. *Microbiome* **6**, 41. (doi:10.1186/s40168-018-

0420-9)

70.  Pecoraro V, Zerulla K, Lange C, Soppa J. 2011 Quantification of Ploidy in Proteobacteria

Revealed the Existence of Monoploid, (Mero-)Oligoploid and Polyploid Species. *PLoS ONE* **6**,

e16392. (doi:10.1371/journal.pone.0016392)

71.  Koga R *et al.* 2022 Single mutation makes Escherichia coli an insect mutualist. *Nat. Microbiol.*

**7**, 1141–1150. (doi:10.1038/s41564-022-01179-9)

72.  Medina Munoz M, Spencer N, Enomoto S, Dale C, Rio RVM. 2020 Quorum sensing sets the

stage for the establishment and vertical transmission of Sodalis praecaptivus in tsetse flies. *PLOS*

*Genet.* **16**, e1008992. (doi:10.1371/journal.pgen.1008992)

73.  Bailly-Bechet M, Martins-Simões P, Szöllősi GJ, Mialdea G, Sagot M-F, Charlat S. 2017 How

Long Does Wolbachia Remain on Board? *Mol. Biol. Evol.* **34**, 1183–1193.

(doi:10.1093/molbev/msx073)

74.  Turelli M *et al.* 2018 Rapid Global Spread of wRi-like Wolbachia across Multiple Drosophila.

*Curr. Biol.* **28**, 963-971.e8. (doi:10.1016/j.cub.2018.02.015)

75.  Sanaei E, Charlat S, Engelstädter J. 2021 *Wolbachia* host shifts: routes, mechanisms, constraints

and evolutionary consequences. *Biol. Rev.* **96**, 433–453. (doi:10.1111/brv.12663)

76.  Engelstädter J, Hurst GDD. 2009 The Ecology and Evolution of Microbes that Manipulate Host

Reproduction. *Annu. Rev. Ecol. Evol. Syst.* **40**, 127–149.

(doi:10.1146/annurev.ecolsys.110308.120206)

77.  Lalucat J, Gomila M, Mulet M, Zaruma A, García-Valdés E. 2022 Past, present and future of the

boundaries of the Pseudomonas genus: Proposal of Stutzerimonas gen. Nov. *Syst. Appl.*

*Microbiol.* **45**, 126289. (doi:10.1016/j.syapm.2021.126289)

78.  Williams DJ *et al.* 2022 The genus Serratia revisited by genomics. *Nat. Commun.* **13**, 5195.

(doi:10.1038/s41467-022-32929-2)

79.  Leisner JJ, Laursen BG, Prévost H, Drider D, Dalgaard P. 2007 *Carnobacterium:* positive and

negative effects in the environment and in foods. *FEMS Microbiol. Rev.* **31**, 592–613.

(doi:10.1111/j.1574-6976.2007.00080.x)

80.  Dong X, Jia H, Yu Y, Xiang Y, Zhang Y. 2024 Genomic revisitation and reclassification of the

genus *Providencia*. *mSphere* **9**, e00731-23. (doi:10.1128/msphere.00731-23)

81.  Ochman H, Elwyn S, Moran NA. 1999 Calibrating bacterial evolution. *Proc. Natl. Acad. Sci.* **96**,

12638–12643. (doi:10.1073/pnas.96.22.12638)

82.  Perreau J, Patel DJ, Anderson H, Maeda GP, Elston KM, Barrick JE, Moran NA. 2021 Vertical

Transmission at the Pathogen-Symbiont Interface: Serratia symbiotica and Aphids. *mBio* **12**,

e00359-21. (doi:10.1128/mBio.00359-21)

83.  Shropshire D, Conner WR, Vanderpool D, Hoffmann AA, Turelli M, Cooper BS. 2023 Rapid

host switching of Wolbachia and even more rapid turnover of their phages and incompatibility-

causing loci. (doi:10.1101/2023.12.04.569981)

84.  Vancaester E, Blaxter M. 2023 Phylogenomic analysis of Wolbachia genomes from the Darwin

Tree of Life biodiversity genomics project. *PLOS Biol.* **21**, e3001972.

(doi:10.1371/journal.pbio.3001972)

85.  Himler AG *et al.* 2011 Rapid Spread of a Bacterial Symbiont in an Invasive Whitefly Is Driven

by Fitness Benefits and Female Bias. *Science* **332**, 254–256. (doi:10.1126/science.1199410)

86.  Zhao D, Zhang Z, Niu H, Guo H. 2023 Pathogens are an important driving force for the rapid

spread of symbionts in an insect host. *Nat. Ecol. Evol.* **7**, 1667–1681. (doi:10.1038/s41559-023-

02160-3)

87.  Ferrari J, Vavre F. 2011 Bacterial symbionts in insects or the story of communities affecting

communities. *Philos. Trans. R. Soc. B Biol. Sci.* **366**, 1389–1400. (doi:10.1098/rstb.2010.0226)

88.  Ravenscraft A, Berry M, Hammer T, Peay K, Boggs C. 2019 Structure and function of the

bacterial and fungal gut microbiota of Neotropical butterflies. *Ecol. Monogr.* **89**, e01346.

(doi:10.1002/ecm.1346)

89.  Song SJ *et al.* 2020 Comparative Analyses of Vertebrate Gut Microbiomes Reveal Convergence

between Birds and Bats. *mBio* **11**, e02901-19. (doi:10.1128/mBio.02901-19)

90.  Engel P, Moran NA. 2013 The gut microbiota of insects – diversity in structure and function.

*FEMS Microbiol. Rev.* **37**, 699–735. (doi:10.1111/1574-6976.12025)

91.  Fisher RM, Henry LM, Cornwallis CK, Kiers ET, West SA. 2017 The evolution of host-

symbiont dependence. *Nat. Commun.* **8**, 15973. (doi:10.1038/ncomms15973)

92.  Perreau J, Moran NA. 2022 Genetic innovations in animal–microbe symbioses. *Nat. Rev. Genet.*

**23**, 23–39. (doi:10.1038/s41576-021-00395-z)

93.  Dedeine F, Boulétreau M, Vavre F. 2005 Wolbachia requirement for oogenesis: occurrence

within the genus Asobara (Hymenoptera, Braconidae) and evidence for intraspecific variation in

A. tabida. *Heredity* **95**, 394–400. (doi:10.1038/sj.hdy.6800739)

94.  Hosokawa T, Koga R, Kikuchi Y, Meng X-Y, Fukatsu T. 2010 *Wolbachia* as a bacteriocyte-

associated nutritional mutualist. *Proc. Natl. Acad. Sci.* **107**, 769–774.

(doi:10.1073/pnas.0911476107)

95.  Meier R, Lawniczak MKN, Srivathsan A. 2024 Illuminating Entomological Dark Matter with

DNA Barcodes in an Era of Insect Decline, Deep Learning, and Genomics. *Annu. Rev. Entomol.*

(doi:10.1146/annurev-ento-040124-014001)

96.  Kolasa M, Kajtoch Ł, Michalik A, Maryańska☐Nadachowska A, Łukasik P. 2023 Till evolution

do us part: The diversity of symbiotic associations across populations of *Philaenus* spittlebugs.

*Environ. Microbiol.* **25**, 2431–2446. (doi:10.1111/1462-2920.16473)

97.  Baldo L *et al.* 2006 Multilocus Sequence Typing System for the Endosymbiont *Wolbachia*

*pipientis*. *Appl. Environ. Microbiol.* **72**, 7098–7110. (doi:10.1128/AEM.00731-06)

98.  Smith MA *et al.* 2012 Wolbachia and DNA Barcoding Insects: Patterns, Potential, and Problems.

*PLoS ONE* **7**, e36514. (doi:10.1371/journal.pone.0036514)

99.  Bleidorn C, Henze K. 2021 A new primer pair for barcoding of bees (Hymenoptera: Anthophila)

without amplifying the orthologous coxA gene of Wolbachia bacteria. *BMC Res. Notes* **14**, 427.

(doi:10.1186/s13104-021-05845-9)

100. Minard G, Tikhonov G, Ovaskainen O, Saastamoinen M. 2019 The microbiome of the *Melitaea*

*cinxia* butterfly shows marked variation but is only little explained by the traits of the butterfly or

its host plant. *Environ. Microbiol.* **21**, 4253–4269. (doi:10.1111/1462-2920.14786)

101. Parisy B *et al.* 2024 Arctic plant-fungus interaction networks show major rewiring with

environmental variation. *Commun. Earth Environ.* **5**, 735. (doi:10.1038/s43247-024-01902-w)

102. Davis NM, Proctor DM, Holmes SP, Relman DA, Callahan BJ. 2018 Simple statistical

identification and removal of contaminant sequences in marker-gene and metagenomics data.

*Microbiome* **6**, 226. (doi:10.1186/s40168-018-0605-2)

103. Andriienko V, Buczek M, Meier R, Srivathsan A, Łukasik P, Kolasa MR. 2024 Implementing

high-throughput insect barcoding in microbiome studies: impact of non-destructive DNA

extraction on microbiome reconstruction. *PeerJ* **12**, e18025. (doi:10.7717/peerj.18025)

16S rRNA OTUs

Mean R² = 20%

16S rRNA ZOTUs

Mean R² = 18%

COI *Wolbachia* ZOTUs

Mean R² = 34%

**Residual associations of bacterial OTUs**

host species effect · host genotype effect · host individual effect · site effect

positive association · negative association

presence-absence model

Alphaproteobacteria
Betaproteobacteria
Gammaproteobacteria
Sphingobacteriia
Flavobacteriia
Mollicutes
Bacilli
Actinomycetes
unassigned

abundance model

Alphaproteobacteria
Betaproteobacteria
Gammaproteobacteria
Sphingobacteriia
Flavobacteriia
Mollicutes
Bacilli
Actinomycetes
unassigned

**(b)** Residual associations of bacterial ZOTUs

host species effect · host genotype effect · host individual effect · site effect

presence-absence model

Alphaproteobacteria
Betaproteobacteria
Gammaproteobacteria
Sphingobacteriia
Flavobacteriia
Mollicutes
Bacilli
Actinomycetes
unassigned

abundance model

Alphaproteobacteria
Betaproteobacteria
Gammaproteobacteria
Sphingobacteriia
Flavobacteriia
Mollicutes
Bacilli
Actinomycetes
unassigned

**(c)** Residual associations of *Wolbachia* COI ZOTUs

host species effect · host genotype effect · host individual effect · site effect

presence-absence model

OTU105
OTU143
OTU148
OTU152
OTU154
OTU175
OTU180
OTU189
OTU308

abundance model

OTU105
OTU143
OTU148
OTU152
OTU154
OTU175
OTU180
OTU189
OTU308