Foraging niche partitioning of three *Myotis* bat species and marine fish

consumption by *Myotis pilosus* in a subtropical East Asian region

Authors:

Xiaodong Wei[1], Emily Shui Kei Poon[1], John Chun Ting Chung[1], David Tsz Chung

Chan[1], Chung Tong Shek[2], Wing Chi Tsui[2], Huabin Zhao[3], Simon Yung Wa Sin[1],*

Contact information:

[1] School of Biological Sciences, The University of Hong Kong, Pok Fu Lam Road,

Hong Kong, China.

[2] Agriculture, Fisheries and Conservation Department, Hong Kong SAR Government,

Hong Kong, China.

[3] Department of Ecology, College of Life Sciences, Wuhan University, Wuhan

430072, Hubei, China

* Correspondence:

Simon Yung Wa Sin

Email: yungwa.sin@gmail.com

Telephone number: (852)39173825

ORCID id: 0000-0003-2484-2897

Keywords:

Evolution of piscivory; Fecal DNA metabarcoding; Fish catching behavior;

Interspecific competition; Piscivorous bat diet; Rickett's big-footed myotis.

Abstract

Most bats are insectivorous, but some species have evolved the ability to prey on fish.

Although piscivory has been confirmed in the Rickett's big-footed myotis (*Myotis*

pilosus), the extent of piscivory of other cohabiting *Myotis* species is uncertain. This

study aims to explore the dietary niches and fish consumption of three *Myotis*

species in a subtropical East Asian region, and specifically the fish diet of *M.*

pilosus. Our findings reveal, for the first time, that *M. pilosus* consumes marine

fishes, in contrast to previous research conducted in inland regions that suggested

year-round consumption of cyprinids in freshwater habitats. We also observed

seasonal variation in the diets of *M. pilosus*. It predominately hunted wide-banded

hardyhead silverside, sailfin flying fish, and shorthead anchovy during the wet

season, while mainly preying upon mullets during the dry months. In more inland

areas, *M. pilosus* was found to primarily feed on invasive freshwater poeciliids.

Furthermore, *M. pilosus* consumed more fish during the dry season, while there was

a greater consumption of insects during the wet months. Most notably among our

findings is the consumption of fish by two individuals of Horsfield's myotis (*M.*

horsfieldii), indicating that the species is potentially piscivorous. We revealed that

both *M. horsfieldii* and *M. pilosus* consumed water striders, suggesting that foraging

of aquatic insects could be driving the evolution of fishing behavior. Our findings

have also shed light on the flexibility of foraging behavior in piscivorous bats.

Keywords:

Evolution of piscivory; Fecal DNA metabarcoding; Fish catching behavior;

Interspecific competition; Piscivorous bat diet; Rickett's big-footed myotis.

1. Introduction

Bats are the only mammals capable of sustained flight. They use echolocation during

flight to navigate and forage for food nocturnally. There are over 1400 known bat

species worldwide, forming a diverse group that have adapted to a wide range of diets

(Taylor, 2019). The majority of bats are insectivorous, feeding primarily on insects,

while others are frugivorous and nectarivorous bats, which consume fruit and nectar,

respectively (Ramírez-Fráncel et al., 2022). In addition to insects, certain bat species

have adapted to feed on other animals; for instance, sanguivorous vampire bats

consume blood (Riskin and Carter, 2023), and piscivorous bats prey on fish

(Aizpurua and Alberdi, 2018). Piscivorous bats can be found among two distinct

genera: *Noctilio* (bulldog bats) and *Myotis* (mouse-eared bats). Among these, only

the *N. leporinus* (greater bulldog bat) in Latin America has been confirmed as

mainly piscivorous (Brooke, 1994). Three other species display 'limited' fishing

behaviors, including *M. vivesi* (fish-eating myotis) in the Gulf of California (Otálora-

Ardila et al., 2013), *M. pilosus* (Rickett's big-footed myotis) in southern and eastern

China, Vietnam, and Laos (Ma et al., 2003), and *M. capaccinii* (long-fingered myotis)

in the Mediterranean Sea (Aihartza et al., 2003). Additionally, several species have

'unconfirmed' fishing behaviors, such as *N. albiventris* (lesser bulldog bat), *M.*

daubentonii (Daubenton's myotis), and *M. macropus* (large-footed myotis).

The evolution of fishing behavior in bats has long fascinated scientists, and

comprehending the dietary compositions of various piscivorous bats is crucial for

understanding their foraging ecology and unravelling the intricacies of these

evolutionary processes. While fishing behavior is an important hunting strategy, it is

worth noting that none of these species are obligate piscivorous (Aizpurua et al.,

2014). Fish-eating bats also prey upon arthropods to varying extents, and the

prevalence and intensity of fishing or hunting arthropods can be influenced by

different factors such as their morphological traits, prey availability, and interspecific

competition within their habitats (Chang et al., 2019).

In Asia, *M. pilosus* is relatively large for its genus, exhibiting a body length

ranging from 51 to 65 mm (Wilson and Mittermeier 2019). They possess remarkably

enormous feet with enlarged, laterally compressed claws, resembling those of *N.*

leporinus (Fish et al., 1991, Ospina-Garcés et al., 2016). However, it was not until

2003 that the piscivorous nature of *M. pilosus* was confirmed (Ma et al., 2003). The

bats capture insects in mid-air and use their hind limbs to employ a trawling method

for catching fish (Jiang et al., 2003). While fish remains have been discovered in their

diets in some regions, regular fish consumption has only been verified in Beijing in

northern China (Ma et al., 2006). Studies conducted in Beijing, Shandong, and

Guizhou have shed light on the dietary habits of *M. pilosus* (Ma et al., 2006, Chang et

al., 2019, Ma et al., 2003, Wang et al., 2024). These studies reported on the

piscivorous and insectivorous feeding behaviors of *M. pilosus*, with species from the

order Cypriniformes (cyprinids) being their primary fish prey. In Beijing, they found

that *M. pilosus* consumed grass carp (*Ctenopharyngodon idella*), Eurasian carp

(*Cyprinus carpio*), pale chub (*Zacco* spp.), and Amur minnow (*Rhynchocypris* spp.)

(Ma et al., 2006, Chang et al., 2019). In Shandong, in addition to grass carp and

Eurasian carp, they also found bitterlings (*Rhodeus* spp.) to be their primary prey

(Chang et al., 2019). These fish species are similarly found in freshwater habitats

within inland areas (Froese and Pauly, 2024). Additionally, *M. pilosus* was found to

consume at least seven orders of insects, such as Coleoptera (beetles), Lepidoptera

(moths and butterflies), Hemiptera, Diptera, and more (Ma et al., 2006, Chang et al.,

2019).

However, discrepancies in the extent of fishing incidence have been observed

among these studies. One possible factor contributing to these discrepancies could be

the variations in spatiotemporal prey availability across different sampling locations

and seasons. Further research on their fish and insect consumption is crucial to gain a

better understanding of how resource variation or resource partitioning is associated

with its adaptations to the environment through foraging strategy. Thus, it is necessary

to investigate the dietary compositions of *M. pilosus* in various locations or habitats

across seasons to provide further insights into their fishing behavior. However,

beyond the few populations studied, the dietary composition of *M. pilosus* remains

largely unknown.

There have been speculations that several other Asiatic *Myotis* species, such as

M. macrotarsus (pallid large-footed myotis), *M. horsfieldii* (Horsfield's myotis), *M.*

hasseltii (lesser large-footed myotis), *M. stalkeri* (Kei myotis), *M.* adversus (large-

footed bat), *M. macrodactylus* (big-footed myotis), and *M. macropus*, might be

piscivorous due to their large hind feet and claws (Aizpurua and Alberdi, 2018).

Notably, *M. horsfieldii* is a native bat species in South Asia. Although it is smaller

compared to *M. pilosus*, with a body length ranging from 44 to 51 mm, its capability

to directly capture insects from water surfaces (Wilson and Mittermeier 2019)

suggests a potential for piscivory. It is widely believed that the fishing behavior in

bats evolved from their insectivorous habits, with the hypothesis that capturing

insects from water represents an intermediate stage in this behavior (Aizpurua et al.,

2013). Despite the relatively large hind feet of *M. horsfieldii*, exceeding half the

length of the tibia, and their observed circular flight patterns above open water

surfaces at a close distance of a few centimeters to search for insects, there is

currently a lack of evidence for fishing behavior in *M. horsfieldii* (Aizpurua and

Alberdi, 2018). Furthermore, their dietary composition has never been reported to

date, leaving unanswered questions regarding whether fish forms part of its diet and

what insect species they prey upon.

Hong Kong, situated at the southern coast of China, is the distribution range of

Myotis species (Shek, 2006). It is a semi-island known for its diverse habitats,

including various types of forests, coastal areas, wetlands, marine, and freshwater

habitats, each hosting a wide array of arthropod and fish species (Dudgeon and

Corlett, 2004). In this region, four *Myotis* species coexist, including *M. pilosus*, *M.*

horsfieldii, *M. chinensis* (Chinese myotis), and *M. muricola* (whiskered myotis,

locally rare). *Myotis chinensis* is bigger in size than *M. pilosus*, with a body length

ranging from 91 to 97 mm (Wilson and Mittermeier 2019), and possesses relatively

large hind feet compared to some other bat species. However, there are only a few

studies conducted in China reporting its diets. Despite its large size, it is believed that

M. chinensis exclusively feeds on insects. They are capable of preying on larger

insects and capturing ground-dwelling insects through picking behavior, primarily

consuming species in Coleoptera (e.g., carabid beetles), Orthoptera (grasshoppers),

and Diptera (flies) according to an earlier study (Ma et al., 2008). Given the

uncertainties surrounding the extent of piscivory by different *Myotis* species in the

region, this study seeks to investigate the diets of *M. pilosus*, *M. horsfieldii*, and *M.*

chinensis in Hong Kong by analyzing their fecal compositions using DNA

metabarcoding.

Traditional microscopy for species identification of fecal or stomach contents

presents several challenges. The identification of partially digested prey body parts

under a microscope relies heavily on the expertise and experience of the analyst.

Moreover, the process can be a time-consuming, subjective, and may have limitations

in taxonomic resolution, potentially introducing biases or inconsistencies in prey

identification (Svendsen et al., 2023, Massey et al., 2021). In contrast, DNA

metabarcoding overcomes some of these challenges. It is a high-throughput technique

that enables the analysis of a large number of samples and provides a higher

taxonomic resolution even with small amounts of prey DNA (de Sousa et al., 2019,

Monterroso et al., 2019).

By analyzing the prey identities and relative quantities of the three *Myotis*

species, we can gain insights into the spectra of habitats bat individuals forage and

prey they target in a habitat-diverse environment. Notably, both *M. chinensis* and *M.*

pilosus are listed as Near Threatened species in the Red List of China's Vertebrates

(Jiang et al., 2016), with *M. pilosus* also being classified as a Vulnerable species in the

IUCN Red List of Threatened Species (IUCN, 2023, Jiang et al., 2019). Hence, it

would be important to further explore the dietary niche partitioning of these sympatric

species. The findings would inform us their trophic relationships and what types of

food resources and habitats are critical to their population sustainability, and

significantly contribute to the conservation efforts for each of these species in the

region.

In this study, we aim 1**)** to investigate the dietary compositions of the three

Myotis species at both individual and species levels, 2) to estimate the dietary

diversity within individuals and populations, 3) to assess the effect of environmental

and host factors on the dietary compositions of these species, and 4) to determine the

patterns of dietary niche partitioning among the *Myotis* species. We hypothesize that

the three species might demonstrate dietary niche partitioning. We predict distinct

dietary compositions for each species, resulting in minimal dietary overlap between

species. Specifically, we predict that *M. pilosus* might include diverse fish taxa in its

diet due to the diverse aquatic habitats close to the roosting sites. We also hypothesize

that there is seasonal variation in the dietary compositions of the bat species. We

predict a shift in food consumption pattern across different seasons.

2. Materials and Methods

2.1 Sample collection

Between 2018 and 2021, we visited water tunnels or caves in Hong Kong to capture

Myotis bats during wet (April to September) and dry (October to March) season. We

refrained from visiting roosting sites during the months when bats were overwintering

or breeding/nursing, thus minimizing any potential disturbance. The locations include

Lin Ma Hang (LMH01), Tai Lam Chung (TLC01 and TLC02), Pak Tam Chung

(PTC), and Sai Kung (SK0, SK02, and SK03) (Fig. S1 and Table S1a-c) (Shek, 2004).

We captured the bats using hand-held hoop nets and placed them in breathable

sterilized bags. To prevent cross-contamination, each individual bat was kept in a

separate sterilized bag. We identified the species and sex of each bat based on its

morphology (Poon et al., 2023, Shek, 2006). The fresh feces from each bat were

collected and placed in individual 2mL tubes. They were preserved in 100% ethanol

and then kept at -80°C until DNA extraction. In total, we have collected fecal samples

from 62 *M. pilosus*, 51 *M. horsfieldii*, and 43 *M. chinensis*. Physical parameters of

each bat, such as body weight and forearm length, were measured. After collecting

samples and recording their physical parameters, all bats were released immediately

back to the wild. Approvals for animal experiments were granted by the Department

of Health (ref. 19-177 in DH/SHS/8/2/3 Pt. 30), the Committee on the Use of Live

Animals (ref. 4963-19), and the Agriculture, Fisheries, and Conservation Department

(AFCD; ref. 35 in AF GR CON 09/51 Pt.8).

2.2 DNA extraction and metabarcoding

The QIAamp Fast DNA Stool Mini Kit (Qiagen, Hilden, Germany) was used to

extract the fecal DNA. We included negative controls and mock communities (Table

S2a) during DNA extraction to later check for contamination during PCR. We

quantified the fecal DNA using the Qubit dsDNA high-sensitivity (HS) assays on an

Invitrogen Qubit 4 Fluorometer (Thermo Fisher Scientific, Massachusetts, US)

(Huang et al., 2021).

All fecal DNA samples, mock communities (Table S2a), and negative controls

were used for library preparation using three genetic markers through two-step PCR

(Huang et al., 2022, Huang et al., 2021) (Supplementary Materials and Methods). The

first pair of markers used was a universal pair (18s_SSU3_F:

5'GGTCTGTGATGCCCTTAGATG3' and 18s_SSU3_R:

5'GGTGTGTACAAAGGGCAGGG3') which targets the V7 region of 18S small

subunit ribosomal DNA (rDNA; ca. 174bp) (McInnes et al., 2017). This pair of

markers provides an overview of the dietary compositions of *Myotis* bats. To

investigate whether fish was consumed by the three *Myotis* species, we used the

mitochondrial 12S rDNA specific primers (12S_AcMDB07_HK_F:

5'GCCTATATACCRCCGTCG3' and 12S_AcMDB07_ R:

5'GTACACTTACCATGTTACGACTT3'; ca. 283bp) to amplify fish DNA (Bylemans

et al., 2018). Considering the important prey of *Myotis* bats, which consists of

arthropods, we used a third pair of markers which amplifies the mitochondrial

cytochrome c oxidase subunit I gene (COI; COI_Fwh2_F: 5'

GGDACWGGWTGAACWGTWTAYCCHCC3' and COI_Fwh2_R: 5'

GTRATWGCHCCDGCTARWACWGG3'; ca. 219bp) (Vamos et al., 2017). This pair

of markers offers a higher resolution in identifying the macroinvertebrate species

consumed by the bats. For each marker, we generated a library multiplex by pooling

libraries from different samples in an equimolar ratio. Subsequently, all three libraries

were sequenced on an Illumina NovaSeq (PE 150bp) by Novogene (Hong Kong).

2.3 Sequencing data preprocessing

Raw paired-end DNA reads were merged by using the -fastq_mergepairs function in

USEARCH v11.0.667 (Edgar, 2010). Primer sequences were removed with

CUTADAPT v2.5 (Martin, 2011). The assessment of trimmed reads quality was

completed with FastQC v0.11.9 (Wingett and Andrews, 2018) and VSEARCH

v2.18.0 (Rognes et al., 2016). Only high-quality trimmed reads within the target

lengths (18S: 130-180bp; COI: 130-210bp; 12S: 140-290bp) were retained for later

analysis. These pre-processed reads were then dereplicated by using the -

derep_fulllength command in VSEARCH. Chimeras and singletons were removed

from the dereplicated reads by using USEARCH. Using the -usearch_global function

in VSEARCH, all pre-processed reads were clustered into amplicon sequence variants

(ASVs) based on 99% similarity.

Using the SINTAX algorithm in USEARCH, each ASV was assigned to the

lowest identifiable taxonomic level with a confidential cutoff of 0.7 (Edgar, 2016).

The ribosomal RNA database SILVA (Glöckner et al., 2017), the mitochondrial

database MIDORI for eukaryotes (Leray et al., 2022), and MitoFish for fish (Zhu et

al., 2023) were used as taxonomic classification reference databases for 18S, COI, and

12S sequences, respectively. We also assigned ASVs against the NCBI non-redundant

nucleotide sequences database to obtain the best 1000 blast hits with a similarity

higher than 99% and e-value lesser than 1e-50. The BASTA with the lowest common

ancestor (LCA) algorithm (Kahlke and Ralph, 2019) helped us assign the lowest

common taxonomic level shared by 80% of blast hits. To obtain a high taxonomic

classification resolution, the results from SINTAX and LCA were combined by

assigning ASVs with lower ranks of taxonomies.

Potential false-positive and contaminant ASVs were eliminated by comparing

them with mock communities and negative controls (Table S2b-d). Non-diet ASVs

(e.g., Fungi, Bacteria, and Algae) and unclassifiable ASVs (e.g., ASVs that could not

be identified beyond the domain level) were discarded from the analysis. After ASVs

cleaning, samples with reads number lower than 100 were subsequently excluded. To

better present our taxonomic data in figures, we classified each identified taxon as

high and low abundance taxon according to whether it occupies over 0.1% number of

reads. High abundance taxa were grouped to the lowest taxonomic level, while low

abundance taxa were grouped into higher taxonomic level. In this way, the taxa were

classified into taxonomic categories (Table S3a-c) (Huang et al., 2022).

2.4 Data analysis

After data preprocessing, we used 18S data of 144 samples, 12S data of 48 samples,

and COI data of 100 samples for downstream data analysis. In the 18S dataset, a total

of 11 taxonomic categories were classified from 91 ASVs. The identifiable taxonomic

levels categorized included three classes, eight orders, two families, and one genus

(Table S3a). In the 12S dataset, 22 taxonomic categories were classified from 24

ASVs. The taxonomic levels included nine families and 12 species (Table S3b). In the

COI dataset, 29 taxonomic categories were classified from 654 ASVs. The taxonomic

levels included two classes, 18 orders, 13 families, 19 genera, and 18 species (Table

S3c). In the statistical analysis, we calculated (1) the percentage of read count for each

taxon in a sample (relative read abundance, RRA), (2) the percentage of occurrence

for each taxon in a sample (weighted percentage of occurrence, wPOO), and (3) the

proportion of samples in which a taxon is detected (frequency of occurrence, FOO).

The RRA or wPOO at the population level is presented as the mean of RRA or wPOO

of all individual samples of a bat species (Lee et al., 2021). We conducted these

analysis in R v4.2.1 and visualized with the R package ggplot2 v3.4.2 (Hadley, 2016).

2.4.1 Diet diversity analysis

To determine the relationships between sample numbers and dietary species Chao2

diversity, we generated rarefaction curves using hill numbers from ASVs or taxa

(q=0) via the R package iNext (Hsieh et al., 2016) (Fig. S2). Using hill numbers based

on variant q values (the order of diversity), we estimated the diet diversity of three

Myotis species at both individual (alpha diversity) and population/species (gamma

diversity) levels. Hill number only considers the occurrence of each ASV when q=0.

The weight of species abundance in hill number increases when q value increases.

The hill number is equivalent to the exponential of Shannon's diversity index and

inversion of Simpson's dominance at q=1 and q=2, respectively (Alberdi and Gilbert,

2019). Pairwise diversity comparisons at the individual level (alpha) between *Myotis*

species were carried out using the Kruskal–Wallis test, followed by Benjamini-

Hochberg correction ($p<0.05$).

Beta diversity refers to the differences in dietary taxa compositions between

individual samples. To analyze beta diversity, we calculated pairwise binary Jaccard

dissimilarity distances and pairwise Bray–Curtis dissimilarity distances using the

occurrence of each ASV and the fourth root transformed RRA of each ASV,

respectively. We visualized the dissimilarities with principal coordinates analysis

(PCoA) with the plot_ordination function in R package phyloseq v1.42.0 (McMurdie

and Holmes, 2019), and fitted the effect of each taxonomic category into the PCoA

result using the envfit and ordiArrowMul function in R package vegan v2.6.4

(Oksanen et al., 2022). Pairwise permutational multivariate analysis of variance

(PERMANOVA) tests were performed to investigate the separation of dietary

compositions between the *Myotis* species. Significant interspecific composition

variants observed in PERMANOVA tests are premised on the intraspecific

homogeneity, which were tested by betadisper function. We also carried out similarity

percentage (SIMPER) analyses to assess the contribution of each taxon to the

difference between *Myotis* species by using R scripts simper_pretty.R (Steinberger et

al., 2016), which were then checked by the Kruskal-Wallis test by using R scripts

R_krusk.R (Steinberger, 2020).

Within each species, we performed multiple analyses to determine the

contribution of host variables (i.e., sex) and environmental variables (i.e., sampling

seasons and sampling locations) to the diversity variation. Data from SK01-03 and

PTC were combined for analyses due to their close geographic proximity. Generalized

linear models were performed to evaluate the effect of variables on alpha dietary

diversity by using the logarithmic hill number of order $q=1$ as the dependent variable.

We visualized the beta diversity dissimilarities of variable groups using PCoA based

on Bray-Curtis and Jaccard dissimilarity distances with fitted taxonomic categories.

PERMANOVA and corresponding beta-dispersion tests were used to assess the beta

diversity variations among variable groups. We identified taxa that significantly

contributed to the composition variation among variable groups, using the SIMPER

test, which was checked by the Kruskal–Wallis test. The dietary composition overlap

between bat individuals was measured by Pianka's niche overlap index using the R

package "spaa" v0.2.2 (Zhang et al., 2016). A network was generated to visualize

individual-level diet overlap by using the "qgraph" package v1.9.5 (Pedersen et al.,

2017).

3. Results

3.1 Dietary compositions of the three* Myotis *bat species

Based on 18S data, the three *Myotis* species primarily preyed upon insects, arachnids,

and/or ray-finned fish (Actinopteri) (Fig. 1 and S3). Specifically, flies (dipterans),

moths and/or butterflies (lepidopterans), orthopterans, and true bugs (hemipterans),

were the most commonly consumed insect groups. Arachnids such as spiders

(Araneae) and mites (Mesostigmata), as well as fish also formed part of their diet.

The three *Myotis* species exhibited noticeable interspecific variations in their

primary food sources (Fig. 1b-d). *Myotis chinensis* (18S, n=40) primarily consumed

insects (72% RRA), particularly bush crickets (*Sasima* spp. in order Orthoptera at

39%) and moths and/or butterflies (22%). Additionally, spiders accounted for 27% of

their diet (Fig. 1b and Table S4). *Myotis horsfieldii* (18S, n=49) mainly hunted smaller

size insects (91%), with flies (55%) and true bugs (16%) being the primary prey. They

consumed arachnids (7%) but in smaller proportions (Fig. 1c; Table S4).

Fish was mainly found in the diets of *M. pilosus* (n=55; Fig. 1d; Table S4). The

18S data also revealed the presence of fish contents in two samples of *M. horsfieldii*,

which were not detected by the 12S (Fig. 1a and 1c; Table S5a-b). To validate the

presence of fish in the samples of *M. horsfieldii*, we first confirmed the bat species

identity by DNA barcoding using primers SFF_145f and SFF_351r (Walker et al.,

2016), and then we confirmed the fish contents of these samples by performing PCR

using the 18S and 12S primers (section 2.2) and Sanger sequencing. We concluded

that the results were consistent with the DNA metabarcoding finding that fish was

identified in these two samples.

Myotis pilosus (18S, n=55) displayed significant intraspecific differences in diets

(Fig. 1 and 2a; Table 6a-b). Approximately one-fifth (9/55) of the individuals were

found to exclusively feed on macroinvertebrates, with a majority of these individuals

found in the wet season. These individuals primarily prey on moths and/or butterflies.

In contrast, the remaining individuals preyed on fish in varying degrees, constituting

6-100% of their individual diets (Fig. 1a and 1d; Table S5a-b).

Based on the 12S metabarcoding data, all individuals that consumed fish were *M.*

pilosus (Fig. 2a and S4; Table S6a-c). Both freshwater, marine, or brackish fish were

found in the diet of *M. pilosus*, at least twelve fish species were detected. *Myotis*

pilosus primarily hunted marine fish, with Old World silversides (Atherinidae)

making up 33% of their diet, including wide-banded hardyhead silverside

(*Atherinomorus lacunosus*, max. size <25cm) at 31%. Mullets (Mugilidae) comprised

31% of their diet, with flathead grey mullet (*Mugil cephalus*, common max. size

<50cm) making up 17%, and eastern keelback mullet (*Planiliza affinis*, max. size

<40cm) and kanda mullet (*Osteomugil engeli*, max. size <48cm) each contributing

about 6%. Herrings and sprats (Clupeidae) accounted for 12% of their diet, with

silver-stripe round herring (*Spratelloides gracilis*, max. size <11cm) being commonly

consumed at 10%. Mosquitofish (*Gambusia affinis*, max. size <5.1cm, from the

family Poeciliidae) was the dominant freshwater fish at 13% (Froese and Pauly,

2024). Notably, mosquitofish was mainly consumed by individuals from Tai Lam

Chung (Fig. 2; Table S6a-b).

The COI data offered a more detailed revelation of the macroinvertebrate prey

consumed by the *Myotis* species (Fig. 3 and S5; Table S7a-c). Overall, the majority of

prey consisted of dipterans (about 32%), such as lake flies in the family Chironomidae

(e.g., *Procladius culiciformis*, *Glyptotendipes tokunagai*, *Chironomus flaviplumus*,

etc.); lepidopterans (14%), including grass moths like *Syntonarcha iriastis* and

Cirrhochrista brizoalis; orthopterans (13%), including crickets like *Mecopoda* spp.

(bush crickets) and *Gryllotalpa* spp. (mole cricket); and spiders (12%), such as

northern golden orb weaver (*Nephila pilipes*) (Fig. 3; Table S7a).

The macroinvertebrate orders in diets revealed by COI were consistent with those

unveiled by 18S. Notable variations were observed in the dietary compositions among

Myotis species, as revealed by COI. *Myotis chinensis* (COI, n=34) primarily preyed

on arthropods in several orders, such as bush cricket *Mecopoda elongata* (23%) from

Orthoptera (30%), *S. iriastis* (8%) from Lepidoptera (16%), and *N. pilipes* (7.5%)

from Araneae (25%). Spiders were found in more than 50% of the *M. chinensis*

samples (Fig. S5b; Table S7a). In *M. horsfieldii* (COI, n=44), smaller size insects like

dipterans accounted for more than 66% of their total diet, with *P. culiciformis* (24%)

and *G. tokunagai* (15%) being the main contributors. More than 50% of the *M.*

horsfieldii samples contained various lake fly species. Lepidopterans and

hemipterans, on the other hand, made up 9% and 10% respectively (Fig. 3c; Table

S7a). Although a larger proportion of 30% of insects remained unidentified compared

to the other two bat species, the primary prey of *M. pilosus* (COI, n=22) consisted of

arthropods from various orders, including Lepidoptera (20%), Araneae (15%),

Orthoptera (7%), Hemiptera (8%), and Trichoptera (caddisflies, 7%) (Fig. 3d; Table

S7a).

3.2 Alpha and gamma diversity of consumed taxa

The alpha diversity of individual diets indicated that the diets of *M. horsfieldii*

exhibits the highest diversity compared to the other two *Myotis* species, as shown by

18S and COI data (Fig. 4 and 5; Table S8a-b). Notably, there is a distinct decrease in

hill numbers as the *q* value increases, particularly observed in the macroinvertebrate

(COI) composition of *M. horsfieldii* and *M. chinensis* diets (Fig. 5; Table S8b). This

suggests that although these bats consume a greater variety of macroinvertebrates

compared to *M. pilosus*, their prey compositions are highly uneven, with certain taxa

dominating their diets. Furthermore, the macroinvertebrate (COI) compositions of *M.*

chinensis were more unevenly distributed compared to *M. horsfieldii* (Fig. 5; Table

S8b). In the case of individuals of *M. pilosus* that consumed fish, mostly two to three

fish species were detected in their diets (12S data; Fig. 2a; Fig. 5). A similar pattern of

prey consumption based on 18S data was also observed at the population level as

reflected by gamma diversity, with the dietary composition of *M. horsfieldii* being the

most diverse (Fig. 5; Table S8a). Despite having the lowest macroinvertebrate (COI)

diversity in their diets, *M. pilosus* displayed comparable diversity of overall (18S)

diets to *M. chinensis* at the population level (Fig. 5; Table S8a-b).

3.3 Effects of environmental and host factors on the dietary compositions

According to the PERMANOVA tests conducted using Bray–Curtis and Jaccard

dissimilarity distances, the diets of *M. pilosus* were found to differ significantly

between seasons and locations based on 12S (Table S9a-b) and 18S (Table S10a-b)

data. Specifically, during the wet season, there was a higher consumption of

macroinvertebrates by *M. pilosus*, mainly lepidopterans as evidenced by the 18S data,

while in the dry season more fish was consumed. Moreover, the fish compositions of

M. pilosus from SK, where a majority of samples that were collected from, exhibited

notable variations between the wet and dry seasons (Fig. 2a). The SIMPER analysis

results further identified the main contributors to location difference. Old World

silverside (Atherinidae) were consumed more at SK, while mosquitofish were

consumed more at TLC (Fig. 6a-b and Table S11a-b) (Froese and Pauly, 2024).

Similarly, the diets of *M. chinensis* also differed significantly between the two

seasons based on both 18S (Fig. 1a and Table S10a-b) and COI (Fig. 3a and Table

S12a-b) data. The 18S data showed that during the wet season, *M. chinensis* mainly

consumed moths and/or butterflies (Lepidoptera), while in the dry season, they

consumed a large proportion of *Sasima* bush crickets (Orthoptera) (Fig. 1a).

Although both GLM and PERMANOVA analyses consistently demonstrated a

significant divergence in the seasonal diets of *M. horsfieldii* (Table S12a-b and S13),

it's important to note that the small sample size (n=4) from the wet season in the

COI dataset may limit the accuracy of these findings. Further investigations are

required in the future to confirm this result. Furthermore, the PERMANOVA

analysis revealed an additional noteworthy disparity in the diets of the two sexes of

M. horsfieldii (Table S12a-b), which was mainly attributed to the distinct

consumption of dipterans, lepidopterans, and hemipterans by each sex (Table S14a-

d).

3.4 Dietary niche partitioning between the three* Myotis *species

Based on the PCoA analysis, 18S and COI data revealed highly similar patterns of

dietary niche partitioning among the three *Myotis* species (Fig. 6c-f). The PCoA

analysis using Bray–Curtis and Jaccard distances within each of the 18S and COI

dataset also revealed consistent patterns. The overall (18S) and macroinvertebrate

(COI) diets of *M. horsfieldii* were distinctly differentiated from those of *M. pilosus*

and *M. chinensis*, while the dietary compositions of *M. pilosus* and *M. chinensis*

were more overlapping with each other (Fig. 6c-f). The SIMPER analysis results

indicated the main contributors to this differentiation in *M. horsfieldii* were

predominantly dipterans and hemipterans, which constituted a significant portion of

their diet (Fig. 6c-d; Table S15a-b). Specifically, the true bugs included water

striders (e.g., *Asclepios apicalis*) while dipterans included a group of lake flies (e.g.,

P. culiciformis, *G. tokunagai*, *Kiefferulus tainanus*, *Polypedilum nubifer*,

Chironomus spp. etc.) (Fig. 6e-f; Table S16a-b).

Myotis chinensis also differentiated from *M. horsfieldii* and *M. pilosus* by

consuming a considerable proportion of bush crickets (Orthoptera), such as *Sasima*

spp. and *M. elongata*, as well as spiders (Araneae, including *N. pilipes*) (Fig. 6c-f;

Table S15a-b and S16a-b). On the other hand, the dietary niche of *M. pilosus*

differed from the other two species by including a large portion of fish (Actinoptera)

in their diets (Fig. 6c-d and Table S15a-b). *M. pilosus* also differentiated from *M.*

chinensis by consuming caddisflies (Trichoptera) (Fig. 6e-f; Table S16a-b).

At the individual level, *M. horsfieldii* showed greater intraspecific overlap of

individual diets compared to those of *M. chinensis* and *M. pilosus* (Fig. S6). This

suggests the dietary compositions among *M. horsfieldii* individuals were more similar,

and the dietary compositions among individuals of *M. chinensis* and *M. pilosus* were

more variable.

4. Discussion

Fish eating behavior of *Myotis*

This study provides new insights into the dietary compositions of *M. pilosus* and *M.*

chinensis in coastal habitats, expanding on previous research that primarily focused

on inland regions of China. Additionally, our research contributes to the

understanding of the dietary compositions of *M. horsfieldii*, a species found in

southeast and south Asia that has not been previously studied in terms of its

foraging ecology. Contrary to earlier studies conducted in inland regions of China,

which suggested that *M. pilosus* primarily fed on cyprinids in freshwater habitats

year-round (Ma et al., 2006, Chang et al., 2019), our research in Hong Kong

unveiled a different dietary pattern. Our findings show that *M. pilosus* in Hong

Kong did not prey on cyprinids but instead exhibited a diverse diet, targeting a wide

variety of fish prey species from nine different families. These prey species

encompassed marine, brackish, and freshwater fish, highlighting the versatility of

M. pilosus in its feeding habits. Moreover, we observed variations in the species

compositions consumed by *M. pilosus* across different seasons. For instance, marine

species such as *A. lacunosus*, *Doboatherina duodecimalis* (tropical silverside, max.

size 11cm), *Parexocoetus brachypterus* (sailfin flying fish, max. size 13cm), and *E.*

heteroloba were only fished by *M. pilosus* during the wet season at Sai Kung. This

was observed specifically at sampling site SK01 but not the other sites. Worth noting

is that SK01 is located 1.7 km or less away from the nearest shore at the southwest,

making it a likely site for *M. pilosus* to fish rather than flying north- or eastward to

reach the shore, which is located over 3 km away.

Atherinomorus lacunosus, which forms large schools along sandy shorelines,

serves as an important food source for *M. pilosus*. This fish species forages

nocturnally to capture zooplankton that migrate vertically to the upper water column

during nocturnal hours (Skibinski, 2005). This feeding behavior creates large amount

of ripples that trawling bats like *M. pilosus* can detect. Another noteworthy prey

species is *P. brachypterus*. Sailfin flying fish is commonly found in coastal waters

in large shoals and possessing elongated, wing-like pectoral fins, which enable it to

leap out of the water and glide rapidly for considerable distances above the surface,

an adaptation to evade underwater predators. During the breeding season, spawning

flying fish aggregate in abundance near the surface at night, with many vigorously

jumping and flying out of the water to release ripe eggs and sperm (Stevens et al.,

2003, Digo et al., 2015, Lewis, 1961), exposing them to attack by aerial predators,

such as *M. pilosus*.

During the dry season in Sai Kung, *M. pilosus* captured the marine fish species

Spratelloides gracilis and *Sardinella melanura* (blacktip sardinella, common size

10cm). Clupeids, such as *S. gracilis* and *S. melanura*, are primarily forage fish known

for their high egg production. For instance, *S. gracilis* spawns near the water surface,

releasing approximately 1600 eggs per unit body mass (Dalzell and Wankowski,

1980). Although previous studies in China have reported crustaceans as prey for *M.*

pilosus and other bat species (Wang et al., 2024), our study did not identify any

crustaceans in the diet of *M. pilosus*. Based on our results, one notable target species

for *M. pilosus* is *M. cephalus*, along with other mullet species, during dry months.

Mugil cephalus is known for congregating in schools over sand or mud in coastal and

brackish waters. In Hong Kong, this species exhibits two short spawning peaks in

winter (dry) months, resulting in a large numbers of juveniles appearing inshore,

especially in estuaries, during February and March (Sadovy and Cornish, 2000).

In the Tai Lam Chung woodland area, which is an inland, *M. pilosus* primarily

preys on the exotic freshwater poeciliid, *G. affinis*. The southern shore is obstructed

by highland with an elevation of at least 400m, and it is at least 9 km away from the

nearest shore in any direction (Fig. S1). *Gambusia affinis* is a freshwater species and a

widespread non-native fish in Hong Kong, originating from North America and

released by the Hong Kong Government for mosquito control purposes around 1940

(Dudgeon and Corlett, 2004). Surveys on the distribution of *G. affinis* in Hong Kong

have reported its presence in the lower reaches of the closest streams within less than

1km of TLC forest (Tsang and Dudgeon, 2021a). Experiments conducted on the

ecological effects of *G. affinis* have shown that it reduces the abundance and richness

of invertebrates and alters assemblage compositions in Hong Kong wetland (Tsang

and Dudgeon, 2021b). Our findings suggest that *M. pilosus* may play a potential role

in controlling invasive poeciliids in the local ecosystem. Similar findings have been

reported in *M. capaccinii* in northwest Israel and eastern Iberian Peninsula, where *M.*

capaccinii fed on exotic *Gambusia* spp. (Aizpurua et al., 2014, Levin et al., 2006).

Mosquito fish frequently swim close to water surface, using their upturned mouth to

break the surface and capture floating insects. Furthermore, the decrease in oxygen

during the night may compel these fish to come to the water surface for breathing

(Aizpurua et al., 2014, Levin et al., 2006).

Our study has made an intriguing discovery regarding the feeding habits of *M.*

horsfieldii. We reveal two individuals *M. horsfieldii* from Sai Kung have included

fish in their diet. Interestingly, we have found the first case of *M. horsfieldii* preyed

on *Asclepios apicalis*, a water strider species that is also targeted by *M. pilosus*.

While most water striders are known for their capacity to glide effortlessly on the

calm freshwater surface, *Asclepios* spp. are considered as sea skaters which are

found primarily in brackish water along coasts (Poolprasert et al., 2022, Andersen

and Foster, 1992). *Myotis horsfieldii* might be drawn to this food source due to the

consistent ripples produced by *Asclepios* as they manoeuvre across water's surface.

Reports on *M. horsfieldii*'s foraging behavior are limited, but one literature mentions

that *M. horsfieldii* typically flies in circles around 10 cm above water to skim

insects, and they roost near water sources (Wilson and Mittermeier 2019). Our

finding suggests that *M. horsfieldii* may capture fish while foraging insects over

waters. Despite this finding, it should be noted that fish was only detected in very few

individuals of *M. horsfieldii*. Therefore, further investigation is needed to determine

whether *M. horsfieldii* actively engage in fishing or if those individuals simply

mistakenly catch live or dead fish on the water surface while foraging for insects. Our

discovery that *M. horsfieldii*, which was not previously known to consume fish, does

in fact include fish in their diets has significant implications for the evolution of

fishing behavior. The active foraging of aquatic insects by bats may lead to occasional

consumption of fish, which in turn creates the selective pressure that drives the

evolution of fishing behavior.

Foraging niches of the three *Myotis* species

We observed a higher proportion of fish in the diet of *M. pilosus* during the dry

season compared to the wet season, with a significant increase in insect components

during the wet season, which is consistent with Ma et. al. (2006) but different from

Chang et al. (2018). Our results showed that this increase in insect consumption was

due to a 43% rise in the number of individuals preyed on macroinvertebrates during

the wet season (22/35) compared to the dry season (4/20). Furthermore, while there

were some variations in the insect dietary compositions compared to previous

research, our study revealed a similar finding to Chang et al. (2019) in terms of *M.*

pilosus primarily preying on lepidopterans in the wet season. Based on the nocturnal

behavior of most moths and the fact that the lepidopterans identified in our study are

all moth species, such as *S. iriastis* and *C. brizoalis* consumed by *M. pilosus*, we

suggest that most of the lepidopterans preyed upon by *M. pilosus* are moths. Surveys

on the abundance of forest invertebrates in Hong Kong have shown that local

lepidopterans vary seasonally, with both number and biomass experiencing winter

(dry season) lows and early summer (in May during wet season) maxima (Kai and

Corlett, 2002). During the wet season, local *M. pilosus* also targeted trichopterans.

Trichopterans are moth-like and closely related to lepidopterans, which are nocturnal

and commonly associated with freshwater bodies (De Moor and Ivanov, 2008).

Here, we provide additional evidence for the specialization of *M. pilosus* in

fishing, as we have consistently found fish in its diet throughout the year. It is worth

noting that the abundance of lepidopterans in local terrestrial habitat is higher

during summer (Kai and Corlett, 2002), this high abundance likely explains why

some individuals of *M. pilosus* have shifted their target prey from fish to insects

during this time. This shift may be driven by higher energetic profit associated with

capturing a larger number of moths compared to capturing fish. Despite the higher

nutritional value of a single fish compared to a single moth, catching fish likely

incurs a greater energy cost (Aizpurua et al., 2013). This is due to the heavier body

mass of fish and the increased effort required to accurately detect fish underwater

and subsequently pull it out of the water (Aihartza et al., 2008). This results in lower

capture efficiency when targeting fish (Altenbach, 1989). Therefore, our study

highlights the adaptive feeding behavior of *M. pilosus*, which adjusts its diet based

on prey availability, underscoring the significance of this feeding strategy.

Our results show that there is a greater overlap in the macroinvertebrate diet

between *M. pilosus* and *M. chinensis*, while the diet of *M. horsfieldii* is distinct from

the other two species. We found that local *M. chinensis*, similar to *M. pilosus*, also

hunted a higher proportion of lepidopterans, such as grass moth, *S. iriastis*, during the

wet season. Another noteworthy foraging habit shared between *M. chinensis* and *M.*

pilosus is their considerable consumption of non-flying nocturnal arthropods,

specifically spiders like the northern golden orb weaver (*N. pilipes*). *Nephila pilipes*

typically build intricate webs to capture prey on bushes and trees near water sources

(Harvey et al., 2007). Spiders have been documented as part of the diet of several

other *Myotis* species, such as *M. myotis*, *M. emarginatus*, *M. lucifugus*, *M. nattereri*,

M. evotis, and *M. septentrionalis* (Goiti et al., 2011, Maucieri and Barclay, 2021,

Swift and Racey, 2002, Kaupas and Barclay, 2018). However, previous studies did

not report spiders as a food source for *Myotis* bats in China (Ma et al., 2003, Ma et

al., 2008, Ma et al., 2006). Moreover, *M. chinensis* mainly preyed on orthopterans,

especially bush crickets, such as *Mecopoda* and *Sasima* spp., during the dry season.

The consumption of spiders and orthopterans suggests that *M. chinensis* is capable

of foraging in cluttered environments of woodlands and gleans arthropods from

various substrate surfaces, such as spider webs and/or foliage in bushes. While there

are some geographical variations in dietary composition, our findings agree with Ma

et al. (2008), which remarked that conspecifics in Beijing prey on 'non-wing

beating' insects such as coleopterans (e.g., ground beetles in Carabidae) and

orthopterans while lacking hymenopterans (sawflies, wasps, bees, and ants) and

lepidopterans (Ma et al., 2008).

Based on our results, we found that the diet of *M. horsfieldii* is distinctive,

which prominently includes a high consumption of smaller flying insects,

specifically dipterans. Notably, *M. horsfieldii* preys on a wide variety of lake fly

species from the Chironomidae family, leading to a higher dietary diversity

compared to *M. chinensis* and *M. pilosus*. Conversely, our observations indicate that

spiders, orthopterans, and coleopterans were seldom found in their excrement. In the

forests of Hong Kong, dipterans constitute the majority of the insect biomass, and

their abundance did not exhibit seasonal variation (Kai and Corlett, 2002).

Numerous chironomid species bear a striking resemblance to mosquitoes and adults

swarms in terrestrial habitats serve as vital sources of food for bats (Puig-Montserrat

et al., 2020, Beck, 1995). Local *M. horsfieldii* also targets hemipterans and

lepidopterans, though to a lesser extent compared to dipterans.

Our findings indicate that *M. pilous* is primarily a fish-eating bat, while *M.*

chinensis and *M. horsfieldii* primarily feed on insects. One plausible reason for this

dietary differentiation pattern could be differences in their morphological

characteristics. Previous study suggested that the body size of aerial hunting bats

limits their weight-carrying capacity, and fishing bat species tend to have relatively

larger body sizes and longer hind feet that give them morphological advantages to

fish hunting (Chang et al., 2019). Despite *M. chinensis* (91-97mm body length, 25-

30g) (Wilson and Mittermeier 2019) and *M. pilosus* (51-65mm, 11.7-32.5g)

exhibiting larger body sizes and heavier body masses, *M. pilosus* share a similar

hind foot to forearm length ratio (0.31±0.014) to other fishing bats (0.31±0.051)

(Chang et al., 2019), while the ratio for *M. chinensis* (0.26) is more similar to those

of other non-fishing bats (0.25±0.038). *Myotis horsfieldii* (44-51mm, 5.0-7.5g)

(Wilson and Mittermeier 2019) is smaller in size compared to the other two species.

It has a low value ratio (0.23), which is also more akin to other non-fishing trawling

bats (Chang et al., 2019). Apart from the influence of morphological adaptations on

prey capturing ability, other factors contributing to the dietary differentiation pattern

could include differences in prey detection ability through variations in echolocation

call structures during foraging (Aizpurua and Alberdi, 2018).

This dietary study uncovered the overlapping macroinvertebrate diets of *M.*

pilosus and *M. chinensis*, as well as the distinct diet of *M. horsfieldii*, with the latter

being documented for the first time. Both *M. pilosus* and *M. chinensis* are

considered regionally threatened. The potential interspecific competition observed

in our findings, particularly between *M. pilosus* and *M. chinensis* during wet season,

underscores the importance of protecting macroinvertebrate food sources in local

terrestrial ecosystem to prevent intensifying competition among the congenerics. To

safeguard the sustainability of vulnerable *M. pilosus*, it is also crucial to preserve the

integrity of marine, brackish, and freshwater ecosystems in Hong Kong, as these

diverse environments serve as essential foraging grounds for *M. pilosus* to capture

the diverse range of fish species.

5. Conflict of Interest

The authors declare that the research was conducted in the absence of any commercial

or financial relationships that could be construed as a potential conflict of interest.

6. Data Availability Statement

Raw sequencing data will be uploaded to the NCBI Sequence Read Archive (SRA).

Accession number will be provided after the manuscript is accepted.

Author contributions:

S.Y.W.S. Conceptualization; E.S.K.P. and S.Y.W.S. Methodology; E.S.K.P., J.C.T.C.,

D.T.C.C., C.T.S., W.C.T., and S.Y.W.S. Investigation; X.W. Formal analysis; C.T.S.,

W.C.T., and S.Y.W.S. Resources; X.W. Visualization; X.W. and E.S.K.P. Writing -

Original Draft; E.S.K.P., S.Y.W.S. and all authors Writing - Review & Editing;

S.Y.W.S. Supervision; S.Y.W.S. Project administration; S.Y.W.S. Funding acquisition.

Statement on ethics:

Acknowledgments:

This study was funded by the Environment and Conservation Fund (ref. ECF

94/2019) in Hong Kong, China. We express our gratitude to the AFCD of the

HKSAR Government for their generous support. Special thanks are extended to Chi

Pan Tong and Alex Wai Kit Lo for their assistance during our study. The High-

Performance Computing (HPC) service provided by the Information Technology

Services at the University of Hong Kong supported our data analysis.

7. Figures Legends and Tables

Fig. 1. Dietary composition of *Myotis* bat species based on taxa detected in fecal

samples using the 18S rDNA marker. The relative read abundance of each taxon were

visualized at both (a) individual level (n=144) and (b-d) species level for (b) *M.*

chinensis (n=40), (c) *M. horsfieldii* (n=49), and (d) *M. pilosus* (n=55). Prey in classes

with read abundance lower than 0.1% of all taxa detected were grouped into "Others".

Only taxa with abundance higher than 1% in relative read abundance are shown. F,

Family. O, Order. Bat photos © Agriculture, Fisheries and Conservation Department.

Fig. 2. Dietary composition of *Myotis pilosus* based on taxa detected in fecal samples

using the 12S rDNA marker. The relative read abundance of each taxon were

visualized at both (a) individual (n=48) and (b) species levels. Individual bars were

grouped by sampling locations and seasons. Prey in classes with read abundance

lower than 0.1% of all taxa detected were grouped into "Others". Only taxa with

abundance higher than 1% in relative read abundance are shown. Refer to Fig. S1 for

locations. SK, Sai Kung (including Pak Tam Chung); TLC, Tai Lam Chung; LMH,

Lin Ma Hang.

Fig. 3. Dietary composition of *Myotis* bat species based on taxa detected in fecal

samples using the COI marker. The relative read abundance of each taxon was

visualized at both (a) individual level (n=102) and (b-d) species level for (b) *M.*

chinensis (n=34), (c) *M. horsfieldii* (n=44), and (d) *M. pilosus* (n=22). Prey in classes

with read abundance lower than 0.1% of all taxa detected were grouped into "Others".

Only taxa with abundance higher than 1% in relative read abundance are shown. G,

Genus. F, Family. O, Order.

Fig. 4. Pairwise alpha diversity comparison between *Myotis* species. Alpha diversity

is represented by using the hill numbers of order $q=0$ (upper panel), $q=1$ (middle

panel), and $q=2$ (lower panel), which were calculated based on the ASVs abundance

detected by18S rDNA (left panel) and COI (right panel) marker at individual level.

Only the *p*-values of species pairs which have significant different alpha diversity

were shown ($p<0.05$).

Fig. 5 Dietary diversity for each *Myotis* represented in the hill numbers of variant

order *q*. The hill numbers were calculated at individual level and species level to

estimate the (a) alpha diversity and (b) gamma diversity, respectively. The ASVs

abundance of each sample were detected by using 18S rDNA (left panel), COI

(middle panel), and 12S rDNA (right panel) marker gene.

Fig. 6 Principal coordinate analysis (PCoA) of dietary compositions in (a, b) *Myotis*

pilosus and (c-f) three *Myotis* species based on (a, c, e) the pairwise Bray–Curtis

dissimilarity distances calculated from the fourth root transformed RRA of each ASV,

and (b, d, f) the pairwise Jaccard dissimilarity distances calculated from the

occurrence of each ASV. Each point corresponds to one sample, with the color

indicates (a, b) sampling locations, seasons, and (c-f) *Myotis* species. The ASVs were

identified by the (a, b) 12S rDNA, (c, d) 18S rDNA, and (e, f) COI marker gene.

Taxonomic categories significantly contributing to the difference between groups

were shown (see Table S11, S15, and S16). Refer to Fig. S1 for locations. SK, Sai

Kung (including Pak Tam Chung); TLC, Tai Lam Chung; LMH, Lin Ma Hang.

8. References

AIHARTZA, J., ALMENAR, D., SALSAMENDI, E., GOITI, U. & GARIN, I. 2008.
Fishing behaviour in the long-fingered bat Myotis capaccinii (Bonaparte,
1837): an experimental approach. *Acta Chiropterologica,* 10**,** 287-301.
AIHARTZA, J. R., GOITI, U., ALMENAR, D. & GARIN, I. 2003. Evidences of
piscivory by Myotis capaccinii (Bonaparte, 1837) in southern Iberian
Peninsula. *Acta chiropterologica,* 5**,** 193-198.
AIZPURUA, O., AIHARTZA, J., ALBERDI, A., BAAGØE, H. J. & GARIN, I. 2014.
Fine-tuned echolocation and capture-flight of Myotis capaccinii when facing
different-sized insect and fish prey. *Journal of Experimental Biology,* 217**,**
3318-3325.
AIZPURUA, O. & ALBERDI, A. 2018. Ecology and evolutionary biology of fishing
bats. *Mammal Review,* 48**,** 284-297.
AIZPURUA, O., GARIN, I., ALBERDI, A., SALSAMENDI, E., BAAGØE, H. &
AIHARTZA, J. 2013. Fishing long-fingered bats (Myotis capaccinii) prey
regularly upon exotic fish. *PLoS One,* 8**,** e80163.
ALBERDI, A. & GILBERT, M. T. P. 2019. A guide to the application of Hill numbers
to DNA‐based diversity analyses. *Molecular ecology resources,* 19**,** 804-817.
ALTENBACH, J. S. 1989. Prey capture by the fishing bats Noctilio leporinus and
Myotis vivesi. *Journal of Mammalogy,* 70**,** 421-424.
ANDERSEN, N. M. & FOSTER, W. 1992. Sea skaters of India, Sri Lanka, and the
Maldives, with a new species and a revised key to Indian Ocean species of
Halobates and Asclepios (Hemiptera, Gerridae). *Journal of Natural History,*
26**,** 533-553.
BECK, A. 1995. Fecal analyses of European bat species. *Myotis,* 32**,** 109-119.
BROOKE, A. P. 1994. Diet of the fishing bat, Noctilio leporinus (Chiroptera:
Noctilionidae). *Journal of Mammalogy,* 75**,** 212-218.
BYLEMANS, J., GLEESON, D. M., HARDY, C. M. & FURLAN, E. 2018. Toward
an ecoregion scale evaluation of eDNA metabarcoding primers: A case study
for the freshwater fish biodiversity of the Murray–Darling Basin (Australia).
Ecology and evolution, 8**,** 8697-8712.
CARPENTER, R. E. 1968. Salt and water metabolism in the marine fish-eating bat,
Pizonyx vivesi. *Comparative Biochemistry and Physiology,* 24**,** 951-964.
CHANG, Y., SONG, S., LI, A., ZHANG, Y., LI, Z., XIAO, Y., JIANG, T., FENG, J. &
LIN, A. 2019. The roles of morphological traits, resource variation and
resource partitioning associated with the dietary niche expansion in the fish‐
eating bat Myotis pilosus. *Molecular Ecology,* 28**,** 2944-2954.
DALZELL, P. & WANKOWSKI, J. 1980. *The Biology, Population Dynamics, and*
Fisheries Dynamics of Exploited Stocks of Three Baitfish Species, Stolephorus
Heterolobus, S. Devisi, and Spratelloides Gracilis, in Ysabel Passage, New
Ireland Province, Papua New Guinea, Department of Primary Industry.
DE MOOR, F. & IVANOV, V. 2008. Global diversity of caddisflies (Trichoptera:
Insecta) in freshwater. *Freshwater animal diversity assessment***,** 393-407.

DE SOUSA, L. L., SILVA, S. M. & XAVIER, R. 2019. DNA metabarcoding in diet
studies: Unveiling ecological aspects in aquatic and terrestrial ecosystems.
Environmental DNA, 1**,** 199-214.
DIGO, E. O., ABAD, K. L. M., GUINO-O, I. J. B., SAMILLANO, L. K. C.,
EDUQUE JR, R. M., TORRES, M. A. J. & REQUIERON, E. A. 2015.
Application of geometric morphometrics in the body shapes of flying fish
(Parexocoetus brachypterus) in Maitum, Sarangani Province. *Aquaculture,*
Aquarium, Conservation & Legislation, 8**,** 1027-1034.
DUDGEON, D. & CORLETT, R. 2004. *The ecology and biodiversity of Hong Kong,*
Joint Publishing (Hong Kong) Ltd.
EDGAR, R. C. 2010. Search and clustering orders of magnitude faster than BLAST.
Bioinformatics, 26**,** 2460-2461.
EDGAR, R. C. 2016. SINTAX: a simple non-Bayesian taxonomy classifier for 16S
and ITS sequences. *biorxiv***,** 074161.
FISH, F. E., BLOOD, B. R. & CLARK, B. D. 1991. Hydrodynamics of the feet of
fish-catching bats: influence of the water surface on drag and morphological
design. *Journal of Experimental Zoology,* 258**,** 164-173.
FROESE, R. & PAULY, D. 2024. FishBase. World Wide Web electronic publication.
February 2024 ed.
GLÖCKNER, F. O., YILMAZ, P., QUAST, C., GERKEN, J., BECCATI, A.,
CIUPRINA, A., BRUNS, G., YARZA, P., PEPLIES, J. & WESTRAM, R.
2017. 25 years of serving the community with ribosomal RNA gene reference
databases and tools. *Journal of biotechnology,* 261**,** 169-176.
GOITI, U., AIHARTZA, J., GUIU, M., SALSAMENDI, E., ALMENAR, D.,
NAPAL, M. & GARIN, I. 2011. Geoffroy's bat, Myotis emarginatus, preys
preferentially on spiders in multistratified dense habitats: a study of foraging
bats in the Mediterranean. *Folia Zoologica,* 60**,** 17-24.
GWYNNE, D. T. & EDWARDS, E. 1986. Ultrasound production by genital
stridulation in Syntonarcha iriastis (Lepidoptera: Pyralidae): long-distance
signalling by male moths? *Zoological Journal of the Linnean Society,* 88**,** 363-
376.
HADLEY, W. 2016. *Ggplot2: Elegrant graphics for data analysis,* Springer.
HARVEY, M. S., AUSTIN, A. D. & ADAMS, M. 2007. The systematics and biology
of the spider genus Nephila (Araneae: Nephilidae) in the Australasian region.
Invertebrate Systematics, 21**,** 407-451.
HSIEH, T., MA, K. & CHAO, A. 2016. iNEXT: an R package for rarefaction and
extrapolation of species diversity (H ill numbers). *Methods in ecology and*
evolution, 7**,** 1451-1456.
HUANG, P.-Y., POON, E. S. K., WONG, A. T. C., SO, I. W. Y., SUNG, Y.-H. & SIN,
S. Y. W. 2021. DNA metabarcoding reveals the dietary composition in the
endangered black-faced spoonbill. *Scientific Reports,* 11**,** 18773.
HUANG, P. Y., POON, E. S. K., CHAN, L. Y., CHAN, D. T. C., HUYNH, S., SO, I.
W. Y., SUNG, Y. H. & SIN, S. Y. W. 2022. Dietary diversity of multiple

shorebird species in an Asian subtropical wetland unveiled by DNA metabarcoding. *Environmental DNA,* 4**,** 1381-1396.

IUCN 2023. The IUCN Red List of Threatened Species. Version 2023-1.

JIANG, F., ZHENXIN, L., MIN, C., YING, L., XICHEN, Z., JIANG, Z. & SHUYI, Z. 2003. Morphological features, echolocation calls and foraging strategy in the trawling piscivorous bat: Rickett's big-footed bat Myotis ricketti. *Acta Ecologica Sinica,* 23**,** 1712-1718.

JIANG, T. L., FENG, J., CSORBA, G. & BATES, P. 2019. Myotis pilosus. The IUCN Red List of Threatened Species 2019: e.T14193A22062554.

JIANG, Z., JIANG, J., WANG, Y., ZHANG, E., ZHANG, Y., LI, L., XIE, F., CAI, B., CAO, L. & ZHENG, G. 2016. Red list of China's vertebrates. *Biodiversity Science,* 24**,** 500.

KAHLKE, T. & RALPH, P. J. 2019. BASTA–Taxonomic classification of sequences and sequence bins using last common ancestor estimations. *Methods in Ecology and Evolution,* 10**,** 100-103.

KAI, K. H. & CORLETT, R. T. 2002. Seasonality of forest invertebrates in Hong Kong, south China. *Journal of tropical Ecology,* 18**,** 637-644.

KAUPAS, L. A. & BARCLAY, R. M. 2018. Temperature-dependent consumption of spiders by little brown bats (Myotis lucifugus), but not northern long-eared bats (Myotis septentrionalis), in northern Canada. *Canadian Journal of Zoology,* 96**,** 261-268.

LEE, B. Y., WAKABAYASHI, K., SIN, S. Y. W., OHTSUKA, S. & TSANG, L. M. 2021. DNA Metabarcoding Revealed Interspecific Dietary Difference and Prey Selectivity in Juvenile Horseshoe Crabs Carcinoscorpius rotundicauda and Tachypleus tridentatus From Hong Kong. *Frontiers in Marine Science,* 8**,** 752806.

LERAY, M., KNOWLTON, N. & MACHIDA, R. J. 2022. MIDORI2: A collection of quality controlled, preformatted, and regularly updated reference databases for taxonomic assignment of eukaryotic mitochondrial sequences. *Environmental DNA,* 4**,** 894-907.

LEVIN, E., BARNEA, A., YOVEL, Y. & YOM-TOV, Y. 2006. Have introduced fish initiated piscivory among the long-fingered bat? *Mammalian Biology,* 71**,** 139-143.

LEWIS, J. B. 1961. The growth, breeding cycle and food of the flyingfish Parexocoetus brachypterus hillianus (Gosse). *Bulletin of Marine Science,* 11**,** 258-266.

LIU, Y., HE, G., XU, H., HAN, X., JONES, G., ROSSITER, S. J. & ZHANG, S. 2014. Adaptive functional diversification of lysozyme in insectivorous bats. *Molecular biology and evolution,* 31**,** 2829-2835.

MA, J., JONES, G., ZHANG, S., SHEN, J., METZNER, W., ZHANG, L. & LIANG, B. 2003. Dietary analysis confirms that Rickett's big-footed bat (Myotis ricketti) is a piscivore. *Journal of Zoology,* 261**,** 245-248.

MA, J., LIANG, B., ZHANG, S. & METZNER, W. 2008. Dietary composition and echolocation call design of three sympatric insectivorous bat species from

China. *Ecological Research,* 23**,** 113-119.

MA, J., ZHANG, J., LIANG, B., ZHANG, L., ZHANG, S. & METZNER, W. 2006.
Dietary characteristics of Myotis ricketti in Beijing, north China. *Journal of*
Mammalogy, 87**,** 339-344.

MARTIN, M. 2011. Cutadapt removes adapter sequences from high-throughput
sequencing reads. *EMBnet. journal,* 17**,** 10-12.

MASSEY, A. L., ROFFLER, G. H., VERMEUL, T., ALLEN, J. M. & LEVI, T. 2021.
Comparison of mechanical sorting and DNA metabarcoding for diet analysis
with fresh and degraded wolf scats. *Ecosphere,* 12**,** e03557.

MAUCIERI, D. & BARCLAY, R. 2021. Consumption of spiders by the little brown
bat (Myotis lucifugus) and the long-eared myotis (Myotis evotis) in the Rocky
Mountains of Alberta, Canada. *Canadian Journal of Zoology,* 99**,** 221-226.

MCINNES, J. C., ALDERMAN, R., DEAGLE, B. E., LEA, M. A., RAYMOND, B.
& JARMAN, S. N. 2017. Optimised scat collection protocols for dietary DNA
metabarcoding in vertebrates. *Methods in Ecology and Evolution,* 8**,** 192-202.

MCMURDIE, P. J. & HOLMES, S. 2019. Package 'phyloseq'. Tech. Rep. Stanford
Univ, Stanford, CA, USA.

MONTERROSO, P., GODINHO, R., OLIVEIRA, T., FERRERAS, P., KELLY, M. J.,
MORIN, D. J., WAITS, L. P., ALVES, P. C. & MILLS, L. S. 2019. Feeding
ecological knowledge: the underutilised power of faecal DNA approaches for
carnivore diet analysis. *Mammal Review,* 49**,** 97-112.

OKSANEN, J., SIMPSON, G., BLANCHET, F., KINDT, R., LEGENDRE, P.,

MINCHIN, P., O′ HARA, R., SOLYMOS, P., STEVENS, H. & SZOECS, E.

2022. Vegan Community Ecology Package Version 2.6—2 April 2022. The
Comprehensive R Archive Network.

OSPINA-GARCÉS, S. M., DE LUNA, E., HERRERA M, L. G. & FLORES-
MARTÍNEZ, J. J. 2016. Cranial shape and diet variation in Myotis species
(Chiroptera: Vespertilionidae): testing the relationship between form and
function. *Acta Chiropterologica,* 18**,** 163-180.

OTÁLORA-ARDILA, A., HERRERA M, L. G., FLORES-MARTÍNEZ, J. J. &
VOIGT, C. C. 2013. Marine and terrestrial food sources in the diet of the fish-
eating myotis (Myotis vivesi). *Journal of Mammalogy,* 94**,** 1102-1110.

PEDERSEN, T. L., PEDERSEN, M., LAZYDATA, T., RCPP, I. & RCPP, L. 2017.
Package 'ggraph'. *Retrieved January,* 1**,** 2018.

POOLPRASERT, P., SENARAT, S., KETTRATAD, J., KANEKO, G.,
MONGKOLCHAICHANA, E., CHAROENPHON, N. & THAOCHAN, N.
2022. Comprehensive Structure of the Female Marine Water-Strider Asclepios
annandalei Distant, 1915 from Pranburi River Estuary, Thailand: New
Information for the Genus Asclepios. *Tropical life sciences research,* 33**,** 47.

POON, E. S. K., CHEN, G., TSANG, H. Y., SHEK, C. T., TSUI, W. C., ZHAO, H.,
GUÉNARD, B. & SIN, S. Y. W. 2023. Species richness of bat flies and their
associations with host bats in a subtropical East Asian region. *Parasites &*
Vectors, 16**,** 37.

PUIG-MONTSERRAT, X., FLAQUER, C., GÓMEZ-AGUILERA, N., BURGAS,

A., MAS, M., TUNEU, C., MARQUÈS, E. & LÓPEZ-BAUCELLS, A. 2020.

Bats actively prey on mosquitoes and other deleterious insects in rice paddies:
Potential impact on human health and agriculture. *Pest Management Science,*
76**,** 3759-3769.

RAMÍREZ-FRÁNCEL, L. A., GARCÍA-HERRERA, L. V., LOSADA-PRADO, S.,

REINOSO-FLÓREZ, G., SÁNCHEZ-HERNÁNDEZ, A., ESTRADA-
VILLEGAS, S., LIM, B. K. & GUEVARA, G. 2022. Bats and their vital
ecosystem services: a global review. *Integrative Zoology,* 17**,** 2-23.

RAZGOUR, O., CLARE, E. L., ZEALE, M. R., HANMER, J., SCHNELL, I. B.,

RASMUSSEN, M., GILBERT, T. P. & JONES, G. 2011. High-throughput

sequencing offers insight into mechanisms of resource partitioning in cryptic
bat species. *Ecology and evolution,* 1**,** 556-570.

RISKIN, D. K. & CARTER, G. G. 2023. The evolution of sanguivory in vampire
bats: origins and convergences. *Canadian Journal of Zoology,* 101**,** 207-221.

ROGNES, T., FLOURI, T., NICHOLS, B., QUINCE, C. & MAHÉ, F. 2016.
VSEARCH: a versatile open source tool for metagenomics. *PeerJ,* 4**,** e2584.

SADOVY, Y. & CORNISH, A. S. 2000. *Reef fishes of Hong Kong*, Hong Kong
University Press.

SHEK, C. 2004. Bats of Hong Kong: An introduction of Hong Kong bats with an
illustrative identification key. *Hong Kong Biodiversity,* 7**,** 1-9.

SHEK, C. T. 2006. *Field guide to the terrestrial mammals of Hong Kong*, Cosmos
Books.

SKIBINSKI, H. 2005. *Light positively affects foraging success of the night feeding*
zooplanktivorous fish hardyhead silverside (Atherinomorus lacunosus). BSc
thesis, Uppsala University.

STEINBERGER, A. 2020. kruskal.pretty: Automates kruskal.test execution on
simper_pretty.R output. Suen Lab, University of Wisconsin-Madison.

STEINBERGER, A., DILL-MCFARLAND, K. & COX, M. 2016. simper.pretty:
automates simper exectution for comparisons of interest. Suen Lab, University
of Wisconsin-Madison.

STEVENS, P. W., BENNETT, C. K. & BERG, J. J. 2003. Flyingfish spawning
(Parexocoetus brachypterus) in the northeastern Gulf of Mexico.
Environmental biology of fishes, 67**,** 71-76.

SVENDSEN, A.-S. L., NIELSEN, L. B., SCHMIDT, J. B., BRUHN, D.,
ANDERSEN, L. H. & PERTOLDI, C. 2023. eDNA Metabarcoding-and
Microscopic Analysis for Diet Determination in Waterfowl, a Comparative
Study in Vejlerne, Denmark. *Biology,* 12**,** 1272.

SWIFT, S. & RACEY, P. 2002. Gleaning as a foraging strategy in Natterer's bat
Myotis nattereri. *Behavioral Ecology and Sociobiology,* 52**,** 408-416.

TAYLOR, M. 2019. *Bats: an illustrated guide to all species*, Ivy Press.

TSANG, A. H. F. & DUDGEON, D. 2021a. Do exotic poeciliids affect the
distribution or trophic niche of native fishes? Absence of evidence from Hong
Kong streams. *Freshwater Biology,* 66**,** 1751-1764.

TSANG, A. H. F. & DUDGEON, D. 2021b. A manipulative field experiment reveals
the ecological effects of invasive mosquitofish (Gambusia affinis) in a tropical
wetland. *Freshwater Biology,* 66**,** 869-883.

VAMOS, E. E., ELBRECHT, V. & LEESE, F. 2017. Short COI markers for
freshwater macroinvertebrate metabarcoding. PeerJ Preprints.

WALKER, F. M., WILLIAMSON, C. H., SANCHEZ, D. E., SOBEK, C. J. &
CHAMBERS, C. L. 2016. Species from feces: order-wide identification of
Chiroptera from guano and other non-invasive genetic samples. *Plos one,* 11**,**
e0162342.

WANG, Q., FENG, J., WU, H. & JIANG, T. 2024. Insectivorous bats provide more
pest suppression services than disservices-a case study in China. *Biological*
Control, 188**,** 105435.

WILSON, D. E. & MITTERMEIER , R. A. 2019. *Handbook of the Mammals of the*
World, Lynx Edicions.

WINGETT, S. W. & ANDREWS, S. 2018. FastQ Screen: A tool for multi-genome
mapping and quality control. *F1000Research,* 7.

ZHANG, J., DING, Q. & HUANG, J. 2016. spaa: Species association analysis. *R*
package version 0.2, 2**,** 33.

ZHU, T., SATO, Y., SADO, T., MIYA, M. & IWASAKI, W. 2023. MitoFish,
MitoAnnotator, and MiFish pipeline: updates in 10 years. *Molecular Biology*
and Evolution, 40**,** msad035.

Figure 1

975
976

977   Figure 2

978
979

Figure 3

983   Figure 4

Figure 5

989   Figure 6

Supplementary Materials

Foraging niche partitioning of three *Myotis* bat species and marine fish consumption by *Myotis pilosus* in a subtropical East Asian region

Supplementary materials and methods

Supplementary figures S1-S6

Supplementary Materials and Methods

Preparation of DNA metabarcoding libraries through 2-step PCRs

For 18S rDNA libraries, 1st step PCRs were carried out in 30 μl reactions containing 6 μl of 5X GoTaq Flexi Buffer, 0.6 μl of 10 mM dNTP Mix, 3.6 μl of 25 mM MgCl$_2$, 0.15 μl of 5 U/μl GoTaq G2 Flexi DNA Polymerase, 1 μl of extracted DNA, 0.6 μl of each assigned 10 μM forward and reverse primer uniquely tagged with heterogeneity spacer (Cruaud et al. 2017), 6 μl of 10 % DMSO (Sigma), 0.15 μl of 20 mg/ml BSA (NEB) and ultrapure water. Thermal cycling condition was 95 °C for 2 min; 25 cycles of 95 °C for 30 sec, 50 °C for 30 sec and 72 °C for 30 sec; and final extension at 72 °C for 5 min.

For COI libraries, the 1st step PCRs were carried out in 6 μl of 5X GoTaq Flexi Buffer, 0.6 μl of 10 mM dNTP Mix, 3.6 μl of 25 mM MgCl$_2$, 0.15 μl of 5 U/μl GoTaq G2 Flexi DNA Polymerase, 1 μl of extracted DNA, 1.5 μl of each assigned 10 μM forward and reverse primer uniquely tagged with heterogeneity spacer (Cruaud et al. 2017), 6 μl of 10 % DMSO (Sigma), 0.15 μl of 20 mg/ml BSA (NEB) and ultrapure water. Thermal cycling condition was 95 °C for 2 min; 25 cycles of 95 °C for 30 sec, 48 °C for 30 sec and 72 °C for 30 sec; and final extension at 72 °C for 5 min. Target PCR products generated from COI primers were gel extracted to eliminate non-targets.

For 12S rDNA libraries, 1st step PCRs were carried out in 30 μl reactions containing 6 μl of 5X GoTaq Flexi Buffer, 0.6 μl of 10 mM dNTP Mix, 3.6 μl of 25 mM MgCl$_2$, 0.15 μl of 5 U/μl GoTaq G2 Flexi DNA Polymerase, 1 μl of extracted DNA, 0.6 μl of each assigned 10 μM forward and reverse primer uniquely tagged with heterogeneity spacer (Cruaud et al. 2017), 6 μl of 10 % DMSO (Sigma), 0.15 μl of 20 mg/ml BSA (NEB) and ultrapure water. Each of the 1st step PCR products was eluted in 20 μl elution buffer and used as the templates for the 2nd step PCRs.

The 2nd step PCRs of both markers were performed with the same conditions: 2nd step PCRs were performed in 45 μl reactions comprising 9 μl of 5X GoTaq Flexi Buffer, 0.9 μl of 10 mM dNTP Mix, 5.4 μl of 25 mM MgCl$_2$, 0.225 μl of 5 U/μl GoTaq G2 Flexi DNA Polymerase, 20 μl of 1st step PCR products, 0.9 μl of each assigned 10 μM forward and reverse index primers, 4.5 μl of 10 % DMSO (Sigma) and ultrapure water. PCR condition was 95 °C for 2 min; 15 cycles of 95 °C for 30 sec, 55 °C for 30 sec, and 72 °C for 30 sec; and final extension at 72 °C for 5 min. We used the PureLink PCR Purification Kit (Invitrogen, Carlsbad, CA) to clean up all PCR products. The band sizes of all 1st and 2nd PCR products were checked on 1.5% agarose gels by gel electrophoresis.

Reference

Cruaud P, Rasplus J-Y, Rodriguez LJ, Cruaud A (2017) High-throughput sequencing of multiple amplicons for barcoding and integrative taxonomy Scientific reports 7:41948

Supplementary Figures

Figure S1.

The fecal samples of *Myotis* bats were collected from seven sites in Hong Kong. The small map at the upper left corner represents Hong Kong at the south coast of China, with the area enclosed in a red box magnified for ease of viewing. The pie charts in the magnified map represent the relative quantity (not to scale) of samples collected at each location, with larger pie charts indicating a larger sample size. Sites in close proximity were grouped as sampling locations for analyses (see Table S1). The locations include LMH (indicated in black), TLC (indicated in dark blue, including TLC01 and TLC02), and SK (indicated in dark red, including PTC, SK01, SK02, and SK03). LMH, Lin Ma Hang. TLC, Tai Lam Chung. PTC, Pak Tam Chung Camp. SK, Sai Kung.

Figure S2.
Rarefaction curves based on Chao2 species richness. The Chao2 species richness was estimated from hill numbers of diversity order q=0, which were evaluated based on ASVs (left panel) or taxa (b) in each *Myotis* species. Dietary ASVs or taxa were identified by (a) 18S rDNA, (b) COI, and (c) 12S rDNA marker gene. The shading area represent the 95% confidence intervals, which were estimated using a bootstrap method based on 100 replicates.

18S rDNA

Figure S3.
Dietary composition of three *Myotis* bat species based on taxa detected in fecal samples using the 18S rDNA marker. The weighted percentages of occurrence of each taxon were visualized at both (a) bat individual level (n=144) and (b-d) bat species level for (b) *M. chinensis* (n=40), (c) *M. horsfieldii* (n=49), and (d) *M. pilosus* (n=55). Prey in classes with read abundance lower than 0.1% of all taxa detected were grouped into "Others". Only taxa with abundance higher than 1% in relative read abundance are shown. F, Family. O, Order.

Figure S4.
Dietary composition of *Myotis pilosus* based on taxa detected in fecal samples using the 12S rDNA marker. The weighted percentages of occurrence of each taxon were visualized at both (a) individual (n=48) and (b) species levels. Individual bars were grouped by sampling locations and seasons. Prey in classes with read abundance lower than 0.1% of all taxa detected were grouped into "Others". Only taxa with abundance higher than 1% in relative read abundance are shown. SK, Sai Kung (including Pak Tam Chung); TLC, Tai Lam Chung; LMH, Lin Ma Hang. Refer to Fig. S1 for locations.

Figure S5.

Dietary composition of three *Myotis* bat species based on taxa detected in fecal samples using the COI marker. The weighted percentages of occurrence of each taxon were visualized at both (a) individual level (n=102) and (b-d) species level for (b) *M. chinensis* (n=34), (c) *M. horsfieldii* (n=44), and (d) *M. pilosus* (n=22). Prey in classes with read abundance lower than 0.1% of all taxa detected were grouped into "Others". Only taxa with abundance higher than 1% in relative read abundance are shown. G, Genus. F, Family.

(a)

Myotis chinensis
Myotis horsfieldii
Myotis pilosus

Interspecific overlap
Intraspecific overlap

(b)

Figure S6.
Dietary overlap of three sympatric *Myotis* bat species. Each colored circle represents a single bat sample. Dietary overlap between individual pairs was calculated based on their diet composition determined by (a) 18S [*M. chinensis* (n=40), *M. horsfieldii* (n=49), and *M. pilosus* (n=55)] and (b) COI [*M. chinensis* (n=34), *M. horsfieldii* (n=44), and *M. pilosus* (n=22)] markers. The thickness of the line connecting circles represents the extent of dietary overlap between individuals.