Acute Activation of Genes Through Transcriptional Condensates Impact Non-target Genes in a Chromatin Domain

Darshika Bohra[1,#] , Zubairul Islam[2,3,#], Sundarraj Nidharshan[2,4], Aprotim Mazumder[1]*, Dimple Notani[2]*

[1]Tata Institute of Fundamental Research Hyderabad, Hyderabad-500046, Telangana, India.

[2]National Centre for Biological Sciences, Tata Institute for Fundamental Research, Bangalore-560065, Karnataka, India.

[3]Sastra Deemed University, Thanjavur - 613401, Tamil Nadu, India.

[4]School of Biotechnology, Amrita Vishwa Vidyapeetham, Kollam - 690525, Kerala, India

#These authors contributed equally to this work

*Correspondence: aprotim@tifrh.res.in, dnotani@ncbs.res.in,

Abstract

Transcription activation of genes by estrogens is driven by enhancers, which are often located within the same Topologically Associating Domain (TAD) as non-targeted promoters. We investigated how acute enhancer-driven activation affects neighbouring non-target genes within the same TAD. Using single-molecule RNA FISH (smFISH), we tracked the transcription of *TFF1* (enhancer-targeted) and *TFF3* (non-targeted) during estrogen stimulation. We observed mutually exclusive expression patterns: *TFF1* expression peaked at 1 hour, while *TFF3* reached its peak at 3 hours, after *TFF1*'s activation had diminished. Chromatin looping data indicated that the enhancer loops with *TFF1* but not *TFF3*, suggesting that *TFF3* upregulation is not due to direct enhancer-promoter interactions. CRISPR deletion of the *TFF1* enhancer and 1,6-hexanediol (HD) exposure revealed that the *TFF1* enhancer:promoter undergo Liquid-Liquid Phase Separation (LLPS), which sequesters the transcriptional machinery and inhibits *TFF3* expression. As estrogen signalling wanes or LLPS is disrupted, *TFF1* expression declines while *TFF3* expression increases. Our findings reveal that enhancer-driven activation can indirectly influence neighbouring genes, highlighting a dynamic shift in gene expression as signalling progresses.

Introduction

Acute transcriptional activation of ligand-induced genes drive downstream signalling response. Similar to development-specific genes, signalling induced genes are also driven by enhancers (Hah et al., 2013; Li et al., 2013; Liu et al., 2014; Uyehara & Apostolou, 2023). Upon binding with ligand induced transcription factors (TFs), these enhancers loop with their target promoters mostly, in the same TAD (Buecker et al., 2014; Bulger & Groudine, 1999; Chepelev et al., 2012; Furlong & Levine, 2018; Oh et al., 2021; Panigrahi & O'Malley, 2021; Ptashne, 1986; Sanyal et al., 2012; Yan et al., 2018). Enhancer: promoter pairing is thought to be specific and forms the basis of noise-free gene activation of a subset of genes crucial for signalling response (Bojcsuk et al., 2017; Chen et al., 2018; Friedman et al., 2024; Galouzis & Furlong, 2022; Zabidi et al., 2015). Due to such specificity, gene transcription occurs in waves of early and late responsive genes (Fowler et al., 2011; Yamamoto & Alberts, 1976) Often, the protein factors translated from early genes, regulate the expression of late responsive genes (Dixon et al., 1996; Freter et al., 1996; Herschman, 1991; Williams et al., 1999; Winkles, 1997; Winston & Pledger, 1993). However, it is not known if the genes that are activated early in signalling time course are spatially related to late activating genes. Further, if spatial proximity of any gene to an early gene is enough to cause the temporal differential transcription due to sequestration of the transcriptional machinery from late gene to early gene is poorly understood. Transcription factors (TFs), after binding to cognate DNA motif, recruit RNA polymerase machinery and co-activators for gene activation (Levine & Tjian, 2003; Liu et al., 2014). These machinery are limited in supply, and may in effect be sequestered from other genomic regions (Koşar & Erbaş, 2022). Further, acute activation of genes is linked with phase-separation of TFs, polymerases, mediators and other co-factors/activators, most of which harbour low complexity regions to promote weak protein-protein interactions driving formation of TF-condensates (Boehning et al., 2018; Boija et al., 2018; Cai et al., 2019; Cho et al., 2018; Chong et al., 2018, 2022; Mann & Notani, 2023; Sabari et al., 2018; Shrinivas et al., 2019; Stortz et al., 2020, 2024). Though the

precise stoichiometry of protein molecules in these condensates is unknown, such structures do
involve multiple molecules of each transcriptional protein. The formation of such phase-
separated compartments potentially can act as a sink for transcriptional machinery depriving
neighbouring promoters and enhancers that are not part of the compartment. Such sequestration
would cause indirect suppression of these genes that are spatially proximal in the same TAD but
not the part of the same condensate.
99
In order to investigate the effect of an enhancer: promoter pair on a neighbouring gene within the
same TAD, we looked at the paradigmatic model of estrogen signalling in mammary epithelial
cells namely MCF7 (Levenson & Jordan, 1997; Masiakowski et al., 1982). We selected E2-
induced E-P pair *TFF1* and its neighbouring gene *TFF3* on Chr21 located at a distance of 43kb
from each other, as discussed below (Chinery et al., 1996). We chose to answer these questions
using an integrated approach consisting of single molecule RNA FISH (smFISH), genome-wide
conformation capture and sequencing (4C-seq), and perturbation of cis-acting regulatory
elements by CRISPR. E2-signalling can be induced within minutes by treating the cells with
estradiol (E2) which causes activation and repression of several genes across the genome. The
peak of E2-mediated signalling occurs at 40 minutes post-induction and starts to decay by 160
minutes (Hah et al., 2011). This occurs via the binding of estrogen receptor-alpha (ERα) on the
enhancers (Li et al., 2013). *TFF1* is found in a topologically associated domain (TAD) along
with a few other genes including *TFF2*, *TFF3*, *TMPRSS3*, and *UBASH3A* (Oh et al., 2021;
Quintin et al., 2014; Rao et al., 2014). *TFF1* and *TFF3* are located at 10kb and 60kb from
enhancer respectively (Fig. 1A) and this locus has been useful in answering the questions about
promoter-enhancer interactions and multi-gene regulation (Oh et al., 2021; Quintin et al., 2014;).
116
Using this system, we identified that *TFF1* and *TFF3* genes located within the same TAD show
distinct and opposing expression profiles over the course of E2-mediated signalling. In
agreement with this, the enhancer also showed increased lopping interaction with the cognate
gene promoter during peak expression at 1h of signalling. We also identified the role of TF-
mediated LLPS on the regulation of these two genes during signalling. Hence, we propose that
ligand dependent condensation of ERα can support the expression of the cognate gene in concert
with the enhancer while negatively affecting the expression of a neighbouring gene.
124
125   **Results**
126
127   **ERα binding and ligand-induced gene expression of *TFF1* and *TFF3* change over the**
128   **course of estrogen signaling**
129
130   In order to understand how acute activation of one gene in a TAD affects the transcription of a
131   proximate gene, we chose to study *TFF1* and its neighbouring gene *TFF3* at a 43kb distance
132   within same TAD. *TFF1* expression is linked with an enhancer located 10 kb downstream (Li et
133   al., 2013; Saravanan et al., 2020; Oh et al., 2021) and the distance between *TFF1* enhancer and
134   *TFF3* gene is 53kb (Fig. 1A).
135
17-β-oestradiol exposure leads to ERα binding on regulatory regions leading to gene activation.
Gene transcription of E2-regulated genes was shown to peak at 1h and significantly reduces at 3h
due to the rapid degradation of ERα (Hah et al., 2011). The binding of ERα in the genome also

follows this temporal kinetics where it peaks at 1h and reduces at 3h post ligand stimulation (Hah
et al., 2011; Li et al., 2013; Liu et al., 2014). At the *TFF1* locus, the *TFF1* enhancer and to some
extent, its promoter, were bound by ERα even in the absence of E2 induction, while it increased
in strength at these regions and also at various other regions in the locus at 1h of ligand
stimulation (Saravanan et al., 2020). Notably, the binding at these regions substantially decreased
at 3h. On the other hand, ERα did not bind on *TFF3* promoter throughout the course of
signalling (Fig. 1A and Fig. S1B). We have previously shown that such estrogen-induced
clustered binding of ERα on *TFF1* region is associated with its acute activation post estrogen
stimulation (Saravanan et al., 2020). We then tested the level of their expression by nascent
RNA-seq (GRO-seq). We observed a dramatic transcriptional activation of *TFF1* at 1h which
reduced substantially at 3h. The expression level of *TFF3* was comparatively low and did not
fluctuate upon signalling (Fig. 1A). However, qRT-PCR on unspliced *TFF3* exhibited
upregulation at 1h and down-regulation at 3h. (Fig. 1B) While this broad trend is similar to *TFF1*
as shown by both GRO-seq and qRT-PCR, the relative changes for *TFF3* was smaller than
TFF1. We reasoned that these differences in *TFF3* could be because of its low baseline
expression (Clark et al., 2015; Conesa et al., 2016; Sha et al., 2015; Svensson et al., 2017;
Tarazona et al., 2011), and lower yet expression of unspliced transcripts. Therefore, in order to
test the relative induction of these genes and their co-regulation, we decided to perform single
cell measurements of RNA (Fig. S1A) that do not rely on PCR amplification (sc-RNA-seq) and
can provide absolute numbers of *TFF1* and *TFF3* transcripts in a given cell. Towards this, we
chose to employ smFISH (Femino et al., 1998; Haimovich & Gerst, 2018; Kwon, 2013; Raj et
al., 2008).
161
162 ***TFF1* and *TFF3* exhibit opposite trends during the E2 signalling time-course**
163
MCF7 cells are hypertriploid to hypotetraploid, and each of the three alleles within a nucleus can
behave differently, and smFISH is pre-eminently suited to capturing cell- and allele-specific
heterogeneity of expression even for low expressing genes, compared to bulk studies that
average over cell populations. Briefly, smFISH allows to visualize single RNA molecules in
fixed cells using multiple fluorescently labelled oligonucleotide probes targeted to the RNA of
interest and therefore, it can be used to image the transcription and the localization of multiple
gene transcripts at same and different time points after signalling (Femino et al., 1998; Raj et al.,
2008).
172
We designed the smFISH probes targeting the intronic region of *TFF1* and *TFF3* to measure
their nascent transcripts (referred to as InTFF1 and InTFF3 hereon) whereas the exonic probes
were used to primarily measure mature mRNA transcripts (referred to as ExTFF1 and ExTFF3
hereon). Intronic probes are particularly suited for investigating nascent transcriptional status at
the time of fixation, while probes against mature mRNA reflects on more steady-state levels of
functional mRNA due to finite mRNA lifetimes (Skinner et al., 2016). Additionally, the intronic
probes can help determine the localisation and number of alleles that are transcribing at a given
time thus allowing for additional interpretations regarding the expression of multiple genes
(Skinner et al., 2016).
182
Representative images from the smFISH experiment probing InTFF1 and InTFF3 are depicted
(Fig. 2A). The smaller foci represent individual intronic/nascent transcripts while the larger foci

represent the site of transcription (Raj et al., 2006; Zenklusen et al., 2008). Since MCF7 cells are
hypertriploid in nature, we expected to see 1-3 large foci per cell representing sites of
transcription. Additionally, we observed many individual transcripts labelled by intronic probes
(Fig. 2A, D) within the same nuclei (probes against mature mRNA are usually more
cytoplasmic). This is suggestive of the fact that the transcripts undergo non-co-transcriptional
splicing as the transcripts labelled by intronic probes are localised away from the site of
transcription. This is not surprising as several reports have shown that nascent transcript can
undergo splicing well after transcription (Coulon et al., 2014; Drexler et al., 2020; Khodor et al.,
2012). Indeed, a very recent study has shown that such post-transcriptional splicing occurs in a
small zone around genes including *TFF1*(Coté et al., 2023). Therefore, observation of nascent
transcripts away from the site of transcription as we see for *TFF1* and *TFF3* (Fig. 2A, D) is not
unexpected. We quantified the number of such transcripts per nuclei as a proxy for ongoing
transcription.
198
At 1h post E2 induction, *TFF1* mean transcript counts increased significantly compared to
uninduced, whereas, the increment was less pronounced for *TFF3* transcripts. In contrast, the
mean transcript counts for *TFF3* increased significantly at 3h post induction while *TFF1*
transcription showed a decrease at 3h compared to 1h (Fig. 2B). To check the transcriptional
status of *TFF1* and *TFF3* in the same cell, we plotted the transcript counts from individual cells
(Fig. 2C). The data suggested that the transcript counts for *TFF1* increased at 1h and cells that
showed high counts for *TFF1* were likely to have low counts for *TFF3*. But the RNA counts for
TFF1 decreased at 3h and increased for *TFF3*. Overall, it was evident that the transcriptional
profile for these two genes located in the same TAD were negatively correlated as they peaked at
different time points. To further confirm this, smFISH using probes targeting both the intronic
and exonic transcripts in the same experiment was conducted. Intronic probes represent active
transcription, while exonic probes show accumulated mature mRNA from past transcriptional
events even in the absence of active transcription (Skinner et al., 2016). Representative images
from the smFISH experiment probing InTFF1 and ExTFF1 are depicted (Fig. 2D). We observed
that the exonic transcript counts for both *TFF1* and *TFF3* increased at 1h compared to uninduced
(Fig. 2E-F). Strikingly, at 3h, the exonic transcripts for *TFF1* continued to increase even while
the intronic transcript counts reduced, though the fold increase (1.12 ±0.02) was less compared
to that between uninduced and 1h (1.63 ± 0.28), indicating a plateauing of steady state levels.
Exonic transcripts for *TFF3* at 3h remained comparable to 1h while the intronic counts increased
suggesting, *TFF1* transcription reduces at 3h whereas, *TFF3* expression increases. We reasoned
that the ratio of intronic transcript number (InTFF1) to exonic transcripts number (ExTFF)
should represent the status of transcription as an increase in the number of intronic transcript due
to expression would result in a higher ratio while also taking into consideration the number of
mature transcripts. As expected, the ratio of intronic transcripts to exonic transcripts also showed
that transcription is active at 1h for *TFF1* as the ratio is higher compared to uninduced and 3h
(Fig. 2G and Fig. S2A). Contrastingly, the ratio was highest at 3h for *TFF3* suggesting that
active transcription takes place much after the peak of E2-mediated signalling and maximal
TFF1 expression (Fig. 2H and Fig. S2B).
227
228 **TFF1 enhancer does not change target promoters during signalling time-course**
229

In order to identify the molecular players behind differential expression of these two genes that
are in the same TAD, we asked if the ERα-bound enhancer downstream to *TFF1* gene, loops
with *TFF1* at 1h, and with *TFF3* at 3h. This enhancer acutely activates *TFF1* at 1h post estrogen
stimulation (Saravanan et al., 2020; Ho et al., 2021). We interrogated the looping using enhancer
as a viewpoint by 4C-seq in uninduced, 1 and 3h post E2 stimulation. We observed robust
interactions between enhancer and *TFF1* promoter at 1h post induction which reduced at 3h. On
the other hand, its interaction with *TFF3* promoter exhibited very low counts in uninduced as
well as E2-induced conditions at both time points and in both replicates (Fig. 3A, Fig. S3A). This
suggests that the looping of enhancer was potentially inducing the expression of *TFF1* gene at 1h
and loss of interactions at 3h resulted in weak *TFF1* transcription. However, lack of interactions
between enhancer and *TFF3* did not explain the gain of *TFF3* expression at 3h post ligand
stimulation.

This could mean that enhancer interaction is critical for *TFF1* expression, but is less important
for *TFF3* expression. Thus, *TFF1* expression should be affected more severely upon deletion of
the enhancer than *TFF3*. To test this, we investigated the expression of *TFF1* and *TFF3* in MCF-
7 cells where the enhancer downstream to *TFF1* was homozygously deleted using CRISPR-Cas9
(referred to as the Δ*TFF1*e from hereon) (Saravanan et al., 2021). Using smFISH, we quantified
the intronic transcripts for *TFF1* and *TFF3* in the Δ*TFF1*e cells, compared to WT cells. We
observed that the mean number of *TFF1* transcripts was reduced drastically in the Δ*TFF1*e
compared to the WT (Fig. 3C). The reduction was far more substantial (49.23 ±2.6 at 1hr and
40.75 ±7.8 at 3h) for *TFF1* than *TFF3* (6.03 ±1.8 at 1hr and 7.6 ±0.7 at 3h) (Fig. 3D). The
absolute transcript counts for *TFF1* and *TFF3* in the Δ*TFF1*e cells have also been shown for
clearer visualization (Fig. 3E), as these are obscured when compared to WT. To get a sense of
fold changes, we took a ratio of mean transcript counts in WT cells to Δ*TFF1*e cells at each time
point. We observed that the drop in gene transcripts between WT and Δ*TFF1*e were several folds
higher for *TFF1* compared to *TFF3* (Fig. 3F). This suggests that enhancer deletion has a more
robust impact on the transcription of *TFF1* compared to *TFF3*. Nonetheless, *TFF3* was also
affected even though, it does not loop with the enhancer. This is in accordance with the 4C-seq
data where we observed prominent looping between the enhancer and *TFF1* but less so with
TFF3. These results suggest that the enhancer plays a more important role in the expression of
the primary gene while it has less impact on a gene located more distally but within the same
TAD during course of signalling.

Levels of ERα in the nucleus dictate the extent of *TFF1* and *TFF3* inductions

After ruling out enhancer looping as the determinant of differential expression, we looked for
other candidates that could regulate the differential gene expression. As discussed above,
globally, E2-mediated gene expression is known to peak at 1h after stimulation and drop
significantly by 3h (Hah et al., 2011). Upon ligand stimulation, ERα translocates into nucleus,
increasing mean ERα intensity in the nucleus which is high at 1h and then decreases significantly
by 3h due to degradation. These changes in intensities have been captured using
immunofluorescence for ERα (Saravanan et al., 2020). We tested if the intensities of ERα in
individual cells correlate with the expression of *TFF1* and *TFF3* at 1h and 3h of E2 signalling.
Towards this, we combined smFISH with immunofluorescence for ERα. To improve contrast for
the ERα signal, we additionally performed a chromatin retention assay to get rid of any

chromatin unbound ERα (Fig. 4A). The representative images show that the cells with very high
levels of nuclear ERα (blue circle) exhibited low counts of both *TFF1* and *TFF3* while the cells
with medium levels of ERα (red circle) possessed higher *TFF1* than *TFF3*. Similarly, the cells
with the lower levels of ERα (grey circle) showed higher *TFF3* expression as compared to *TFF1*
(Fig. 4A). Histograms depicting the ERα mean intensities across individual cells, showed that the
nuclear level of ERα increases post 1h of induction and then goes down at 3h (Fig. 4B, C),
similar to *TFF1* expression (Fig. 4D).
283
To further corroborate this, we parsed the ERα population cells into 3 categories namely low,
medium, and high within the cells imaged at 1 and 3h. We plotted the mean counts of *TFF1* in
each of these bins at different time points and observed that the mean count was higher in the
mid-category (Fig. 4D). While for *TFF3*, the mean count was significantly higher in the low bin
(Fig. 4E). The transcript counts for *TFF1* and *TFF3* against ERα intensities on a cell-by-cell
basis (Fig. 4F), also showed this feature where very high levels of ERα in the nucleus were not
conducive to the expression of either gene (Fig. 4F). To test if ERα had a causal role in intensity
based expression of *TFF1* and *TFF3*, we increased the levels of ERα by over expression of ERα-
GFP (Fig. S4A,C). The representative images from smFISH experiment in cells overexpressing
ERα-GFP confirmed that *TFF1* and *TFF3* were downregulated in transfected cells while these
genes were not perturbed in non-transfected cells in neighbourhood (Fig. S4A, C). Meanwhile,
the cells overexpressing ERα-GFP do not show any impairment in the expression of GAPDH
(housekeeping gene) (Fig. S4A,B). As another control, we transfected the cells with EGFP-C1
(same backbone as ERα-GFP construct) and observed no effect on *TFF1* or *TFF3* (Fig. S4D,E).
The data suggest that loss of *TFF1* and *TFF3* expression upon ERα overexpression was not a
general effect of transfection stress, but rather specific to ERα overexpression. These results
indicate that high nuclear levels of ERα can be detrimental to the expression of genes it
regulates. This could be due to widespread condensate formation which in turn could sequester
the transcriptional protein complexes and competitively abrogate transcription across the
multiple loci. Thus, the global level of ERα in the nucleus can be predictive of transcriptional
status of specific genes. The binding of ERα at 1 and 3h is proportional to its nuclear levels (Fig.
3A); suggesting, its overexpression would lead to more binding in the genome which is
detrimental to gene expression.
307
308   **LLPS perturbation down-regulates *TFF1* but supports *TFF3* expression**
309
The data obtained from combined smFISH and ERα immunofluorescence indicates that ERα
could be a determining factor in controlling the differential gene expression of *TFF1* and *TFF3*.
Existing literature indicates that ERα-mediated condensate plays a role in E2-induced gene
expression (Boija et al., 2018; Nair et al, 2019; Sabari et al., 2018; Saravanan et al., 2020). It led
us to hypothesize that ERα-mediated condensate at the *TFF1* locus could be sequestering all the
factors required for active transcription and thus preventing activation of the *TFF3* locus at the
active phase of E2 signalling. To validate this, we treated the cells with 3% 1,6-Hexanediol for 5
minutes which is known to disrupt LLPS (Gamliel et al., 2022; Kroschwald et al., 2017).
Following the treatment, we performed smFISH to look at the transcription of *TFF1* and *TFF3* in
the same cell and observed a dramatic reduction in *TFF1* transcripts, whereas statistically
significant increase in *TFF3* was noted (Fig. 5A). Since, ERα forms condensates only after
estrogen stimulation, 1,6-Hexanediol had no effect on *TFF1* in the absence of E2 signalling (Fig.

S5A). Together, these results suggest that the functional loss of *TFF1* promoter transcription due
to the dissolution of ERα condensate allowed *TFF3* promoter in the neighbourhood to gain
access to transcriptional machinery leading to its upregulation. In order to test the generality of
this observation, beyond the *TFF1*, *TFF3* locus, we divided E2-responsive genes in three
categories, low, moderate, and high based on their expression upon E2-40m stimulation. We
observed the significant increase at 40m and down-regulation at 160m when the binding of ERα
is reduced in the genome (Fig. 5B). Next, we tested the expression of their nearby upregulated
genes at 160m compared to 40m post E2 stimulation. The nearby genes are within a distance of
1Mb from the E2-responsive genes which is approximately the average size of a TAD. Indeed,
we observed the significant up regulation of nearby genes at 160m when the expression of highly
induced genes dropped. Additionally, the expression of these genes was reduced at the peak of
signalling (Fig. 5B), showing that at 40m of signalling, the acute activation of primary genes and
sequestration of transcription machinery by these genes leads to the loss of expression of nearby
genes.
336
337
338 **Discussion**
339
340 smFISH allowed us to simultaneously capture allele level transcription of two genes in the same
341 TAD, *TFF1* and *TFF3* at single cell level during the peak and fall of signalling. We were able to
342 capture the anti-correlated expression of these two genes revealing an intricate regulatory
343 feedback between acutely activated enhancer-dependent gene, *TFF1* which caused the
344 dysregulation of non-enhancer targeted gene, *TFF3* within the same TAD (Fig. 5B).
345
Our data suggests that while condensate formation on enhancer allows robust activation of target
gene however, it negatively impacts the expression of other neighbouring genes potentially due
to the sequestration of transcriptional machinery from these genes. When the condensates
dissolve, the locally enriched transcriptional machinery is available to the other loci, potentially
allowing for increased transcription of the these genes, that are not the direct target of enhancer
(Fig 5C). Forced dissolution of condensates allows the expression of non-enhancer target genes
in the cells although at low levels. The data also suggests that globally, the neighbouring genes to
ligand induced genes are upregulated at the fall of signalling as an indirect consequence of
excess polymerase availability in the neighbourhood.
355
356
357 **Non-enhancer target genes are also regulated indirectly in the same TAD**
358
ERα binding strength increases at the enhancer of *TFF1* at 1h of ligand stimulation (Saravanan et
al., 2020). Such estrogen-induced binding of ERα leads to robust ligand induced activation of
TFF1 gene. While, as *TFF3* promoter remains unbound by ERα (Fig. 3A) and does not interact
with *TFF1* enhancer throughout the course of signalling. These data suggest that *TFF3*
expression is both ERα and enhancer independent. Despite non-dependence, *TFF3* expression
was mutually exclusive to *TFF1* expression and was dampened in the absence of enhancer. The
latter could be due to the enhancer mediated repositioning of the entire TAD that benefitted the
expression of *TFF3*.
367

The mean expression of *TFF1* was many fold higher than *TFF3* thus explaining its dependence
on an active enhancer to its upstream. These cells that expressed *TFF1*, did not express *TFF3* as
efficiently suggesting Pol2 was potentially sequestered from *TFF3* promoter. However, at 3h
when liganded ERα degrades leading to dissolution of condensates on enhancer, *TFF3*
expression increased suggesting a local redistribution of active transcription machinery to other
genes within the same chromatin domain. Even at 1h, when condensates were perturbed by 1,6
HD, the expression of *TFF3* increased suggesting the access to free Pol2 pool by *TFF3*
promoter. The data explains negative-correlation between *TFF1* and *TFF3* expression at the peak
of signalling and robust activation of *TFF3* at 3h when *TFF1* expression decreased at single cell
level.
378
379 **High and low levels of ERα were detrimental to enhancer mediated activation of TFF1:**
380
The cells with very high nuclear levels of ERα do not support optimum transcription of either
gene. Further, *TFF1* expression was more in cells showing mid-ERα levels at 3h while, *TFF3*
peaked in cells with low ERα levels. The reason for this observation could lie in the fact that
very low levels of TF do not reach or exceed the critical concentration required for LLPS (for
TFF1) but this poor transcriptional state of *TFF1* was favourable for *TFF3*. Alternatively, a very
high concentration of the TF could result in the formation of extremely dense homogenous
condensates effectively reducing transcription (Chong et al., 2022; Ryu et al., 2024). Our data
suggests that such a mechanism may be general and not just applicable to the *TFF1-TFF3* pair
(Fig. 5B). Together, our study underscores the indirect effects of ligand-induced chronic gene
transcription that is dependent on enhancer activation and phase-separation within a TAD.
391
392
393 **Materials and Methods**
394
395 **Cell culture:**
396
MCF-7 cells (WT scramble control and Δ*TFF1*e were generated in Saravanan et al., 2020) cells
were cultured in non-stripping media consisting of DMEM (Gibco, 12100-046) supplemented
with 10% FBS (Gibco, 16000-044) and 1% Penicillin-Streptomycin-Glutamine (Gibco, 10378-
016) in a 5% CO_2 humidified incubator at 37 °C (unstripped MCF-7). Cells were passaged and
seeded into glass bottom dishes in non-stripping media for 24h and allowed to reach 60–80%
confluency. These cells were then hormone-stripped for three days in stripping media containing
phenol red-free DMEM (Gibco, 21063-029) supplemented with 5% charcoal–stripped
FBS(Gibco,12676029) and maintained in the humidified incubator at 37 °C (stripped MCF-7).
In order to induce the estrogen transcriptional responses, on the third day cells were treated with
β-estradiol (E2758, Sigma-Aldrich) at 100nM concentration for various periods as mentioned in
the respective Figures. For untreated control, cells were either treated with equal microliters of
ethanol on the third day or with ERα inhibitor ICI182780 (1047, Tocris Biosciences) at 100nM
concentration for 24 h after two days of stripping.
For perturbation of LLPS in E2-induced condition, cells were incubated with media containing
3% 1,6-Hexanediol for 5 minutes (after 30 minutes of E2 treatment) followed by removal of 1,6-
Hexanediol and recovery for 25 mins in E2 containing media.

For experiments with overexpression of EGFP- ERα or EGFP, cells were transfected 24 hours
before E2 induction using X-tremeGENE™ HP DNA Transfection Reagent (XTGHP-RO) in
stripping media.
416
417   **Probe for smFISH:**
418
419   Probes were designed, custom-made, and tagged with indicated fluorophores (Quasar 570 or Cal
420   610) by Biosearch Technologies (https://www.biosearchtech.com/products/rna-fish). Probes
421   were made to target the intronic or exonic regions of the *TFF1* and *TFF3* genes. Probe sequences
422   can be found in supplementary table 1-4 in supplementary materials.
423
424   **Chromatin retention assay and smFISH:**
425
For smFISH experiments without chromatin retention assay cells, the cells were treated with
estradiol or vehicular control for indicated times. Following this, the cells were washed with
nuclease-free 1X PBS (Ambion, AM9624) twice. This was followed by fixation using 4%
paraformaldehyde (PFA, Sigma, P6148) in 1X PBS for 10 minutes at room temperature. The
fixative was removed and cells were washed twice with 1X PBS. The cells were permeabilized
using 70% ethanol at 4°C overnight. On the next day, the permeabilizing agent was removed and
the cells were washed twice with 1X PBS. The cells were washed once with a wash buffer (10%
Formamide (Ambion, 9342) and 2X SSC (Ambion, AM9763) in nuclease-free water) for 5
minutes at room temperature. The wash buffer was aspirated and the cells were incubated with a
hybridization mix (100 μl of mix contains 10% formamide, 89 μl hybridization buffer, and 1 μl
each of indicated probes) overnight at 37°C in a humid chamber. The next day, the solution was
removed and cells were washed with the wash buffer at 37°C for 30 minutes. To stain the nuclei,
DAPI (Invitrogen, D1306; 2μg/ml) in wash buffer was added to the cells and incubated for 10
minutes at 37°C. The cells were then washed with 2X SSC for 5 minutes at 4°C. The solution
was removed and the cells were covered with a few drops of the mountant Vectashield (Vector
Labs). The plates were imaged after at least 1h.
For smFISH experiments with chromatin retention assay cells were treated with estradiol or
vehicular control for indicated times. Following this, the cells were treated with CSK buffer
(consisting of 10 mM PIPES buffer, 100 mM NaCl, 3 mM MgCl2, 300 mM sucrose, and 0.7%
Triton-X 100) for 15 minutes at room temperature. After this, the cells were washed with
nuclease-free 1X PBS twice. This was followed by fixation using 4% paraformaldehyde in 1X
PBS for 10 minutes at room temperature. The fixative was removed and cells were washed twice
with 1X PBS. The cells were permeabilized using 0.3% Triton-X 100 (Sigma, T8787) in 1X PBS
for 10 minutes at room temperature. The permeabilizing agent was aspirated and the cells were
washed twice with 1X PBS. This was followed by a washing step using the wash buffer for 5
minutes at room temperature. Following this, the cells were incubated with the hybridization mix
to which the antibody of interest was also added at appropriate dilution. The next day, the
hybridization mix containing probes and antibodies was removed. Cells were washed twice with
1X PBS. Then the cells were incubated with 1X PBS containing a secondary antibody against the
antibody of interest at room temperature for 2h. This was followed by incubation with the wash
buffer at 37°C for 30 minutes. To stain the nuclei, DAPI in wash buffer was added to the cells
and incubated for 10 minutes at 37°C. The cells were then washed with 2X SSC for 5 minutes at

458 4°C. The solution was removed and the cells were covered with a few drops of the mountant
459 Vectashield. The plates were imaged after at least 1h.
460

Antibody staining/ immunofluorescence:

463 The primary antibody against ERα (Santa Cruz, sc8002(F10)) was added to the hybridization
464 mix at the dilution of 1:400 and incubated overnight at 37°C. The next day, an anti-mouse Alexa
465 Fluor™ 488 secondary antibody (Invitrogen, A11029) was added to 1X PBS at the dilution of
466 1:1000. Cells were incubated for 2h followed by the continuation of the smFISH protocol as
467 indicated above.
468

Image acquisition:

471 The plates were imaged on an Olympus IX83 inverted widefield fluorescence microscope with a
472 Retiga 6000 CCD monochrome camera (QImaging). The images were acquired using a 60X,
473 1.42 N.A. oil immersion objective or a 100X, 1.4 N.A. oil immersion objective. The z-step size
474 was 0.3μm and 35 slices were acquired. The resolution at which the images were acquired is
475 2752 x 2208. Narrow band-pass filters were used to distinguish the signal from Quasar 570 and
476 Cal 610 labeled probes (ChromaTechnology- 49309 and 49310).
477

Image analysis and representation:

480 The images of mRNA channels were subjected to background subtraction using a rolling ball
481 radius of 10 pixels across the entire stack using Fiji. For representative images, the stacks were
482 Z-projected in Fiji and shown. Transcripts were counted using either Rajlabimagetools (courtesy
483 of Arjun Raj lab) or an in-house MATLAB (Mathworks) script based on earlier work (Femino et
484 al., 1998; Raj et al., 2008). Briefly, the code uses a nuclear mask to count the intron-containing
485 RNA present in the nucleus (InTFF), and another mask for the whole cell to count the mature
486 mRNA, detected by exonic probes (ExTFF), that are present in the nucleus as well as the
487 cytoplasm. The mean intensity for the ERα immunofluorescence channel was also quantified
488 using the MATLAB script on a cell-by-cell basis using the nuclear mask.
489

Graphing and Statistics

491 The graphs were plotted using Python 3, MATLAB, and Origin Pro and edited using Adobe
492 Photoshop. To perform a t-test for data combined from three repeats, GraphPad
493 (https://www.graphpad.com/quickcalcs/ttest1/) was used. Non-parametric tests were performed
494 in case of single cell data. For this Mann-Whitney test was used. Significance is represented as :
495 *** denotes $p < 0.001$,** denotes $p < 0.01$,* denotes $p < 0.05$, ns denotes $p > 0.05$.

Circular Chromatin Conformation Capture-seq:

497 4C was performed as per the protocol described in van de Werken et al., 2012 with minor
498 variations. MCF7 cells were fixed with fresh formaldehyde (1.5%) and quenched with glycine
499 (125mM) followed by washes with ice-cold 1XPBS (2X) and scraped, pelleted, and stored at -
500 80^0C. Lysis buffer [Tris-Cl pH 8.0 (10mM), NaCl (10mM), NP-40 (0.2%), PIC (1X)] was added
501 to the pellets and homogenized by Dounce homogenizer (20 stroked with pestle A followed by
502 pestle B). The 3C digestion was performed with DpnII (200 units, NEB) and ligation was

performed by T4 DNA ligase and ligation mix [Triton X-100 (1%), 1x Ligation buffer (10X Ligation buffer- Tris-Cl pH 7.5 (500mM), MgCl2 (100mM), DTT (100mM), BSA (0.105mg/ml), ATP (1.05mM)]. The ligated samples were purified by PCI and subjected to ethanol precipitation. The pellet was eluted in 1X TE (pH 8.0) to obtain the 3C library. The 4C digestion was performed by NlaIII (50 units, NEB), and the samples were ligated, purified, and precipitated similar to the 3C library to obtain the 4C library. The 4C library was subjected to RNaseA treatment and purified with the QIAquick PCR purification kit. The concentration of the library was then measured by Nanodrop and subjected to PCRs using the oligos for the enhancer viewpoint. The samples were next PCR purified using the same kit and subjected to next-generation sequencing with Illumina HiSeq2500/ NOVA seq. The 4C oligos are listed in supplementary table 5.

4C Data Analysis:

The sequenced reads in fastq file were demultiplexed by matching the appropriate primer sequences for each condition without allowing for any mismatches. Demultiplexed reads were processed using 4cseqpipe software. Restriction site tracks were created for the hg38 human genome by mentioning the restriction sites of the first cutter as GATC and that of the second cutter as CATG. The phred scores of demultiplexed reads were changed to phred64 format and the fastq files were converted into raw format. Further, valid 4C reads were mapped to the generated restriction site tracks. Unique fragment ends/non-unique fragment ends were used. The mapped reads were normalized and near cis domainograms at a maximum height of 0.1 were created by using the truncated log mean statistic with a trend resolution of 1kb for the genomic region chr11:42300000-42400000.

RNA Isolation, cDNA synthesis and PCR:

Cells were lysed in 1 ml of Trizol (Thermo Fisher Inc.). 200 ul chloroform was added to the sample, briefly vortexed and centrifuged at 12K rpm for 12 mins. The aqueous phase was carefully collected and transferred to the fresh tube. 1 volume of isopropanol was added to the sample and incubated at room temperature for 10 mins to precipitate the RNA. The samples were centrifuged at 12K rpm for 12 mins, supernatant was discarded without disturbing the pellet. The pellet obtained was washed with 75% ethanol. Pellet was air dried and dissolved in RNase free water. RNA obtained was treated with ezDNase (Invitrogen) to remove the traces of contaminating DNA. 1µg of RNA was used for each cDNA synthesis reaction by Superscript IV (Invitrogen) and random hexamers as per manufacturer's recommendation. The CFX96 touch (Biorad) real time PCR was used for qRT-PCRs. The fold changes were calculated by the ΔΔCt method and individual expression data was normalized to GAPDH mRNA. The *p*-values were calculated by Student's unpaired two-tailed *t*-test from independent four biological replicates. qRT-PCR primers are listed in Supplementary Table 6.

GRO-Seq analysis:

Fastq files from GEO accession number GSE43836 were downloaded from European Nucleotide Archive. Reads with base quality <20 in a sliding window of 4 bases and with a length of <36 were removed using Trimmomatic 0.39. Trimmed reads were aligned using bowtie2 2.5.1 with default parameters. Duplicate reads from the alignment files were removed using samtools

1.16.1. De-duplicated aligned reads were assigned in a strand-specific manner to transcript feature of the hg38.ncbiRefSeq.gtf.gz file by allowing multi-mapping reads and considering the largest overlap in case of overlapping features using featureCounts v2.0.3. CPM normalised strand-specific signal files with a bin size of 1 bp were generated using bamCoverage 3.5.1. Differential gene expression analysis was performed with default parameters using Deseq2 1.36.0. Upregulated genes were defined as those with adjusted p-value <0.05 and log2(FC) >1 in their respective conditions. Top, Middle and bottom 10% of upregulated genes based on their base mean value in E2-40m vs VEH condition were subsetted as highly expressing, moderately expressing and low expressing upregulated genes. The closest upregulated genes in E2-160m vs E2-40m near highly expressing, moderately expressing and low expressing upregulated genes in E2-40m vs VEH condition were identified using bedtools closest v2.30.0. Only genes within a genomic distance of 1Mb were considered. Log2 transformed DEseq2 normalised counts with the addition of the arbitrary value 1 was used to compare the gene expression trends across time points and categories of upregulated genes. All plots were generated using R 4.2.2.

Funding and Acknowledgements: This project was supported by intramural funds at TIFR Hyderabad from the Department of Atomic Energy, Government of India (Project Identification No. RTI 4007). We further acknowledge support of the Department of Atomic Energy, Government of India, under project no. 12-R&D-TFR-5.04-0800 and intramural funds from National Centre for Biological Sciences–Tata Institute of Fundamental Research (NCBS-TIFR). D.N. is an EMBO Global Investigator. We also acknowledge the funding support from Wellcome-IA (IA/1/14/2/501539). DB is supported by TIFR-Hyderabad PhD program. ZI acknowledges funding support from ICMR, India. SN acknowledges funding support from DBT-JRF, India.

Contribution: The project was conceived by DN, AM, DB and ZI. Experiments were performed by DB and ZI. 4C and GRO-seq data analysis was performed by SN. The manuscript was written by DB, ZI, DN and AM. All authors approved the manuscript.

Competing Interest: Authors declare no competing interest.

References

Ahmed, J., Meszaros, A., Lazar, T., & Tompa, P. (2021). DNA-binding domain as the minimal region driving RNA-dependent liquid-liquid phase separation of androgen receptor. Protein Science, 30(7), 1380–1392. https://doi.org/10.1002/pro.4100

Boehning, M., Dugast-Darzacq, C., Rankovic, M., Hansen, A. S., Yu, T., Marie-Nelly, H., McSwiggen, D. T., Kokic, G., Dailey, G. M., Cramer, P., Darzacq, X., & Zweckstetter, M. (2018). RNA polymerase II clustering through carboxy-terminal domain phase separation. *Nature Structural & Molecular Biology*, *25*(9), 833–840. https://doi.org/10.1038/s41594-018-0112-y

Boija, A., Klein, I. A., Sabari, B. R., Dall'Agnese, A., Coffey, E. L., Zamudio, A. V., Li, C. H., Shrinivas, K., Manteiga, J. C., Hannett, N. M., Abraham, B. J., Afeyan, L. K., Guo, Y. E., Rimel,

J. K., Fant, C. B., Schuijers, J., Lee, T. I., Taatjes, D. J., & Young, R. A. (2018). Transcription Factors Activate Genes through the Phase-Separation Capacity of Their Activation Domains. *Cell*, *175*(7), 1842-1855.e16. https://doi.org/10.1016/j.cell.2018.10.042

Bojcsuk, D., Nagy, G., & Balint, B. L. (2017). Inducible super-enhancers are organized based on canonical signal-specific transcription factor binding elements. Nucleic Acids Research, 45(7), 3693–3706. https://doi.org/10.1093/nar/gkw1283

Brown, A. M., Jeltsch, J. M., Roberts, M., & Chambon, P. (1984). Activation of pS2 gene transcription is a primary response to estrogen in the human breast cancer cell line MCF-7. Proceedings of the National Academy of Sciences, 81(20), 6344–6348. https://doi.org/10.1073/pnas.81.20.6344

Buecker, C., Srinivasan, R., Wu, Z., Calo, E., Acampora, D., Faial, T., Simeone, A., Tan, M., Swigut, T., & Wysocka, J. (2014). Reorganization of enhancer patterns in transition from naive to primed pluripotency. *Cell Stem Cell*, *14*(6), 838–853. https://doi.org/10.1016/j.stem.2014.04.003

Bulger, M., & Groudine, M. (1999). Looping versus linking: Toward a model for long-distance gene activation. *Genes & Development*, *13*(19), 2465–2477.

Cai, D., Feliciano, D., Dong, P., Flores, E., Gruebele, M., Porat-Shliom, N., Sukenik, S., Liu, Z., & Lippincott-Schwartz, J. (2019). Phase separation of YAP reorganizes genome topology for long-term YAP target gene expression. *Nature Cell Biology*, *21*(12), 1578–1589. https://doi.org/10.1038/s41556-019-0433-z

Chalbos, D., Philips, A., Galtier, F., & Rochefort, H. (1993). Synthetic antiestrogens modulate induction of pS2 and cathepsin-D messenger ribonucleic acid by growth factors and adenosine 3',5'-monophosphate in MCF7 cells. Endocrinology, 133(2), 571–576. https://doi.org/10.1210/en.133.2.571

Chen, H., Levo, M., Barinov, L., Fujioka, M., Jaynes, J. B., & Gregor, T. (2018). Dynamic interplay between enhancer-promoter topology and gene activity. *Nature Genetics*, *50*(9), 1296–1303. https://doi.org/10.1038/s41588-018-0175-z

Chepelev, I., Wei, G., Wangsa, D., Tang, Q., & Zhao, K. (2012). Characterization of genome-wide enhancer-promoter interactions reveals co-expression of interacting genes and modes of higher order chromatin organization. *Cell Research*, *22*(3), 490–503. https://doi.org/10.1038/cr.2012.15

Chinery, R., Williamson, J., & Poulsom, R. (1996). The Gene Encoding Human Intestinal Trefoil Factor (*TFF3*) Is Located on Chromosome 21q22.3 Clustered with Other Members of the Trefoil Peptide Family. Genomics, 32(2), 281–284. https://doi.org/10.1006/geno.1996.0117

Cho, W.-K., Spille, J.-H., Hecht, M., Lee, C., Li, C., Grube, V., & Cisse, I. I. (2018). Mediator and RNA polymerase II clusters associate in transcription-dependent condensates. Science, 361(6400), 412–415. https://doi.org/10.1126/science.aar4199

Chong, S., Dugast-Darzacq, C., Liu, Z., Dong, P., Dailey, G. M., Cattoglio, C., Heckert, A.,
Banala, S., Lavis, L., Darzacq, X., & Tjian, R. (2018). Imaging dynamic and selective low-
complexity domain interactions that control gene transcription. *Science*, *361*(6400), eaar2555.
https://doi.org/10.1126/science.aar2555

Chong, S., Graham, T. G. W., Dugast-Darzacq, C., Dailey, G. M., Darzacq, X., & Tjian, R.
(2022). Tuning levels of low-complexity domain interactions to modulate endogenous oncogenic
transcription.   *Molecular   Cell*,   *82*(11),   2084-2097.e5.
https://doi.org/10.1016/j.molcel.2022.04.007

Clark, M. B., Mercer, T. R., Bussotti, G., Leonardi, T., Haynes, K. R., Crawford, J., Brunck, M.
E., Cao, K.-A. L., Thomas, G. P., Chen, W. Y., Taft, R. J., Nielsen, L. K., Enright, A. J., Mattick,
J. S., & Dinger, M. E. (2015). Quantitative gene profiling of long noncoding RNAs with targeted
RNA sequencing. *Nature Methods*, *12*(4), 339–342. https://doi.org/10.1038/nmeth.3321
636
637 Conesa, A., Madrigal, P., Tarazona, S., Gomez-Cabrero, D., Cervera, A., McPherson, A.,
638 Szcześniak, M. W., Gaffney, D. J., Elo, L. L., Zhang, X., & Mortazavi, A. (2016). A survey of
639 best   practices   for   RNA-seq   data   analysis.   *Genome   Biology*,   *17*(1),   13.
640 https://doi.org/10.1186/s13059-016-0881-8
641
Coté, A., O'Farrell, A., Dardani, I., Dunagin, M., Coté, C., Wan, Y., Bayatpour, S., Drexler, H.
L., Alexander, K. A., Chen, F., Wassie, A. T., Patel, R., Pham, K., Boyden, E. S., Berger, S.,
Phillips-Cremins, J., Churchman, L. S., & Raj, A. (2023). Post-transcriptional splicing can occur
in a slow-moving zone around the gene. *eLife*, *12*. https://doi.org/10.7554/eLife.91357.2

Coulon, A., Ferguson, M. L., de Turris, V., Palangat, M., Chow, C. C., & Larson, D. R. (2014).
Kinetic competition during the transcription cycle results in stochastic RNA processing. *eLife*, *3*,
e03939. https://doi.org/10.7554/eLife.03939

Dixon, B. S., Sharma, R. V., & Dennis, M. J. (1996). The Bradykinin B2 Receptor Is a Delayed
Early Response Gene for Platelet-derived Growth Factor in Arterial Smooth Muscle Cells *.
Journal   of   Biological   Chemistry,   *271*(23),   13324–13332.
https://doi.org/10.1074/jbc.271.23.13324

Drexler, H. L., Choquet, K., & Churchman, L. S. (2020). Splicing Kinetics and Coordination
Revealed by Direct Nascent RNA Sequencing through Nanopores. *Molecular Cell*, *77*(5), 985-
998.e8. https://doi.org/10.1016/j.molcel.2019.11.017

Femino, A. M., Fay, F. S., Fogarty, K., & Singer, R. H. (1998). Visualization of single RNA
transcripts   in   situ.   *Science   (New   York,   N.Y.)*,   *280*(5363),   585–590.
https://doi.org/10.1126/science.280.5363.585

Fowler, T., Sen, R., & Roy, A. L. (2011). Regulation of Primary Response Genes. *Molecular
Cell*, *44*(3), 348–360. https://doi.org/10.1016/j.molcel.2011.09.014

Freter, R. R., Alberta, J. A., Hwang, G. Y., Wrentmore, A. L., & Stiles, C. D. (1996). Platelet-
derived Growth Factor Induction of the Immediate-early gene MCP-1 Is Mediated by NF-κB and
a 90-kDa Phosphoprotein Coactivator *. *Journal of Biological Chemistry*, *271*(29), 17417–
17424. https://doi.org/10.1074/jbc.271.29.17417

Friedman, M. J., Wagner, T., Lee, H., Rosenfeld, M. G., & Oh, S. (2024). Enhancer–promoter
specificity in gene transcription: Molecular mechanisms and disease associations. *Experimental*
& Molecular Medicine, *56*(4), 772–787. https://doi.org/10.1038/s12276-024-01233-y

Fullwood, M. J., Liu, M. H., Pan, Y. F., Liu, J., Xu, H., Mohamed, Y. B., Orlov, Y. L., Velkov,
S., Ho, A., Mei, P. H., Chew, E. G. Y., Huang, P. Y. H., Welboren, W.-J., Han, Y., Ooi, H. S.,
Ariyaratne, P. N., Vega, V. B., Luo, Y., Tan, P. Y., … Ruan, Y. (2009). An oestrogen-receptor-
α-bound human chromatin interactome. Nature, 462(7269), Article 7269.
https://doi.org/10.1038/nature08497

Furlong, E. E. M., & Levine, M. (2018). Developmental enhancers and chromosome topology.
Science, *361*(6409), 1341–1345. https://doi.org/10.1126/science.aau0320

Galouzis, C. C., & Furlong, E. E. M. (2022). Regulating specificity in enhancer-promoter
communication. *Current Opinion in Cell Biology*, *75*, 102065.
https://doi.org/10.1016/j.ceb.2022.01.010

Gamliel, A., Meluzzi, D., Oh, S., Jiang, N., Destici, E., Rosenfeld, M. G., & Nair, S. J. (2022).
Long-distance association of topological boundaries through nuclear condensates. Proceedings of
the National Academy of Sciences of the United States of America, 119(32), e2206216119.
https://doi.org/10.1073/pnas.2206216119

Hah, N., Danko, C. G., Core, L., Waterfall, J. J., Siepel, A., Lis, J. T., & Kraus, W. L. (2011). A
rapid, extensive, and transient transcriptional response to estrogen signaling in breast cancer
cells. Cell, 145(4), 622–634. https://doi.org/10.1016/j.cell.2011.03.042

Hah, N., Murakami, S., Nagari, A., Danko, C. G., & Kraus, W. L. (2013). Enhancer transcripts
mark active estrogen receptor binding sites. *Genome Research*, *23*(8), 1210–1223.
https://doi.org/10.1101/gr.152306.112

Haimovich, G., & Gerst, J. E. (2018). Single-molecule Fluorescence in situ Hybridization
(smFISH) for RNA Detection in Adherent Animal Cells. *Bio-Protocol*, *8*(21), e3070.
https://doi.org/10.21769/BioProtoc.3070
691

Herschman, H. R. (1991). Primary response genes induced by growth factors and tumor
promoters. *Annual Review of Biochemistry*, *60*(Volume 60, 1991), 281–319.
https://doi.org/10.1146/annurev.bi.60.070191.001433

Jeltsch, J. M., Roberts, M., Schatz, C., Garnier, J. M., Brown, A. M., & Chambon, P. (1987).
Structure of the human oestrogen-responsive gene pS2. Nucleic Acids Research, 15(4), 1401–
1414.

Khodor, Y. L., Menet, J. S., Tolan, M., & Rosbash, M. (2012). Cotranscriptional splicing
efficiency differs dramatically between Drosophila and mouse. *RNA*, *18*(12), 2174–2186.
https://doi.org/10.1261/rna.034090.112

Kim, J., Petz, L. N., Ziegler, Y. S., Wood, J. R., Potthoff, S. J., & Nardulli, A. M. (2000).
Regulation of the estrogen-responsive pS2 gene in MCF-7 human breast cancer cells. The

Journal of Steroid Biochemistry and Molecular Biology, 74(4), 157–168. https://doi.org/10.1016/S0960-0760(00)00119-9

Koşar, Z., & Erbaş, A. (2022). Can the Concentration of a Transcription Factor Affect Gene Expression? *Frontiers in Soft Matter*, *2*. https://www.frontiersin.org/articles/10.3389/frsfm.2022.914494

Kroschwald, S., Maharana, S., & Simon, A. (2017, May 22). Hexanediol: A chemical probe to investigate the material properties of membrane-less compartments. Matters. https://doi.org/10.19185/matters.201702000010

Kwon, S. (2013). Single-molecule fluorescence in situ hybridization: Quantitative imaging of single RNA molecules. *BMB Reports*, *46*(2), 65–72. https://doi.org/10.5483/BMBRep.2013.46.2.016

Levenson, A. S., & Jordan, V. C. (1997). MCF-7: The First Hormone-responsive Breast Cancer Cell Line1. Cancer Research, 57(15), 3071–3078.

Levine, M., & Tjian, R. (2003). Transcription regulation and animal diversity. *Nature*, *424*(6945), 147–151. https://doi.org/10.1038/nature01763

Li, W., Notani, D., Ma, Q., Tanasa, B., Nunez, E., Chen, A. Y., Merkurjev, D., Zhang, J., Ohgi, K., Song, X., Oh, S., Kim, H.-S., Glass, C. K., & Rosenfeld, M. G. (2013). Functional roles of enhancer RNAs for oestrogen-dependent transcriptional activation. *Nature*, *498*(7455), 516–520. https://doi.org/10.1038/nature12210

Liu, Z., Merkurjev, D., Yang, F., Li, W., Oh, S., Friedman, M. J., Song, X., Zhang, F., Ma, Q., Ohgi, K. A., Krones, A., & Rosenfeld, M. G. (2014). Enhancer Activation Requires trans-Recruitment of a Mega Transcription Factor Complex. *Cell*, *159*(2), 358–373. https://doi.org/10.1016/j.cell.2014.08.027

Mann, R., & Notani, D. (2023). Transcription factor condensates and signaling driven transcription. *Nucleus*, *14*(1), 2205758. https://doi.org/10.1080/19491034.2023.2205758

Masiakowski, P., Breathnach, R., Bloch, J., Gannon, F., Krust, A., & Chambon, P. (1982). Cloning of cDNA sequences of hormone-regulated genes from the MCF-7 human breast cancer cell line. Nucleic Acids Research, 10(24), 7895–7903. https://doi.org/10.1093/nar/10.24.7895

Nair, S. J., Yang, L., Meluzzi, D., Oh, S., Yang, F., Friedman, M. J., Wang, S., Suter, T., Alshareedah, I., Gamliel, A., Ma, Q., Zhang, J., Hu, Y., Tan, Y., Ohgi, K. A., Jayani, R. S., Banerjee, P. R., Aggarwal, A. K., & Rosenfeld, M. G. (2019). Phase separation of ligand-activated enhancers licenses cooperative chromosomal enhancer assembly. Nature Structural & Molecular Biology, 26(3), Article 3. https://doi.org/10.1038/s41594-019-0190-5

Nunez, A. M., Berry, M., Imler, J. L., & Chambon, P. (1989). The 5' flanking region of the pS2 gene contains a complex enhancer region responsive to oestrogens, epidermal growth factor, a tumour promoter (TPA), the c-Ha-ras oncoprotein and the c-jun protein. The EMBO Journal, 8(3), 823–829.

Oh, S., Shao, J., Mitra, J., Xiong, F., D'Antonio, M., Wang, R., Garcia-Bassets, I., Ma, Q., Zhu, X., Lee, J.-H., Nair, S. J., Yang, F., Ohgi, K., Frazer, K. A., Zhang, Z. D., Li, W., & Rosenfeld, M. G. (2021). Enhancer release and retargeting activates disease-susceptibility genes. *Nature*, *595*(7869), Article 7869. https://doi.org/10.1038/s41586-021-03577-1

Panigrahi, A., & O'Malley, B. W. (2021). Mechanisms of enhancer action: The known and the unknown. *Genome Biology*, *22*(1), Article 1. https://doi.org/10.1186/s13059-021-02322-1

Ptashne, M. (1986). Gene regulation by proteins acting nearby and at a distance. *Nature*, *322*(6081), 697–701. https://doi.org/10.1038/322697a0

Quintin, J., Le Péron, C., Palierne, G., Bizot, M., Cunha, S., Sérandour, A. A., Avner, S., Henry, C., Percevault, F., Belaud-Rotureau, M.-A., Huet, S., Watrin, E., Eeckhoute, J., Legagneux, V., Salbert, G., & Métivier, R. (2014). Dynamic Estrogen Receptor Interactomes Control Estrogen-Responsive Trefoil Factor (TFF) Locus Cell-Specific Activities. Molecular and Cellular Biology, 34(13), 2418–2436. https://doi.org/10.1128/MCB.00918-13

Raj, A., Peskin, C. S., Tranchina, D., Vargas, D. Y., & Tyagi, S. (2006). Stochastic mRNA Synthesis in Mammalian Cells. *PLOS Biology*, *4*(10), e309. https://doi.org/10.1371/journal.pbio.0040309

Raj, A., van den Bogaard, P., Rifkin, S. A., van Oudenaarden, A., & Tyagi, S. (2008). Imaging individual mRNA molecules using multiple singly labeled probes. *Nature Methods*, *5*(10), Article 10. https://doi.org/10.1038/nmeth.1253

Rao, S.S.P. *et al.* (2014) 'A 3D map of the human genome at kilobase resolution reveals principles of chromatin looping', *Cell*, 159(7), pp. 1665–1680. doi:10.1016/j.cell.2014.11.021.

Richter, W. F., Nayak, S., Iwasa, J., & Taatjes, D. J. (2022). The Mediator complex as a master regulator of transcription by RNA polymerase II. Nature Reviews Molecular Cell Biology, 23(11), Article 11. https://doi.org/10.1038/s41580-022-00498-3

Ryu, K., Park, G., & Cho, W.-K. (2024). Emerging insights into transcriptional condensates. *Experimental & Molecular Medicine*, *56*(4), 820–826. https://doi.org/10.1038/s12276-024-01228-9

Sabari, B. R., Dall'Agnese, A., Boija, A., Klein, I. A., Coffey, E. L., Shrinivas, K., Abraham, B. J., Hannett, N. M., Zamudio, A. V., Manteiga, J. C., Li, C. H., Guo, Y. E., Day, D. S., Schuijers, J., Vasile, E., Malik, S., Hnisz, D., Lee, T. I., Cisse, I. I., … Young, R. A. (2018). Coactivator condensation at super-enhancers links phase separation and gene control. *Science*, *361*(6400), eaar3958. https://doi.org/10.1126/science.aar3958

Sanyal, A., Lajoie, B. R., Jain, G., & Dekker, J. (2012). The long-range interaction landscape of gene promoters. *Nature*, *489*(7414), 109–113. https://doi.org/10.1038/nature11279

Saravanan, B., Soota, D., Islam, Z., Majumdar, S., Mann, R., Meel, S., Farooq, U., Walavalkar, K., Gayen, S., Singh, A. K., Hannenhalli, S., & Notani, D. (2020). Ligand dependent gene regulation by transient ERα clustered enhancers. *PLOS Genetics*, *16*(1), e1008516. https://doi.org/10.1371/journal.pgen.1008516

779 Sha, Y., Phan, J. H., & Wang, M. D. (2015). Effect of low-expression gene filtering on detection
780 of differentially expressed genes in RNA-seq data. *Conference Proceedings : ... Annual*
781 *International Conference of the IEEE Engineering in Medicine and Biology Society. IEEE*
782 *Engineering in Medicine and Biology Society. Annual Conference*, *2015*, 6461.
783 https://doi.org/10.1109/EMBC.2015.7319872
784

Shi, Y., Chen, J., Zeng, W.-J., Li, M., Zhao, W., Zhang, X.-D., & Yao, J. (2021). Formation of
nuclear condensates by the Mediator complex subunit Med15 in mammalian cells. BMC
Biology, 19(1), 245. https://doi.org/10.1186/s12915-021-01178-y

Shrinivas, K., Sabari, B. R., Coffey, E. L., Klein, I. A., Boija, A., Zamudio, A. V., Schuijers, J.,
Hannett, N. M., Sharp, P. A., Young, R. A., & Chakraborty, A. K. (2019). Enhancer Features
that Drive Formation of Transcriptional Condensates. *Molecular Cell*, *75*(3), 549-561.e7.
https://doi.org/10.1016/j.molcel.2019.07.009

Skinner, S. O., Xu, H., Nagarkar-Jaiswal, S., Freire, P. R., Zwaka, T. P., & Golding, I. (2016).
Single-cell analysis of transcription kinetics across the cell cycle. *eLife*, *5*, e12175.
https://doi.org/10.7554/eLife.12175

Stortz, M., Pecci, A., Presman, D. M., & Levi, V. (2020). Unraveling the molecular interactions
involved in phase separation of glucocorticoid receptor. *BMC Biology*, *18*(1), 59.
https://doi.org/10.1186/s12915-020-00788-2

Stortz, M., Presman, D. M., & Levi, V. (2024). Transcriptional condensates: A blessing or a
curse for gene regulation? *Communications Biology*, *7*(1), 1–10. https://doi.org/10.1038/s42003-
024-05892-5

Sun, J.-M., Spencer, V. A., Li, L., Yu Chen, H., Yu, J., & Davie, J. R. (2005). Estrogen
regulation of trefoil factor 1 expression by estrogen receptor α and Sp proteins. Experimental
Cell Research, 302(1), 96–107. https://doi.org/10.1016/j.yexcr.2004.08.015

Svensson, V., Natarajan, K. N., Ly, L.-H., Miragaia, R. J., Labalette, C., Macaulay, I. C., Cvejic,
A., & Teichmann, S. A. (2017). Power analysis of single-cell RNA-sequencing experiments.
Nature Methods, *14*(4), 381–387. https://doi.org/10.1038/nmeth.4220
807

808 Tarazona, S., García-Alcalde, F., Dopazo, J., Ferrer, A., & Conesa, A. (2011). Differential
809 expression in RNA-seq: A matter of depth. *Genome Research*, *21*(12), 2213–2223.
810 https://doi.org/10.1101/gr.124321.111
811

Uyehara, C. M., & Apostolou, E. (2023). 3D enhancer-promoter interactions and multi-
connected hubs: Organizational principles and functional roles. *Cell Reports*, *42*(4), 112068.
https://doi.org/10.1016/j.celrep.2023.112068

van de Werken HJ, de Vree PJ, Splinter E, Holwerda SJ, Klous P, de Wit E, de Laat W. 4C
technology: protocols and data analysis. Methods Enzymol. 2012;513:89-112. doi:
10.1016/B978-0-12-391938-0.00004-5. PMID: 22929766.

Williams, M., Lyu, M.-S., Yang, Y.-L., Lin, E. P., Dunbrack, R., Birren, B., Cunningham, J., &
Hunter, K. (1999). *Ier5,*a Novel Member of the Slow-Kinetics Immediate-Early Genes.
Genomics, *55*(3), 327–334. https://doi.org/10.1006/geno.1998.5679

Winkles, J. A. (1997). Serum- and Polypeptide Growth Factor-Inducible Gene Expression in
Mouse Fibroblasts. In K. Moldave (Ed.), *Progress in Nucleic Acid Research and Molecular*
Biology (Vol. 58, pp. 41–78). Academic Press. https://doi.org/10.1016/S0079-6603(08)60033-1

Winston, J. T., & Pledger, W. J. (1993). Growth factor regulation of cyclin D1 mRNA
expression through protein synthesis-dependent and -independent mechanisms. *Molecular*
Biology of the Cell, *4*(11), 1133–1144. https://doi.org/10.1091/mbc.4.11.1133

Yamamoto, K. R., & Alberts, B. M. (1976). Steroid Receptors: Elements for Modulation of
Eukaryotic Transcription. *Annual Review of Biochemistry*, *45*(Volume 45, 1976), 721–746.
https://doi.org/10.1146/annurev.bi.45.070176.003445

Yan, J., Chen, S.-A. A., Local, A., Liu, T., Qiu, Y., Dorighi, K. M., Preissl, S., Rivera, C. M.,
Wang, C., Ye, Z., Ge, K., Hu, M., Wysocka, J., & Ren, B. (2018). Histone H3 lysine 4
monomethylation modulates long-range chromatin interactions at enhancers. *Cell Research*,
28(2), 204–220. https://doi.org/10.1038/cr.2018.1

Zabidi, M. A., Arnold, C. D., Schernhuber, K., Pagani, M., Rath, M., Frank, O., & Stark, A.
(2015). Enhancer–core-promoter specificity separates developmental and housekeeping gene
regulation. *Nature*, *518*(7540), 556–559. https://doi.org/10.1038/nature13994

Zenklusen, D., Larson, D. R., & Singer, R. H. (2008). Single-RNA counting reveals alternative
modes of gene expression in yeast. *Nature Structural & Molecular Biology*, *15*(12), 1263–1271.
https://doi.org/10.1038/nsmb.1514

Zhang, F., Biswas, M., Massah, S., Lee, J., Lingadahalli, S., Wong, S., Wells, C., Foo, J., Khan,
N., Morin, H., Saxena, N., Kung, S. H. Y., Sun, B., Parra Nuñez, A. K., Sanchez, C., Chan, N.,
Ung, L., Altıntaş, U. B., Bui, J. M., … Lallous, N. (2023). Dynamic phase separation of the
androgen receptor and its coactivators key to regulate gene expression. Nucleic Acids Research,
51(1), 99–116. https://doi.org/10.1093/nar/gkac1158

Zhou, T., & Feng, Q. (2022). Androgen receptor signaling and spatial chromatin organization in
castration-resistant prostate cancer. Frontiers in Medicine, 9.
https://www.frontiersin.org/articles/10.3389/fmed.2022.924087

Figure 1

Figure 1. ERa binding and ligand-induced gene expression of *TFF1* and *TFF3* change over the course of estrogen signaling

A. Schematic depicting TFF1 locus, UCSC genome browser snapshots showing the binding of ERα, H3K27ac status, H3K4me3 signal, and Gro-seq signal for robustly E2-induced TFF1 locus. First, second and third ERα ChIP-seq and Gro-seq tracks are from vehicle-treated, E2-1h and E2-3h in WT cells, respectively.

B. qRT-PCR showing the changes in expression of TFF1 and TFF3 genes during the E2 signaling time course. Error bars denote SEM from four biological replicates. Each dot represents a replicate. p-values were calculated by Student's two-tailed unpaired t-test, and the significance is represented as: *** denotes $p < 0.001$,** denotes $p < 0.01$,* denotes $p < 0.05$, ns denotes $p > 0.05$.

Figure 2

Figure 2. TFF1 and TFF3 expressions show opposite trends during the E2 signaling time-course

A. 60X Representative images from single molecule RNA FISH experiment showing transcripts for TFF1 and TFF3. The probe was designed against the unspliced RNA containing the intronic region. The scale bar is 5 microns.

B. The mean RNA numbers are depicted. These are counted using an in-house MATLAB code which uses the DAPI-stained nuclei as the mask to count the RNA present in the nucleus. The graph shows the mean of means from three different repeats of the experiment, and error bars denote SEM (n= 665, N=3). p-values were calculated by Student's two-tailed unpaired t-test, and the significance is represented as: *** denotes $p < 0.001$,** denotes $p < 0.01$,* denotes $p < 0.05$, ns denotes $p > 0.05$.

C. Scatter plots showing the distribution of InTFF1 and InTFF3 on a cell-by-cell basis (n= 665, N=3). The absolute RNA numbers are combined from three different repeats. Density plots have been used to clearly visualize overlapping data points.

D. 60X Representative images from single molecule RNA FISH experiment showing transcripts for InTFF1 and ExTFF1. Scale bar is 5 microns.

E. The mean RNA numbers for InTFF1 and ExTFF1 are depicted. Separate probes were used to target unspliced (InTFF1) and mature (ExTFF1) RNA. These are counted using an in-house MATLAB code which uses the DAPI-stained nuclei as the mask to count the intronic RNA present in the nucleus and a free-drawn region to designate the cell to count the exonic RNA present in the nucleus as well as the cytoplasm. The graph shows the mean of means from three different repeats of the experiment, and error bars denote SEM (n>360, N=3). p-values were calculated by Student's two-tailed unpaired t-test, and the significance is represented as: *** denotes $p < 0.001$,** denotes $p < 0.01$,* denotes $p < 0.05$, ns denotes $p > 0.05$.

F. The mean RNA numbers for InTFF3 and ExTFF3 are depicted. Separate probes were used to target unspliced (InTFF3) and mature (ExTFF3) RNA. The graph shows the mean of means from three different repeats of the experiment, and error bars denote SEM (n>210, N=3). p-values were calculated by Student's two-tailed unpaired t-test, and the significance is represented as: *** denotes $p < 0.001$,** denotes $p < 0.01$,* denotes $p < 0.05$, ns denotes $p > 0.05$.

G. Violin plots showing the ratio of intronic to exonic TFF1 counts are depicted. The graph shows the distribution of ratios combined from three different repeats (n>360, N=3). p-values were calculated by the Mann-Whitney test, and the significance is represented as: *** denotes $p < 0.001$,** denotes $p < 0.01$,* denotes $p < 0.05$, ns denotes $p > 0.05$.

H. Violin plots showing the ratio of intronic to exonic TFF3 counts are depicted. The graph shows the distribution of ratios combined from three different repeats (n>210, N=3). p- values were calculated by the Mann-Whitney test, and the significance is represented as: *** denotes p < 0.001,** denotes p < 0.01,* denotes p < 0.05, ns denotes p > 0.05.

Figure 3

Figure 3. Enhancer looping does not account for the differential expression of TFF1 and TFF3 genes

A. 4C-seq plot at TFF1 enhancer viewpoint, the interaction with the promoter is highlighted in yellow. The plot is overlaid with H3K27ac, ERα ChIP signal, and gene annotations.

B. Genome browser snapshot of TFF1 region depicting ERα binding in WT lines. The first, second and third ERα ChIP-seq tracks are from WT cells that are vehicle-treated, E2-1h, and E2-3h, respectively. Blue highlighted regions represent the ΔTFF1e region.

C. The mean RNA numbers for InTFF1 in WT (unshaded) and ΔTFF1e (shaded) MCF7 cells are depicted. The mean of means are shown, and error bars denote SEM from three repeats (n>650, N=3 for each WT and delete line). p-values were calculated by Student's two-tailed unpaired t-test, and the significance is represented as: *** denotes $p < 0.001$,** denotes $p < 0.01$,* denotes $p < 0.05$, ns denotes $p > 0.05$.

D. The mean RNA numbers for InTFF3 in WT (unshaded) and ΔTFF1e (shaded) MCF7 cells are depicted. The mean of means are shown, and error bars denote SEM from three repeats (n>650, N=3 for each WT and delete line). P-values were calculated by Student's two-tailed unpaired t-test, and the significance is represented as: *** denotes $p < 0.001$,** denotes $p < 0.01$,* denotes $p < 0.05$, ns denotes $p > 0.05$.

E. The mean RNA numbers for InTFF1 and InTFF3 in ΔTFF1e MCF7 cells are depicted. The mean of means are shown, and error bars denote SEM from three repeats (n>880, N=3). p-values were calculated by Student's two-tailed unpaired t-test, and the significance is represented as: *** denotes $p < 0.001$,** denotes $p < 0.01$,* denotes $p < 0.05$, ns denotes $p > 0.05$.

F. Ratio of InTFF in WT MCF7 to InTFF in ΔTFF1e MCF7 are depicted. The ratio was obtained by dividing the absolute RNA counts of the WT line by delete lines performed on different days but in the same order (replicate one of WT divided by replicate one of ΔTFF1e). The mean of means are shown, and error bars denote SEM from three repeats (n>650, N=3 for each WT and delete line). p-values were calculated by Student's two-tailed unpaired t-test, and the significance is represented as: *** denotes $p < 0.001$,** denotes $p < 0.01$,* denotes $p < 0.05$, ns denotes $p > 0.05$.

Figure 4

Figure 4. Nuclear levels of the transcription factor ERα dictate the extent of expression of TFF1 and TFF3 genes

A. Representative images showing smFISH for InTFF1 and InTFF3 in combination with immunofluorescence for ERα (along with chromatin retention assay). The blue circle denotes a cell with high ERα and low TFF1 and TFF3, the red circle denotes a cell with medium ERα, high TFF1, and low TFF3, while the cyan circle denotes a cell with low ERα and high TFF3. The scale bar is 5 microns.

B. Histogram representing the distribution of ERα mean intensities in cells under induced, E3-1hr, and E2-3hr conditions (n=210). Intensities at 1hr are the highest while they shift to the left at 3hr and are lowest in uninduced cells. This is plotted from one experimental repeat out of three repeats, as ERα intensities will vary from one immunofluorescence experiment to another.

C. Cumulative histogram representing the distribution of ERα mean intensities in cells under induced, E3-1hr, and E2-3hr conditions (n=210). p-values were calculated by the Mann-Whitney test, and the significance is represented as: *** denotes $p < 0.001$,** denotes $p < 0.01$,* denotes $p < 0.05$, ns denotes $p > 0.05$.

D. ERα intensities were sorted into three categories, namely low (intensities between 0-450 A.U.), mid (intensities between 450-1200 A.U.) and high (intensities between 1200-2100 A.U.). The mean and SEM of transcript count for InTFF1 in the three categories under uninduced, E2-1hr, and E2-3hr were plotted (n=210). Low and mid categories show the highest TFF1 mean. p-values were calculated by the Mann-Whitney test, and the significance is represented as: *** denotes $p < 0.001$,** denotes $p < 0.01$,* denotes $p < 0.05$, and ns denotes $p > 0.05$. This is plotted from one experimental repeat out of three repeats, as ERα intensities will vary from one immunofluorescence experiment to another.

E. ERα intensities were sorted into three categories, namely low (intensities between 0-450 A.U.), mid (intensities between 450-1200 A.U.), and high (intensities between 1200-2100 A.U.). The mean and SEM of transcript count for InTFF3 in the three categories under uninduced, E2-1hr, and E2-3hr were plotted. The low category shows the highest TFF3 mean. p-values were calculated by the Mann-Whitney test, and the significance is represented as: *** denotes $p < 0.001$,** denotes $p < 0.01$,* denotes $p < 0.05$, ns denotes $p > 0.05$. This is plotted from one experimental repeat out of three repeats, as ERα intensities will vary from one immunofluorescence experiment to another.

F. 3D plot representing the distribution of ERα, InTFF1, and InTFF3 on a cell-by-cell basis shows that cells with lower levels of ERα show higher counts for InTFF3. This is plotted from

one experimental repeat out of three repeats, as ERα intensities will vary from one immunofluorescence experiment to another (n=210).

Figure 5

Figure 5. ERα mediated phase separation could govern the differential expression of TFF1 and TFF3 genes

A. Mean transcript counts for InTFF1 and InTFF3 in control, and 3% 1,6-Hexanediol treated cells post 30 minutes of E2-induction. The mean of means are shown, and the error bars denote SEM from three repeats (n>880, N=3). p-values were calculated by Student's two-tailed unpaired t-test, and the significance is represented as: *** denotes $p < 0.001$,** denotes $p < 0.01$,* denotes $p < 0.05$, ns denotes $p > 0.05$.

B. Boxplots showing DESeq2 normalized counts for low expressing, moderately expressing, and highly-expressing genes in the vehicle, E2-40m, and E2-160m respectively (left). Boxplots showing DESeq2 normalized counts for genes near low expressing, moderately expressing, and highly expressing genes in the vehicle, E2-40m, and E2-160m, respectively (right). The p-values in the boxplots were calculated using the Wilcoxon rank-sum test. The boxplots depict the minimum, first quartile, median, third quartile, and maximum values, along with outliers.

C. Model depicting the signaling under uninduced, E2-1hr, and E2-3h conditions- 1. Activation of TFF1 locus by ligand induction. 2. During the active phase, liganded ERα binds on enhancer and promoter. Together these in 3D interact, manifesting as ERα puncta, which results in robust expression of target genes, 3. The phase-separated condensate leads to the sequestration of transcriptional machinery. 4. This favours intra-TAD enhancer: promoter interactions to facilitate robust gene transcription. Upon ERα degradation at 3h, these condensates disappear, and transcriptional machinery becomes available to other promoters leading to their somewhat increased gene transcription.

A

B

Supp. Figure 1

Supp. Figure 1. Experimental design

A. Schematic depicting the experimental design. Cells were cultured in complete media for 24h, followed by stripping for 3 days. Finally, cells were induced for different durations with E2/Vehicle followed by different assays like smFISH, 4C-seq, smFISH-IF, etc.

B. Zoomed in the region around the *TFF3* gene, UCSC genome browser snapshots showing the binding of ERα and H3K4me3 signal. The first, second and third ERα ChIP-seq tracks are in vehicle-treated, E2 treated for 1h, and E2 treated for 3h in WT cells, respectively.

A   B

Supp. Figure 2. *TFF1* **and** *TFF3* **expressions show opposite trends during the E2-signaling time-course**

A. Violin plot showing the ratio of total intronic to absolute exonic *TFF1* counts are depicted. Absolute exonic counts are calculated by subtracting total intronic transcripts from total exonic transcripts. The graph shows the distribution of ratios combined from three different repeats. Error bars denote SEM. p-values were calculated by the Mann-Whitney test, and the significance is represented as: *** denotes $p < 0.001$, ** denotes $p < 0.01$, * denotes $p < 0.05$, and ns denotes $p > 0.05$.

B. Violin plot showing the ratio of total intronic to absolute exonic *TFF3* counts are depicted. Absolute exonic counts are calculated by subtracting total intronic transcripts from total exonic transcripts. The graph shows the distribution of ratios combined from three different repeats. Error bars denote SEM. p-values were calculated by the Mann-Whitney test, and the significance is represented as: *** denotes $p < 0.001$, ** denotes $p < 0.01$, * denotes $p < 0.05$, and ns denotes $p > 0.05$.

Supp. Figure 3

Supp. Figure 3. *TFF1* **enhancer does not change target promoters during signalling time-course**

4C-seq plot at *TFF1* enhancer viewpoint, the interaction with the promoter is highlighted in yellow. The plot is overlaid with H3K27ac, ERα ChIP signal, and gene annotations. There is no substantial interaction between the enhancer and *TFF3* locus at any time point, while the interaction between the enhancer and *TFF1* locus increases at 1hr and decreases at 3hr.

Supp. Figure 4

Supp. Figure 4. Levels of ERα in the nucleus dictate the extent of *TFF1* and *TFF3* inductions

A. Representative images from smFISH experiments showing In*TFF1* and GAPDH (top panel) and In*TFF1* and In*TFF3* (bottom panel) in cells overexpressing ERα-GFP. Yellow circles show cells that are GFP positive, and the transcripts associated with them. Note the visibly fewer *TFF1*, and *TFF3* transcripts in the ERα-GFP positive cells, while GAPDH transcripts remain indistinguishable. This shows that ERα-GFP overexpression specifically affects the transcription of E2-regulated genes like *TFF1* and *TFF3* and not a housekeeping gene like GAPDH. This is further quantified in B and C.

B. Scatter plots showing the distribution of ERα-GFP intensities with GAPDH or In*TFF1* transcript counts on a cell-by-cell basis from the experiment in the first row of A. Cells with high ERα-GFP expression can have high GAPDH expression, but such cells necessarily have low *TFF1* counts.

C. Scatter plots showing the distribution of ERα GFP intensities with In*TFF1* or In*TFF3* counts on a cell-by-cell basis from the experiment in the second row of A. As in B, high ERα-GFP expression leads to low expression of E2-regulated genes like *TFF1* or *TFF3*.

D. Representative images showing smFISH for In*TFF1* and In*TFF3* in cells overexpressing EGFP-C1. Yellow circles show cells that are GFP positive, and the transcripts associated with them. Just expressing EGFP in the same backbone as the ERα-GFP plasmid, does not downregulate *TFF1*, and *TFF3* expression; indicating that the downregulation effect is specific to ERα overexpression and not a generic effect of cell transfection. This is quantified in E.

E. Scatter plots showing the distribution of EGFP-C1 intensities with In*TFF1* or In*TFF3* transcript counts on a cell-by-cell basis. There is no obvious correlation between the levels of EGFP-C1 and *TFF1* or *TFF3*, and a cell with high EGFP levels may well have large *TFF1* or *TFF3* transcript counts.

Supp. Figure 5

Supp. Figure 5. LLPS perturbation down-regulates *TFF1* but supports *TFF3* expression

Mean transcript counts for In*TFF1* and In*TFF3* in control, and 3% 1,6-Hexanediol treated cells in the absence of E2-induction show no difference in *TFF1* counts on 1,6-Hexanediol treatment. The bar denotes the mean of means from two repeats, and the error bars denote SEM. p-values were calculated by Student's two-tailed unpaired t-test, and the significance is represented as: *** denotes p < 0.001, ** denotes p < 0.01, * denotes p < 0.05, ns denotes p > 0.05.

Supp. table 1- In*TFF3* probe sequence

Probe Sequence (5' to 3')	Probe sequence name
catctttccttttctttgt	Int_*TFF3*_1
ctgaagtcagggtttttctc	Int_*TFF3*_2
taaatgaaccttgtgcctgc	Int_*TFF3*_3
ggcagggttgctatgcaatg	Int_*TFF3*_4
acttgggttaatggagcagc	Int_*TFF3*_5
ataaaagccatgccctcatt	Int_*TFF3*_6
tttgaccctgctcagaagac	Int_*TFF3*_7
cgtaattgcttatcctcctg	Int_*TFF3*_8
tcttacattcccttctcttg	Int_*TFF3*_9
ggaaaatggcaatcacgtgc	Int_*TFF3*_10
cagaagttcactgcgtttca	Int_*TFF3*_11
caagcctctatgtctgtatg	Int_*TFF3*_12
ccccaatattttcatgtttc	Int_*TFF3*_13
ctgaattgttgagcatgtcc	Int_*TFF3*_14
tgcttttcacttttttctgc	Int_*TFF3*_15
cgccatagagatcattgcaa	Int_*TFF3*_16
aaaccgtgacaggagaggca	Int_*TFF3*_17
cttgcccctttgttttaaat	Int_*TFF3*_18
ccttcattgtttgaccaaca	Int_*TFF3*_19
cccttgaacccagaaacaaa	Int_*TFF3*_20
cttctcgggaagtcacatac	Int_*TFF3*_21
catttgcctaagtgtctttc	Int_*TFF3*_22
gtctgttaaatgctcagctc	Int_*TFF3*_23
ccttaggaagatgcactttc	Int_*TFF3*_24
cgagttcaaccactgctgaa	Int_*TFF3*_25
agtgtgtttgcttcactttg	Int_*TFF3*_26
agcccagaaaggacagggag	Int_*TFF3*_27
ttggaacaggtgtgtgtgtg	Int_*TFF3*_28
taagtgtgaagcctctgctc	Int_*TFF3*_29
aggggacagaagaggacagc	Int_*TFF3*_30
tgcgatgccatcattttgaa	Int_*TFF3*_31
gttactaaatcagttccctc	Int_*TFF3*_32
gtttccacttctctcaagaa	Int_*TFF3*_33
ggccgatttgcaaaaacagc	Int_*TFF3*_34
ggggatgccacaaatgacaa	Int_*TFF3*_35
cagagagagggagaggagat	Int_*TFF3*_36
cacgatgtggggcagctaag	Int_*TFF3*_37
agatggcttatagccaatcg	Int_*TFF3*_38
tgagctggcacttggacatg	Int_*TFF3*_39

tggatgctggttccaggaag	Int_*TFF3*_40
cctaggacaaagggcaagag	Int_*TFF3*_41
catcattgatcagggagccg	Int_*TFF3*_42
agtggagctgaaggaacagg	Int_*TFF3*_43
cactcagaggagagtgtagt	Int_*TFF3*_44
cagggatcaggcagataaca	Int_*TFF3*_45
tcgggttgctaagcaagata	Int_*TFF3*_46
atgggggaatagccaagatt	Int_*TFF3*_47
acagagcaaaggcttgtgtg	Int_*TFF3*_48

Supp. table 2- Ex*TFF3* probe sequence

Probe Sequence (5' to 3')	Probe sequence name
aaggggagaggtcagttca	Exn_*TFF3*_1
ccagacagtttttctctcc	Exn_*TFF3*_2
ctgcatgcctttgtcaagc	Exn_*TFF3*_3
ttgtttgcttggggaaggc	Exn_*TFF3*_4
tgtttgcacagctgctctg	Exn_*TFF3*_5
aggaggcctcatttatgca	Exn_*TFF3*_6
ctcaggactcgcttcatgg	Exn_*TFF3*_7
cagcatgcagagcgctctg	Exn_*TFF3*_8
aggacagcaaggccaggac	Exn_*TFF3*_9
cacgtactcctcagcagag	Exn_*TFF3*_10
acggcacactggtttgcag	Exn_*TFF3*_11
actccttggggggtgacatg	Exn_*TFF3*_12
cagggatcctggagtcaaa	Exn_*TFF3*_13
ggggcttgaaacaccaagg	Exn_*TFF3*_14
gtgcctcagaaggtgcatt	Exn_*TFF3*_15
caatcacagccgggcaagg	Exn_*TFF3*_16
gctgagatgaacagtgcct	Exn_*TFF3*_17
cgggagcaaagggacagaa	Exn_*TFF3*_18
actttcagcagaagcgctt	Exn_*TFF3*_19
gacatcaggctccagatat	Exn_*TFF3*_20
catgggacctttattcgtt	Exn_*TFF3*_21
gaagaactgtcctcgggtg	Exn_*TFF3*_22
acctcagaaagtctcaggc	Exn_*TFF3*_23
ccacgacgcagcagaaata	Exn_*TFF3*_24
cttagggagtcttttcctg	Exn_*TFF3*_25
atccttgcatgcactgcag	Exn_*TFF3*_26
gtgccagtctggattcaaa	Exn_*TFF3*_27
ctaggctttcctgtgacgt	Exn_*TFF3*_28

gaagcggcacttacagtgt	Exn_*TFF3*_29
gtatttttctgctttccc	Exn_*TFF3*_30
ttattggcttgctgtcttc	Exn_*TFF3*_31
ggaccactttggaagacag	Exn_*TFF3*_32
gtagcgagagtggttgtga	Exn_*TFF3*_33
acccaggaatagtacaagt	Exn_*TFF3*_34
gcctgtttgttcataatgt	Exn_*TFF3*_35

Supp. table 3- In*TFF1* probe sequence

Probe Sequence (5' to 3')	Probe sequence name
acagagcaggaagaagcacg	Int_*TFF1*_1
tgacacttgggaggattgta	Int_*TFF1*_2
ccaacacttcctctttgaaa	Int_*TFF1*_3
aaagcactattctgagaccc	Int_*TFF1*_4
gggagatgttggcatgaaca	Int_*TFF1*_5
catttgggcctatctggatg	Int_*TFF1*_6
caggaacctgtaggagaggg	Int_*TFF1*_7
atgagagaggtggctttgac	Int_*TFF1*_8
cagggaatatgggttgccag	Int_*TFF1*_9
caaccgatccacttgacaac	Int_*TFF1*_10
ttgagatgcaaacacttccc	Int_*TFF1*_11
atccattctgcaggtaaagg	Int_*TFF1*_12
cacacagcatttcctgacag	Int_*TFF1*_13
aaagtgcaagtcgcagatgc	Int_*TFF1*_14
aataagttattcagctcccc	Int_*TFF1*_15
ccaccttgaaactgtaccta	Int_*TFF1*_16
aacgtagaaagccctttctc	Int_*TFF1*_17
ctctggaaacccttgctttg	Int_*TFF1*_18
ccacagcagagatcaagagg	Int_*TFF1*_19
agttcgttctgtacaccgag	Int_*TFF1*_20
aaagacgctctgagccttga	Int_*TFF1*_21
gtaacaggatcatggacctg	Int_*TFF1*_22
ttccacacacgaatgcacta	Int_*TFF1*_23
ctttgtctttgtcgcatgat	Int_*TFF1*_24
tcagggaggaagatttccag	Int_*TFF1*_25
tggtgagggaggatcatcat	Int_*TFF1*_26
cactgtgggggtcaagagag	Int_*TFF1*_27
cgacgtgaaggtgatcatcg	Int_*TFF1*_28
tgggaaggatccgtgttcag	Int_*TFF1*_29
gtctttggagaaagtgttcc	Int_*TFF1*_30

cttgtctgctttgctatcag	Int_*TFF1*_31
ataggaggggaggaactcag	Int_*TFF1*_32
agcaggggggtgaaagaggag	Int_*TFF1*_33
taggatctggtggttgacag	Int_*TFF1*_34
cgcggaatcaaaggtctcag	Int_*TFF1*_35
tctaacaccagagcttcagg	Int_*TFF1*_36
cgcaggatcttatgcaagag	Int_*TFF1*_37
agaggaggaagacacgtgga	Int_*TFF1*_38
aagcagggcagtgaggaaag	Int_*TFF1*_39
cagagggtacaagctgtgaa	Int_*TFF1*_40
attgcaggtaaggtgagagg	Int_*TFF1*_41
gtgtgttcaggaggtggaaa	Int_*TFF1*_42
ttcaccaggtgaagatggag	Int_*TFF1*_43
aggttgaacaaagcaggggg	Int_*TFF1*_44
atgtgcgagacaacgcaagg	Int_*TFF1*_45
tctgatggaggagaaaggca	Int_*TFF1*_46
ttgtggactttgcatctttc	Int_*TFF1*_47
gctcagggaaaatgcaaggt	Int_*TFF1*_48

Supp. table 4- Ex*TFF1* probe sequence

Probe Sequence (5' to 3')	Probe sequence name
gcgacccgagtcagggat	Exn_*TFF1*_1
tgcctcctctctgctccaa	Exn_*TFF1*_2
cttgttctccatggtggcc	Exn_*TFF1*_3
caggaccagggcgcagatc	Exn_*TFF1*_4
gccgagggccagcatggac	Exn_*TFF1*_5
tgtctgggcctcggccagg	Exn_*TFF1*_6
ggggccactgtacacgtct	Exn_*TFF1*_7
ccacaattctgtctttcac	Exn_*TFF1*_8
gagggcgtgacaccaggaa	Exn_*TFF1*_9
cagcccttatttgcacact	Exn_*TFF1*_10
cacgaacggtgtcgtcgaa	Exn_*TFF1*_11
gatagaagcaccaggggac	Exn_*TFF1*_12
gagggacgtcgatggtatt	Exn_*TFF1*_13
tcgtataaaaaggccatac	Exn_*TFF1*_14
attctaaatcttcagaacc	Exn_*TFF1*_15
gtcttaaatgacttttcta	Exn_*TFF1*_16
atgctgatcagagcctctg	Exn_*TFF1*_17
tgtgtaaaggcatagctgg	Exn_*TFF1*_18
caccactggcggccgtgac	Exn_*TFF1*_19

gactcaggctaccccattg	Exn_*TFF1*_20
ttcctggacctgaatgcag	Exn_*TFF1*_21
tcccttagccctgcctttc	Exn_*TFF1*_22
gtctaaaattcacactcct	Exn_*TFF1*_23
caggcagatccctgcagaa	Exn_*TFF1*_24
ggacggcaccgcgtcagga	Exn_*TFF1*_25
tgggactaatcaccgtgct	Exn_*TFF1*_26
ggtggaggtggcagccgag	Exn_*TFF1*_27
agaagcgtgtctgaggtgt	Exn_*TFF1*_28
tgtgagccgaggcacagct	Exn_*TFF1*_29
gtcagagcagtcaatctgt	Exn_*TFF1*_30
ggccaattttgagtagtca	Exn_*TFF1*_31
atcgatctcttttaatttt	Exn_*TFF1*_32

Supp. table 5- 4C Seq oligo sequences

Sequence name	Sequence (5' to 3')
4C *TFF1* Enhancer barcode1 F	AATGATACGGCGACCACCGAGATCTACACTCTTTCCCTACACGACG CTCTTCCGATCTAGAAGTCCTTGAGTGGGGAGATGATC
4C *TFF1* Enhancer barcode2 F	AATGATACGGCGACCACCGAGATCTACACTCTTTCCCTACACGACG CTCTTCCGATCTAGCGACCCTTGAGTGGGGAGATGATC
4C *TFF1* Enhancer barcode3 F	AATGATACGGCGACCACCGAGATCTACACTCTTTCCCTACACGACG CTCTTCCGATCTCCAATTCCTTGAGTGGGGAGATGATC
4C *TFF1* R	CAAGCAGAAGACGGCATACGAGATAGGTCGCAGTGACTGGAGTTC AGACGTGTGCTCTTCCGATCTCCGCTGTTGCGACACACCAGAGAGC

Supp. table 6- qPCR oligo sequences

Sequence name	Sequence (5' to 3')
TFF1 US F	GCCCTTATTTGCACACTGGG
TFF1 US R	CATCAAGGAATTCAGCCCAC
TFF3 US F	CAATCACAGCCGGGCAAGGGT
TFF3 US R	GGGAGGAGGCAGCACTAGGT
GAPDH F	CGCTCTCTGCTCCTCCTGTT
GAPDH R	CCATGGTGTCTGAGCGATGT