A novel iPSC model of Bryant-Li-Bhoj neurodevelopmental/neurodegenerative syndrome demonstrates the role of histone H3.3 in chromatin dynamics, neuronal differentiation, and maturation

Annabel K. Sangree[1,2]*, Rajesh Angireddy[1]*#, Janardhan P. Bhattarai[4], Yingqi Wang[4], Laura M. Bryant[1,3],Elisa A. Waxman[5], Dana E. Layo-Carris[1], Emily E. Durham[1], Kaitlin A Katsura[1], Emily E. Lubin[1,2], Xiao Min Wang[1], Kelly J. Clark[1,2], Minghong Ma[4], Elizabeth J. Bhoj[1,2]#

Author affiliations:
1. Department of Human Genetics, Children's Hospital of Philadelphia, Philadelphia, PA, USA
2. Perelman School of Medicine, University of Pennsylvania, Philadelphia, PA, USA
3. Steve and Cindy Rasmussen Institute for Genomic Medicine, Nationwide Children's Hospital, Columbus, OH, USA
4. Department of Neuroscience, University of Pennsylvania Perelman School of Medicine, Philadelphia, PA 19104, USA
5. Center for Cellular and Molecular Therapeutics, The Children's Hospital of Philadelphia, Philadelphia, PA, USA

* Equal contributors

Co-corresponding author

Corresponding author contact: bhoje@chop.edu

ABSTRACT

Background

Bryant-Li-Bhoj neurodevelopmental syndrome (BLBS) is neurogenetic disorder caused by variants in *H3-3A* and *H3-3B,* the two genes that encode histone H3.3. Ninety-nine percent of individuals with BLBS show developmental delay/intellectual disability, but the mechanism by which variants in H3.3 result in these phenotypes is not yet understood, limiting the therapeutic interventions available to   individuals living with BLBS.

Methods

Here, we investigate how one BLBS-causative variant, *H3-3B* p.Leu48Arg (L48R), affects neurodevelopment using an induced pluripotent stem cell (iPSC) model differentiated to 2D neural progenitor cells (NPCs), 2D forebrain neurons (FBNs), and 3D dorsal forebrain organoids

35 (DFBOs). We employ a multi-omic approach in the 2D models to quantify the resulting changes

36 in gene expression and chromatin accessibility. We used immunofluorescence (IF) staining to

37 define the identities of cells in the 3D DFBO and whole-cell patch clamp to investigate the

38 electrophysiological properties of neurons in DFBOs.

40 **Results**

41 In the 2D systems, we found dysregulated gene expression and chromatin accessibility affecting

42 neuronal fate, adhesion, neurotransmission, and excitatory/inhibitory balance.

43 Immunofluorescence of DFBOs corroborated altered proportions of radial glia and mature

44 neuronal populations. Patch clamp recordings revealed decreased electrical activity in neurons

45 from L48R DFBOs compared to control DFBOs.

48 **Conclusions**

49 These data provide the first mechanistic insights into the pathogenesis of BLBS from a human-

50 derived model of neurodevelopment, which suggest that  H3.3 L48R increases *H3-3B*

51 expression, resulting in the hyper-deposition of H3.3 into the nucleosome which underlies

52 changes in gene expression and chromatin accessibility. Functionally, this causes dysregulation

53 of cell adhesion, neurotransmission, and the balance between excitatory and inhibitory

54 signaling. These results are a crucial step towards preclinical development and testing of

55 targeted therapies for this and related disorders.

57 **Keywords**: Histone H3.3, Mendelian neurodevelopmental disorder, chromatin, iPSC-derived

58 neurons, brain organoid, multi-omic analysis, translational neurogenetics

BACKGROUND

Neurodevelopmental disorders (NDDs) affect 1-8% of the population and are often the result of aberrant components of epigenetic machinery following a genetic insult in the earliest stages of neurodevelopment[1]. Mendelian disorders of epigenetic machinery (MDEMs), caused by germline variants in 85 genes encoding histone writers, readers, erasers, and chaperones, are associated with a wide spectrum of NDDs, highlighting the brain's vulnerability to transcriptional dysregulation[2]. Recently, variants in histones themselves emerged as causative in NDDs with the identification of *de novo* germline variants in the H4 encoding gene cluster[3,4] and the H3.3-encoding genes *H3-3A* and *H3-3B*[5–7].

Bryant-Li-Bhoj neurodevelopmental syndrome (BLBS, Online Mendelian Inheritance in Man ([OMIM](#)) #619720, #619721) is an NDD caused by heterozygous germline variants in *H3-3A* or *H3-3B*. Variants underlying BLBS occur throughout the coding sequences of *H3-3A* and *H3-3B* and are typically observed in just one individual[5,8 (add in Okur citation)]. Phenotypically, affected individuals present almost universally with neurologic and craniofacial features. The most prevalent neurological   feature is developmental delay/intellectual disability, which impacts 99% of published individuals with 51% of individuals exhibiting delayed or no attainment of sitting, 79% of individuals exhibiting delayed or no attainment of independent walking, and more than 40% of individuals not yet having attained their first word at >20 months[8]. From a neuroradiological standpoint, individuals most commonly have delayed myelination or hypomyelination, dysgenesis of the corpus callosum, and/or dilated ventricles[9]. As some individuals age, they develop an abnormal gait or progressive gait ataxia phenotype[5,7,8]. While no targeted therapeutics are available at this time, these neurodegenerative phenotypes suggest that BLBS could be amenable to therapeutic intervention to halt disease progression.

A mechanistic link has not yet been established between the physiologic function of H3.3 and its dysregulation in BLBS. H3.3 is a replication-independent histone that packages DNA and

regulates gene expression. It is preferentially deposited by the chaperones DAXX or HIRA at

enhancers, promoters, and gene bodies[10–18]. Because its deposition is not coupled to the cell

cycle, H3.3 plays a critical epigenetic role in post-mitotic neuronal populations in the brain,

especially in regulating transcription during learning and memory[19]. During early

neurodevelopment in mice, H3.3 accounts for an estimated 31% of the total H3 pool[20]   and

becomes the highly dominant form of H3 (>93%) in adult neurons[21]. Throughout all stages of

neurodevelopment, H3.3 plays a crucial role in establishing chromatin landscapes,

transcriptome, molecular identity, and axon connectivity in newly post-mitotic neurons[10].

The role of H3.3 in disease was first appreciated in cancer through the discovery of somatic

'oncohistone' mutations[22–24].   While extensive work has been done to elucidate the

mechanism of   somatic oncohistone mutations like Gly34Arg/Val, the impact of germline

missense variants, on neurodevelopment remains poorly understood.   .

Past studies into the effects of germline missense perturbation of H3.3 utilized cells from

affected individuals, overexpression systems, and mouse models[5,6,29]. Dermal fibroblasts from

affected individuals hyper-proliferate compared to age- and sex-matched controls[5]. *In silico*

predictions   suggest that certain mutants alter phosphorylation, glycosylation, methylation,

acetylation, and ubiquitination histone post translational modifications (hPTMS) *in cis*[7]. In mice,

three different variants at the same residue (p.Gly34Arg/Val/Trp) cause distinct developmental

defects. p.Gly34Arg mice have severe neurodegeneration, p.Gly34Trp mice have obesity and

bladder obstructions, and p.Gly34Val mice have intermediate phenotypes[29]. These models

capture the same phenotypic heterogeneity recently reported amongst individuals living with

BLBS, including those who share the same causative nucleotide level substitution[8]. These

studies highlight the importance of studying individual variants, as both affected individuals and

models display variable phenotypes. Moreover, there is a gap in knowledge as to how these

variants affect human development in affected organs, most strikingly the brain, highlighting the

necessity of interrogating the effect of BLBS-causing variants in human iPSCs.

Here, we profile the transcriptomic   ,   chromatin accessibility   , and cellular identity

changes induced by one variant observed in our cohort, *H3-3B* p.Leu48Arg (L48R). This variant

lies just beyond the N terminal tail, in the histone core[8], a region of H3.3 that is less well studied

in the context of disease. While the WT leucine has a hydrophobic side chain, arginine has a

positively charged side chain. This L48R substitution is predicted to increase the distance

between itself and other residues on H3.3, perhaps subtly opening the nucleosome at this

position (Fig S1A). The individual harboring the L48R variant presents with neurologic features

that include developmental delay/intellectual disability, microcephaly, hypomyelination,

hypotonia that transitioned to hypertonia, and seizures.

To elucidate how this variant affects neurodevelopment, we utilized 2D and 3D models, allowing

for both a detailed molecular analysis of gene expression and chromatin accessibility, as well as

a platform to study the structural and cellular dynamics of BLBS in a more physiologically

relevant context. In our 2D models of neural progenitor cells (NPCs) and forebrain neurons

(FBNs) we elucidated global dysregulation in gene expression and chromatin accessibility of

genes important for cell adhesion, axon development, and neurotransmission.

The   3D organoid models   , which faithfully recapitulate developing neuroectoderm with the

radial organization of NPC cells into structures that recapitulate the ventricular zone (VZ),

subventricular zone (SVZ), and cortical plate (CP)[30], recapitulated our findings of altered

differentiation capacity of progenitor cell types   Taken together, we find that H3.3 L48R

perturbs neural differentiation   and neuronal maturation in this iPSC model of brain

development. Our results lay the foundation for future studies on BLBS and therapeutic

interventions.

METHODS

Maintenance, editing, and differentiation of 2D cells

iPSCs

iPSCs were edited by the CHOP iPSC core as described[31]. The cells were single cell cloned

and frozen at an early passage. Edited and isogenic control iPSCs were maintained in

mTeSR™ Plus (STEMCELL Technologies 100-02 76) on 10cm plates coated with 1ug

Matrigel® hESC-Qualified Matrix (Corning 354277) and three technical replicates were collected

at passage 30 for RNA-sequencing.

NPCs

Cells were differentiated to NPCs using STEMdiff™ SMADi Neural Induction Kit following the

monolayer culture protocol (STEMCELL Technologies 08582). Briefly, for each well, 2 million

passage 31 iPSCs were resuspended in STEMdiff™ Neural Induction Medium + SMADi + 10

μM Y-27632 and plated in each well of a 6-well plate coated with 1ug of Matrigel. Cells well

given a full medium change with STEMdiff™ Neural Induction Medium + SMADi except on days

when they were passaged (approximately every seven days). Cells were collected at passage

four post differentiation for RNA-sequencing. At the end of differentiation, cells were

cryopreserved in STEMdiff™ Neural Progenitor Freezing Medium. Three separate

differentiations were conducted by two different investigators to control for inter-differentiation

variability.

FBNs

Two biological replicates of NPCs (passage 4) were thawed and differentiated to forebrain

neurons using STEMdiff™ Forebrain Neuron Differentiation Kit (STEMCELL Technologies

08600). Briefly, NPCs were plated in PLO-Laminin coated 6 well plates (1.25 million cells per

well) in STEMdiff™ Neural Induction Medium + SMADi, and after 24 hours, media was changed

to STEMdiff™ Forebrain Neuron Differentiation medium. Cells were cultured in the same

medium until they reached 80-90% confluency and passaged on to 12 well plates coated with

PLO-laminin at a density of 175,000 cells per well in STEMdiff™ Forebrain Neuron Maturation

medium. The FBNs were matured up to three weeks and collected for IF (2 biological replicates)

and RNA-seq (3 technical replicates)

Flow cytometry

Cells were harvested using Accutase, diluted with DPBS, and centrifuged at 3000 xg. They

were then resuspended in 1.6% paraformaldehyde and incubated for 30 minutes at 37 °C with

agitation. After a single wash with DPBS, the cells were resuspended in FACS Buffer (DPBS

with 0.5% BSA and 0.05% sodium azide) and stored at 4 °C until flow cytometry analysis. For

the flow cytometry analysis, cell permeabilization was performed with saponin buffer (diluted to

1× in water; Biolegend, 421,022), followed by a 1-hour incubation with antibodies diluted in

saponin buffer: Forse-1 (DSHB, 1:100), NCAD (Cell Signaling #14215, 1:300), and an IgM

isotype control (Santa Cruz, sc-3881, 1:100). The cells were then washed with either saponin

buffer or FACS buffer and incubated for 1 hour with secondary antibodies: goat anti-rabbit Alexa

488 (Jackson ImmunoResearch, 111–545-144, 1:500) or goat anti-mouse IgM-conjugated Alexa

488 (Thermo Fisher Scientific, A21042, 1:500). Finally, the cells were washed again and

resuspended in FACS buffer. Analysis was conducted using a CytoFLEX flow cytometer

(Beckman Coulter) with the FlowJo software program (BD).

RNA sequencing

RNA extraction, library prep & sequencing

Total RNA (3 replicates of each cell type) was extracted from iPSCs and NPCs using the Zymo

Direct-zol RNA Miniprep (Cat # R2050 or from FBNs with the Zymo Direct-zol RNA Microprep

(Cat # R2062). QC was performed using the Gx Bioanalyzer/Agilent Tapestation for traces and

Qubit for concentration. Libraries were prepared using the Illumina Stranded Total RNA Prep,

Ligation with Ribo-Zero Plus. 200ng of cDNA libraries were amplified using 12 cycles of PCR

and sequenced on a v1.5_SP (200 cycle) flow cell with the NovaSeq6000. Reads were

demultiplexed using Dragen Demultiplexing.

Data analysis

At least 15E6 million r   eads were aligned to Homo_sapiens.GRCh38.cdna.all using

kallisto[32](v0.50.1). Replicates were averaged and d   ifferential expression was calculated

using DeSeq2[33](v1.44.0) in R v 4.4.0 using RStudio 2024.04.1+748. Volcano plots were made

using EnhancedVolcano[34](v1.22.0). We defined genes using adjusted p-values (Benjamini and

Hochberg corrected) and log2fold changes output by DESeq2 with significantly altered

expression with an adjusted p-value ≤ 0.05 and log fold change with an absolute value of ≥

1. GO analyses were performed using clusterProfiler[35] (v4.12.0) using biological pathway

terms. Revigo[36] was used to summarize GO terms and these terms were clustered using a

distance matrix and hierarchical clustering. Each cluster was labeled with one representative

term, obtained by selecting the term within each cluster with the largest LogSize, the Log10

value of the number of annotations for GO Term ID in selected species in the European

Bioinformatics Institute Gene Ontology Annotation database. The list of astrocyte specific genes

used to analyze the FBN data was obtained from PanglaoDB[37],   filtered for human, and a

specificity score ≤ 0.5 (meaning that these genes are expressed in cells other than astrocytes

less than or equal to 50% of the time).

Western Blots

Histones were extracted using the Abcam Histone Extraction Kit (ab113476). Protein

concentration was quantified using the Pierce™ BCA Protein Assay Kit (A65453) and run using

the SkanIt software v6.1 by measuring the absorbance at 550nm. 2ug of histone extract was

loaded per reaction and   separated by electrophoresis using NuPAGE™ 4-12% Bis-Tris

Protein Gel (ThermoFisher Scientific). Proteins were transferred to a polyvinylidene difluoride

(PVDF) membrane using an XCell SureLock Mini-Cell Electrophoresis System (ThermoFisher

Scientific). The membrane was blocked using 5% dry milk in Tris-Buffered Saline (Bio Basic,

Markham) with Tween-20 for 1 hour at room temperature. After briefly rinsing with Tris-Buffered

Saline (Bio Basic, Markham) with Tween-20, the membrane was incubated overnight at 4°C

with a previously validated primary antibody (Table 5). The membrane was then washed with

Tris-Buffered Saline with Tween-20 and incubated in secondary antibody diluted 1:10,000 (Goat

Anti-Rabbit IgG Polyclonal from LI-COR Biosciences) for 1 hour at room temperature. Blots

were visualized using the ODYSSEY® DLx (LI-COR Biosciences) and quantified using ImageJ.

ATAC-sequencing:

Library preparation & sequencing

ATAC-seq libraries were generated using the ATAC-seq Kit from Diagenode (Diagenode, A

Hologic Company) according to manufacturer instructions. Briefly, cells were centrifuged at 500

× g for 10 minutes using a fixed angle refrigerated centrifuge, followed by a wash with 500 µl of

cold 1x PBS and a second centrifugation at 500 × g for 10 minutes. Cells were counted and

approximately 50,000 cells were lysed using cold ATAC lysis + Digitonin (2%). Immediately after

lysis, cell nuclei were spun again at 500 × g for 10 minutes and separated from the lysis buffer.

Immediately following the nuclei extraction, the pellet was resuspended in transposase reaction

mix. The transposition reaction was carried out for 30 minutes at 37 °C. Directly following

transposition the samples were purified using the kit's provided columns. Following purification,

library fragments were amplified using 1x Amplification master mix and the following PCR

conditions: 72°C for 5 minutes, 98°C for 30 seconds, followed by thermocycling at 98°C for 10

seconds, 63°C for 30 seconds and 72°C for 1 minute for 5 cycles, in this step Unique Dual

Indexes Primer Pair were incorporated for multiplexed sequencing. To reduce amplification bias

after the first 5 cycles, qPCR was used to determine the number of PCR cycles needed to

produce the concentration required for sequencing. Each pre-amplified library was further

amplified for 6 cycles as determined by the qPCR step. Final libraries were bead purified

(Beckman Coulter), then assessed for size distribution and concentration using the Bioanalyzer

High Sensitivity DNA Kit (Agilent Technologies). The resulting libraries were pooled in equal

molarity, the pool was diluted to approximately 2 nM using 10 mM Tris-HCl, pH 8.5, denatured,

and loaded onto a P1-100 (2x50) flow cell on an Illumina NextSeq 6000 (Illumina, Inc.)

according to the manufacturer's instructions. De-multiplexed and adapter-trimmed sequencing

reads were generated using bcl2fastq.

Data processing and analysis

Reads were successfully aligned to *Homo* sapiens genome build Genome Reference

Consortium Human Build 38 (GRCh38) with botwie2[38] (v2.5.4). PCR duplicates were removed

using picard[39] (v3.2.0), mitochondrial regions were removed, and blacklisted regions were

identified using ENCODE (GRCh38_unified_blacklist.bed) and removed. Replicates were

merged and subset to 55E6 reads for both Control and L48R NPCs. Peaks were called using

macs3[40] (v3.0.1) with a q-value cutoff of 0.05. Peaks were annotated using

ChIPSeeker[36](v1.40.0). DiffBind[69](v3.14.0) was used to determine differential accessibility

between L48R and control peaks and plotted using the DiffBind   profile plot. For transcription

factor footprinting, reads were shifted to account for the Tn5 overhang using ATACshift from

deeptools[44], and motifs were identified using ATACseqTFEA[55] with human motifs

"best_curated_Human.rds". This packages collects motifs from jasper2018, jolma2013 and

cisbp_1.02 from package motifDB (v 1.28.0) and merged by distance smaller than 1e-9 calculated by

MotIV::motifDistances function (v 1.42.0). Genome browser plots were made using IGV(v2.17.4)

qPCR validation:

Total RNA was extracted from iPSC and NPCs using the Zymo Direct-zol RNA Miniprep (Cat #

R2050 or from FBNs with the Zymo Direct-zol RNA Microprep (Cat # R2062). The concentration

was quantified using the NanoDrop 8000 UV-Vis Spectrophotometer (Thermo Scientific). cDNA

was synthesized from 1ug of RNA using Superscript IV (Thermo Scientific). RT-qPCR reactions

were set up using cDNA, 2X PowerUp™ SYBR® Green Master Mix (Applied Biosystems), and

gene specific primers using GAPDH as an internal control. Samples were run in a 96-well plate

on the QuantStudio™ 3 System using the Comparative C$_T$ ($\Delta\Delta$C$_T$) method. The primers used

are listed in Table 6. Two representative primer pairs are presented below:

GAPDH F: GGCATCCTGGGCTACACTGA, R: GAGTGGGTGTCGCTGTTGAA

H3-3B F: TCGTTGGGCGGTGCTGGTTTTT R: TTAGGCCACTTCTCCGACCGCC

Dorsal Forebrain Organoids (DFBOs)

Forebrain organoids were generated using STEMdiff™ Dorsal Forebrain Organoid

Differentiation Kit (Cat: 08620) according to manufacturer instructions, and this protocol was

adopted from[45]. Briefly, embryoid bodies (EB) were generated in AggreWell™800 24-well plate

in a seeding medium with 10,000 cells per microwell. This method will generate 500 equalized

EBs per one well of a 24-well plate. After 24 hours, the media was changed to a Forebrain

Organoid formation medium, and the EB was maintained for five days. EBs were transferred to

ultra-low attachment 6-well plates (40 organoids per well) in organoid expansion medium.

Organoids were maintained in the same medium for a period of 25 days with frequent media

changes. After the neuroectoderm expansion, the media was changed to the Forebrain

Organoid Differentiation medium and maintained organoids until day 43 with frequent media

changes. For long-term maintenance of DFBO, the media was changed to a Forebrain Organoid

maintenance medium. For the analysis, we harvested organoids at day 25 (Neuroectoderm

expansion) and day 62 (DFBO maturation) and day 90 (Electrophysiology).

Embedding

Organoids were washed with D-PBS then fixed in 5mL fresh 4% paraformaldehyde (PFA)

solution and incubated overnight at 4°C. Organoids were washed in PBS-T and stored in PBS-T

at 4°C for 1 week. Organoids were then incubated in 30% sucrose solution and equilibrated

overnight at 4°C. Equilibration was complete once organoids were no longer floating in the 30%

sucrose solution. Organoids were then incubated at 37°C in porcine gelatin for 1 hour and

embedded in this solution in embedding molds. Six organoids were embedded per block and

allowed to solidify at RT and 4°C before being snap frozen in a dry ice/ethanol slurry (dry ice +

100% ethanol). Frozen samples were stored at -80°C until ready for cryosectioning.

Cryosectioning and Immunoflourescence Staining:

Blocks were removed from the -80°C freezer and allowed to equilibrate to -26°C in the cryostat

(Microm HM 505E, CHOP Pathology Core) for at least 30 minutes before sectioning. Sections

were cut at 16um using a high-profile blade (Leica Biosystems 14035838926) and mounted on

colorfrost Plus Microscope Slides (Fisher Scientific 12-550-17). Slides were stored at -80C until

staining. Slides were allowed to come to room temperature before staining. Hydrophobic

barriers were drawn around sections. Antigen retrieval using sodium citrate pH 6.0, was

performed for the CTIP2 staining. Slides were steamed in a food steamer for 20 minutes in

citrate buffer, washed with PBS-T at 37°C for 10 minutes to fully remove gelatin from slides,

blocked in a humidified chamber for 1 hour, and incubated with primary antibody overnight. The

next day, primary antibody solutions were removed, slides were washed 3 times with PBS-T

and incubated in secondary antibody at room temperature in a humified chamber for 2 hours.

After incubation, secondary antibody solution was removed, slides were washed 3 times with

PBS-T for 30 minutes, allowed to air dry for 5 minutes, then one drop of PermaFluor™ was

added to slides to add coverslip.

Image acquisition and quantification

Images for quantification were acquired using a Keyence BZ-X800 (Itasca, IL) or the Leica

STELLARIS 5 Confocal Microscope. 10x images taken of DFBOs were stitched together using

Keyence BZ-X Series Analysis Software. FIJI (v 2.14.0/1.54f) was used to quantify the fraction

of cells that were positive for cell fate markers. Images were separated by channel, converted to

binary, then particles were analyzed for each image. Positive cells for each marker were

normalized to the total number of cells counted (DAPI) in each image.

Patch Clamp Electrophysiology

Dorsal forebrain organoids at maturation day 90 were prepared for electrophysiological

recordings. Before recording, DFBOs were transferred to a recording chamber and held by a

nylon slice holder and continuously perfused with oxygenated artificial cerebrospinal fluid

(ACSF) containing the following (in mM): 126 mM NaCl, 2.5 mM KCl, 1.2 mM $MgSO_4$,

2.4 mM $CaCl_2$, 25 mM $NaHCO_3$, 1.4 mM NaH_2PO_4, 11 mM glucose, and 0.6 mM sodium L-

ascorbate) and continuously bubbled with 95% O_2 and 5% CO_2). Neurons were visualized with

a 40× water-immersion objective under an Olympus BX51WI upright microscope equipped with

epifluorescence. Recording pipettes were made from borosilicate glass (GC210F-10; Harvard

Apparatus) with a Flaming-Brown P-97 puller (Sutter Instruments; tip resistance 5–8 MΩ). The

pipette solution contained the following (in mM): 120 K-gluconate, 10 NaCl, 1 CaCl2, 10 EGTA,

10 HEPES, 5 Mg-ATP, 0.5 Na-GTP, and 10 phosphocreatine. Whole-cell patch-clamp

recordings were conducted in both current and voltage-clamp modes. Electrophysiological

recordings were controlled by an EPC-10 amplifier combined with Pulse Software (HEKA

Electronic) and analyzed using Igor Pro.

For a subset of neurons, intrinsic membrane properties were measured via current injections

under current-clamp mode. The resting membrane potential was determined when there was no

current injected. For the remaining properties, the cell membrane potential was kept at -70 mV.

The input resistance was calculated as the slope of the current-voltage (I-V) curve (current

steps from -100 pA with 10 pA increments). The action potential (AP) threshold was determined

as the point with the maximum curvature before the AP evoked by the minimum current step.

Statistics

Comparisons were made between Control and L48R cells for each timepoint. For RT-qPCR and

IF, statistics were calculated using Welch Two Sample t-test in R v 4.4.0 using RStudio

2024.04.1+748. A p-value ≤ 0.05 is denoted with *, p-value ≤ 0.01 is denoted with **, and p-

value ≤ 0.001 is denoted with ***. For RNA-seq, statistics were calculated using DESeq2[33], and

for ATAC-seq statistics were calculated using DiffBind[43]. No statistical method was used to

predetermine sample size.

RESULTS

L48R variant affects iPSC differentiation into 2D neural progenitor cells (NPCs) and 2D

forebrain neurons (FBNs)

We introduced the heterozygous c.146T>G *H3-3B* Leu48Arg (L48R) variant into CHOPWT14

iPSCs (from a Hispanic 46XX donor) using CRISPR/Cas9 homology directed repair (HDR)[31]

followed by single cell clone selection and validation by Sanger sequencing (Fig. 1A, Fig. S1A-

C). To validate   pluripotency, we performed RT-qPCR for *OCT4* and *NANOG* in these iPSCs

(Fig. 1B). We did not observe significant changes in the expression of these markers between

L48R and control cells, confirming that this variant did not have a large impact on the identity of

iPSCs.

To begin delineating the effects of the H3.3 L48R variant on brain development and neuronal

function, we differentiated L48R and isogenic control iPSCs into NPCs using a dual SMAD

inhibition protocol[46]. We found that the L48R NPCs had lower expression of   NPC markers

NES (encodes Nestin) and *SOX1* than   control (Fig. 1C), indicating that this variant affects

early neuronal differentiation. FACS analysis on control and L48R NPC stained with progenitor

markers Forse1 and NCAD indicates that the L48R mutation has no effect on the overall NPC

population (Fig. S1D-G). Next, we differentiated the NPCs to FBNs using a protocol that

generates a mixed population of 90% forebrain-type neurons (MAP2+ and TUJ1+) with about

10% glial cells (GFAP+) (Stem Cell Technologies #08600, #08605). We immunostained

FBNs at day 20 of maturation for the neuronal marker TUJ1 and glial marker GFAP. We

observed decreased TUJ1 and increased GFAP intensity, however neither of these reached

significance (Fig. S1H, I). Interestingly, there was no significant change in the total number of

cells expressing TUJ1, and a decrease in L48R cells expressing GFAP compared to control

(Fig. S1J). This suggests that the increased GFAP intensity might be due to an increase in

reactive astrocytes, rather than total cell number.   . Further staining indicated decreased

intensity   of MAP2+ neuronal extensions and   glutamate transporter VGLUT1 in the L48R

FBNs compared to isogenic controls (Fig. 1D, E). We found no change in the number of cells

expressing MAP2, or relative   *MAP2* mRNA expression measured via RT-qPCR between the

two conditions (Fig. 1F, Fig. S1K). We found a significant decrease in VGLUT1 staining and

a significant increase in *GFAP* mRNA levels in L48R FBNs compared to isogenic controls (Fig.

1F,G).   Taken together, we observed disruptions in the differentiation of L48R iPSCs to

NPCs, which impacted subsequent differentiation into FBNs.

Transcriptomic changes in L48R Neural progenitor cells affect differentiation into 2D

Forebrain Neurons

To understand the global transcriptomic changes induced by L48R at various stages of

differentiation   we performed bulk RNA-seq in   3 biological replicates of iPSCs, NPCs, and

FBNs   (Supplementary File 1). Principal component analysis (PCA) on the different cell types

identified three distinct clusters based on cell identity (Fig. S2A). We identified 96 genes with

significantly altered expression (p ≤ 0.05) in L48R iPSCs compared to the isogenic controls (Fig.

2A). Using an additional threshold of absolute value log2-fold change (LFC) ≥ 1, 37 genes were

significantly upregulated, while 20 were downregulated. Using a gene ontology (GO)

analysis, we found no significantly altered pathways in the L48R iPSCs compared to isogenic

controls, but did identify the upregulation of *CDKL5* (LFC = 1.46, adjusted p-value = 6.36E-09)

and *SCML2 (*LFC = 1.04, adjusted p-value = 4.98-E07)   both of which are genes important for

cell cycle progression and differentiation[47,48] (Table 1). We validated this finding using RT-qPCR

in iPSCs, NPCs and FBNS, and found the specific upregulation of *CDKL5* and *SCML2* in L48R

cells at the iPSC stage only (Fig S2   B).   These transcriptomic results indicate that L48R

might impact the ability of iPSCs to proliferate and differentiate into NPCs, but it does not affect

the identity of iPSCs.

Nearly twice as many genes were dysregulated in L48R NPCs. Bulk RNA-seq on NPCs

identified 139 differentially expressed genes in L48R NPCs (p ≤ 0.05) compared to control

NPCs. 53 genes were upregulated (LFC ≥ 1) and 27 downregulated (LFC ≤ -1) (Fig. 2B). GO

analysis for all differentially expressed genes indicate that processes like cell migration and

establishment of synapses are affected, including pathways involved in cell adhesion, synapse

organization and   regulation of axonogenesis,   (Fig. 2C, Table 1). When   focusing

on genes comprising the cell adhesion and axonogenesis pathways   most genes

related to these processes were upregulated in L48R NPCs compared to controls (Fig. 2D)., S

everal protocadherin-encoding genes   which mediate neurite repulsion[49], were upregulated,

while *CDH4* and *WNT7A* were downregulated, indicating an imbalance in cell adhesion and

axon development cues (Table 1).

When L48R NPCs were converted to FBNs, we observed massive transcript-level dysregulation

(773 genes up, 965 down) (Fig. 2E). *EGR1, FOSL1, JUNB,* and *NPAS4,* immediate early genes

which mediate genetic responses to environmental cues[50], were downregulated (Fig. 2E, Table

1). GO analysis identified the upregulation of terms related to synapse organization and activity,

neurotransmitter uptake, and nervous system development (Fig. S2C   ). It also identified

terms related to G/M cell cycle phase transition and mitotic cell cycle phase transition, which are

notable given that neurons are post-mitotic (Fig. S2C   ).   We also found that astrocyte-

associated genes like *AQP4*, *SLC1A3,* and *SLC6A11* are upregulated in L48R FBNs,

suggesting that the   mixed population of cells were transciptomically more similar to

proliferating astrocytes, as opposed to post-mitotic neurons (Fig. S2D   ). To this end, we also

observed increased BrdU staining in L48R FBNs,   indicative of proliferation in the L48R cells

compared to controls following FBN differentiation (Fig. S2E   ).   When we analyzed the GO

terms for down-regulated genes in FBNs, we observed terms like regulation of MAPK cascade

, regulation of apoptotic signaling pathway, and cellular response to external stimulus (Fig. S2F

). Given the dysregulation of fundamental pathways that we observed in NPCs and the

downregulation of immediate early genes, it is likely that the widespread dysregulation observed

in the FBNs is secondary to these primary deficits.

To identify causative genes that might affect the overall differentiation of L48R cells, we looked

for commonly dysregulated genes at the iPSC, NPC, and FBN stage (Fig. 2F). Two genes were

commonly downregulated, and 13 were upregulated in all cell types (Table 2). Restricting the

analysis to genes commonly dysregulated later in the differentiation proccess at the neuronal

NPC and FBN stages, we identified 29 genes that were upregulated in both NPCs and FBNS,

and 8 that were downregulated (Table 3). All of the genes dysregulated early in differentiation

persisted throughout   (Fig. S2G).   **L48R variant increases *H3-3B***

gene expression and has increased protein deposition in 2D neuronal cell types

Because histones orchestrate chromatin compaction, their deposition in the nucleus affects both

genome organization and gene expression. We posited that differential H3.3 deposition

between L48R and control cells might underlie   some of the gene expression changes we

observed. W   e directly queried the altered expression of *H3-3A* and *H3-3B* in cells harboring

the L48R variant. Notably, expression of *H3-3B* was significantly upregulated in NPCs (adjusted

p-value = 1.58E-9), while *H3-3A* expression was not significantly affected (adjusted p-value =

0.99) (Fig. S3A). Using RT-qPCR, w   e   validated this RNA-seq result, finding that *H3-3B*

mRNA levels are upregulated in NPCs, and subsequently observed the same effect in FBNs

(Fig. S3B).   We observed no change in total levels of H3.3 protein between control and L48R

NPCs (Fig. S3C). We next stained FBNs for H3.3 to understand the deposition of H3.3 in these

post-mitotic cells and quantified the corrected total cellular fluorescence of H3.3/DAPI (Fig. S3D

). We observed a significant increase in this metric in L48R cells compared to control,

suggesting that H3.3 is hyper-deposited.

L48R variant alters chromatin accessibility and the occupancy of transcription factors

important for neuronal development in 2D models

To understand how increased H3.3   deposition affected global chromatin accessibility, we

performed Assay for Transposase-Accessible Chromatin with sequencing (ATAC-seq) in two

biological replicates of L48R NPCs and isogenic controls. L48R substantially opens chromatin

compared to the control, specifically at the TSS (Fig. 3A, Fig. S3E) Using DiffBind[43], we

identified 350 differentially accessible sites (FDR ≤0.05, p ≤ 0.01), which were most commonly

within 1kb of a promoter (40.86%) and in distal intergenic regions (21.43%) (Fig. S3E   ,

Supplementary File 2). Of these 350 regions, 253 were significantly more open (log2fold-change

≥ 1) and 97 regions were significantly less open (log2fold-change ≤ -1) in the L48R NPCs

compared to isogenic controls (Fig. 3B). W   e performed a GO analysis using all 350 sites and

found that   he most common terms were related to regulating neuronal development,

axonogenesis, and cell growth (Fig. 3C). *PLXNA3*, a gene important for axon extension[52,53], had

significantly more open chromatin in the L48R NPCs compared to controls (Fig. 3D). *RBFOX1*,

an RNA-binding protein that regulates neuronal splicing and is implicated in a range of NDDs[54]

had regions of more compact chromatin compared to control (Fig. S3H   ). *RBFOX1*

expression is significantly decreased in NPC, validating the ATAC seq (Fig.S3I). Taken

together, we found changes in H3.3 deposition in NPCs   harboring the L48R variant. The

resulting aberrant chromatin accessibility occurred at sites important for early neural

development (*RBFOX1*) as well as later stages of axon and synapse development (*PLXNA3*),

suggesting that some genomic regions may be improperly poised for transcription in cells that

harbor the L48R variant.

Next, we interrogated how changes in chromatin accessibility affected transcription factor (TF)

occupancy. Using ATACseqTFEA[55], we performed an   enrichment analysis and identified 64

TFs predicted to have differential occupancy between L48R and control. This analysis

generates enrichment scores; positive scores indicate   that the motif(s) for a TF are enriched

in L48R cells, and negative scores indicate that the motif(s) are depleted   . Significant factors

were called with an adjusted p-value ≤ 0.05. Of these factors, *ZNF614*, *PBX1,* and *ARX1* had

increased occupancy in L48R   , while *TP63* and *EMX2* had   decreased occupancy in L48R

(Fig. 3E). Increased occupancy of these TFs supports our previous data that transcription is

altered, with a specific effect on neurotransmission. Meanwhile, the factors that were depleted in

L48R support the transcriptomic data that the L48R variant impairs NPC differentiation.

We performed a GO analysis of all factors with changes in occupancy and performed a

dimensional reduction of these terms using REVIGO[36] (Fig. 3F). The terms clustered into four

groups represented by the terms: pattern specification process (cluster 1), mononuclear cell

differentiation (cluster 2), regulation of epithelial cell differentiation (cluster 3), and neural

precursor cell proliferation (cluster 4) (Fig. 3F). Every cluster contains TFs important for fate

specification in the developing nervous system. The most affected terms are important for

transcription of genes required for neural precursor cell proliferation, stem cell differentiation and

fate commitment, and forebrain development. To understand if altered TF occupancy might

account for changes in gene expression, we used TFLink[56] to find the targets of ZNF614, PBX1,

and ARX (increased occupancy) as well as TP63 and EMX2 (decreased occupancy), and

looked for changes in gene expression in our NPC RNA-seq dataset. The TFs with the greatest

number of targets affected in this dataset were *PBX1* (12 up, 5 down) and *TP63* (32 up, 19

down) (Table 4) (Fig.S3G). *DPP6, GABRG3, GABRA5, CDH4, PCDHGA3, WNT7A*, and

PLXNA3, genes with the largest differences in expression in L48R NPCs as compared to

control, are all predicted targets of p63 regulation. While validation is required to show changes

in binding/activity of these factors, these data support that the L48R variant affects TF binding

by driving changes in chromatin accessibility, resulting in aberrant expression of genes required

for NPC differentiation into mature neurons.

Chromatin accessibility changes in L48R NPCs affect expression of genes important for

cell adhesion and neurotransmission in 2D model

We next queried the overlap between ATAC-seq and RNA-seq datasets. First, we determined

the relationship between changes in accessibility and gene expression by comparing the

log2fold change in ATAC-seq reads binned by genes that were down, up, or unchanged in the

RNA-seq. We found that downregulated genes had the greatest decrease in accessibility, while

upregulated genes had the greatest increase in accessibility (Fig 4A).   We next identified six

genes that were down (designation from DESeq2 for RNA-seq and Diffbind for ATAC-seq) in

both datasets (p ≤ 0.05, LFC ≤ -1) (Fig. 4B   ). *CDH4* was the most downregulated gene

across these datasets (Fig. 4C   , Table 1). We performed RT-qPCR gene expression analysis

in iPSCs, NPCs, and FBNs to validate these results, and to understand *CDH4*'s expression

during both development and maturation. In isogenic controls, *CDH4* expression increased as

the iPSCs were differentiated to NPCs and FBNs, but when L48R iPSCs were differentiated to

NPCs and FBNs, *CDH4* had nearly undetectable levels (Fig. 4D   ).

Then   we identified 27 genes that were significantly upregulated in both datasets (p ≤ 0.05,

LFC ≥ 1) (Fig. 4E). Notable genes include *GABRG3* and *DPP6*, two genes involved in

neurotransmission (Table 1). *GABRG3* had more open chromatin at its promoter (Fig. 4F   )

and its expression increased as iPSCs were differentiated to NPCs and FBNs, with a

significantly greater increase in L48R cells compared to control (Fig. 4G   ). A related gene,

GABRA5, was also significantly up in both datasets (Fig. 4E   , Fig. S3G   ). Further, DiffBind

called eight different regions of *DPP6* more accessible in L48R NPCs: 1 exonic, 4 intronic, and

3 within 1kb of its promoter (Fig. 3H   ). When we analyzed the expression of *DPP6* in iPSCs,

NPCs, and FBNs, we observed a 20,000-100,000-fold induction in L48R NPCs and FBNs (Fig.

3I   ). Taken together, our ATAC-seq and RNA-seq identified a set of genes important for

neurotransmission, neuronal identity, and cell cycle progression with expression and chromatin

architecture that are perturbed by the L48R variant in NPCs. The changes in *H3-3B* expression

and H3.3 deposition, coupled with altered TF occupancy, suggest that L48R drives altered

chromatin accessibility, resulting in abnormal gene expression and subsequently affecting NPC

differentiation into neurons and glia.

L48R variant affects radial glia and alters neuronal maturation in 3D dorsal forebrain

organoids

Using an orthogonal model system to further analyze the neurodevelopmental consequences of

the L48R variant, we utilized 3D dorsal forebrain organoids (DFBOs) to understand cortical

development, neuronal maturation, and organization. DFBOs have been used in prior studies of

neurodevelopmental disorders, including models of autism spectrum disorder showing altered

excitatory/inhibitory balance and WNT signaling dysregulation (PMID: 26186191), as well as

investigations into radial glia behavior and mitotic defects in Miller-Dieker syndrome, a severe

lissencephaly (PMID: 28111201).

To study the effects of the L48R variant on early brain development, we converted isogenic

control and L48R 2D iPSCs into 3D DFBOs[45] and analyzed them at different time points (Fig.

S4A). DFBOs were collected at day 25 (D25) to analyze the neuroepithelial expansion (NE) and

at day 62 (D62) to analyze maturation. We analyzed organoid size across stages of

development and observed that L48R organoids were larger at day 25 compared to isogenic

controls, but this difference was not observable at later time points (Fig. 4A). L48R DFBOs at

D62 had fewer NE buds at their edges than isogenic controls (Fig. S4B).

At D25 and D62, we observed autonomously organized radial glia in both sets of organoids by

staining them with SOX2 (Fig. 5B,C). While there was no difference in SOX2 staining at D25

(Fig 5B   ), L48R DFBOs had significantly less expression of this marker at D62 (Fig. 5   C).

Next, we stained organoids with CTIP2, a marker of intermediate progenitor neurons. While not

significant, we observed that L48R DFBOs had more CTIP-positive cells at both time points

(Fig. S4D). During neurodevelopment, radial glia undergo both symmetric division, allowing for

self-renewal, and asymmetric division, which produce intermediate progenitor neurons[57] further

supporting that the pool of radial glia are being depleted through a skewing towards asymmetric

division, leading to increased progenitors committed to a neuronal fate. L48R DFBOs had

decreased TUJ1 staining at D25, but this difference was not present at D62 (Fig. 5B,C). This

may suggest that, while increased CTIP2+ intermediate progenitor cells are available, they are

less able to mature into TUJ1+ neurons within this D25 to D62 timeframe.

Since we observed dysregulation in the radial glia population, we stained D25 and D62

organoids with the cell proliferation marker Ki67. At D25, we observed a slight, but not

significant, increase in Ki67-positive cells in L48R organoids. By D62, however, there was a

significant decrease in Ki67-positive cells, indicating a loss of proliferative activity over time (Fig.

6A). To further probe cell cycle dynamics, we performed a BrdU incorporation assay, which

labels cells in S phase. At day 25, BrdU incorporation was significantly increased in L48R

organoids, suggesting early expansion of the proliferative progenitor pool. By day 62, however,

BrdU incorporation was significantly decreased in L48R organoids compared to controls (Fig.

6B,C). Taken together, these data suggest that the L48R variant promotes early accumulation

of progenitor-like cells that readily enter S phase (increased BrdU at D25), but do not show a

sustained increase in overall cycling (non-significant Ki67 increase at D25). Over time, these

progenitors appear to exit the cell cycle prematurely, as indicated by reduced Ki67 and BrdU

labeling at D62, along with decreased SOX2 expression (Fig. 6D). These results suggest that

L48R radial glia may have a functional deficit in maintaining self-renewing, symmetric divisions,

leading instead to an early shift toward asymmetric divisions and the expansion of intermediate

progenitor populations relative to control.

To understand how the L48R variant affects neuronal maturation, we stained organoids with

mature neuronal marker MAP2, which stains the dendrites of mature post-mitotic neurons. We

stained DFBOs collected at D62 with MAP2 and observed significantly lower levels in the L48R

DFBOs (Fig. 6E,F   ). These results also validate our findings in 2D FBNs   . We also

observed a decrease in PAX6, another radial glia marker in these organoids (Fig 6E,F). To this

point, we identified that L48R DFBOs had fewer proliferating radial glia, an expanded pool of

intermediate progenitors, and fewer   mature TUJ1+ neurons at D62. However, from these

experiments alone, it is not clear whether these differences in cellular composition affected the

function of these organoids.

To begin to answer this question, we looked at the expression of GABA, as our sequencing

experiments elucidated an increased expression of GABA receptors *GABRG3* and *GABRA5*.

We stained for GABA in D62 DFBOs and observed more GABA staining in the L48R condition

(Fig. 7   B). Further, we observed significantly higher mRNA expression of *GABRG3* in L48R

DFBOs (Fig. 7   C), validating our sequencing results in an orthogonal model. Taken together,

our work in DFBOs indicates that the L48R variant affect both symmetric radial glial proliferation

and asymmetric radial glia division, ultimately leading to defects in neuronal maturation. Further,

these results suggest that the L48R mutation altered neuronal subtypes, potentially leading to

an imbalance of excitatory/inhibitory signaling.

Neurons from L48R variant showed decreased electrical activity in 3D dorsal forebrain

organoids

We next investigated the electrophysiological properties of neurons from control and L48R

mutant 3D DFBOs via patch clamp recording (Fig. 7D). A significantly lower percentage of

neurons in the L48R DFBOs fired action potentials in response to positive current injections

compared to the control DFBOs (Fig. 7D a,b & f,g). Among the 45 neurons recorded from the

L48R 3D DFBOs, 17.8% (8/45) were multi-spiking, 24.4% (11/45) single-spiking, and the

remaining 57.8% (26/45) had no spiking (Fig. 7D e). In contrast, of the 48 neurons recorded

from the control organoids, 37.5% (18/48) were multi-spiking, 37.5% (18/48) single-spiking, and

the remaining 25.0% (12/48) neurons were not spiking ($P<0.05$, χ^2-test) (Fig. 7D b).

Interestingly, for the multi-spiking neurons, neurons from L48R mutant DFBOs exhibited a

higher firing frequency than those from control DFBO (Fig.S4E g) (Control: 12.1±1.4 Hz, n=18;

Mutant: 18.0±0.9 Hz, n=8; $P<0.05$; student t-test). No significant differences were observed in

resting membrane potential (RMP) or input resistance between neurons from control and mutant

organoids (Fig. S4E e-i).

Furthermore, a subset of recorded neurons exhibited spontaneous excitatory postsynaptic

currents (sEPSCs), reflecting synaptic inputs to these neurons. The percentage of neurons

showing sEPSCs was significantly lower in the L48R mutant (8.9%; 4/45) compared to the

control DFBOs (35.4%; 17/48) (P<0.05, χ²-test) (Fig. 7D e,j). No significant differences were

found in the frequency or amplitude of sEPSCs (Fig. S4E h,i). These results suggest that the

electrical activity in mutant DFBOs is comparatively less than that of their control counterparts.

To investigate the glial population at day 90 of DFBO maturation, where we observed reduced

electrical activity, we stained the DFBO with MAP2 and GFAP antibodies. We observed very

low levels of glial cell expression in L48R organoids (Fig. 7E). These results indicate that the

L48R mutation delayed neuronal maturation of DFBO by altering different cell types, further

impacting their electrical activity.

DISCUSSION

BLBS is a rare Mendelian syndrome with a mixed neurodevelopmental/neurodegenerative

phenotype for which only palliative interventions are currently available. Prior work in patient-

derived dermal fibroblasts, zebrafish, and mice have facilitated insights into the pathogenic

mechanism of this neurogenetic disorder[5,7]. However, until now   , the neurologic phenotype of

BLBS had never been interrogated in a human model of neurodevelopment. In the first reported

human iPSC-derived model of BLBS, we show that one BLBS causative variant, *H3-3B* p.L48R,

impairs neuronal differentiation and maturation in both 2D iPSC and 3D organoid systems. We

find that the identity of cells at the iPSC stage is largely unaffected by the L48R variant (Fig.

1B), but there   are changes in cell cycle regulators and cell adhesion molecules that affect

differentiation to NPCs (Fig. 2A, Fig. S2B). NPCs have decreased expression of fate markers

like *NES* and *SOX1* (Fig. 1C), and factors important for differentiation including *CDH4, WNT7A*

and *L1CAM* (Fig. 2B-D). Although we saw gene expression changes, total Forse1 and NCAD

positive NPC population in the 2D FBN is not affected in our FACS experiments indicating

subtle gene expression changes in iPSC are not affected overall NPC differentiation. Where as

L48R FBNs express reduced MAP2 (Fig 1D, E) and have widespread transcriptional

dysregulation, including decreased immediate early genes (Fig 2E) and increased astrocyte

markers   (Fig. S2D).   These results indicate that L48R specifically disrupts   NPCs,

further impacting the neuronal differentiation and maturation.   . Given the essential roles of

these processes, such dysregulation could explain the near-universal neurologic features of

BLBS.   . Further, myelin-related anomalies in individuals affected by BLBS account for 46% of

findings on MRIs[9]. Clinical evidence coupled with FBN transcriptomic data suggest a potential

shift in progenitor cell fate towards glia, or a competitive disadvantage for neurons harboring

the mutation, questions warranting further investigation.   .

To further interrogate the drivers of these changes in gene expression in NPCs and FBNs

harboring the L48R variant, we investigated how the expression of *H3-3B* itself was affected,

finding   that *H3-3B* expression was upregulated in both L48R NPCs and FBNs (Fig

S3A,B). There was also increased H3.3 deposition in L48R FBNs (Fig. S3   D). This   effect

on its own gene expression suggests that it might be acting through a gain of function

mechanism   [58], meaning that gene disruption therapies could be a viable therapeutic option for

individuals with this particular variant   . Next, we defined the chromatin landscape of L48R

NPCs using ATAC-seq and identified changes in chromatin accessibility at genomic regions

important for axon guidance, neuron projection, and synapse organization (Fig. 3C, D). We also

found changes in the occupancy of TFs that regulate processes like axon guidance and

GABAergic neuron differentiation (Fig. 3E, F). Previous studies using iPSC-derived neural

progenitor cells and neurons have demonstrated that chromatin accessibility dynamics are

essential to proper neurodevelopment and are often disrupted in neurodevelopmental disorders

such as autism spectrum disorder and epilepsy (PMID: 32732423; PMID: 39190448; PMID:

33545232). Our findings extend these observations by implicating a specific histone variant

mutation in chromatin dysregulation, providing a direct molecular mechanism. Previous work

done in embryonic stem cells showed that H3.3 promotes chromatin accessibility at promoters,

and loss of H3.3 impairs transcription factor binding (PMID: 36782260). The fact that H3.3

L48R increases, rather than decreases, chromatin accessibility, further supports that this variant

is gain of function.

While w   e were able to broadly define the chromatin landscape of L48R NPCs using ATAC-

seq,   , chromatin immunoprecipitation with sequencing (ChIP-seq) for H3.3 and the

distribution of active/repressive hPTMs   will provide more mechanistic insight

underlying the changes in chromatin states.   ChIP-seq specifically for L48R H3.3 would also

indicate changes in chromatin accessibility and gene expression at the sites were due directly to

altered H3.3 deposition.

When integrating our multi-omic data, we found that downregulated genes from RNA-seq had

the greatest decrease in accessibility, while upregulated genes had the greatest increase in

accessibility (Fig 4A).   he RNA- and ATAC-seq datasets overlapped at a few key sites,

and generally had more upregulated genes/open chromatin in L48R NPCs compared to

isogenic controls (Fig. 4E   ). Many genes were related to key neurodevelopmental processes,

such as cell adhesion and neurotransmission   . The consistent downregulation of *CDH4*, a

gene that encodes a cadherin fundamental for cell-cell interactions in the central nervous

system, might be one of the first causes of differentiation deficits in L48R cells, preventing

proper migration during neurodevelopment. The upregulation of *DPP6* and GABA receptors

GABRG3 and *GABRA5* suggest that L48R cells have impaired neurotransmission with

decreased neuronal activity [1]. This altered excitatory/inhibitory balance has been observed in

several NDDs and is considered a unifying mechanism underlying this group of disorders.

Our work in 3D DFBOs recapitulate many of our findings in 2D cultures, and in what is seen

clinically in affected individuals. When we measured the size of the DFBOs as they matured, we

found that they were initially larger, and then normalized in size compared to control (Fig. 5A).

At the time of last evaluation, the individual harboring the L48R variant in the BLBS cohort had

microcephaly[5]. Interestingly, the growth phenotype of individuals with BLBS evolves over

time. The first reported individual with BLBS presented with macrocephaly and overgrowth at

birth but, at re-evaluation a decade later, exhibited microcephaly and severe undergrowth[5,8,62].

We found that this trend applies to other individuals affected by BLBS for whom follow-up data

is available (https://doi.org/10.1016/j.xhgg.2025.100440). L48R DFBOs capture this phenotypic

transition with increased organoid size and proliferation at D25, followed by reduced

proliferation and progenitor cells at D62. Given that , reduced proliferation of neuronal

progenitor cells is the basis for primary microcephaly (PMID: 30936766), we believe that this

L48R DFBO model will be an effective model to study the mechanism driving this

phenotype.

Another significant benefit of this DFBO model is that we could interrogate the impact of the

L48R variant on multiple cell types at once. Our SOX2 and PAX6 staining show that there is a

decrease in radial glia between D25 and D62 (Fig. 5B-C, Fig. S4C). The CTIP2 staining, which

shows an increase in intermediate progenitor neurons during this same window (Fig. S4D),

suggests that this decreased in radial glia is caused by a skewing towards the asymmetric

division of radial glia, away from the expected homeostatic balance between self-renewing

symmetric division and intermediate progenitor neuron-producing asymmetric division[57]. This is

further supported by the Ki67, BrDU, and TUJ1 data, which show a decrease in proliferation

accompanied by an increase in neurons between D25 and D62 (Fig. B-D). The decrease in

radial glia may result from altered TF occupancy affecting pathways that regulate glial fate

commitment and self-renewal. Thus, L48R cells ultimately differentiate into similar numbers of

neurons at the cost of maintaining their progenitor identity. However, the D62 L48R neurons

have delayed maturation (Fig. 6A), potentially because of dysregulated TF occupancy, or cell

death   caused by   grossly dysregulated gene expression and chromatin accessibility   .

Further, our whole-cell patch clamp analysis indicated L48R DFBO has significantly less

electrical activity and these organoids have very low glial cells (Fig. 7D, E). Taken together,

these results indicate that L48R progenitors are less able to proliferate over the longer periods

of organoid maturation and affected maturation of neurons and their electrical activity. The

exhaustion of a replicative pool which only exists for a transient period during

neurodevelopment might also explain why the central nervous system is universally affected but

other organs are not.

We concluded our analyses at day 90 of maturation, where control DFBO started producing the

low levels of GFAP positive glial cells, longer maturation periods are required to understand the

significance of increased glial markers observed in the 2D FBNs, and their effects on glial

production and proliferation in a 3D context will be informative, especially in understanding if

these cells are skewed towards a non-neuronal fate. The use of cerebral organoids, which are

less guided than DFBOs, would enable the study of additional cell types that make up

hippocampal and retinal regions[64]. Oligodendrocyte-containing myelin organoids (myelinoids)[65]

will also be important models to understand the myelin-related anomalies observed in many

individuals affected by BLBS.

Overall, our data suggest a model by which L48R acts through a gain-of-function mechanism

leading to increased *H3-3B* expression that results in the hyper-deposition of H3.3 into the

nucleosome   , disrupting gene expression and chromatin accessibility. Functionally, this

causes dysregulation of cell adhesion, neurotransmission, and the balance between excitatory

and inhibitory signaling.

Alterations in GABAergic signaling, cell adhesion, and radial glia function are central to

neurodevelopmental disorders (NDDs) and converge on core mechanisms of circuit formation.

GABAergic dysfunction, a common feature in disorders like ASD and Rett syndrome, disrupts

the excitatory/inhibitory balance essential for cortical function (PMID: 33772226, 21068835).

Similarly, downregulation of adhesion molecules such as CDH4 can impair neuronal migration

and synaptic organization, as seen in Fragile X syndrome (PMID: 39022311). Radial glia, critical

for both progenitor expansion and neuronal scaffolding, are especially vulnerable to epigenetic

and signaling perturbations, with defects linked to conditions such as microcephaly and

lissencephaly (PMID: 28111201). Together, these observations underscore how epigenetic and

transcriptional alterations like those caused by H3.3 L48R can converge on key developmental

processes to yield circuit-level dysfunctions that underlie NDD pathogenesis.

The ability to define a proposed model with modifiable mRNA and protein level readouts, as well

as functional consequences, establishes the necessary groundwork for future studies into

therapeutic interventions. Specifically, our data show   the upregulation of two GABA$_A$ receptor

encoding genes, *GABRG3* and *GABRA5,* in L48R NPCs. If GABA-mediated dysregulation is

validated in future mechanistic work to underlie   disease pathogenesis, this could be targeted

through the repurposing of an FDA-approved class of GABA$_A$ receptor selective positive

allosteric modular antidepressants[66]. Further, because we found that *H3-3B* itself is

upregulated, suggesting a not previously defined gain-of-function mechanism, targeting the

mutated allele with an antisense oligonucleotide (ASO) or small interfering RNA may be a viable

therapeutic. Finally, if future work identifies the dysregulation of hPTMs, perhaps an existing

FDA-approved HDAC inhibitor could be repurposed for BLBS, which is being explored for

MDEMs[2]. Our results from one model have highlighted a few therapeutic approaches, but

intense future study into the mechanism of disease and the safety of possible interventions is

needed to understand what therapeutics would be most beneficial for individuals affected by

BLBS.

There are noteworthy strengths of our approach, which leverages 2D and 3D iPSC models

coupled with sequencing and IF, but we acknowledge the limitations of this work. First, these

studies were conducted in iPSCs that were derived from a single edited clone. To address this

limitation, we performed at least two separate differentiations of iPSCs to each cell type queried;

however, the gold standard would be to validate these findings in an independently generated

cell line. Further, it is not clear if all findings from our XX (female) model would be recapitulated

in an XY (male) model. Since there is an almost even number of males and females affected by

BLBS[8], these results would be strengthened if recapitulated in both an XX and XY model

system. It is also important to note that 70 unique variants are reported to cause BLBS[8], and we

investigated the effects of only a single BLBS-causing variant. Because of the phenotypic

variability observed in affected individuals and model systems[5,8,29], future studies in isogenic cell

lines from different origins harboring additional BLBS variants will be important in assessing the

generalizability of these findings to have the greatest impact for affected individuals.

In sum, we developed and interrogated a novel isogenic L48R iPSC line differentiated into 2D

NPCs, 2D FBNs, and 3D DFBOs, uncovering critical insights into the pathogenesis of Bryant-Li-

Bhoj syndrome (BLBS). These data support a model in which the L48R variant disrupts NPC

proliferation and differentiation by altering chromatin accessibility at loci regulating axon

guidance, cell adhesion, and synapse organization. These early developmental deficits prevent

neuronal maturation and affect their electrical activity.

Furthermore, the dysregulation of GABA receptor pathways suggests a potential imbalance in excitatory/inhibitory signaling, with implications for neuronal network function. This study represents the first human iPSC-derived model of BLBS and highlights the critical role of chromatin dysregulation in neurodevelopmental disorders. Future work should focus on validating these findings in additional BLBS variants and assessing the therapeutic potential of antisense oligonucleotides or GABA receptor modulators. This model provides a foundation for understanding BLBS pathogenesis and advancing targeted therapeutic strategies for histone-associated neurodevelopmental syndromes.

ACKNOWLEDGMENTS

We would like to sincerely thank the members of the BLBS community for sharing their information and journeys with us. Thank you also to Teodora Orendovici, PhD, Stephen Mahoney and Ashley Bushey from the CHOP High Throughput Sequencing Core for assistance with the ATAC-sequencing; Jean Ann Maguire, PhD from the CHOP iPSC core for generating the iPSC line; and Jonathan Billings, Kyra Booth, Tiancheng Wang and Kayleigh Ostberg from the CHOP Center for Applied Genomics for assistance with RNA-sequencing.

CONTRIBUTIONS

Data Collection: RA, AKS, LMB, DLC, XMW, EAW, JPB, YW

Data Analysis: AKS, RA, ELD

Figure Generation: AKS, RA, EEL

Conceived of study: RA, AKS, EJB

Drafted the manuscript: AKS, RA, EEL

Reviewed manuscript: EEL, ELD, DLC, XMW, KJC, MM

FUNDING

Support for this work was provided by NICHD F30 1F30HD112125 (EEL); NIGMS T32

5T32GM008638 (ELD); NHGRI T32HG009495 and the Eagles Autism Foundation (DELC); the

Chan-Zuckerberg Initiative Neurodegeneration Challenge Network, as well as the Burroughs

Wellcome Fund and the Hartwell Foundation (EJKB).

DATA AVAILABILITY

The datasets generated during the current study will be available in the Gene Expression

Omnibus repository.

COMPETING INTERESTS

The authors declare no competing interests.

Figure Legends:

Figure 1. Generation and initial characterization of L48R 2D iPSCs, NPCs and FBNs. A)

c.146T>G *H3-3B* Leu48Arg (L48R) variant was introduced to CHOPWT14 hiPSCs using

CRISPR HDR and   pure population of edited cells were attained and differentiated into

NPCs   and FBNs   . B-G) **L48R and isogenic control iPSCs showed no significant**

differences in pluripotency markers (OCT4, NANOG). L48R NPCs had reduced

expression of NANOG (p = 0.00836 **) and SOX2 (p = 0.02608) and FBNs exhibited impaired**

neuronal extension (MAP2, VGLUT1) and increased astrocytic marker GFAP (p = 1.104e-

06)   D) Immunofluorescence of FBNs stained with DAPI, MAP2 (neuronal extensions),

VGLUT1 (glutamatergic neurons). E-F) Quantification of images in D) using FIJI. n = 10 images

from 3 replicates for each condition. MAP2 p   = 9.536e-05, VGLUT1 p   = 6.694e-05. G   )

RT-qPCR for astrocyte marker *GFAP*   p   = 1.104e-06.   Comparisons made via t-test and

shown with brackets. *p ≤0.05, ** p ≤0.01, *** p ≤0.001

Figure 2. Transcriptomic changes in L48R progenitor cells affect differentiation into

FBNs. A-C) **RNA-seq reveals dysregulation of cell adhesion (e.g., CDH4, WNT7A) and**

axonogenesis (e.g., PLXNA3) pathways in L48R NPCs. Blue dots represent downregulated

genes, red = upregulated genes. Dotted horizontal line denotes an adjusted p-value of ≤ 0.01

and dotted vertical lines indicate a log2fold change (LFC) with an absolute value of ≥ 1. **GO**

analysis identifies upregulation of genes affecting synapse activity and neurotransmitter

uptake in FBNs, suggesting L48R affects both early differentiation and neuronal function.

D) Heatmap representing the Z-scored LFCs of the genes comprising cell adhesion and

axonogenesis GO terms. These genes appear in 9/12 significant GO terms between L48R and

control NPCs. Descriptions of function and disease association are in Table 1 for these genes.

E) Volcano plot of L48R vs control FBNs formatted as in A-B.   F) Upset plot of shared hits

between the RNA-seq datasets generated in iPSCs, NPCs and FBNs. n = 2 commonly

downregulated genes, n=10 commonly upregulated genes.

Figure 3. L48R variant alters chromatin accessibility and the occupancy of transcription

factors important for neuronal development. A) Metaplot comparison of ATAC-seq

average signal from control and L48R NPCs at all peaks, centered at the transcription

start site (n = 2 Control, 2 L48R biological replicates).   . B) Genomic heatmap of

differentially accessible sites grouped by sites that are more accessible in L48R (top, purple   ),

and those that are less accessible in L48R NPCs (bottom, green   ). C) GO enrichment of

regions with significantly increased chromatin accessibility, annotated by nearest gene, in L48R

NPCs compared to control. D) RNA-seq (top, blue) and ATAC-seq (bottom, pink)   **reveals**

chromatin accessibility changes at loci regulating axon guidance (PLXNA3) in L48R

NPCs. E)   **Transcription factor analysis indicates increased occupancy of PBX1 and**

decreased occupancy of TP63, suggesting dysregulation of differentiation and

neurotransmission pathways. F) REVIGO summary of GO terms associated with transcription

factors in E) clustered by similar GO terms.

Figure 4. Chromatin accessibility changes in L48R 2D NPCs affect expression of genes

important for cell adhesion and neurotransmission. A) ATAC-seq fold change

(L48R/Control) for downregulated, upregulated, and unchanged genes in L48R NPC RNA-seq.

Statistics were calculated using a t-test with Bonferroni correction. B-H) **Overlap analysis**

identifies upregulation of neurotransmission-related genes (e.g., GABRG3, DPP6) and

downregulation of adhesion-related genes (e.g., CDH4) in L48R NPCs. These results

suggest chromatin and transcriptional dysregulation contribute to impaired

differentiation and neuronal maturation.B)   Venn diagram of overlap of significantly

downregulated genes in RNA- and ATAC-seq. C   ) RNA-seq (top, blue) and ATAC-seq

(bottom, pink) tracks for   *CDH4,* a cell adhesion gene downregulated across all L48R cell

types. Control tracks are shown in green, L48R in purple. D   ) RT-qPCR validation of *CDH4*

mRNA levels   from iPSC, NPC, to FBN. ΔΔ Ct values are normalized to *GAPDH*.   p-

values: iPSC = 9.834e-05, NPC   = 0.0008733, FBN   = 0.001522. E   ) Venn diagram of

overlap of significantly upregulated genes in both datasets.   F-G   ) RNA- and ATAC-seq

tracks + RT-qPCR validation of *GABRG3*.   p-values: iPSC = 0.002509, NPC   =

0.008845, FBN   = 0.02047.   H-I   ) RNA- and ATAC-seq tracks + RT-qPCR validation of

DPP6.   p-values: iPSC = 0.0001059, NPC   = 2.596e-05, FBN   = 0.005359.

Comparisons made via t-test and shown with brackets. *p ≤0.05, ** p ≤0.01, *** p ≤0.001

Figure 5. L48R variant perturbs progenitor populations in 3D DFBOs.   A) Left:

Representative images of organoids at D25, 45, and 62.   D25 images were taken at 10X

magnification, D45 and D62 images were taken at 4x magnification, scale bar = 500μm Right:

Quantification of organoid size using Qupath[67]. D25 n = 6 control,   n = 7 L48R   ; D45 n = 6

control and L48R   ; D60 n = 15 control   , n = 16 L48R   . D25 p-value = 0.01684,   . B)

Organoids collected at D25   were stained with neuronal marker TUJ1 (green), progenitor

marker SOX2 (pink) and DAPI (blue). **(C).**   D25 n = 6 control,   n = 7 L48R

. TUJ1 p-value = 0.002887   . C) Organoids collected at D62 were stained for the same

markers as in B). Quantification of staining is shown on the right. D62 n = 8 control, n = 12 L48R

SOX = 0.005652.   Comparisons made via t-test and shown with brackets. *p ≤0.05, ** p

≤0.01, *** p ≤0.001

Figure 6. L48R DFBOs display abnormal proliferation and   fewer mature neurons   . A)

Organoids stained for SOX2 (green) and proliferative marker Ki67 (red) at both day 25 and day

62. Images are quantified on the right. B-C) BrDU incorporation in control (top) and L48R

(bottom) organoids at day 25 (left) and day 62 (right) with quantification (C).   D) Quantification

of SOX2, SOX2 p-values: D25 = 0.08127, D62 = 0.001318. E   ) D62 DFBOs stained with

MAP2 (green) and PAX6 (red). R   epresentative images taken at 10X are displayed **L48R**

DFBOs exhibit reduced mature neuronal marker MAP2.   .   p-value = 0.006039.

PAX6 staining is quantified in Fig S4C. Comparisons made via t-test and shown with brackets.

*p ≤0.05, ** p ≤0.01, *** p ≤0.001

Figure 7. L48R DFBOs display increased GABA abundance

A-B) 10X images of D62 DFBOs stained with GABA (red) and NeuN (green).   . p-value =

0.03561. C) RT-qPCR quantification of *GABRG3* levels in n = 4 DFBOs for both genotypes. p-

value = 3.597e-05. D.a-j) Neurons in L48R DFBOs show less electrical activity than those in

control DFBOs. **a.** Whole-cell patch clamp recording of a neuron from a control DFBO,

showing action potential firing upon current injection (baseline membrane potential at -

70 mV). **b.** Percentages of neurons from the control organoids show various responses

to current injections. **c.** Voltage-clamp recording trace showing spontaneous excitatory

postsynaptic currents (sEPSCs) at a holding potential of -70 mV. **d.** Enlarged trace in

the shaded area in panel C. **e.** Percentages of neurons with or without sEPSCs from the

control organoids. **f.** Whole-cell patch clamp recording of a neuron from an L48R

organoid, showing action potential firing upon current injection (baseline membrane

potential at -70 mV). **g.** Percentages of neurons from the L48R DFBOs showing various

responses to current injections. **h.** Voltage-clamp recording trace showing sEPSCs at a

holding potential of -70 mV. **i.** Enlarged trace in the shaded area in panel H. **j.**

Percentages of neurons with or without sEPSCs from the L48R DFBOs. E) L48R DFBO

have less GFAP positive glial cells. 10X images of D90 DFBOs stained with GFAP (red) and

MAP2 (green). Images were quantified as described in (A). p-value = 0.0003. Data is collected

from two biological replicates and n=16 organoids from the two replicates.

Tables

Table 1 Function and disease associations of the dysregulated genes in L48R compared
to isogenic controls.
942

Category	Data set	Gene name	Function	Disease Associations	Reference
Cell cycle progression / Differentiation	iPSC RNA-seq	*CDKL5*	Encodes a cell cycle inhibitor that promotes the differentiation of neuronal cells	Developmental and Epileptic Encephalopathy 2	47 , OMIM #3006
	iPSC RNA-seq/RNA-seq +	*SCML2*	Encodes polycomb group protein that regulates the cell cycle by controlling the progression of cells from G1 into S phase	No known associations	48

	ATAC-seq overlap				
	NPC RNA-seq + ATAC-seq overlap	CDH4	Encodes a classic cadherin, R-cadherin, a calcium dependent cell adhesion protein. Classic cadherins are important for tissue architecture across organs and specifically in the brain for neurite interactions	No known associations	68,69
	NPC RNA-seq	L1CAM	Encodes L1CAM, which acts statically as glue between cells and promotes cell migration during neurodevelopment	X-linked hydrocephalus, Masa Syndrome	70, OMIM #s 303 307000
Cell adhesion/Cell migration	NPC RNA-seq	FLNA	Encodes Filamin A, an actin binding protein. It is required for neuronal migration to the cerebral cortex. This gene is also important for cardiac morphogenesis.	Periventricular Heterotopia, Otopalatodigital syndrome, Frontometaphyseal dysplasia, Melnick-Needles syndrome	71,72
	NPC-RNA-seq	PCDHGB1, PCDHGB3, PCDHGA4, PCDHGA5, PCDHGB2, PCDHGA3, PCDHGC3, PCDHA6, PCDHB11	Encode members of the clustered protocadherins, which are a family of cell-surface homophilic proteins. Protocadherins regulate axonal tiling and confer identity to neurons, which is required for the distinguishment between self and non-self.	ASD, schizophrenia	49,51
Axon guidance	RNA-seq + ATAC-seq	PLXNA3	Encodes the most widely expressed plexin receptor in fetal brain, plexin-A3[41], a class 3 semaphorin receptor known to be important for axon pathfinding in the developing nervous system and functions directly in signaling repulsion[42]	X-linked intellectual disability syndrome	52

	NPC RNA-seq	*WNT7A*	Required for neural stem/progenitor cell proliferation and self-renewal and adult mouse brains, stimulates the formation and function of excitatory synapses, increases density of dendritic spines	Fuhrmann syndrome, ASD	73–75, OMIM#22893
	NPC RNA-seq	*ARHGAP4*	Encodes a Rho GTPase that negatively regulates cell and axon motility	No known associations	76
	NPC RNA-seq	*SEMA5A*	Encodes semaphorin-5A, a negative regulator of excitatory synapse formation	ASD	77
Neuronal transcription factors	ATAC-seq TF analysis	*PBX1*	Controls organ patterning during embryogenesis. In the brain, it regulates the survival of midbrain dopaminergic neurons and is an early regulator of subventricular zone neurogenesis	Parkinson's disease, CAKUTHED	78,79, OMIM #617641
		ARX	Specifies forebrain patterning and growth, and specifically controls the development of telencephalic GABAergic neurons	Epilepsy and NDDs with or without structural defects	80,81
		TP63	Encodes p63, a protein that works cooperatively with p53 to promote apoptosis during neuronal development, which is important for establishing neuronal connectivity	Ectrodactyly, ectodermal dysplasia, and cleft lip/palate syndrome 3	82, OMIM #6042
		EMX2	Expressed primarily in cortical progenitors where it regulates multiple features of cortical development including arealization and specifying positional identity of cortical cells	Schizencephaly	83, OMIM # 269
Neurotransmission	RNA-seq + ATAC-seq	*DPP6*	*K*nown canonically as an auxiliary subunit of Kv4-mediated A-type K^+ channels, which act to delay excitation and regulate firing frequency. *DPP6* enhances the surface expression of these channels and is thought to accelerate their kinetics. It is also important for dendritic branching and synaptic development during neuronal development.	Schizophrenia	84–86, OMIM #600035
		GABRG3	Encodes a $GABA_A$ receptor- γ3, which binds GABA, the major inhibitory neurotransmitter in the human brain. GABA regulates every step of neurogenesis and regulates NPC proliferation and	Epilepsy, ASD	87–90

		migration through its -A receptors.		
	GABRA5	encodes GABA$_A$ receptor subunit	Early onset epilepsy and developmental delay	[91]
Immediate early genes	FBN RNA-seq	EGR1, FOSL1, JUNB, NPAS4	Immediate early genes are rapidly upregulated in neurons in response to a wide variety of cellular stimuli and are important for synaptic plasticity. They provide the molecular framework for both a rapid and dynamic response to neuronal activity and lasting gene expression changes.	Anxiety, depression, schizophrenia	[50,92]

944 **Table 2 Genes dysregulated across 2D iPSCs, NPCs and FBNs.**

	iPSC		NPC		FBN	
Gene ID	**log2fold change**	**adjusted p-value**	**log2fold change**	**adjusted p-value**	**log2fold change**	**adjusted p-value**
PCDHA6	-2.193450727	2.17E-06	-1.355795213	8.71E-09	-1.308921624	0.045580571
ZNF558[1]	-1.148059517	0.004709638	-1.128578936	8.37E-10	-1.199734433	8.80E-05
GK	1.210861399	7.77E-10	1.030926781	8.37E-10	1.779059583	1.73E-09
SCML21	1.042489231	0.000268	1.115859018	4.57E-12	1.650203502	1.49E-07
SLC16A2	1.001951462	0.000177	1.178133567	1.31E-05	1.771954006	6.49E-10
SLC15A4	3.032307096	6.79E-27	2.640470731	6.30E-25	1.152395578	0.005698916
DPP6	3.988997109	4.29E-09	4.408206465	4.91E-65	6.226215688	1.75E-101
ZNF37A	5.871603628	9.76E-38	4.860032685	3.23E-87	4.325495167	4.14E-22
PCDHGA3	3.097728204	1.15E-05	5.474057123	1.58E-122	4.801058573	3.10E-63
HIST1H1A	2.471946327	8.11E-36	5.848215734	1.80E-18	5.856905392	0.002878737
GLT1D1	8.915844422	1.64E-09	9.139164431	2.33E-10	7.457342153	1.41E-07
TMEM132D	9.629365976	1.03E-11	9.931158475	2.67E-13	8.251489265	1.03E-08

945 [1]There are no clinical phenotypes in OMIM for *SCML2* or *ZNF558* however *SCML2* has been

946 associated with, but is not causative in, Nance-Horan Syndrome

948 **Table 3 Genes dysregulated in both L48R NPCs and FBNs compared to isogenic**

949 **controls.**

Gene ID	NPC		FBN	
	log2fold change	adjusted p-value	log2fold change	adjusted p-value
ANKRD20A1	-1.460915202	2.27E-06	-1.439994876	4.64E-05
CDH41	-3.590286774	2.77E-31	-4.668488495	5.01E-62
PCDHA61	-1.355795213	8.71E-09	-1.308921624	0.045580571
PCDHB8	-1.391417915	0.001425716	-1.546283192	0.005391523
PEG3	-1.908978559	1.35E-20	-2.286904202	6.99E-06
TGM4	-2.716613511	0.008999827	-4.179492072	0.000408431
ZIM2	-2.738228127	0.006096512	-3.135289722	8.30E-07
ZNF5581	-1.128578936	8.37E-10	-1.199734433	8.80E-05
DNASE1L1	1.186058534	0.001772585	1.378797988	0.005401369
DPP61	4.408206465	4.91E-65	6.226215688	1.75E-101
FAM58A	1.12636598	3.36E-12	1.207565785	0.013980524
GABRG3	4.22607955	9.14E-41	1.725418457	1.08E-07
GK1	1.030926781	8.37E-10	1.779059583	1.73E-09
GLT1D11	9.139164431	2.33E-10	7.457342153	1.41E-07
HAUS7	1.077143968	0.000217	1.469187295	0.001673216
HIST1H1A1	5.848215734	1.80E-18	5.856905392	0.002878737
HSD17B10	1.031920999	9.67E-11	1.433723942	6.97E-05
IDH3G	1.001882884	4.29E-11	1.434234676	0.007311695
JADE3	1.079089438	3.51E-12	1.429530196	6.81E-06
NAA10	1.016446794	8.32E-09	1.420341345	0.000765417
NCAM2	1.175437195	0.005286634	1.103948768	2.24E-07
NDUFB11	1.171103978	1.43E-08	1.88856872	0.000684263
PCDHGA31	5.474057123	1.58E-122	4.801058573	3.10E-63
PCDHGA4	2.086201001	2.37E-11	1.512198469	0.016392048
PCDHGB1	1.081975754	0.001407812	1.525777875	0.021808739
PCDHGB2	1.746665418	9.95E-10	1.827544024	0.007254347
PCDHGB3	2.323839531	5.77E-15	1.873734051	0.002575344
PDHA1	1.093747283	3.29E-17	1.509521746	1.13E-19
RPL10	1.023326738	0.00025	1.397910368	0.024534554
SCML21	1.115859018	4.57E-12	1.650203502	1.49E-07
SLC15A41	2.640470731	6.30E-25	1.152395578	0.005698916
SLC16A21	1.178133567	1.31E-05	1.771954006	6.49E-10
SLC30A8	1.506637641	0.0002	1.238069979	0.00434561
TMEM132D1	9.931158475	2.67E-13	8.251489265	1.03E-08
TSPAN7	1.175352888	1.61E-07	2.135303521	5.38E-10

ZNF37A1	4.860032685	3.23E-87	4.325495167	4.14E-22
ZNF41	1.13161708	9.43E-08	1.149731459	0.000458778

950   [1] These genes are dysregulated throughout differentiation.

951   **Table 4 Targets of PBX1 (increased occupancy) and TP63 (decreased occupancy).**

Transcription Factor	Downregulated gene target	Upregulated gene target
PBX1	BLOC1S5-TXNDC5 CDH4 DOCK11 NDUFAF1 ZNF558	ARHGAP4 G6PD GLT1D1 HAUS7 IKBKG SLC10A3 SLC15A4 SLC16A2 SSR4 ZNF41 ZNF492 ZNF726
TP63	AMMECR1L BLOC1S5-TXNDC5 **CDH4** CES1 CHAT DOCK11 GAS8 GOLGA7B INPP5F NDUFAF1 PCDHA6 PCDHB8 PEG3 SETD1A SLITRK4 TGM4 **WNT7A** ZIM2 ZNF558	ARHGAP4 CDC42BP6 DNASE1L1 **DPP6** EMD FAM50A G6PD **GABRA5** **GABRG3** GK GLT1D1 HAUS7 HSD17B10 IDH3G IKBKG JADE3 NAA10 NCAM2 OCA2 **PCDHGA3** PCDHGA5 **PLXNA3** POTEF RPL10 SLC10A3

		SLC15A4
		SLC16A2
		SSR4
		TMEM132D
		TSPAN7
		ZNF37A
		ZNF41

Table 5 Primary antibodies used for protein-based experiments.

Antibody	Species	Catalogue #	Dilution
MAP2	mouse	Millipore mab3418a5	1:400
VGLUT1	Rabbit	Sigma 48-2400	1:1,000
TUJ1 (Neuron-specific beta-III Tubulin)	Mouse	R&D Systems NL1195G	1:200
GFAP	Rabbit	Novus NB120-16997	1:200
BrdU	Mouse	Invitrogen MA3-071	1:100
H3.3	Rabbit	Invitrogen PA5-29602	1:1,000
H4	Mouse	Cell Signaling #2935S	1:500
β-Actin	Mouse	Santa Cruz sc-69879	1:1,000
SOX2	Rabbit	Cell Signaling 35795	1:400
Ki67	Rabbit	Cell Signaling 9129	1:400
PAX6	Mouse	Sigma AMAb91372	1:100
GABA	Rabbit	Sigma A2052	1:500

Table 6 qPCR primers used to validate cell identity and validate RNA-seq results.

Gene	Forward primer[1]	Reverse primer[1]
GAPDH	GGCATCCTGGGCTACACTGA	GAGTGGGTGTCGCTGTTGAA
H3-3B	TCGTTGGGCGGTGCTGGTTTTT	TTAGGCCACTTCTCCGACCGCC
GABRG3	ATGCTACGCCAGCAAGAACAGC	GGCGAAGACAAACAGGAAGCAC
DPP6	GACCGACAGATGCCTAAAGTGG	TGTCGGTGAAGGTTGCTGGCTT

GFAP	CTGGAGAGGAAGATTGAGTCGC	ACGTCAAGCTCCACATGGACCT
OCT4	CCTGAAGCAGAAGAGGATCACC	AAAGCGGCAGATGGTCGTTTGG
NANOG	CTCCAACATCCTGAACCTCAGC	CGTCACACCATTGCTATTCTTCG
SOX1	GAGTGGAAGGTCATGTCCGAGG	CCTTCTTGAGCAGCGTCTTGGT
NES	TCAAGATGTCCCTCAGCCTGGA	AAGCTGAGGGAAGTCTTGGAGC
MAP2	AGGCTGTAGCAGTCCTGAAAGG	CTTCCTCCACTGTGACAGTCTG

[1]All primer sequences were obtained from OriGene Technologies Inc.

Supplementary Figures:

Supplementary Figure 1 Characterization of 2D L48R iPSCs, NPCs and FBNs. A) 3D in

silico structural model of the H3.3-containing nucleosome (PDB: 5X7X) with one of the

two copies of H3.3 in purple; other histones in gray; and DNA in black/grey. The

canonical Leu48 residue is shown on the left, the variant Arg48 is shown on the right in

green. The distance in angstroms between H3.3 T45, H4 and DNA are displayed. B   )

Sanger sequencing verification of isogenic control (top) and heterozygous c.146T>C edit in

CHOPWT14 cells (bottom). C   ) Description of CHOPWT14 iPSCs. Method = reprogramming

method. D-G) Analysis of representative markers comparing iPSC and NPC

differentiation using flow cytometry. D & F) Representative dot plots of flow cytometry

results with antibodies specific to Forse1 and NCAD. At Day 0, iPSCs are devoid of

Forse1 and NCAD, whereas by 20 days of NPC differentiation, cells are positive for

both Forse1 and NCAD. E & G) Box plots showing the percentage of Forse1 and

NCAD positive cells in four biological replicates. Samples were treated with secondary

conjugated to Alexa 488 and data is represented as FITC signal against SSC-A. Data

are analyzed from four biological replicates, each with a minimum of 10,000 cells

analyzed for each condition.   H   ) IF analysis in FBNs of TUJ1 (neurons) and GFAP (glia)

collected at day 20. I   ) Quantification of n = 5 z-stacks for control, and n = 6 z-stacks for L48R

FBNs. Corrected total fluorescence was calculated by dividing the fluorescence intensity of

TUJ1 and GFAP by the fluorescence intensity of DAPI for each image. J) Quantification of the

number of cells per image expressing each marker of interest.   K) RT-

qPCR analysis of *MAP2* mRNA levels in iPSCs and FBNs.   p-values iPSCs = 0.4915   , FBNs

= 0.2222   .

Supplementary Figure 2 Transcriptomic dysregulation induced by L48R variant A) PCA of

L48R and control iPSCS, NPCs and FBNS, colored by genotype. B) RT-qPCR for SCML2 and

CDKL5 to validate findings from bulk RNA-Seq. C) REVIGO summary of GO terms associated

with significantly upregulated genes in FBNs, clustered by similarity. D   ) Heatmap of z-

scored LFCs of astrocyte genes (from [37]). E   ) BrdU labeling of control and L48R FBNs. F   )

GO plot of top 15 pathways downregulated in L48R FBNs compared to control FBNs. F) GO

plot of significantly dysregulated genes in both L48R NPCs and FBNs compared to control.

Supplementary Figure 3 H3.3 deposition is altered in L48R cells, resulting in changes in

chromatin landscape. A) log2 normalized counts of *H3-3B* and *H3-3B* from the NPC RNA-seq.

B) RT-qPCR analysis of *H3-3B* in iPSCs (p-value = 0.3348), NPCs (p-value = 0.04321), and

FBNs (p-value = 0.0006555). B) RT-qPCR validation of *H3-3B* expression in iPSCs, NPCs and

FBNs. p = 0.051 in iPSCs, 0.0015 NPCs, p=0.4472 in FBNS. C)   Western blot on nuclear

extracted H3.3 and H4 as a loading control. C) Western blot probing for H3.3 in cytoplasmic and

chromatin-bound fractions. D) IF for H3.3 and MAP2 in Control and L48R FBNs (left).

Quantification of H3.3 fluorescence (right). p-value < 2.2e-16. E) Metaplot comparison of

ATAC-seq average signal from control and L48R NPCs at all peaks, showing the region

spanning +/- 2kb from the TSS ($n = 2$ control, 2 L48R biological replicates. F) Pie plot of

the genomic annotations of sites that are differentially accessible in L48R vs control NPCs

(identified using DiffBind) G) log2-normalized counts for selection of genes that were affected in

both ATAC- and RNA-seq. Genes to the left of the dashed line were down and those to the right

were upregulated. H) RNA- and ATAC-seq tracks for *RBFOX1*. I) RT-qPCR validation of

gene expression changes in iPSCs, NPCs and FBNs. p-values: iPSC, NPC = 0.0153, FBN =

0.1113 .

1006

Supplementary Figure 4 Cell identity analysis in 3D DFBOs corroborates 2D findings. A)

Schematic of the dorsal forebrain organoid protocol employed in this study. Organoids were

collected for sectioning at D25, and for RT-qPCR analysis and sectioning at D62. B) Images

taken at 4X magnification of control (top) and L48R (bottom) organoids at D62 showing

differences in NE buds (denoted by errors). C) Quantification of IF for PAX6 (red) at D62 (top)

shown in Fig 6E. p-value = 0.3291 D) CTIP2 staining at D25 and D62, along with corrected

total fluorescence quantifications of n = 6 organoids (control) or n = 4 organoids (L48R). E)

Quantification of CTIP2 shown in D. D25 p-value = 0.1045, D62 p-value = 0.138. E. a-i)

Intrinsic electrophysiological properties of neurons in control and L48R DFBOs. **a.**

neuron from a control **(a)** and an L48R DFBO **(b)** filled with Alexa Fluor 488, depicting a

bright soma with processes. **c, d.** Whole-cell patch clamp recording of neurons showing

spontaneous activity from control and L48R DFBOs in current-clamp mode without

current injection. **e-i**. Bar graphs showing mean (+SEM) and scatter plots comparison of

resting membrane potential (**e**), input resistance (**f**), maximum spike frequency during

current injection (**g**), sEPSC amplitude **(h)**, and sEPSC frequency **(i)** between control

and L48R DFBOs. (*$P < 0.05$, [NS] $P > 0.05$; student t test). Data is collected from two

biological replicates, 45 neurons from Control and 45 neutrons from L48R are analyzed.

1024

1025

1026 **Supplementary Files**

Supplementary File 1. DESeq analysis of RNA-seq experiments in iPSCs, NPCs, and FBNs.

Supplementary File 2. DiffBind results with significant changes in chromatin accessibility from the ATAC-seq in NPCs. Genomic regions were annotated with ChIPSeeker.

REFERENCES

1. Ropers, H.H. (2010). Genetics of early onset cognitive impairment. Annu. Rev. Genomics Hum. Genet. *11*, 161–187. https://doi.org/10.1146/annurev-genom-082509-141640.

2. Harris, J.R., Gao, C.W., Britton, J.F., Applegate, C.D., Bjornsson, H.T., and Fahrner, J.A. (2024). Five years of experience in the Epigenetics and Chromatin Clinic: what have we learned and where do we go from here? Hum. Genet. *143*, 607–624. https://doi.org/10.1007/s00439-023-02537-1.

3. Tessadori, F., Giltay, J.C., Hurst, J.A., Massink, M.P., Duran, K., Vos, H.R., van Es, R.M., Deciphering Developmental Disorders Study, Scott, R.H., van Gassen, K.L.I., et al. (2017). Germline mutations affecting the histone H4 core cause a developmental syndrome by altering DNA damage response and cell cycle control. Nat. Genet. *49*, 1642–1646. https://doi.org/10.1038/ng.3956.

4. Tessadori, F., Duran, K., Knapp, K., Fellner, M., Deciphering Developmental Disorders Study, Smithson, S., Beleza Meireles, A., Elting, M.W., Waisfisz, Q., O'Donnell-Luria, A., et al. (2022). Recurrent de novo missense variants across multiple histone H4 genes underlie a neurodevelopmental syndrome. Am. J. Hum. Genet. *109*, 750–758. https://doi.org/10.1016/j.ajhg.2022.02.003.

5. Bryant, L., Li, D., Cox, S.G., Marchione, D., Joiner, E.F., Wilson, K., Janssen, K., Lee, P., March, M.E., Nair, D., et al. (2020). Histone H3.3 beyond cancer: Germline mutations in Histone 3 Family 3A and 3B cause a previously unidentified neurodegenerative disorder in 46 patients. Sci. Adv. *6*, eabc9207. https://doi.org/10.1126/sciadv.abc9207.

6. Maver, A., Čuturilo, G., Ruml, S.J., and Peterlin, B. (2019). Clinical Next Generation Sequencing Reveals an H3F3A Gene as a New Potential Gene Candidate for Microcephaly Associated with Severe Developmental Delay, Intellectual Disability and Growth Retardation. Balk. J. Med. Genet. BJMG *22*, 65–68. https://doi.org/10.2478/bjmg-2019-0028.

7. Okur, V., Chen, Z., Vossaert, L., Peacock, S., Rosenfeld, J., Zhao, L., Du, H., Calamaro, E., Gerard, A., Zhao, S., et al. (2021). De novo variants in H3-3A and H3-3B are associated with neurodevelopmental delay, dysmorphic features, and structural brain abnormalities. NPJ Genomic Med. *6*, 104. https://doi.org/10.1038/s41525-021-00268-8.

8. Layo-Carris, D.E., Lubin, E.E., Sangree, A.K., Clark, K.J., Durham, E.L., Gonzalez,
E.M., Smith, S., Angireddy, R., Wang, X.M., Weiss, E., et al. (2024). Expanded
phenotypic spectrum of neurodevelopmental and neurodegenerative disorder
Bryant-Li-Bhoj syndrome with 38 additional individuals. Eur. J. Hum. Genet.
https://doi.org/10.1038/s41431-024-01610-1.

9. Alves, C. a. P.F., Sherbini, O., D'Arco, F., Steel, D., Kurian, M.A., Radio, F.C.,
Ferrero, G.B., Carli, D., Tartaglia, M., Balci, T.B., et al. (2022). Brain Abnormalities
in Patients with Germline Variants in H3F3: Novel Imaging Findings and Neurologic
Symptoms Beyond Somatic Variants and Brain Tumors. AJNR Am. J. Neuroradiol.
43, 1048–1053. https://doi.org/10.3174/ajnr.A7555.

10. Funk, O.H., Qalieh, Y., Doyle, D.Z., Lam, M.M., and Kwan, K.Y. (2022). Postmitotic
accumulation of histone varíant H3.3 in new cortical neurons establishes neuronal
chromatin, transcriptome, and identity. Proc. Natl. Acad. Sci. U. S. A. *119*,
e2116956119. https://doi.org/10.1073/pnas.2116956119.

11. Jin, C., Zang, C., Wei, G., Cui, K., Peng, W., Zhao, K., and Felsenfeld, G. (2009).
H3.3/H2A.Z double variant-containing nucleosomes mark "nucleosome-free regions"
of active promoters and other regulatory regions. Nat. Genet. *41*, 941–945.
https://doi.org/10.1038/ng.409.

12. Goldberg, A.D., Banaszynski, L.A., Noh, K.-M., Lewis, P.W., Elsaesser, S.J.,
Stadler, S., Dewell, S., Law, M., Guo, X., Li, X., et al. (2010). Distinct factors control
histone variant H3.3 localization at specific genomic regions. Cell *140*, 678–691.
https://doi.org/10.1016/j.cell.2010.01.003.

13. Chen, P., Zhao, J., Wang, Y., Wang, M., Long, H., Liang, D., Huang, L., Wen, Z., Li,
W., Li, X., et al. (2013). H3.3 actively marks enhancers and primes gene
transcription via opening higher-ordered chromatin. Genes Dev. *27*, 2109–2124.
https://doi.org/10.1101/gad.222174.113.

14. Jin, C., and Felsenfeld, G. (2007). Nucleosome stability mediated by histone
variants H3.3 and H2A.Z. Genes Dev. *21*, 1519–1529.
https://doi.org/10.1101/gad.1547707.

15. Chow, C.-M., Georgiou, A., Szutorisz, H., Maia e Silva, A., Pombo, A., Barahona, I.,
Dargelos, E., Canzonetta, C., and Dillon, N. (2005). Variant histone H3.3 marks
promoters of transcriptionally active genes during mammalian cell division. EMBO
Rep. *6*, 354–360. https://doi.org/10.1038/sj.embor.7400366.

16. Jin, C., and Felsenfeld, G. (2006). Distribution of histone H3.3 in hematopoietic cell
lineages. Proc. Natl. Acad. Sci. U. S. A. *103*, 574–579.
https://doi.org/10.1073/pnas.0509974103.

17. Deaton, A.M., Gómez-Rodríguez, M., Mieczkowski, J., Tolstorukov, M.Y., Kundu, S.,
Sadreyev, R.I., Jansen, L.E., and Kingston, R.E. (2016). Enhancer regions show

high histone H3.3 turnover that changes during differentiation. eLife *5*, e15316.
https://doi.org/10.7554/eLife.15316.

18. Kraushaar, D.C., Jin, W., Maunakea, A., Abraham, B., Ha, M., and Zhao, K. (2013).
Genome-wide incorporation dynamics reveal distinct categories of turnover for the
histone variant H3.3. Genome Biol. *14*, R121. https://doi.org/10.1186/gb-2013-14-
10-r121.

19. Herre, M., and Korb, E. (2019). The chromatin landscape of neuronal plasticity. Curr.
Opin. Neurobiol. *59*, 79–86. https://doi.org/10.1016/j.conb.2019.04.006.

20. Maze, I., Wenderski, W., Noh, K.-M., Bagot, R.C., Tzavaras, N., Purushothaman, I.,
Elsässer, S.J., Guo, Y., Ionete, C., Hurd, Y.L., et al. (2015). Critical Role of Histone
Turnover in Neuronal Transcription and Plasticity. Neuron *87*, 77–94.
https://doi.org/10.1016/j.neuron.2015.06.014.

21. Piña, B., and Suau, P. (1987). Changes in histones H2A and H3 variant composition
in differentiating and mature rat brain cortical neurons. Dev. Biol. *123*, 51–58.
https://doi.org/10.1016/0012-1606(87)90426-x.

22. Lan, F., and Shi, Y. (2015). Histone H3.3 and cancer: A potential reader connection.
Proc. Natl. Acad. Sci. U. S. A. *112*, 6814–6819.
https://doi.org/10.1073/pnas.1418996111.

23. Kallappagoudar, S., Yadav, R.K., Lowe, B.R., and Partridge, J.F. (2015). Histone H3
mutations--a special role for H3.3 in tumorigenesis? Chromosoma *124*, 177–189.
https://doi.org/10.1007/s00412-015-0510-4.

24. Deshmukh, S., Ptack, A., Krug, B., and Jabado, N. (2022). Oncohistones: a
roadmap to stalled development. FEBS J. *289*, 1315–1328.
https://doi.org/10.1111/febs.15963.

25. Schwartzentruber, J., Korshunov, A., Liu, X.-Y., Jones, D.T.W., Pfaff, E., Jacob, K.,
Sturm, D., Fontebasso, A.M., Quang, D.-A.K., Tönjes, M., et al. (2012). Driver
mutations in histone H3.3 and chromatin remodelling genes in paediatric
glioblastoma. Nature *482*, 226–231. https://doi.org/10.1038/nature10833.

26. Khuong-Quang, D.-A., Buczkowicz, P., Rakopoulos, P., Liu, X.-Y., Fontebasso,
A.M., Bouffet, E., Bartels, U., Albrecht, S., Schwartzentruber, J., Letourneau, L., et
al. (2012). K27M mutation in histone H3.3 defines clinically and biologically distinct
subgroups of pediatric diffuse intrinsic pontine gliomas. Acta Neuropathol. (Berl.)
124, 439–447. https://doi.org/10.1007/s00401-012-0998-0.

27. Wu, G., Broniscer, A., McEachron, T.A., Lu, C., Paugh, B.S., Becksfort, J., Qu, C.,
Ding, L., Huether, R., Parker, M., et al. (2012). Somatic histone H3 alterations in
pediatric diffuse intrinsic pontine gliomas and non-brainstem glioblastomas. Nat.
Genet. *44*, 251–253. https://doi.org/10.1038/ng.1102.

28. Behjati, S., Tarpey, P.S., Presneau, N., Scheipl, S., Pillay, N., Van Loo, P., Wedge,
D.C., Cooke, S.L., Gundem, G., Davies, H., et al. (2013). Distinct H3F3A and
H3F3B driver mutations define chondroblastoma and giant cell tumor of bone. Nat.
Genet. *45*, 1479–1482. https://doi.org/10.1038/ng.2814.

29. Khazaei, S., Chen, C.C.L., Andrade, A.F., Kabir, N., Azarafshar, P., Morcos, S.M.,
França, J.A., Lopes, M., Lund, P.J., Danieau, G., et al. (2023). Single substitution in
H3.3G34 alters DNMT3A recruitment to cause progressive neurodegeneration. Cell
186, 1162-1178.e20. https://doi.org/10.1016/j.cell.2023.02.023.

30. Yang, Q., Hong, Y., Zhao, T., Song, H., and Ming, G.-L. (2022). What Makes
Organoids Good Models of Human Neurogenesis? Front. Neurosci. *16*, 872794.
https://doi.org/10.3389/fnins.2022.872794.

31. Maguire, J.A., Cardenas-Diaz, F.L., Gadue, P., and French, D.L. (2019). Highly
Efficient CRISPR-Cas9-Mediated Genome Editing in Human Pluripotent Stem Cells.
Curr. Protoc. Stem Cell Biol. *48*, e64. https://doi.org/10.1002/cpsc.64.

32. Bray, N.L., Pimentel, H., Melsted, P., and Pachter, L. (2016). Near-optimal
probabilistic RNA-seq quantification. Nat. Biotechnol. *34*, 525–527.
https://doi.org/10.1038/nbt.3519.

33. Love, M.I., Huber, W., and Anders, S. (2014). Moderated estimation of fold change
and dispersion for RNA-seq data with DESeq2. Genome Biol. *15*, 550.
https://doi.org/10.1186/s13059-014-0550-8.

34. Blighe, Kevin, Nitesh, Turaga, Grionia, Andrea, Turkes, Emir, and Benjamin,
Ostendorf (2018). EnhancedVolcano: Publication-ready volcano plots with enhanced
colouring and labeling.

35. Yu, G., Wang, L.-G., Han, Y., and He, Q.-Y. (2012). clusterProfiler: an R package for
comparing biological themes among gene clusters. Omics J. Integr. Biol. *16*, 284–
287. https://doi.org/10.1089/omi.2011.0118.

36. Supek, F., Bošnjak, M., Škunca, N., and Šmuc, T. (2011). REVIGO summarizes and
visualizes long lists of gene ontology terms. PloS One *6*, e21800.
https://doi.org/10.1371/journal.pone.0021800.

37. Franzén, O., Gan, L.-M., and Björkegren, J.L.M. (2019). PanglaoDB: a web server
for exploration of mouse and human single-cell RNA sequencing data. Database J.
Biol. Databases Curation *2019*, baz046. https://doi.org/10.1093/database/baz046.

38. Langmead, B., and Salzberg, S.L. (2012). Fast gapped-read alignment with Bowtie
2. Nat. Methods *9*, 357–359. https://doi.org/10.1038/nmeth.1923.

39. Broad Institute Picard Toolkit.

40. Zhang, Y., Liu, T., Meyer, C.A., Eeckhoute, J., Johnson, D.S., Bernstein, B.E.,
Nusbaum, C., Myers, R.M., Brown, M., Li, W., et al. (2008). Model-based analysis of
ChIP-Seq (MACS). Genome Biol. *9*, R137. https://doi.org/10.1186/gb-2008-9-9-
r137.

41. Yu, G., Wang, L.-G., and He, Q.-Y. (2015). ChIPseeker: an R/Bioconductor package
for ChIP peak annotation, comparison and visualization. Bioinformatics *31*, 2382–
2383. https://doi.org/10.1093/bioinformatics/btv145.

42. Wang, Q., Li, M., Wu, T., Zhan, L., Li, L., Chen, M., Xie, W., Xie, Z., Hu, E., Xu, S.,
et al. (2022). Exploring Epigenomic Datasets by ChIPseeker. Curr. Protoc. *2*, e585.
https://doi.org/10.1002/cpz1.585.

43. Ross-Innes, C.S., Stark, R., Teschendorff, A.E., Holmes, K.A., Ali, H.R., Dunning,
M.J., Brown, G.D., Gojis, O., Ellis, I.O., Green, A.R., et al. (2012). Differential
oestrogen receptor binding is associated with clinical outcome in breast cancer.
Nature *481*, 389–393. https://doi.org/10.1038/nature10730.

44. Ramírez, F., Ryan, D.P., Grüning, B., Bhardwaj, V., Kilpert, F., Richter, A.S., Heyne,
S., Dündar, F., and Manke, T. (2016). deepTools2: a next generation web server for
deep-sequencing data analysis. Nucleic Acids Res. *44*, W160-165.
https://doi.org/10.1093/nar/gkw257.

45. Birey, F., Andersen, J., Makinson, C.D., Islam, S., Wei, W., Huber, N., Fan, H.C.,
Metzler, K.R.C., Panagiotakos, G., Thom, N., et al. (2017). Assembly of functionally
integrated human forebrain spheroids. Nature *545*, 54–59.
https://doi.org/10.1038/nature22330.

46. Chambers, S.M., Fasano, C.A., Papapetrou, E.P., Tomishima, M., Sadelain, M., and
Studer, L. (2009). Highly efficient neural conversion of human ES and iPS cells by
dual inhibition of SMAD signaling. Nat. Biotechnol. *27*, 275–280.
https://doi.org/10.1038/nbt.1529.

47. Valli, E., Trazzi, S., Fuchs, C., Erriquez, D., Bartesaghi, R., Perini, G., and Ciani, E.
(2012). CDKL5, a novel MYCN-repressed gene, blocks cell cycle and promotes
differentiation of neuronal cells. Biochim. Biophys. Acta *1819*, 1173–1185.
https://doi.org/10.1016/j.bbagrm.2012.08.001.

48. Lecona, E., Rojas, L.A., Bonasio, R., Johnston, A., Fernández-Capetillo, O., and
Reinberg, D. (2013). Polycomb protein SCML2 regulates the cell cycle by binding
and modulating CDK/CYCLIN/p21 complexes. PLoS Biol. *11*, e1001737.
https://doi.org/10.1371/journal.pbio.1001737.

49. Flaherty, E., and Maniatis, T. (2020). The role of clustered protocadherins in
neurodevelopment and neuropsychiatric diseases. Curr. Opin. Genet. Dev. *65*, 144–
150. https://doi.org/10.1016/j.gde.2020.05.041.

50. Duclot, F., and Kabbaj, M. (2017). The Role of Early Growth Response 1 (EGR1) in
Brain Plasticity and Neuropsychiatric Disorders. Front. Behav. Neurosci. *11*, 35.
https://doi.org/10.3389/fnbeh.2017.00035.

51. Moreland, T., and Poulain, F.E. (2022). To Stick or Not to Stick: The Multiple Roles
of Cell Adhesion Molecules in Neural Circuit Assembly. Front. Neurosci. *16*, 889155.
https://doi.org/10.3389/fnins.2022.889155.

52. Steele, J.L., Morrow, M.M., Sarnat, H.B., Alkhunaizi, E., Brandt, T., Chitayat, D.A.,
DeFilippo, C.P., Douglas, G.V., Dubbs, H.A., Elloumi, H.Z., et al. (2022).
Semaphorin-Plexin Signaling: From Axonal Guidance to a New X-Linked Intellectual
Disability Syndrome. Pediatr. Neurol. *126*, 65–73.
https://doi.org/10.1016/j.pediatrneurol.2021.10.008.

53. Cheng, H.J., Bagri, A., Yaron, A., Stein, E., Pleasure, S.J., and Tessier-Lavigne, M.
(2001). Plexin-A3 mediates semaphorin signaling and regulates the development of
hippocampal axonal projections. Neuron *32*, 249–263.
https://doi.org/10.1016/s0896-6273(01)00478-0.

54. Fogel, B.L., Wexler, E., Wahnich, A., Friedrich, T., Vijayendran, C., Gao, F.,
Parikshak, N., Konopka, G., and Geschwind, D.H. (2012). RBFOX1 regulates both
splicing and transcriptional networks in human neuronal development. Hum. Mol.
Genet. *21*, 4171–4186. https://doi.org/10.1093/hmg/dds240.

55. Ou, Jianhong ATACseqTFEA: Transcription Factor Enrichment Analysis for ATAC-
seq. Version R package version 1.6.0.

56. Liska, O., Bohár, B., Hidas, A., Korcsmáros, T., Papp, B., Fazekas, D., and Ari, E.
(2022). TFLink: an integrated gateway to access transcription factor-target gene
interactions for multiple species. Database J. Biol. Databases Curation *2022*,
baac083. https://doi.org/10.1093/database/baac083.

57. Penisson, M., Ladewig, J., Belvindrah, R., and Francis, F. (2019). Genes and
Mechanisms Involved in the Generation and Amplification of Basal Radial Glial
Cells. Front. Cell. Neurosci. *13*, 381. https://doi.org/10.3389/fncel.2019.00381.

58. Lynch, D.C., Revil, T., Schwartzentruber, J., Bhoj, E.J., Innes, A.M., Lamont, R.E.,
Lemire, E.G., Chodirker, B.N., Taylor, J.P., Zackai, E.H., et al. (2014). Disrupted
auto-regulation of the spliceosomal gene SNRPB causes cerebro-costo-mandibular
syndrome. Nat. Commun. *5*, 4483. https://doi.org/10.1038/ncomms5483.

59. Lewis, P.W., Müller, M.M., Koletsky, M.S., Cordero, F., Lin, S., Banaszynski, L.A.,
Garcia, B.A., Muir, T.W., Becher, O.J., and Allis, C.D. (2013). Inhibition of PRC2
activity by a gain-of-function H3 mutation found in pediatric glioblastoma. Science
340, 857–861. https://doi.org/10.1126/science.1232245.

60. Shi, L., Shi, J., Shi, X., Li, W., and Wen, H. (2018). Histone H3.3 G34 Mutations
Alter Histone H3K36 and H3K27 Methylation In Cis. J. Mol. Biol. *430*, 1562–1565.
https://doi.org/10.1016/j.jmb.2018.04.014.

61. Parenti, I., Rabaneda, L.G., Schoen, H., and Novarino, G. (2020).
Neurodevelopmental Disorders: From Genetics to Functional Pathways. Trends
Neurosci. *43*, 608–621. https://doi.org/10.1016/j.tins.2020.05.004.

62. Shah, P.S., Weksberg, R., and Chitayat, D. (2006). Overgrowth with severe
developmental delay following IVF/ICSI: A newly recognized syndrome? Am. J.
Med. Genet. A. *140*, 1312–1315. https://doi.org/10.1002/ajmg.a.31274.

63. Wenzel, T.J., and Mousseau, D.D. (2024). Brain organoids engineered to give rise
to glia and neural networks after 90 days in culture exhibit human-specific
proteoforms. Front. Cell. Neurosci. *18*, 1383688.
https://doi.org/10.3389/fncel.2024.1383688.

64. Mulder, L.A., Depla, J.A., Sridhar, A., Wolthers, K., Pajkrt, D., and Vieira de Sá, R.
(2023). A beginner's guide on the use of brain organoids for neuroscientists: a
systematic review. Stem Cell Res. Ther. *14*, 87. https://doi.org/10.1186/s13287-023-
03302-x.

65. James, O.G., Selvaraj, B.T., Magnani, D., Burr, K., Connick, P., Barton, S.K.,
Vasistha, N.A., Hampton, D.W., Story, D., Smigiel, R., et al. (2021). iPSC-derived
myelinoids to study myelin biology of humans. Dev. Cell *56*, 1346-1358.e6.
https://doi.org/10.1016/j.devcel.2021.04.006.

66. Fasipe, O.J., Agede, O.A., and Enikuomehin, A.C. (2020). Announcing the novel
class of GABA-A receptor selective positive allosteric modulator antidepressants.
Future Sci. OA *7*, FSO654. https://doi.org/10.2144/fsoa-2020-0108.

67. Bankhead, P., Loughrey, M.B., Fernández, J.A., Dombrowski, Y., McArt, D.G.,
Dunne, P.D., McQuaid, S., Gray, R.T., Murray, L.J., Coleman, H.G., et al. (2017).
QuPath: Open source software for digital pathology image analysis. Sci. Rep. *7*,
16878. https://doi.org/10.1038/s41598-017-17204-5.

68. Takeichi, M. (2007). The cadherin superfamily in neuronal connections and
interactions. Nat. Rev. Neurosci. *8*, 11–20. https://doi.org/10.1038/nrn2043.

69. Polanco, J., Reyes-Vigil, F., Weisberg, S.D., Dhimitruka, I., and Brusés, J.L. (2021).
Differential Spatiotemporal Expression of Type I and Type II Cadherins Associated
With the Segmentation of the Central Nervous System and Formation of Brain
Nuclei in the Developing Mouse. Front. Mol. Neurosci. *14*, 633719.
https://doi.org/10.3389/fnmol.2021.633719.

70. Kiefel, H., Bondong, S., Hazin, J., Ridinger, J., Schirmer, U., Riedle, S., and
Altevogt, P. (2012). L1CAM: a major driver for tumor cell invasion and motility. Cell
Adhes. Migr. *6*, 374–384. https://doi.org/10.4161/cam.20832.

71. Robertson, S.P., Twigg, S.R.F., Sutherland-Smith, A.J., Biancalana, V., Gorlin, R.J.,
Horn, D., Kenwrick, S.J., Kim, C.A., Morava, E., Newbury-Ecob, R., et al. (2003).
Localized mutations in the gene encoding the cytoskeletal protein filamin A cause
diverse malformations in humans. Nat. Genet. *33*, 487–491.
https://doi.org/10.1038/ng1119.

72. Feng, Y., Chen, M.H., Moskowitz, I.P., Mendonza, A.M., Vidali, L., Nakamura, F.,
Kwiatkowski, D.J., and Walsh, C.A. (2006). Filamin A (FLNA) is required for cell-cell
contact in vascular development and cardiac morphogenesis. Proc. Natl. Acad. Sci.
U. S. A. *103*, 19836–19841. https://doi.org/10.1073/pnas.0609628104.

73. Ciani, L., Boyle, K.A., Dickins, E., Sahores, M., Anane, D., Lopes, D.M., Gibb, A.J.,
and Salinas, P.C. (2011). Wnt7a signaling promotes dendritic spine growth and
synaptic strength through Ca^{2+}/Calmodulin-dependent protein kinase II. Proc. Natl.
Acad. Sci. U. S. A. *108*, 10732–10737. https://doi.org/10.1073/pnas.1018132108.

74. Caracci, M.O., Avila, M.E., Espinoza-Cavieres, F.A., López, H.R., Ugarte, G.D., and
De Ferrari, G.V. (2021). Wnt/β-Catenin-Dependent Transcription in Autism
Spectrum Disorders. Front. Mol. Neurosci. *14*, 764756.
https://doi.org/10.3389/fnmol.2021.764756.

75. Qu, Q., Sun, G., Murai, K., Ye, P., Li, W., Asuelime, G., Cheung, Y.-T., and Shi, Y.
(2013). Wnt7a regulates multiple steps of neurogenesis. Mol. Cell. Biol. *33*, 2551–
2559. https://doi.org/10.1128/MCB.00325-13.

76. Vogt, D.L., Gray, C.D., Young, W.S., Orellana, S.A., and Malouf, A.T. (2007).
ARHGAP4 is a novel RhoGAP that mediates inhibition of cell motility and axon
outgrowth. Mol. Cell. Neurosci. *36*, 332–342.
https://doi.org/10.1016/j.mcn.2007.07.004.

77. Duan, Y., Wang, S.-H., Song, J., Mironova, Y., Ming, G., Kolodkin, A.L., and Giger,
R.J. (2014). Semaphorin 5A inhibits synaptogenesis in early postnatal- and adult-
born hippocampal dentate granule cells. eLife *3*, e04390.
https://doi.org/10.7554/eLife.04390.

78. Grebbin, B.M., Hau, A.-C., Groß, A., Anders-Maurer, M., Schramm, J., Koss, M.,
Wille, C., Mittelbronn, M., Selleri, L., and Schulte, D. (2016). Pbx1 is required for
adult subventricular zone neurogenesis. Dev. Camb. Engl. *143*, 2281–2291.
https://doi.org/10.1242/dev.128033.

79. Villaescusa, J.C., Li, B., Toledo, E.M., Rivetti di Val Cervo, P., Yang, S., Stott, S.R.,
Kaiser, K., Islam, S., Gyllborg, D., Laguna-Goya, R., et al. (2016). A PBX1
transcriptional network controls dopaminergic neuron development and is impaired
in Parkinson's disease. EMBO J. *35*, 1963–1978.
https://doi.org/10.15252/embj.201593725.

80. Colasante, G., Collombat, P., Raimondi, V., Bonanomi, D., Ferrai, C., Maira, M.,
Yoshikawa, K., Mansouri, A., Valtorta, F., Rubenstein, J.L.R., et al. (2008). Arx is a

direct target of Dlx2 and thereby contributes to the tangential migration of
GABAergic interneurons. J. Neurosci. Off. J. Soc. Neurosci. *28*, 10674–10686.
https://doi.org/10.1523/JNEUROSCI.1283-08.2008.

81. Lim, Y., Cho, I.-T., Shi, X., Grinspan, J.B., Cho, G., and Golden, J.A. (2019). Arx
Expression Suppresses Ventralization of the Developing Dorsal Forebrain. Sci. Rep.
9, 226. https://doi.org/10.1038/s41598-018-36194-6.

82. Jacobs, W.B., Govoni, G., Ho, D., Atwal, J.K., Barnabe-Heider, F., Keyes, W.M.,
Mills, A.A., Miller, F.D., and Kaplan, D.R. (2005). p63 is an essential proapoptotic
protein during neural development. Neuron *48*, 743–756.
https://doi.org/10.1016/j.neuron.2005.10.027.

83. Bishop, K.M., Rubenstein, J.L.R., and O'Leary, D.D.M. (2002). Distinct actions of
Emx1, Emx2, and Pax6 in regulating the specification of areas in the developing
neocortex. J. Neurosci. Off. J. Soc. Neurosci. *22*, 7627–7638.
https://doi.org/10.1523/JNEUROSCI.22-17-07627.2002.

84. Lin, L., Sun, W., Throesch, B., Kung, F., Decoster, J.T., Berner, C.J., Cheney, R.E.,
Rudy, B., and Hoffman, D.A. (2013). DPP6 regulation of dendritic morphogenesis
impacts hippocampal synaptic development. Nat. Commun. *4*, 2270.
https://doi.org/10.1038/ncomms3270.

85. Malloy, C., Ahern, M., Lin, L., and Hoffman, D.A. (2022). Neuronal Roles of the
Multifunctional Protein Dipeptidyl Peptidase-like 6 (DPP6). Int. J. Mol. Sci. *23*, 9184.
https://doi.org/10.3390/ijms23169184.

86. Naujock, M., Speidel, A., Fischer, S., Kizner, V., Dorner-Ciossek, C., and Gillardon,
F. (2020). Neuronal Differentiation of Induced Pluripotent Stem Cells from
Schizophrenia Patients in Two-Dimensional and in Three-Dimensional Cultures
Reveals Increased Expression of the Kv4.2 Subunit DPP6 That Contributes to
Decreased Neuronal Activity. Stem Cells Dev. *29*, 1577–1587.
https://doi.org/10.1089/scd.2020.0082.

87. Karayannis, T., Au, E., Patel, J.C., Kruglikov, I., Markx, S., Delorme, R., Héron, D.,
Salomon, D., Glessner, J., Restituito, S., et al. (2014). Cntnap4 differentially
contributes to GABAergic and dopaminergic synaptic transmission. Nature *511*,
236–240. https://doi.org/10.1038/nature13248.

88. Young, S.Z., Lafourcade, C.A., Platel, J.-C., Lin, T.V., and Bordey, A. (2014).
GABAergic striatal neurons project dendrites and axons into the postnatal
subventricular zone leading to calcium activity. Front. Cell. Neurosci. *8*, 10.
https://doi.org/10.3389/fncel.2014.00010.

89. Hotz, A.L., Jamali, A., Rieser, N.N., Niklaus, S., Aydin, E., Myren-Svelstad, S., Lalla,
L., Jurisch-Yaksi, N., Yaksi, E., and Neuhauss, S.C.F. (2022). Loss of glutamate
transporter eaat2a leads to aberrant neuronal excitability, recurrent epileptic

seizures, and basal hypoactivity. Glia *70*, 196–214.
https://doi.org/10.1002/glia.24106.

90. Rylaarsdam, L., Rakotomamonjy, J., Pope, E., and Guemez-Gamboa, A. (2024).
iPSC-derived models of PACS1 syndrome reveal transcriptional and functional
deficits in neuron activity. Nat. Commun. *15*, 827. https://doi.org/10.1038/s41467-
024-44989-7.

91. Butler, K.M., Moody, O.A., Schuler, E., Coryell, J., Alexander, J.J., Jenkins, A., and
Escayg, A. (2018). De novo variants in GABRA2 and GABRA5 alter receptor
function and contribute to early-onset epilepsy. Brain J. Neurol. *141*, 2392–2405.
https://doi.org/10.1093/brain/awy171.

92. Kim, S., Kim, H., and Um, J.W. (2018). Synapse development organized by
neuronal activity-regulated immediate-early genes. Exp. Mol. Med. *50*, 1–7.
https://doi.org/10.1038/s12276-018-0025-1.

1375

1376

1377

Graphical Abstract

Figure 1

Figure S1

Figure 2

Figure S2

Figure 3

Figure S3

Figure 4

Figure 5

Figure 6

Placeholder, need 2nd rep

Figure 7

A

Day 25
Collect organoids
for sectioning

Day 62
Collect organoids
for sectioning +
RNA for RT-qPCR

DFBO formation
media

DFBO expansion media

DFBO differentiation
media

DFBO maintenance
media

B

Control

L48R

Day 25

Day 62

C

Control

L48R

CTIP2

Figure S4