Title: Water volume, biological and PCR replicates influence the characterization of deep-sea

pelagic fish communities.

Running title: Sampling eDNA in the deep sea.

Authors:

Pedro A. Peres[1], Heather Bracken-Grissom[1,2]

1 Institute of Environment, Department of Biological Sciences, Florida International University,

3000 NE 151st St, North Miami, FL, 33181 USA.

2 Department of Invertebrate Zoology, Smithsonian Institution, National Museum of Natural

History, Washington, WA, 20013-7012 USA.

Contact information: Pedro A. Peres: pedro.peres27@gmail.com ; Heather Bracken-Grissom:

hbracken@fiu.edu

Author contributions: PAP and HBG – conception and design of the study; PAP and HBG –

data collection; PAP – formal analyses; HBG – supervision; PAP and HBG – funding

acquisition; PAP – writing original draft; PAP and HBG – writing review and editing.

Data availability statement: All data used is available in Supplemental Materials.

Funding statement: This material is based upon work supported by the National Science

Foundation under Grant No. HRD-1547798 and Grant No. HRD-2111661. These NSF Grants

were awarded to Florida International University as part of the Centers of Research Excellence in

Science and Technology (CREST) Program. This research was also made possible by a grant

from the Office of Naval Research (#98288/1184488/2), the Gulf of Mexico Research Initiative

(DEEPEND|RESTORE consortium), and the National Oceanic and Atmospheric

Administration's RESTORE Science Program under award NA19NOS4510193 to Florida

International University.

Conflict of interest disclosure: Authors declare no conflict of interest.

Ethics statement: Authors confirm that the manuscript has not been submitted elsewhere, and

that all research meets the ethical guidelines of the study country.

Acknowledgments: We thank all researchers involved in the DEEPEND|RESTORE consortium

for their support in sample collections. This is contribution #X from the Coastlines and Oceans

Division of the Institute of Environment at Florida International University.

ABSTRACT

The pelagic deep sea is challenging to investigate due to logistical constraints regarding

access and collection of samples, however environmental DNA (eDNA) can potentially

revolutionize our understanding of this ecosystem. Although advancements are being made

regarding technology and eDNA samplers, many countries and research groups still do not have

access to modern equipment. At the same time, no investigation has been performed to assess the

impact of different experimental designs using common gear found on research vessels (i.e.,

CTD mounted with Niskin bottles). Here, we investigated the effects of sampled water volume,

biological and PCR replicates in characterizing deep-sea pelagic biodiversity. Samples were

collected at 450m depth at night in the northern Gulf of Mexico using Niskin bottles, and we

targeted the fish community using the MiFish primer (12S rRNA). Our results show that 1L is

insufficient to characterize deep-sea pelagic fish communities. The treatments 5L and 10L

detected similar communities and numbers of species, but 10L detected more exact sequence

variants (ESVs). Five biological replicates can detect up to 80% of the species detected in the

water collected in both 5L and 10L treatments. Extrapolation analysis indicates that 10 replicates

would also potentially recover 100% of the species. More biological replicates would be

necessary to detect all ESVs. PCR replicates also had an important role in taxa and ESV

detection, being the best strategy to perform more replicates if you filter less water (optimal

number inversely proportional to the water volume filtered). We suggest that future studies

collect at least 5L, 5 field replicates, and 5-10 PCR replicates to adequately investigate deep-sea

pelagic biodiversity using eDNA, considering the limitations we mention. Our study provides

guidance for future eDNA studies and a potential route to democratize eDNA studies at a global

scale.

Key-words: deep sea, replicates, MiFish, Gulf of Mexico, ESV

INTRODUCTION

The use of DNA extracted from environmental samples (environmental DNA, eDNA) has

become a common method to describe biodiversity and test ecological hypotheses (Dickie et al.,

2018; Ruppert et al., 2019). Because this is a non-invasive approach, it has also gained

popularity to inform conservation and management practices (Barnes & Turner, 2016; Beng &

Corlett, 2020). eDNA can be particularly useful for the documentation of species diversity when

a habitat is hard to access, or the target organisms are very small (Stat et al., 2017). Although this

method is becoming a popular tool for coastal systems, it is less frequently employed to

characterize deep-sea pelagic ecosystems (>200 m depth). The deep-sea, including the midwater,

is the largest habitat on Earth but one of the least understood (Ramirez-Llodra et al., 2010).

Deep-pelagic animals participate in diel vertical migration, the largest migration on the planet in

terms of biomass, and play a critical role in the biological pump and transport of nutrients, acting

as the lifeline between the surface and deep ecosystems. In addition, there is potential economic

importance represented by natural resources exploration, from fisheries to deep-sea mining

(Thurber et al., 2014). Investigations of the deep sea and associated diversity are challenging due

to logistical constraints regarding access, collecting, and filtering water samples. In this context,

eDNA has the potential to revolutionize our understanding of this enigmatic and vast habitat.

Advances in robotics are improving sampling strategies, as seen using autonomous vehicles in

combination with *in situ* filtration for both surface and deep-pelagic communities (Billings et al.,

2017; Yamahara et al., 2019; Yoerger et al., 2021; Govindarajan et al., 2022). However, this type

of technology might not be accessible for many scientific groups due to the costs associated with

specialized equipment or trained personnel, limiting the global use of this approach

(Govindarajan et al., 2022). Additionally, the use of specialized equipment and/or numerous

materials for eDNA sampling/filtering is not possible on all vessels due to limited space and

technological capabilities. Until advanced technologies are available for many research groups,

there is a need to further investigate experimental designs that use common equipment with

limited resources. This can facilitate the number of research groups performing eDNA in the

deep sea, thus enabling large-scale collaborative studies.

Many eDNA studies, including the ones investigating the deep-pelagic habitat, use Niskin bottles

to collect water samples, a common piece of equipment on research vessels. Usually, Niskin

bottles are coupled with a conductivity temperature depth (CTD) rosette and triggered at a

specific depth and location (McClenaghan et al., 2020; Govindarajan et al., 2021; Visser et al.,

2021; Govindarajan et al., 2023). After collection, water samples are filtered, and the filters are

preserved for future processing. Although there is some general guidance for eDNA studies (e.g.,

Dickie et al., 2018), little is discussed about specific sampling schemes focused on mid- and

deep-water pelagic systems. If using Niskin bottles to collect water is a viable option, the amount

of water and the number of biological replicates becomes a limitation because these factors are

constrained by the size and number of bottles available. For instance, filtering many replicates of

large amounts of water can be a laborious and sometimes impossible task (i.e. time constraints,

filter clogging, and pump capacity). These limitations should be central in discussing deep-

pelagic eDNA explorations as they will dictate the experimental design, representing a first,

critical step in eDNA studies.

Besides equipment limitations, the pelagic deep-sea environment presents other unique

challenges to consider when designing eDNA-based investigations. For instance, the magnitude

of this habitat and often low density of individuals, can result in low levels of eDNA. Previous

work has suggested that higher volumes of water, biological and PCR replicates might be

necessary in offshore environments, where DNA can be sparsely distributed (Kumar et al., 2021;

Cote et al., 2023); however, no experimental study has proposed or investigated potential

universal standards (water volume, biological and PCR replication) for deep-pelagic biodiversity

studies. Although obvious, one should not assume that standard sampling methods from coastal

systems are sufficient or translatable for deep-pelagic waters.

Here, we investigate the current methodological knowledge gaps and provide guidance for future

deep-pelagic eDNA studies. We test the impact of sampled water volume, biological and PCR

replicates in assessing deep-sea fish pelagic biodiversity using common equipment found on

research vessels (CTD rosettes and Niskin bottles). We collected 1L, 5L, and 10L simultaneously

within the deep-scattering layer (450 m depth) in the northern Gulf of Mexico which was

targeted using acoustic methods. Each treatment was represented by five biological replicates (15

samples total). We used MiFish primer set and aimed for 10 PCR replicates per biological

replicate and investigated species and exact sequence variant (ESV) detection. This experimental

design allowed us to detect the optimal sampling scheme, acknowledging the challenges

associated with pelagic deep-sea sampling and the limitations of using accessible gear.

MATERIALS AND METHODS

Experimental design

We investigated the effects of water volume and biological and PCR replicates by examining

samples collected simultaneously at 450m depth (see below). Our experimental design consisted

of three treatments: 1L, 5L, 10L. Each treatment consisted of five biological replicates (n = 15)

plus one negative control. The negative control is represented by 1L of ultrapure water filtered

using the same apparatus that was used for the biological samples (see below). To investigate the

effects of technical replicates, 10 PCR replicates were run for each biological replicate and the

negative control when possible (n = 160).

Sampling and shipboard processing

Water samples were collected during a cruise on R/V Point Sur in the Northern Gulf of Mexico

(29°00.175N, 87°31.131W) as part of the DEEPEND|RESTORE consortium (Cook et al., 2020).

Samples were collected at 450m depth (nighttime) during a cast of a CTD (Sea-Bird 911plus)

mounted with eleven 10-L Niskin Bottles. To maximize the detection of individuals,

multifrequency acoustic backscatter data was collected with a calibrated pole-mounted

echosounder system (Simrad EK60 and EK80). The transducers were mounted in an enclosed

housing and suspended 2.5 m below the water surface (see Boswell et al., 2020 for additional

details). Backscatter data were collected simultaneously at four frequencies (18, 38, 70, and 120

kHz). This was used to detect the deep scattering layer (DSL) in real-time and guide us in the

precise moment to trigger the Niskin Bottles. Because one of our objectives was to investigate

the effect of filtering different water volumes, all samples were collected simultaneously during

the same CTD cast to avoid any temporal or spatial variation in our results.

Once on board, seawater was transferred from Niskin bottles to 1-L sterile Nalgene bottles and

immediately transported to the laboratory and filtered. We used a MilliporeSigma™ Chemical

Duty Vacuum Pressure Pump (WP6111560) for filtration of all biological replicates and a

negative control. Each replicate of specific seawater volume (1L, 5L, or 10L) was filtered

through a sterile hydrophilic polyethersulfone (PES) filter (Pall Corporation; 47-mm diameter;

0.45-μm pore size). To prevent contamination, the filtering station was cleaned with 10% bleach,

and nitrile gloves were worn throughout the filtering protocol and were changed frequently.

Following sample filtration, the filters were placed in falcon tubes with silica beads, which

function as a desiccator, drying out the filters and reducing DNA degradation (Bakker et al.,

2017). Samples were stored at -80°C in a freezer onboard. Upon arrival, samples were

transported in dry ice to Florida International University and stored in -80ºC freezer until sample

processing.

eDNA: Sample Processing, DNA extraction, PCR, sequencing

Genomic DNA from samples was extracted using the DNeasy Blood & Tissue Kit (250) (Cat.

No. / ID: 69506) according to the manufacturer's protocol. Whole filters (47 mm) were used for

genomic DNA extraction. Genomic DNA was eluted into 200µl and frozen at -20°C.

We decided to perform this investigation focusing on the fish community based on their

abundance and ecological relevance in the deep-pelagic environment (Irigoien et al., 2014;

Iglesias et al., 2023), and because of the availability of primers capable of giving species

resolution in deep-sea fish communities (McClenaghan et al., 2020; Canals et al., 2021; Polanco

F. et al., 2021). Portions of hyper-variable regions of the mitochondrial 12S ribosomal RNA

(rRNA) gene were PCR amplified from each genomic DNA sample using the MiFishUF and

MiFishUR primers with spacer regions (Miya et al., 2015). Both forward and reverse primers

also contained a 5' adaptor sequence to allow for subsequent indexing and Illumina sequencing.

Each 25 µL PCR reaction was mixed according to the Promega PCR Master Mix specifications

(Promega catalog # M5133, Madison, WI) which included 12.5ul Master Mix, 0.5 µM of each

primer, 1.0 µl of gDNA, and 10.5 µl DNase/RNase-free H2O. DNA was PCR amplified using

the following conditions: initial denaturation at 95ºC for 3 minutes, followed by 45 cycles of 20

seconds at 98ºC, 30 seconds at 60ºC, and 30 seconds at 72ºC, and a final elongation at 72ºC for

10 minutes. PCR amplification was performed in replicates of ten (PCR replicates) per biological

replicate. We had problems amplifying some 1L samples, resulting in few PCR replicates for two

of the biological replicates. We decided to reallocate the resources by increasing one PCR

replicate to the other biological replicates. PCR replicates were not pooled and were kept

separate. Each reaction was visually inspected using a 2% agarose gel with 5μl of each sample as

input to determine amplicon size and PCR efficiency. Successful PCRs were cleaned by

incubating amplicons with Exo1/SAP for 30 minutes at 37°C following by inactivation at 95°C

for 5 minutes and stored at -20°C.

A second round of PCR was performed to complete the sequencing library construct, appending

with the final Illumina sequencing adapters and integrating a sample-specific,12-nucleotide

index sequence. The indexing PCR included Promega Master mix, 0.5 μM of each primer and 2

μl of template DNA (cleaned amplicon from the first PCR reaction) and consisted of an initial

denaturation of 95°C for 3 minutes followed by 8 cycles of 95°C for 30 sec, 55°C for 30

seconds and 72°C for 30 seconds. Final indexed amplicons from each sample were cleaned and

normalized using SequalPrep Normalization Plates (Life Technologies, Carlsbad, CA). 25μl of

PCR amplicon is purified and normalize using the Life Technologies SequalPrep Normalization

kit (cat#A10510-01) according to the manufacturer's protocol. Samples are then pooled together

by adding 5μl of each normalized sample to the pool. Library preparation was performed by

Jonah Ventures (Boulder, CO).

Sample library pools were sent for sequencing on an Illumina MiSeq (San Diego, CA) at the

Texas A&M Agrilife Genomics and Bioinformatics Sequencing Core facility using the v2 500-

cycle kit (cat# MS-102-2003). Necessary quality control measures were performed at the

sequencing center prior to sequencing.

eDNA: Bioinformatics

Raw sequence data were demultiplexed using pheniqs v2.1.0 (Galanti et al., 2021), enforcing

strict matching of sample barcode indices (i.e, no errors). Cutadapt v3.4 (Martin, 2011) was then

used remove gene primers from the forward and reverse reads, discarding any read pairs where

one or both primers (including a 6 bp, fully degenerate prefix) were not found at the expected

location (5') with an error rate < 0.15. Read pairs were then merged using vsearch v2.15.2

(Torbjørn et al., 2016), discarding resulting sequences with a length of < 130 bp, > 210 bp, or

with a maximum expected error rate > 0.5 bp (Edgar & Flyvbjerg, 2015). For each sample, reads

were then clustered using the unoise3 denoising algorithm (Edgar, 2016) as implemented in

vsearch, using an alpha value of 5 and discarding unique raw sequences observed less than 8

times. Counts of the resulting exact sequence variants (ESVs) were then compiled and putative

chimeras were removed using the uchime3 algorithm, as implemented in vsearch. For each final

ESV, a consensus taxonomy was assigned using a custom best-hits algorithm and a reference

database consisting of publicly available sequences (GenBank, Benson et al., 2005) as well as

Jonah Ventures voucher sequences records. Reference database searching used an exhaustive

semi-global pairwise alignment with vsearch, and match quality was quantified using a custom,

query-centric approach, where the % match ignores terminal gaps in the target sequence, but not

the query sequence. The consensus taxonomy was then generated using either all 100% matching

reference sequences or all reference sequences within 1% of the top match, accepting the

reference taxonomy for any taxonomic level with > 90% agreement across the top hits. Non-fish

species were removed from the final dataset. For downstream analyses, we kept only ESVs that

could be confidently assigned to a species (>98% similarity, Polanco F. et al.,2021). We kept

species that were identified at the genus level if they had a reference deposited in public

databases and an associated voucher number. The number of reads was coded as

presence/absence. The final datasets consisted of 1) presence/absence of each species per

biological replicate per treatment; 2) presence/absence of each species per PCR replicate per

biological replicate per treatment; 3) presence/absence of each ESV per biological replicate per

treatment; 4) presence/absence of each ESV per PCR replicate per biological replicate per

treatment.

Biodiversity analyses

We compared the number of species (richness) and ESVs detected in each treatment (1L, 5L,

10L) by performing a 1-way analysis of variance (ANOVA) followed by a posthoc Tukey test.

For comparisons of species and ESVs composition across treatments, we performed a

Permutation Multivariate Analysis of Variance (PERMANOVA; Anderson, 2001). First, we

built a dissimilarity matrix based on the Bray-Curtis index using the vegdist function in the

vegan package (Oksanen et al., 2022). Because the package's default is to use quantitative data

for all indices, we set binary = TRUE to account for the presence/absence structure of our

dataset. The resulting matrix was used as the response variable and volume as the explanatory

variable in the PERMANOVA model using the adonis2 function. Significance was accessed

using 9999 permutations. To investigate the effects of each treatment, we performed a pairwise

permutational test for multiple comparisons set to 9999 permutations using the pairwise.adonis2

function in the pairwiseAdonis (Arbizu, 2017). The relationships among each biological replicate

per treatment were visually represented by a Non-metric Multidimensional Scaling (nMDS) plot.

This analysis was done using the metaMDS function implemented in the vegan package.

Because convergence was not reached under default parameters, we used the argument trymax =

1000 to increase the number of random starts to find a stable solution. A permutation-based

similarity percentage (PER-SIMPER) analysis was performed to identify the taxa that most

contributed to the significant differences among treatments on both species and ESV datasets

(Gilbert & Escarguel, 2019). We also visualized the species and ESVs detected in each treatment

by plotting a Venn Diagram using ggvenn package (Yan, 2023), and used the gplots package

(Warnes et al., 2024) to find the exclusive and shared species/ESVs among treatments.

We investigated the efficiency of the number of biological replicates and PCR replicates in

detecting biodiversity by constructing rarefaction and extrapolation (R/E) plots using the

package iNterpolation/EXTrapolation (iNEXT, Hsieth et al., 2016). We followed a non-

asymptotic approach to plot a coverage-based R/E sampling completeness curve based on

sampling-unit-based incidence data. Depending on the analysis, the sampling unit was

considered the number of biological replicates (n = 5) per treatment (1L, 5L, 10L) or the number

of PCR replicates (n = 10) per biological replicate (n = 5) per treatment (1L, 5L, 10L). R/E plots

were constructed for both species and ESV datasets.

All analyses were run in R software (version 4.1.3).

RESULTS

We retained 2,061,989 reads that were assigned to the Class Actinopteri. After all filtering steps,

we retained 1,487,217 reads in total that could be assigned to a species (>98% similarity). The

number of reads per biological treatment was 1L = 48,793.4 ± 49,311.5; 5L = 100,506.2 ±

27,211.7; 10L = 148,143.8 ± 51,240.5. The number of reads per replicate per biological

treatment was 1L = 4,900.3 ± 4,847.5; 5L = 9,136.9 ± 2,473.8; 10L = 13,467.6 ± 4,658.2. One

biological replicate of the 1L treatment detected zero species. Negative controls had negligible

amounts of contamination, with an average of 408.4 ± 1890.8 across PCR replicates. Only 6

species were found in the negative control, usually represented by one ESV in just one of the

PCR replicates. We detected 98 taxa that could be confidently assigned to a species, representing

a total of 661 ESVs (Supplementary Table 1).

The richness detected by the 1L treatment is lower than the number detected by the treatment

filtering 10L (Tukey post-hoc, $p = 0.004$), but the number of species detected in 5L and 10L was

not statistically different ($p = 0.109$; Table 1; Figure 1A). ESVs detected by 1L treatments are

lower than the number of ESVs detected by the 10L treatment ($p = 0.004$); ESVs detected by 5L

treatments are marginally lower than the 10L ($p = 0.048$; Table 1; Figure 1B). In both cases, 1L

and 5L are not different (species, $p = 0.206$; ESVs, $p = 0.399$).

Table 1. Analysis of Variance (ANOVA) results comparing number of species (richness) found
in each treatment. Permutation Multivariate Analysis of Variance (PERMANOVA) results
comparing community structure among biological replicates per treatment (1L, 5L, 10L). DF:
degrees of freedom; SS: sum of squares; F: F-value; Pseudo-F: pseudo-F-value; P: p-value.

			ANOVA - Richness					
		Species					**ESVs**	
Source of Variation	*DF*	*SS*	*F*	*P*	*DF*	*SS*	*F*	*P*
Volume	2	1493	8.112	0.005	2	1493	8.112	0.005
Residuals	12	1104			12	1104		
			PERMANOVA - Community Structure					
		Species					**ESVs**	
Source of Variation	*DF*	*SS*	*Pseudo-F*	*P*	*DF*	*SS*	*Pseudo-F*	*P*
Volume	2	0.71959	1.8027	0.004	2	0.9926	1.3744	0.002
Residuals	12	2.395			12	5.3258		

Figure 1. Box-plot representing the number of species (A) and ESVs (B) detected in each

treatment (1L, 5L, 10L). Letters represent the result of the *post-hoc* Tukey test (similar letters

are not statistically different). Colors represent different treatments.

Similarly, the PERMANOVA analysis indicated that volume affects community composition

(Table 1). Considering the species dataset, the community detected by 1L marginally differs

from the 5L (pairwise permutational test for multiple comparisons, p = 0.053) and differs from

the 10L communities (p = 0.017), but 5L and 10L are not different (p = 0.113; Figure 2A).

Considering the ESVs dataset, each treatment detected a different community composition (1L

vs. 5L, p = 0.04; 1L vs. 10L, p = 0.017; 5L vs. 10L, p = 0.031, Figure 2B). When looking at the

species contributing to the significant differences found in each comparison, the PER-SIMPER

analysis indicates the top 15 species contributed 46.8% to the dissimilarity between 1L vs. 5L,

and 34.2% between 1L vs. 10L (Table 2). Although each species contributes little to the

dissimilarity found, PER-SIMPER indicates that 8 out of the 15 species had a role in both cases:

Barbourisia rufa, Benthosema suborbitale, Caranx crysos, Cyclothone pallida, Diaphus lucidus,

Gonostoma elongatum, Lampanyctus alatus, and *Rondeletia loricata.* Twenty-two species

(22.4%) were found in all treatments (Figure 3, Supplementary Table 1). Exclusive species

represent species detected in only one biological replicate in 100% (1L = 11, and 5L = 7) or 68%

(10L = 22) of the cases, with 10L being the treatment that had the highest number of exclusive

species (32)(Figure 3A, Supplementary Table 1). Regarding ESVs, the top 15 ESVs contribute to

33.6% (1L vs. 5L), 30.7% (1L vs. 10L), 25.6% (5L vs. 10L) (Supplementary Table 1). Most of

the ESVs contributing to the differences between 1L vs. 5L and 10L indicated by PER-SIMPER

are ESVs from the species detected in the species-level analysis (Table 2, Supplementary Table

1). In the case of the 5L vs. 10L comparison, all ESVs represent species shared between the two

treatments (Supplementary Table 1). Only twenty-seven ESVs were shared among all treatments

(4.1%), and the 10L treatment had the highest amount of exclusive ESVs (351, 53.3%) (Figure

3B)

Table 2. Permutation-based similarity percentage (PER-SIMPER) results showing the top 15

species contributing to the differences found between statistically different communities (see

PERMANOVA results).

1L vs. 5L		1 vs. 10L	
Species	%	Species	%
Cyclothone pallida	4.3	*Lagocephalus laevigatus*	3.2
Gonostoma elongatum	4.3	*Canthigaster rostrata*	2.8
Diaphus lucidus	3.6	*Cyclothone pallida*	2.8
Caranx crysos	3.3	*Diaphus lucidus*	2.8
Cyclothone pseudopallida	3.3	*Gonostoma elongatum*	2.8
Barbourisia rufa	3.3	*Benthosema suborbitale*	2.3
Lampanyctus alatus	3.1	*Notolychnus valdiviae*	2.3
Melamphaes simus	3.1	*Caranx crysos*	2.1
Diaphus perspicillatus	3.0	*Lampanyctus alatus*	2.1

Cyclothone obscura	2.8	*Barbourisia rufa*	2.0
Rondeletia loricata	2.8	*Cyclothone acclinidens*	1.9
Benthosema suborbitale	2.5	*Rondeletia loricata*	1.9
Lepidophanes guentheri	2.5	*Nannobrachium lineatum*	1.9
Seriola rivoliana	2.4	*Caranx hippos*	1.8
Diaphus rafinesquii	2.4	*Lutjanus campechanus*	1.8
Total	46.8		34.3

346 Figure 2. Non-metric multi (A) and

347 ESVs (B) composition found in each treatment (1L, 5L, 10L). The 1L replicate that detected zero

348 species/ESVs was excluded to facilitate visualization of the spatial distribution of samples.

349 Colors represent different treatments.

Figure 3. Venn diagram indicating the number of exclusive species in each treatment (1L, 5L,

10L), and shared species between two treatments or shared among all treatments (A); Venn

diagram indicating the number of exclusive ESVs in each treatment (1L, 5L, 10L), and shared

ESVs between two treatments or shared among all treatments.

When analyzing the impact of the number of biological replicates in detecting the total

biodiversity across all of our samples, we show that 5 biological replicates could detect up to

80% of all 98 species in both 5L and 10L treatments. The extrapolation analysis indicates that 10

replicates would recover 100% of the species detected in the water collected (note: not total

biodiversity from the region, but all 98 species). In contrast, 5 biological replicates of 1L

treatment detected only 50% of the total biodiversity, and the extrapolation analysis indicates

that more than 10 biological replicates would be necessary to detect 100% (Figure 4A). A

different scenario is found when analyzing ESVs, showing that 5 biological replicates detected

~40% of the 611 ESVs detected. Extrapolation analyses indicated that more than 10 biological

replicates would be necessary to detect all ESVs (Figure 4B).

Figure 4. Rarefaction and extrapolation plot depicting the increase of species (A) or ESVs (B)

detected as more biological replicates were added per treatment (1L, 5L, 10L).

When analyzing the impact of the number of PCR replicates per biological replicate, we find that

10 PCR replicates show variable biodiversity coverage for 5L, detecting between 60-85% of total

number of species depending on the biological replicate (Figure 5B). Similarly, 10L biological

replicates show a similar pattern, but detecting around 80-90% of the total number of species

(Figure 5C). In contrast, results concerning PCR replicates of 1L treatments are extremely

variable (Figure 5A). Similar to the previous results, PCR replicates R/E plots analyzing ESVs

indicate large variation among 1L biological replicates (Figure 5D), and moderate/low variation

among 5L and 10L biological replicates. In these cases, 10 PCR replicates detected between 20-

60% of the ESVs sampled in this study (Figure 5E-F). One of the biological replicates in the 1L

treatment did not have enough DNA for running all PCR replicates.

Figure 5. Rarefaction and extrapolation plot depicting the increase of species/ESVs detected as

more PCR replicates were added biological replicate per treatment (A: species-1L; B: species-

5L; C: species10L; D: ESVs-1L; E: ESVs-5L; F: ESVs-10L).

DISCUSSION

Experimental design is a crucial part of any ecological study, but there are no golden rules when

implementing eDNA approaches (Deiner et al., 2017; Dickie et al., 2018; Ruppert et al., 2019;

Cote et al., 2023). Sampling design considerations must be carefully considered before going to

sea, especially for deep-pelagic eDNA investigations. As far as we know, previous investigations

have not tested how different sampling schemes (water volume and replication) using common

equipment (e.g., Niskin Bottles coupled with CTD) can impact deep-sea biodiversity

assessments. Our study shows that common practices adopted in coastal waters (filtering 1L of

water) are not ideal for characterizing deep-pelagic fish biodiversity. Filtering 5L or 10L did not

result in statistically different species compositions but detected different ESV compositions. In

terms of replication, 5 or more biological replicates should be the standard for similar

experimental designs using Niskin bottles, and the number of PCR replicates should be higher

when the volume of water filtered is lower. Overall, results are similar when investigating

species and ESVs. We recommend 5-10 PCR replicates per biological replicate to maximize the

detection of taxa and different ESVs. Our results have implications for future eDNA studies

targeting deep-pelagic communities and for future eDNA studies comparison.

Water volume

In aquatic eDNA studies, a standing question is: how much water should be filtered? Data from

studies conducted in shallow systems indicate that filtered water volume is known to impact taxa

detection (Muha et al., 2019; Kumar et al., 2022). In these cases, filtered water volume usually

varies from 0.5 mL to 2L (Rees et al., 2014; Bessey et al., 2020; Kumar et al., 2022). Directly

transposing shallow water standards to mid- and deep-pelagic ecosystems eDNA investigations

might not be ideal considering the differences between environments. The open ocean is the

largest environment on Earth, covering 70% of our planet, and the deep sea represents 80-90% of

our oceans, making it the largest habitat on Earth (Ramirez-Llodra et al., 2010). Therefore,

species abundance and distribution are expected to differ substantially across spatial scales when

comparing shallow- and deep-pelagic ecosystems. Few studies have investigated the impacts of

different filtering protocols, resulting in similar water volumes being filtered between shallow

water and deep-pelagic eDNA studies (McClenaghan et al., 2020; Diao et al., 2023). Our results

indicate that 1L is far from ideal for characterizing deep-pelagic fish communities. Although

some comparisons are not statistically significant, NMDS plots and rarefactions curves suggest

large variation among 1L biological replicates in comparison to 5L and 10L treatments. This can

be explained by the low eDNA abundance per volume of water in large habitats, leading to

stochastic detection of eDNA molecules in 1L samples. Interestingly, species composition

differences between 1L vs. 5L and 10L can be explained by the omission of rare species, but also

species known to be very common in deep-pelagic habitats within the Gulf of Mexico including

species from Gonostomatidae (*Cyclothone pallida*, *Gonostoma elongatum*) and Myctophidae

(*Lampanyctus alatus*) (Ross et al., 2010). Therefore, we conclude that 1L does not represent the

eDNA available nor describe the local fish diversity, indicating that more water must be filtered

when investigating deep-pelagic communities.

Surprisingly, 5L and 10L detected similar numbers of fish species per biological replicate and

showed the same species composition, even though the mean number of reads in the 10L

treatment was almost 50% larger than in the 5L treatment. A comparison of the number of

operational taxonomic units (OTUs) found in different filtered water volumes (~5-60L) also

indicated no changes in total OTUs, although this varied according to location (Yoshida et al.,

2023). It should be noted that species detection based on eDNA is constrained by the persistence

of DNA molecules in the water, correct amplification via PCR, and reference libraries that

contain the necessary information to identify species correctly. Our results are relevant when

considering the effort-time-cost balance related to eDNA fieldwork that does not involve an

autonomous water sampler and specialized filtering apparatus. Collecting and filtering water can

be laborious and time-consuming and involve multiple researchers when using commonly

available gear, like Niskin bottles and vacuum pumps. Here, we suggest that, when limitations to

water collection exist, efforts toward increasing filtered water volume above 10L may not impact

the recovery of deep-pelagic species. A similar conclusion was shown when comparing 250mL

and 1.5L collected using a CTD (McClenaghan et al., 2020), but also when comparing CTD

samples (~2L) and a recently developed large-volume autonomous eDNA sampler with in-situ

filtration mounted on the midwater robot Mesobot (~40-60L) (Govindarajan et al., 2022). In the

latter study, the community composition detected was not different when comparing the CTD

samples and the Mesobot, but the number of sequence reads and ESV was higher when using the

autonomous samples (Govindarajan et al., 2022). This is similar to our findings, as the 10L

treatment detected more ESVs and a different ESVs community composition than the 1L and 5L

treatments. A striking conclusion is that 10L did not detect more species, but detected more

intraspecific variation (i.e., more ESVs per species). We believe both 5L and 10L treatments

reflected the natural proportion of fish eDNA that could be detected because filtering more water

did not detect more species. Therefore, choosing how many liters of water will be filtered should

also consider whether the detection of more ESVs is part of the goal of the study. For instance,

recent analytical developments have suggested that the number of intraspecific ESVs can be used

to estimate species abundance (Yoshitake et al., 2019). The understanding of the impacts of

filtered volumes on the characterization of deep-pelagic communities is still under development,

and more studies will illuminate whether there is an ideal volume that can be filtered. We should

also note that Govindarajan et al. (2022) targeted the 18S rRNA region and many metazoan

groups at lower taxonomic resolution, while we targeted 12S rRNA and fish species at a higher

taxonomic resolution. The taxonomic resolution and efficiency vary according to the primer used

and the region targeted (Ruppert et al., 2019; Zhang et al., 2020; Polanco et al., 2021; Kumar et

al., 2022), being an important step during experimental design. We used a taxon-specific primer

(MiFish) to better understand the nuances of sampling schemes at the highest taxonomic

resolution (i.e., species). It is not possible to say if the results of Govindarajan et al. (2022)

would be different whether they had targeted a different region (i.e., used a different primer) or

investigated higher taxonomic resolution. It is important to notice that we are discussing these

data under a community composition framework, with the goal of comparing samples across

time and space. When the main goal of an investigation is to detect one or a few specific and/or

rare taxa, higher volumes should be aimed as DNA detection likely increases (Sepulveda et al.,

2019; Bessey et al., 2020; Govindarajan et al., 2022). Our study also reports evidence for this

latter statement: the 10L detected an absolute number of species higher than 1L and 5L

treatments (but the number of species per biological replicate is not different from the 5L

treatment). These rare species have a minor role when discussing communities'

similarities/differences. We should notice that this is related to the eDNA signal detected, but

some species might not be rare in terms of abundance at the organismal level. As mentioned

before, this could be explained by species shedding less DNA or primer incompatibilities.

Biological and PCR replicates

Another crucial factor to consider when performing eDNA investigations is the number of

biological and PCR replicates. Our work is one of the first to address the variation in these

factors when characterizing deep-sea systems. Variation among 1L biological and PCR replicates

was higher than other treatments for both species and ESVs datasets. Five biological and 10 PCR

replicates of the 1L treatment were insufficient to assess the local fish biodiversity (i.e., the

species and ESVs detected in this study). In fact, 1L treatments detected a different species/ESV

composition and fewer species/ESV than the other treatments, and even our extrapolation

analyses could not reach levels compared to the other treatments. This means that not only the

volume of water but also the number of replicates is important when considering eDNA applied

to mid- and deep-water pelagic systems. Our results indicate that, under certain circumstances, it

could be better to favor less volume and more replicates if one is limited on the amount of water

that they can sample using their available gear (e.g., Niskin bottles coupled with CTD can only

sample a specific total volume of water). We provide empirical data to the literature reinforcing

previous suggestions for other systems recommending more replicates to improve the detection

probabilities and to estimate fish biodiversity reliably (Bessey et al., 2020; Stauffer et al., 2021;

Cote et al., 2023; Yoshida et al., 2023). Considering species detection, five biological replicates

of the 5L and 10L treatments did not yield different results, but biodiversity sampling was

maximized by adding more biological replicates in our study. Therefore, an optimal deep-sea

eDNA experimental design could aim for a minimum of five biological replicates of a minimum

of 5L, which is doable considering most traditional water samplers (e.g., McClenaghan et al.,

2020; Feng et al., 2022; Govindarajan et al., 2023). However, future studies should address this

topic because the trade-offs between water filtered and the number of biological replicates using

autonomous samplers have yet to explored.

Similarly, the investigation of the ideal number of PCR replicates is still scarce in the literature.

Amplification is a stochastic process during PCR reactions, and more reactions will probably

result in the detection of more taxa by chance. The exact number needed will vary depending on

the probability of DNA detection, which varies according to species abundance, rate of shedding

DNA, environmental conditions, and other factors (Ruppert et al., 2019). A simulation study

showed that four PCR replicates might result in high levels of false negatives, and six to eight

PCR replicates should be performed depending on the probability of DNA detection (Ficetola et

al., 2015). As discussed before, it is feasible to assume that DNA is sparsely distributed in the

deep sea, considering this is the largest habitat on Earth and individuals' density might not be

high. Still, PCR replicates are often low in studies investigating deep waters (2-4 PCR replicates;

e.g., McClenaghan et al., 2020; Canals et al., 2021; Govindarajan et al., 2022; Cote et al. 2023;

Yoshida et al., 2023). Our results show that all samples benefited from more PCR replicates

when considering species and ESVs detection. Besides biological replicates, the ideal sampling

scheme should balance the volume of water filtered with the number of PCR replicates. Our

results indicate that at least 5-10 PCR replicates should be the goal, considering the water

volume filtered. That is, if limited by the water volume, the best strategy is to increase the

number of PCR replicates; if not limited by water volume, fewer PCR replicates can be

performed. Ideally, future studies should first find the optimal design to maximize water volume

and PCR replicates accordingly.

Limitations

It is important to mention the limitations of using common equipment in deep-sea eDNA studies.

The maximum water volume is limited by the capacity and number of Niskin Bottles attached to

the CTD and the number of deployments within a scientific mission. Our study leverages

common gear found on research vessels and could not test if filtering more than 10L would

change the biodiversity characterization, especially at the species level, as ESV diversity is

favored by filtering more water. Previous studies did not find different species composition when

comparing filtering 2.5L using Niskin Bottles and 40-60L using an autonomous vehicle in the

Gulf of Mexico and targeting the 18S region (Govindarajan et al., 2022). However, community

characterization changed in some locations near Japan after filtering >25L of water per

biological replicate and targeting the 12S region (Yoshida et al., 2023). A potential future

standard in this type of study could be first to investigate the ideal water volume needed to

characterize the community at the specific ocean basin using a specific primer. Considering our

results and Govindarajan et al. (2022), 5L-10L might be sufficient for the northern Gulf of

Mexico when targeting the 18S and 12S rRNA markers. We should note that some taxa might

shed less DNA than others (Andruszkiewicz Allan et al., 2021), which should also be considered

when planning an eDNA investigation. For instance, we could not detect substantial decapod

eDNA using the MiDecap primer set (Komai et al., 2019) using the volume of water and

replicates used in this study (data and results not shown). This is surprising, considering the

diversity and abundance of pelagic decapods in the Gulf of Mexico collected using nets and with

available reference sequences (Burdett et al., 2017; Varela et al., 2021). We predict that primer

compatibility and shedding may have been involved in the lack of recovery, but further studies

should explore the impact of water volume and reference libraries on the recovery of deep-sea

crustaceans and other taxa. Simulation analysis indicates that increasing the water volume

filtered could lead to similar species being detected using eDNA and traditional methods,

reinforcing this idea (Cote et al., 2023). Additionally, if detecting specific species (instead of

community composition) is the goal, one might consider maximizing the water volume collected,

as mentioned before (Sepulveda et al., 2019; Bessey et al., 2020; Govindarajan et al., 2022).

Still, the exact volume might be taxa-specific and should also be a first step in this type of

investigation. Another factor to be considered is that eDNA might degrade while samples are

waiting to be or while they are being processed (Holman et al., 2022). In our case, all replicates

were filtered in a randomized order to avoid this pitfall. However, the decay rate of deep-sea

eDNA samples is still unknown for many taxa, and future studies should address this knowledge

gap (but see McCartin et al., 2022 for insights).

Besides the technical limitations mentioned, the primary pitfall of all deep-water eDNA

investigations is the catastrophic lack of reliable reference libraries. eDNA approaches are 100%

reliant on the completeness and quality of available reference databases for properly identifying

taxa (Bylemans et al., 2018; Collins et al., 2019; Schenekar et al., 2020). In our study, ~50% of

our reads could not be confidently assigned to a fish species. Another situation that highlights the

importance of reliable reference libraries is that we also detected a few Indo-Pacific species in

our final dataset. We cannot confirm if they represent real occurrences or problems in reference

sequences available in public databases. This study, among others, highlights the urgent need for

deep-pelagic reference libraries that are linked to curated, voucher specimens deposited in

museums that can be verified by taxonomic experts. Current efforts, including our own, are

expanding deep-sea reference libraries (e.g., Bucklin et al., 2010; Weigand et al., 2019; Varela et

al., 2021; Govindarajan et al., 2022), which need to be standardized, shared and extended to as

many taxa and locations as possible.

Conclusions

The application of eDNA to mid- and deep-water ecosystems is gaining popularity, however few

studies have experimentally tested the effects of different sampling protocols on the recovery of

species biodiversity (Canals et al., 2021; Easson et al., 2020; Govindarajan et al., 2021; Laroche

et al., 2020; Merten et al., 2021). Our study provides empirical results on how experimental

design can impact the characterization of deep-pelagic communities. Specifically, we indicate

that a minimum of 5L-10L of water should be filtered when targeting the deep-scattering layer

and paired with a minimum of 5 or more biological replicates for deep-sea eDNA studies when

investigating species composition. Additionally, we show the importance of more PCR replicates

when analyzing deep-sea samples. Technological advances in eDNA samplers are crucial and are

expanding our understanding of deep-sea ecosystems. But, until high-powered technology

becomes assessable for all, using Niskin bottles coupled with a CTD can be a way to

democratize the field and increase our local and global-scale observations. Most importantly,

because deep-ocean explorations are constrained by logistics and budget, we believe our work

can help in providing some guidance on potential standards and how eDNA data generated by

different research groups across the globe could be comparable when the focus is mid- or deep-

water community characterization using common gear.

REFERENCES

Billings, A., Kaiser, C., Young, C.M., Hiebert, L.S., Cole, E., Wagner, J.K.S., Van Dover, C.

L., 2017. SyPRID sampler: a large-volume, high-resolution, autonomous, deep-ocean

precision plankton sampling system. Deep Sea Research Part II: topical Studies in

Oceanography. Advances in deep-sea biology: biodiversity, ecosystem functioning

and conservation 137, 297–306. https://doi.org/10.1016/j.dsr2.2016.05.007.

Yamahara, K.M., Preston, C.M., Birch, J., Walz, K., Marin, R., Jensen, S., Pargett, D.,

Roman, B., Ussler, W., Zhang, Y., Ryan, J., Hobson, B., Kieft, B., Raanan, B.,

Goodwin, K.D., Chavez, F.P., Scholin, C., 2019. In situ autonomous acquisition and

preservation of marine environmental DNA using an autonomous underwater

vehicle. Front. Mar. Sci. 6, 373. https://doi.org/10.3389/fmars.2019.00373.

Yoerger, D.R., Govindarajan, A.F., Howland, J.C., Llopiz, J.K., Wiebe, P.H., Curran, M.,

Fujii, J., Gomez-Ibanez, D., Katija, K., Robison, B.H., Hobson, B.W., Risi, M., Rock, S.

M., 2021. A hybrid underwater robot for multidisciplinary investigation of the ocean

twilight zone. Sci. Robot. 6 (55), eabe1901 https://doi.org/10.1126/scirobotics.

abe1901.

Cook, A. B., Bernard, A. M., Boswell, K. M., Bracken-Grissom, H., D'Elia, M., DeRada, S., ...

& Sutton, T. T. (2020). A Multidisciplinary approach to investigate deep-pelagic ecosystem

dynamics in the Gulf of Mexico following Deepwater Horizon. Frontiers in Marine Science, 7,

548880.

Galanti, Shasha and Gunsalus (2021). Pheniqs 2.0: accurate, high-performance Bayesian

decoding and confidence estimation for combinatorial barcode indexing. BMC Bioinformatics

22.

Martin (2011), Cutadapt removes adapter sequences from high-throughput sequencing reads.

EMBnet. journal 17.

Torbjørn et al. VSEARCH: a versatile open source tool for metagenomics. PeerJ 4 e2584 (2016).

Edgar and Flyvbjerg (2015), Error filtering, pair assembly and error correction for next-

generation sequencing reads. Bioinformatics 31.

Edgar (2016), UNOISE2: improved error-correction for Illumina 16S and ITS amplicon

sequencing. doi: https://doi.org/10.1101/081257.

Benson et al. (2005), GenBank. Nucleic Acids Res 33.

McClenaghan, B., Fahner, N., Cote, D., Chawarski, J., McCarthy, A., Rajabi, H., Singer, G. and

Hajibabaei, M., 2020. Harnessing the power of eDNA metabarcoding for the detection of deep-

sea fishes. PLoS One, 15(11), p.e0236540.

Canals, O., Mendibil, I., Santos, M., Irigoien, X., & Rodríguez-Ezpeleta, N. (2021). Vertical

stratification of environmental DNA in the open ocean captures ecological patterns and behavior

of deep-sea fishes. Limnology and Oceanography Letters, 6(6), 339-347.

Polanco F, A., Richards, E., Flück, B., Valentini, A., Altermatt, F., Brosse, S., Walser, J.C., Eme,

D., Marques, V., Manel, S., & Albouy, C. (2021). Comparing the performance of 12S

mitochondrial primers for fish environmental DNA across ecosystems. Environmental DNA,

3(6), 1113-1127.

Kumar, G., Farrell, E., Reaume, A. M., Eble, J. A., & Gaither, M. R. (2022). One size does not

fit all: Tuning eDNA protocols for high-and low-turbidity water sampling. Environmental DNA,

4(1), 167-180.

Ruppert, K. M., Kline, R. J., & Rahman, M. S. (2019). Past, present, and future perspectives of

environmental DNA (eDNA) metabarcoding: A systematic review in methods, monitoring, and

applications of global eDNA. Global Ecology and Conservation, 17, e00547.

Govindarajan, A. F., McCartin, L., Adams, A., Allan, E., Belani, A., Francolini, R., ... & Herrera,

S. (2022). Improved biodiversity detection using a large-volume environmental DNA sampler

with in situ filtration and implications for marine eDNA sampling strategies. Deep Sea Research

Part I: Oceanographic Research Papers, 189, 103871.

McClenaghan, B., Fahner, N., Cote, D., Chawarski, J., McCarthy, A., Rajabi, H., ... &

Hajibabaei, M. (2020). Harnessing the power of eDNA metabarcoding for the detection of deep-

sea fishes. PLoS One, 15(11), e0236540.

Diao, C., Wang, M., Zhong, Z., Li, Y., Xian, W., & Zhang, H. (2023). Biodiversity exploration

of Formosa Ridge cold seep in the South China Sea using an eDNA metabarcoding approach.

Marine Environmental Research, 190, 106109.

Yoshida, T., Kawato, M., Fujiwara, Y., Nagano, Y., Tsuchida, S., & Yabuki, A. (2023).

Optimization of environmental DNA analysis using pumped deep-sea water for the monitoring

of fish biodiversity. Frontiers in Marine Science, 9, 965800.

Bessey, C., Jarman, S. N., Berry, O., Olsen, Y. S., Bunce, M., Simpson, T., ... & Keesing, J.

(2020). Maximizing fish detection with eDNA metabarcoding. Environmental DNA, 2(4), 493-

504.

Collins, R. A., Bakker, J., Wangensteen, O. S., Soto, A. Z., Corrigan, L., Sims, D. W., ... &

Mariani, S. (2019). Non-specific amplification compromises environmental DNA metabarcoding

with COI. Methods in Ecology and Evolution, 10(11), 1985-2001.

Schenekar, T., Schletterer, M., Lecaudey, L. A., & Weiss, S. J. (2020). Reference databases,

primer choice, and assay sensitivity for environmental metabarcoding: Lessons learnt from a re-

evaluation of an eDNA fish assessment in the Volga headwaters. River Research and

Applications, 36(7), 1004-1013.

Stauffer, S., Jucker, M., Keggin, T., Marques, V., Andrello, M., Bessudo, S., ... & Waldock, C.

(2021). How many replicates to accurately estimate fish biodiversity using environmental DNA

on coral reefs?. Ecology and Evolution, 11(21), 14630-14643.

Canals, O., Mendibil, I., Santos, M., Irigoien, X., & Rodríguez-Ezpeleta, N. (2021). Vertical

stratification of environmental DNA in the open ocean captures ecological patterns and behavior

of deep-sea fishes. Limnology and Oceanography Letters, 6(6), 339-347.

Varela, C., Golightly, C., Timm, L. E., Wilkins, B., Frank, T., Fenolio, D., ... & Bracken-

Grissom, H. D. (2021). DNA barcoding enhances large-scale biodiversity initiatives for deep-

pelagic crustaceans within the Gulf of Mexico and adjacent waters. The Journal of Crustacean

Biology, 41(1), ruab005.

Komai, T., Gotoh, R. O., Sado, T., & Miya, M. (2019). Development of a new set of PCR

primers for eDNA metabarcoding decapod crustaceans. Metabarcoding and Metagenomics, 3,

e33835.

McCartin, L. J., Vohsen, S. A., Ambrose, S. W., Layden, M., McFadden, C. S., Cordes, E. E., ...

& Herrera, S. (2022). Temperature controls eDNA persistence across physicochemical

conditions in seawater. Environmental Science & Technology, 56(12), 8629-8639.

Andruszkiewicz Allan, E., Zhang, W. G., Lavery, A., & Govindarajan, A. (2021). Environmental

DNA shedding and decay rates from diverse animal forms and thermal regimes. Environmental

DNA 3 (2): 492–514.

Holman, L. E., Chng, Y., & Rius, M. (2022). How does eDNA decay affect metabarcoding

experiments?. Environmental DNA, 4(1), 108-116.

Kumar, G., Reaume, A. M., Farrell, E., & Gaither, M. R. (2022). Comparing eDNA

metabarcoding primers for assessing fish communities in a biodiverse estuary. PloS one, 17(6),

e0266720.

Zhang, S., Zhao, J., & Yao, M. (2020). A comprehensive and comparative evaluation of primers

for metabarcoding eDNA from fish. Methods in Ecology and Evolution, 11(12), 1609-1625.

Muha, T. P., Robinson, C. V., Garcia de Leaniz, C., & Consuegra, S. (2019). An optimised

eDNA protocol for detecting fish in lentic and lotic freshwaters using a small water volume.

PLoS One, 14(7), e0219218.

Feng, Y., Sun, D., Shao, Q., Fang, C., & Wang, C. (2022). Mesozooplankton biodiversity,

vertical assemblages, and diel migration in the western tropical Pacific Ocean revealed by eDNA

metabarcoding and morphological methods. Frontiers in Marine Science, 9, 1004410.

Govindarajan, A. F., Llopiz, J. K., Caiger, P. E., Jech, J. M., Lavery, A. C., McMonagle, H., ... &

Zhang, W. G. (2023). Assessing mesopelagic fish diversity and diel vertical migration with

environmental DNA.

Ramirez-Llodra, E., Brandt, A., Danovaro, R., De Mol, B., Escobar, E., German, C. R., ... &

Vecchione, M. (2010). Deep, diverse and definitely different: unique attributes of the world's

largest ecosystem. Biogeosciences, 7(9), 2851-2899.

Rees, H. C., Maddison, B. C., Middleditch, D. J., Patmore, J. R., & Gough, K. C. (2014). The

detection of aquatic animal species using environmental DNA–a review of eDNA as a survey

tool in ecology. Journal of applied ecology, 51(5), 1450-1459.

Weigand, H., Beermann, A. J., Čiampor, F., Costa, F. O., Csabai, Z., Duarte, S., ... & Ekrem, T.

(2019). DNA barcode reference libraries for the monitoring of aquatic biota in Europe: Gap-

analysis and recommendations for future work. Science of the Total Environment, 678, 499-524.

Bucklin, A., Ortman, B. D., Jennings, R. M., Nigro, L. M., Sweetman, C. J., Copley, N. J., ... &

Wiebe, P. H. (2010). A "Rosetta Stone" for metazoan zooplankton: DNA barcode analysis of

species diversity of the Sargasso Sea (Northwest Atlantic Ocean). Deep Sea Research Part II:

Topical Studies in Oceanography, 57(24-26), 2234-2247.

Iglesias, I.S., Santora, J.A., Fiechter, J. & Field, J.C. (2023). Mesopelagic fishes are important

prey for a diversity of predators. Frontiers in Marine Sciences, 10, 10.3389/fmars.2023.1220088

Irigoien, X., Klevjer, T. A., Røstad, A., Martinez, U., Boyra, G., Acuña, J. L., ... & Kaartvedt, S.

(2014). Large mesopelagic fishes biomass and trophic efficiency in the open ocean. Nature

communications, 5(1), 3271.

Gibert, C., & Escarguel, G. (2019). PER-SIMPER—A new tool for inferring community

assembly processes from taxon occurrences. Global Ecology and Biogeography, 28(3), 374-385.

Hsieh, T. C., Ma, K. H., & Chao, A. (2016). iNEXT: an R package for rarefaction and

extrapolation of species diversity (H ill numbers). Methods in Ecology and Evolution, 7(12),

1451-1456.

Govindarajan, A. F., Francolini, R. D., Jech, J. M., Lavery, A. C., Llopiz, J. K., Wiebe, P. H., &

Zhang, W. (2021). Exploring the use of environmental DNA (eDNA) to detect animal taxa in the

mesopelagic zone. Frontiers in Ecology and Evolution, 9, 574877.

Visser, F., Merten, V. J., Bayer, T., Oudejans, M. G., De Jonge, D. S. W., Puebla, O., ... &

Hoving, H. J. T. (2021). Deep-sea predator niche segregation revealed by combined cetacean

biologging and eDNA analysis of cephalopod prey. Science Advances, 7(14), eabf5908.

Beng, K. C., & Corlett, R. T. (2020). Applications of environmental DNA (eDNA) in ecology

and conservation: opportunities, challenges and prospects. Biodiversity and Conservation, 29,

2089-2121.

Barnes, M. A., & Turner, C. R. (2016). The ecology of environmental DNA and implications for

conservation genetics. Conservation genetics, 17(1), 1-17.

Stat, M., Huggett, M. J., Bernasconi, R., DiBattista, J. D., Berry, T. E., Newman, S. J., ... &

Bunce, M. (2017). Ecosystem biomonitoring with eDNA: metabarcoding across the tree of life in

a tropical marine environment. Scientific reports, 7(1), 12240.

Thurber, A. R., Sweetman, A. K., Narayanaswamy, B. E., Jones, D. O., Ingels, J., & Hansman,

R. L. (2014). Ecosystem function and services provided by the deep sea. Biogeosciences, 11(14),

3941-3963.

Ross, S. W., Quattrini, A. M., Roa-Varón, A. Y., & McClain, J. P. (2010). Species composition

and distributions of mesopelagic fishes over the slope of the north-central Gulf of Mexico. Deep

Sea Research Part II: Topical Studies in Oceanography, 57(21-23), 1926-1956.

Yoshitake, K., Yoshinaga, T., Tanaka, C., Mizusawa, N., Reza, M. S., Tsujimoto, A., Kobayashi,

A. & Watabe, S. (2019). HaCeD-Seq: a novel method for reliable and easy estimation about the

fish population using haplotype count from eDNA. Marine Biotechnology, 21, 813-820.

Cote, D., McClenaghan, B., Desforges, J., Fahner, N. A., Hajibabaei, M., Chawarski, J., Roul, S.,

Singer, G., Aubry, C, & Geoffroy, M. (2023). Comparing eDNA metabarcoding and

conventional pelagic netting to inform biodiversity monitoring in deep ocean environments.

ICES Journal of Marine Science, 80(10), 2545-2562.

Burdett, E. A., Fine, C. D., Sutton, T. T., Cook, A. B., & Frank, T. M. (2017). Geographic and

depth distributions, ontogeny, and reproductive seasonality of decapod shrimps (Caridea:

Oplophoridae) from the northeastern Gulf of Mexico. Bulletin of Marine Science, 93(3), 743-

767.

806