Persistent legacy effects on soil microbiota facilitate plant adaptive responses to drought

Nichole A. Ginnan[1,2*], Valéria Custódio[3*], David Gopaulchan[3*], Natalie Ford[1,4], Isai Salas-
González[5], Dylan H. Jones[3], Darren M. Wells[3], Ângela Moreno[6], Gabriel Castrillo[3#], Maggie R.
Wagner[1,4#]

[1] Kansas Biological Survey and Center for Ecological Research, University of Kansas, Lawrence,
KS, USA 66045
[2] One Health Microbiome Center, Huck Institutes of the Life Sciences, Penn State University,
State College, PA, USA
[3] School of Biosciences, University of Nottingham, Sutton Bonington, UK
[4] Department of Ecology and Evolutionary Biology, University of Kansas, Lawrence, KS, USA
66045
[5] Centre for Genomics Sciences. Universidad Nacional Autónoma de México
[6] Ministério da Agricultura e Ambiente, Cabo Verde
14
15 [*]These authors contributed equally
16 [#]Corresponding authors (gabriel.castrillo@nottingham.ac.uk and maggie.r.wagner@ku.edu)

Abstract

Both chronic and acute drought alter the composition and physiology of the soil microbiota,

favoring the selection of microbes with functional traits that preserve fitness in these challenging

conditions[1]. This drought-adapted microbiota may influence water-use efficiency mechanisms in

host plants. Currently, it is largely unknown how this soil microbial drought legacy manifests at

the molecular and physiological levels and how it influences microbe-dependent plant responses

to drought in diverse natural soils. Here, to infer how drought legacy affects soil microbiota

composition and function, we collected soils spanning two distinct and geographically distant

precipitation gradients. By controlling for variation in numerous other soil properties including

porosity and elemental profiles, we identified a group of bacterial taxa that are highly sensitive to

water availability and critical for the formation of soil drought memory. Next, by imposing an acute

experimental drought we show that this microbial drought legacy is robust to short-term

perturbations. Furthermore, we reveal that soil microbiota from historically dry climates buffered

a wild grass species native to the precipitation gradient, but not the domesticated crop species

maize, from the negative effects of subsequent acute drought. In particular, microbiota with a legacy of chronic water limitation altered the expression of a small subset of host genes that mediate the effect of acute drought on transpiration and intrinsic water use efficiency. Our results reveal how long-term exposure to water stress alters soil microbial communities and demonstrate the resulting "legacy effects" on the drought responses of neighboring plants.

Main Text

Rapid global changes, such as escalating temperatures and changing precipitation patterns, intensify pressure on both natural and agricultural ecosystems. Among these changes, increasing frequency and intensity of droughts are particularly formidable challenges, with devastating consequences for plant health and survival. However, soil microbial communities, with their remarkable ability to quickly adapt to environmental shifts[2], may play a crucial role in bolstering the resilience of plants and ecosystems to drought stress[3]. Additionally, previous exposure to stress can shape the future trajectories of microbial communities and influence their responses to subsequent environmental disturbances, a phenomenon referred to as legacy effects or ecological memory[4]. Despite the growing recognition of microbial legacy effects, little is known about the mechanisms driving them, their long-term persistence, and whether their effects extend uniformly across different plant species. Furthermore, soil properties such as porosity or mineral nutrient content can also govern soil microbiomes[5], and vice versa, making it difficult to isolate the effects of precipitation on the microbiome.

Here, to investigate microbial drought legacy effects and isolate the drivers and impacts of microbial adaptations to water limitation, we (1) evaluate natural microbiome variation across two independent precipitation gradients, (2) test the ability of legacy effects to persist through experimental perturbation, and (3) evaluate the impacts of microbiome precipitation history on plant responses to acute drought at the molecular and physiological levels. Finally, we assess the

extent to which microbiome legacy effects are transferable across plant species. Our results

demonstrate that legacy effects of historical exposure to dry conditions are particularly salient at

the metatranscriptomic level and trigger transcriptional changes in plant roots that improve

resistance to subsequent acute droughts, at least in some plant species.

Mineral nutrient accumulation and historical precipitation impact soil microbiota

composition

To explore the relationship between free-living soil microbiota and water availability, we analysed

microbiome compositional and functional traits in soils exposed to a gradient of water availability.

First, we identified six never-irrigated remnant prairies along a steep precipitation gradient in

Kansas, USA (Fig. 1a, b and Supplementary Table S1). We sequenced the metagenomes of

these six soils to evaluate genetic variation within and among the bacterial communities along the

precipitation gradient. In general, we did not observe significant differences in bacterial alpha

diversity in the soils examined (Extended Data Fig.1a). In line with previous findings[6],

Actinomycetota and Bacillota were enriched in historically drier soils, while Pseudomonadota and

Acidobacteriota were more prevalent in wetter soils (Extended Data Fig. 1b). Supporting these

findings, between-sample variation showed a strong biogeographic signature (PERMANOVA, R^2

= 0.11, p = 0.001) and site precipitation explained 5.3% of the variation (Fig. 1c). The first axis of

principal coordinate analysis (PCoA) separated the bacterial communities of the two soils

exposed to the highest precipitation levels (CWR, WEL) from the rest of soils (Fig. 1c).

To disentangle the influence of soil edaphic properties and precipitation history on the microbiota

compositional changes detected across the gradient of soils, we examined 24 mineral nutrient

and trace element profiles (K, Fe, Mg, Ca, Zn, etc.) across all soils using ICP-MS. Soil nutrients

are known drivers of microbial taxonomic and functional composition, particularly bacterial

communities[7,8]. Consistent with the pattern seen for microbial taxonomic composition, we

observed significant mineral nutrient differences, with site explaining 27.9% of the variation and

precipitation explaining 28.6% (Extended Data Fig. 1c). The first principal coordinate axis of the

soil mineral nutrient profiles separated the three low-precipitation sites (TRI, SVR, and HAY), from

the three higher-precipitation sites (KNZ, CWR, and WEL) (Extended Data Fig. 1c).

Concentrations of several individual mineral nutrients were correlated with mean annual

precipitation (Extended Data Fig. 1d and e), and the overall mineral nutrient dissimilarities were

correlated with the microbiota composition dissimilarities across the gradient of soils (Mantel, r =

0.384; *p*-value = 1e-04; Extended Data Fig. 1f). This suggests that the accumulation of mineral

nutrients in these soils might be influenced by the precipitation patterns, and both may impact the

composition of the microbiota in the soil collection.

Next, we used X-ray computed tomography to quantify soil porosity of undisturbed soil cores. Soil

porosity, the percent of space or pores among soil aggregates, is directly related to soil hydraulic

properties such as water retention and infiltration[9]. In general, lower porosity results in lower water

retention and infiltration. Therefore, it is a good indicator for assessing the effect of precipitation

on the actual water content in the soil. In all cases, soil porosity decreased with depth and

increased with precipitation (Extended Data Fig. 2a-b). These results suggest that precipitation

history might affect soil porosity in surface layers of soils[10], which could affect soil niche properties

and, consequently, the abundance of specific microbiota members. Nevertheless, we found no

correlation between the soil porosity dissimilarities and the microbiota composition dissimilarities

in these soils (Extended Data Fig. 2c). This suggests that soil porosity is not a key element

controlling the overall microbial community composition in these soils.

Fig. 1. Water-related soil memory traits are preserved across distant precipitation gradients. a. Map of Kansas, USA, showing the locations selected for collecting the soils used in this work. **b.** The graph shows the precipitation (mm/year) recorded for the Kansas soils over the period 1981 to 2021 in the NASA POWER database. Statistical significance between soils was determined via an ANOVA on a fit linear model and letters represent the compact letter display of a Tukey *post-hoc* test ($p < 0.05$). **c.** Principal coordinate analysis (PCoA) of the soil microbiota projection from Kansas soils representing a gradient of precipitation. Colors represent the individual soils. The bar on the left denotes the percentage of the variance explained by the different experimental variables. **d.** Bar graph showing the correlation coefficient calculated between each mineral nutrient abundance and the precipitation gradient across Kansas soils. Bars in orange are significant ($p < 0.05$). **e and f**. Comparative analysis of relative abundances of the water legacy bacterial markers identified in soils from Kansas (**e**) and Santiago Island (**f**) exposed to low or high

123 precipitation levels. Colored points represent the average value of standardized relative abundance for each
124 bacterial taxon. The line connecting both points is the difference between low and high-precipitation soils.
125 A black line indicates statistical significance ($q < 0.05$). **g**. Heatmap showing the standardized relative
126 abundance of validated bacterial markers of water legacy across both collections of soils exposed to distinct
127 precipitation levels (Bottom panel).
128

Soil bacterial taxonomic and functional enrichments suggest local adaptations to water

availability

Next, to identify microbial biomarkers of precipitation legacy, we modelled the relative

abundances of bacterial taxa in relation to precipitation levels while controlling for soil porosity

and soil elemental composition. This analysis revealed distinct clusters of bacterial taxa at multiple

taxonomic levels (NCBI taxID, species, and family) whose relative abundances varied significantly

along the precipitation gradient (Extended Data Fig. 2d-f). Across the soils, 19 bacterial taxa

(NCBI taxIDs) were positively correlated and nine were negatively correlated with precipitation

(Fig. 1d). Taken together, these results indicate that water availability in the soil likely modulates

the composition of soil bacterial communities, possibly through the selection of functions

necessary to adapt to water availability in the soil.

We used assembled metagenomic contigs to explore the enrichment of bacterial biological

processes in the soils spanning the precipitation gradient. To focus on functions that are

associated with water availability rather than site-specific variation, we collapsed our six soils into

two groups representing our sites with low-precipitation histories (TRI, SVR, HAY) and high-

precipitation histories (KNZ, CWR, WEL). These groupings preserved the observed similarities in

microbiome taxonomic composition, mineral nutrient content, and porosity (Fig. 1c and Extended

Data Fig. 1c, 2a). We identified 62 Gene Ontology (GO) categories that were differentially

abundant between these groups (Extended Data Fig. 2g). Biological processes typically

associated with drought tolerance[11], such as nitrogen cycle metabolic processes, fatty acid (a

major cell membrane structural component) biosynthesis, DNA repair, and glucan metabolism

were enriched in low-precipitation soils (Extended Data Fig. 2g)[12–14], while processes related to

ion transport, lipid catabolism, and cellular aldehyde and ketone metabolism were depleted

(Extended Data Fig. 2g). These results confirm that the bacterial communities from the low-

precipitation soils are functionally different from those associated with high-precipitation soils, and

even with multiple soil factors influencing these communities, there is still clear evidence of

adaptations to osmotic stress. From this, we can conclude that the bacterial communities across

the Kansas precipitation gradient are functionally adapted to local precipitation levels, making

them excellent candidates for exploring how precipitation legacy of the soil microbiome affects

drought tolerance in plants.

Lastly, to isolate bacterial biomarkers of water availability in soils, we fit the relative abundances

of the most abundant (>0.1%) and prevalent (>20%) taxa to five statistical models that

incorporated mean annual precipitation, soil mineral nutrient composition, and soil porosity as

predictors. This analysis revealed seven bacterial taxa (NCBI taxaIDs) belonging to

Bradyrhizobiaceae, *Enterobacteriaceae*, *Polyangiaceae*, *Pseudomonadaceae*,

Rubrobacteraceae, *Streptomycetaceae*, and *Vicinamibacteraceae* enriched in soils with low

precipitation history (Fig. 1e). We also identified 7 taxa from *Acidobacteriaceae*, *Bacillaceae*,

Bradyrhizobiaceae, *Chitinophagaceae*, and *Enterobacteriaceae* that were enriched in soils with

high precipitation history (Fig. 1e). Therefore, we postulated that these precipitation-sensitive

bacteria can serve as biomarkers of water availability and likely contribute to the formation of

drought memory in these soils.

Distant natural precipitation gradients have conserved metagenomic properties

To validate that these bacterial community changes are generally associated with precipitation

patterns, rather than being specific to Kansas, we repeated our analyses for nine never-irrigated

(rain-fed) natural soils from a second precipitation gradient in Santiago Island, Cabo Verde

(Extended data Fig. 3a). Compared to the steep Kansas macro-gradient, these soils represent a

stable micro-gradient of precipitation (Extended Data Fig. 3b). Like in Kansas, bacterial alpha

diversity was similar across all soil samples (Extended Data Fig. 1a and 3c), and Actinomycetota

was enriched, while Pseudomonadota was depleted in soils with the lowest precipitation

(Extended Data Fig. 3d). We observed small differences in bacterial beta diversity among the

soils, partially explained by their geographical location (Extended Data Fig. 3e). As expected due

to the smaller geographic area covered by these samples, a principal coordinates analysis

showed a reduced biogeographic signature in the bacterial community composition of Santiago

soils compared to Kansas soils (Fig. 1c and Extended Data Fig. 3e).

Unlike in the Kansas soils (Extended Data Fig. 1d–f), neither the mineral nutrient profiles

(Extended Data Fig. 3f and g) nor the porosity profiles (Extended Data Fig. 4a) of the Santiago

soils corresponded to high- and low-precipitation groups. However, several individual mineral

nutrients were correlated with precipitation (Extended Data Fig. 3h) and mineral nutrient

dissimilarities were correlated with microbiota dissimilarities in these soils (r = 0.541; *p*-value =

1e-04; Extended Data Fig. 3i).

Soil porosity was correlated with neither mean annual precipitation (Extended Data Fig. 4b) nor

mineral nutrient content in the soils (Extended Data Fig. 4c). Consistent with findings from Kansas

(Extended Data Fig. 2c), soil porosity dissimilarities were uncorrelated with microbiota

composition dissimilarities in the Santiago soils (Extended Data Fig. 4d). Despite the reduced

biogeographic signature in the Santiago soils (Extended Data Fig. 3e), we speculated that a

reduced set of bacterial biomarkers that are highly sensitive to minor but consistent differences in

water availability may be detectable.

To determine whether small variations in precipitation affect specific members of the bacterial

communities, we identified clusters of bacterial taxa with differential relative abundance in high-

precipitation *vs.* low-precipitation soils (Extended Data Fig. 4e). These differences were most

pronounced when comparing soils at the extremes of the precipitation micro-gradient (Extended

Data Fig. 4e) and were consistent across taxonomic levels (Extended Data Fig. 4e-g). We found

22 bacterial taxa that were positively correlated with mean annual precipitation, and 10 that were

negatively correlated (Extended Data Fig. 4h). The patterns of bacterial GO term

enrichment/depletion due to precipitation, such as ornithine metabolic processes and purine

nucleoside triphosphate metabolic processes (enriched in drier soils) and lipid catabolic process

and ionic homeostasis (important for maintaining cell turgor, enriched in wetter soils), were similar

to those observed in the Kansas gradient (Extended Fig. 2g and Extended Data Fig. 4i). This

indicates that the strategies used by soil bacteria to cope with water limitation are robust to

variation in other edaphic properties and are conserved across geographically distant regions.

By applying the same procedures used to identify biomarkers in the Kansas soils, we identified

19 bacterial taxa in Santiago soils whose relative abundances responded to mean annual

precipitation after controlling for soil mineral nutrient content and porosity. Nine taxa (from

Archangiaceae,   *Bacillaceae*,   *Hymenobacteraceae*,   *Methylobacteriaceae*,   and

Rubrobacteraceae) were enriched in low-precipitation soils while ten (from *Bradyrhizobiaceae*,

Chitinophagaceae,   *Enterobacteriaceae*,   *Methylobacteriaceae*,   *Nitrosomonadaceae*,

Polyangiaceae, *Rhizobiaceae*, *Vicinamibacteraceae*, *Xanthomonadaceae*) were depleted in low-

precipitation soils (Fig 1f and Extended Data Fig. 3b). Notably, seven of these families—

Bradyrhizobiaceae,   *Enterobacteriaceae*,   *Polyangiaceae*,   *Rubrobacteraceae*,

Vicinamibacteraceae, Bacillaceae, and *Chitinophagaceae*—were also identified as biomarkers of

soil precipitation legacy in the Kansas soils. To validate the consistency of these bacterial

biomarkers across different geographic regions, we examined whether the enrichment patterns

of the Kansas biomarkers were conserved in Santiago soils and vice versa (Extended Data Fig.

4j-k and Supplementary Table S2). We validated eight bacterial markers from Kansas and eleven

biomarkers from Santiago as showing the same pattern of enrichment across both precipitation

gradients (Extended Data Fig. 4j-k). In particular, three bacterial taxa (*Rubrobacteraceae*

2653851, *Bradyrhizobiaceae* 1355477, *Enterobacteriaceae* 573) were consistently identified in

both soil gradients, validating them as robust bacterial biomarkers of soil water legacy (Fig. 1g).

Thus, we conclude that the functional and taxonomic differences among our Kansas soils

accurately represent the legacy effects of chronic water limitation and are likely generalizable to

other water-limited soil microbiomes.

Strain-level genetic analysis identifies candidate genes for bacterial drought tolerance

Finally, we investigated whether there were genetic differences between bacterial lineages across

the precipitation gradient, focusing on the identified bacterial markers and other highly abundant

and prevalent taxa from both Kansas and Santiago soils. We mapped the shotgun metagenomic

sequence reads from each sample to reference genomes of the focal species, and subsequently

identified sequence variants across the genomes (Fig. 1e-g). Analysis of the genetic distances

showed variations in strain-level microbiome structure across both the Kansas and Santiago

precipitation gradients (Extended Data Fig. 5a), between precipitation levels (Extended Data Fig.

5b), and between geographical regions (Extended Data Fig. 5c).

Subsequently, we conducted a genotype-environment association analysis to identify genetic

variants associated with mean annual precipitation. Significant associations were found in the

genomes of the biomarkers *Luteitalea* (ID1855912), *Bradyrhizobium* (ID1355477),

Bradyrhizobium (ID1325100), *Edaphobacter* (ID2703788), *Candidatus* Solibacter (ID332163),

and *Microvirga* (ID2807101) (Extended Data Fig. 5d). All identified variants were located within

protein-coding regions, specifically the LuPra_RS24355 (DNA-directed RNA polymerase subunit

beta), Bdiaspc4_RS17620 (epoxide hydrolase family protein), XH93_RS32435 (response

regulator transcription factor), GWR55_RS00680 (NADP-dependent oxidoreductase),

GWR55_RS05605 (glucose-6-phosphate isomerase), HYX25_05330 (carbamoyl-phosphate

synthase large subunit), and JO965_13190 (IS630 family transposase) (Extended Data Fig. 5d).

These associations suggest candidate genes related to drought tolerance in free-living bacteria

(Extended Data Fig. 5d). These results indicate that in addition to the bacterial biomarkers of

precipitation history identified at species or higher taxonomic level (Supplementary Table S2),

precipitation legacy effects likely manifest through genetic differentiation of bacterial lineages

within taxa. These groups of water-sensitive taxa might be critical for the formation of soil memory

traits, and their study could reveal new ecological mechanisms by which soil drought legacy drives

responses to water stress in plants.

Metagenomic and especially metatranscriptomic precipitation legacies are resilient to

short-term perturbations

To assess the effects of short-term perturbations on soil microbiome legacy, we exposed the six

Kansas soils to a five-month-long conditioning phase in a growth chamber. Replicate pots of each

soil were either left unplanted or planted with a seedling of the grass species *Tripsacum*

dactyloides (eastern gamagrass, which is native to Kansas), and were either drought-challenged

or well-watered, in a factorial design (Extended Data Fig. 6a). To confirm that our experimental

acute drought applied had physiological consequences in our system, we quantified various

morphological and anatomical features of the eastern gamagrass plants (Fig. 2a and Extended

Data Fig. 6b-d). Compared to well-watered plants, the plants exposed to acute drought treatments

were shorter and had more root aerenchyma (Fig. 2a and Extended Data Fig. 6b-c), which are

both indicative of drought stress[15]. This indicates that our experimental drought was sufficient to

induce differential responses in the plants and, presumably, in the soil-dwelling microbes.

Next, we explored how precipitation legacy affected the responses of free-living (bulk soil) and

gamagrass-associated (rhizosphere) microbial communities when exposed to intermittent

drought and watering events. Congruent with the pattern seen across the natural precipitation

gradient, bacterial community alpha diversity was similar across treatment groups (Extended Data

Fig. 6e). There were also no major taxonomic profile shifts across treatments at the phylum level

(Extended Data Fig. 6f). However, canonical analysis of principal coordinates of metagenomic

sequences indicated that the conditioning water and host treatments impacted metagenome

content but only explained 4.3% and 1.1% of the variation in the data, respectively (Extended

Data Fig. 6g). In contrast, precipitation history explained 14.1% of the variation in bacterial

community composition, confirming that the conditioning treatments did not erase the ecological

memory of the soils (Fig. 2b). In fact, we still detected significant differences in bacterial

enrichment between the two soil groups with different precipitation legacies across taxonomic

levels, regardless of the treatment applied (Extended Data Fig. 6h). For instance,

Ktedonosporobacteraceae and *Rubrobacteraceae* were consistently depleted and enriched,

respectively, in all low-precipitation soils compared to high-precipitation soils across all treatments

(Extended Data Fig. 6h). Furthermore, the identified bacterial markers for drought legacy (Fig. 1d-

f) generally followed the expected enrichment and depletion patterns (Extended Data Fig. 6i),

which remained robust despite the applied treatments (Extended Data Fig. 6i). This resilience

was particularly evident in soils with low-precipitation legacies (Fig. 2c and Extended Data Fig.

6i).

To further validate the robustness of the water soil legacy to experimental perturbation, we

investigated patterns of strain-level genetic variation in the identified bacterial markers and other

prevalent species across the samples, as described previously (Fig. 1e-g). We aligned the

shotgun metagenomic sequence reads to the reference genomes of the focal species to identify

sequence variants, then conducted a principal coordinates analysis of standardized pairwise

Fig. 2. Soil drought microbial legacy is resilient to short-term perturbations. a. Box plot showing the quantification of different *Tripsacum dactyloides* phenotypes in response to the treatments applied (drought conditions or regular watering) during the conditioning phase. In the graph, the data from soils with low precipitation legacy (low ppt legacy) is colored in brown while the data from soils with high precipitation legacy (high ppt legacy) is colored in blue (see also Data extended Fig. 6b). ANOVA was used to discern significant differences among groups and means were separated using Tukey's post hoc test. **b.** Canonical analysis of principal coordinate showing the projection of the microbiota of each soil with wet and dry legacy after the different treatments. Notice that the soil legacy separates the samples into two groups on the first axis and explains a significant proportion of the observed variance (left bar). **c.** Heatmaps showing the soil enrichment patterns (in relation to soil baseline (wet or dry legacy)) of different drought legacy markers across the collection of soils exposed to the different treatments. The tiles of the heatmaps are colored based on the log2 fold change estimated from a generalized linear model contrasting the abundance of each bacterial marker (y-axis) in a given soil (x-axis) against their corresponding soil baseline. The bacterial markers (y-axis) are clustered. Squares outlined in black represent bacterial markers significantly enriched (red) and depleted (blue) with respect to the corresponding soil baseline ($q < 0.1$). **d.** Canonical analysis of principal coordinates showing the projection of the metatranscriptome of each soil with wet and dry legacy after the different conditioning treatments. Notice that the soil legacy (wet and dry) separates the samples into two groups on the first axis and explains a significant proportion of the observed variance (left bar). **e.** Principal coordinate analysis based on standardized pairwise genetic distances showing the projected identified SNPs found in the genome of the drought legacy markers. Notice that despite the conditioning perturbations applied there is a clear separation of the samples on the first axis based on the soil water legacy (drought and precipitation).

genetic distances based on those variants. This revealed a clear separation of the samples in the

first axis based on the soil water legacy (Fig. 2e) rather than the experimental treatments.

Consistent with our previous findings (Fig. 2), our experimental perturbations did not affect allele

frequencies (Extended Data Fig. 7a,b). Additionally, a genome-wide association analysis showed

significant associations with genes from the bacterial markers *Luteitalea* (ID1855912) and

Bradyrhizobium (ID1355477), which had been previously identified across the precipitation

gradients (Extended Data Fig. 5d, 7c). Additional associations were also detected with the

biomarker *Bradyrhizobium*: ID1325107 (Extended Data Fig. 7c).

Next, to investigate the robustness of drought legacy effects on bacterial functions, we tested

whether enrichment of GO terms between low-precipitation and high-precipitation soils was

maintained after experimental perturbation (Extended Data Fig. 8a). Again, we found a consistent

enrichment of GO terms related to the nitrogen cycle metabolic process, including

purine−containing compound metabolic process and pyrimidine−containing compound metabolic

process, and a depletion of ion transport and amino acid catabolic process. Some of these GO

categories were also identified in the free-living communities from Kansas and Santiago soils

(Extended Fig. 2g, 4i, and 8a). These results show that the legacy effect of precipitation on the

functional content of soil metagenomes was robust to the perturbations applied during the

conditioning phase.

To further validate this conclusion, we analysed metatranscriptome data from the same samples

to confirm which biological processes were actually active across the different treatments and

soils. A canonical analysis of principal coordinates showed that precipitation history explained

24.9% of the variation in microbial gene expression while conditioning phase water treatment and

host treatment explained only 12.3% and 1.8% of the variation, respectively (Fig. 2d and Extended

Data Fig. 8b). Indeed, we confirmed that the conditioning phase treatments did not alter the

functional characteristics of the main bacterial phyla in these soils: even after five months of

experimental perturbation, functionally active Actinomycetota and Bacillota remained enriched in

all historically dry soils while Acidobacteriota, Planctomycetota, and Pseudomonadota remained

enriched in all historically wet soils (Extended Data Fig. 8c). Furthermore, in agreement with our

previous observations (Extended Data Fig. 6h), conserved patterns of functional activation within

multiple taxonomic groups were detected across both soil groups, regardless of the conditioning

treatments applied (Extended Data Fig. 8d-e). Also, the metatranscriptome analysis revealed GO

categories that remained enriched in low-precipitation soils after the applied conditioning

treatments, such as tetrapyrrole metabolic process, response to osmotic stress, liposaccharide

metabolic process, heme metabolic process, and trehalose catabolic process (Extended Data

Fig. 8f). Therefore, we confirmed that the long-term legacy of precipitation on soil bacterial

community function is robust to short-term (five months) perturbations, including acute drought

and the introduction of a host plant. This functional memory of the soil microbial community could

influence host responses to water changes in natural ecosystems, potentially providing protection

against subsequent acute drought.

Microbiome precipitation legacy alters host plant responses to subsequent acute drought

despite taxonomic convergence of root microbiomes

During the conditioning phase, we observed that the *T. dactyloides* plants grown in low-

precipitation-legacy soils had increased shoot height, root and shoot mass, and total aerenchyma

area under drought treatment compared to plants grown in high-precipitation-legacy soils (Fig. 2a

and Extended Data Fig. 6c). To confirm that these effects of soil precipitation legacy on plant

phenotype (Fig. 2a) were conferred by the microbiome rather than co-varying soil properties, we

extracted the experimentally-conditioned microbial communities and used them to inoculate a

new generation of *T. dactyloides* cv. "Pete". These "test phase" plants were divided between

water-limited conditions (⅔ of all plants) and well-watered control conditions (⅓ of plants;

Extended Data Fig. 6a). We confirmed that our test phase drought treatment impaired plant

growth (Extended Data Fig. 9a). After five weeks we uprooted gamagrass plants for phenotyping

and sampled crown roots, which are highly active in water acquisition[16,17], for 16S rRNA gene

microbiome profiling and root gene expression analysis. Our focus only on bacteria was motivated

by past work showing that root-colonizing fungi from these soils were insensitive to drought[18] and

by our observation that >97% of all metagenomic reads in this study were bacterial. To capture

the effects of the precipitation gradient extremes, we measured plant gene expression only for

plants that received inocula derived from the two easternmost (wet-legacy) and two westernmost

(dry-legacy) soils.

We found that precipitation legacy did not affect the alpha diversity (Shannon diversity) or the

taxonomic composition of the gamagrass root bacterial community as measured by 16S rRNA

gene sequencing (Fig. 3a). The taxonomic composition of the crown root bacterial community

was only impacted by the test phase water treatment (Fig. 3b; ANOVA.CCA, $F_{1,139}$=2.57,

R^2=0.016, p = 0.004), not legacy (Fig. 3c), which only explained 1.6% of the variation in the data.

We further assessed root microbiome stability by calculating beta-dispersion, a measure of within-

group variability. Neither test phase water treatment, inoculum legacy, nor their interaction

impacted gamagrass root bacterial community beta-dispersion (Fig. 3d; ANOVA $F_{1,144}$=0.571, p =

0.45). Altogether, these results suggest that gamagrass exerts a strong homogenizing influence

during root microbiome formation, resulting in the assembly of a stable and drought-resistant

microbiota. The stability of the gamagrass root bacterial microbiome, even after receiving

compositionally different inocula, indicates a deterministic process (not stochastic) of community

assembly. Because eastern gamagrass (and the "Pete" cultivar in particular[19]) shares a long

evolutionary history with Kansas soil microbes, this result supports previous findings that co-

evolution promotes stable community assembly[20,21].

Fig. 3. Test phase gamagrass root microbiome is remarkably stable. **a.** Shannon diversity was not impacted by precipitation legacy or test phase water treatment. Points are EMMs and error bars represent the standard error. ANOVA was used to discern significant differences among groups and means were separated using Tukey's post hoc test. **b**. A canonical analysis of principal coordinates indicates that test phase water treatment, but not legacy (**c**) significantly impacted the gamagrass root bacterial microbiome composition. Ellipses represent the 95% confidence intervals. **d.** Drought treatment did not impact the root microbiome within group variation (beta-dispersion). Points are EMMs and error bars represent the standard error. **e**. Maize root microbiome Shannon diversity was significantly impacted by test phase water treatment, but not legacy. Points are EMMs and error bars represent the standard error. ANOVA was used to discern significant differences among groups and means were separated using Tukey's post hoc test. **f**. Legacy and Test phase water treatment (**g**) significantly impacted the maize root bacterial microbiome composition. Ellipses represent the 95% confidence intervals. **h.** Acute drought treatment also increased maize root microbiome within group variation. Points are EMMs and error bars represent the standard error.

Although precipitation legacy did not significantly shape 16S rRNA gene diversity within the *T. dactyloides* root microbiome, its clear effect on metagenome content and microbial gene expression (Fig. 2e, 2f, and 2g and Extended Data Fig. 8f) suggested high potential to influence plant phenotypes. Indeed, RNA-seq analysis confirmed that microbial inocula with contrasting precipitation legacies differentially influenced host gene expression in the crown roots. In addition

to 15 *T. dactyloides* genes that were differentially expressed depending on whether the inoculum

originated from a high-precipitation or low-precipitation region (Fig. 4a; Supplementary Table S3),

we observed that microbial legacy influenced the responses of 183 genes to acute drought (Fig.

4b). The majority of these (55%) were unresponsive to acute drought in plants grown with a dry-

legacy microbiota, but were significantly down-regulated or up-regulated in plants grown with a

wet-legacy microbiota (Fig. 4b, gene sets I and II, respectively; Supplementary Table S3). This

strongly suggests that the ecological memory of drought transmitted through soil microbiota

subdued the overall transcriptional response of gamagrass to acute drought. The 50 *T.*

dactyloides genes that decreased expression under acute drought only in the presence of a wet-

legacy microbiome (Fig. 4b gene set I) included five that are orthologous to maize genes predicted

to be involved in ethylene- or ABA-mediated signalling of water stress (Td00002ba004498,

Td00002ba024351, Td00002ba011993, Td00002ba005402, Td00002ba000033) and a heat

shock protein linked to temperature stress (Td00002ba042486) (Supplementary Data 1). Notably,

the latter three genes were not identified as drought-sensitive when averaging across inocula,

demonstrating that the microbial context is necessary for a complete understanding of molecular

plant drought responses. We note that these signatures of microbiota legacy on the plant

transcriptome were detectable even when averaging across all treatments applied during the

conditioning phase (Extended data Fig. 6a), again confirming that microbiome legacy effects with

implications for plant fitness are robust to short-term perturbations.

Next, we phenotyped the test phase plants to determine whether microbial precipitation legacy

altered host physiological and morphological responses to a subsequent acute drought. For each

of 63 traits, including mineral and nutrient content, mass and height, photosynthesis and gas

exchange rates, crown root anatomy, and root system architecture, we calculated a drought

susceptibility index (S-index) that measures the stability of a trait under drought relative to control

plants that received the same inoculum type (Supplementary Table S4). We used a random forest

Fig. 4. Drought legacy of microbial inocula alters the response to drought in eastern gamagrass. a. 15 genes were up-regulated in plants inoculated with microbiota from a low-precipitation region, relative to plants inoculated with microbiota from a high-precipitation region. **b.** In total, 183 *T. dactyloides* genes responded to acute drought in a manner that was dependent on the drought legacy of the soil microbiota, regardless of the inoculum's treatment during the conditioning phase (Supplementary Data 1). Only genes with a drought response of ≥ 4-fold and with annotated maize ortholog(s) are shown here for illustration purposes. Each pair of points shows one gene; the relative position of the two points illustrates how the gene's expression changed in response to acute drought during the Test Phase, depending on whether the plant had been inoculated with microbiota derived from a low-precipitation or high-precipitation environment. Genes are grouped into sets according to the pattern of how inoculum legacy altered their drought responses. **c.** The model used for mediation analysis to test whether the expression of the 198 legacy-sensitive genes (see panels a-b) contributed to the overall effect of acute drought on plant phenotype. **d.** Mediation analysis confirmed that the expression of legacy-sensitive genes is part of the plant response to acute drought resulting in decreased intrinsic water-use efficiency (iWUE, units: μmol/mol) and transpiration (E, units: mol m⁻² s⁻¹). Error bars indicate 95% confidence intervals. **e.** Microbiomes with a low-precipitation legacy caused increased iWUE and E during acute drought, relative to inocula derived from high-precipitation-legacy soils. The s-index describes trait values scaled by the mean value of well-watered control plants, such that an s-index of 1 indicates that droughted plants were phenotypically identical to non-droughted plants.

model to rank and identify the ten most important traits for describing the phenotypic effects of

microbiome precipitation legacy (Extended Data Fig. 10a,b). We then summarized the variation

in these top traits using constrained redundancy analysis, which revealed that microbiome

precipitation legacy explained 5% of the total variation in host phenotypic response to acute

drought (ANOVA.CCA, $F_{1,187}$ = 10.1, R^2 = 0.05, p = 0.001) (Extended Data Fig. 11a). Microbiota

precipitation legacy significantly impacted six of the top eight non-correlated traits: transpiration,

leaf chlorophyll content, max root width, root total metaxylem area, median root diameter, Cu, K,

and intrinsic water use efficiency (iWUE) – an important plant drought response trait[22,23] (Extended

Data Fig. 11b-j).

To solidify the mechanisms linking microbiome legacy to plant phenotype, we used mediation

analysis to determine whether the 198 genes that were differentially expressed depending on

inoculum legacy (Fig. 4a-b) were causally involved in trait responses to acute drought (Fig. 4c).

Expression patterns of this subset of genes were summarized in two dimensions (MDS1 and

MDS2) using non-metric multidimensional scaling of Bray-Curtis dissimilarities among individual

plants. MDS1 mediated 18% of the total drought-induced decrease in iWUE (p < 0.001), and 8.9%

of the total decrease in transpiration (p = 0.018; Fig. 4d). In contrast, MDS2 mediated 11% of the

iWUE response to drought (p = 0.03), but in the opposite direction, meaning that the activity of

these genes counteracted the negative effects of acute drought on iWUE. Four of the top eight

genes with the strongest positive loadings on MDS2 were orthologs of the maize gene ZmNAS7,

which encodes the enzyme nicotianamine synthase; two of these were also in the top 5% of genes

with the strongest negative loadings on MDS1 (Extended Data Fig. 12), indicating that they have

a protective function that minimizes the drop in iWUE during acute drought. All four ZmNAS7

orthologs were up-regulated between 4-fold and 12-fold in response to drought, but only in plants

that were inoculated with dry-legacy microbiota (Fig. 4b, gene set IV), suggesting a possible

mechanism by which microbiota with a legacy of chronic water stress confer drought tolerance.

Consistent with this, salicylic acid–a key regulator of root microbiome assembly[24]–is reported to

induce expression of ZmNAS7[25]. Furthermore, although nicotianamine is best known for its role

in metal transport, the overexpression of NAS genes in rice conferred tolerance of acute drought,

including maintenance of photochemical efficiency at pre-drought levels[26]. Overall, these results

show that microbiomes with a history of water-stress improve the drought response of eastern

gamagrass, and that a complete understanding of plant drought response requires consideration

of the microbial environment, including legacy effects of water scarcity.

Beneficial microbiome drought legacy effects do not extend to the crop species *Zea mays*

To explore whether the observed microbial legacy effects were transferable across hosts, we

simultaneously applied the same inocula to seedlings of the maize (*Zea mays* ssp. *mays*)

genotype B73. Maize is genetically closely related to gamagrass, making it a good candidate for

testing whether microbiome legacy effects are transferable to domesticated, non-native species.

Maize plants were four weeks old at the time of sampling, and the drought treatment was sufficient

to hinder maize growth (Extended Data Fig. 9b). Unlike in eastern gamagrass, the taxonomic

composition of bacterial communities in maize crown roots retained clear signatures of the

inoculum precipitation legacy (Fig. 3g., ANOVA.CCA, $F_{1,255}$ = 4.254, R^2 = 0.014, p = 0.001), in

addition to being impacted by acute drought treatment (Fig. 3f). Also, unlike gamagrass, acute

drought significantly increased beta-dispersion compared to the well-watered control, indicating

that water deprivation gave rise to more variability in the maize microbiome (Fig. 3h; ANOVA $F_{1,262}$

= 18.17, p < 0.001). Like in eastern gamagrass, however, all inocula resulted in root communities

with a similar mean Shannon diversity, but drought treatment significantly decreased alpha

diversity (Fig. 3e). Next, we identified bacterial amplicon sequence variants (ASVs) that were

differentially abundant in droughted *vs.* control plants in a manner that depended on the

precipitation legacy of the microbiome. This included only two ASVs in gamagrass roots,

ASV1598 (*Azospirillum* sp.) and ASV2352 (*Enterobacteriaceae*) (Extended Data Fig. 13a;

Supplementary Table S5), but 100 ASVs in maize roots (Extended Data Fig. 13b; Supplementary

Table S5), again indicating that gamagrass root microbiomes are more stable than those of maize.

The maize ASV enrichments did not show a phylogenetic pattern. We speculated that these

drought-responsive ASVs could contribute to plant phenotypic responses to drought.

To validate this, we also characterized the transcriptional response of maize and found that 23

genes were up-regulated in plants inoculated with dry-legacy microbiota, relative to those that

received wet-legacy inocula (Extended Data Fig. 14a). These included six defense-related genes,

particularly those related to jasmonic acid signaling, suggesting that soil microbiota from low-

precipitation sites were perceived as pathogens by maize, but not eastern gamagrass. Other

notable functions that were up-regulated by the dry-legacy inocula in maize include root

development and iron acquisition (Supplementary Table S3). We also identified 109 maize genes

that responded to acute drought in a manner that was dependent on microbial precipitation

legacy, of which only 30 were identified as drought-responsive when averaging across inocula

(Extended Data Fig. 14b). Like in eastern gamagrass, most of these were drought-responsive

only in wet-legacy-inoculated or only in dry-legacy-inoculated plants, with relatively few genes

reversing the direction of their drought response depending on inoculum legacy (Extended Data

Fig. 14c). For example, 30 genes were up-regulated in response to drought only in plants grown

with wet-legacy microbiota (Extended Data Fig. 14c, gene set I). These genes included several

related to pathogen defense (Zm00001eb116230, Zm00001eb150050, Zm00001eb222540) and

one involved in the response to symbiotic fungi (Zm00001eb033580), suggesting that soils from

high-precipitation regions contain both harmful and beneficial microorganisms and that

interactions with both types are activated under water deprivation in maize (Supplementary Table

S3). Notably, one of the 18 genes that reversed its drought response depending on the inoculum

was TIP3, which encodes an aquaporin that regulates water storage in the vacuole[27]. In the

presence of wet-legacy microbiota, TIP3 expression increased 7-fold during acute drought,

whereas in the presence of dry-legacy microbiota TIP3 expression decreased nearly 3-fold

(Extended Data Fig. 14c, gene set VI). Averaged across the two inocula, however, TIP3 was not

differentially expressed between droughted and well-watered plants; thus, its role in drought

response was apparent only when accounting for microbiota legacy.

We identified four sets of orthologous genes that were sensitive to microbiota legacy in both host

species. Zm00001eb076200 encodes a peroxidase that reversed its drought response depending

on inoculum in maize; however, its *T. dactyloides* ortholog, Td00002ba025285, was up-regulated

4-fold due to drought but only in plants grown with wet-legacy inoculum. The gene

Zm00001eb077640, encoding L-allo-threonine aldolase, also reversed its drought response in

maize. In contrast, its ortholog Td00002ba026366 was consistently up-regulated in response to

drought, but more so in plants inoculated with dry-legacy biota. The maize pathogenesis-related

protein-like gene Zm00001eb150050, which has been linked to both biotic and abiotic stress

responses, was up-regulated 7-fold in response to drought only in plants inoculated with wet-

legacy microbiota; however, both of its *T. dactyloides* orthologs were down-regulated 8-fold in

plants inoculated with microbiota from low-precipitation sites, relative to inocula from high-

precipitation sites, regardless of water availability. Finally, Zm00001eb301540 encodes a

tryptophan synthase that was up-regulated 1.7-fold in the presence of wet-legacy but not dry-

legacy microbes; its ortholog Td00002ba012570, in contrast, was down-regulated in the presence

of wet-legacy microbes. Thus, none of these four sets of orthologous genes showed congruent

responses to drought in maize and eastern gamagrass.

Finally, we investigated whether the beneficial effects of microbiome legacy on *T. dactyloides*

physiological and morphological phenotypes (Fig. 4d-e) were also conferred on maize. Overall,

microbiome precipitation legacy explained only 2.0% of the phenotypic variation in maize

(Extended Data Fig. 15a., ANOVA.CCA, $F_{1,175}$ = 3.72, R^2 = 0.02, p = 0.001), a weaker effect than

observed in eastern gamagrass. In fact, in general, we observed no significant phenotypic

differences attributable to microbiome legacy (Extended Data Fig. 15b,c). Thus, the capacity to

integrate the beneficial effect of soil drought legacy is host specific and it is related to the ability

of the host to maintain a stable microbiota in response to drought stress.

Conclusions

Microbially-mediated legacy effects of past water-limited conditions can affect plant performance

during subsequent droughts[28,29]. Our results reveal the importance of these stable signatures of

historical precipitation in microbiota composition and function not only for microbial adaptation,

but also for host survival under water-scarce conditions. We identified bacterial markers and

numerous microbial genes and functional pathways that are associated with mean annual

precipitation, including those suggesting an increased nitrogen-metabolizing and DNA repair

capacity in dry-legacy microbiota. We validated these findings by controlling for other soil

properties that influence microbial communities, namely soil porosity and mineral nutrient content.

Furthermore, we confirmed that these metagenomic patterns are shared by two independent

natural precipitation gradients separated by more than 7,000 km, suggesting that these signatures

of historic precipitation levels may generalize to soils across the globe. Thus, we provide evidence

for the ecological mechanisms involved in the formation of soil drought memory and demonstrate

its robustness to a five-month perturbation, which could mimic a particularly dry or wet season.

We have shown that these metagenomic precipitation legacies have significant implications for

plant response to drought. Inoculation with dry-legacy microbiota altered the transcription of key

genes controlling drought responses in the plant. In the prairie grass *Tripsacum dactyloides*, dry-

and wet-legacy soil microbiomes give rise to taxonomically similar root endosphere microbial

communities yet have strikingly different effects on plant gene expression and phenotypes during

a subsequent acute drought. This effect was less obvious in maize, which has a relatively unstable

root microbiome and reduced physiological and morphological sensitivity to soil microbiome

legacy during drought. Based on these findings, root microbiome stability appears to play a role

in adaptive plant responses to a subsequent acute drought. Importantly, differences in gamagrass

and maize responses to microbiome history indicate that crops may not reap the same benefits

as native plant species from potentially beneficial microbial communities. Therefore, our

discoveries significantly contribute to our mechanistic understanding of microbiome drought

legacy effects, their resistance to perturbation, and their role in plant drought responses, with

implications for agricultural and ecosystem management.

Data availability

The 16S rRNA amplicon sequencing and metagenomic data associated with this study have been

deposited in the NCBI Sequence Read Archive under the BioProject IDs PRJNA1186942 and

PRJNA1179931, respectively. The raw transcriptomic and metatranscriptomic data have been

deposited in the Gene Expression Omnibus under accession XXXXX and XXXX, respectively.

We deposited all scripts and additional data structures required to reproduce the results of this

study in the following Zenodo repository: doi: 10.5281/zenodo.13821006. Source data are

provided with this paper.

Acknowledgements

We thank Arun Seetharam, Matthew Hufford, and the PanAnd consortium for granting early

access to the *Tripsacum dactyloides* genome and annotation. We thank Dr. Mitch Greer, Michaela

VonLintel, Justin Roemer, Dr. Alan Schlegel, and the Nature Conservancy for granting access to

soil collection sites in Kansas, and Dr. Peter Balint-Kurti for providing maize seeds. We thank Dr

Craig J. Sturrock and Dr Brian S. Atkinson from the Hounsfield Facility at the University of

Nottingham for their assistance with the analysis of the CT images. We are grateful to the

undergraduate student researchers and technicians who helped to set up experiments and collect

data: Felicity Tso, Hannah Reid, and Carmen Rodriguez. Thank you to Dr. Joel Swift, Ceyda

Kural-Rendon, Dr. Ben Sikes (University of Kansas), Dr. Manuel Kleiner (North Carolina State

University), Dr Omri Finkel (The Hebrew University of Jerusalem, Israel), and Dr Nicholas Girkin

(University of Nottingham, UK) for critical reading of the manuscript. This work was supported by

a UKRI/BBSRC - NSF/BIO Lead Agency Opportunity, National Science Foundation award #IOS-

2016351 and BBSRC grant #BB/V011294/1. Maize root transcriptome library preps and

sequencing were performed in the RTSF Genomics Core at Michigan State University. *T.*

dactyloides library preps and RNA-seq were performed at the NC State University Genomic

Sciences Laboratory (Raleigh, NC, USA). Research reported in this publication was made

possible in part by the services of the KU Genome Sequencing Core. This lab is supported by the

National Institute of General Medical Sciences (NIGMS) of the National Institutes of Health under

award number P30GM145499. We thank the University of Kansas Medical Center – Genomics

Core for generating the 16S amplicon sequence data sets. The Genomics Core is supported by

the Kansas Intellectual and Developmental Disabilities Research Center (NIH U54 HD 090216),

the Molecular Regulation of Cell Development and Differentiation – COBRE (P30 GM122731),

The NIH S10 High End Instrumentation Grant (NIH S10OD021743) and the Frontiers CTSA Grant

(UL1TR002366).

Author contributions

Conceptualization: N.A.G., V.C., D.G., G.C., and M.R.W. Data curation: N.A.G., V.C., D.G., and

M.R.W.; Formal analysis: N.A.G., V.C., D.G., I.S.-G., and M.R.W.; Investigation: N.A.G., V.C.,

D.G., N.F., D.D.J., D.M.W., A.M., G.C., and M.R.W.; Methodology: G.C. and M.R.W.; Project

administration: G.C. and M.R.W.; Software: N.A.G., V.C., D.G., and M.R.W.; Supervision: G.C.

and M.R.W.; Visualization: N.A.G., V.C., D.G., and M.R.W.; Writing – original draft: N.A.G., G.C.,

and M.R.W.; Writing – review & editing: V.C. and D.G.

Competing interests: The authors declare no competing interests.

References

1.   Schimel, J., Balser, T. C. & Wallenstein, M. Microbial stress-response physiology and its

implications for ecosystem function. *Ecology* **88**, 1386–1394 (2007).

2.   Chase, A. B., Weihe, C. & Martiny, J. B. H. Adaptive differentiation and rapid evolution of a

soil bacterium along a climate gradient. *Proc. Natl. Acad. Sci. U. S. A.* **118**, (2021).

3.   Lau, J. A. & Lennon, J. T. Rapid responses of soil microorganisms improve plant fitness in

novel environments. *Proc. Natl. Acad. Sci. U. S. A.* **109**, 14058–14062 (2012).

4.   Canarini, A. *et al.* Ecological memory of recurrent drought modifies soil processes via

changes in soil microbial community. *Nat. Commun.* **12**, 5308 (2021).

5.   Custódio, V. *et al.* Sculpting the soil microbiota. *Plant J.* **109**, 508–522 (2022).

6.   Naylor, D., DeGraaf, S., Purdom, E. & Coleman-Derr, D. Drought and host selection

influence bacterial community dynamics in the grass root microbiome. *ISME J.* **11**, 2691–

2704 (2017).

7.   Liu, Z. *et al.* Shifts in microbial communities and networks are correlated with the soil

ionome in a kiwifruit orchard under different fertilization regimes. *Appl. Soil Ecol.* **149**,

103517 (2020).

8.   Dai, Z. *et al.* Metallic micronutrients are associated with the structure and function of the

soil microbiome. *Nat. Commun.* **14**, 8456 (2023).

9.   Guellouz, L., Askri, B., Jaffré, J. & Bouhlila, R. Estimation of the soil hydraulic properties

from field data by solving an inverse problem. *Sci. Rep.* **10**, 9359 (2020).

10. Rousseva, S., Torri, D. & Pagliai, M. Effect of rain on the macroporosity at the soil surface.

Eur. J. Soil Sci. **53**, 83–93 (2002).

11. Naylor, D. & Coleman-Derr, D. Drought Stress and Root-Associated Bacterial

Communities. *Front. Plant Sci.* **8**, 2223 (2017).

12. Sun, Y., Wilkinson, B. J., Standiford, T. J., Akinbi, H. T. & O'Riordan, M. X. D. Fatty acids

regulate stress resistance and virulence factor production for Listeria monocytogenes. *J.*

Bacteriol. **194**, 5274–5284 (2012).

13. Woodcock, S. D. *et al.* Trehalose and α-glucan mediate distinct abiotic stress responses in

Pseudomonas aeruginosa. *PLoS Genet.* **17**, e1009524 (2021).

14. Gil-Quintana, E. *et al.* Local inhibition of nitrogen fixation and nodule metabolism in

drought-stressed soybean. *J. Exp. Bot.* **64**, 2171–2182 (2013).

15. Zhu, J., Brown, K. M. & Lynch, J. P. Root cortical aerenchyma improves the drought

tolerance of maize (*Zea mays* L.). *Plant Cell Environ.* **33**, 740–749 (2010).

16. Gao, Y. & Lynch, J. P. Reduced crown root number improves water acquisition under water

deficit stress in maize (Zea mays L.). *J. Exp. Bot.* **67**, 4545–4557 (2016).

17. Ahmed, M. A. *et al.* Root type matters: measurement of water uptake by seminal, crown,

and lateral roots in maize. *J. Exp. Bot.* **69**, 1199–1206 (2018).

18. Swift, J. F. *et al.* Drought stress homogenizes maize growth responses to diverse natural

soil microbiomes. *Plant Soil* (2024) doi:10.1007/s11104-024-06853-x.

19. Springer, T. L. & Dewald, C. L. Eastern gamagrass and other*Tripsacum*species. in *Warm-*

Season (C4) Grasses 955–973 (American Society of Agronomy, Crop Science Society of

America, Soil Science Society of America, Madison, WI, USA, 2016).

20. Batstone, R. T., O'Brien, A. M., Harrison, T. L. & Frederickson, M. E. Experimental

evolution makes microbes more cooperative with their local host genotype. *Science* **370**,

476–478 (2020).

21. Morella, N. M. *et al.* Successive passaging of a plant-associated microbiome reveals robust

habitat and host genotype-dependent selection. *Proceedings of the National Academy of*

Sciences **117**, 1148–1159 (2020).

22. Hund, A., Ruta, N. & Liedgens, M. Rooting depth and water use efficiency of tropical maize

inbred lines, differing in drought tolerance. *Plant Soil* **318**, 311–325 (2009).

23. Schulz, P. *et al.* Improving plant drought tolerance and growth under water limitation

through combinatorial engineering of signalling networks. *Plant Biotechnol. J.* **19**, 74–86

(2021).

24. Lebeis, S. L. *et al.* PLANT MICROBIOME. Salicylic acid modulates colonization of the root

microbiome by specific bacterial taxa. *Science* **349**, 860–864 (2015).

25. Zhou, M.-L. *et al.* Nicotianamine synthase gene family as central components in heavy

metal and phytohormone response in maize. *Funct. Integr. Genomics* **13**, 229–239 (2013).

26. Lee, D.-K. *et al.* The rice OsNAC6 transcription factor orchestrates multiple molecular

mechanisms involving root structural adaptions and nicotianamine biosynthesis for drought

tolerance. *Plant Biotechnol. J.* **15**, 754–764 (2017).

27. Chaumont, F., Barrieu, F., Wojcik, E., Chrispeels, M. J. & Jung, R. Aquaporins constitute a

large and highly divergent protein family in maize. *Plant Physiol.* **125**, 1206–1215 (2001).

28. de Vries, F., Lau, J., Hawkes, C. & Semchenko, M. Plant–soil feedback under drought:

does history shape the future? *Trends Ecol. Evol.* **38**, 708–718 (2023).

29. Evans, S., Allison, S. & Hawkes, C. Microbes, memory and moisture: Predicting microbial

moisture responses and their impact on carbon cycling. *Funct. Ecol.* (2022)

doi:10.1111/1365-2435.14034.

1 **Supplementary Materials for**
2
3 **Persistent legacy effects on soil microbiota facilitate plant adaptive responses to drought**
4
5
6 Nichole A. Ginnan, Valéria Custódio, David Gopaulchan, Natalie Ford, Isai Salas-González, Dylan
7 H. Jones, Darren M. Wells, Ângela Moreno, Gabriel Castrillo, Maggie R. Wagner.
8
9
10
11

12 **This PDF file includes:**

Materials and Methods

No statistical methods were applied to predetermine sample size. The experiments were randomized, and investigators were blinded to allocation during experiments and outcome assessment.

1.   Legacy phase: Characterization of soils across precipitation gradients

1.1   Region selection and soil collection

Soils exposed to the macrogradient of precipitation were sampled from six never-irrigated native prairies across Kansas, USA in October 2020. This selection includes eastern Kansas tallgrass prairies: Welda Prairie (WEL), Clinton Wildlife Reserve (CWR), and Konza Wildlife Reserve (KNZ), as well as western Kansas shortgrass prairies: Hays Prairie (HAY), Smoky Valley Ranch (SVR), and Kansas State University's Tribune Southwest Research Center (TRI). The GPS coordinates of each collection site are available in the Supplemental Table S1. For soil collection, each site was split into three subplots. In each subplot, the surface soil (≈10 cm) containing thick plant root masses was removed with a bleach-sterilized metal shovel. Then, approximately 2.5 L of soil was collected from each site and pooled into a clean plastic bag, for a total of ≈7.5 L of soil collected per geographical location. Soil was held at room temperature for transport back to the laboratory where it was then stored at 4°C until use (~ 1 month).

Soils exposed to the microgradient of precipitation were selected from nine regions in Santiago Island, Cabo Verde in March 2022. These regions included Porto Gouveia, Mato Correia, Colonato, Ribeira Riba, Ribeira de São Domingos (Nora), Tomba Toiro, Serra Malagueta, Achada Leitão and Pico de Antónia. GPS coordinates are available in the Supplemental Table S1. For the soil collection, each site was divided into six equal parts (subplots). Within each subplot, six samples were collected following a typical zig-zag pattern with approximately 3 m of separation between the harvesting points and avoiding field edges, roads, and pathways. In each collection

point, the top 2 cm layer of soil was discarded using a handheld soil auger and pooled to produce

a unique composite sample per subplot. Thus, for each region, we collected six composite

samples. All tools used for the soil collection were previously disinfected with 70% ethanol, and

the operators wore gloves the whole time. Soil samples were transported in clean plastic bags to

the Instituto Nacional de Investigação e Desenvolvimento Agrário (INIDA).

For both Kansas and Cabo Verde soils, sub-samples for downstream metagenomic sequencing

and nutrient mineral content analyses were air-dried in sterile plastic trays at room temperature

for 1 week and then sieved using a 2 mm sieve to remove rocks, big soil particles, and vegetable

debris. The Kansas sub-samples were shipped to the University of Nottingham for further

processing, and all sub-samples were stored at 4 ºC until use.

1.2 Precipitation data collection

For all collection sites in Kansas and Santiago Island, precipitation values were retrieved from

publicly available sources with global coverage. Briefly, for Kansas, daily precipitation from 1981

to 2021 was extracted from the NASA POWER database based on the latitude and longitude of

each site[1]. For Santiago Island, we used three different approaches to compile rainfall data. First,

we compiled data from the available literature[2–4]. Briefly, the files containing the precipitation data

were downloaded directly from the supplemental material of each publication or were kindly

provided by the authors of the publications. Then, we extracted the rainfall data from stations in

Colonato, Ribeira de São Domingos, and Serra Malagueta. Next, for Porto Gouveia, Ribeira Riba,

Achada Leitão, and Pico de Antónia, the methodology of the World Meteorological Organization

was applied, which advises that on a small scale, we can consider reliable data collected from

stations located 10 km or less from the study region[5]. Finally, for Mato Correia and Tomba Toiro

we extracted the data from NASA POWER database based on the latitude and longitude[1]. The

dataset used in each case included the monthly precipitation average data from 1981 to 2021 for

Kansas and from 1981 to 2009 for Santiago.

1.3 Soil elemental content analysis

The soil mineral nutrients and trace elements profiles were determined using inductively coupled

plasma mass spectrometry (ICP-MS). The soil samples were dried using plastic weighing boats

in the fume hood for approximately 72 h at ambient temperature. Five grams of soil were weighed

in 50 mL conical tubes with a four-decimal balance, and treated with 20 mL of 1 M NH_4HCO_3, 5

mM diamine-triamine-penta-acetic acid (DTPA), and 5 mL 18.2 MΩcm Milli-Q Direct water (Merck

Millipore), for 1 h at 150 r.p.m. in a rotary shaker (adapted from[6]) to extract available elements.

Each treated sample was gravity-filtered through a quantitative filter paper (Whatman 42-

WHA1442070) to obtain approximately 5 mL of filtrate. Prior to the digestion, 20 µg/L of Indium

(In) was added to the nitric acid Primar Plus (Fisher Chemicals) as an internal standard for

assessing error in dilution, variations in sample introduction and plasma stability in the ICP-MS

instrument. Next, 0.5 mL of the soil filtrates were open-air digested in glass Pyrex tubes using 1

mL of concentrated trace metal grade nitric acid spiked indium internal standard for 2 h at 115 ºC

in dry block heater (DigiPREP MS, SCP Science; QMX Laboratories, Essex, UK). After cooling,

the digests were diluted to 10 mL with 18.2 MΩcm Milli-Q Direct water and elemental analysis

was performed using an ICP-MS, PerkinElmer NexION 2000 equipped with Elemental Scientific

Inc 4DXX FAST Dual Rinse autosampler, FAST valve and peristaltic pump. The instrument was

fitted with PFA-ST3 MicroFlow nebulizer, baffled cyclonic C3 high sensitivity glass spray chamber

cooled to 2ºC with PC3X Peltier heated/cooled inlet system, 2.0 mm i.d. quartz injector torch and

a set of nickel cones. Twenty-three elements were monitored including following stable isotopes:

^{7}Li, ^{11}B, ^{23}Na, ^{24}Mg, ^{31}P, ^{34}S, ^{39}K, ^{43}Ca, ^{52}Cr, ^{55}Mn, ^{56}Fe, ^{59}Co, ^{60}Ni, ^{63}Cu, ^{66}Zn, ^{75}As, ^{82}Se, ^{85}Rb,

^{88}Sr, ^{98}Mo, ^{111}Cd, ^{208}Pb and ^{115}In. Helium was used as a collision gas in Kinetic Energy

Discrimination mode (KED) at a flow rate of 4.5 mL/min while measuring Na, Mg, P, S, K, Ca, Cr,

Mn, Fe, Ni, Cu, Zn, As, Se and Pb to exclude possible polyatomic interferences. The remaining

elements were measured in the standard mode. Any isobaric interferences were automatically

corrected by the instrument Syngistix™ software for ICP-MS v.2.3 (Perkin Elmer). The ICP-MS

measurements were performed in peak hopping scan mode with dwell times ranging from 25 to

50 ms depending on the element, 20 sweeps per reading and three replicates. The ICP-MS

conditions were as follows: RF power – 1600 Watts, auxiliary gas flow rate 1.20 L/min. Torch

alignment, nebulizer gas flow and quadrupole ion deflector (QID) voltages (in standard and KED

mode) were optimized before analysis for highest intensities and lowest interferences (oxides and

doubly charged ions levels lower than 2.5%) with NexION Setup Solution containing 1 µg/L of

Be, Ce, Fe, In, Li, Mg, Pb and U in 1% nitric acid using a standard built-in software procedure. To

correct for variation between and within ICP-MS analysis runs, liquid reference material was

prepared using pooled digested samples, and run after the instrument calibration, and then after

every nine samples in all ICP-MS sample sets. Equipment calibration was performed at the

beginning of each analytical run using seven multi-element calibration standards (containing 2

µg/L In internal standard) prepared by diluting 1000 mg/L single element standards solutions

(Inorganic Ventures; Essex Scientific Laboratory Supplies Ltd) with 10% nitric acid. As a

calibration blank, 10% nitric acid containing 2 µg/L In internal standard was used and it was run

throughout the course of the analysis.

Sample concentrations were calculated using the external calibration method within the

instrument software. Further data processing, including the calculation of final elements

concentrations was performed in Microsoft Excel. First, sample sets run at different times were

connected as an extension of the single-run drift correction. Linear interpolation between each

pair of liquid reference material standards was used to generate a theoretical standard for each

sample that was then used to correct the drift by simple proportion to the first liquid reference

material standard analysed in the first run. Liquid reference material composed of pooled samples

was used instead of the CRM to match the chemical matrix of the samples as closely as possible,

thereby emulating the sample drift. Second, the blank concentrations were subtracted from the

sample concentrations and each final element concentration was obtained by multiplying by the

dilution factor and normalizing the element concentrations to the sample's dry weight.

1.4 Soil porosity analysis

To quantify soil porosity, we used X-ray computed tomography. First, soils were collected from

the selected locations in Kansas (October 2020) and Santiago (March 2022), according to

material and methods 1.1. Briefly, each site was divided into three and six subplots in the case of

Kansas and Santiago, respectively. Within each subplot the top layer of soil was removed (10 cm

in Kansas and 2 cm in Santiago), and the soil samples were collected using polyvinylchloride

(PVC) columns of 5 cm internal diameter and 7 cm (Kansas) or 5 cm (Santiago) length. After the

soil collection, the bottom of each column was sealed with tape to retain the soil in the column.

We collected six soil columns per location.

The undisturbed soil columns were non-destructively imaged using Phoenix v|tome|x MDT

(Waygate Technologies GmbH, Wunstorf, Germany) at The Hounsfield Facility, University of

Nottingham. Scans were acquired by collecting 2695 projection images at 180 kV X-ray energy,

200 µA current and 334 ms detector exposure time in fast mode (15 min total scan time per

column). Scan resolution was 55 µm for Kansas columns and 40 µm for Santiago columns.

1.5 Metagenomic analysis of free-living soil microbiota - DNA extraction

For metagenomic analysis, approximately 5 g of each soil sample was transferred into 50-mL

conical tubes containing 20 mL of sterile distilled water. To remove large plant debris and soil

particles, the samples were shaken thoroughly and filtered into new sterile 50-mL tubes using

100-µm nylon mesh cell strainers. The filtered soil solutions were centrifuged at high speed for

20 min in a centrifuge Eppendorf 5810R and most of the supernatants were discarded. The

remaining 1-2 mL of supernatants were used to dissolve the soil pellets. The resulting

suspensions were transferred to sterile 1.5-mL Eppendorf tubes. Samples were centrifuged again

at high speed in a benchtop centrifuge and the supernatants were discarded. The remaining

pellets were stored at - 80ºC for DNA extraction. For DNA isolation we used 96-well-format MoBio

PowerSoil Kit (MOBIO Laboratories; QIAGEN) following the manufacturer's instructions. Before

starting the extraction, all samples were manually randomized by placing them in a plastic bag

that was shaken several times. Samples were then taken individually from the bag and loaded in

the DNA extraction plates. This random distribution was maintained throughout library preparation

and sequencing.

1.6 Metagenomic library preparation and sequencing

DNA sequencing libraries were prepared using the Rapid PCR Barcoding Kit (SQK-RPB004) from

Oxford Nanopore Technologies, UK. In brief, 1 µL Fragmentation Mix (FRM) was added to 3 µL

DNA (2-10 ng/µL), and the reaction was mixed by gently finger-flicking. The DNA was fragmented

using the following conditions: 30°C for 1 min, then 80°C for 1 min in an Applied Biosystems Veriti

96-Well Thermal Cycler (Applied Biosystems, CA, USA). The fragmented DNA was cooled, then

barcoded and amplified in a PCR reaction containing 20 µL nuclease-free water, 25 µL LongAmp

Taq 2X master mix (New England Biolabs, MA, USA), 4 µL fragmented DNA, and 1 µL barcode

adaptor. The reaction was gently mixed and amplified using the following conditions: 95°C for 3

min, 20 cycles of denaturation at 95°C for 15 s, annealing at 56°C for 15 s and extension at 65°C

for 6 min, and a final extension of 65°C for 6 min. The resulting DNA library was purified using

0.6X Agencourt AMPure XP beads (Beckman Coulter, CA, USA) and eluted in 10 µL 10 mM Tris-

HCl pH 8.0, 50 mM NaCl. The library concentration was determined using a Qubit 4 Fluorometer

with the Qubit dsDNA HS Assay Kit (Thermo Fisher Scientific, USA). Equimolar quantities of

individual barcoded sample libraries were pooled and the volume was adjusted to 10 µL using 10

mM Tris-HCl pH 8.0, 50 mM NaCl. Subsequently, 1 μL rapid sequencing adapter (RAP) was

added to the pooled library and the tube was incubated at room temperature for five minutes.

Then, 34 μL Sequencing Buffer, 25.5 μL Loading Beads, and 4.5 μL nuclease-free water were

added to the tube, and the contents were mixed gently. The prepared pooled library was added

to a verified and primed FLO-MIN106 R9.4.1 flow cell (Oxford Nanopore Technologies, UK) in a

MinION (Oxford Nanopore Technologies, UK) following the manufacturer's instructions. DNA

sequencing was conducted with default parameters using a MinIT (Oxford Nanopore

Technologies, UK) with MinKNOW v2.1.12 (Oxford Nanopore Technologies, UK). Fast5 files were

base-called with Guppy v4.0.15 using the template_r9.4.1_450bps_hac.jsn high accuracy model

(Oxford Nanopore Technologies, UK).

1.7 Metagenomic sequences processing

The initial dataset underwent demultiplexing, and primer and barcode sequences were trimmed

using qcat v1.1.0 (Oxford Nanopore Technologies Ltd., UK). Reads with ambiguous barcode

assignments were excluded from further analysis. The reads were filtered with NanoFilt v2.8.0[7]

to discard low-quality sequences (Q-score < 9) and sequences < 100 bp. We used the Kraken

v2.1.2 pipeline[8] to classify the whole metagenome shotgun sequencing reads. The reads were

classified using the Kraken 2 archaea, bacteria, viral, plasmid, human, UniVec_Core, protozoa,

and fungi reference database (k2_pluspf_20220607). To estimate relative abundances, the

Bracken v2.7 pipeline[9] was applied to the classification results. Subsequently, Pavian v1.0[10] was

used to extract abundance and taxonomic tables.

1.8 Data Analysis

1.8.1 Precipitation data analysis

To determine the differences in precipitation levels across regions, we compared the mean annual

precipitation in each region by fitting a linear model with the following formula:

Mean annual precipitation ~ Region

Differences between regions were indicated using the confidence letter display derived from

Tukey's *post hoc* test implemented in the package multcomp v.1.4.25[11]. We inspected the

normality and variance homogeneity (here and elsewhere) using Q-Q plots and the Levene test,

respectively. We visualized the results via a point range plot using the ggplot2 v3.5.1 R package[12].

1.8.2 Soil elemental content analysis

For the soil elemental profile, we created a matrix in which each cell contained the calculated

element concentration in one sample. Then, we applied a z-score transformation to each ion

across the samples in the matrix. Afterward, we applied a principal component analysis (PCA)

using the Euclidean distance between samples and the z-score matrix as input to compare the

elemental profiles of soils. Additionally, we estimated the variance explained by porosity,

precipitation, region, and the interaction between them by performing a PERMANOVA via the

function *adonis2* from the vegan v.2.6-4 R package[13]. The significant variables were visualized

via a stacked bar plot using the *ggplot2* v3.5.1 R package.

To visualize the mineral content in each region, the z-score matrix created above was

hierarchically clustered (method *ward.D*, function *hclust*), and we visualized the results using a

heatmap. The rows in the heatmap were ordered according to the dendrogram order obtained

from the clustered ions, and the regions were ordered according to the precipitation gradient (low

precipitation to high). The heatmap was coloured based on the z-score.

To explore the relationship between ion concentrations and the precipitation gradient, we

performed a correlation test using *cor* function from *stats* v.4.3.1 package in R[14] of the average z-

score measurement of each ion against the precipitation gradient. Afterward, we plotted the

correlation coefficient for each ion in a barplot.

1.8.3 Soil porosity analysis

The soil core images taken using x-ray computed tomography were analyzed using ImageJ

software[15]. First, XY slice projection images were filtered using a median function to remove any

noise from the raw data then an automatic threshold (Li method) was applied to produce binary

images. In the binary images pores and solid particles were represented by black and white pixels,

respectively. Afterwards, a region of interest (ROI) was defined in the central part of each

projection to remove any potential border effect, cropping to a 600x600 pixel area. From the ROI

defined in each image we extracted soil features, including particle area, perimeter, circularity,

roundness, solidity, compactness, percentage of pores and pore size using the *measurement*

function.

To remove the variability in the topsoil due to transportation and handling, we used two strategies.

Firstly, we plotted the pore average size (mm) and soil porosity (%) for each sample. Then, we

excluded from the analysis all projections having a value of soil porosity > 40% and considered

the topsoil of those samples as the first projections after the exclusion. The second strategy was

applied to samples with an irregular shape (e.g., mountain-like shape) at the topsoil level. We

discarded all the projections with an irregular shape until we found projections with a regular

distribution of soil layers. For each collection of soils defined in Kansas and Santiago, we created

a data frame with the projection number (or slice) and the soil depth (projection number multiplied

by the resolution). The soil porosity for each soil type was visualized via a point plot using ggplot2

v3.5.1 R package[12].

To compare the soil elemental profiles against soil porosity across the samples in Kansas and

Santiago, first, we applied a z-score transformation of each ion and soil porosity across the

samples. Then, we estimated the distance between samples using Euclidean distance.

Afterwards, we contrasted the dissimilarity matrices of each pair of datasets (soil elemental profile

vs. soil porosity) using the *mantel* test implemented in the vegan v.2.6-4 R package[13]. Finally, we

computed the significance of the correlation between matrices by permuting the matrices 10,000

times[11].

1.8.4 Taxonomic data analysis

To compare the alpha diversity across regions, we calculated the Shannon diversity index using

the diversity function from the vegan v.2.6-4 package in R[13]. We used an ANOVA to test the alpha

diversity differences between regions. Differences between regions were indicated using the

confidence letter display derived from Tukey's *post hoc* test implemented in the R package

multcomp v.1.4.25[11].

The beta diversity analysis (Principal Coordinates Analysis, PCo) was based on Bray-Curtis

dissimilarity matrices calculated using the rarefied relative abundance tables. Additionally, we

estimated the variance explained by soil porosity, precipitation, regions, and the interaction

between them by performing a PERMANOVA via the function adonis2 from the vegan v.2.6-4 R

package[13]. The significant variables were visualized via a stacked bar plot using ggplot2 v3.5.1 R

package[12].

The relative abundance of bacterial phyla was depicted using a stacked bar plot using ggplot2

v3.5.1 package.

To compare the microbiome composition against the elemental profiles and soil porosity, we

contrasted the dissimilarity matrices of each pair of datasets (soil microbiome vs. soil elemental

profile and soil microbiome vs. soil porosity) using the *mantel* test implemented in the vegan v.2.6-

4 R package[13]. Briefly, we calculate the microbiome dissimilarity matrix using Bray-Curtis

distance. Next, we applied a z-score transformation of each ion and soil porosity across the

samples. Afterward, we calculated the soil elemental dissimilarity matrix and soil porosity

dissimilarity matrix using Euclidean distance. Finally, we used the *mantel* test to compare and test

the significance of the correlation between matrices by permuting the matrices 10,000 times[11].

1.8.5 Heatmap and enrichment analysis

We used the R package DESeq2 v.1.40.2[16] to compute the bacterial enrichment profiles in the

soils across the precipitation gradient. For each taxID (NCBI's taxonomic identifier assigned to

the taxa) in the rarefied table, as well as at the species, and family levels, we estimated difference

in abundance compared to the wettest region in Kansas and Santiago using a generalized linear

model (GLM) with the following design:

303   Abundance ~ Region

We extracted the following comparisons from the fitted model CWR vs WEL, HAY vs WEL, KNZ

vs WEL, SVR vs WEL, TRI vs WEL for Kansas samples and Achada Leitão vs Pico de Antónia,

Colonato vs Pico de Antónia, Mato Correia vs Pico de Antónia, Porto Gouveia vs Pico de Antónia,

Nora vs Pico de Antónia, Ribeira Riba vs Pico de Antónia, Serra Malagueta vs Pico de Antónia,

Tomba Toiro vs Pico de Antónia for Santiago samples. A taxID, species, or family was considered

statistically significant if it had a *p*-value < 0.05. We visualized the results for the GLM using a

heatmap. The rows in the heatmap were ordering according to the dendrogram obtained from the

tax ID, species and family analysis. The relative abundance matrix was standardized across the

significant tax ID, species, and family by using the z-score and the heatmap was coloured based

on this value.

1.8.6 Identification of marker taxa associated with precipitation gradients

To identify the corresponding bacterial isolates considered as "biomarker" taxa associated with

precipitation gradients, we identified the principal components (PCs) that explain more than 80%

of the variance in the data. These identified PCs were used to control the effects of soil elemental

profile on the taxa abundances. Then, we fit five models: Poisson, negative binomial, two zero-

inflated, and a multiple regression model.

For Poisson and negative binomial models, we fitted the following design:

Abundance ~ precipitation + offset (porosity) + offset (soil elemental profile: six first PCs for

324   Kansas and seven first PCs for Santiago)

326   In parallel, we fitted the zero-inflated models using the following design:

327   Poisson: Abundance ~ precipitation + offset (porosity) | 1 + offset (soil elemental profile) | 1

328   Negative binomial: Abundance ~ precipitation + porosity | porosity + soil elemental profile | soil

329   elemental profile

331   Next, to assess the statistical significance, we applied ANOVA to the best-performing model for

332   each taxon according to the Akaike Information Criterion (AIC). Additionally, we applied a multiple

333   regression model with the following design:

334   Abundance ~ precipitation + porosity + soil elemental profile

Then, we applied ANOVA to find taxIDs with a significant partial regression coefficient for

precipitation.

A taxa ID was considered a "marker" if it had a relative abundance > 0.01 and a prevalence >

20%. We visualized the average standardized relative abundance (z-score) of the significant taxa

ID in a point plot using ggplot2 v3.5.1 package[12].

1.8.7 Enrichment of bacteria biological functions associated with precipitation gradients

To identify biological processes enriched within the microbial communities, the sequence reads

were assembled into contigs for each sample using metaFlye from the Flye v2.9 package[17] with

default mode. The contigs generated were then grouped and deduplicated using the dedupe.sh

tool in BBTools v38.76[18] to eliminate redundancies. Next, we determined the relative abundance

of the contigs by mapping the reads from the samples to the contigs using minimap2 v2.17[19], and

extracting the relative abundance counts using CoverM v0.6.1[20] in the 'contig' mode and

reads_per_base coverage method. Taxonomic classification of the contigs was performed using

the CAT v8.22 taxonomic classification pipeline[21]. Subsequently, the contigs were filtered to retain

only bacteria sequences. DESeq2 v1.40.0[16] was used to determine the contig enrichment profiles

by fitting a GLM with the following design:

354   Abundance ~ Legacy + Biological Replicate

The low-precipitation soil versus the high-precipitation soil contrast was extracted from the fitted

model. Contigs meeting the criteria of an FDR-adjusted *p*-value (*q*-value) <0.05 and a $\log_2$-

transformed fold change >2 were selected for further analysis. Open reading frames encoded

within the contigs were predicted using FragGeneScanRs v1.1.0[22] with default settings. This was

followed by functional annotation of the predicted proteins using the eggNOG-mapper v2.1.9[23]

pipeline with the eggNOG v5.0.2 database[24] with Diamond v2.0.11[25] and MMseqs2[26].The genes

annotated with Gene Ontology (GO) classifications were subsequently extracted, and a GO

enrichment analysis focusing on biological processes was conducted. This involved employing

adaptive GO clustering in conjunction with Mann–Whitney U testing, using the GO_MWU tool[27],

which evaluates the enrichment of each GO category based on whether genes linked to the GO

category are significantly clustered at either the top or bottom of a globally ranked gene list. First,

genes were ranked based on the signed $\log_2$-transformed fold change values. For each gene,

any missing parental terms for specific GO categories were then automatically added. Next, fully

redundant categories (those containing identical sets of genes), were collapsed into the more

specific GO term. To further streamline the analysis, highly similar categories were grouped using

complete linkage clustering based on the fraction of shared genes. We used default settings,

where GO categories were merged if the most dissimilar pair within a group, shared more than

75% of the genes in the smaller category. The merged group was named after the largest

category. Significantly enriched and depleted GO categories were then determined by an adjusted

p-value of <0.05. This approach simplified the GO hierarchy and addressed multiple testing which

improved the statistical power of the GO enrichment analysis. The most prominent enriched and

depleted GO categories shared across comparisons were visualized in ggplot2 v3.4.2 and

coloured based on the square root transformed delta rank values (enrichment score) of the GO

categories. GO enrichment analysis was repeated on the Santiago soil samples using the same

methodology, and comparisons between the following geographical regions were extracted: Serra

Malagueta vs Pico de Antonia, Tomba Toiro vs Pico de Antonia, Mato Correia vs Pico de Antonia,

Colonato vs Pico de Antonia, Ribeira de Sao Domingos (Nora) vs Pico de Antonia, Porto Gouveia

vs Pico de Antonia, Ribeira Riba vs Pico de Antonia.

1.8.8 Analysis of genetic variation among bacterial lineages along the precipitation gradient

To assess genetic differences between bacteria lineages along the precipitation gradient, we

focused on 22 of the identified bacterial markers (1:Pseudomonas ID287, 2:Salmonella ID28901,

3:Rubrobacter ID49319, 4:Sorangium ID56, 6:Bradyrhizobium ID722472, 7:Luteitalea

ID1855912, 9:Bradyrhizobium ID1355477, 10:Flavisolibacter ID661481, 11:Bradyrhizobium

ID858422, 13:Rubrobacter ID2653851, 16:Bradyrhizobium ID1437360, 17:Candidatus Koribacter

ID658062, 18:Streptomyces ID1916, 19:Klebsiella ID573, 20:Bradyrhizobium ID1325107,

21:Edaphobacter ID2703788, 24:Flavisolibacter ID1492898, 27:Rhizobium ID384,

28:Rubrobacter ID2653852, 29:Microvirga ID2807101, 30:Archangium ID83451, 33:Nitrosospira

ID1231), as well as 11 additional abundant and prevalent species (5:Bacillus ID1428,

8:Bradyrhizobium ID1274631, 12:Priestia ID1404, 14:Lacibacter ID2760713, 15:Bradyrhizobium

ID1325100, 22:Candidatus Solibacter ID332163, 23:Burkholderia ID28450, 25:Bacillus ID1396,

26:Escherichia ID562, 31:Pseudomonas ID303, 32:Paenibacillus ID1464) across the samples, as

proxies for the broader bacterial communities. Together, these taxa were selected for their high

genome coverage across samples, enabling more precise allele frequency estimates. Reference

genomes for each bacterium was retrieved from the NCBI Genome database, and the filtered

shotgun metagenomic reads were aligned to these genomes using minimap2 v2.17-r941[19].

Alignments were sorted and indexed with SAMtools v1.18[28], followed by variant calling using

BCFtools v1.18[29]. This process identified 32,358,569 sequence variants, which were then filtered

using VCFtools v0.1.16[30], to retain only biallelic single nucleotide polymorphisms (SNPs) for

further analysis. SNP filtering criteria included a variant quality score >20, a minor allele frequency

(MAF) >0.01, less than 50% missing data, and a minimum sequencing depth of 10x in each

sample. After filtering, 15,365 high-quality biallelic SNPs were retained. Genetic distances were

computed using PLINK v1.90p[31], and reduced to two dimensions through classical

multidimensional scaling with the stats v4.3.0 package. PCoA plots were created with ggplot2

v3.4.2[12], colored by soil type, precipitation levels and geographical region.

To identify genes potentially under selection across the precipitation gradient, we conducted a

genetic-environment association (GEA) analysis. For this, SNPs were re-filtered using VCFtools

v0.1.16[30] with the same criteria, but allowing <70% missing data and a minimum sequencing

depth of 3x per sample. After filtering, 885,990 biallelic SNPs were retained. Subsequently, GEA

analysis was performed using a general linear model in the rMVP v1.1.1 package[32], with native

SNP data imputation, and average precipitation at each sampling location as the environmental

variable. The genetic structure in the data was corrected using the first 10 principal components

(PCs). Manhattan plots were generated using CMplot v4.5.1[33] and significant associations were

identified using the permutation method within the rMVP package.

2.   Soil drought legacy is resilient to short-term perturbations

2.1 Experimental design

The six soils collected from across the Kansas precipitation gradient, as described in materials

and methods 1.1, were used in this experiment. The conditioning perturbations imposed in these

experiments took place over approximately 20 weeks at the University of Kansas from December

17th, 2020 to May 5th, 2021. Each of the six input soils remained independent throughout the

experiment. Mesocosms consisted of a 1:1 (v/v) mixture of field-collected soil to sterile turface

MVP (Turface Athletics, Buffalo Grove, IL). A total of 192 sterile 100 mL pots were filled with the

six soils and which were then randomly assigned to one of four conditions in a fully-factorial

design: with or without a host, and either water-stressed or well-watered. Half the pots were

planted with seedlings of the native prairie grass *T. dactyloides* (Eastern gamagrass, cultivar

"Pete"); the rest remained unplanted. Thus, the replication was *N*=8 per soil per treatment. All

mesocosms were allowed to adapt to their watering regimes in a growth chamber set to a 12-hour

day cycle, 27°C/23°C, and ambient humidity. Well-watered control pots were watered every 1-2

days and water-stressed plants were watered every 3-5 days when plants displayed drought

symptoms (e.g., leaf curling). All pots were fertilized with 35 mL of 1mL/L concentration of Bonide

10-10-10 plant food (Bonide Products LLC, Oriskany, NY) on week 8 and week 12.

2.2 Sample collection

To characterize the baseline microbial communities going into the conditioning treatments, we

sampled four replicates of each soil/treatment combination one week after beginning the

experiment. To collect the samples, the top centimeter of soil was discarded and the remaining

soil was homogenized to ensure even sampling of the top, middle, and bottom of the pot. Two

grams of this homogenized soil were placed in a 15mL tube, flash-frozen in liquid nitrogen, and

stored at -80°C for microbial DNA and RNA extraction. To characterize the effects of the four

treatments on the microbial communities, we sampled all remaining replicates at the end of the

20-week conditioning phase. Soil samples were collected as described for the baseline

communities, but an additional 6 g of homogenized soil was preserved at 4°C in 50 mL conical

tubes for use as inocula in a downstream experiment. Additionally, for the planted pots, we

measured *T. dactyloides* shoot height before uprooting the plants. We collected samples of a

crown root from each plant (3 cm long each, beginning 2 cm from the base of the plant) and stored

them in 50% ethanol at 4°C for downstream laser ablation tomography analysis. The remaining

roots and shoots were dried in an oven at 225°C for 12 hours and then weighed separately.

2.3 Changes in bacterial community structure associated with drought and well-watered

conditioning with and without a host.

2.3.1 DNA extraction

Total DNA was extracted from baseline and post-conditioning soil sub-samples using the DNA

Set for NucleoBond RNA Soil Kit (Macherey-Nagel, Düren, Germany) according to the

manufacturer's instructions.

2.3.2 Library preparation and sequencing

DNA sequencing libraries were prepared using the Rapid PCR Barcoding Kit (SQK-RPB004) from

Oxford Nanopore Technologies, UK, and sequenced on a FLO-MIN106 R9.4.1 flow cell (Oxford

Nanopore Technologies, UK) in a MinION (Oxford Nanopore Technologies, UK), with a MinIT

(Oxford Nanopore Technologies, UK) using MinKNOW v2.1.12[34] (Oxford Nanopore

Technologies, UK) as described in materials and methods section 1.6.

2.3.3 Sequence processing

Raw sequence data were demultiplexed and primer and barcode sequences were trimmed using

qcat v1.1.0 (Oxford Nanopore Technologies Ltd., UK). Reads with ambiguous barcode

assignments were excluded from further analysis. The reads were filtered with NanoFilt v2.8.0[7]

to discard low-quality sequences (Q-score <9) and sequences <100 bp. We used the Kraken

v2.1.2 pipeline[8] for classifying the whole metagenome shotgun sequencing reads. The reads were

classified using the Kraken 2 archaea, bacteria, viral, plasmid, human, UniVec_Core, protozoa

and fungi reference database (k2_pluspf_20220607). To estimate relative abundances, the

Bracken v2.7 pipeline[9] was applied to the classification results. Subsequently, Pavian v1.0[10]

facilitated the extraction of abundance and taxonomic tables. Functions in phyloseq v1.44.0[35] with

microbiome v1.22.0 and microbiomeutilities v1.0.17[36] were used to filter the dataset and remove

samples with low read depth (<1000 reads), remove unidentified taxa and singletons, transform

abundance values using rarefaction, subset and merge sample and taxonomic groups and

perform other data frame manipulations.

2.4 Plant biomass

Root and shoots were detached, dried in an oven at 225˚C for 12 hours, and then weighed.

2.5 Root Laser Ablation Tomography (LAT) analysis

We collected samples of a crown root from each plant (3 cm long each, beginning 2 cm from the

base of the plant) and stored them in 50% ethanol at 4°C. The samples were shipped to the

University of Nottingham for downstream LAT analysis. Briefly, root segments were dehydrated

in 100% methanol for 48 hours, transferred to 100% ethanol for 48 hours, then dried with an

automated critical point dryer (CPD, Leica EM CPD 300, Leica Microsystem). Root anatomical

images were acquired using a laser ablation tomograph (LATScan, Lasers for Innovative

Solutions LLC). This utilises a combination of precise positioning stages with a guided pulsed UV

(355 nm) laser, to thermally vaporise thin sections of the root, and then to illuminate the exposed

surface. The tomograph was retrofitted with a microscopic imaging system, using a machine

vision camera unit (Model Grasshopper3, FLIR) and infinity-corrected long working distance

magnifying objectives (Mitutoyo (UK) Ltd.)

2.6 Changes in gamagrass morphological features under drought and well-water conditioning

To identify morphological features in gamagrass that changed with the conditioning (drought and

well-watered) treatments, we used Image J to quantify the average area aerenchyma, number

aerenchyma, number metaxylem, cortical cell layers, total area metaxylem, average area

metaxylem, stele min. diameter, stele max. diameter, stele area, stele perimeter, total area

aerenchyma, adjusted cortex area, root min. diameter, cortex min. diameter, root max. diameter,

cortex max. diameter, total perimeter, cortex perimeter, root total area, and cortex area.

Additionally, we quantified the root-shoot ratio, number of leaves, number of green leaves, root

mass, shoot mass, and shoot height.

2.7 Metatranscriptome analysis

2.7.1 RNA isolation

Total RNA was extracted from baseline and post-conditioning soil sub-samples with the

NucleoBond RNA Soil Kit (Macherey-Nagel, Düren, Germany) using the manufacturer's

instructions. Isolated RNA was treated with Turbo DNA-free (Applied Biosystems, Waltham, MA,

USA) to remove contaminating DNA, following the manufacturer's instructions.

2.7.2 Library preparation and sequencing

RNA libraries were prepared using 1 µg of total RNA according to established protocols with

modifications. Briefly, poly(A)-tail-containing RNA was removed from the RNA samples using

Sera-mag oligo(dT) magnetic beads (GE Healthcare Life Sciences, Marlborough, MA, USA) and

then the samples were subjected to ribodepletion with the NEBNext rRNA Depletion Kit (New

England Biolabs, MA, USA), following the manufacturers' instructions. The purified RNA was

resuspended in a fragmentation mix consisting of 6.25 µL Milli-Q water, 5 µL 5x First strand buffer,

and 1.25 µL random primers (3 µg/µL) and fragmented at 94°C for 6 min. First-strand cDNA

synthesis was performed using a mixture of 0.8 µL reverse transcriptase, 2 µL 100 mM DTT, 0.4

µL 25 mM dNTP, 0.5 µL RNAseOUT (40U/µL), 10 µL RNA, and 6.3 µL Milli-Q water. The reactions

were incubated at 25°C for 10 min, 42°C for 50 min, and 70°C for 15 min. Second-strand cDNA

synthesis was performed by adding a master mix of 18.4 µL Milli-Q water, 5 µL 10X second strand

buffer, 1.2 µL 25 mM dNTP, 0.4 µL RNAse H (5U/µL), and 5 µL DNA Pol I (10U/µL) to the sample,

followed by incubation at 16°C for 1 h. The samples were then purified using Agencourt AMPure

XP beads. Subsequently, the libraries were end-repaired with a mixture of 30 µL sample, 2.5 µL

of 3 U/µL T4 DNA polymerase, 0.5 µL of 5 U/µL Klenow DNA polymerase, 2.5 µL of 10 U/µL T4

PNK, 5 µL of 10X T4 DNA ligase buffer with 10 mM ATP, 0.8 µL of 25 mM dNTP mix, and 8.7 µL

Milli-Q water, incubated at 20°C for 30 min, and purified again using Agencourt AMPure XP

beads. Following this, the RNA libraries were adenylated in a mix containing 34 µL of the end-

repaired sample, 3 µL of 5 U/µL Klenow exo-, 5 µL of 10X Enzymatics Blue Buffer, 1 µL of 10 mM

dATP, and 9 µL of Milli-Q water. The mixture was incubated at 37°C for 30 min, followed by 70°C

for 5 min, and then purified using Agencourt AMPure XP beads. Individual samples were indexed

through ligation using a mix of 10.25 µL sample, 1 µL of 600 U/µL T4 DNA ligase, 12.5 µL of 2x

Rapid Ligation Buffer, and 1.25 µL of 2.5 µM indexing adapter from the KAPA Dual-Indexed

Adapter Kit (Kapa Biosystems, MA, USA). The samples were incubated at 25°C for 15 min,

followed by the addition of 5 µL of 0.5 M EDTA pH 8. The libraries were purified twice with

Agencourt AMPure XP beads. The libraries were enriched in a reaction containing 20 µL sample,

25 µL of 2X KAPA HiFi HS Mix (Kapa Biosystems, MA, USA), 2.5 µL of 5 µM I5 primer, and 2.5

µL of 5 µM I7 primer. The reactions were initially heated to 98°C for 45 seconds, followed by 14

cycles of 98°C for 15 seconds, 60°C for 30 seconds, and 72°C for 30 seconds, with a final

extension at 72°C for 1 minute. The resulting RNA libraries were purified using Agencourt AMPure

XP beads, quantified on a Qubit 4 Fluorometer (Thermo Fisher Scientific, USA), and the library

size was assessed using High Sensitivity D1000 ScreenTape on the Agilent 4200 TapeStation

(Agilent Technologies, Santa Clara, CA). Equimolar quantities of individual barcoded RNA

libraries were pooled in a randomized manner and shipped on dry ice to Beijing Genomics Institute

(BGI, Shenzhen, China). Each library pool was sequenced on an MGI Tech MGISEQ-2000

sequencing platform to generate a minimum of 10 million 100 bp paired-end reads per sample.

2.7.3 Taxonomic classification of transcripts

Cutadapt v4.6[37] was used to remove primer and barcode sequences and low-quality sequences

from the paired-end reads of the sequenced RNA libraries. To identify taxa with enriched gene

expression activity, the reads were classified using the Kraken v2.1.2 pipeline[8] with the archaea,

bacteria, viral, plasmid, human, UniVec_Core, protozoa and fungi reference database

(k2_pluspf_20220607), and the Bracken v2.7 pipeline[9] was applied to the classification results to

estimate the relative abundances. The counts table was generated from Pavian v1.0[10]. Data

filtering and statistical analysis were then performed as before using phyloseq v1.44.0[35] with

microbiome v1.22.0[38] and microbiomeutilities v1.0.17[36].

2.8 Data Analysis

2.8.1 Changes in bacterial community structure associated with drought and well-watered

conditioning with and without a host

To assess the alpha diversity across the samples, we calculated the Shannon Diversity Index

using phyloseq v1.44.0[35]. We used ANOVA to test for significant differences in Shannon Diversity

indices between groups and means were separated using Tukey's honestly significant difference

(HSD) test from the agricolae v1.3.5 R package[39]. For beta diversity, Bray-Curtis dissimilarity

matrices were calculated using phyloseq v1.44.0 and the variance explained by legacy,

conditioning, and host were estimated by performing permutational multivariate analysis of

variance (PERMANOVA) using the adonis2 function in vegan v2.6.4 R package[13]. Constrained

ordination of beta-diversity was plotted using canonical analysis of principal coordinates (CAP)

based on Bray-Curtis dissimilarity matrices calculated with vegan v2.6.4. We visualized

differences with the CAP analysis, using the following models:

579   ~ legacy + Condition (conditioning + host + biological replicate)

580   ~ conditioning + Condition (legacy + host + biological replicate)

581   ~ host + Condition (legacy + conditioning + biological replicate)

583   The relative abundance of taxa was plotted as a stacked bar representation using phyloseq

584   v1.44.0. The tax_glom function in phyloseq v1.44.0 was used to agglomerate taxa, and the

585   aggregate_rare function in microbiome v1.22.0 was used to aggregate rare groups. We used

586   DESeq2 v1.40.0[16] to calculate the enrichment profiles by fitting a generalized linear model (GLM)

587   with the following design:

588   Abundance ~ Legacy + Conditioning + Biological Replicate

We extracted the following comparisons from the fitted model: wet soil legacy with watered

conditioning vs wet soil legacy (baseline), wet soil legacy with drought conditioning vs wet soil

legacy (baseline), dry soil legacy (baseline) vs wet soil legacy (baseline), dry soil legacy with

watered conditioning vs wet soil legacy (baseline), and dry soil legacy with drought conditioning

vs wet soil legacy (baseline). Taxa were considered significant if they had an FDR-adjusted p-

value (q-value) <0.05. The results of the GLM analysis were rendered in heatmaps, coloured

based on the $\log_2$-transformed fold change output by the GLM. Significant differences between

comparisons with a q-value <0.05 with $\log_2$-transformed fold change >2 were highlighted with

black squares.

Relative abundances of the taxonomic markers were extracted, and an ANOVA was performed

to assess significant differences between treatment groups. Tukey's Honest Significant Difference

(HSD) test, implemented using the agricolae v1.3.5 R package[39], was used for post-hoc pairwise

comparisons. To further explore the effects of watering and drought treatments, with and without

a host, on the relative abundances of the taxonomic markers, we subset the data and applied a

generalized linear model (GLM) using DESeq2 v1.40.0[16]. The model was structured as follows:

606   Abundance ~ Legacy + Conditioning + Host + Biological Replicate

We then extracted the following comparisons from the fitted model for each soil legacy: water

conditioning vs baseline, drought conditioning vs baseline, water conditioning with host vs

baseline, drought conditioning with host vs baseline. Markers were considered significant if the

FDR-adjusted p-value (q-value) was < 0.05. Results from the GLM analysis were visualized in a

heatmap, with colours representing log2-transformed fold changes. Comparisons showing

significant differences (q-value <0.05 and log2-transformed fold change >2) were highlighted with

black squares.

2.8.2 Gene Ontology (GO) term enrichment analysis

To identify enriched biological processes within the microbial communities, the sequence reads

from individual samples were assembled into contigs, their relative abundance counts

determined, and were subjected to taxonomic classification as described before. DESeq2

v1.40.0[16] was utilized to determine the bacterial contig enrichment profiles by fitting a generalized

linear model (GLM) with the following design:

622   Abundance ~ Legacy + Conditioning + Biological Replica

We extracted the following comparisons from the fitted model: wet soil legacy with watered

conditioning vs wet soil legacy (baseline), wet soil legacy with drought conditioning vs wet soil

legacy (baseline), dry soil legacy (baseline) vs wet soil legacy (baseline), dry soil legacy with

watered conditioning vs wet soil legacy (baseline), and dry soil legacy with drought conditioning

vs wet soil legacy (baseline). Contigs meeting the criteria of an FDR-adjusted p-value (q-value) <

0.05 and a $\log_2$-transformed fold change > 2 were selected. Open reading frames were predicted

and functionally annotated, and genes with GO classifications were subjected to GO enrichment

analysis with the GO_MWU tool[27].

2.8.3 Changes in gamagrass morphological features under drought and well-water condition

For each root feature identified, we used ANOVA to test for significant differences between groups

and means were separated using Tukey's honestly significant difference (HSD) test from the

agricolae v1.3.5 R package[39] Subsequently, the feature values were normalized using the rescale

function from the scales v1.2.1 R package. The mean normalized feature values were then

visually represented on a heatmap using ggplot2 v3.4.2. Subsequently, Pearson correlation

coefficients between these features and corresponding p-values were computed using the rcorr

function from the Hmisc v5.0.1 package[40]. The results of the correlation analysis were graphically

presented using ggplot2 v3.4.2, where the colour of the plots reflected the correlation coefficient

values. Significant correlations (p <0.05) were emphasized with black squares on the plots.

Furthermore, the coefficient of variation for the feature values was calculated and depicted using

ggplot2 v3.4.2. The three plots were integrated based on the hierarchical clustering of the Pearson

correlation coefficients of the features. The clustering employed the ward.D2 method within the

hclust function in R, utilizing Euclidean distances calculated using the dist function.

2.8.4 Metatranscriptome sequence analysis

To assess transcriptional differences in the activity of the bacterial community, Bray-Curtis

dissimilarity matrices were calculated. The variance explained by legacy, conditioning, and host

was estimated using permutational multivariate analysis of variance (PERMANOVA) with the

adonis2 function from the vegan v2.6.4 R package[13]. Beta-diversity patterns were visualized

through constrained ordination using canonical analysis of principal coordinates (CAP). We

applied CAP analysis to visualize differences using the following models:

~ Legacy + Condition (conditioning + host + biological replicate)

~ Conditioning + Condition (legacy + host + biological replicate)

~ Host + Condition (legacy + conditioning + biological replicate)

Transcriptional activity among taxa was displayed as a stacked bar plot using phyloseq v1.44.0.

We employed DESeq2 v1.40.0[16] to calculate enrichment profiles by fitting a generalized linear

model (GLM) with the following design:

Abundance ~ Legacy + Conditioning + Biological Replicate

From the fitted model, we extracted the following comparisons: wet soil legacy with watered

conditioning vs wet soil legacy (baseline), wet soil legacy with drought conditioning vs wet soil

legacy (baseline), dry soil legacy (baseline) vs wet soil legacy (baseline), dry soil legacy with

watered conditioning vs wet soil legacy (baseline), and dry soil legacy with drought conditioning

vs wet soil legacy (baseline). Taxa were considered differentially abundant if the FDR-adjusted

p-value (*q*-value) was <0.05. Results from the GLM analysis were visualized in heatmaps, where

colours represent log2-transformed fold changes. Comparisons showing significant differences

(*q*-value <0.05 and log2-transformed fold change >2) were highlighted with black squares.

Additionally, high-quality filtered reads of the transcriptome were de novo assembled into a

reference metatranscriptome using Trinity v2.15.1[41] with default parameters. Open reading

frames in transcripts were predicted with TransDecoder v5.7.1 with default settings. Functional

annotation of the predicted proteins was performed using the eggNOG-mapper v2.1.9[23] pipeline,

utilising the eggNOG v5.0.2 database[24] with Diamond v2.0.11[25] and MMseqs2[26]. The taxonomic

classification of transcripts was conducted using the CAT v8.22 taxonomic classification

pipeline[21]. Sequence reads were further filtered using SortMeRNA v4.3.6[42] with the

smr_v4.3_default_db.fasta database to remove residual amplified rRNA sequences. Transcript

quantification analysis was performed using Salmon v1.10.0[43] in the mapping-based mode with

the de novo assembled reference metatranscriptome. Subsequently, the transcript-level

abundance estimates from salmon were extracted as transcripts per million (TPM) for the

identified transcripts using the R package tximport v1.28.0[44]. DESeq2 v1.40.0[16] was utilized to

determine the bacterial transcript enrichment profiles by fitting a generalised linear model (GLM)

as described before. Genes meeting the criteria of an FDR-adjusted p-value (q-value) <0.05, a

$\log_2$-transformed fold change >2, and had GO classifications, were subjected to GO enrichment

analysis with the GO_MWU tool[27].

2.8.5 Analysis of genetic variation among bacterial lineages

Filtered shotgun metagenomic reads were aligned to the reference genomes of 33 selected taxa,

including 22 identified bacterial markers and 11 additional abundant and prevalent species (see

Supplementary Materials and Methods 1.8.8), using Minimap2 v2.17-r941[19]. The resulting

alignments were sorted and indexed with SAMtools v1.18[28]. Variant calling was performed using

BCFtools v1.18[29], and variants were filtered with VCFtools v0.1.16[30]. Filtering criteria included a

variant quality score >20, a minor allele frequency (MAF) >0.01, <50% missing data, and a

minimum sequencing depth of 10x in each sample. After filtering, a total of 8,293 high-quality

biallelic SNPs were available for further analysis. To assess genetic variation within and between

groups, we analysed molecular variance (AMOVA) using poppr v2.9.6[45]. The significance of the

AMOVA results was determined with a permutation test using the randtest function in ade4

v1.7.22[46]. To determine the extent of genetic differentiation between groups, we calculated the

Fixation index (FST) values using the hierfstat v0.5.11[47] package. Principal coordinates analysis

(PCoA) plots were generated as previously described in Supplementary Materials and Methods

1.8.8, and coloured to reflect soil legacy, drought/ well-watered and host/ no-host treatments. To

identify genes associated with soil legacy, we conducted a genome-wide association study

(GWAS). SNPs were re-filtered using VCFtools v0.1.16[30], applying the same criteria but allowing

for up to 70% missing data and a minimum sequencing depth of 3x in each sample. GWAS was

conducted using a general linear model in the rMVP v1.1.1 package[32]. Associations were

identified by comparing bacterial lineages from dry legacy soil to those from wet legacy soil.

Genetic structure was accounted for by incorporating the first 10 principal components. Significant

associations were identified through permutation testing within the rMVP package, and Manhattan

plots were generated using CMplot v4.5.1[33].

3. Effects of soil microbiome legacy on plant tolerance to drought

3.1 Experimental design and non-destructive phenotypic measurements

At the end of the "Conditioning Phase", homogenized soil was collected from each pot by

discarding the top one centimeter of soil, mixing the soil in the pot with a clean plastic spatula,

and placing six grams in a sterile 50 mL conical tube. For rhizosphere samples (i.e., planted pots),

plants were gently pulled from the pots and the soil particles adhered to and within the root bundle

were shaken into a sterile 50 mL conical tube. The rhizosphere particles were homogenized with

a clean plastic spatula and particles were poured out of the tube until six grams remained. Soil

and rhizosphere samples were stored at 4˚C overnight.

Soil microbes were extracted from the 6 g soil or rhizosphere sample the following day by adding

25 mL of autoclaved 1X PBS with 0.0001% Tween-89 to the 50 mL tube containing the sample.

Tubes were vigorously shaken to mix and break up large soil aggregates. Large particles were

allowed to settle to the bottom of the tube; the supernatant was then filtered through autoclaved

Miracloth (Sigma-Aldrich) into a new sterile 50 mL conical tube. Filtered samples were then

centrifuged at 3600g for 25 min at 4°C. The supernatant was discarded, and the microbial pellet

was resuspended in 6 mL of 1X PBS buffer using a vortex. The resuspended pellets were stored

at 4°C until used for inoculations a few hours later.

The microbial extract from each of the 192 Conditioning Phase pots (as well as the 24

uninoculated control pots) was used to inoculate 4 pots for the "Test Phase": one pot per

combination of watering treatment (droughted or control) and host species (maize or gamagrass).

To create the inoculum, the resuspended pellet was inverted three times to mix and 1 mL of the

sample was added to 100 mL of sterile 0.5X MS liquid medium, for a microbial titer equivalent to

0.01 g soil per mL. The "mock inoculation" controls were created by substituting 1 mL of sterile

PBS for the resuspended microbial pellet. Finally, 25 mL of this suspension was inoculated onto

the soil surface of each Test Phase pot. Thus, each microbial community extracted from one of

the 192 conditioning phase pots was used to inoculate four plants in the Test Phase. To maintain

statistical independence of the experimental replicates from the conditioning phase, no pooling

was performed.

Before inoculation, pots were planted with 3-4-day old gamagrass or maize germinants.

Gamagrass seeds were soaked in 3% hydrogen peroxide for 24 hours and germinated in seed

trays filled with sterile clay. Maize seeds were soaked in 70% ethanol for 3 minutes, then soaked

in 5% NaClO on a rotator for 2 min, and then rinsed with sterile DI water three times. Treated

maize seeds were germinated on sterile damp paper towels inside sealed plastic bags.

Pots were fully randomized and, prior to inoculation, were filled with a homogenized 5:1 (w/w)

mixture of all-purpose sand (TechMix All-purpose 110241) and calcined clay (Pro's choice rapid

dry) that had been sterilized by autoclaving on a one-hour liquid cycle. Pots were autoclaved on

a 30 min liquid cycle and then filled with the sand:clay mixture, leaving one inch of room at the

top of the pot. To help keep the mixture from falling out of the drain holes in the bottom of the

pots, a sterile filter paper was shaped into a cone, pushed to the bottom of the pot, and a sterile

marble was used to weigh the paper down. This effectively blocked the substrate, but still allowed

water to exit the drainage holes. Plants were grown under 12-h days, 27°C/23°C (day/night), and

ambient humidity, with the light setting set to 1, which is equivalent to 312 μmol/m^{2}s. Three-day

old gamagrass and maize leaf photosynthetic rates and gas exchange were measured using the

LI-6800 (LI-COR, Lincoln, NE, USA), across 3 days for each host. The LI-6800 aperture was 2

cm, warmup tests were performed at the start of each measurement session, the Fluorometry

was set to "on", the APD_Leaf to set to 1.5 kpa. The newest fully emerged leaf, which was most

commonly the 4th leaf, on each plant was clamped in the chamber and allowed to stabilize until

all measurements were stable for at least 30 secs before the measurements were recorded, which

took approximately five minutes per leaf. Similarly, the leaf chlorophyll content was also measured

using the MC-100 Chlorophyll Concentration Meter (apogee instruments, Logan, UT, USA).

Maize plants were sampled four weeks after planting and gamagrass was sampled five

weeks after planting. In total, we measured 286 *T. dactyloides* plants (192 droughted, 94 well-

watered) and 268 maize plants (182 droughted, 86 well-watered).

At the end of the Test Phase, uprooted plants were gently shaken to remove the soil attached to

roots, prior to the collection of phenotypic, transcriptomic, and microbiome data as described

below. One crown root was cut off with a ceramic blade and placed in a 1.7 mL tube on dry ice

for downstream DNA extraction for *16S rRNA* gene sequencing. Another crown root (0.15 - 0.2

g) was cut off with a ceramic blade, placed in a 1.7 mL tube, and flash frozen in liquid nitrogen for

downstream RNA extraction. In between plants, the ceramic blade, plastic tweezers, plastic

cutting board, and gloves were cleaned with 30% bleach. All samples were held on dry ice and

then transferred to a -80˚C freezer for storage. Next, the root and shoot were separated with a

ceramic blade. Three cm of crown root beginning ~2 cm from the base of the shoot was cut with

a ceramic blade and submerged in 50% EtOH for LAT analysis. The rest of the root system was

submerged in 70% EtOH in a 50 mL centrifuge tube for downstream root architecture scanning.

Shoot height and number of leaves were recorded. Shoots were placed in individual paper bags,

dried in an oven at 225˚C for 12 hours, and the shoot dry weight was recorded.

3.2 Root system architecture and root biomass analyses

A Perfection V600 flatbed scanner (Epson, Nagano, Japan) was used to scan intact maize and

gamagrass root systems collected from Test Phase plants. The scanner was set to professional

mode, reflective, document, black and white mode, 600 dpi, with a threshold of 55. A clear plastic

tray filled with clean water was placed on the scanning bed. Each root system was placed in the

tray with water and the tangled roots were gently pulled apart using plastic tweezers until they

were no longer overlapping. A small amount of fine fibrous roots that fell off during this process

were pushed to the corner of the tray and not included in the root cluster scan. The scanned root

images were then analyzed using Rhizovision Explorer software[48] in "whole root" mode and

converted to 600 dpi. The region of interest tool was used to outline the main root bundle before

pressing play to collect feature measurements. Finally, after collection of root system architecture

data, the roots were dried in an oven at 225˚C for 12 hours and then weighed.

3.3 Root Laser Ablation Tomography (LAT) analysis

We collected one crown root from each plant (3 cm long each, beginning 2 cm from the base of

the plant) and stored them in 50% ethanol at 4°C. These samples were shipped to the University

of Nottingham for LAT analysis. Briefly, root segments were dried with an automated critical point

dryer (CPD, Leica EM CPD 300, Leica Microsystem). Then, samples were ablated by a laser

beam (Avia 7000, 355 nm pulsed laser) to vaporize the root tissue at the camera focal plane

ahead of an imaging stage and cross-sectional images were taken using a Canon T3i camera

with a 5×micro-lens (MP-E 65 mm) on the laser-illuminated surface. ImageJ software[15] was used

to measure root anatomical traits captured in the high-quality LAT images.

3.4 Leaf ionome and shoot biomass analyses

The elemental profiles of the shoots were measured using Inductively Coupled Plasma Mass

Spectrometry (ICP-MS). The shoot biomass from all uprooted plants was dried in an oven at

225°C for 12 h and then weighed. The dried biomass samples were cut into small pieces using a

clean ceramic scalpel and placed in 5 mL Eppendorf tubes with 3 zirconium oxide beads. Shoots

were pulverized using a Tissue Lyzer II (Qiagen) using 2 cycles of 60 seconds at the frequency

of 30 s^{-1}. Next, 5-10 mg of pulverized shoot samples were weighted on a Mettler five-decimal

analytical scale, and 1-3 mL (depending on the sample dry weight) of concentrated trace metal

grade nitric acid Primar Plus (Fisher Chemicals) was added to each tube. Prior to the digestion,

20 µg/L of Indium (In) was added to the nitric acid as an internal standard to assess putative errors

in dilution or variations in sample introduction and plasma stability in the ICP-MS instrument. The

samples were then digested in DigiPREP MS dry block heaters (SCP Science; QMX Laboratories)

for 4 h at 115°C. After cooling down, the digested samples were diluted to 10-30 mL (depending

on the volume of the nitric acid added) with 18.2 MΩcm Milli-Q Direct water. The elemental

analysis was performed using an ICP-MS, PerkinElmer NexION 2000 equipped with Elemental

Scientific Inc 4DXX FAST Dual Rinse autosampler, FAST valve and peristaltic pump. The

instrument was fitted with a PFA-ST3 MicroFlow nebulizer, baffled cyclonic C3 high sensitivity

glass spray chamber cooled to 2 ºC with PC3X Peltier heated/cooled inlet system, 2.0 mm i.d.

quartz injector torch and a set of nickel cones. Twenty-four elements were monitored including

the following stable isotopes: ^{7}Li, ^{11}B, ^{23}Na, ^{24}Mg, ^{31}P, ^{34}S, ^{39}K, ^{43}Ca, ^{48}Ti, ^{52}Cr, ^{55}Mn, ^{56}Fe, ^{59}Co,

^{60}Ni, ^{63}Cu, ^{66}Zn, ^{75}As, ^{82}Se, ^{85}Rb, ^{88}Sr, ^{98}Mo, ^{111}Cd, ^{208}Pb and ^{115}In. Helium was used as a collision

gas in Kinetic Energy Discrimination mode (KED) at a flow rate of 4.5 mL/min while measuring

Na, Mg, P, S, K, Ca, Ti, Cr, Mn, Fe, Ni, Cu, Zn, As, Se and Pb to exclude possible polyatomic

interferences.

The remaining elements were measured in the standard mode. The instrument Syngistix™

software for ICP-MS v.2.3 (Perkin Elmer) automatically corrected any isobaric interferences. The

ICP-MS measurements were performed in peak hopping scan mode with dwell times ranging

from 25 to 50 ms depending on the element, 20 sweeps per reading and three replicates. The

ICP-MS conditions were adjusted to an RF power of 1600 Watts and an auxiliary gas flow rate of

1.20 L/min. Torch alignment, nebuliser gas flow and quadrupole ion deflector (QID) voltages (in

standard and KED mode) were optimized before analysis for highest intensities and lowest

interferences (oxides and doubly charged ions levels lower than 2.5%) with NexION Setup

Solution containing 1 µg/L of Be, Ce, Fe, In, Li, Mg, Pb and U in 1% nitric acid using a standard

built-in software procedure. To correct for variation between and within ICP-MS analysis runs,

liquid reference material was prepared using pooled digested samples and run after the

instrument calibration and then after every nine samples in all ICP-MS sample sets. Equipment

calibration was performed at the beginning of each analytical run using seven multi-element

calibration standards (containing 2 µg/L In internal standard) prepared by diluting 1000 mg/L

single-element standards solutions (Inorganic Ventures; Essex Scientific Laboratory Supplies Ltd)

with 10% nitric acid. As a calibration blank, 10% nitric acid containing 2 µg/L In internal standard

was used, and it was run throughout the analysis. Sample concentrations were calculated using

the external calibration method within the instrument software. Further data processing, including

the calculation of final element concentrations, was performed in Microsoft Excel.

3.5 Crown root transcriptomics

3.5.1 RNA extraction and sequencing

For RNA extractions and sequencing, flash-frozen crown roots were freeze-dried for 48 hours and

finely ground with pellet pestles. The RNA extraction protocol was carried out according to the

NucleoSpin RNA Plant kit (Macherey-Nagel, Düren, Germany).

For the 132 maize samples, remnant DNA was removed from purified RNA using the DNA-Free

kit (Invitrogen, Carlsbad, CA, USA). RNA-seq libraries were prepared using the QuantSeq 3'

mRNA-Seq V2 kit with unique dual sequences and the unique molecular identifier (UMI) module

(Lexogen, Vienna, Austria) following the manufacturer's recommendations. Libraries were pooled

at equimolar concentrations and then sequenced (2x150bp, but reverse reads were not used) on

a NovaSeq S4 flow cell (Illumina, San Diego, CA, USA) along with a 25% PhiX spike-in. Maize

RNA-seq library preparations and sequencing were performed by the RTSF Genomics Core at

Michigan State University. For the 132 *T. dactyloides* samples, RNA-seq libraries were prepared

using the NEBNext Ultra II Directional Library Kit with the oligo-dT magnetic isolation module

(New England Biolabs, Ipswich, MA, USA) and sequenced on the Illumina NovaSeq 6000

platform at the Genomic Sciences Laboratory at North Carolina State University to generate a

minimum of 40M read pairs (2x150bp) per sample.

3.5.2 Sequence processing

For the maize sequence reads, UMIs were removed from all sequences and added to the read

headers using UMI-tools[49]. Next, cutadapt version 4.2[37] was used to remove the first four bases

of each read, remove poly-A tails (if present), remove spurious poly-G runs using the --nextseq-

trim=10 parameter, and remove adapter sequences. Reads that were <10 bp long or that aligned

to maize rRNA gene sequences were removed; the remaining reads were aligned to the maize

reference genome B73 RefGen_v5[50] using HISAT2 version 2.2.1[51]. Aligned reads were converted

to BAM format, sorted, and indexed using samtools version 1.9[28]. We then used UMI-tools[49] to

de-duplicate reads that both shared a UMI and had identical mapping coordinates. Finally, we

used the FeatureCounts function of the subread package version 2.0.5[52] with the maize genome

annotation version Zm00001eb.1 and parameters -O --fraction -M --primary -g ID -t gene to

generate a table of gene expression estimates.

For the *T. dactyloides* sequence reads, we first used cutadapt to remove NEBNext adapter

sequences, poly-A tails, spurious poly-G runs, and low-quality tails using the -q 20,20 parameter

and other default parameters. The cleaned reads were aligned to the *T. dactyloides* reference

genome (Td-KS_B6_1-REFERENCE-PanAnd-2.0a[28] using HISAT2. The alignments were name-

sorted so that mate-pairs could be fixed using the fixmate function of samtools[28], and then re-

sorted based on coordinates, de-duplicated, and converted to indexed BAM files using the same

software. Finally, a table of transcript expression estimates was generated using the

FeatureCounts function of the subread package with parameters -p -O --fraction -M --primary -g

ID -t gene and the *T. dactyloides* genome annotation version Td-KS_B6_1-REFERENCE-

PanAnd-2.0a_Td00002ba.2.

3.5.3 Statistical analyses

Because the maize and *T. dactyloides* RNA-seq datasets were generated using different

approaches (3' tag sequencing *vs.* full-length sequencing, respectively) we analyzed them in

parallel rather than comparing them directly. For each species, we used DESeq2[16] to identify

genes that were differentially expressed between plants inoculated with microbiomes from a low-

precipitation climate ("dry legacy") *vs.* those inoculated with microbiomes from a high-precipitation

climate ("wet legacy"). A single negative binomial model with default parameters was used to

estimate $\log_2$-fold changes in gene expression due to inoculum legacy, while also controlling for

the other experimental factors, using the model:

Counts ~ Legacy + ConditioningPhaseWater + ConditioningPhaseHost + TestPhaseWater

Statistical support was obtained using the Wald test with Benjamini-Hochberg FDR correction. All

available samples were included in each analysis; thus, these results should be interpreted as

the gene expression response to microbiome precipitation legacy, averaged across all levels of

the other experimental factors.

In addition, we investigated whether plants' gene expression responses to limited *vs.* ample water

during the Test Phase were affected by inoculum precipitation legacy. To do so, we used DESeq2

to fit a model with the formula:

Counts ~TestPhaseWater*Legacy

Then, we extracted the estimated $\log_2$-fold changes due to acute drought (relative to well-watered

conditions) for both dry-legacy-inoculated and wet-legacy-inoculated plants. We inferred a

meaningful interaction between these variables when the 95% confidence intervals of the two

drought-induced $\log_2$-fold changes did not overlap at all. These interacting genes are candidates

for linking real-time plant drought response to the microbiome's historical environmental

conditions.

Finally, we conducted a mediation analysis to determine whether the *T. dactyloides* genes that

were sensitive to inoculum legacy were implicated in phenotypic responses to subsequent acute

drought. A gene was considered legacy-sensitive if its expression was significantly affected by

the main effect of inoculum legacy, or if it was affected by the interaction between inoculum legacy

and test phase drought treatment, as described above. We summarized expression patterns of

this subset of genes (normalized as transcripts per million, calculated using the full set of

expressed genes) using non-metric multidimensional scaling of the Bray-Curtis distances among

all *T. dactyloides* plants, which resulted in two axes of variation: MDS1 and MDS2. Next, we used

the mediation package in R[53] to compare the direct effects of test phase drought treatment on

each focal plant trait (see section *3.7.1. Plant trait feature selection*) to the indirect effects of the

drought treatment mediated through MDS1 and MDS2. Each mediation analysis used the linear

models:

Trait value ~ MDS + Test phase drought treatment

MDS ~ Test phase drought treatment

where MDS represents the "site score" of each individual plant on either MDS1 or MDS2.

Separate models were fit to test the potential roles of MDS1 and MDS2 as mediator variables;

however, a follow-up analysis using both MDS1 and MDS2 as simultaneous mediators,

implemented in lavaan[54], yielded equivalent conclusions.

3.5.4 Gene annotation

We downloaded maize and eastern gamagrass genome assemblies and annotations from

MaizeGDB[50,55]. Functional information for maize genes was taken from the *Zea mays* genome

annotation version Zm-B73-REFERENCE-NAM-5.0_Zm00001eb.1 and accessed using

MaizeGDB's MaizeMine tool[50,55–57]. For *Tripsacum dactyloides* genes, we relied on DNA

sequence homology with annotated maize genes to infer function. We used BEDtools[58] to extract

gene coordinates and protein-coding sequences from the *T. dactyloides* reference genome Td-

KS_B6_1-REFERENCE-PanAnd-2.0a, and then used OrthoFinder[59] to compare coding

sequences from the two species. OrthoFinder identified 32,785 *T. dactyloides* genes (71.5% of

the total) as homologs of maize genes, grouping them into 21,658 distinct orthogroups. All

differential gene expression analyses, however, considered the entire set of expressed genes;

those without maize orthologs, or with unannotated maize orthologs, were considered to be of

unknown function.

3.6 16S rRNA amplicon sequencing

3.6.1 DNA extractions and library preparation

One crown root from each plant was cut off with a ceramic blade and placed in a 1.7 mL tube and

flash frozen in liquid nitrogen for downstream amplicon library preparation. After collection, root

sub-samples were kept on dry ice and ceramic tweezers were used to transfer the whole root to

1.1 mL cluster tubes (USA Scientific, Ocala, FL, USA). Tweezers were sterilized with 80% EtOH

between samples. Roots were selected at random for placement in cluster tubes and were stored

at -80ºC until DNA extraction, at which time roots were freeze-dried for 48 hours in a FreeZone

lyophilizer (Labconco, Kansas City, MO, USA). After freeze-drying, roots were flash frozen in

liquid nitrogen. To break apart thick roots, sterile forceps and a dissecting needle were used

before placing the rack of cluster tubes in an HT Lysing Homogenizer (OHAUS, Parsippany, NJ,

USA) with two clean 5/32" steel balls in each tube. Samples were homogenized at 25 Hz for 1

min. Root material was then transferred to 2 mL bead-beating 96-well plates containing sterile 1

mm garnet beads with 850 µL of lysis buffer (1M Tris, pH = 8.0; 100 mM NaCl; 10 mM EDTA).

A positive (ZymoBiomics Microbial Community Standard) and negative control, (800 µL of lysis

buffer) were included on each plate. Bead-beating plates were stored at -20ºC until extraction.

After thawing, 10 µL of 20% sodium dodecyl sulfate (SDS) was added to each well before

homogenizing for a total of 20 min at 20 Hz. Plates were incubated in a water bath (55ºC for 90

min) and centrifuged (6 min at 4500 x g). 400 µL of the resulting supernatant was transferred to

new 1 mL 96-well plates containing 120 µL of 5 M potassium acetate in each well and incubated

overnight at -20ºC. After thawing the plates, they were centrifuged (6 min at 4500 x g) and 400

µL of the supernatant was transferred to a new 1 mL 96-well plate containing 600 µL of diluted

SPRI-bead solution (protocol derived from[60]). These plates were mixed thoroughly for 5 min at

1000 r.p.m. on an orbital plate shaker. Samples were allowed to incubate for 10 min so DNA could

bind to beads, after which the plate was centrifuged (6 min at 4500 x g) and placed on a magnet

rack for 10 min. The supernatant was removed, and the beads were washed twice with 900 µL of

80% EtOH. After washing, the supernatant was decanted and the beads were air-dried. DNA was

eluted in 75 µL of pre-heated 1x TE (pH = 7.5; 37ºC) and transferred to clean 0.45 mL plates and

stored at -20ºC.

16S-v4 rRNA gene amplification was performed using paired 515f/806r primers[61,62]. PCR

reactions contained DreamTaq Master Mix (Thermo Fisher Scientific, Waltham, MA, USA), 10

mg/µL bovine serum albumin (BSA), 100 µM peptide nucleic acid (PNA), and PCR-grade water.

BSA was used to enhance PCR amplification while PNA was used to suppress primer binding

and subsequent amplification of mitochondrial and chloroplast 16S regions[63]. The PCR included

an initial denaturation step at 95ºC for 2 min, followed by 27 cycles of an additional denaturation

at 95ºC for 20 s, PNA annealing at 78ºC for 5 s, primer annealing at 52ºC for 20 s, and extension

at 72ºC for 50 s. This was followed by a final extension step for 10 min at 72ºC. PCR products

were purified by incubating at 37ºC for 20 min and then 15 min at 80ºC after mixing with 0.78 µL

of PCR-grade water, 0.02 µL of 10 U/µL exonuclease I (Applied Biosystems), and 0.2 µL of 1U/µL

shrimp alkaline phosphatase (Applied Biosystems) per 10 µL PCR product. Two µL of the purified

amplicons were used as template DNA in an indexing PCR to attach barcoded P5 and P7 Illumina

adaptors. The 10 µL reaction included 5 µM each of P5 and P7 adaptors, 1x DreamTaq Master

Mix (Thermo Fisher Scientific), 10 mg/mL BSA, 100 µM PNA, and PCR-grade water. An initial

denaturation step was performed at 95ºC for 2 min. For 8 cycles, an additional denaturation was

carried out at 95ºC for 20 s, PNA annealing at 78ºC for 5 s, primer annealing at 52ºC for 20 s,

and extension at 72ºC for 50 s. A final extension step was performed at 72ºC for 10 min. PCR

products were verified via 2% agarose gel electrophoresis and then pooled by 96-well plate using

the "Just-a-Plate" PCR cleanup and normalization kit (Charm Biotech, St. Louis, MO, USA). Pools

were size selected and combined in equimolar concentrations before being sequenced on an

Illumina SP flow cell on the NovaSeq 6000 platform (2x250bp reads).

1010

3.6.2 Amplicon data processing

We removed primer sequences from raw 16S-v4 Illumina reads using cutadapt[37], requiring at

least 5 nucleotides of overlap. Additional quality control and processing was performed with the

DADA2 software[64]. Forward reads were discarded if they had more than 6 expected errors,

otherwise they were truncated at 200 nucleotides; for reverse reads the parameters were 7

expected errors and 170 nucleotides. Error rates were estimated separately for forward and

reverse reads based on a sample of $1x10^8$ bases, then used to denoise and dereplicate reads

using the standard DADA2 functions. Chimeric sequences were detected and removed using the

"consensus" procedure in DADA2. Each individual sample was processed in parallel, after which

all of the resulting amplicon sequence variant (ASV) tables were merged. Finally, taxonomy of

each ASV was assigned by comparison to the RDP database[65].

1022

3.6.3 Root bacterial community diversity analysis

All statistical analyses were performed using R (v4.4.0). The cleaned and prepared microbiome

data, as phyloseq objects, were loaded into R[35]. The phyloseq object sample data table was

replaced with an updated metadata file. The "mock" samples were removed from the dataset and

the data was subset by the test phase host, maize and gamagrass using

phyloseq::subset_samples (v1.48.0). Maize and gamagrass root bacterial microbiome Shannon

Diversity was fit to a mixed-effects model using the lme4 (v1.1-35.4) package[66] and lmer()

function, using the following model:

1031   expShannon~TestWater+Legacy + (1|DNAplate)

1032

1033 A type III ANOVA was performed using stats::anova()[67] to test for significance, followed by

1034 pairwise comparisons and significance tests using emmeans (v1.10.2) package[68] with false

1035 discovery rate adjustments. Data visualizations were generated using ggplot2 (v3.5.1)[12], the error

1036   bars represent standard error and the points are estimated marginal means. All plots were saved

1037   in PDF format. Composite figures were created in Adobe Illustrator.

1038

1039   Bacterial community beta diversity was accessed with a Constrained Analysis of Principal

1040   Coordinates, using phyloseq::ordinate, method="CAP", and distance="Euclidean" with the

1041   following formulas:

1042   ~Legacy+Condition(DNAplate)+Condition(CondHost)+

1043   Condition(CondWater)+Condition(TestWater)

1044

1045   ~TestWater+Condition(DNAplate)+Condition(CondHost)+

1046   Condition(CondWater)+Condition(Legacy)

1047

An ANOVA-like permutation test was used to test for model significance using the

vegan::anova.cca() function[13]. CAP1 and MDS1 axes were plotted using ggplot2, including 95%

confidence interval ellipse (stat_ellipse()). To evaluate treatment group beta-dispersion,

Euclidean distances were calculated for the centered-log ratio transformed counts using

phyloseq::distance(). Both "Legacy" and "TestWater" terms were tested for beta-dispersion

differences using the Euclidean distances matrix and vegan::betadisper() and vegan::permutest()

to test for significance[13]. Distances were then extracted from the results output and fit to a linear

model. Estimated marginal means were calculated and plotted using the same methods as above.

The error bars represent standard errors.

1057

1058   Then, an ASV differential abundance analysis was performed on gamagrass- and maize-

1059   associated ASVs separately. Centered-log ratio transformed counts of ASVs were each fit to a

1060   linear model using an iterative for-loop. Each ASV was subset using phyloseq:subset_taxa and

1061   the phyloseq::psmelt() function was used to reformat the phyloseq object into a data frame. Then

1062   the ASV abundance was fit to the following model:

1063   Abundance ~ Legacy*TestWater

1064

1065   Then, stats::anova() was used to test for significance, followed by an FDR p-value adjustment

1066   using stats::p.adjust(). ASVs with significant (≤0.05) adjusted p-values for the Legacy*TestWater

1067   interaction terms were extracted and z-scores of the center-log ration transformed counts were

1068   plotted in a heatmap using pheatmap::pheatmap()[69] with clustering_method="complete".

1069

3.7 Plant trait data analyses

3.7.1 Plant trait feature selection

All plant data was loaded into RStudio. The intrinsic water use efficiency (iWUE) was calculated

using the ratio of photosynthetic rate (A) and stomatal conductance (gs) or A/gs ((μmol m^{-2}

s^{-1})/(mol m^{-2} s^{-1})). Data was subset by Test Phase host treatment groups (gamagrass and maize).

For the feature selection, the "mock" treatment was removed. The 67 features that were selected

for testing were subset into a data frame. Three sample rows were removed from the dataset

because they had missing data. The drought susceptibility index (S-index) was calculated for

each of these features using the following formula:

1079 $S = Trait_{drought}/mean(Trait_{control})$

1080

Then, for each host, a random forest model was used to select the top 10 most important features

from the 67 total s-index measurements using randomForest()[70], with "Legacy" as the predictor.

Correlations (stats::cor()) were estimated for the top ten traits and if any two of the top 10 traits

were highly correlated ($r \geq 0.7$), the trait that ranked lower in the random forest model was

removed. Non-correlative top features from the test water treatment groups (drought vs. watered

control) were combined for each test phase host (maize and gamagrass). Vegan:rda() was used

to perform a redundancy analysis on the top features using the following formula:

Top_features~ Legacy + Condition(CondGroup), scale=TRUE

1089

1090   *3.7.2 Feature analyses*

1091   An ANOVA-like permutation test was used to test for model significance using the

1092   vegan::anova.cca() function. A biplot was created using ggplot2 plotting RDA1 and PC1 axes, as

1093   well as the species scores (represented by arrows), and stat_ellipse was used to plot 95%

1094   confidence intervals of the site scores.

1095

Finally, s-indices for each of the top features and iWUE were fit to mixed-effects models, when

possible, or a fixed-effects model if overfitting of the mixed effects model occurred. Each feature

or trait was visually assessed for outliers, which were removed. Removing the outliers did not

impact the interpretation of any of the results. Formulas used for the mixed effects or linear model,

respectively, include:

lmer(s-index ~ Legacy + (1|CondGroup))

lm(s-index ~ Legacy + CondGroup)

1103

1104   The fit of the model was accessed and if needed, the s-index was transformed using sqrt(), exp(),

1105   or log() to improve the fit. ANOVA was used to assess significance and estimated marginal means

1106   were calculated and plotted as described above.

1107

REFERENCES

1. Stackhouse, P. Nasa power. https://power.larc.nasa.gov/.

2. Neves, D. J. D., Silva, V. de P. R., Almeida, R. S. R., Sousa, F. de A. S. de & Silva, B. B.

da. General aspects of the climate in the Cabo verde archipelago. *Rev. Ambiência* **13**,

(2017).

3. Varela-Lopes, G. E. & Molion, L. C. B. Precipitation patterns in Cape Verde islands:

Santiago island case study. *Atmos. Clim. Sci.* **04**, 854–865 (2014).

4. Moreno, Â., Ramos, T. B., Gonçalves, M. C. & Pereira, L. S. Estimating soil hydraulic

properties from limited data to improve irrigation management in agricultural soils of

Santiago island, Cape Verde: Soil hydraulic properties in Cape Verde soils. *Irrig. Drain.* **63**,

405–415 (2014).

5. Unknown, (:unkn). Guide to Meteorological Instruments and Methods of Observation:

(CIMO guide). 2014 edition, updated in 2017.[SUPERSEDED]. Preprint at

https://doi.org/10.25607/OBP-432 (2017).

6. Soltanpour, P. N. & Schwab, A. P. A new soil test for simultaneous extraction of macro-

and micro-nutrients in alkaline soils. *Commun. Soil Sci. Plant Anal.* **8**, 195–207 (1977).

7. De Coster, W., D'Hert, S., Schultz, D. T., Cruts, M. & Van Broeckhoven, C. NanoPack:

visualizing and processing long-read sequencing data. *Bioinformatics* **34**, 2666–2669

(2018).

8. Wood, D. E., Lu, J. & Langmead, B. Improved metagenomic analysis with Kraken 2.

(2019) doi:10.1101/762302.

9. Lu, J., Breitwieser, F. P., Thielen, P. & Salzberg, S. L. Bracken: estimating species

abundance in metagenomics data. *PeerJ Comput. Sci.* **3**, e104 (2017).

10.   Breitwieser, F. P. & Salzberg, S. L. Pavian: interactive analysis of metagenomics

data for microbiome studies and pathogen identification. *Bioinformatics* **36**, 1303–1304

(2020).

11.   Hothorn, T., Bretz, F. & Westfall, P. Simultaneous inference in general

parametric models. *Biom. J.* **50**, 346–363 (2008).

12.   Wickham, H. *Ggplot2: Elegant Graphics for Data Analysis*. (Springer, Berlin,

Germany, 2016).

13.   Oksanen, J. *et al.* vegan : Community Ecology Package. R package version 1.8-
1139   5. (2007).

14.   The R Project for Statistical Computing. http://www.R-project.org/.

15.   Schneider, C. A., Rasband, W. S. & Eliceiri, K. W. NIH Image to ImageJ: 25
1142   years of image analysis. *Nat. Methods* **9**, 671–675 (2012).

16.   Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and
1144   dispersion for RNA-seq data with DESeq2. *Genome Biol.* **15**, 550 (2014).

17.   Kolmogorov, M. *et al.* metaFlye: scalable long-read metagenome assembly using
1146   repeat graphs. *Nat. Methods* **17**, 1103–1110 (2020).

18.   BBTools. *DOE Joint Genome Institute* https://jgi.doe.gov/data-and-tools/bbtools/
1148   (2016).

19.   Li, H. Minimap2: pairwise alignment for nucleotide sequences. *Bioinformatics* **34**,
1150   3094–3100 (2018).

20.   Woodcroft, B. J. *CoverM: Read Coverage Calculator for Metagenomics*. (Github).

21.   von Meijenfeldt, F. A. B., Arkhipova, K., Cambuy, D. D., Coutinho, F. H. & Dutilh,
1153   B. E. Robust taxonomic classification of uncharted microbial sequences and bins with CAT
1154   and BAT. *Genome Biol.* **20**, 217 (2019).

22.   Van der Jeugt, F., Dawyndt, P. & Mesuere, B. FragGeneScanRs: faster gene
1156   prediction for short reads. *BMC Bioinformatics* **23**, 198 (2022).

23.   Cantalapiedra, C. P., Hernández-Plaza, A., Letunic, I., Bork, P. & Huerta-Cepas,
1158   J. eggNOG-mapper v2: Functional Annotation, Orthology Assignments, and Domain
1159   Prediction at the Metagenomic Scale. *Mol. Biol. Evol.* **38**, 5825–5829 (2021).

24.   Huerta-Cepas, J. *et al.* eggNOG 5.0: a hierarchical, functionally and
phylogenetically annotated orthology resource based on 5090 organisms and 2502 viruses.
Nucleic Acids Res. **47**, D309–D314 (2019).

25.   Buchfink, B., Reuter, K. & Drost, H.-G. Sensitive protein alignments at tree-of-life

scale using DIAMOND. *Nat. Methods* **18**, 366–368 (2021).

26.   Steinegger, M. & Söding, J. MMseqs2 enables sensitive protein sequence

searching for the analysis of massive data sets. *Nat. Biotechnol.* **35**, 1026–1028 (2017).

27.   Wright, R. M., Aglyamova, G. V., Meyer, E. & Matz, M. V. Gene expression

associated with white syndromes in a reef building coral, Acropora hyacinthus. *BMC*

Genomics **16**, 371 (2015).

28.   Li, H. *et al.* The Sequence Alignment/Map format and SAMtools. *Bioinformatics*

25, 2078–2079 (2009).

29.   Danecek, P. *et al.* Twelve years of SAMtools and BCFtools. *Gigascience* **10**,

(2021).

30.   Danecek, P. *et al.* The variant call format and VCFtools. *Bioinformatics* **27**, 2156–

2158 (2011).

31.   Chang, C. C. *et al.* Second-generation PLINK: rising to the challenge of larger

and richer datasets. *Gigascience* **4**, 7 (2015).

32.   Yin, L. *et al.* RMVP: A memory-efficient, visualization-enhanced, and parallel-

accelerated tool for genome-wide association study. *Genomics Proteomics Bioinformatics*

19, 619–628 (2021).

33.   LiLin-Yin. CMplot: Circle Manhattan Plot. *CRAN: Contributed Packages* The R

Foundation https://doi.org/10.32614/cran.package.cmplot (2015).

34.   Faisal, M., Silander, O. & Freed, N. Starting a MinION sequencing run using

MinKNOW v1. (2020) doi:10.17504/protocols.io.bfwqjpdw.

35.   McMurdie, P. J. & Holmes, S. phyloseq: an R package for reproducible

interactive analysis and graphics of microbiome census data. *PLoS One* **8**, e61217 (2013).

36.   Shetty, S. Lahti L. microbiomeutilities: microbiomeutilities: Utilities for Microbiome

Analytics. 2022. Preprint at (2023).

37.   Martin, M. Cutadapt removes adapter sequences from high-throughput

sequencing reads. *EMBnet.journal* **17**, 10–12 (2011).

38.   Shetty, S. A. & Lahti, L. Microbiome data science. *J. Biosci.* **44**, (2019).

39.   De Mendiburu, F. & Simon, R. Agricolae - Ten years of an open source statistical

tool for experiments in breeding, agriculture and biology. *PeerJ* (2015)

doi:10.7287/peerj.preprints.1404.

40.   The Hmisc and rms Packages. in *Biostatistics and Computer-based Analysis of*

Health Data using R 173–186 (Elsevier, 2016).

41.   Grabherr, M. G. *et al.* Full-length transcriptome assembly from RNA-Seq data

without a reference genome. *Nat. Biotechnol.* **29**, 644–652 (2011).

42.   Kopylova, E., Noé, L. & Touzet, H. SortMeRNA: fast and accurate filtering of

ribosomal RNAs in metatranscriptomic data. *Bioinformatics* **28**, 3211–3217 (2012).

43.   Patro, R., Duggal, G., Love, M. I., Irizarry, R. A. & Kingsford, C. Salmon provides

fast and bias-aware quantification of transcript expression. *Nat. Methods* **14**, 417–419

(2017).

44.   Soneson, C., Love, M. I. & Robinson, M. D. Differential analyses for RNA-seq:

transcript-level estimates improve gene-level inferences. *F1000Res.* **4**, 1521 (2015).

45.   Kamvar, Z. N., Tabima, J. F. & Grünwald, N. J. Poppr: an R package for genetic

analysis of populations with clonal, partially clonal, and/or sexual reproduction. *PeerJ* **2**,

e281 (2014).

46.   Thioulouse, J. *et al. Multivariate Analysis of Ecological Data with ade4.* (Springer,

New York, NY, 2019).

47.   Goudet, J. hierfstat, a package for r to compute and test hierarchical *F*-statistics.

Mol. Ecol. Notes **5**, 184–186 (2005).

48.   Seethepalli, A. *et al.* RhizoVision Explorer: open-source software for root image

analysis and measurement standardization. *AoB Plants* **13**, lab056 (2021).

49.   Smith, T., Heger, A. & Sudbery, I. UMI-tools: modeling sequencing errors in

Unique Molecular Identifiers to improve quantification accuracy. *Genome Res.* **27**, 491–499

(2017).

50.   Hufford, M. B. *et al.* De novo assembly, annotation, and comparative analysis of

26 diverse maize genomes. *Science* **373**, 655–662 (2021).

51.   Guo, J., Gao, J. & Liu, Z. HISAT2 parallelization method based on Spark cluster.

J. Phys. Conf. Ser. **2179**, 012038 (2022).

52.   Liao, Y., Smyth, G. K. & Shi, W. featureCounts: an efficient general purpose

program for assigning sequence reads to genomic features. *Bioinformatics* **30**, 923–930

(2014).

53.   Yimer, B. B., Lunt, M., Beasley, M., Macfarlane, G. J. & McBeth, J. BayesGmed:

An R-package for Bayesian causal mediation analysis. *PLoS One* **18**, e0287037 (2023).

54.   Rosseel, Y. lavaan: An R Package for Structural Equation Modeling. *J. Stat.*

Softw. **48**, 1–36 (2012).

55.   Portwood, J. L., 2nd *et al.* MaizeGDB 2018: the maize multi-genome genetics

and genomics database. *Nucleic Acids Res.* **47**, D1146–D1154 (2019).

56.   Woodhouse, M. R. *et al.* A pan-genomic approach to genome databases using

maize as a model system. *BMC Plant Biol.* **21**, 385 (2021).

57.   Shamimuzzaman, M. *et al.* MaizeMine: A Data Mining Warehouse for the Maize

Genetics and Genomics Database. *Front. Plant Sci.* **11**, 592730 (2020).

58.   Quinlan, A. R. BEDTools: The Swiss-Army Tool for Genome Feature Analysis.

Curr. Protoc. Bioinformatics **47**, 11.12.1–34 (2014).

59.   Emms, D. M. & Kelly, S. OrthoFinder: phylogenetic orthology inference for

comparative genomics. *Genome Biol.* **20**, 238 (2019).

60.   Rohland, N. & Reich, D. Cost-effective, high-throughput DNA sequencing

libraries for multiplexed target capture. *Genome Res.* **22**, 939–946 (2012).

61.   Apprill, A., McNally, S., Parsons, R. & Weber, L. Minor revision to V4 region SSU

rRNA 806R gene primer greatly increases detection of SAR11 bacterioplankton. *Aquat.*

Microb. Ecol. **75**, 129–137 (2015).

62.   Parada, A. E., Needham, D. M. & Fuhrman, J. A. Every base matters: assessing

small subunit rRNA primers for marine microbiomes with mock communities, time series

and global field samples. *Environ. Microbiol.* **18**, 1403–1414 (2016).

63.   Lundberg, D. S., Yourstone, S., Mieczkowski, P., Jones, C. D. & Dangl, J. L.

Practical innovations for high-throughput amplicon sequencing. *Nat. Methods* **10**, 999–1002

(2013).

64.   Callahan, B. J. *et al.* DADA2: High-resolution sample inference from Illumina

amplicon data. *Nat. Methods* **13**, 581–583 (2016).

65.   Cole, J. R. *et al.* Ribosomal Database Project: data and tools for high throughput

rRNA analysis. *Nucleic Acids Res.* **42**, D633–42 (2014).

66.   Bates, D., Mächler, M., Bolker, B. & Walker, S. Fitting linear mixed-effects

models Usinglme4. *J. Stat. Softw.* **67**, (2015).

67.   *R: A Language and Environment for Statistical Computing : Reference Index.*

(2010).

68.   https://CRAN.R-project.org/package=emmeans.

69.   pheatmap: Pretty Heatmaps. *Comprehensive R Archive Network (CRAN)*

https://CRAN.R-project.org/package=pheatmap.

70.   CRAN: R News. https://CRAN.R-project.org/doc/Rnews/.

1262

1263

1264
1265
1266
1267

Figures

Extended Data Fig. 1. Precipitation level and mineral nutrient content in Kansas soils
correlate with soil microbiota composition. a. Alpha diversity (estimated using the Shannon
diversity index) did not differ across soil samples. Soils are ordered according to the gradient of
precipitation (Fig 1a). **b**. Phylogram showing the relative abundance profiles of the main bacterial
phyla across soils exposed to different precipitation levels. **c**. Principal component analysis
showing the projected ionomic profiles among the different soils exposed to the gradient of
precipitation. The bar on the left denotes the percentage of the variance explained by the different
experimental variables. **d**. Heatmap showing the standardized concentration (z-score) of each
mineral nutrient (rows) in the collection of soil exposed to different precipitation levels. The values
were clustered according to the ion concentration and the region was ordered according to the
precipitation gradient (lower to high). **e**. Bar graph showing the Pearson correlation coefficient
between each mineral nutrient abundance and the level of precipitation across the collection of
soils used. Coloured bars are significant (*q* < 0.05). **f**. Pairwise correlation analysis between soil
mineral nutrient dissimilarities and soil microbiome composition dissimilarities in soils
representing a gradient of precipitation. Panel shows the Mantel *r* statistic and its *p*-value.

1286
1287

1288
Extended Data Fig. 2. Precipitation levels, but not soil porosity, explain microbiota
composition across the Kansas soil collection. a. Percent soil porosity changes with depth in
soils exposed to the precipitation gradient. The data were obtained from soil images generated
by X-ray computed tomography. **b**. Pearson correlation analysis between the soil porosity and
the precipitation gradient. The correlation coefficient r and its *p*-value are shown. **c**. Pairwise
correlation analysis between soil microbiota dissimilarities and soil porosity dissimilarities. The

panel show the Mantel *r* statistic and its *p*-value. **d-f**. Heatmaps showing changes in the relative abundance of bacterial taxa (d), species (e), and family (f) that were found to be significantly different from the wetter region in the Kansas soil collection. In all cases, the values have been clustered according to taxonomic categories and soils. Note that this soil collection represents a precipitation gradient. **g.** The heatmap illustrates the enriched or depleted biological processes between soils from three wet regions (high precipitation: WEL, CWR, and KNZ) and three dry regions (low precipitation: TRI, SVR, and HAY). Gene enrichment analysis was conducted using a generalized linear model, followed by Gene Ontology (GO) classification. Significantly enriched and depleted GO categories were identified based on an adjusted *p*-value threshold of < 0.05. The colours (red and blue) represent the enrichment score, calculated using square root-transformed delta rank values of the GO categories. The line graph on the right displays the -log10 adjusted *p*-value of the enriched or depleted GO categories in the comparison.

Extended Data Fig. 3. The microgradient of precipitation identified in Santiago island and the soil mineral content influence the general soil microbiota structure. a. Map of Santiago island, Cabo Verde, showing the locations selected for collecting the soils used in this work. **b.**

The graph shows the precipitation (mm/year) recorded or estimated for the Santiago soils over
the period 1981 to 2009 across diverse databases. Statistical significance between soils was
determined via linear modelling and letters represent the compact letter display of a Tukey *post-*
hoc test. **c**. Alpha diversity, measured using the Shannon diversity index, did not differ across soil
samples. Soils are ordered according to the gradient of precipitation (panel b)**. d**. Phylogram
showing the relative abundance profiles of the most abundant bacterial phyla across Santiago
soils. **e**. Principal component analysis showing the projected microbiota composition among the
different soils (colours). The bar on the left denotes the percentage of the variance explained by
the different experimental variables. **f**. Principal component analysis showing the projected
mineral nutrient profiles among the different soils (colours). The bar on the left denotes the
percentage of the variance explained by the different experimental variables. **g**. Heatmap showing
the standardized concentration (z-score) of each mineral nutrient (rows) in the collection of
Santiago soils. The values have been clustered according to the ion concentration and the region
h. Bar graph showing the Pearson correlation coefficient between each mineral nutrient
abundance and the level of precipitation across the collection of soils. Coloured bars are
statistically significant ($q < 0.05$). **i**. Pairwise correlation analysis between soil mineral nutrient
content dissimilarities and soil microbiota composition dissimilarities. Panel shows the
Mantel *r* statistic and its *p*-value.

1445 **Extended Data Fig. 4. The microgradient of precipitation identified in Santiago explains the**
1446 **soil microbiota composition. a.** Percentage of soil porosity changes with the soil depth in soils

across the precipitation microgradient. The data were obtained from soil images generated by X-ray computed tomography. The colours represent different soils. **b**. Pearson correlation analysis between the soil porosity and the precipitation microgradient. The correlation coefficient r and its *p*-value are shown. **c.** Pairwise correlation analysis between soil porosity dissimilarities and soil mineral nutrient content (ionome) dissimilarities. **d.** Pairwise correlation analysis between soil porosity dissimilarities and soil microbiota composition dissimilarities. In both panels, c and d, the Mantel *r* statistic and its *p*-value are shown. **e-g.** Heatmaps showing changes in relative abundance of **(e)** bacterial taxa, **(f)** species and **(g)** families across the soil collection. Only bacterial taxa with a significantly different relative abundance from the wetter region in the rainfall microgradient were used for this analysis. In all cases, values have been grouped by taxonomic categories and soils. **h.** Bar graph showing the Pearson correlation coefficient between the relative abundance of each bacterial taxa and the level of precipitation. Coloured bars are significant (*q* < 0.05). **i.** The heatmap depicts enriched or depleted biological processes between soils from the region with high annual precipitation (Pico de Antónia) and the remaining soils (Achada Leitão, Serra Malagueta, Mato Correia, Ribeira de São Domingos (Nora), Porto Gouveia, Tomba Toiro, Colonato, and Ribeira Riba). Gene enrichment analysis was performed using a generalized linear model, followed by Gene Ontology (GO) classification. Significantly enriched and depleted GO categories were identified based on an adjusted *p*-value threshold of < 0.05. Enriched (red) and depleted (green) GO terms represent the enrichment score, calculated using square root-transformed delta rank values of the GO categories. **j-k.** Comparative analysis of the relative abundances of bacterial markers identified in soils from **(j)** Kansas and **(k)** Santiago Island in the corresponding pool of soils with high or low levels of precipitation. Coloured points represent the average value of standardized relative abundance for each region (Kansas and Santiago). The line connecting both points is the difference between low and high precipitation soils. A black line indicates statistical significance (*q* < 0.05).

Extended Data Fig. 5. Genetic variation among bacterial lineages along the Kansas and Santiago precipitation gradients. To assess genetic differences between bacteria lineages along the precipitation gradient, we selected 33 bacterial species (22 of the identified bacterial markers and 11 additional abundant and prevalent taxa) that exhibited high genome coverage in the soil samples from Kansas, USA and Santiago Island, Cabo Verde, as proxies for the broader bacterial communities. Reference genomes for each species were retrieved from the NCBI Genome database, and filtered shotgun metagenomic reads were mapped to these genomes to identify high-quality biallelic single nucleotide polymorphisms (SNPs). Genetic distances between the bacterial lineages were calculated based on the identified SNPs and PCoA plots were generated and coloured by **a**. soil type, **b**. precipitation levels and **c**. geographical regions of Kansas (Kansas State University's Tribune Southwest Research Center (TRI), Smoky Valley Ranch (SVR), Hays Prairie (HAY), Konza Wildlife Reserve (KNZ), Clinton Wildlife Reserve (CWR), Welda Prairie (WEL)) and Santiago Island (Ribeira Riba, Porto Gouveia, Ribeira de São Domingos (Nora), Colonato, Mato Correia, Tomba Toiro, Achada Leitão, Serra Malagueta, Pico de Antónia). The variance explained by each axis is indicated. Manhattan plot **d**. illustrates significant SNPs linked to the precipitation gradient, derived from the genetic-environment association (GEA) analysis. The GEA analysis was conducted using a general linear model, with average precipitation at each sampling location as the environmental variable. Significant associations were identified using the permutation method. The x-axis of the plot represents the SNP positions along the genomes of the selected bacterial species, while the y-axis displays the -$\log_{10}$ p-values from the association model. The horizontal line indicates the statistical significance threshold, as determined by the permutation test. SNPs above this threshold, highlighted in red, were significantly associated with the precipitation gradient. Bacterial taxa with fewer than 1,000 high-quality biallelic SNPs after filtering, and with no significant SNPs detected from the GEA analysis, are not shown in the plot.

1529

Extended Data Fig. 6. Soil water legacy is resilient to short-term water- and host-related perturbations. a. Schematic representation of the experimental pipeline used to evaluate the

resilience of the soil water legacy to perturbations and their functional importance to plant drought
response. The six soils from Kansas, representing a gradient of precipitation with clear ecological
memory of water availability (Legacy phase), were either left unplanted or planted with seedlings
of the native grass species *Tripsacum dactyloides* (eastern gamagrass), and then subjected to
either drought conditions or regular watering in a factorial design (Conditioning phase). To
evaluate the functional effects of soil precipitation legacy on plants, and to isolate the role of the
microbiota from possible roles of non-microbial soil properties, we then used the experimentally-
conditioned microbial communities to inoculate a new generation of *T. dactyloides* and *Z. mays*
plants. These "Test Phase" plants were divided between water-limited conditions and well-
watered control conditions. **b**. Box plot showing the quantification of different *Tripsacum*
dactyloides phenotypes in response to the treatments applied (drought conditions or regular
watering) during the conditioning phase. ANOVA was used to discern significant differences
among groups and estimated marginal means were compared using Tukey's post hoc test. **c**.
Exemplary pictures taken by laser ablation tomography (LAT) of crown roots of *Tripsacum*
dactyloides grown in soils with a high-precipitation (CWR) and low-precipitation (SVR) history
after exposure to acute drought during. Note that the number of aerenchyma in the root, important
for the drought response, in plants exposed to a low precipitation legacy (SVR) is higher compared
to that found in plants grown in soils with high precipitation history (CWR). **d.** Phenotypic traits of
Tripsacum dactyloides in response to the conditioning phase treatments (drought or regular
watering). The heatmap on the left depicts key features extracted from the plant, colored based
on mean normalised values. Pearson correlation coefficients (r) between these features are
presented in the centre heatmap, with significant correlations ($p < 0.05$) highlighted by black
squares. The bar plot on the right indicates the coefficient of variation of the feature values, with
highly variable features highlighted in blue. The heatmaps are clustered based on Pearson
correlation coefficient values. **e**. Alpha diversity of bacterial communities, measured using the
Shannon diversity index, was not affected by the different conditioning treatments (drought or
well-watered, with or without host). **f**. Phylograms show that the different conditioning treatments
(drought or well-watered, with or without host) did not impact the relative abundance profiles of
main bacterial phyla. **g**. Canonical analysis of principal coordinates (CAP) showing the effects of
conditioning phase treatments on microbiome assembly: (left) soil legacy (baseline), well-
watered, drought, and (right) host or no host. PERMANOVA R^2 and *p*-value are shown within the
CAP plot. The bar on the left describes the percentage of the variance explained by the
experimental variables. **h**. The heatmap displays the enrichment of bacterial families across soils
with wet and dry legacies, exposed to either drought or well-watered treatments, in comparison
to soils with a wet legacy. Colors in the heatmap represent $\log_2$ fold changes, calculated using a
generalized linear model that contrasts the abundance of each bacterial family in the respective
treatments with that of the wet soil legacy. Tiles outlined in black denote statistically significant
enrichment (red) or depletion (blue) ($q < 0.05$) with a $\log_2$ fold change > ±2 with respect to the
soils with wet legacy. The heatmap was clustered based on taxonomic classification (represented
by the dendrogram on the left). **i**. Boxplots showing the relative abundance of bacterial markers
of water legacy across soils with dry or wet legacy (baseline) after exposure to conditioning phase
treatments (drought or well-watered, with or without host). ANOVA was performed to detect
significant differences among the groups, with Tukey's post hoc test used to compare the
estimated marginal means. Notice that in most cases the conditioning treatments did not affect
the relative abundance of bacterial markers compared to the baseline soils.

Extended Data Fig. 7. Genetic variation among bacterial lineages driven by long-term water
legacy remains stable despite acute drought/water perturbations and host conditioning.
Filtered shotgun metagenomic reads were aligned to the reference genomes of 33 selected taxa,
including 22 identified bacterial markers and 11 additional abundant and prevalent taxa. Genetic
distances between these bacterial lineages were calculated based on identified single nucleotide
polymorphisms (SNPs). Principal Coordinate Analysis (PCoA) plots were generated and color to
depict **a.** acute drought and well-watered treatments, and **b.** host and no-host conditioning
perturbations. Phi values, derived from Analysis of Molecular Variance (AMOVA), indicate the
genetic variance between groups, while the *p*-value reflects the significance of the AMOVA results
based on permutation testing. Fixation Index (F_{ST}) values were used to measure the degree of
genetic differentiation between groups. The variance explained by each axis is displayed on the
PCoA plots. **c.** A Manhattan plot illustrated significant SNPs linked to the wet and dry soil legacies,
derived from the genetic-environment association (GEA) analysis. This analysis was conducted
using a general linear model and significant associations were identified using the permutation
method. In the Manhattan plot, the x-axis represents the SNP positions along the genomes of the

selected bacterial species, while the y-axis displays the -log10 p-values from the association
model. A horizontal line represents the statistical significance threshold, determined by the
permutation test. SNPs exceeding this threshold (highlighted in red) were significantly associated
with soil legacy effects. Bacterial taxa with fewer than 1,000 high-quality biallelic SNPs after
filtering, or with no significant SNPs detected from the GEA analysis, are excluded from the plot.

Extended Data Fig. 8. Precipitation legacy effects on microbial community function are
robust to short-term (five months) acute perturbations. a. The heatmap illustrates enriched
or depleted Gene Ontology (GO) categories in soils with wet or dry legacies, subjected to drought

or well-watered treatments, in comparison to soils with a wet legacy (baseline). Gene enrichment
analysis was conducted using a generalized linear model, followed by GO classification.
Significantly enriched and depleted GO categories were identified using an adjusted p-value
threshold of < 0.05. Colors (red for enrichment, green for depletion) represent enrichment scores,
calculated based on square root-transformed delta rank values of the GO categories. The
heatmap is clustered by soil treatments and GO terms. **b**. Canonical analysis of principal
coordinates (CAP) showing the projected metatranscriptome differences across the conditioning
treatments: (top) baseline soil legacy, baseline exposed to well-watered or drought treatment, and
(bottom) host or no host treatment. The bar on the left describes the percentage of the variance
explained by the experimental variables. **c**. Phylogram showing the main bacterial phyla
functionally active across soils with different water legacy subjected to drought or well-watered
treatments. **d-e**. Heatmaps showing enrichment in functionally active bacterial (**d**) families and (**e**)
phyla across soils with different water legacy and exposed to drought or well-watered treatments
compared to soils with wet legacy. Heatmaps were colored based on $\log_2$ fold changes derived
from a generalised linear model contrasting the abundance of each family in a given treatment
against soils with wet legacy. Tiles outlined in black denote statistically significant enrichment
(red) or depletion (blue) (q < 0.05) with a $\log_2$ fold change > ±2 with respect to the soils with wet
legacy. Heatmaps were clustered based on taxonomic classification (tree on the left). **f**. The
heatmap displays biological processes that are enriched or depleted based on metatranscriptomic
analysis of soils with wet or dry legacies, subjected to drought or well-watered treatments,
compared to soils with a wet legacy (baseline). Gene enrichment analysis was performed using
a generalized linear model, followed by GO classification. Significantly enriched and depleted
Gene Ontology (GO) categories were identified with an adjusted p-value threshold of < 0.05.
Colors (red for enrichment and green for depletion) indicate enrichment scores, calculated from
square root-transformed delta rank values of the GO categories. The heatmap is clustered by soil
treatments and GO terms.

1746
1747
1748 **Extended Data Figure 9. Test phase gamagrass and maize drought treatments effectively**
1749 **reduce plant growth**. Test phase drought treatment significantly reduced gamagrass shoot
1750 height (cm) (**a**) and maize shoot height (cm) (**b**).

1781
1782 **Extended Data Figure 10. Plant trait feature selection. a-b.** Gamagrass and **c-d.** maize traits
1783 important in explaining legacy effects were ranked using a random forest model. The top ten traits
1784 were tested for significant correlations (≥ 0.7).

1794
Extended Data Figure 11. Microbiome legacy impacts gamagrass phenotypic drought
response. a. A constrained redundancy analysis on the top non-correlated traits found that legacy
explains 5.0% of gamagrass phenotypic response to acute drought. **b-j**. Assessment of individual
traits indicates that microbiome drought legacy significantly improved gamagrass performance
under drought. Points are EMMs and error bars represent the standard error, and significant p-
values (≤0.05) are bolded.

Extended Data Figure 12. Contributions of legacy-sensitive *Tripsacum dactyloides* genes
to NMDS axes that mediate drought responses. **a**. Histograms showing the distributions of
gene loadings onto both axes. The top 5% of genes with the strongest positive and negative
loadings onto each axis are in red. **b**. Breakdown of the number of genes with positive/negative
loadings onto each axis. **c**. Lists of genes comprising the 5% tails of the distributions in **a**. Detailed
information about the expression responses of these genes is available in Supplementary Table
S3. 'Gene_set' corresponds to the patterns of response to microbiome legacy presented in Figure
4a-b.

1823

Extended Data Figure 13. Test phase water and legacy have a large impact on ASV abundances in maize, but not gamagrass. a. Gamagrass root bacterial and **b.** maize root bacterial centered-log ratio transformed ASV counts were individually fit to a linear model to identify ASVs that were differentially abundant across test phase water treatments in the context of precipitation legacy. Dark green represents ASV enrichment and light green to white represents depletion. Multi-colored squares indicate phylum-level taxonomic assignments, which do not have an obvious phylogenetic pattern.

1831
Extended Data Figure 14. Drought legacy of microbial inocula alters the transcriptional
response to drought in the maize crown root. a. 23 genes were up-regulated in plants
inoculated with microbiota from a low-precipitation region, relative to plants inoculated with
microbiota from a high-precipitation region. **b.** The sets of genes that responded to the main
effects of inoculum legacy and Test Phase drought stress had minimal overlap with each other or
with the set of genes that were sensitive to the interaction between the two. **c.** In total, 109 maize
genes responded to drought in a manner that was dependent on the drought legacy of the soil
microbiota, regardless of the inoculum's treatment during the conditioning phase. For illustration
purposes, only annotated genes with |log$_2$FoldChange| > 1 in at least one microbial context are
shown here; the full list is available in Supplementary Table S3. Each pair of points shows one
gene; the relative position of the two points illustrates how the gene's expression changed in
response to drought stress during the Test Phase, depending on whether the plant had been
inoculated with microbiota derived from a low-precipitation or dry-precipitation environment.
Genes are grouped into sets according to the pattern of how inoculum legacy altered their drought
responses. Note: the names of these gene sets are not meant to correspond to the names of the
T. dactyloides gene sets shown in Figure 4b; in each species, Gene set I contains the most genes,
Gene set II contains the next most, and so on.

1855

Extended Data Figure 15. Microbiome Legacy had minimal impact on maize phenotypic drought response. a. A constrained redundancy analysis on the top non-correlated traits found that legacy only explains 2.0% of maize phenotypic response to acute drought. **b-j.** Assessment of individual traits indicates that microbiome drought legacy did not significantly improve maize performance under drought, but did impact several mineral nutrient concentrations. Points are EMMs and error bars represent the standard error, and significant *p*-values (≤ 0.05) are bolded.

Supplementary Table Legends.

Supplementary Table S1. List of identified regions in the United States and Cabo Verde representing precipitation gradients. The soils used in this work were collected from these regions. The table also shows the coordinates associated with these regions, the state, municipalities, and country where these regions are located.

Supplementary Table S2. List of bacterial markers of soil water legacy identified in this work using soils from the two precipitation gradients from Kansas and Santiago. In addition to the bacterial taxa ID (taxID), the table also shows taxonomical attributes (Family, Genus, label Family, and Label genus) of the bacterial markers identified. The column F shows the direction of the marker enrichment: High to low is the marker is enriched in the soils with high precipitation levels as compared to soils with low precipitation levels and vice versa. The localization of the precipitation gradient is shown in column G.

Supplementary Table S3. Differentially expressed genes of test phase gamagrass and maize roots. Tab one provides information for interpreting the column names and four other tabs. Tab two lists gamagrass genes that are significantly up or down-regulated based on the main effects of microbial inoculum legacy. Tab three lists gamagrass genes that are significantly up or down-regulated based on the interaction of test phase water treatment and microbial inoculum legacy. Tab four lists maize genes that are significantly up or down-regulated based on the main effects of microbial inoculum legacy. Tab five lists maize genes that are significantly up or down-regulated based on the interaction of test phase water treatment and microbial inoculum legacy.

Supplementary Table S4. Test phase plant phenotypic data and microbiome metadata. Tab one provides information for interpreting the column names. Tab two provides the measurements for all plants used in the plant phenotypic analyses, including both Gamagrass and maize. Tab three provides the measurements for the plant samples used in the root bacterial microbiome data analyses, with sequencing data that passed all qualify checks and filtering.

Supplementary Table S5. Differentially abundant test phase gamagrass and maize root microbiome ASVs. Tab one provides information for interpreting the column names. Tab two lists the gamagrass root microbiome ASVs, including their full taxonomic assignment that where significantly differentially abundant based on legacy in the context of test phase water treatment. Tab three lists the maize root microbiome ASVs, including their full taxonomic assignment that where significantly differentially abundant based on legacy in the context of test phase water treatment.