Title

Genome-wide Search for Gene Mutations likely Conferring Insecticide Resistance in the Common Bed Bug, *Cimex lectularius*

Authors

Kouhei Toga [1], Fumiko Kimoto[2], Hiroki Fujii[2] and Hidemasa Bono [3,*]

Affiliations

1 . Laboratory of BioDX, PtBio Co-Creation Research Center, Genome Editing Innovation Center, Hiroshima University, 3-10-23 Kagamiyama, Higashi-Hiroshima city, Hiroshima, 739-0046, Japan
2 . Research & Development Division, Fumakilla Limited, 1-11-13 Umehara, Hatsukaichi city, Hiroshima, 739-0494, Japan
3 . Laboratory of Genome Informatics, Graduate School of Integrated Sciences for Life, Hiroshima University, 3-10-23 Kagamiyama, Higashi-Hiroshima city, Hiroshima, 739-0046, Japan
*Correspondence: bonohu@hiroshima-u.ac.jp; Tel.: +81 82 424 4013

Simple Summary: Bedbugs have expanded globally over the past two decades, causing several health risks. Mutations in their genes allow bed bugs to develop insecticide resistance. However, the extent to which gene mutations exist in the bug genome remains largely unknown because the genomes of resistant strains have not been determined. We accurately sequenced the genomes of both susceptible and resistant strains and compared the gene sequences between the two strains. Several genes with resistance-specific mutations have been identified. These mutations can alter gene function and lead to insecticide resistance.

Abstract: Insecticide resistance in the bedbug *Cimex lectularius* is poorly understood due to the lack of genome sequences for resistant strains. In Japan, we identified a resistant strain of *C. lectularius* that exhibits a higher pyrethroid resistance ratio compared to many previously discovered strains. We sequenced genomes of the pyrethroid resistant and susceptible strains using long-read sequencing, resulting in the construction of highly contiguous genomes (N50 of resistant strain: 2.1 Mb and N50 of susceptible strain: 1.5 Mb). Gene prediction was performed by BRAKER3 and Functional annotation was performed by Fanflow4insects workflow. Next, we compared their amino acid sequences to identify gene mutations, identifying 729 mutated transcripts that were specific to the resistant strain. Among them, those defined previously as resistance genes were included. Additionally, enrichment analysis implicated DNA damage response, cell cycle regulation, insulin metabolism, and lysosomes in the development of pyrethroid resistance. Genome editing of these genes can provide insights into the evolution and mechanisms of insecticide resistance. This study expanded the target genes to monitor allele distribution and frequency changes, which will likely contribute to the assessment of resistance levels. These findings highlight the potential of genome-wide approaches to understand insecticide resistance in bed bugs.

Keywords: insecticide resistance; *Cimex lectularius*; genome sequencing; genome editing

1. Introduction

Over the past 20 years, bed bugs (*Cimex lectularius* and *Cimex hemipterus*) have spread rapidly globally [1,2]. Bedbugs are obligate hematophagous insects [3,4]. Bedbug bites can cause pruritic rashes [5], folliculitis, and cellulitis due to secondary infections [6]. Although approximately 40 pathogens have been detected in bedbugs from natural populations, no evidence exists that they have been transmitted to humans [7]. Histamine, an aggregated pheromone released by bedbugs, is deposited in house dust and may pose health risks [8,9]

Histamine poisoning can be occurred when ingestion of histamine included in food, leading to rash, urticaria, and edema [10]. Dermatitis occurs by histamine exposure to skin, primally in atopic patients [11].

Insecticide resistance is believed to be a factor contributing to the recent resurgence of bed bugs [10,11]. Infestation was rare in the 1960s because of the use of dichlorodiphenyltrichloroethane (DDT) or other synthetic organic insecticides, such as pyrethroids [2,12]. More recently, spraying with mixtures of pyrethroids and neonicotinoids has been shown to be effective for controlling bed bugs [2,13]. The physiological mechanisms conferring insecticide resistance can be divided into several types: metabolic, penetration, and knockdown resistance [14]. The major groups of enzymes associated with metabolic resistance include cytochrome P450s, glutathione S-transferase (GST), and ATP-binding cassette (ABC) transporters [15]. Cuticle thickening has been implicated in penetration resistance [16], and RNA-seq analysis confirmed that several genes encoding P450s, cuticle proteins (CPRs), and ABC transporters were upregulated in pyrethroid-resistant strains [17–19]. Target site insensitivity causes knockdown resistance. To date, amino acid mutations in voltage-gated sodium channels have been observed in the common bed bug *C. lectularius* with pyrethroid resistance [18,20]. Amino acid substitutions from "L to I" or "V to L" in voltage-gated sodium channel have been observed in *C. lectularius* as well as in other insects [21,22]. Recently, resistant strains of *C. lectularius* for organophosphates and carbamates have been found in Japan, and their acetylcholinesterase (AChE, target genes of these two insecticides) possess mutations [23]. However, the genomic information involved in the development of insecticide resistance relies heavily on information from other insects.

Genome sequencing of the common bed bug, *C. lectularius* has revealed gene sets related to its ecology [24,25]. Previous studies have used short-read sequencing to identify SNPs associated with pyrethroid resistance [26,27]. These studies have identified multiple genomic regions that are likely involved in pyrethroid resistance. However, no study has accurately determined the whole-genome sequence of the resistant strain, and the full extent of mutations in resistant strains remains unknown. In Japan, inquiries into bedbugs have increased since 2000 [28]. In this study, we constructed highly contiguous genomes of susceptible and resistant strains isolated from Japan using long-read sequencing. Next, gene prediction and functional annotation of the assembled genome were performed. We identified amino acid mutations that are potentially related to pyrethroid resistance. This study expanded the target genes to monitor allele distribution and frequency changes, which will likely contribute to the assessment of resistance levels. These findings could highlight the potential of genome-wide approaches to understand insecticide resistance in bed bugs.

2. Materials and Methods

2.1. Insects

Susceptible strains (Teikyo-university strain) of *C. lectularius* were collected from fields at the Department of Medical Zoology, Research Institute of Endemics, Nagasaki University (Isahaya City, Nagasaki, Japan) by Prof. N. Omori on May 22, 1956 [29]. This strain was maintained at Nagasaki University and Teikyo University and bred and maintained at the Japan Environmental Sanitation Center (Kawasaki City, Kanagawa, Japan) from approximately 1972. This strain was considered insecticide susceptible and was kindly provided to Fumakilla Limited (Chiyoda-ku, Tokyo, Japan) in 2011. The resistant strain (Hiroshima strain) was collected from a hotel in Hiroshima City (Hiroshima, Japan) in July 2010 and maintained by Fumakilla Limited. Bed bugs were reared by feeding mouse blood. All samples used in this study were collected in Japan. Therefore, the international ABS (Access and Benefit-Sharing)

framework under the Convention on Biological Diversity and the Nagoya Protocol is not applicable.

2.2. Insecticide Efficacy Tests

Technical-grade insecticides were used to test the susceptibility. Permethrin (Lot. No. 191001, 97.1%, Sumitomo Chemical Co., Ltd., Tokyo, Japan) is a pyrethroid insecticide, and dinotefuran (Lot. No. K16530422, 99.1%; Mitsui Chemicals Crop & Life Solutions, Inc., Tokyo, Japan) is a neonicotinoid insecticide.

Simplified bioassays were conducted to evaluate the susceptibility of adult *C. lectularius* to the representative insecticides. Bioassays were performed on adults 7–10 days after blood feeding. Adult *C. lectularius* (male: females = 1:1) were held on a piece of tape applied to a white paper. Following the topical application method described in WHO [30], an acetone solution (0.4125 µL) of the test insecticide was topically applied to the dorsal mesothorax of each test *C. lectularius* using a hand microapplicator (Burkard Manufacturing Co., Ltd., Rickmansworth, UK). The control group was administered only acetone. Two insecticides, Permethrin and Dinotefuran, were tested using six doses in insecticide efficacy test, and each replicate contained at least 20 insects. For Permethrin, the Teikyo-u strain was exposed to doses of 0.000391, 0.000782, 0.00156, 0.00313, 0.00625, and 0.0125 µg/individual. Higher doses of 15.6, 31.3, 62.5, 125, 250, and 500 µg/individual were applied to the Hiroshima strain. For Dinotefuran, identical doses of 0.00754, 0.0151, 0.0301, 0.0603, 0.121, and 0.241 µg/individual were used for both susceptible and resistant strains. The treated insects were kept in a plastic cup, the bottom of which was lined with filter paper. Mortality was observed 24 h after treatment, and the half-lethal dose (LD$_{50}$) was calculated using the Bliss' probit method [31]. A moribund insect that was unable to walk was considered dead. The resistance ratio (RR) was calculated by dividing the LD$_{50}$ of the resistant strain by that of the susceptible strain. Fieller's equation was used for calculating the 95% confidence intervals [32].

2.3. DNA extraction and genome sequencing

The genomic DNA of *C. lectularius* was extracted from the whole body of an adult male using a MagAttract HMW DNA Kit (Qiagen Co. Ltd., Valencia, CA, USA). *C.lectularius* genomic DNA was fragmented to a target size of 10-15 kb using the g-TUBE (Covaris, Woburn, MA, USA). Fragmented DNA was then purified using SMRTbell cleanup beads (PacBio, Menlo Park, CA, USA). A sequence library was prepared using the SMRTbell Prep Kit 3.0 (PacBio),and were loaded on the Sequel II sequencing plate 2.0 with on-plate loading concentrations 90 pM. The libraries were sequenced using Sequel IIe (PacBio). Samples were sequenced using the SMRTCell 8 M tray on the Sequel IIe platform with 30h movie time.

2.4. De novo assembly and quality assessment

To remove mouse genome contamination from HiFi reads of *C. lectularius,* HiFi reads were mapped to the mouse genome sequences (GRCm39) of GENCODE (accessed on January 10, 2024) using Minimap2 [33], and the unmapped reads were used in the subsequent assembly processes. HiFi reads generated from PacBio sequencers were assembled using Hifiasm v0.19.7 using the '-primary' option [34]. Quast v 5.2.0 was used to assess the assembly statistics [35]. Gene set completeness was assessed using the Benchmark of Universal Single-Copy Orthologs (BUSCOs) v5.2.2, with '-l insecta_odb10 (2020-09-10)' or with '-l hemiptera_odb10' (2024-01-08). Merqury v1.3 and Meryl db (k=21) were used for k-mer-based validation of the genome assembly [36].

Minimap2 was used to align the genome sequences between strains with '-cx asm5' option. Dot plots were generated using the dotPlotly program with '-s -t -m 500 -q 500000 -l'

(https://github.com/tpoorten/dotPlotly, accessed on March 8, 2024) to visualize genomic differences.

2.5. RNA sequencing

Public RNA sequencing data were searched from NCBI using the keywords *Cimex lectularius*. Metadata was downloaded using NCBI SRA Run Selector (https://swww.ncbi.nlm.nih.gov/Traces/study/, accessed on 26 Feb 2024) (Table S1).

Total RNA was extracted from one adult that was used as a control in 2.2 Insecticide Efficacy Tests using RNeasy Mini Kit (Qiagen, Hilden, Germany). The extraction was performed in triplicate for both the susceptible and resistant strains. Libraries were prepared using TruSeq stranded mRNA (Illumina, San Diego, CA, USA) and sequenced using NovaSeqX (Illumina), generating 150 bp pair-end reads. Accession numbers are listed in Table S2.

SRA Toolkit (v3.0.6) (https://github.com/ncbi/sra-tools) was used for SRA retrieval and conversion to FASTQ files. Trim Galore! (v0.6.10) [37] was run for trimming and quality control with option '--paired' options. Low-quality base calls were trimmed using Cutadapt (v3.4) [38] included Trim Galore!. RNA-Seq reads were mapped to the genome assembly using HISAT2 (v2.2.1) with '-q --dta' option [39]. The SAM files were converted into BAM files using SAM tools [40]. The obtained BAM files were used for gene prediction.

2.6. Gene prediction and functional annotation

Repeat regions were identified and soft-masked using RepeatModeler v 2.0.3 and RepeatMasker v4.1.6, in Dfam TEtools v1.88 (https://github.com/Dfam-consortium/TETools). Gene prediction was performed by BRAKER3 [41] using the soft-masked genome.Besides BAM files in which RNA-Seq reads were mapped to genome assembly, OrthoDB protein sets of Arthropoda [42] (https://bioinf.uni-greifswald.de/bioinf/partitioned_odb11/, accessed on June 17, 2023) were used as extrinsic evidence when performing BRAKER3. The transcript and amino acid sequences were extracted from the GTF files using GFFread v0.12.7 [43]. Gene set completeness was assessed using BUSCO with '-l insecta_odb' (2020-09-10) and '-l hemiptera_odb10' (2024-01-08). The functional annotation workflow, Fanflow4Insects [44], was modified and used against the reference protein datasets of *Homo sapiens, Mus musculus, Caenorhabditis elegans, Drosophila melanogaster,* hemipteran insects (*Cimex lectularius* (current reference genome Clec2.1), *Acyrthosiphon pisum, Daktulosphaira vitifoliae, Homalodisca vitripennis, Nilaparvata lugens, Rhodnius prolixus*), and UniProtKB/Swiss-Prot. Reference protein datasets were retrieved from Ensembl or Ensembl Metazoa (accessed on May 5, 2024). Sequence similarities were searched using GGSEARCH v36.3.8g in the FASTA package (https://fasta.bioch.virginia.edu/; accessed July 19, 2023). A domain search for the Pfam database v35.0 was performed using HMMSCAN in HMMER v3.3.2 (http://hmmer.org/).

Gene models of susceptible strains were transferred into the genome assembly of resistant strains using Liftoff [45]. The transcripts with different CDS between susceptible and resistant strains were identified by searching for 'matches_ref_protein: False' in the GFF files generated by Liftoff. Amino acid sequences of corresponding transcript IDs between susceptible and resistant strains were aligned using MAFFT v 7.525 with '--auto' option [46]. The protein sequence of Clec2.1 were also included in the MAFFT alignment. The corresponding amino acid sequences of Clec2.1 were identified from the results of Fanflow4Insects, as mentioned above. Alignment files were imported using the AlignIO module in Biopython 1.83, and only transcripts with resistant-specific mutations were selected. On the list, if transcripts were missing start and/or stop codons after liftoff, the correctness of the missing codons was manually verified by identifying contig fragmentation or liftoff mistakes. Differences in amino acid sites between the susceptible and resistant strains were visualized on a Tree [47]. To characterize the

list of transcripts with resistance-specific mutations, enrichment analysis was performed using Metascape (https://metascape.org/, accessed on June 24, 2024), with default settings [48].

3. Results

3.1. Lethal effects of pyrethroid for the susceptible and resistant strains

We evaluated the resistance status of the Hiroshima strain. (Table 1). LD_{50} of susceptible strain (Teikyo university strain) showed 0.002 μg of permethrin per individual, whereas LD_{50} of Hiroshima strain showed 40.1 μg of permethrin per individual. The resistance ratio (RR) was 19,859, which was higher than that of many previously discovered strains [49]. Hereafter, we refer to the Teikyo University strain as the susceptible strain, and the Hiroshima strain as the resistant strain.

Table 1. The LD_{50} value and resistance ratio of each strain to two insecticides.

Insecticide	Strain	Number of specimens	Slope ± SE	LD_{50} (μg/individual)	LD_{50} 95%CI	df	P-value	RR	RR 95%C
Permethrin	Teikyo-u	120	3.1 ± 1.08	0.0020	0.0015 - 0.0028	4	0.77	-	-
	Hiroshima	120	1.3 ± 0.39	40.1	24.4 - 66.0	4	0.73	19,859	10,559 38,99

SE : Standard error

LD_{50} : The dose of insecticide that is lethal to 50 % of the test specimens

df : degree of freedom

RR : LD_{50} (Hiroshima strain)/LD_{50} (Teikyo-u strain)
CI : Confidence Interval

3.2. De novo genome assembly

We generated primary contigs with genome sizes of 644,683,587 bp and 614,893,704 bp for the susceptible and resistant strains, respectively (Table 2). Indicators for genome contiguity were of higher quality than those of the current reference genome Clec2.1 (Table 2). The contig N50 lengths for the susceptible and resistant strains were 1,530,193 bp and 2,132,290 bp. These assembly metrics were of higher quality than those of the current reference genome Clec2.1 (Table 2). The number of contigs in both strains was lower than that in Clec2.1 reference genome. The BUSCO completeness in both strains was comparable to that of the Clec2.1 reference genome. K-mer-based evaluation demonstrated high base accuracy and completeness of primary contigs in susceptible and resistant strains The susceptible strain showed a quality value (QV) of 57.0 and completeness of 97.8%, while resistant strain had QV of 56.9 and completeness of 94.9%. The K-mer spectrum analysis revealed two peaks in both strains, indicating the construction of a diploid genome assembly (Figure 1a and b). No counts were observed that could be considered sequence errors. We examined the genomic differences between susceptible and resistant strains using dot-plot visualization (Figure 1c). Linear plots confirmed the high similarity between the susceptible and resistant strains.

Table 2. Comparison of assembly statistics with the current reference genome.

Parameter	Susceptible	Resistant	Clec2.1

	Length(bp)	644,683,587	614,893,704	510,848,842
	contig N50(bp)	1,530,193	2,132,290	0.5524
Assembly statistics	contig L50	124	85	285
	No. of contigs	1324	1092	2,868
	Largest contig	12,214,012	9,042,003	6,600,950
	GC%	35.08	35.19	34.82
BUSCO results (OrthoDB v10 insect database)	Complete [single, duplicated]	99.3%[98.1%,1.2%]	99.2%[97.8%,1.4%]	99.3% [98.1%, 1.2%]
	Fragmented	0.3%	0.2%	0.3%
	Missing	0.4%	0.6%	0.4%
BUSCO results (OrthoDB v10 hemiptera database)	Complete (single, duplicated)	99.4%[97.8% ,1.6%]	99.1%[97.4%, 1.7%]	99.6% [98.2%, 1.4%]
	Fragmented	0.2%	0.2%	0.1%
	Missing	0.4%	0.7%	0.3%

3.3. Gene prediction and functional annotation

Because repetitive sequences may affect gene prediction accuracy [50], these sequences in the genome assembly were searched using RepeatModeler and RepeatMasker. In both strains, approximately 50% of the genomic regions contained repeat sequences (Table S3, shown in bases masked). These values are slightly higher than those of the current reference genome, Clec 2.1. BRAKER3 predicted 13116 and 13104 protein-coding genes in the susceptible and resistant strains, respectively (Table 3). Gene annotations and protein sequences are available at figsShare (https://doi.org/10.6084/m9.figshare.26362135.v2). The BUSCO completeness values for both strains were comparable to Clec2.1. Gene annotations of the susceptible strains were transferred to the resistant strain using Liftoff [45], identifying the corresponding genes between both strains (12724 protein-coding genes and 20151 transcripts) (Table 3). The BUSCO completeness of the resistant strains after liftoff did not decline substantially, indicating that the transfer of gene predictions was successful (Table 3). Functional annotation was added to the protein-coding genes of the susceptible strains using Fanflow4Insects, which included GGSEARCH and HMMSCAN. The number of gene hits for each reference protein is shown in Table 4. Of 13116 genes in susceptible strains, 99% protein-coding genes (13010/13116) were successfully annotated by GGSEARCH and HMMSCAN. The genes that could not be annotated were classified as hypothetical proteins (Table 4). The complete results of Fanflow4Insects are presented in Table S4. We performed a BUSCO analysis using databases for glires and primates. If significant contamination were present in the assembly constructed in this study, the BUSCO completeness scores would be higher. However, the BUSCO analysis of assemblies from

susceptible strains showed completeness values of 4.9% for glires and 3.5% for primates. These results suggest that the assemblies of susceptible strains are not composed of genomes derived from either mice or humans.

Table 3. Statistics of protein-coding gene prediction.

BRAKER and Liftoff		Susceptible	Resistant	Clec2.1*	Resistant (after liftoff)
	The number of protein-coding gene	13116	13104	11949	12724
	The number of transcript	20928	20890	24194	20151
BUSCO results (OrthoDB v10 insect database)	Complete [single, duplicated]	98.4% [69.1%,29.3%]	98.3%[68.0% ,30.3%]	99.7%[65.9%,3 3.8%]	97.3%[68.5% ,28.8%]
	Fragmented	0.1%	0.3%	0.3%	0.4%
	Missing	1.5%	1.4%	0.0%	2.3%
BUSCO results (OrthoDB v10 hemiptera database)	Complete (single, duplicated)	98.1%[65.3%, 32.8%]	97.8%[64.9% ,32.9%]	99.5%[60.3%,3 9.2%]	96.9%[65.7% ,31.2%]
	Fragmented	0.0%	0.1%	0.0%	0.1%
	Missing	1.9%	2.1%	0.5%	3.0%

* The data of Clec2.1 were referred to Ensembl statistics and protein fasta file.

Table 4. Summary of functional annotation results.

Annotation category	Reference (protein ID)	Number of gene hits
Gene annotation by top hit (GGSEARCH)	*Homo_sapiens*	9913
	Mus_musculus	9738
	Caenorhabditis_elegans	8323
	Drosophila_melanogaster	9585
	Cimex_lectularius (Clec2.1)	11933
	Acyrthosiphon_pisum	10269
	Daktulosphaira_vitifoliae	10151
	Homalodisca_vitripennis	10481
	Nilaparvata_lugens	10578
	Rhodnius_prolixus	10098
	UniProtKB/Swiss-Prot	10828
	at least one of the above	12993
Genes annotated only by HMMSCAN	Pfam domain database	17
All genes annotated at protein level		13010
Hypothetical protein	No hit	106
Total		13116

3.4. The search for mutated genes related to insecticide resistance

As previous report suggested, point mutations are related to the acquisition of pyrethroid resistance in bed bugs [21]. These genes can be identified on a genome-wide scale by comparing the protein sequences of susceptible and resistant strains. Liftoff revealed 3938 transcripts with amino acid mismatches between susceptible and resistant strains. These amino acid sequences were aligned with those of Clec2.1, using MAFFT, and the alignments among the three strains were compared. Clec2.1 genome comes from a susceptible strain [24]. The MAFFT alignment revealed 729 transcripts with resistance-specific mutations (Table S5). We investigated the number of mutation sites in the amino acid sequence of each transcript (Figure 2A, Table S5). The majority of the transcripts (496 of 729) had one mutation in their amino acid sequences. *laminin subunit beta 1* had the largest number of mutated sites (54). To verify the reliability of the gene list, we searched for known or candidate genes with resistance-specific mutations, including sodium voltage-gated channels, AChE, Gamma-aminobutyric acid (GABA) receptors, nicotinic acetylcholine receptors (nAChR), P450s, ABC transporters, GSTs, and CPRs) [14]. Transcripts with resistance-specific mutations were detected in all these genes, except for the GABA receptor (Figure 2B-G). No studies have identified mutations in *C. lectularius* nAChR, a target gene of neonicotinoid insecticides. Although we determined the resistance ratio of the neonicotinoid insecticide (dinotefuran) between the susceptible and resistant strains (Table S6), the Hiroshima strain did not show neonicotinoid resistance (resistance ratio: 1.11).

Enrichment analysis was performed on 729 transcripts to characterize the genes (Figure 3, Table S7, and Table S8). Enrichment analysis using Metascape was conducted using *Homo sapiens* gene ID or *D. melanogaster* gene ID, which were described in Table S5. The molecular functions of a large number of genes are evolutionarily conserved among species, and there is abundant and detailed information on human gene function. For this reason, we used

human gene IDs for enrichment analysis to infer molecular functions in insects. Using *Homo sapiens* gene ID, we identified several GO terms potentially related to the development of insecticide resistance, as described below. DNA metabolic process, mitotic cell cycle process, insulin metabolic process and Lysosome were enriched (-log10 (p-value) > 6) (Figure 3A). Gene lists associated with the GO terms are shown in Table S7. Forty-two genes were included in the "DNA metabolic process" category. Metascape clustered several similar ontologies, and the summarized results are shown in the GroupID column in Table S7. The "DNA metabolic process" category included multiple terms related to DNA damage and repair, such as "DNA replication," "DNA repair," "DNA damage response," "double-strand break repair," and "recombinational repair" (Table S7). Fifty -two genes were clustered under GO term "mitotic cell cycle process" (Table S7). Among these genes, CCNB2 (cyclin B) and ORC1 are activated during M and G1 phases of the cell cycle, respectively [51,52]. Nine genes clustered under the GO term "insulin metabolic process" (Table S7). Among these genes, CPE (Carboxypeptidase E) is involved in the proteolytic processing of proinsulin peptides in humans [53]. Thirteen genes were clustered under the GO term "Lysosome" (Table S7). Intracellular proteases respond to insecticide stress in resistant strains to maintain cellular health [54]. As expected, the cathepsin genes (CTSB and CTSD) were included among the 13 genes. Additionally, components of lysosomes, such as SLC11A2, were included. SLC11A2 is a divalent metal transporter that is a component of lysosomes [55].

When using *D. melanogaster* gene ID, "Lysosome" was enriched, similar to the results obtained using human gene IDs (Figure 3B, Table S8). GO terms included cathepsin. Tetraspanins (Tsp29Fa and Tsp42Ee) were also identified. The enriched GO term "glucose transmembrane transport" was related to the "insulin metabolic process" enriched when using human gene ID, as these were associated with nutrient response [56].

4. Discussion

We successfully constructed a highly contiguous genome assembly of susceptible and resistant strains of *C. lectularius* and identified mutations that may be related to the development of insecticide resistance. The present study confirmed the mutations conferring pyrethroid resistance in the amino acids of the sodium voltage-gated channel, as expected from previous reports in *C. lectularius*. We confirmed mutations in other target genes that confer neonicotinoid or OPs resistance (AChE and nAChR). Although mutations in AChE have been found in *Cimex* species [23], no reports have described mutations in the nAChR. However, we were unable to confirm neonicotinoid resistance in this study. Mutations found in nAChR may not be associated with neonicotinoid reception. Genes related to metabolic resistance (CYP450 and ABC transporters) or penetration resistance (CPRs) were highly expressed in resistant strains. To the best of our knowledge, this is the first study to report mutations in these genes.

A total of 729 transcripts with resistance-specific mutations were identified. We found "DNA metabolic process," "Lysosome," "Insulin metabolic process" and "glucose transmembrane transport" and "Mitotic cell cycle process" as enriched GO terms in the 729 transcripts. These biological processes may be related to the physiological adaptation of the resistant strains, as discussed below.

4.1. DNA metabolic process

Enrichment analysis revealed that resistance-specific mutations were present in many genes associated with DNA damage, repair, and replication. Bifenthrin, a pyrethroid insecticide, cause the DNA damage in Sf9 cells [57]. In *Anopheles coluzzii*, the expression of DNA repair-related genes increased after exposure to pyrethroids [58]. Mutations identified in this study may promote DNA damage repair in resistant strains, leading to adaptations to insecticide exposure.

4.2. Lysosome

In the house fly *Musca domestica* and red flour beetle *Tribolium castaneum*, lysosomal protease activity generally increases in response to exposure to pyrethroids and DDT [54]. Actually, protease functioning in the lysosome, cathepsin B, was present in the result of enrichment analysis. Mutations were also observed in the lysosomal components of tetraspanin (*Tsp29Fa* and *Tsp42Ee*). In *D. melanogaster*, Tsp42Ee is primarily located in retina [59], and its mutation leads to light-induced retinal degeneration. Whether the tetraspanin mutation found in this study is related to insecticide resistance through the visual system may provide a target for understanding the ecology and evolution of resistant strains.

4.3. Insulin metabolic process and glucose transmembrane transport

Enrichment analysis Insulin signaling is conserved between mammals and insects, and regulates circulating sugar levels [56]. Therefore, we discuss insulin metabolic processes and glucose transmembrane transport in the same section.

Mutations were present in glucose and trehalose transporters (Tret1-1, pippin, and nebu) (Table S8). *Tret1-1* expression is upregulated in response to starvation in *D. melanogaster* and *T. castaneum* [60–62], leading to the efficient uptake of trehalose and glucose. *Pippin* regulate the selective transport of carbohydrates into the brain [63], but its response to starvation remains unknown. Mutations in these transporters may contribute to resistance to starvation.

When the human gene ID was used for enrichment analysis, the insulin metabolic process was enriched. Components of insulin signaling are regulated in response to starvation in *D. melanogaster* [56]. Therefore, "insulin metabolic process" and "glucose transmembrane transport" may be associated with each other. Additionally, in *D. melanogaster*, 114 mutations in insulin-signaling genes have been found in DDT-resistant strains [64]. In cotton leafworm *Spodoptera litura*, exposure of azadirachtin, a [tetranortriterpenoid] found in neem [tree] (*Azadirachta indica*), deactivate insulin signaling [65]. Oral treatment of pyridalyl upregulated the expression of insulin receptor in the olive fruit fly *Bacterocera oleae* [66]. These results suggest the connection between insulin signaling and insecticide exposure. In conclusion, modifying insulin signaling may be a common feature in conferring insecticide resistance in insects.

4.4. Mitotic Cell Cycle Process

DNA damage is monitored during cell cycle progression, and the cell cycle is arrested until the DNA damage is repaired [67]. Mutations were observed in 52 genes associated with the mitotic cell cycle. Among these, ORC1 has been extensively studied with respect to cell cycle regulation [68]. ORC1 plays a central role in the assembly of the pre-replication complex to initiate DNA replication and its activation is regulated during the cell cycle [69]. Mutations observed in insecticide-resistant strains may allow strict regulation of the cell cycle, preventing daughter cells from inheriting damaged DNA in response to insecticide exposure.

4.5. The relationships among GO terms enriched in this study

The GO terms identified in this study (DNA metabolic process, insulin metabolic process, glucose transmembrane transport, lysosomes, and mitotic cell cycle process) may be related to cell cycle regulation. DNA metabolic processes play crucial roles in cell-cycle progression and arrest of cell cycle [67]. Cell cycle progression depends on sugars such as glucose [70]. Insulin signaling regulates the cell cycle (i.e., cell growth and cell cycle progression) by activating mTOR signaling [71]. Furthermore, lysosomes function as nutrient sensors in response to amino acid inputs [72], suggesting an association between lysosomes and the response to nutrients in *C. lectularus*. Investigating the phenotypic functions of the mutated genes will be a challenge in the future. Genome editing is an effective method for elucidating molecular mechanisms underlying insecticide resistance.

4.6. Limitations of this study

Mutation identified in this study may be rare depending on the populations. In bed bugs, population genetics to detect alleles with mutation have been carried out on voltage-gated sodium channel α-subunit gene [20]. The functional contributions of the mutations to pyrethroid resistance levels could be better elucidated by comparing genes from a larger number of common bedbug strains. Furthermore, strict evaluation for pyrethroid resistance levels may need population genetic approach to clarify the allele frequencies.

At this stage, we can only link the mutated genes identified in this study to functions known in other insects, as it has not yet been demonstrated how these mutations contribute to changes in insect physiology. While these mutations likely enhance efficiency to relevant physiological processes, gene functional analyses are necessary to fully understand how these mutations contribute to the development of insecticide resistance.

5. Conclusions

Through genome-wide search for gene mutations in resistant strains of *C. lectularius*, we identified 729 mutant transcripts that have scarcely been studied in common bedbugs by comparing the sequences of protein-coding genes. These genes may enable the development of insecticide resistance in the resistant strains. Population genetics approach may clarify the extent of functional contributions of these mutations. In bed bugs, population genetics to detect alleles with mutation have been carried out on voltage-gated sodium channel α-subunit gene [20]. This study contributes to the expansion of target genes to detect the distribution and changes in allele frequencies within populations. Distribution and changes in allele frequency may also be helpful for assessing the degree of insecticide resistance. In addition, genome editing of these genes may be an effective method for understanding the evolution and physiological mechanisms involved in the acquisition of insecticide resistance.

Supplementary Materials: The following supporting information can be downloaded from figshare: https://doi.org/10.6084/m9.figshare.26362135.v2, Table S1: Public RNA sequencing data used in this study; Table S2: Metadata of RNA sequencing data in susceptible and resistant strains; Table S3: The counts and ratio of repeats annotated by RepeatModeler and RepeatMasker; Table S4: The full result of Fanflow4Insects; Table S5:729 transcripts with resistance-specific mutations; Table S6: The LD50 value and resistance ratio of each strain to Dinotefuran; Table S7: Detail results of Metascape when using *Homo sapiens* gene ID; Table S8: Detail results of Metascape when using *Drosophila melanogaster* gene ID. Gene annotation and protein sequences are available in FigShare (https://doi.org/10.6084/m9.figshare.26362135.v2).

Author Contributions: Conceptualization, K. T. and H. B.; methodology, K. T. and H. B.; software, K. T.; validation, K. T. and H. B.; formal analysis, K. T.; investigation, K. T.; resources, H. B.; data curation, K. T.; writing—original draft preparation, K. T. and F. K.; writing—review and editing, K. T., F.K., H. F., and H. B.; visualization, K. T. and F. K.; supervision, H. B.; project administration, H. B.; funding acquisition, F. K., H. F. and H. B. All authors have read and agreed to the published version of the manuscript.".

Funding: This study was supported by the Center of Innovation for Bio-Digital Transformation (BioDX), an open innovation platform for industry-academia co-creation of JST (COI-NEXT, grant number JPMJPF2010 to H.B.), and the Subsidy for Bioeconomy Industry Creation Support Project (Hiroshima Prefecture).

Data Availability Statement: Assembled sequences and raw sequence data were deposited in DDBJ under the umbrella BioProjects PRJDB18520 and PRJDB18521 for susceptible and resistant strains, respectively. Primary and alternate contigs of the susceptible strain were deposited under the accession numbers BAAFRD010000001-BAAFRD010001324 and BAAFRE010000001-BAAFRE010006538, respectively. Primary and alternate contigs of the resistant strain were deposited in the DDBJ under the accession numbers BAAFRF010000001-BAAFRF010001092 and BAAFRG010000001-BAAFRG010004661, respectively. Raw sequence reads were deposited in the DDBJ Sequence Read Archive under the accession numbers DRA018917 (HiFi reads of susceptible strains), DRA018918 (HiFi reads of resistant strains), and DRA018916

(RNA-Seq reads). The results of gene prediction and functional annotation are available at figshare (https://doi.org/10.6084/m9.figshare.26362135.v2).

Acknowledgments: We would like to thank all laboratory members at Hiroshima University for their valuable comments. Computations were performed on the computers at the Hiroshima University Genome Editing Innovation Center. Computations were partially performed on the NIG supercomputer at ROIS National Institute of Genetics.

Conflicts of Interest: K.T. and H.B. declare no conflicts of interest. F.K. and H.F. are employees of Fumakilla Limited.

Figure legends

Figure 1. K-mer assembly evaluation by Merqury and dot plot between susceptible and resistant strains. (a) K-mer spectrum assembly plot generated by Merqury of the susceptible strain and (b) resistant strain. Primary or alternate contig are indicated by different colors. (c) Dot-plot between susceptible strain and resistant strain. Color legend indicate the mean percent identity (per query)

Figure 2. Transcripts with mutations in the resistant strain. (a) The number of mutation sites per transcript is shown by circles. (b-g) Mutation sites of candidate resistance genes are shown with different amino acids. "Susceptible" refers to susceptible strain sequenced in this study and Clec2.1. Gene IDs are indicated by 'g' followed by a number, and transcript variations are denoted by 't' followed by a number. In (b), two mutated sites corresponded to the sites of 925 in housefly *Musca domestica*.

Figure 3. Results of enrichment analysis by Metascape. (a) Results using human gene ID and (b) results using fly gene ID are shown respectively. Bar graphs are colored differently according to the value of $-\log10(P)$.

References

1. Akhoundi, M.; Sereno, D.; Durand, R.; Mirzaei, A.; Bruel, C.; Delaunay, P.; Marty, P.; Izri, A. Bed Bugs (Hemiptera, Cimicidae): Overview of Classification, Evolution and Dispersion. *Int. J. Environ. Res. Public Health* **2020**, *17*, 4576, doi:10.3390/ijerph17124576.

2. Doggett, S.L.; Lee, C.-Y. Historical and Contemporary Control Options Against Bed Bugs, *Cimex* Spp. *Annu. Rev. Entomol.* **2023**, *68*, 169–190, doi:10.1146/annurev-ento-120220-015010.

3. Usinger, R.L. *Monograph of Cimicidae (Hemiptera-Heteroptera)*; The Thomas Say Foundation-Entomological Society of America: Lanham, MD, USA, 1966.

4. Reinhardt, K.; Siva-Jothy, M.T. Biology of the Bed Bugs (Cimicidae). *Annu. Rev. Entomol.* **2007**, *52*, 351–374, doi:10.1146/annurev.ento.52.040306.133913.

5. Sheele, J.M.; Crandall, C.; Chang, B.F.; Arko, B.L.; Dunn, C.; Negrete, A. Cimicosis in Persons Previously Fed Upon by Bed Bugs. *Cureus* **2019**, doi:10.7759/cureus.5941.

6. Goddard, J. Bed Bugs (Cimex Lectularius) and Clinical Consequences of Their Bites. *JAMA* **2009**, *301*, 1358, doi:10.1001/jama.2009.405.

7. Doggett, S.L.; Dwyer, D.E.; Peñas, P.F.; Russell, R.C. Bed Bugs: Clinical Relevance and Control Options. *Clin. Microbiol. Rev.* **2012**, *25*, 164–192, doi:10.1128/CMR.05015-11.

8.   DeVries, Z.C.; Santangelo, R.G.; Barbarin, A.M.; Schal, C. Histamine as an Emergent Indoor Contaminant: Accumulation and Persistence in Bed Bug Infested Homes. *PLoS ONE* **2018**, *13*, e0192462, doi:10.1371/journal.pone.0192462.

9.   Gries, R.; Britton, R.; Holmes, M.; Zhai, H.; Draper, J.; Gries, G. Bed Bug Aggregation Pheromone Finally Identified. *Angew. Chem.* **2015**, *127*, 1151–1154, doi:10.1002/ange.201409890.

10.   Shalaby, A.R. Significance of Biogenic Amines to Food Safety and Human Health. *Food Res. Int.* **1996**, *29*, 675–690, doi:10.1016/S0963-9969(96)00066-X.

11.   Heyer, G.; Hornstein, O.P.; Handwerker, H.O. Skin Reactions and Itch Sensation Induced by Epicutaneous Histamine Application in Atopic Dermatitis and Controls. *J. Invest. Dermatol.* **1989**, *93*, 492–496, doi:10.1111/1523-1747.ep12284051.

12.   Boase, C. Bedbugs - Back from the Brink. *Pest. Outlook* **2001**, *12*, 159–162, doi:10.1039/b106301b.

13.   Wang, C.; Singh, N.; Cooper, R. Field Study of the Comparative Efficacy of Three Pyrethroid/Neonicotinoid Mixture Products for the Control of the Common Bed Bug, Cimex Lectularius. *Insects* **2015**, *6*, 197–205, doi:10.3390/insects6010197.

14.   Dang, K.; Doggett, S.L.; Veera Singham, G.; Lee, C.-Y. Insecticide Resistance and Resistance Mechanisms in Bed Bugs, Cimex Spp. (Hemiptera: Cimicidae). *Parasit. Vectors* **2017**, *10*, 318, doi:10.1186/s13071-017-2232-3.

15.   Mamidala, P.; Jones, S.C.; Mittapalli, O. Metabolic Resistance in Bed Bugs. *Insects* **2011**, *2*, 36–48, doi:10.3390/insects2010036.

16.   Lilly, D.G.; Latham, S.L.; Webb, C.E.; Doggett, S.L. Cuticle Thickening in a Pyrethroid-Resistant Strain of the Common Bed Bug, Cimex Lectularius L. (Hemiptera: Cimicidae). *PLoS ONE* **2016**, *11*, e0153302, doi:10.1371/journal.pone.0153302.

17.   Adelman, Z.N.; Kilcullen, K.A.; Koganemaru, R.; Anderson, M.A.E.; Anderson, T.D.; Miller, D.M. Deep Sequencing of Pyrethroid-Resistant Bed Bugs Reveals Multiple Mechanisms of Resistance within a Single Population. *PLoS ONE* **2011**, *6*, e26228, doi:10.1371/journal.pone.0026228.

18.   Zhu, F.; Gujar, H.; Gordon, J.R.; Haynes, K.F.; Potter, M.F.; Palli, S.R. Bed Bugs Evolved Unique Adaptive Strategy to Resist Pyrethroid Insecticides. *Sci. Rep.* **2013**, *3*, 1456, doi:10.1038/srep01456.

19.   Mamidala, P.; Wijeratne, A.J.; Wijeratne, S.; Kornacker, K.; Sudhamalla, B.; Rivera-Vega, L.J.; Hoelmer, A.; Meulia, T.; Jones, S.C.; Mittapalli, O. RNA-Seq and Molecular Docking Reveal Multi-Level Pesticide Resistance in the Bed Bug. *BMC Genom.* **2012**, *13*, 6, doi:10.1186/1471-2164-13-6.

20.   Zhu, F.; Wigginton, J.; Romero, A.; Moore, A.; Ferguson, K.; Palli, R.; Potter, M.F.; Haynes, K.F.; Palli, S.R. Widespread Distribution of Knockdown Resistance Mutations in the Bed Bug, *Cimex*

Lectularius (Hemiptera: Cimicidae), Populations in the United States. *Arch. Insect Biochem. Physiol.* **2010**, *73*, 245–257, doi:10.1002/arch.20355.

21.   Seong, K.M.; Lee, D.-Y.; Yoon, K.S.; Kwon, D.H.; Kim, H.C.; Klein, T.A.; Clark, J.M.; Lee, S.H. Establishment of Quantitative Sequencing and Filter Contact Vial Bioassay for Monitoring Pyrethroid Resistance in the Common Bed Bug, Cimex Lectularius. *J. of Med. Entomol.* **2010**, *47*, 592–599, doi:10.1093/jmedent/47.4.592.

22.   Soderlund, D.M.; Knipple, D.C. The Molecular Biology of Knockdown Resistance to Pyrethroid Insecticides. *Insect Biochem. Mol. Biol.* **2003**, *33*, 563–577, doi:10.1016/S0965-1748(03)00023-7.

23.   Komagata, O.; Kasai, S.; Itokawa, K.; Minagawa, K.; Kazuma, T.; Mizutani, K.; Muto, A.; Tanikawa, T.; Adachi, M.; Komatsu, N.; et al. Common Substitution Mutation F348Y of Acetylcholinesterase Gene Contributes to Organophosphate and Carbamate Resistance in Cimex Lectularius and C. Hemipterus. *Insect Biochem. Mol. Biol.* **2021**, *138*, 103637, doi:10.1016/j.ibmb.2021.103637.

24.   Benoit, J.B.; Adelman, Z.N.; Reinhardt, K.; Dolan, A.; Poelchau, M.; Jennings, E.C.; Szuter, E.M.; Hagan, R.W.; Gujar, H.; Shukla, J.N.; et al. Unique Features of a Global Human Ectoparasite Identified through Sequencing of the Bed Bug Genome. *Nat. Commun.* **2016**, *7*, 10165, doi:10.1038/ncomms10165.

25.   Rosenfeld, J.A.; Reeves, D.; Brugler, M.R.; Narechania, A.; Simon, S.; Durrett, R.; Foox, J.; Shianna, K.; Schatz, M.C.; Gandara, J.; et al. Genome Assembly and Geospatial Phylogenomics of the Bed Bug Cimex Lectularius. *Nat. Commun.* **2016**, *7*, 10164, doi:10.1038/ncomms10164.

26.   Fountain, T.; Ravinet, M.; Naylor, R.; Reinhardt, K.; Butlin, R.K. A Linkage Map and QTL Analysis for Pyrethroid Resistance in the Bed Bug Cimex Lectularius. *G3 (Bethesda)* **2016**, *6*, 4059–4066, doi:10.1534/g3.116.033092.

27.   Haberkorn, C.; David, J.; Henri, H.; Delpuech, J.; Lasseur, R.; Vavre, F.; Varaldi, J. A Major 6 Mb Superlocus Is Involved in Pyrethroid Resistance in the Common Bed Bug *Cimex Lectularius. Evol. Appl.*   **2023**, *16*, 1012–1028, doi:10.1111/eva.13550.

28.   Hirao, M. Recent Resurgence of Bedbug and Its Management. *Jap. J. Sanit. Zool.* **2010**, *61*, 211–221, doi:10.7601/mez.61.211.

29.   Yokoo, H. Experimental Studies on the Ecology and Chemical Control of Cimex Jectularius and C. Hemipterus. *Endemic diseases bulletin of Nagasaki University* **1959**, 1, 398–406

30.   World Health Organization World Health Statistics 2009. *Estadísticas sanitarias mundiales 2009* **2009**, 149.

31.   Bliss, C.I. The Method of Probits. *Science* **1934**, *79*, 38–39, doi:10.1126/science.79.2037.38.

32.   Fieller, E.C. The Biological Standardization of Insulin. *Journal of the Royal Statistical Society Series B: Statistical Methodology* **1941**, *7*, 1–54, doi:10.2307/2983630.

33. Li, H. Minimap2: Pairwise Alignment for Nucleotide Sequences. *Bioinformatics* **2018**, *34*, 3094–3100, doi:10.1093/bioinformatics/bty191.

34. Cheng, H.; Concepcion, G.T.; Feng, X.; Zhang, H.; Li, H. Haplotype-Resolved de Novo Assembly Using Phased Assembly Graphs with Hifiasm. *Nat Methods* **2021**, *18*, 170–175, doi:10.1038/s41592-020-01056-5.

35. Gurevich, A.; Saveliev, V.; Vyahhi, N.; Tesler, G. QUAST: Quality Assessment Tool for Genome Assemblies. *Bioinformatics* **2013**, *29*, 1072–1075, doi:10.1093/bioinformatics/btt086.

36. Rhie, A.; Walenz, B.P.; Koren, S.; Phillippy, A.M. Merqury: Reference-Free Quality, Completeness, and Phasing Assessment for Genome Assemblies. *Genome Biol* **2020**, *21*, 245, doi:10.1186/s13059-020-02134-9.

37. Krueger, F.; James, F.; Ewels, P.; Afyounian, E.; Weinstein, M.; Schuster-Boeckler, B.; Hulselmans, G.; Sclamons FelixKrueger/TrimGalore: V0.6.10 - Add Default Decompression Path 2023.

38. Martin, M. Cutadapt Removes Adapter Sequences from High-Throughput Sequencing Reads. *EMBnet j.* **2011**, *17*, 10, doi:10.14806/ej.17.1.200.

39. Kim, D.; Paggi, J.M.; Park, C.; Bennett, C.; Salzberg, S.L. Graph-Based Genome Alignment and Genotyping with HISAT2 and HISAT-Genotype. *Nat Biotechnol* **2019**, *37*, 907–915, doi:10.1038/s41587-019-0201-4.

40. Li, H.; Handsaker, B.; Wysoker, A.; Fennell, T.; Ruan, J.; Homer, N.; Marth, G.; Abecasis, G.; Durbin, R.; 1000 Genome Project Data Processing Subgroup The Sequence Alignment/Map Format and SAMtools. *Bioinformatics* **2009**, *25*, 2078–2079, doi:10.1093/bioinformatics/btp352.

41. Gabriel, L.; Brůna, T.; Hoff, K.J.; Ebel, M.; Lomsadze, A.; Borodovsky, M.; Stanke, M. BRAKER3: Fully Automated Genome Annotation Using RNA-Seq and Protein Evidence with GeneMark-ETP, AUGUSTUS, and TSEBRA. *Genome Res.* **2024**, genome;gr.278090.123v1, doi:10.1101/gr.278090.123.

42. Kuznetsov, D.; Tegenfeldt, F.; Manni, M.; Seppey, M.; Berkeley, M.; Kriventseva, E.V.; Zdobnov, E.M. OrthoDB V11: Annotation of Orthologs in the Widest Sampling of Organismal Diversity. *Nucleic Acids Research* **2023**, *51*, D445–D451, doi:10.1093/nar/gkac998.

43. Pertea, G.; Pertea, M. GFF Utilities: GffRead and GffCompare. *F1000Res* **2020**, *9*, 304, doi:10.12688/f1000research.23297.2.

44. Bono, H.; Sakamoto, T.; Kasukawa, T.; Tabunoki, H. Systematic Functional Annotation Workflow for Insects. *Insects* **2022**, *13*, 586, doi:10.3390/insects13070586.

45. Shumate, A.; Salzberg, S.L. Liftoff: Accurate Mapping of Gene Annotations. *Bioinformatics* **2021**, *37*, 1639–1643, doi:10.1093/bioinformatics/btaa1016.

46. Katoh, K. MAFFT: A Novel Method for Rapid Multiple Sequence Alignment Based on Fast Fourier Transform. *Nucleic Acids Research* **2002**, *30*, 3059–3066, doi:10.1093/nar/gkf436.

47.	Price, M.N.; Arkin, A.P. Interactive Analysis of Functional Residues in Protein Families. *mSystems* **2022**, *7*, e00705-22, doi:10.1128/msystems.00705-22.

48.	Zhou, Y.; Zhou, B.; Pache, L.; Chang, M.; Khodabakhshi, A.H.; Tanaseichuk, O.; Benner, C.; Chanda, S.K. Metascape Provides a Biologist-Oriented Resource for the Analysis of Systems-Level Datasets. *Nat Commun* **2019**, *10*, 1523, doi:10.1038/s41467-019-09234-6.

49.	Duan, J.J.; Larson, K.; Watt, T.; Gould, J.; Lelito, J.P. Effects of Host Plant and Larval Density on Intraspecific Competition in Larvae of the Emerald Ash Borer (Coleoptera: Buprestidae). *Environ Entomol* **2013**, *42*, 1193–1200, doi:10.1603/EN13209.

50.	Hoff, K.J.; Lomsadze, A.; Borodovsky, M.; Stanke, M. Whole-Genome Annotation with BRAKER. In *Gene Prediction*; Kollmar, M., Ed.; Methods in Molecular Biology; Springer New York: New York, NY, 2019; Vol. 1962, pp. 65–95 ISBN 978-1-4939-9172-3.

51.	Ohta, S.; Tatsumi, Y.; Fujita, M.; Tsurimoto, T.; Obuse, C. The ORC1 Cycle in Human Cells. *Journal of Biological Chemistry* **2003**, *278*, 41535–41540, doi:10.1074/jbc.M307535200.

52.	Ding, L.; Cao, J.; Lin, W.; Chen, H.; Xiong, X.; Ao, H.; Yu, M.; Lin, J.; Cui, Q. The Roles of Cyclin-Dependent Kinases in Cell-Cycle Progression and Therapeutic Strategies in Human Breast Cancer. *IJMS* **2020**, *21*, 1960, doi:10.3390/ijms21061960.

53.	Steiner, D.F. The Proprotein Convertases. *Current Opinion in Chemical Biology* **1998**, *2*, 31–39, doi:10.1016/S1367-5931(98)80033-1.

54.	Wilkins, R.M. Insecticide Resistance and Intracellular Proteases. *Pest Management Science* **2017**, *73*, 2403–2412, doi:10.1002/ps.4646.

55.	Montalbetti, N.; Simonin, A.; Kovacs, G.; Hediger, M.A. Mammalian Iron Transporters: Families SLC11 and SLC40. *Molecular Aspects of Medicine* **2013**, *34*, 270–287, doi:10.1016/j.mam.2013.01.002.

56.	Teleman, A.A. Molecular Mechanisms of Metabolic Regulation by Insulin in *Drosophila*. *Biochemical Journal* **2010**, *425*, 13–26, doi:10.1042/BJ20091181.

57.	Xu, Z.; Zhu, L.; Yang, Y.; Zhang, Y.; Lu, M.; Tao, L.; Xu, W. Bifenthrin Induces DNA Damage and Autophagy in Spodoptera Frugiperda (Sf9) Insect Cells. *In Vitro Cell.Dev.Biol.-Animal* **2021**, *57*, 264–271, doi:10.1007/s11626-021-00554-w.

58.	Ingham, V.A.; Tennessen, J.A.; Lucas, E.R.; Elg, S.; Yates, H.C.; Carson, J.; Guelbeogo, W.M.; Sagnon, N.; Hughes, G.L.; Heinz, E.; et al. Integration of Whole Genome Sequencing and Transcriptomics Reveals a Complex Picture of the Reestablishment of Insecticide Resistance in the Major Malaria Vector Anopheles Coluzzii. *PLoS Genet* **2021**, *17*, e1009970, doi:10.1371/journal.pgen.1009970.

59.	Xu, H.; Lee, S.-J.; Suzuki, E.; Dugan, K.D.; Stoddard, A.; Li, H.-S.; Chodosh, L.A.; Montell, C. A Lysosomal Tetraspanin Associated with Retinal Degeneration Identified via a Genome-Wide Screen. *EMBO J* **2004**, *23*, 811–822, doi:10.1038/sj.emboj.7600112.

60.   Hertenstein, H.; McMullen, E.; Weiler, A.; Volkenhoff, A.; Becker, H.M.; Schirmeier, S. Starvation-Induced Regulation of Carbohydrate Transport at the Blood–Brain Barrier Is TGF-β-Signaling Dependent. *eLife* **2021**, *10*, e62503, doi:10.7554/eLife.62503.

61.   Xu, J.; Sheng, Z.; Palli, S.R. Juvenile Hormone and Insulin Regulate Trehalose Homeostasis in the Red Flour Beetle, Tribolium Castaneum. *PLoS Genet* **2013**, *9*, e1003535, doi:10.1371/journal.pgen.1003535.

62.   Gao, Y.; Zhu, C.; Li, K.; Cheng, X.; Du, Y.; Yang, D.; Fan, X.; Gaur, U.; Yang, M. Comparative Proteomics Analysis of Dietary Restriction in Drosophila. *PLoS ONE* **2020**, *15*, e0240596, doi:10.1371/journal.pone.0240596.

63.   McMullen, E.; Weiler, A.; Becker, H.M.; Schirmeier, S. Plasticity of Carbohydrate Transport at the Blood-Brain Barrier. *Front. Behav. Neurosci.* **2021**, *14*, 612430, doi:10.3389/fnbeh.2020.612430.

64.   Zhang, C.; Seong, K.M.; Sun, W.; Mittapalli, O.; Qiu, B.; Clark, J.M.; Pittendrigh, B.R. The Insulin Signaling Pathway in Drosophila Melanogaster: A Nexus Revealing an "Achilles' Heel" in DDT Resistance. *Pesticide Biochemistry and Physiology* **2021**, *171*, 104727, doi:10.1016/j.pestbp.2020.104727.

65.   Sun, R.; Jiang, L.; Chen, W.; Xu, Y.; Yi, X.; Zhong, G. Azadirachtin Exposure Inhibit Ovary Development of Spodoptera Litura (Lepidoptera: Noctuidae) by Altering Lipids Metabolism Event and Inhibiting Insulin Signaling Pathways. *Ecotoxicology and Environmental Safety* **2023**, *262*, 115151, doi:10.1016/j.ecoenv.2023.115151.

66.   Abbasi-Mojdehi, M.R.; Hajizadeh, J.; Zibaee, A.; Keyhanian, A.A.; Rajaei, F. Alterations in Some Physiological Processes of Bactrocera Oleae Rossi (Diptera: Tephritidae) Following Pyridalyl Treatment. *Pesticide Biochemistry and Physiology* **2020**, *164*, 85–90, doi:10.1016/j.pestbp.2019.12.014.

67.   Russell, P. Checkpoints on the Road to Mitosis. *Trends in Biochemical Sciences* **1998**, *23*, 399–402, doi:10.1016/S0968-0004(98)01291-2.

68.   DePamphilis, M.L. Cell Cycle Dependent Regulation of the Origin Recognition Complex. *Cell Cycle* **2005**, *4*, 70–79, doi:10.4161/cc.4.1.1333.

69.   Li, C.-J.; DePamphilis, M.L. Mammalian Orc1 Protein Is Selectively Released from Chromatin and Ubiquitinated during the S-to-M Transition in the Cell Division Cycle. *Molecular and Cellular Biology* **2002**, *22*, 105–116, doi:10.1128/MCB.22.1.105-116.2002.

70.   Icard, P.; Simula, L. Metabolic Oscillations during Cell-Cycle Progression. *Trends in Endocrinology & Metabolism* **2022**, *33*, 447–450, doi:10.1016/j.tem.2022.04.006.

71.   Fingar, D.C.; Blenis, J. Target of Rapamycin (TOR): An Integrator of Nutrient and Growth Factor Signals and Coordinator of Cell Growth and Cell Cycle Progression. *Oncogene* **2004**, *23*, 3151–3171, doi:10.1038/sj.onc.1207542.

72.   Mukherjee, A.B.; Appu, A.P.; Sadhukhan, T.; Casey, S.; Mondal, A.; Zhang, Z.; Bagh, M.B. Emerging New Roles of the Lysosome and Neuronal Ceroid Lipofuscinoses. *Mol Neurodegeneration* **2019**, *14*, 4, doi:10.1186/s13024-018-0300-6.

(a)

GO:0006259: DNA metabolic process
GO:1903047: mitotic cell cycle process
GO:1901142: insulin metabolic process
hsa04142: Lysosome
GO:0034660: ncRNA metabolic process
GO:0055086: nucleobase-containing small molecule metabolic process
GO:0006397: mRNA processing
R-HSA-5696397: Gap-filling DNA repair synthesis and ligation in GG-NER
GO:0010256: endomembrane system organization
GO:1901657: glycosyl compound metabolic process
R-HSA-428157: Sphingolipid metabolism
R-HSA-382551: Transport of small molecules
GO:0034976: response to endoplasmic reticulum stress
GO:0033365: protein localization to organelle
GO:0030163: protein catabolic process
GO:0060080: inhibitory postsynaptic potential
GO:0010817: regulation of hormone levels
GO:0044282: small molecule catabolic process
GO:0007040: lysosome organization
GO:0006664: glycolipid metabolic process

-log10(P)

(b)

dme04142: Lysosome - Drosophila melanogaster (fruit fly)
GO:1904659: glucose transmembrane transport
GO:1901292: nucleoside phosphate catabolic process
R-DME-5656169: Termination of translesion DNA synthesis
R-DME-381119: Unfolded Protein Response (UPR)
R-DME-5218859: Regulated Necrosis
GO:0016192: vesicle-mediated transport
dme04141: Protein processing in endoplasmic reticulum - Drosophil
GO:0006644: phospholipid metabolic process
GO:0033363: secretory granule organization
GO:0042335: cuticle development
GO:0065009: regulation of molecular function
GO:0006986: response to unfolded protein
GO:0035041: sperm DNA decondensation
GO:0007445: determination of imaginal disc primordium
GO:0019321: pentose metabolic process
GO:0000070: mitotic sister chromatid segregation
GO:0008104: protein localization
R-DME-917729: Endosomal Sorting Complex Required For Transport
R-DME-9840310: Glycosphingolipid catabolism

-log10(P)