Abundance and distribution of ringed and bearded seals in the Chukchi Sea: a reference for future trends

Peter L. Boveng[1]*, Vladimir I. Chernook[2], Erin E. Moreland[1], Paul B. Conn[1], Irina S. Trukhanova[3,1], Michael F. Cameron[1], Cynthia L. Christman[4,1], Justin A. Crawford[5], Lois Harwood[6], Benjamin X. Hou[1], Stacie M. Koslovsky[1], Jessica M. Lindsay[1,7], Denis I. Litovka[8], Josh M. London[1], Brett T. McClintock[1], Nikita Platonov[9], Lori Quakenbush[5], Erin L. Richmond[1], Alexander Vasiliev[2], Andrew L. Von Duyke[10], and Amy Willoughby[4,1]

[1]Marine Mammal Laboratory, Alaska Fisheries Science Center, National Marine Fisheries Service, National Oceanic and Atmospheric Administration, Seattle, Washington, USA

[2]RPO "Marine Mammal Council", Moscow, Russia

[3]North Pacific Wildlife Consulting, Seattle, Washington, USA

[4]Cooperative Institute for Climate, Ocean, and Ecosystem Studies, University of Washington, Seattle, Washington, USA

[5]Arctic Marine Mammal Program, Alaska Department of Fish and Game, Fairbanks, Alaska, USA

[6]Fisheries Joint Management Committee, Inuvik, Northwest Territories, Canada

[7]School of Aquatic and Fishery Sciences, University of Washington, Seattle, Washington, USA

[8]Chukotka Arctic Scientific Center, Anadyr, Chukotka, Russia

[9]Severtsov Institute of Ecology and Evolution, Russian Academy of Sciences, Moscow, Russia

[10]Department of Wildlife Management, North Slope Borough, Utqiaġvik, Alaska, USA

*   Correspondence: peter.boveng@noaa.gov; ORCID 0000-0003-2732-4833

Abundance and distribution of ringed and bearded seals in the Chukchi Sea: a reference for future trends

Abstract: Ringed (*Pusa hispida*) and bearded (*Erignathus barbatus*) seals are vulnerable to loss of their sea ice habitat in the rapidly warming Arctic. In April and May of 2016, we conducted an aerial survey over the ice-covered areas of the Chukchi Sea using thermal and color cameras to detect and count these seals on sea ice. We related the seal counts to environmental variables, and used the relationships to estimate the species' distributions and abundance throughout the Chukchi Sea. We accounted for incomplete detection due to seals missed by sensors or image processing errors, behavioral responses to aircraft, or incomplete availability (i.e. seals that are in water or in snow dens on the ice, called lairs). For the latter, we used satellite-linked time-depth recorders and remotely sensed snow melt indices to estimate the proportion of ringed seal individuals that are visible on ice for each day and location surveyed. To our knowledge, this is the first study where use of lairs by ringed seals has been formally addressed while estimating abundance from aerial surveys. Ringed seal abundance was estimated as $\widehat{N}_{rd} = 592{,}577$ (95% CI: 478,448–733,929), with highest densities near Kotzebue Sound, Alaska, USA. Bearded seal abundance was estimated as $\widehat{N}_{bd} = 147{,}421$ (95% CI: 114,155–190,380), with highest densities in broken pack ice near Bering Strait. The influence of spatial variables, such as snow depth and ice type, was consistent with prior studies of the species' natural history, particularly ringed seals' preference for snow of adequate depth for lairs. Our study provides the first comprehensive abundance estimates for ringed and bearded seals in the Chukchi Sea and establishes a reference for monitoring how their populations respond to Arctic warming.

Keywords: Beringia, wildlife remote sensing, infrared imaging, multispectral imaging, *Phocidae*, *Phoca hispida*, shorefast ice, snow on sea ice

1 Introduction

Arctic ringed seals (*Pusa hispida hispida*) and Pacific bearded seals (*Erignathus barbatus nauticus*) are denizens of the sea ice, traditional staple resources for Arctic Indigenous communities, and crucial prey for polar bears (*Ursus maritimus*). They are considered ice-associated species because they rely on sea ice to support vital aspects of their life history. Climate-driven declines in seasonal sea ice (e.g. Stroeve and Notz 2018; Stabeno and Bell 2019; Kim et al. 2023) have prompted concern for the viability of many Arctic marine mammal species (Laidre et al. 2015), including ice-associated seals. In 2012, the Arctic subspecies of ringed seals and the Beringia Distinct Population Segment (DPS) of Pacific bearded seals were listed as Threatened under the U.S. Endangered Species Act (National Marine Fisheries Service 2012a; National Marine Fisheries Service 2012b). The listing decisions were primarily based on concerns of future habitat loss (i.e. sea ice) rather than documented population declines, as trends in population status of these currently abundant species remain largely unknown (Laidre et al. 2015).

Despite concerns about ringed and bearded seal long-term persistence, it remains difficult to reliably estimate abundance and trends due to the challenges of surveys over vast and remote ranges. Different survey methods have been used over several decades, further making temporal comparisons difficult. Early aerial surveys for ringed seals in Alaska waters (e.g., the Chukchi and Beaufort seas; see Frost et al. 1988; Frost et al. 2004, and references therein) used fixed-wing aircraft at low altitude, with human observers visually counting and identifying seals in predetermined survey strips over the landfast ice zone. A shared set of survey protocols allowed these surveys to be compared to one another as indices of ringed seal abundance. However, variation in detection probability, observer experience, and aircraft disturbance, as well as a lack of correction for availability (i.e. the proportion of seals exposed on the ice and available for detection, versus in the water or under snow and not available; Marsh and Sinclair 1989), make it difficult to compare those early surveys with more recent surveys that used different aircraft, observers, and protocols for counting seals.

For aerial surveys of ringed and bearded seals in 1999 and 2000, Bengtson et al. (2005) used distance sampling methods (Buckland et al. 2001) and wet/dry data from satellite-linked time-depth recorders (TDRs) to estimate and correct for the proportion of ringed seals that were not available because they were in the water; there were no TDR data for bearded seals. The ringed seal abundance estimates, corrected for availability, are likely more robust for comparison purposes but, like the earlier surveys, they remained uncorrected for the detection probability on the trackline (g(0) in distance sampling terminology; e.g., Conn et al. 2012; Burt et al. 2014). In the mid-2000s, helicopter-based surveys of ice-associated seals in the Bering

Sea attempted to account for g(0) < 1.0 by using mark-recapture distance sampling (Borchers et al. 2006) to estimate detection probability, and TDR-based corrections for availability (Conn et al. 2013; Ver Hoef et al. 2014). However, due to seals' response to noise, helicopter surveys caused disturbance that negatively biased abundance estimates (M. Cameron, pers. obs.) and contributed to substantial species identification errors (Conn et al. 2013).

In 2012 and 2013, our team of U.S. and Russian Federation scientists conducted the first comprehensive surveys of ice-associated seals in the Bering Sea and Sea of Okhotsk (Boveng et al. In prep., Chernook et al. 2014; Conn et al. 2014; Sigler et al. 2015; Chernook et al. 2018). These sensor-based aerial surveys used fixed-wing aircraft, but instead of human observers they used thermal imaging to detect the warm seals resting on the cold sea ice. Thermal detections were then matched with higher-resolution digital color photographs to determine species. As far as we are aware, this was the first analytical effort to estimate absolute seal abundance while accounting simultaneously for incomplete detection by the sensors and image processing tools, species misidentification, and availability less than 1.0. However, estimates of ringed seal abundance from Conn et al. (2014) could not be corrected for availability because of uncertainty about the proportion of ringed seals that were using snow dens on the sea ice ('subnivean lairs'; Kelly et al. 2006) while surveys were conducted.

In 2016, our team conducted a second joint U.S.–Russian Federation survey, for ringed and bearded seals in the Chukchi Sea and a small portion of the East Siberian Sea, again using sensor-based protocols in fixed-wing aircraft. That survey is the focus of this paper, in which we report results of the Chukchi and East Siberian Surveys (ChESS), using a spatio-temporal statistical model to estimate the abundance of ringed and bearded seals while accounting for incomplete detection, behavioral response of seals to survey aircraft, and availability less than 1.0. Importantly, our approach incorporates a novel conceptual model for seal availability that relies on TDR data for both ringed and bearded seals, and on the timing of snow melt onset to help quantify ringed seals' transition from hauling out in lairs to hauling out on the exposed ice or snow surface—an important component of their availability.

2 Methods and Materials

2.1 Overview

In our aerial survey, we obtained counts of ringed and bearded seals in the area sampled by our sensors. With the survey area divided into grid cells, seal counts within grid cells were then related to habitat variables (covariates) and used to predict abundances in unsampled locations. Our approach relied on maximum likelihood, and took into account several nuisance processes that affected the detectability of seals, such as

incomplete detection by the sensors, behavioral response of seals to aircraft (i.e. escape behavior), and avail-
ability. Our main quantities of interest were the values of ringed and bearded seal abundance that maximized
this likelihood, given the observed counts, but the covariate relationships are also informative and help to
explain habitat use. In the remainder of this section, we describe the survey and image processing methods,
our choices of explanatory variables, and our approaches to modeling ringed and bearded seal availability
and density.

2.2 Survey methods

Two teams of researchers, one in the Russian Federation exclusive economic zone (EEZ) and one in the
U.S. EEZ, conducted visual and instrument-based aerial surveys over the Chukchi Sea during April and May
of 2016 (Figure 1). Surveys in the Russian EEZ (Chernook et al. 2019) were conducted from an AN-26 Arctica
aircraft equipped with a Malakhit-M cooled long-wavelength infrared (LWIR; 8–14 μm) scanner; cooled in-
frared sensors provide greater sensitivity than uncooled sensors. At a target altitude of 250 m, the LWIR
scanner had a 'ground sampling distance' (GSD; i.e. resolution) of 11 cm/pixel on the center of the continu-
ously recorded image strip. Three Nikon D800 digital color 36-megapixel cameras with 50-mm lenses were
mounted at angles such that their combined fields of view completely covered the strip width of the LWIR
scanner, which was 483 m when flying at a target altitude of 250 m. The center-mounted color camera had a
GSD of 2.27 cm/pixel at the center of its image.

Surveys in the U.S. EEZ were conducted from a King Air A90 aircraft configured with three downward
facing, cooled LWIR cameras (FLIR A6751sc SLS) fitted with 25-mm lenses mounted in the open-air belly
port. One uncooled LWIR camera (FLIR A645sc) was used in place of a cooled camera for 13 flights while the
cooled camera was temporarily inoperable. Both types of LWIR camera provided 20–23 cm GSD at the target
survey altitude of 305 m. We paired each LWIR (hereafter referred to as 'thermal') camera with a 29-mega-
pixel machine-vision color camera (Prosilica GT6600c) fitted with a 100-mm Zeiss lens to provide a GSD of
1.71–2.13 cm/pixel at survey altitude. Paired color and thermal images were collected continuously while
on survey effort at a rate of two frames per second. Dedicated on-board computers running custom software
(Skeyes; MoviTHERM, Irvine, CA) managed image acquisition and synchronized data logging from an inte-
grated Arduino GPS. At the target altitude, the thermal cameras covered a strip width of 470 m. The color
images, which determined the area surveyed, had a footprint width equal to 83% of the thermal strip width,
or 390 m. The target survey ground speed of approximately 260km/h was set to maximize the range of the
aircraft, and resulted in 50% forward overlap of images. Side camera pairs were set at an internal angle of
25.5° to allow a slight gap between sea ice coverage in the images collected from the center and side cameras,

maximizing survey swath while eliminating image processing and analysis complications from duplicate de-
tections.

Variable weather and sea ice conditions make it difficult to implement a pre-defined sampling design.
Therefore, the survey teams were instructed to spread survey effort over space and time, as is commonly
done in space-filling designs developed for spatial modelling (Diggle and Lophaven 2006). This model-based
approach exhibited satisfactory performance in initial simulation testing, and higher performance with sur-
vey effort distributed over space and time, compared to stratified approaches used in previous surveys (Conn
et al. 2016). Survey teams were instructed to avoid consideration of expected seal density or habitat quality
when laying out or modifying survey track lines, to minimize bias due to preferential sampling (Diggle et al.
2010; Conn et al. 2017).

We divided our study area into 1,354 grid cells of approximately 25 × 25 km (625 km^2) each (Figure 1),
commensurate with the resolution of sea ice concentration data that we related to seal densities (see **2.4**
Explanatory variables). We tallied seal counts and survey effort at this scale, though with slightly different
methods for U.S. and Russian survey flights. In the United States, we quantified effort as the total area pho-
tographed with the color cameras, and did not attempt to model thermal detections that fell outside the color
images. For the Russian surveys, we used the area of the thermal scanner swath to quantify effort, because
photographs were only triggered when thermal signatures characteristic of seals on ice ('hot spots') were
detected on the thermal scanner; not all hot spots were photographed, which led to different probabilities of
identifying detected seals to species, compared to the U.S. surveys. Methods to account for these differences
are described in the section on abundance modelling below. The seal counts and metadata we used from the
Russian surveys were published in Chernook et al. (2019). Our analysis combines the U.S. and Russian data
under a single, more comprehensive model for estimation of seal abundance and distribution across the
Chukchi Sea.

Figure 1. The ringed and bearded seal aerial survey area in the Chukchi Sea, 2016. The study area is divided into 25 × 25 km (625 km^2) grid cells (blue lines). Black lines represent 'on effort' flight tracks over U.S. waters and dark gray lines represent flight tracks over Russian Federation waters. Gaps in lines represent 'off effort' segments, primarily due to fog. Blue shading represents the 1999 and 2000 seal survey area (Bengtson et al. 2005).

2.3 Processing images for seal detection

2.3.1 Russian Federation surveys

The thermal scanner recording was processed with custom software to automatically mark seals on the ice using a combination of temperature contrast, size, and shape of hot spots. Operators adjusted software settings during the process to ensure the reliable identification of thermal signatures from seals while excluding false positives, for example caused by seal water access holes or dirt (Chernook et al. 2000a;

Chernook et al. 2000b). Operators reviewed and manually corrected the results of automatic thermal pro-
cessing and used detection times and coordinates to match thermal detections with the corresponding color
images. On color images, seals detected as hot spots from the thermal scanner were marked by operators,
and remaining false-positive hot spots were eliminated by verification with the content of the color images.
Combined thermal and color image processing was independently conducted by two counters. Results from
both counts were compared for discrepancies and resolved. Once a seal was found in both thermal and color
images, a digital portrait was provided to biologists for species identification and determination of age class
(adult or pup). This processing resulted in a database containing unique seal identifiers, species information,
geographical coordinates, detection times, and thermal and color image file names.

2.3.2 U.S. surveys

After surveys were completed, we used a partially automated process to detect and classify seals in the
survey images. In the first step, the same software used for image acquisition was used to apply an automated
algorithm to find hot spots in the thermal images. The algorithm was designed to exclude hot spots that were
either too small in area (< 6 pixels) or too large (>300 pixels), and to use simple measures of the hot spot
temperature profile to further eliminate false positive detections. Specifically, a curve was fit to the grayscale
pixel values along a single line bisecting the warmest region of each hot spot. The peak value, the maximum
slope along the curve, and the second derivative of the curve were compared to threshold values that had
been empirically derived from thermal signatures of known seals in aerial imagery collected prior to our
survey. This algorithm identified 316,099 candidate hot spots from 997,642 processed thermal images. In
the second step, all candidate hot spots were then reviewed visually to remove those that were unlikely to
be from seals, reducing the number that required further processing. For each of the remaining 15,454 hot
spots, the corresponding color image was examined by experienced observers to determine the hot spot's
source and to classify species and age class (pup vs. non-pup) of detected seals. To quantify the seal detection
rate of this process, a systematic random sample of 28,052 color images was visually searched for the pres-
ence of seals by an observer without access to the thermal images. The partially automated process detected
247 of the 258 seals found in the visual search, for an overall detection rate of 0.96 (Supplement 1). Lacking
a similar study of the sensors and detection process used in the Russian Federation surveys, we applied the
same detection probability to the Russian data.

2.4 Explanatory variables

Counts of seals present on sea ice are a function of seal density and availability to be detected (Marsh and
Sinclair 1989). We selected 14 environmental, temporal, and geographic covariates to help explain variation

in density and availability (Table 1). Based on Indigenous knowledge (e.g., Oceana and Kawerak Inc. 2014; Huntington et al. 2015), natural history observations (e.g., Burns 1981), previous aerial surveys (e.g., Bengtson et al. 2005), and satellite tracking results (e.g., Crawford et al. 2012; Olnes et al. 2020; Von Duyke et al. 2020), we anticipated that seal densities would vary by sea ice concentration and type, with ringed seals typically selecting landfast and nearshore pack ice, and bearded seals typically selecting pack ice. We also anticipated that both species would occur at higher densities south of Point Hope (Bengtson et al. 2005; Von Duyke et al. 2020). Ringed seals require sufficient snow to construct subnivean lairs, so we included a remotely sensed snow depth covariate. We included wind speed, air temperature, precipitation, atmospheric pressure, day of year, and solar hour as availability covariates because more ringed and bearded seals are known to haul out on ice in calm, warm, dry conditions, later in the spring, and at hours closer to solar noon and midnight (Burns and Harbo 1972; Frost et al. 2004; London et al. 2022).

Accounting for ringed seals' availability is further complicated by their seasonal use of lairs, wherein they are not visible to the aerial survey sensors. Their transition from hauling out in lairs to hauling out on the exposed surface of the ice or snow (i.e. basking) is often not abrupt. This transition is partly influenced by increasing ambient temperatures that support the seals' molt (Feltz and Fay 1966) and cause snow melt, eventually leading to lair collapse (Kelly et al. 2006). Individuals may haul out on the ice to bask near breathing holes prior to lair collapse, and may return to using lairs between basking bouts. Ringed seals will also dig out before lairs collapse when conditions (i.e. air temperature, wind speed, and solar energy) are favorable for basking, but some lairs remain in use late into spring (Kelly and Quakenbush 1990). Thus, a seal's transition away from hauling out in lairs can be a gradual process—under normal conditions—when lairs remain intact, or more abrupt when rain, thin snow, or early snow melt cause lairs to collapse prematurely. We refer to this transition as emergence (Lindsay et al. 2021). To explore a relationship between emergence and snow melt we tested two remotely sensed covariates (Table 1) associated with snow melt onset, the date on which the snow layer first becomes saturated with moisture.

Table 1. Explanatory covariates used in the analysis of ringed and bearded seal availability and density (abundance).

Covariate name	Model component	Type	Description; source
Hour	Availability	Continuous	Solar hour, calculated as a function of time and location using the SolaR package (Lamigueiro 2012)

Day of Year	Availability	Integer	Day since December 31, 2016 (Alaska time), included as linear, quadratic, and cubic effects
Precip	Availability	Continuous	Convective precipitation (kg/m^2); North American Regional Reanalysis (NARR), National Centers for Environmental Prediction (NCEP), Mesinger et al. (2006)
Pressure	Availability	Continuous	Atmospheric pressure at sea level (kPa); NARR, NCEP, Mesinger et al. (2006)
Temp	Availability	Continuous	Air temperatures predicted at 2 m above the Earth's surface; NARR, NCEP, Mesinger et al. (2006)
Wind	Availability	Continuous	Northerly and easterly vector components combined into a single wind speed; NARR, NCEP, Mesinger et al. (2006)
Melt1	Availability (ringed seals)	Integer	Date of snow melt onset; Estimated for each grid cell from Special Sensor Microwave/Imager (SSM/I) data, using the Mean Differences and Standard Deviation Analysis (MDSDA) algorithm (Belchansky et al. 2004)
Melt2	Availability (ringed seals)	Integer	Same as Melt1, but estimated with the continuous melt onset algorithm in Markus et al. (2009) and downloaded from https://earth.gsfc.nasa.gov/cryo/data/arctic-sea-ice-melt
Sea_ice	Seal density	Continuous	Proportion of each grid cell covered in sea ice; Sea Ice Concentrations from Nimbus-7 SMMR and DMSP SSM/I-SSMIS Passive Microwave Data, Version 2, https://nsidc.org/data/nsidc-0051/versions/2#anchor-2, (Cavalieri et al. 1997)
I_no_ice	Seal density	Binary	A binary indicator for whether a grid cell had 0% sea ice
Fast_ice	Seal density	Continuous	Proportion of a grid cell covered in landfast ice. Derived from U.S. National Ice Center, Weekly Arctic Shapefile, https://usicecenter.gov/Products/ArcticData (accessed 2016).
Snow_depth	Seal density	Continuous	Estimated average snow depth in the last 5 days of March; computed from SSM/I data (Markus and Cavalieri 1998; Comiso et al. 2003),

Easting	Seal density	Continuous	East-west scale on a WGS 84/ North Pole Lambert azimuthal equal-area (LAEA) Bering projection (EPSG: 3571). This covariate was mean-standardized before inclusion in seal abundance models.
Northing	Seal density	Continuous	North-south scale on same projection as above. This covariate was also mean standardized.

2.5 Availability modeling

Our surveys only had the ability to detect seals that were basking on ice and visible to the airborne sen-
sors. Thus, to estimate seal abundance from aerial survey counts, we needed to account for the proportion
of seals that were available, a_{ist} (Bengtson et al. 2005). Here, the subscripts indicate that availability has the
potential to vary by species (i), location (i.e. grid cell; s), and time (t). An alphabetical list of notation used in
model development is provided in Table 2.

Table 2. Notation for models fitted to ringed and bearded seal aerial survey count data.

Notation	Definition
$\hat{a}$	Average estimate of availability used to correct historical bearded seal abundance estimates from Bengtson et al. (2005).
a_{ist}	Predicted probability that a seal of species i is available to be detected (i.e. is on ice and visible to aircraft) during surveys of grid cell s at time t. On occasion we drop the i subscript when the species is apparent.
A	Study area size (km^2) for Bengtson et al. (2005) survey
A_{st}	Proportional area of grid cell s that was surveyed by aircraft at time t. For Russian surveys, A_{st} was calculated relative to the width of the thermal survey swath; for U.S. surveys, it was calculated relative to the total area photographed.
α_{st}	Probability that a hauled out ringed seal is not in a subnivean lair.
b_i	Probability that a seal of species i remains on ice and does not flush into water as aircraft passes overhead (i.e. the complement of disturbance probability).

$\boldsymbol{\beta}_i$	Vector of regression coefficients used to explain abundance-covariate relationships for species i.
c_{st}	Probability that a seal, detected in cell s at time t during Russian surveys, is identified to species.
c^{US}	Probability that a seal, detected during U.S. surveys, is identified to species.
C_{ist}	Count of seals of species i that are photographed during aerial surveys in grid cell s at time t Note that i=1 is ringed seals; i=2 is bearded seals; and i=3 denotes that species could not be determined.
d	Probability that a seal basking on ice is detected by the detection algorithm applied to thermal imagery.
Day_t	Julian Day associated with survey time t
$\widehat{D}$	Overall density of bearded seals (#/km²) from Bengtson et al. (2005) aerial surveys.
δ_{st}	Difference between Julian day associated with survey time t and the day of melt onset for cell s.
h_{ist}	Predicted probability that a seal of species i is hauled out of the water in surveys of grid cell s at time t.
μ_{ist}	Expected count of seal species i at time t in grid cell s.
$\widehat{N}$	Corrected bearded seal abundance estimate associated with Bengtson et al. (2005) aerial surveys (correcting for availability probability)
N_i	Total abundance of species i in our survey area (i=1: ringed seals; i=2: bearded seals).
N_{ist}	Abundance of species i in grid cell s at time t.
p_{ist}	Overall detection probability of species i at time t in grid cell s.
ρ	Power parameter for the Tweedie distribution.
U_{st}	Number of unphotographed seals in grid cell s at time t (Russian surveys only).
w_s	Proportion of ocean habitat in grid cell s (i.e. omitting land).
$\boldsymbol{x}_{ist}$	Vector of regression covariates associated with species i during surveys in grid cell s at time t.

2.5.1 Ringed seal availability

A ringed seal that is hauled out on the ice but concealed within a lair is not available to be detected in

aerial surveys. Therefore, we treated ringed seal availability as a product of two components: (1) the proba-

bility that a seal is out of the water (h_{st}) and (2) the probability that it is visible on the snow or ice surface

(i.e. not in a lair), given that it is out of the water (α_{st}), such that $a_{st} = h_{st}\alpha_{st}$.

To estimate h_{st} for ringed seals we used hourly wet/dry data obtained during 2005–2019 from TDRs on

65 individuals in the Bering, Chukchi, and Beaufort seas (Harwood et al. 2007; Kelly et al. 2010a; Crawford

et al. 2012; Crawford et al. 2019; Von Duyke et al. 2020). Using only the data from April–June, we fit gener-

alized additive models (GAMs) in the mgcv package (Wood 2017) to the proportion of ringed seals hauled

out during aerial survey hours (09:00–15:00 local solar time) in relation to day-of-year, spring air tempera-

ture, days from snow melt onset, and sea (Bering, Chukchi, or Beaufort). For additional details on how ringed

seal haul-out probabilities (h_{st}) were modeled, see Supplement section 2.

We used temporal trends in seal counts (Lindsay et al. 2021) to estimate the shape of α_{st}. Kelly et al.

(2006) showed that the proportion of radio-tagged ringed seals visible on the ice-snow surface increases as

the snow becomes saturated with water, likely because lairs are melting and becoming unusable, or un-

needed due to warmer ambient temperatures. Therefore, we used snow melt onset (i.e. saturation) dates

estimated with the MDSDA algorithm (Mean Differences and Standard Deviation Analysis of passive

microwave brightness temperatures; Belchansky et al. 2004) as an indicator of emergence from lairs. Defin-

ing

$$\delta_{st} = Day_t - Melt1_s$$

to be the difference between the survey day (Day_t) and the day of snow melt onset ($Melt1_s$) indicated by

the MDSDA algorithm for cell s (such that negative values indicate days prior to snow melt onset and positive

values indicate days after snow melt onset), we used this finding to set up a logistic model for the proportion

of hauled-out ringed seals that have emerged from lairs:

$$\alpha_{st} = \begin{cases} \left(1 + \exp\left(-f(\delta_{st})\right)\right)^{-1} & \text{if } \delta_{st} < 0 \\ 1.0 & \delta_{st} \geq 0 \end{cases}.$$

We included this emergence model in our larger model for ringed seal counts, using a penalized spline basis

for $f(\delta_{st})$ to estimate its (approximately sigmoid) shape, and an additional likelihood penalty to force α_{st} to

approach 1.0 by the time the MDSDA algorithm indicated that the snow melt onset had occurred. For addi-

tional information on this conceptual model, see Supplement section 2.

2.5.2 Bearded seal availability

London et al. (2022) estimated availability of bearded, ribbon, and spotted seals in the Bering and Chukchi
seas. Specifically, they applied generalized linear mixed pseudo-models (Ver Hoef et al. 2010) to data from
wet/dry sensors on TDRs attached to seals. These hourly data identified periods when seals were hauled
out of the water and, therefore—because these species rest on the ice surface and do not obscure themselves
in snow lairs—available to be detected during surveys. We used London et al.'s fitted models directly to
predict bearded seal availability for each grid cell and time that we surveyed. Predictions depended on day-
of-year, hour-of-day, temperature, wind speed, precipitation, and barometric pressure (Table 1). London et
al.'s models also used latitude to describe changes in haul-out behavior from the Bering Sea, north through
the southern Chukchi Sea, though because there were few bearded seal telemetry records from the middle
and northern Chukchi Sea during April and May, we used a single latitude value consistent with the southern
Chukchi Sea for all predictions. We also used estimated variances of predictions to propagate uncertainty
about availability into standard errors of abundance estimates (see Supplement section 1).

2.6 Abundance (density) modelling

To estimate seal abundance, we fit a statistical model that expressed the expected count of each species
as a function of (a) true, underlying abundance, (b) area surveyed, (c) the expected proportion of seals that
were basking (as a function of environmental conditions), and (d) detection probability for seals that were
basking on ice. We also evaluated species misclassification probabilities (Conn et al. 2013; McClintock et al.
2015), but we found that they were negligible and did not include them in the model. Our model is spatially
and temporally explicit, allowing expected abundance, and thus expected counts, to change by grid cell and
by day as ice and other conditions changed. We write the overall model hierarchically so that each compo-
nent can be examined in sequence.

2.6.1 Seal densities over space and time

To model how seal densities vary over time and space, we let N_i give the total abundance in the survey
area of species i. We follow the suggestion of Conn et al. (2015) to stabilize estimation by assuming that
abundance in the survey area remains constant over the course of the survey, though we allow spatial distri-
butions to shift within the survey area as sea ice changes. Given total abundance, we write the abundance of
species i, in spatial location (grid cell) s, at time (day) t, as

$N_{ist} = N_i \pi_{ist}$,

where π_{ist} is the proportion of species i that is in grid cell s at time t. We then write π_{ist} as a function of
habitat and landscape covariates using a multinomial logit link function to constrain $0 \leq \pi_{ist} \leq 1$ and
$\sum_s \pi_{ist} = 1$:

$$\pi_{ist} = \frac{w_s \exp(x_{ist}\boldsymbol{\beta})}{\sum_j w_j \exp(x_{ijt}\boldsymbol{\beta}) \sum_j w_j exp(x_{ijt}\boldsymbol{\beta})}.$$

Here, w_s is the proportion of grid cell s that is composed of marine habitat (i.e. omitting land), x_{ist} is a row
vector of explanatory variables for species i in cell s at time t, and $\boldsymbol{\beta}$ is a column vector of regression coeffi-
cients. We used a parameterization that allowed for smooth effects of continuous covariates by imbedding
penalized splines into estimation. This was done with the mgcv package (Wood 2017) to formulate x_{ist} in
terms of cubic spline bases and then penalizing $\boldsymbol{\beta}$ during estimation; for more detail, see Supplement section
1. We used sea ice concentration, proportion of landfast ice, easting, northing, and snow depth as covariates
for seal densities. Because it is difficult to determine the effective number of parameters for this class of
models, we did not conduct a formal model selection process. We provide further justification for the set of
covariate predictors in **4. Discussion**.

2.6.2 Expected seal counts related to abundance

Given a model for how abundance varies over space, we next describe how abundance relates to seal
counts. We start by noting that (1) transects only sample a small fraction of a grid cell, (2) not all seals are
detected, and (3) not all seal observations are identified to species. We thus write the expected count of each
observation type (i=1: bearded seal; i=2: ringed seal; i=3: unknown seal species) as

$$\mu_{ist} = \begin{cases} N_{ist}A_{st}p_{ist}c_{st} & \text{for } i = 1,2 \\ (1 - c_{st})\sum_{k=1}^{2} N_{kst}A_{st}p_{kst} & \text{for } i = 3 \end{cases},$$

where A_{st} is the proportion of ocean habitat in grid cell s that is surveyed at time t, p_{ist} is the probability of
detecting a seal of species i that is at location s and time t, and c_{st} is the probability of identifying the seal to
species. We further decompose p_{ist} as

$$p_{ist} = a_{ist}b_i d,$$

where a_{ist} is the probability that a seal is basking on ice during the survey (i.e. availability, as defined in
sections 2.4–2.5), b_i is a non-disturbance probability (i.e. the probability that a basking seal remains on ice
and is not flushed into the water before the sensors pass overhead), and d is the probability that a basking
seal is detected by thermal sensors and semi-automated image processing steps (Supplement section 1). We

developed an informed prior distribution for p_{ist} based on auxiliary data sources. In particular, we used
disturbance trials to provide information on b_i and double sampling to provide information on d (Conn et al.
2014).

2.6.4 Classification of counts to species

Count data from the aerial surveys consisted of counts of ringed seals (C_{1st}), counts of bearded seals
(C_{2st}), and counts of seals of unknown species (U_{st}). The probability of identifying seals to species (c_{st})
differed dramatically between U.S. and Russian surveys. In the United States, we based effort on area photo-
graphed, and only 10 out of 5,720 seal observations (0.17%) were classified as 'unknown' species. We used
a single value for c_{st}, c^{US}, where $c^{US} = \sum_s \sum_t \sum_i C_{ist} / \sum_s \sum_t \sum_i (C_{ist} + U_{st})$ (here, the summation over s only
includes cells surveyed in U.S. transects). In Russian surveys, a variable proportion of seals had clear photo-
graphs, such that 148 out of 318 seal observations (46.5%) had species information. We thus used different
approaches for parameterizing c_{st}: We set $c_{st} = \sum_i C_{ist} / (\sum_i C_{ist} + U_{st})$ for cases where $(\sum_i C_{ist} + U_{st}) > 0$.
For times and cells where no seals were counted, we set $c_{st} = \sum_s \sum_t \sum_i C_{ist} / \sum_s \sum_t \sum_i (C_{ist} + U_{st})$, where the
summations apply only to cells (s) surveyed in Russian transects (i.e. the mean proportion of seals identified
to species).

We assumed that counts (i.e., C_{ist}, U_{st}) followed a compound Poisson-Gamma (CPG) distribution with
mean μ_{ist}. The CPG distribution is a version of the Tweedie distribution where the power parameter (ρ) is
limited to the range $1 < \rho < 2$ (Jørgensen 1987). This three-parameter distribution is flexible, accommodat-
ing both zero inflation and overdispersion, and multiple authors have shown it to be appropriate for marine
mammal count data (Miller et al. 2013; Sigourney et al. 2020). Separate power and dispersion parameters
were estimated for each observation type (i.e. ringed, bearded, or unknown species counts) and survey plat-
form (i.e. U.S. vs. Russian aircraft).

2.7 Model fitting and computing

We used maximum marginal likelihood to fit our model, with a joint pseudo-likelihood that included com-
ponents for (1) U.S. and Russian count data, (2) prior distributions for regression parameters (including
spline parameters), and (3) a penalty that forced the proportion of ringed seals basking after MDSDA melt
dates to be 1.0. We used a parametric bootstrap procedure to propagate detection uncertainty into abun-
dance estimates. For more detail on likelihood components, penalties, and bootstrap procedures see Sup-
plement section 1. We coded our joint pseudo-likelihood function using Template Model Builder (TMB;

Kristensen et al. 2016) syntax, and used the 'nlminb' function in R (R Core Team 2017) to optimize the like-

lihood. Importantly, TMB allowed us to use the Laplace approximation to integrate over prior distributions

of smooth effects (which may be interpreted as random effects). We used randomized quantile residuals

(RQR; Dunn and Smyth 1996) to assess goodness-of-fit (Supplement section 1). We compiled a standalone R

package, 'CHESSseal,' that includes all data, R scripts, and TMB code needed to reproduce our analyses. It is

available on GitHub at https://github.com/pconn/CHESSseal and will be published and archived at a long-

term, publicly available repository upon manuscript acceptance.

2.8 Comparison to historical seal density estimates

Bengtson et al. (2005) estimated eastern Chukchi Sea ringed seal abundance (corrected for availability)

and bearded seal density (not corrected for detection probability) from aerial surveys in 1999 and 2000. To

compare our estimates to theirs, we attempted to match nearshore and offshore strata used in their surveys

(Figure 1). We also used the mean availability predictions for our flights in late May to correct strata-specific

Bengtson et al. (2005) bearded seal density estimates, reconstructing a survey-area-wide estimate as

$\qquad \widehat{N} = \widehat{D}A/\hat{a}$,

where $\widehat{D}$ represents a density estimate, A is the area (km^2) of sea ice habitat, and $\hat{a}$ is the mean availability

prediction. The surveys reported in Bengtson et al. (2005) occurred between May 21 and June 6. To compare

our estimates to theirs, we summed predicted abundance for grid cells with centroids overlapping their sur-

vey areas. Because our procedure results in an estimate for each day of our survey, we took the mean value

from May 21 through the end of our survey on May 31. We used the coefficient of variation from our previous

bootstrap procedure to construct log-based confidence intervals (Burnham et al. 1987; Buckland et al. 2001)

for our estimates. We used the delta method (Dorfman 1938) to calculate standard errors for abundance

estimates associated with Bengtson et al. (2005) surveys.

Although there were surveys of seals in the Chukchi and Beaufort seas prior to 1999 (e.g., Frost et al. 1988;

Frost et al. 2004), we do not compare our results to those surveys because those estimates were uncorrected

for availability and detection probability.

2.9 Ethics approval

This research was conducted in conformance with all U.S., Russian Federation, and Canadian laws appli-

cable to ringed and bearded seals. Alaska Fisheries Science Center (AFSC) conducted aerial surveys and de-

ployed TDRs on bearded seals under National Marine Fisheries Service (NMFS) Research Permits 782-1765

and 15126, and Letters of Assurance of Compliance with Animal Welfare Act regulations, A/NW 2010-3 and
A/NW 2016-1 from the AFSC/Northwest Fisheries Science Center Institutional Animal Care and Use Com-
mittee. Ringed and bearded seals were tagged with TDRs by Alaska Department of Fish and Game (ADFG)
and North Slope Borough Department of Wildlife Management (NSB-DWM) under NMFS Research Permits
358-1787-01, 15324, and 30466 issued to ADFG, and under approved protocols by the ADFG Animal Care
and Use Committee 2006-16, 2014–03, 2015-25, 2016-23, 0027-2017-27, and 0027-2018-29. Ringed seal
tagging with TDRs in Canada was conducted under Department of Fisheries and Oceans Scientific License
number SLE-04/05-328 and SLE-05/06-322, and Animal Care and Use Protocol UFWI-ACC-2004-2005-
001U.

3 Results

The U.S. survey aircraft completed 13,685 km of track line on survey effort between 7 April and 31 May
2016; the Russian Federation aircraft completed 13,961 km during 19–26 April and 12–18 May (Figure 1).
The total area over which ringed and bearded seal abundances were estimated was 846,250 km^2, 35% within
the U.S. EEZ and 65% within the Russian Federation EEZ. The total numbers of individual seals photographed
were 4,764 ringed seals, 1,094 bearded seals, and 180 unknown species. A total of 5,720 seals were observed
in US waters, while 318 were observed in Russian waters. This difference in encounter rate was mostly due
to differences in seal densities, although the Russian aircraft also had a lower detection probability for ringed
seals due to higher behavioral response to the aircraft and fewer seals predicted to be basking during the
Russian survey flights. For instance, the mean predicted detection probability for ringed seals was $\bar{p} = 0.24$
and $\bar{p} = 0.12$ for U.S. and Russian surveys, respectively, but for bearded seals it was $\bar{p} = 0.23$ for each coun-
try.

We estimated that abundances of ringed and bearded seals in the Chukchi Sea were $\widehat{N}_{rd} = 592{,}577$ (95%
CI: 478,448–733,929) and $\widehat{N}_{bd} = 147{,}421$ (95% CI: 114,155–190,380), respectively. Over the course of the
study, the highest densities of ringed seals were observed in landfast ice in and around Kotzebue Sound,
Alaska (Figure 2; see place names in Figure 1). Bearded seal densities were highest in offshore pack ice just
north of the Bering Strait (Figure 2). Although the distribution of seals varied somewhat as ice conditions
changed, time-averaged estimates of abundance suggested that 80% of ringed seals (point estimate 474,908)
and 64% of bearded seals (point estimate 93,766) were in U.S. waters. The corresponding numbers for Rus-
sian waters, 117,669 ringed seals and 53,655 bearded seals, differ from those obtained by Chernook et al.
(2019) because that initial analysis did not account for availability, among several other analytical differ-
ences. P-values from RQR goodness-of-fit tests were all greater than 0.10 for ringed seals, bearded seals, and

unknown seal species in both U.S. and Russian surveys, suggesting that our models fit the count data reason-
ably well (Supplement section 1).

Ringed seal densities were dependent on northing and easting predictors—latitude and longitude effects,
respectively—with densities (Figure 3) predicted to be the highest in the southeast quadrant of our study
area, near Kotzebue Sound, and in the western edge of our study area, in the East Siberian Sea near
Chaunskaya Bay, Chukotka (Figure 2). Densities increased slightly with sea ice concentrations above ≈0.6,
but there was also a negative relationship (and high uncertainty) with lower ice concentrations. Ringed seal
densities were high in cells with small proportions of landfast ice (<0.25) and in cells with intermediate to
high proportions of landfast ice (>0.5), but lower in between these values (Figure 3). It should be noted that
grid cells along much of the Alaska coastline (e.g. in locations north of Kotzebue Sound; Figure 1) had small
proportions of landfast ice, simply because the size of the grid cells (≈625 km^2) was large relative to the
typically narrow band of landfast ice in each cell (Mahoney et al. 2014). Also, predicted ringed seal densities
were high in Kotzebue Sound, where many cells had high or complete landfast ice cover. Densities increased
strongly with snow depth values up to 20 cm, then reached a plateau between 20 and 30 cm (Figure 3). The
modelled probability that a ringed seal was visible on the snow or ice surface (i.e. not in a lair), given that it
was out of the water, increased from 0 at the beginning of the survey to 0.25 at about 15 days prior to the
onset of snow melt indicated by the MDSDA index, (Figure 4); this probability then increased rapidly to 1.0
by about 5 days after snow melt onset.

The relationships between bearded seal abundance and predictive covariates (Figure 5) suggested that
bearded seal densities were highest in grid cells without landfast ice, in cells with sea ice concentration near
0.8, and in cells with low snow depths. However, the most important predictor (as judged from the magni-
tude of y-axis value changes) was northing. Along with easting, these two covariates do not imply specific
drivers of distributions. They may, however, highlight areas of low or high abundance that are not well pre-
dicted by the other, presumably more ecologically relevant covariates such as sea ice concentration.

Figure 2. Remotely sensed sea ice concentration and estimates of ringed and bearded seal abundance (individuals per 25 × 25-km grid cell) in the Chukchi Sea on three dates at the beginning, middle, and end of the April–May, 2016 aerial surveys. Each column represents a different survey date, while rows correspond to sea ice concentration, ringed seal abundance, and bearded seal abundance. Note that the color scaling for seal abundance is nonlinear, depicting densities of zero in the darkest (indigo) color tone and very low densities in the second tone.

Figure 3. Estimated smooth functions (dark lines) for the relationship of ringed seal abundance to environmental and physiographic variables, on the multinomial logit scale, together with 95% confidence intervals (shaded areas). Note that the northing variable has larger values for locations farther south. Northing and easting variables were standardized to their means (so that a value of 1.0 roughly corresponds to the centroid of the study area). Landfast ice and sea ice values are proportions of area covered.

Figure 4. Proportion of ringed seals estimated to be basking on ice (i.e. not hidden in a subnivean lair) given that they were hauled out while aerial surveys were being conducted, together with 95% confidence intervals (shaded area). The model was penalized if the curve did not reach 1.0 by the date on which the MDSDA algorithm indicated the snow layer was saturated ('snow melt onset', 0 on x-axis).

Figure 5. Estimated smooth functions (dark lines) for the relationship of bearded seal abundance to environmental and physiographic variables, on the multinomial logit scale, together with 95% confidence intervals (shaded areas). Note that the northing variable has larger values for locations farther south. Northing and easting variables were standardized to their means (so that a value of 1.0 roughly corresponds to the centroid of the study area). Landfast ice and sea ice values are proportions of area covered.

3.1 Comparison with historical estimates

Ringed seal abundance estimates reported for 1999 and 2000 in the eastern Chukchi Sea (blue shaded area in Figure 1) were 252,488 (SE 47,204) and 208,857 (SE 25,502), respectively (Bengtson et al. 2005). We re-constructed 95% log-based confidence intervals for those estimates as (175,576–363,091) and (164,550–265,095). For comparison, our ringed seal abundance estimate from the 2016 survey, in just the Bengtson et al. (2005) survey region (blue shaded area in Figure 1) at the end of May, was 394,411 (95% log-based CI: 318,449–448,493).

Bearded seal density estimates (Bengtson et al. 2005), adjusted with our availability estimates for late May ($\hat{a} = 0.36$, SE 0.03) led to abundance estimates of 25,076 (95% CI: 7,480–84,073) and 50,992 (95% CI: 30,207–86,079) for 1999 and 2000, respectively. Restricting abundance estimates from our 2016 surveys to grid cells with centroids in the Bengtson et al. (2005) study area (blue shaded area in Figure 1) led to a point estimate for late May of 72,182 (95% CI: 55,894–93,216).

4 Discussion

Our effort was the first to survey ringed and bearded seals over the entire Chukchi Sea, which was only possible through international cooperation. Conducting a sensor-based survey (as opposed to a survey with human observers) allowed us to fly faster and at higher altitudes, increasing the aircraft range and decreasing seal disturbance. We also used auxiliary data from TDRs to estimate availability, and employed protocols that allowed us to directly estimate the detection probability of our thermal sensing method and the proportion of seals that fled into the water before they could be detected. These are all improvements over previous surveys in the Chukchi Sea, which were limited in spatial scope and were able to only partially address detection and availability bias (e.g., Frost et al. 1988; Frost et al. 2004; Bengtson et al. 2005). In addition to providing a better abundance estimate, the results improve our understanding of ringed and bearded seal habitat composition, ecology, and prospects for persistence in rapidly warming Arctic marine ecosystems.

4.1 Factors influencing ringed and bearded seal distribution

Ringed seals were primarily found in the southern Chukchi Sea, concentrated in landfast ice in and around Kotzebue Sound; there were more moderate ringed seal densities in landfast ice elsewhere, along the northwest coast of Alaska and northeast coast of Russia near Chaunskaya Bay, Chukotka. Bearded seals also were primarily found in the southern Chukchi Sea, occupying broken pack ice with leads, similar to where they had been observed in past surveys. The broad geographic distribution patterns for both species in the U.S. portion of the Chukchi Sea match those found during surveys in 1999 and 2000 (Bengtson et al. 2005).

4.1.1 Ringed seals

Influence of snow depth on ringed seal density

We found that ringed seal densities in spring were related to remotely sensed snow depth values, con-
sistent with *in situ* studies of snow depth and the distribution of lairs. Resting lairs—also called simple, or
haul-out lairs—are single-chambered structures used mostly by adults. Pups are typically found in multi-
chambered structures—called pupping, pup, or birth lairs—that may include small tunnels excavated by the
pups during their rearing period (e.g., Kelly et al. 1986). An analysis of ringed seal structures surveyed along
the Alaskan Chukchi Sea coast during the 1980s found that the minimum snow depth at intact pupping lairs
was 37 cm at Ledyard Bay and 58 cm in Kotzebue Sound, and the average depth at pupping lairs over the two
sites was 75 cm (Hauser et al. 2021). Kelly et al. (2010b) reviewed other studies from Alaska, Canada, Russia,
and Svalbard in which snow depths were measured at different types of ringed seal structures. Snow drifts
to depths of 45 cm or more were found at resting lairs, and at least 50 cm—or more commonly, over 65 cm—
were found at pupping lairs. Kelly et al. (2010b) concluded that average snow depths on flat ice of at least 20
to 30 cm are required for drifts to form along ice deformations and reach adequate depths for excavation of
lairs. In our results, ringed seal densities increased strongly with snow depth values up to 20 cm and plat-
eaued between 20 and 30 cm, in close concurrence with the *in situ* findings that less than 20 cm average
snow depth is suboptimal for viable lairs. The broad confidence intervals for ringed seal densities at average
snow depths greater than 25 cm may reflect that the seals use areas of greater snow depths but are not
necessarily found at high densities in all cells with deep snow; other habitat dimensions are also likely to be
important. However, the narrow intervals and strong increase in seal densities as average snow depths in-
creased up to 25 cm seem supportive of ringed seals' selection of breeding habitat with sufficient snow for
lair construction and integrity.

Remotely sensed snow depth products that provide information over the entire range of ringed seal hab-
itat have only recently been tested for consistency over broad regions with the average snow depth require-
ments for lairs as derived from local, *in situ* studies. Lindsay et al. (2021) related the NASA Cryosphere data-
base of 25-km gridded snow depths on sea ice (Markus and Cavalieri 1998; Comiso et al. 2003) to the raw
counts of ringed seals detected in our surveys. There were modest but imprecise increases in counts of seals
as with snow depths increased to about 25 cm for pups, and 35 cm for all ages combined. Lindsay et al. (2023)
used an uncrewed aircraft to conduct photographic surveys of ringed seals on landfast ice near Kotzebue,
Alaska in the spring of 2019. By relating seal counts to Landsat 8 210-m gridded pixel brightness values (an
index of surface roughness) from May 6, 2019, and relating the brightness values to *in situ* April snow depths,

they found indirectly that the highest seal counts occurred in areas where average April snow depth was 25 cm. In the current study, we related our ringed seal density estimates, rather than raw counts alone, to the NASA snow depth index, an estimate of the average depth over each 25 × 25-km grid cell. Our finding that ringed seal densities increased strongly with remotely sensed average snow depth up to 25 cm supports the conclusion from *in situ* measurements for this important habitat feature across the Chukchi Sea, an extensive region of the Pacific Arctic.

The capability to remotely monitor a snow-depth index of lair habitat quality over broad regions of the Arctic ringed seal's range is likely to be helpful for assessing the subspecies' status in the warming climate. A more complete understanding of lair habitat though, would include consideration of the timing and extent of sea ice formation, and patterns in ice surface structure, wind, air temperature, and precipitation. These factors interact to influence the blowing snow flux and the drift formation in the lee of ice surface structures (Kovacs et al. 2024); they are likely to vary among broad regions such as the Bering, Chukchi, and Beaufort Seas. We suspect that additional examples of aerial surveys for ringed seals, coupled with remotely sensed snow depths, may reveal regional variation in the relationship between ringed seal density and average snow depth alone.

Possible influence of covariates not in our model

Our results include the geographical covariates easting and northing that may identify low or high abundance areas that are not well predicted by other, presumably more ecologically relevant covariates in the model. They may also reflect influential covariates that are missing. We had no data to represent the distributions of ringed or bearded seal prey, which could influence seals' distribution and density. We note, though, that the high densities of ringed seals we observed in Kotzebue Sound are a regular winter and spring occurrence reported by Indigenous hunters (Lindsay et al. 2023) that may reflect dependable winter and spring prey availability. Saffron cod (*Eleginus gracilis*) is highly prevalent in Alaska ringed seal diets (Quakenbush et al. 2020). Although saffron cod spawning grounds are not well known in Alaska, Kotzebue Sound and the adjacent southeast Chukchi Sea include large areas of shallow, nearshore waters that are typically ice covered from December to March—consistent with saffron cod spawning habitat (Wolotira 1985; Vestfals et al. 2019). Using survey catches of saffron cod larvae in the Chukchi Sea, coupled with biophysical transport models, Deary et al. (2021) suggested that Kotzebue Sound is an important saffron cod spawning ground. With the peak of spawning occurring in winter, gravid saffron cod would be ideal prey for ringed seals, especially pregnant females, which give birth to pups in March and April. The combination of dependable seasonal ice cover—much of which is landfast—and a reliable prey base could explain the high densities of ringed seals observed in Kotzebue Sound and southeast Chukchi Sea surveys over several decades (Frost

527 et al. 1988; Bengtson et al. 2005, this study), and the reliance of Indigenous peoples of the region for winter
528 sustenance over millennia (Lucier and VanStone 1991).

4.1.2 Bearded seals

In past aerial surveys during March–May, primarily in the Bering Sea, bearded seal densities were highest
in sea ice concentrations greater than 25%. There were density peaks at concentrations of 70–90%
(Simpkins et al. 2003) and 67% (Conn et al. 2014), though Ver Hoef et al. (2014) found only a weak increase
in the probability of occurrence in concentrations above 25% and no evidence for an effect of ice concentra-
tion on bearded seal density. A sample of juvenile bearded seals tagged in 2004–2009 and tracked during
spring in the Bering Sea seemed to prefer ice concentrations similar to those identified from aerial surveys
(~80%; Cameron et al. 2018) or lower (~50–60%; Breed et al. 2018). For juvenile bearded seals tagged in a
more recent and warmer period of 2014–2018, there was only weak selection by juveniles in spring, for ice
concentration <50%, perhaps reflecting a shift in the distribution of ice concentrations or an indication that
other habitat features such as foraging quality are more important (Olnes et al. 2021). Our finding of a strong
increase in bearded seal densities with sea ice concentration up to a peak at 80%, and a slight decrease there-
after, agrees more closely with previous aerial survey results (based on all age classes) than with results
from tracking juveniles. Together with the strong decline we observed in bearded seal density with increas-
ing proportions of landfast ice in a survey grid cell, our results are also consistent with the many qualitative
natural history descriptions of bearded seals' preference for broken, drifting ice, as well as their typical (but
not complete) avoidance of the heaviest concentrations of pack ice and extensive landfast ice (Fedoseev
1965; Burns and Harbo 1977; Burns and Frost 1979; Burns 1981; Smith 1981; Fedoseev 1984; Nelson et al.
1984; Kingsley et al. 1985; Cameron et al. 2010). While we also observed a decline in bearded seal density
with increasing snow depth, we are not aware of prior studies documenting a negative association with snow
depth. Lacking an ecological explanation based on bearded seal natural history, this result may serve as a
reminder that models with multiple covariates sometimes yield illusory associations. For example, if there is
a latitudinal (northing) gradient in snow depth, these two variables would not be independent and thus dif-
ficult to interpret. Deeper snow depth over areas of landfast ice could also contribute to the apparent asso-
ciation between bearded seal densities and areas of shallower snow depth.

4.2 Assumptions and limitations

555   We modeled ringed seal availability as a function of a snow melt index (i.e., Belchansky et al. 2004) that
556 was previously found to be related to an increase in the proportion of seals basking (Kelly et al. 2006). Our

model could accommodate a gradual transition away from lair use—including intermittent lair use by individuals—but it could not account for a portion of ringed seals that may continue to spend some time in lairs after the snow melt index reaches saturation. We had no means for estimating the uncertainty associated with this assumption, and we caution against interpreting our confidence intervals too rigidly; they are likely too narrow for ringed seals despite having included several more sources of uncertainty in our model than previous examples of aerial surveys for seals. There may also be negative bias in abundance stemming from this assumption because some ringed seals have been observed using lairs after snow melt was under way. Future research should be devoted to refining ringed seal availability estimates through analysis of existing data (e.g. Lindsay et al. 2021) and field work (e.g. applying additional TDRs, investigating relationships between internal lair conditions and snow melt onset, and examining variations in emergence timing as a function of environmental covariates).

The timing of our surveys of the Chukchi Sea in 2016 was, in part, a result of our desire to combine Chukchi Sea estimates with Bering Sea estimates from a previous survey (e.g. Conn et al. 2014) and estimates from surveys planned for the Beaufort Sea, in order to develop total estimates for the Beringia DPS of bearded seals and for ringed and bearded seals in U.S. waters (i.e. the Alaska stocks for assessment under the Marine Mammal Protection Act; Young et al. 2023). In particular, we wanted to avoid surveying in June when large numbers of seals migrate north from the Bering Sea to the Chukchi Sea (Melnikov 2017; Melnikov 2022). However, this approach may not have been the best timing for ringed seals, as VHF radio-tagging data and monitoring of ringed seal lairs in the Beaufort Sea (Kelly et al. 2006; Kelly et al. 2010a; Kelly et al. 2010b) suggest that a large proportion of seals were in lairs and thus were unavailable for counting when our surveys began in early April. Smith (1981) suggested that optimal timing of bearded seal surveys might be late April–May when mothers are accompanied by pups, at least a month earlier than the best time for ringed seal surveys. If ringed seals are of higher priority in future surveys, we suggest that more intensive surveys in the latter half of May and early June may be more useful, as it would remove a large source of uncertainty about availability in April and early May. Also, aircraft capable of surveying at sufficient speed could allow coverage of both the Bering and Chukchi seas in a single survey period, reducing the potential for bias induced by seals migrating north during the survey.

We avoided surveying small portions of the study area that occurred within 48 km (30 miles) of the Alaskan villages of Utqiaġvik, Wainwright, Point Lay, and Point Hope, to reduce the potential for disturbance of traditional bowhead whale (*Balaena mysticetus*) harvest activities. Coastal Indigenous village sites in the Arctic are recognized as places of dependable, high densities of marine resources. Therefore, not surveying near these villages could potentially have biased our estimates low. Our analysis estimated that 4% of the

totals for both ringed and bearded seals occurred within these avoided areas, which composed 2.1% of the

total study area. Therefore, because our estimated seal densities in these avoided areas were roughly twice

the overall average for each species, and because these areas were very small in comparison to the total study

area, we consider that excluding them was unlikely to have caused significant bias in our overall abundance

estimates. Nevertheless, it would be beneficial in future surveys to incorporate survey effort in those areas

if it can be done in coordination with the communities to avoid interfering with traditional subsistence ac-

tivities.

4.3 Implications for future status of ringed and bearded seals

Relationships between snow depth and ringed seal vital rates

Several studies have found associations between regional snow depth and ringed seal survival, reproduc-

tion, or indices of condition (e.g. see review by Kelly et al. 2010b). Hammill and Smith (1991) found that a

decrease from 23 cm to 10 cm in average snow depth in Barrow Strait, Canada in the mid-1980s coincided

with an increase in the predation rate by polar bears from 0.1 to 0.4 seals/km^2. An index of recruitment from

seals harvested in western Hudson Bay during 1991–2001 was sharply lower for cohorts born in years with

spring average snow depths less than about 32 cm (Ferguson et al. 2005). Chambellant et al. (2012) found

that an index of pup survival was optimized between about 25 and 52 cm of snow depth (peak at 38 cm)

measured on the ground near their study area in western Hudson Bay during 1991–2006. Iacozza and

Ferguson (2014) found that variability in remotely sensed spring snow depth in western Hudson Bay from

2003–2010 was negatively related to the proportion of young-of-the-year in the autumn harvest at Arviat,

Canada; the number of 'dry snow cover days', which is an index of the duration of winter snow cover, was

positively related to pup survival, and together, the two indices accounted for 82% of the variation in the

young-of-the-year harvest numbers. Hamilton et al. (2018) found that ringed seals hauled out less frequently

and—during winter—in shorter bouts and for less time overall, after the cessation of annual formation of

landfast ice that occurred in 2006 around Svalbard. Although the changes were not directly linked to vital

rates, the implication was that without landfast ice and its associated snow cover, ringed seals hauled out

less in lairs, with likely impacts on their energy budgets and predation risk. But also in Svalbard, where large

areas have not had regular seasonal sea ice since 2006, an interdecadal synthesis of several vital rate indices

found no clear pattern consistent with climate-related hypotheses (Andersen et al. 2021). Potential explana-

tions are 'life history plasticity' or that small-scale variations in breeding habitat remain suitable in the lim-

ited area where the hunted samples were collected (Andersen et al. 2021).

Around Alaska and the western Canadian Arctic, associations among ringed seal vital rates or condition and snow depth on sea ice have not been assessed, though some studies have made comparisons to ice-covered area, the length of the ice-covered period, or other statistics that may be related to spring snow depth. Crawford et al. (2015) concluded that young ringed seals in the Chukchi and northern Bering seas grew faster during 2003–2012, which was characterized as a period of reduced sea ice, compared to an earlier period with more ice, 1974–1984. Ringed seals harvested at three northern Bering Sea and one southern Chukchi Sea villages included higher proportions of pups during the recent period, as well. A continuation of the study, comparing the period 2000–2015 with 2016–2020 (Quakenbush et al. 2020), found that growth of ringed seals born in the most recent period did not vary from average, except those born in 2017. Blubber thickness (i.e. body condition) was below average in 2017–2018 but returned to average in 2019. The years of 2017 and 2018 were extremely warm in the Pacific Arctic, with low sea ice coverage and impacts documented throughout the ecosystem (Huntington et al. 2020).

In the western Canadian Arctic, ringed seals were monitored in the Amundsen Gulf and western Prince Albert Sound during 36 years between 1971 and 2019 (Harwood et al. 2012; Harwood et al. 2020). These areas were characterized as prime habitat, with sea ice that typically persists well beyond the ringed seal pup rearing period. Despite long-term trends toward later autumn freeze-up and earlier spring sea ice clearance that coincided with a declining trend in seal body condition, the poorest body condition and lowest indices of reproduction occurred in the years with the latest ice break-up, suggesting more complex processes than a simple directional influence of Arctic warming on the seals' vital rates.

In general, the studies summarized above have examined suites of indices for vital rates but have not had the means to detect overall demographic impacts, such as changes in abundance or loss of significant portions of breeding range. Although monitoring indices of vital rates or other demographic parameters provides insight about mechanisms of response to environmental variability and climatic trends, sampling considerations are often complex, as are interpretations of the results; the need remains for complementary, direct means of monitoring abundance and distribution of breeding populations, such as count-based surveys.

Trends in snow depth on sea ice

Snow depth on Arctic sea ice has decreased, prompting the question of whether ringed seal populations have already been impacted by diminished or degraded habitat. Warming in the upper layer of Arctic waters has caused later onset of sea ice formation in autumn, which in turn reduces snow accumulation on the ice because more snow falls into open water (Webster et al. 2018). The result has been a clear multidecadal

decline in average snow depth on sea ice. For example, Webster et al. (2014) found that snow on Chukchi and Beaufort sea ice in 2009–2013 was about 56% thinner than during the latter half of the 20th century. March–May snow depth on sea ice at three long-term monitoring sites in the Canadian Arctic declined by 0.1, 1.7, and 0.6 cm/decade from 1960–2020 (but increased by 0.1 cm/decade at a fourth site; Lam et al. 2022). Lee et al. (2021) found an Arctic-wide average reduction in snow depth from the period 1954–1991 to the period 2003–2020, but with a statistically significant positive trend of ~0.6 cm/year on multi-year ice, and negative trends (~−0.4 cm/year) over mixed and first-year ice, the habitat for most Arctic ringed seals. Stroeve et al. (2020), using a detailed physical model of snow depth driven by weather reanalysis products, estimated a loss of 1.9–3.0 cm/decade from 1980–2016 in the Chukchi Sea. A version of the same model was used to estimate declining trends over the past 2–3 decades in snow depth, blowing snow flux, and ringed seal lair habitat around Svalbard (Kovacs et al. 2024). Dou et al. (2021) showed that rain-on-snow events in the ERA5 reanalysis have shifted as much as 4–6 days earlier per decade in some regions; there have been more rain-on-snow events in spring; and the increased rain-on-snow events have caused a reduction in spring snow depth of 0.5 cm/decade over the Arctic Ocean since 1980 and up to 2 cm/decade in the Kara-Barents seas and Canadian Arctic Archipelago. The observed open-water period in the Chukchi Sea has increased by about 2 days/year over the period 1979–2013 (Crawford et al. 2021). The length of the open water period is particularly relevant to habitat quality for Arctic ringed seals because snow depth and quality is influenced by the ice freeze-up and melt onset dates.

Reference levels for climate change impacts on ringed and bearded seals in the Chukchi Sea

Given the observed and ongoing declines in Arctic sea ice and snow depth, do our estimates in 2016 provide a reference for assessing climate-driven population trends of ringed and bearded seals in the Chukchi Sea? The large numbers of both species that we estimated and the apparent increases since a previous, smaller scale survey are encouraging, especially given the current designations of both species as 'Threatened' under the U.S. Endangered Species Act (National Marine Fisheries Service 2012a; National Marine Fisheries Service 2012b). Our overall ringed and bearded seal densities in 2016 were 2–3 times and 1.9 times higher, respectively, than the average of those observed during 1999 and 2000 in a smaller scale study in the eastern Chukchi Sea (Bengtson et al. 2005). However, increases in Chukchi Sea ringed and bearded seal estimates could be due to factors other than population growth, such as shifting of breeding locales north from the Bering Sea, shifts to earlier northward spring migration from the Bering Sea, or differences in survey methods. For instance, although ringed seal densities reported by Bengtson et al. (2005) were adjusted for availability, the double-observer method they used to correct for incomplete detection is now recognized to be biased if assumptions of independence are not met, which is often the case in aerial surveys (e.g., Conn et

al. 2012; Burt et al. 2014). Further, their estimates did not take into account disturbance of seals into the water by the aircraft, which flew at an altitude of 91 m, compared to altitudes of 250–300 m during our survey. Both of these factors would negatively bias abundance estimators from Bengtson et al. (2005). We took steps to accommodate such sources of detection bias, making future comparisons more straightforward. However, the differences in survey protocols between our sensor-based surveys and prior human-observer surveys currently pose difficulties for comparing with past estimates.

The studies of vital rates that we summarized earlier indicate that Arctic ringed seals in some parts of their range—Hudson Bay and around Svalbard—have responded to years of poor snow cover on sea ice in ways that would be expected to impact vital rates and reduce population size under long-term Arctic warming. Both of those regions are peripheral, rather than core ringed seal habitat, due to low latitude (Hudson Bay) or oceanographic features that mediate Arctic conditions (Svalbard). In contrast, where ringed seals have been studied in the core areas of their range, they have yet to exhibit consistent negative impacts from climatic trends. In the Alaska Chukchi Sea and in the western Canadian Arctic, some of the most notable reductions in indices of vital rates and condition have actually come during colder periods or years with higher ice coverage and later ice breakup (Crawford et al. 2015; Harwood et al. 2020). We suggest that our spring 2016 estimate of ringed seal abundance in the Chukchi Sea represents a baseline reference level for assessing future climate-related changes in abundance. Similarly, bearded seals in Alaska have not exhibited declines in body condition, growth, or reproduction compared to earlier periods (Crawford et al. 2015; Quakenbush 2020), and they are not dependent upon snow cover for shelter of their young. We are not aware of other scientific evidence for climate-driven impacts on bearded seals of the Chukchi Sea, nor of impacts documented by Indigenous hunters and observers in the region. Therefore, we suggest that our bearded seal abundance estimate is also a reasonable reference point for assessment of climate-related impacts to that species.

How ringed and bearded seals will respond to future climate change remains uncertain. Will seals migrate northward earlier as spring sea ice diminishes in the Bering Sea? Will there be enough prey of sufficient quality to accommodate them? Will snow depths on sea ice, which are predicted to continue declining over the next century (Hezel et al. 2012; Webster et al. 2021), be sufficient for ringed seal lair construction? Although current populations in the Chukchi Sea appear healthy, there is cause for concern about how they will cope with a rapidly warming Arctic. The upper and lower bounds of our 95% confidence intervals for ringed and bearded seal abundance were about 25% above and 20% below the point estimates, respectively. These levels of precision provide reasonable expectations for detecting substantial changes in Arctic seal populations over broad regions using methods like those we have demonstrated in the Chukchi Sea. For example,

with a coefficient of variation (CV) of 13% that reflects the precision of our bearded seal estimate, a 25% decline in the population would have a 33% chance of being detected in a subsequent survey, whereas a 50% decline would be detected 97% of the time (95% two-sided z-test). Our ringed seal estimate, with a CV of 11%, was nominally more precise, but was based on additional assumptions that are not reflected in the precision estimate. We consider the two species to be similar for the purposes of detecting future trends. Increasing the number of subsequent surveys would increase the power to detect changes in a regression framework. Future instrument-based surveys in the Chukchi Sea and other areas (e.g. Bering and Beaufort seas) will help us measure ringed and bearded seals' responses to the dramatic changes occurring in the Arctic.

5. Acknowledgements

We are grateful to Gavin Brady, Sarah Brown, Shawn Dahle, and Carter Johnson for species identification and detection rate estimation; John Bengtson for project facilitation as co-Convenor of the Working Group on Marine Mammals under Area V of the U.S.-Russia Agreement on Cooperation in the Field of Environmental Protection and Natural Resources; Capt. Robert Dodson, 1st Officer Sarah Morris and Aircraft Mechanic James DeLuca ("JD") of Dynamic Aviation for safe operation of the King Air A90; John Cabrer of MoviTHERM for collaboration on Skeyes software and the hotspot detection method; the late Craig George for logistical support and coordination with whaling activities in Utqiaġvik; Paul Wade for helpful comments on the manuscript; Dave Anderson and Rich Lomire of the National Weather Service for assistance with housing; the U.S. Fish and Wildlife Service for providing cooled LWIR cameras; the Russian non-profit organization, Marine Mammal Council, provided logistical support for aerial surveys; N.A. Chernook and V.V. Asyutenko for processing Russian survey images; and D. M. Glazov for contribution to the field effort. The organization of this international collaboration would not have been possible without the facilitation by Vladimir N. Burkanov of North Pacific Wildlife Consultants.

The projects to deploy satellite-linked TDRs by ADFG, and NSB-DWM were successful because of support and cooperation of the Ice Seal Committee, the NSB Fish and Game Management Committee, and coastal Native communities, especially the Native Village of Barrow, Native Village of Kotzebue, Native Village of Koyuk, and the Native Village of Saint Michael. The success of the studies was due to assistance from subsistence hunters—especially John Goodwin, Merlin Henry, Joe Skin, Isaac Leavitt, Aaron Morris, Bobby Sarren, and Billy Adams—and biologists, especially Mark Nelson, Anna Bryan, Ryan Adam, Rowenna Gryba, and Jason Herreman. Expertise and leadership in the field for tagging seals in Canadian waters was provided by Tom Smith; community crews from Tuktoyaktuk, Inuvik and Paulatuk, Northwest Territories; and Brendan Kelly and John Moran of the University of Alaska.

6. Competing interests

The authors declare there are no competing interests. Reference to trade names does not imply endorsement by the National Marine Fisheries Service, NOAA.

7. Author contributions

Conceptualization: Peter Boveng, Vladimir Chernook; Data curation: Stacie Koslovsky; Formal analysis: Paul Conn, Joshua London, Brett McClintock, Irina Trukhanova; Funding acquisition: Peter Boveng, Vladimir Chernook; Investigation: Vladimir Chernook, Erin Moreland, Paul Conn, Cynthia Christman, Benjamin Hou, Denis Litovka, Erin Richmond, Alexander Vasiliev, Amy Willoughby; Methodology: Peter Boveng, Vladimir Chernook, Paul Conn, Benjamin Hou, Jessica Lindsay, Erin Moreland, Alexander Vasiliev; Project administration: Michael Cameron, Erin Moreland; Resources: Justin Crawford, Lois Harwood, Nikita Platonov, Lori Quakenbush, Andrew Von Duyke; Software: Benjamin Hou, Stacie Koslovsky; Supervision: Peter Boveng, Vladimir Chernook; Validation: Paul Conn, Stacie Koslovsky, Erin Moreland; Visualization: Paul Conn, Joshua London, Erin Richmond; Writing - original draft: Peter Boveng, Paul Conn; Writing - review and editing: All authors.

8. Funding statement

The primary funding for the ChESS surveys and analyses was provided by NMFS, including a portion supported by the Joint Institute for the Study of the Atmosphere and Ocean (JISAO) under National Oceanic and Atmospheric Administration (NOAA) Cooperative Agreement NA15OAR4320063, Contribution No. 2024-1368. The North Pacific Research Board, grant NA17NMF4720289, project 1813, provided support for post-processing of Russian survey data. The projects to deploy satellite-linked TDRs by ADFG, and NSB-DWM were funded by a U.S. Department of the Interior, Tribal Wildlife Grant for Federally Recognized Tribes; Shell Exploration and Production Co.; NSB-Shell Baseline Studies Program; National Fish and Wildlife Foundation with funding from Conoco Phillips; the Native Village of Kotzebue; NMFS Alaska Region; NOAA [NA05NMF4391187 and NA08NMF4390544]; and the Bureau of Ocean Energy Management [M13PC00015]. In Canadian waters, tagging studies were funded by Environmental Studies Research Fund; Dept. of Indian Affairs and Northern Development; Dept. of Fisheries and Oceans (DFO), the Fisheries Joint Management Committee; and DFO Program of Energy Research and Development (PERD-NPOL #122-22 & #122-25), Polar Continental Shelf Project.

9. Data Availability

All data and code will be made freely available as a GitHub repository upon acceptance for publication in
a peer-reviewed journal.

10. References

Andersen, M., Kovacs, K.M., and Lydersen, C. 2021. Stable ringed seal (*Pusa hispida*) demography despite
significant habitat change in Svalbard, Norway. Polar Research 40:1-14.

Belchansky, G.I., Douglas, D.C., Mordvintsev, I.N., and Platonov, N.G. 2004. Estimating the time of melt onset
and freeze onset over Arctic sea-ice area using active and passive microwave data. Remote Sensing of
Environment 92:21-39. doi:10.1016/j.rse.2004.05.001.

Bengtson, J.L., Hiruki-Raring, L.M., Simpkins, M.A., and Boveng, P.L. 2005. Ringed and bearded seal densities
in the eastern Chukchi Sea, 1999–2000. Polar Biology 28:833-845.

Borchers, D.L., Laake, J.L., Southwell, C., and Paxton, C.G.M. 2006. Accommodating unmodeled heterogeneity
in double-observer distance sampling surveys. Biometrics 62:372-378. doi:10.1111/j.1541-
0420.2005.00493.x.

Breed, G.A., Cameron, M.F., Ver Hoef, J.M., Boveng, P.L., Whiting, A., and Frost, K.J. 2018. Seasonal sea ice
dynamics drive movement and migration of juvenile bearded seals *Erignathus barbatus*. Marine Ecology
Progress Series 600:223-237.

Buckland, S.T., Anderson, D.R., Burnham, K.P., Laake, J.L., Borchers, D.L., and Thomas, L. 2001. Introduction
to distance sampling. Oxford University Press, Oxford, UK. 432 pp.

Burnham, K.P., Anderson, D.R., White, G.C., Brownie, C., and Pollock, K.H. 1987. Design and analysis methods
for fish survival experiments based on release-recapture. American Fisheries Society Monograph 5.

Burns, J.J. 1981. Bearded seal *Erignatus barbatus* Erxleben, 1777. *In* Handbook of Marine Mammals Volume
2: Seals. *Edited by* S.H. Ridgway and R.J. Harrison. Academic Press, New York, NY. pp. 145-170.

Burns, J.J., and Frost, K.J. 1979. The natural history and ecology of the bearded seal, *Erignathus barbatus*.
Contract #02-5-002-53. Alaska Department of Fish and Game, Fairbanks, AK. 77 pp.

Burns, J.J., and Harbo, S.J. 1972. An aerial census of ringed seals, northern coast of Alaska. Arctic 25:179-290.

Burns, J.J., and Harbo, S.J. 1977. An aerial census of spotted seal, *Phoca vitulina largha*, and walruses,
Odobenus rosmarus, in the ice front of Bering Sea. *In* Environmental Assessment of the Alaskan
Continental Shelf. Quarterly Reports of Principal Investigators April-June 1977. Vol. 1. U.S. Department
of Commerce, NOAA and the U.S. Department of Interior, Bureau of Land Management, Boulder, CO. pp.
58-132.

Burt, M.L., Borchers, D.L., Jenkins, K.J., and Marques, T.A. 2014. Using mark–recapture distance sampling
methods on line transect surveys. Methods in Ecology and Evolution 5:1180-1191. doi:10.1111/2041-
210X.12294.

Cameron, M.F., Bengtson, J.L., Boveng, P.L., Jansen, J.K., Kelly, B.P., Dahle, S.P., Logerwell, E.A., Overland, J.E.,
Sabine, C.L., Waring, G.T., and Wilder, J.M. 2010. Status review of the bearded seal (*Erignathus barbatus)*.
NOAA Technical Memorandum NMFS-AFSC-211. U.S. Department of Commerce, Seattle, WA. 246 pp.
http://www.alaskafisheries.noaa.gov/protectedresources/seals/ice/bearded/statusreview10.pdf.

Cameron, M.F., Frost, K.J., Ver Hoef, J.M., Breed, G.A., Whiting, A.V., Goodwin, J., and Boveng, P.L. 2018. Habitat
selection and seasonal movements of young bearded seals (*Erignathus barbatus*) in the Bering Sea. PLoS
ONE 13:e0192743. doi:10.1371/journal.pone.0192743.

Cavalieri, D.J., Parkinson, C.L., Gloersen, P., and Zwally, H.J. 1997. Arctic and Antarctic Sea Ice Concentrations
from Multichannel Passive-Microwave Satellite Data Sets: October 1978-September 1995 - User's Guide.
NASA Technical Memorandum. 17 pp.
https://nsidc.org/sites/default/files/nasa20technical20memorandum20104647_0.pdf.

Chambellant, M., Stirling, I., Gough, W.A., and Ferguson, S.H. 2012. Temporal variations in Hudson Bay ringed
seal (*Phoca hispida*) life-history parameters in relation to environment. Journal of Mammalogy 93:267-
281. doi:10.1644/10-mamm-a-253.1.

Chernook, V.I., Grachev, A.I., Vasiliev, A.N., Trukhanova, I.S., Burkanov, V.N., and Solovyev, B.A. 2014. Results
of instrumental aerial survey of ice-associated seals on the ice in the Okhotsk Sea in May 2013. Izvestiya
TINRO 179:158-176.

Chernook, V.I., Kuznetsov, N.V., Shafikov, I.N., Krivushchenko, P.A., and Egorov, S.A. 2000a. Methods of
automated counting of harp seals on aerial imagery ICES/NAFO Working Group on Harp and Hooded
Seals Copenhagen. 5 pp.

Chernook, V.I., Timoshenko, Y.K., Meisenheimer, P., Innes, S., Kuznetsov, N.V., and Egorov, S.A. 2000b.
Assessment of the White Sea harp seal pups in 1998. ICES/NAFO Working Group on Harp and Hooded
Seals Copenhagen, Denmark. 10 pp.

Chernook, V.I., Trukhanova, I.S., Vasiliev, A.N., Grachev, A.I., Litovka, D.I., Burkanov, V.N., and Zagrebelny, S.V.
2018. Abundance and distribution of phocid seals on ice in the western Bering Sea in spring, 2012–2013.
Izvestiya TINRO 192:74-88. doi:10.26428/1606-9919-2018-192-74-88.

Chernook, V.I., Trukhanova, I.S., Vasiliev, A.N., Litovka, D.I., Glazov, D.M., and Burkanov, V.N. 2019. First
instrumental aerial survey of ringed seals (*Pusa hispida*) and bearded seals (*Erignathus barbatus*) in the
Russian zone of the Chukchi and East-Siberian Seas in spring 2016 Izvestiya TINRO 199:152-162.

Comiso, J.C., Cavalieri, D.J., and Markus, T. 2003. Sea ice concentration, ice temperature, and snow depth using
AMSR-E data. IEEE Transactions on Geoscience and Remote Sensing 41:243-252.

Conn, P.B., Johnson, D.S., Hoef, J.M.V., Hooten, M.B., London, J.M., and Boveng, P.L. 2015. Using spatiotemporal
statistical models to estimate animal abundance and infer ecological dynamics from survey counts.
Ecological Monographs 85:235-252. doi:10.1890/14-0959.1.

Conn, P.B., Laake, J.L., and Johnson, D.S. 2012. A hierarchical modeling framework for multiple observer
transect surveys. PLoS ONE 7:e42294. doi:10.1371/journal.pone.0042294.

Conn, P.B., McClintock, B.T., Cameron, M.F., Johnson, D.S., Moreland, E.E., and Boveng, P.L. 2013.
Accommodating species identification errors in transect surveys. Ecology 94:2607-2618.
doi:10.1890/12-2124.1.

Conn, P.B., Moreland, E.E., Regehr, E.V., Richmond, E.L., Cameron, M.F., and Boveng, P.L. 2016. Using
simulation to evaluate wildlife survey designs: polar bears and seals in the Chukchi Sea. Royal Society
Open Science 3:150561. doi:10.1098/rsos.150561.

Conn, P.B., Thorson, J.T., and Johnson, D.S. 2017. Confronting preferential sampling when analysing
population distributions: diagnosis and model-based triage. Methods in Ecology and Evolution 8:1535-
1546. doi:10.1111/2041-210X.12803.

Conn, P.B., Ver Hoef, J.M., McClintock, B.T., Moreland, E.E., London, J.M., Cameron, M.F., Dahle, S.P., and Boveng,
P.L. 2014. Estimating multispecies abundance using automated detection systems: ice-associated seals
in the Bering Sea. Methods in Ecology and Evolution 5:1280-1293. doi:10.1111/2041-210X.12127.

Crawford, A., Stroeve, J., Smith, A., and Jahn, A. 2021. Arctic open-water periods are projected to lengthen
dramatically by 2100. Communications Earth & Environment 2:1-10.

Crawford, J.A., Frost, K.J., Quakenbush, L.T., and Whiting, A. 2012. Different habitat use strategies by subadult
and adult ringed seals (*Phoca hispida*) in the Bering and Chukchi seas. Polar Biology 35:241-255.
doi:10.1007/s00300-011-1067-1.

Crawford, J.A., Frost, K.J., Quakenbush, L.T., and Whiting, A. 2019. Seasonal and diel differences in dive and
haul-out behavior of adult and subadult ringed seals (*Pusa hispida*) in the Bering and Chukchi seas. Polar
Biology 42:65-80. doi:10.1007/s00300-018-2399-x.

Crawford, J.A., Quakenbush, L.T., and Citta, J.J. 2015. A comparison of ringed and bearded seal diet, condition
and productivity between historical (1975-1984) and recent (2003-2012) periods in the Alaskan Bering
and Chukchi seas. Progress in Oceanography 136:133-150. doi:10.1016/j.pocean.2015.05.011.

Deary, A., Vestfals, C., Mueter, F., Logerwell, E., Goldstein, E., Stabeno, P., Danielson, S., Hopcroft, R., and Duffy-
Anderson, J. 2021. Seasonal abundance, distribution, and growth of the early life stages of polar cod
(*Boreogadus saida*) and saffron cod (*Eleginus gracilis*) in the US Arctic. Polar Biology 1-22.

Diggle, P., and Lophaven, S. 2006. Bayesian geostatistical design. Scandinavian Journal of Statistics 33:53-64.

Diggle, P.J., Menezes, R., and Su, T.-l. 2010. Geostatistical inference under preferential sampling. Journal of
the Royal Statistical Society Series C: Applied Statistics 59:191-232.

Dou, T., Xiao, C., Liu, J., Wang, Q., Pan, S., Su, J., Yuan, X., Ding, M., Zhang, F., and Xue, K. 2021. Trends and spatial
variation in rain-on-snow events over the Arctic Ocean during the early melt season. The Cryosphere
15:883-895.

Dunn, P.K., and Smyth, G.K. 1996. Randomized quantile residuals. Journal of Computational and Graphical
Statistics 5:236-244.

Fedoseev, G.A. 1965. The ecology of the reproduction of seals on the northern part of the Sea of Okhotsk.
Izvestiya TINRO 65:212-216.

Fedoseev, G.A. 1984. Population structure, current status, and perspective for utilization of the ice-inhabiting
forms of pinnipeds in the northern part of the Pacific Ocean *In* Marine Mammals. *Edited by* A.V. Yablokov.
Nauka, Moscow, Russia. pp. 130-146.

Feltz, E.T., and Fay, F.H. 1966. Thermal requirements in vitro of epidermal cells from seals. Cryobiology
3:261-264. doi:10.1016/S0011-2240(66)80020-2.

Ferguson, S.H., Stirling, I., and McLoughlin, P. 2005. Climate change and ringed seal (*Phoca hispida*)
recruitment in western Hudson Bay. Marine Mammal Science 21:121-135.

Frost, K.J., Lowry, L.F., Gilbert, J.R., and Burns, J.J. 1988. Ringed seal monitoring: relationships of distribution
and abundance to habitat attributes and industrial activities. Final Report, Outer Continental Shelf
Environmental Assessment Program, Research Unit 667. Department of the Interior, Department of
Commerce, 101 pp. https://www.govinfo.gov/app/details/GOVPUB-I-
3b7466d9154853e3d74dea7885c39b59.

Frost, K.J., Lowry, L.F., Pendleton, G., and Nute, H.R. 2004. Factors affecting the observed densities of ringed
seals, *Phoca hispida*, in the Alaskan Beaufort Sea, 1996-99. Arctic 57:115-128. doi:10.14430/arctic489.

Hamilton, C.D., Kovacs, K.M., Ims, R.A., and Lydersen, C. 2018. Haul-out behaviour of Arctic ringed seals (*Pusa*
hispida): inter-annual patterns and impacts of current environmental change. Polar Biology 41:1063-
1082. doi:10.1007/s00300-018-2260-2.

Hammill, M.O., and Smith, T.G. 1991. The role of predation in the ecology of the ringed seal in Barrow Strait,
Northwest Territories, Canada. Marine Mammal Science 7:123-135. doi:10.1111/j.1748-
7692.1991.tb00559.x.

Harwood, L., Smith, T.G., and Melling, H. 2007. Assessing the potential effects of near shore hydrocarbon
exploration on ringed seals in the Beaufort Sea region 2003-2006. Environmental Research Studies
Funds Report No. 162. 103 pp.

Harwood, L.A., Smith, T.G., Alikamik, J., Alikamik, E., Lea, E.V., Stirling, I., Wright, H., Melling, H., and Zhu, X.
2020. Long-term, harvest-based monitoring of ringed seal body condition and reproduction in Canada's
western arctic: an update through 2019. Arctic 73:206-220. doi:10.14430/arctic70428.

Harwood, L.A., Smith, T.G., Melling, H., Alikamik, J., and Kingsley, M.C.S. 2012. Ringed seals and sea ice in
Canada's Western Arctic: Harvest-based monitoring 1992-2011. Arctic 377-390.

Hauser, D.D.W., Frost, K.J., and Burns, J.J. 2021. Ringed seal (*Pusa hispida*) breeding habitat on the landfast ice
in northwest Alaska during spring 1983 and 1984. PLoS ONE 16:e0260644.
doi:10.1371/journal.pone.0260644.

Hezel, P.J., Zhang, X., Bitz, C.M., Kelly, B.P., and Massonnet, F. 2012. Projected decline in spring snow depth on
Arctic sea ice caused by progressively later autumn open ocean freeze-up this century. Geophysical
Research Letters 39:L17505. doi:10.1029/2012gl052794.

Huntington, H.P., Danielson, S.L., Wiese, F.K., Baker, M., Boveng, P., Citta, J.J., De Robertis, A., Dickson, D.M.S.,
Farley, E., George, J.C., Iken, K., Kimmel, D.G., Kuletz, K., Ladd, C., Levine, R., Quakenbush, L., Stabeno, P.,
Stafford, K.M., Stockwell, D., and Wilson, C. 2020. Evidence suggests potential transformation of the
Pacific Arctic ecosystem is underway. Nature Climate Change 342-348. doi:10.1038/s41558-020-0695-
2.

Huntington, H.P., Nelson, M., and Quakenbush, L.T. 2015. Traditional knowledge regarding walrus, ringed
seals, and bearded seals near Barrow, Alaska. Final report to the Eskimo Walrus Commission, the Ice Seal
Committee, and the Bureau of Ocean Energy Management for contract #M13PC00015. 8 pp.

Iacozza, J., and Ferguson, S.H. 2014. Spatio-temporal variability of snow over sea ice in western Hudson Bay,
with reference to ringed seal pup survival. Polar Biology 37:817-832. doi:10.1007/s00300-014-1484-z.

Jørgensen, B. 1987. Exponential dispersion models. Journal of the Royal Statistical Society: Series B
(Methodological) 49:127-145.

Kelly, B.P., Badajos, O.H., Kunnasranta, M., and Moran, J. 2006. Timing and re-interpretation of ringed seal
surveys. Final Report. Coastal Marine Institute University of Alaska Fairbanks, Fairbanks, AK. 60 pp.

Kelly, B.P., Badajos, O.H., Kunnasranta, M., Moran, J.R., Martinez-Bakker, M., Wartzok, D., and Boveng, P.
2010a. Seasonal home ranges and fidelity to breeding sites among ringed seals. Polar Biology 33:1095-
1109. doi:10.1007/s00300-010-0796-x.

Kelly, B.P., Bengtson, J.L., Boveng, P.L., Cameron, M.F., Dahle, S.P., Jansen, J.K., Logerwell, E.A., Overland, J.E.,
Sabine, C.L., Waring, G.T., and Wilder, J.M. 2010b. Status review of the ringed seal (*Phoca hispida*). NOAA
Technical Memorandum NMFS-AFSC-212. U.S. Department of Commerce, Seattle, WA. 250 pp.
http://www.afsc.noaa.gov/publications/AFSC-TM/NOAA-TM-AFSC-212.pdf.

Kelly, B.P., and Quakenbush, L.T. 1990. Spatiotemporal use of lairs by ringed seals (*Phoca hispida*). Canadian
Journal of Zoology 68:2503-2512. doi:10.1139/z90-350.

Kelly, B.P., Quakenbush, L.T., and Rose, J.R. 1986. Ringed seal winter ecology and effects of noise disturbance.
In Outer Continental Shelf Environmental Assessment Program. Final Reports of Principal Investigators,

Volume 61. Minerals Management Service, Alaska Outer Continental Shelf Office, Anchorage, AK. pp. 447-536.

Kim, Y.-H., Min, S.-K., Gillett, N.P., Notz, D., and Malinina, E. 2023. Observationally-constrained projections of an ice-free Arctic even under a low emission scenario. Nature Communications 14:3139.

Kingsley, M.C.S., Stirling, I., and Calvert, W. 1985. The distribution and abundance of seals in the Canadian high Arctic, 1980-82. Canadian Journal of Fisheries and Aquatic Sciences 42:1189-1210.

Kovacs, K.M., Liston, G.E., Reinking, A.K., Gerland, S., and Lydersen, C. 2024. Climate warming impacts on ringed seal breeding habitat in Svalbard. Ecological Modelling 495:110790. doi:10.1016/j.ecolmodel.2024.110790.

Kristensen, K., Nielsen, A., Berg, C.W., Skaug, H., and Bell, B. 2016. TMB: automatic differentiation and Laplace approximation. Journal of Statistical Software 70. doi:10.18637/jss.v070.i05.

Laidre, K., Stern, H., Kovacs, K.M., Lowry, L., Moore, S.E., Regehr, E.V., Ferguson, S.H., Wiig, Ø., Boveng, P., Angliss, R.P., Born, E.W., Litovka, D., Quakenbush, L., Lydersen, C., Vongraven, D., and Ugarte, F. 2015. Arctic marine mammal population status, sea ice habitat loss, and conservation recommendations for the 21st century. Conservation Biology 29:724-737. doi:10.1111/cobi.12474.

Lam, H.M., Geldsetzer, T., Howell, S.E.L., and Yackel, J. 2022. Snow Depth on Sea Ice and on Land in the Canadian Arctic from Long-Term Observations. Atmosphere-Ocean 1-17.

Lamigueiro, O.P. 2012. solaR: solar radiation and photovoltaic systems with R. Journal of Statistical Software 50:1-32.

Lee, S.M., Shi, H., Sohn, B.J., Gasiewski, A.J., Meier, W.N., and Dybkjær, G. 2021. Winter Snow Depth on Arctic Sea Ice From Satellite Radiometer Measurements (2003–2020): Regional Patterns and Trends. Geophysical Research Letters 48:e2021GL094541.

Lindsay, J.M., Hauser, D.D.W., Mahoney, A.R., Laidre, K.L., Goodwin, J., Harris, C., Schaeffer, R.J., Schaeffer, R., Sr., Whiting, A.V., Boveng, P.L., Laxague, N.J.M., Betcher, S., Subramaniam, A., Witte, C.R., and Zappa, C.J. 2023. Characteristics of ringed seal *Pusa hispida* (natchiq) denning habitat in Kotzebue Sound, Alaska, during a year of limited sea ice and snow. Marine Ecology Progress Series 705:1-20. doi:10.3354/meps14252.

Lindsay, J.M., Laidre, K.L., Conn, P.B., Moreland, E.E., and Boveng, P.L. 2021. Modeling ringed seal Pusa hispida
habitat and lair emergence timing in the eastern Bering and Chukchi Seas. Endangered Species Research
46:1-17. doi:10.3354/esr01140.

London, J.M., Conn, P.B., Hardy, S.K., Richmond, E.L., Ver Hoef, J.M., Cameron, M.F., Crawford, J.A., Von Duyke,
A.L., Quakenbush, L.T., and Boveng, P.L. 2022. Spring haul-out behavior of seals in the Bering and Chukchi
seas. bioRxiv 2022-2004. doi:10.1101/2022.04.07.487572.

Lucier, C.V., and VanStone, J.W. 1991. Winter and spring fast ice seal hunting by Kangiġmiut and other Iñupiat
of Kotzebue Sound, Alaska Estudes Inuit Studies 15:30-49.

Mahoney, A.R., Eicken, H., Gaylord, A.G., and Gens, R. 2014. Landfast sea ice extent in the Chukchi and Beaufort
Seas: the annual cycle and decadal variability. Cold Regions Science and Technology 103:41-56.
doi:10.1016/j.coldregions.2014.03.003.

Markus, T., and Cavalieri, D.J. 1998. Snow depth distribution over sea ice in the Southern Ocean from satellite
passive microwave data. Antarctic sea ice: physical processes, interactions and variability 74:19-39.

Markus, T., Stroeve, J.C., and Miller, J. 2009. Recent changes in Arctic sea ice melt onset, freezeup, and melt
season length. Journal of Geophysical Research: Oceans 114. doi:10.1029/2009JC005436.

Marsh, H., and Sinclair, D.F. 1989. Correcting for visibility bias in strip transect aerial surveys of aquatic fauna.
The Journal of Wildlife Management 1017-1024.

McClintock, B.T., Moreland, E.E., London, J.M., Dahle, S.P., Brady, G.M., Richmond, E.L., Yano, K.M., and Boveng,
P.L. 2015. Quantitative assessment of species identification in aerial transect surveys for ice-associated
seals. Marine Mammal Science 21:1057-1076. doi:10.1111/mms.12206.

Melnikov, V.V. 2017. Seasonal movements and relative abundance of bearded seals (*Erignathus barbatus*) in
the coastal waters of the Chukotka Peninsula. Arctic 70:403-413. doi:10.14430/arctic4682.

Melnikov, V.V. 2022. Observations of Ringed Seal (*Phoca hispida*) in the Nearshore Waters of the Chukotka
Peninsula. Arctic 75:431-448. doi:10.14430/arctic76257.

Mesinger, F., DiMego, G., Kalnay, E., Mitchell, K., Shafran, P.C., Ebisuzaki, W., Jović, D., Woollen, J., Rogers, E.,
and Berbery, E.H. 2006. North American regional reanalysis. Bulletin of the American Meteorological
Society 87:343-360.

Miller, D.L., Burt, M.L., Rexstad, E.A., and Thomas, L. 2013. Spatial models for distance sampling data: recent
developments and future directions. Methods in Ecology and Evolution 4:1001-1010.

National Marine Fisheries Service. 2012a. Threatened status for the Arctic, Okhotsk, and Baltic subspecies of
the ringed seal and endangered status for the Ladoga subspecies of the ringed seal. Federal Register 77
FR 76706. 29 pp. https://www.federalregister.gov/documents/2012/12/28/2012-31066/endangered-
and-threatened-species-threatened-status-for-the-arctic-okhotsk-and-baltic-subspecies-of

National Marine Fisheries Service. 2012b. Threatened status for the Beringia and Okhotsk distinct population
segments of the *Erignathus barbatus nauticus* subspecies of the bearded seal. Federal Register 77 FR
76740. 29 pp. https://www.federalregister.gov/documents/2012/12/28/2012-31068/endangered-
and-threatened-species-threatened-status-for-the-beringia-and-okhotsk-distinct-population

Nelson, R.R., Burns, J.J., and Frost, K.J. 1984. The bearded seal (*Erignathus barbatus*). *In* Marine Mammal
Species Accounts, Wildlife Technical Bulletin No. 7. Vol. 7. *Edited by* J.J. Burns. Alaska Department of Fish
and Game, Juneau, AK. pp. 1-6.

Oceana and Kawerak Inc. 2014. Bering Strait Marine Life and Subsistence Data Synthesis. Oceana,
Washington, DC. 499 pp. https://oceana.org/reports/the-bering-strait-marine-life-and-subsistence-
data-synthesis/.

Olnes, J., Breed, G.A., Druckenmiller, M.L., Citta, J.J., Crawford, J.A., Von Duyke, A.L., and Quakenbush, L. 2021.
Juvenile bearded seal response to a decade of sea ice change in the Bering, Chukchi, and Beaufort seas.
Marine Ecology Progress Series 661:229-242. doi:10.3354/meps13609.

Olnes, J., Crawford, J., Citta, J.J., Druckenmiller, M.L., Von Duyke, A.L., and Quakenbush, L. 2020. Movement,
diving, and haul-out behaviors of juvenile bearded seals in the Bering, Chukchi and Beaufort seas, 2014–
2018. Polar Biology. doi:10.1007/s00300-020-02710-6.

Quakenbush, L. 2020. Biological monitoring of ice seals in Alaska to determine health and status of
populations — diet, disease, contaminants, reproduction, body condition, growth, and age at maturity.
Edited by L. Polasek. Alaska Pinniped Research Final Report to NMFS for NA16NMF4390029. Alaska
Department of Fish and Game. pp. 1-282.
https://www.adfg.alaska.gov/static/research/programs/marinemammals/pdfs/biomonitoring_biolog
y_adfg_2020_pinniped_research_ice_seals_report.pdf.

Quakenbush, L., Bryan, A., Crawford, J., and Olnes, J. 2020. Biological monitoring of ringed seals in the Bering
1023  and Chukchi seas. Alaska Department of Fish and Game, Arctic Marine Mammal Program, Final report to
1024  National Oceanographic and Atmospheric Administration for Award # NA16NMF4720079. 1-33 pp.
1025  https://adfg.alaska.gov/static/research/programs/marinemammals/pdfs/biomonitoring_biology_qua
1026  kenbush_2020_ringed_seal_report.pdf.

1027 R Core Team. 2017. R: A language and environment for statistical computing. Vienna, Austria: R Foundation
1028  for Statistical Computing. https://www.R-project.org/.

1029 Sigler, M., DeMaster, D., Boveng, P., Cameron, M., Moreland, E., Williams, K., and Towler, R. 2015. Advances in
1030  methods for marine mammal and fish stock assessments: thermal imagery and CamTrawl. Marine
1031  Technology Society Journal 49:99-106. doi:10.4031/MTSJ.49.2.10.

1032 Sigourney, D.B., Chavez-Rosales, S., Conn, P.B., Garrison, L., Josephson, E., and Palka, D. 2020. Developing and
1033  assessing a density surface model in a Bayesian hierarchical framework with a focus on uncertainty:
1034  insights from simulations and an application to fin whales (*Balaenoptera physalus*). PeerJ 8:e8226.
1035  doi:10.7717/peerj.8226.

1036 Simpkins, M.A., Hiruki-Raring, L.M., Sheffield, G., Grebmeier, J.M., and Bengtson, J.L. 2003. Habitat selection
1037  by ice-associated pinnipeds near St. Lawrence Island, Alaska in March 2001. Polar Biology 26:577-586.
1038  doi:10.1007/s00300-003-0527-7

1039 Smith, T.G. 1981. Notes on the bearded seal, *Erignathus barbatus*, in the Canadian Arctic. Canadian Technical
1040  Report of Fisheries and Aquatic Sciences No. 1042. 1042. Department of Fisheries and Oceans, Arctic
1041  Biological Station, Quebec, Canada. 49 pp.

1042 Stabeno, P.J., and Bell, S.W. 2019. Extreme conditions in the Bering Sea (2017–2018): record-breaking low
1043  sea-ice extent. Geophysical Research Letters 46:8952-8959. doi:10.1029/2019GL083816.

1044 Stroeve, J., Liston, G.E., Buzzard, S., Zhou, L., Mallett, R., Barrett, A., Tschudi, M., Tsamados, M., Itkin, P., and
1045  Stewart, J.S. 2020. A Lagrangian snow evolution system for sea ice applications (SnowModel‐LG): Part
1046  II—Analyses. Journal of Geophysical Research: Oceans 125:e2019JC015900.

1047 Stroeve, J., and Notz, D. 2018. Changing state of Arctic sea ice across all seasons. Environmental Research
1048  Letters 13:103001. doi:10.1088/1748-9326/aade56.

Ver Hoef, J.M., Cameron, M.F., Boveng, P.L., London, J.M., and Moreland, E.E. 2014. A spatial hierarchical model
for abundance of three ice-associated seal species in the eastern Bering Sea. Statistical Methodology
17:46-66. doi:10.1016/j.stamet.2013.03.001.

Ver Hoef, J.M., London, J.M., and Boveng, P.L. 2010. Fast computing of some generalized linear mixed pseudo-
models with temporal autocorrelation. Computational Statistics 25:39-55. doi:10.1007/s00180-009-
0160-1.

Vestfals, C.D., Mueter, F.J., Duffy-Anderson, J.T., Busby, M.S., and De Robertis, A. 2019. Spatio-temporal
distribution of polar cod (*Boreogadus saida*) and saffron cod (*Eleginus gracilis*) early life stages in the
Pacific Arctic. Polar Biology 42:969-990. doi:10.1007/s00300-019-02494-4.

Von Duyke, A.L., Douglas, D.C., Herreman, J.K., and Crawford, J.A. 2020. Ringed seal (*Pusa hispida*) seasonal
movements, diving, and haul-out behavior in the Beaufort, Chukchi, and Bering Seas (2011–2017).
Ecology and Evolution 10:5595-5616. doi:10.1002/ece3.6302.

Webster, M., DuVivier, A., Holland, M., and Bailey, D. 2021. Snow on Arctic sea ice in a warming climate as
simulated in CESM. Journal of Geophysical Research: Oceans 126:e2020JC016308.

Webster, M., Gerland, S., Holland, M., Hunke, E., Kwok, R., Lecomte, O., Massom, R., Perovich, D., and Sturm, M.
2018. Snow in the changing sea-ice systems. Nature Climate Change 8:946-953.

Webster, M.A., Rigor, I.G., Nghiem, S.V., Kurtz, N.T., Farrell, S.L., Perovich, D.K., and Sturm, M. 2014.
Interdecadal changes in snow depth on Arctic sea ice. Journal of Geophysical Research: Oceans 119:5395-
5406. doi:10.1002/2014JC009985.

Wolotira, R.J., Jr. 1985. Saffron cod (*Eleginus gracilis*) in western Alaska: the resource and its potential. NOAA
Technical Memorandum NMFS F/NWC-79. https://repository.library.noaa.gov/view/noaa/5647.

Wood, S.N. 2017. Generalized additive models: An introduction with R. 2 ed. Chapman & Hall/CRC press, New
York. 496 pp. doi:10.1201/9781315370279.

Young, N.C., Brower, A.A., Muto, M.M., Freed, J.C., Angliss, R.P., Friday, N.A., Boveng, P.L., Brost, B.M., Cameron,
M.F., Crance, J.L., Dahle, S.P., Fadely, B.S., Ferguson, M.C., Goetz, K.T., London, J.M., Oleson, E.M., Ream, R.R.,
Richmond, E.L., Shelden, K.E.W., Sweeney, K.L., Towell, R.G., Wade, P.R., Waite, J.M., and Zerbini, A.N. 2023.
Alaska marine mammal stock assessments, 2022. National Marine Fisheries Service. 316 pp.

https://www.fisheries.noaa.gov/national/marine-mammal-protection/marine-mammal-stock-

assessment-reports-region.

1078