# Focused Ultrasound Augments the Delivery and Penetration of Model Therapeutics into Cerebral Cavernous Malformations

**Delaney G. Fisher[1,#], Matthew R. Hoch[1,#], Catherine M. Gorick[1], Claire Huchthausen[2],**

**Victoria R. Breza[1], Khadijeh A. Sharifi[3,4], Petr Tvrdik[3,4], G. Wilson Miller[1,5],**

**and Richard J. Price[1,5]**

# Authors contributed equally

1. Department of Biomedical Engineering, University of Virginia, Charlottesville, VA

2. Department of Physics, University of Virginia, Charlottesville, VA

3. Department of Neuroscience, University of Virginia, Charlottesville, VA

4. Department of Neurosurgery, University of Virginia Health System, Charlottesville, VA

5. Department of Radiology & Medical Imaging, University of Virginia, Charlottesville, VA

Corresponding Author:

Richard J. Price, Ph.D.

415 Lane Road, MR-5, Room 2316

Charlottesville, VA 22908.

Phone: 434-924-0020

Email: rprice@virginia.edu

**Abstract**

**BACKGROUND:** Cerebral cavernous malformations (CCMs) are vascular neoplasms in the brain that can cause debilitating symptoms. Current treatments pose significant risks to some patients, motivating the development of new nonsurgical options. We recently discovered that focused ultrasound-microbubble treatment (FUS) arrests CCM formation and growth. Here, we build on this discovery and assess the ability of FUS to deliver model therapeutics into CCMs.

**METHODS:** Quantitative T1 mapping MRI sequences were used with 1 kDa (MultiHance; MH) and 17 kDa (GadoSpin D; GDS) contrast agents to assess the FUS-mediated delivery and penetration of model small molecule drugs and biologics, respectively, into CCMs of Krit1 mutant mice.

**RESULTS:** FUS elevated the rate of MH delivery to both the lesion core (4.6-fold) and perilesional space (6.7-fold). Total MH delivery more than doubled in the lesion core and tripled in the perilesional space when FUS was applied immediately prior to MH injection. For the model biologic drug (i.e. GDS), FUS was of greater relative benefit, resulting in 21.7-fold and 3.8-fold delivery increases to the intralesional and perilesional spaces, respectively

**CONCLUSIONS:** FUS augments the delivery and penetration of therapeutics into the complex and disorganized CCM microenvironment. Benefits to small molecule drug delivery are more evident in the perilesional space, while benefits to biologic delivery are more evident in CCM cores. These findings, when combined with ability of FUS alone to control CCMs, highlight the potential of FUS to serve as a powerful non-invasive therapeutic platform for CCM.

**Introduction**

Cerebral cavernous malformation (CCM) is a vascular disorder characterized by the development of abnormal, dilated clusters of blood vessels in the brain[1]. These malformations are prone to repetitive hemorrhages, inducing debilitating symptoms, such as neurological deficits, seizures, and stroke, in affected individuals[2–4]. Presently, the prevailing recourse for treating symptomatic CCMs is surgical resection. However, surgical excision of CCMs poses an elevated risk of complications and morbidity[5,6].

Despite multiple studies investigating therapeutic targets and screening pharmacological treatments for CCM[7–18], no approved drug treatments exist for CCM. The majority of tested pharmacological agents for CCM are small molecules. In comparison, larger biologic molecules, such as antibodies and gene therapies, have not been as thoroughly explored. Additionally, drugs showing promise in acute CCM models often demonstrate limited efficacy in more clinically-representative chronic models, suggesting a potential need for greater local doses of these therapies[19,20]. Indeed, though CCMs are known to be more permeable than healthy cerebrovasculature[21–24], delivery of systemically administered drugs to these complex lesions is poorly understood.

Focused ultrasound-mediated blood-brain barrier opening (FUS) has emerged as a promising non-invasive drug delivery technology[25–27]. With FUS, acoustic energy is concentrated into a confined volume, facilitating the oscillation of intravenously administered gas-filled microbubbles within blood vessels of the targeted region. These microbubble oscillations induce a transient disruption of endothelial tight junctions[28] and increased active transport[29], enabling therapeutic delivery across the blood-brain barrier (BBB). Magnetic resonance imaging (MRI) guidance permits spatial targeting of FUS to specific brain regions and BBB opening confirmation through the accumulation of gadolinium-based MRI contrast agents.

Recently, our group demonstrated that FUS, in the absence of therapeutic delivery, arrests the formation and growth of CCMs[30]. This remarkable observation prompts the exploration of the

combined impact of FUS-mediated lesion stabilization and therapeutic delivery on CCMs. While

our previous study also confirmed that FUS enhanced MRI contrast agent delivery beyond the

natural permeability of CCMs, the MRI sequences only provided qualitative assessments. In

particular, this qualitative MRI approach was sub-optimal for visualizing contrast agent delivery to

the lesion core. Indeed, the cellular and molecular composition within the lesion core, including

mutated endothelium, red blood cells, and their byproducts, differs substantially from the

perilesional space, characterized by dense populations of astrocytes and microglia[30,31]. This

difference not only affects MRI signal but may also have important implications for drug delivery

to these distinct regions. Consequently, to facilitate comprehensive measurements of potential

enhanced therapeutic delivery with FUS in the intricate CCM microenvironment, quantitative MRI

methods are needed.

Building on our recent observations[30], the objective of this study was to establish a

foundation for therapeutic delivery approaches that harness and synergize with this potent

bioeffect. We have previously demonstrated that T1-contrast mapping can enable longitudinal,

quantitative concentration measurements of gadolinium-based molecules in CCMs[31]. Thus, this

is an ideal method to measure FUS-induced changes for therapeutic delivery to CCMs. To this

end, we employed T1-contrast mapping MRI to quantitatively evaluate the delivery of 1 kDa and

17 kDa molecules to CCMs, comparing outcomes with and without FUS. This study lays the

groundwork for treatment regimens of FUS-delivered molecules capable of inducing CCM

regression and clearance.

## Results

### FUS Enhances Delivery Rate of MultiHance in CCMs

We first tested if FUS would increase the delivery rate of a model small molecule drug to the CCM microenvironment. To this end, we employed T1 mapping MRI to measure the concentration of the MRI contrast agent MultiHance (MH; gadobenate dimeglumine; ~1 nm; ~1 kDa) before and after the application of FUS in CCM mice. One frontal hemisphere received FUS

**Figure 1. FUS Enhances Delivery Rate of MultiHance to CCMs. A, B)** PNP histories (**A**) and integrated acoustic emissions (**B**) for FUS treatments (n=6). f = fundamental frequency; BB = broadband. **C)** T1 mapping MRIs illustrating MH accumulation, in and around 3 CCMs, over a 40 min time period. Perilesional and intralesional regions are denoted. FUS was applied to 2 of the 3 CCMs at the 20 min timepoint. A marked increase in MH concentration is evident in and around FUS-treated CCMs at 40 min. **D**) Temporal fold change in intralesional MH concentration over the average initial concentration for FUS⁻ and FUS⁺ CCMs. FUS⁺ CCMS were treated at 20 min after MH injection (blue shading). **E)** Slope ratios (Post-FUS/Pre-FUS) derived from intralesional MH concentration plots. *p=0.0221; Mann-Whitney test. **F)** Temporal fold change in perilesional MH concentration over the average initial concentration for FUS⁻ and FUS⁺ CCMs. **G)** Slope ratios (Post-FUS/Pre-FUS) derived from perilesional MH concentration plots. ****p<0.0001; Mann-Whitney test.

(n=6 sonication targets) with passive cavitation detection (PCD) feedback control 20 minutes after

intravenous (i.v.) MH injection. During FUS, peak-negative pressures (PNPs) settled into a range

of 0.3 to 0.4 MPa (**Figure 1A**), yielding integrated acoustic emissions shown in **Figure 1B**. The

contralateral hemisphere was not sonicated (i.e., FUS⁻ control) to illustrate baseline CCM

permeability. As expected, prior to FUS, CCMs in the non-sonicated and sonicated hemispheres

displayed similar rates of MH accumulation (**Figure 1C, D, F**). After FUS, the rate of MH

accumulation in the lesion core was enhanced (**Figure 1D**), increasing to well-above (4.6-fold)

the rate of MH accumulation in FUS⁻ CCMs (p=0.0221; **Figure 1E**). Predictably, the perilesional

space of these CCMs also displayed the same permeability rate prior to FUS in both groups

(**Figure 1F**). FUS then increased perilesional MH delivery rate by 6.7-fold over the rate of MH

accumulation in FUS⁻ CCMs (p<0.0001; **Figure 1G**). These results indicate that FUS enhances

the delivery rate of a model small molecule drug to both the lesion core and the surrounding CCM

microenvironment.

**FUS Enhances Total Delivery of MultiHance in CCMs**

We then tested the ability of FUS to augment model small molecule drug delivery to CCMs

when applied concurrently with model drug injection, reflecting the clinical staging to maximize

overall delivery. On day 1, T1 mapping MRI was conducted on CCM mice following i.v. MH

injection to measure baseline permeability (**Figure 2A**). On day 2, FUS was applied to one frontal

hemisphere of the same CCM mice immediately before i.v. MH injection. T1 mapping MRI was

conducted for 20 mins thereafter (**Figure 2B**). FUS markedly boosted the intralesional MH

delivery rate, as well as mean intralesional MH concentration (p=0.0070; **Figure 2C**), with a 2.5-

fold enhancement evident at 20 minutes. Area under the curve (AUC) analysis, representing the

integrated exposure of CCM tissue to the model drug through time, indicates that FUS enhances

intralesional model drug exposure by 1.9-fold (p=0.0122; **Figure 2D**). Regarding the perilesional

**Figure 2. FUS Enhances Total Delivery of MultiHance to CCMs. A)** T1 mapping MRI illustrating baseline permeability to MH, in and around 2 CCMs, 20 min after MH injection on day 1. **B)** T1 mapping MRI illustrating enhanced MH accumulation, in and around 2 CCMs, 20 min after MH injection and FUS treatment on day 2. **C**) Intralesion MH concentration as a function of time after MH injection. Baseline permeability to MH (FUS⁻) was measured on day 1, with FUS (FUS⁺) measurements made on paired CCMs on day 2. **p=0.0070; Repeated measures two-way ANOVA with Geisser-Greenhouse correction. **D**) Area under the curve (AUC) metric derived from intralesion concentration data in C. *p=0.0122; Wilcoxon matched-pairs signed rank test. **E**) Perilesion MH concentration as a function of time after MH injection. ***p=0.0005; Repeated measures two-way ANOVA with Geisser-Greenhouse correction. **F**) Area under the curve (AUC) metric derived from perilesional concentration data in E. ***p=0.0007; Wilcoxon matched-pairs signed rank test.

space, MH concentration was also markedly elevated with FUS (p=0.0005; **Figure 2E**), with a

3.1-fold enhancement evident at 20 minutes. AUC yielded a 2.9-fold increase in model drug

exposure over FUS⁻ CCMs (p=0.0007; **Figure 2F**). Notably, MH delivery after FUS becomes

evident in the perilesional space (**Figure 2E**) before the intralesional space (**Figure 2C**) (0.040

mM versus 0.029 mM, respectively, after 5 minutes), yet both locations plateau to the same mean

concentration by 20 mins post-injection (0.069 mM each). These results reveal that FUS can more

than double the amount of a small molecule delivered to the lesion core and triple the amount in

the surrounding CCM microenvironment.

### FUS Enhances Total Delivery of GadoSpin D in CCMs

Next, we tested the potential for FUS to enhance the total delivery and penetration of a

biologic, which are typically >1 kDa, to CCMs. To this end, we employed the MRI contrast agent

GadoSpin D (GDS; dendritic Gd-chelate; ~5 nm; ~17 kDa) as a model biologic. As in the MH

experiments (**Figure 2**), baseline permeability of CCMs to GDS was measured on day 1 (**Figure**

**3A**). On day 2, FUS was applied to paired CCMs from day 1. FUS improved total GDS delivery in

both the intralesional and perilesional spaces compared to baseline CCM permeability (**Figure**

**3B**). FUS elicited a striking increase in GDS delivery to the lesion core (p=0.0106; **Figure 3C**),

reaching 21.7-fold at 20 minutes. AUC was increased 4.8-fold in CCM cores with FUS (p=0.0078;

**Figure 3D**). Meanwhile, perilesional delivery of GDS was also enhanced with FUS (p= 0.0021;

**Figure 3. FUS Enhances Total Delivery of Gadospin D in CCMs. A)** T1 mapping MRI illustrating baseline permeability to GDS, in and around a CCM, 20 min after GDS injection on day 1. **B)** T1 mapping MRI illustrating enhanced GDS accumulation, in and around a CCM, 20 min after GDS injection and FUS treatment on day 2. **C)** Intralesion GDS concentration as a function of time after GDS injection. Baseline permeability to GDS (FUS⁻) was measured on day 1, with FUS (FUS⁺) measurements made on paired CCMs on day 2. *p=0.0106; Repeated measures two-way ANOVA with Geisser-Greenhouse correction. **D)** Area under the curve (AUC) metric derived from intralesion concentration data in C. **p=0.0078; Wilcoxon matched-pairs signed rank test. **E)** Perilesion GDS concentration as a function of time after GD injection. **p=0.0021; Repeated measures two-way ANOVA with Geisser-Greenhouse correction. **F)** Area under the curve (AUC) metric derived from perilesional concentration data in E. *p=0.0195; Wilcoxon matched-pairs signed rank test.

**Figure 3E**), reaching a 3.8-fold increase at 20 minutes. For GDS in the perilesional space,

integrated tissue-drug exposure increased 2.2-fold (p=0.0195; **Figure 3F**). The lesion core and

perilesional space followed a similar temporal pattern of GDS enhancement following FUS, but

the intralesional space peaked at a higher concentration than the perilesional space (0.010 mM

versus 0.0076 mM, respectively).

**Comparison of FUS-Mediated MultiHance and GadoSpin D Delivery to Intralesion and**

**Perilesion CCM Compartments**

We also investigated whether FUS differentially affects the delivery of MH and GDS to

intralesional and perilesional regions of CCMs. To this end, we first needed to verify that the

**Figure 4. Focused Ultrasound Application in Multihance and Gadospin D Delivery Experiments was Comparable. A, B)** PNP histories during BBB opening by acoustic emissions feedback control for MH (n=5) (A) and GDS (n=4) (B) treatments. **C, D)** Average (C) and maximum (D) PNPs for MH and GDS delivery experiments. Mann-Whitney tests. **E)** Integrated acoustic emissions from key spectral domains for MH and GDS delivery experiments. f = fundamental frequency; BB = broadband. Mann-Whitney tests.

153 applied FUS PNP, as well as the resultant MB activity, were equivalent in the MH and GDS

154 experiments. The PNP histories for the MH (**Figure 4A**) and GDS (**Figure 4B**) experiments

155 followed similar trajectories, and there were no differences in average (**Figure 4C**) and maximum

**Figure 5. Comparison of FUS-Mediated MultiHance and GadoSpin D Delivery to Intralesion and Perilesion CCM Compartments. A, B**) Intralesion (A) and perilesion (B) AUC ratios (FUS$^+$/FUS$^-$) for MH and GDS. Mann-Whitney tests. **C)** Intralesion/perilesion ratios of AUC ratios for MH and GDS. *p=0.045; Mann-Whitney test. **D, E)** Intralesion (D) and peilesion (E) maximum concentration ratios (FUS$^+$/FUS$^-$) for MH and GDS. Mann-Whitney tests. **F)** Intralesion/perilesion ratios of maximum concentration ratios for MH and GDS. *p=0.025; Mann-Whitney test. **G, H)** Intralesion (G) and perilesion (H) post-FUS time to maximum concentration for MH and GDS. ***p<0.001; Mann-Whitney tests.

(**Figure 4D**) applied PNP. Moreover, MB activity, as assessed by acoustic emissions across

several key spectral domains (i.e. sub-harmonic, harmonic, ultra-harmonic, and broadband), was

equivalent for the MH and GDS experiments. Thus, any differences between MH and GDS

delivery were not due to differences in FUS application and/or MB response.

When comparing GDS to MH delivery using the AUC metric, similar levels of FUS-

mediated delivery enhancement (i.e. $FUS^+/FUS^-$) to both the intralesional (**Figure 5A**) and

perilesional spaces (**Figure 5B**) were observed, with GDS exhibiting a slight trend (p=0.23) over

MH in intralesional AUC augmentation (**Figure 5A**). To then examine whether intralesional or

perilesional AUC augmentation might be favored for one or both of the contrast agents, we

calculated the ratio of intralesional FUS-mediated AUC enhancement over perilesional FUS-

mediated AUC enhancement. Resultant values >1 suggest greater relative intralesional

amplification (**Figure 5C**). By this metric, GDS exhibited greater relative FUS-mediated

augmentation of delivery to the intralesional space when compared to MH (**Figure 5C**). We then

repeated this analysis using maximum concentration as the key metric. As with the AUC

comparisons, there was no difference between the 2 contrast agents with respect to FUS-

mediated intralesional (**Figure 5D**) and perilesional (**Figure 5E**) delivery augmentation, but there

was greater relative amplification of delivery to the intralesional space for GDS (**Figure 5F**).

Finally, we compared post-FUS times to maximum concentration in the intralesional and

perilesional spaces for MH and GDS (**Figure 5G and 5H**). For both regions, GDS reached its

maximum concentration in about 10 min after FUS, while MH concentration was typically still

increasing at the final (20 min) timepoint.

**Discussion**

We previously elucidated that FUS can arrest CCM growth and formation, even in the

absence of therapeutic delivery[30]. Here, we aimed to advance the synergistic potential for

concurrent therapeutic delivery with this approach. Utilizing longitudinal T1 mapping MRI, we

quantified the impact of FUS on therapeutic delivery of model small molecule drugs and biologics

to CCMs. Our findings revealed a significant enhancement in the delivery rate of a 1 kDa small

molecule, exhibiting a 4.6-fold increase in the lesion core and a 6.7-fold increase in the

perilesional space. Moreover, FUS augmented overall delivery of both the 1 kDa small molecule

and a 17 kDa model biologic to CCMs, with a 2.5-fold increase for the model small molecule drug

and an impressive 22-fold increase for the model biologic in the lesion core. In the perilesional

space, there was a 3.1-fold increase for the model small molecule drug and a 3.8-fold increase

for the model biologic. GDS reached its post-FUS maximum concentration sooner than MH,

suggesting there may be a more transient delivery window for biologics. Finally, our analysis

uncovered a nuanced aspect of FUS enhancement, wherein the relative FUS-mediated effect is

more pronounced for the small molecule in the perilesional space and for the model biologic in

the lesion core. These results collectively establish a robust foundation for employing FUS in

targeted therapeutic delivery regimens to effectively mitigate CCMs.

**T1 Mapping MRI Enables Spatiotemporal, Intra-CCM, Delivery Comparisons**

Given the notable heterogeneity in baseline CCM permeability[23,31,32], methods allowing for

comparative measurements in the same CCMs over time are important for generating statistical

power and robust conclusions. We have previously shown that T1 mapping MRI enables

longitudinal and quantitative assessments of contrast agent deposition in individual CCMs[31].

Thus, it was reasonable to leverage this MRI approach to measure model drug delivery to CCMs

with FUS. Yet another advantage of T1 mapping MRI is that it has sufficient spatial resolution to

discern differences in discrete CCM tissue compartments. Indeed, the lesion core harbors

mutated, cavernous vessels filled with clotted blood components, while the perilesional space

surrounds the core with dense populations of astrocytes, microglia, and macrophages[30,31,33].

These regional differences in the CCM microenvironment pose varying biotransport challenges

that can influence the efficacy of different delivery approaches and molecule sizes. T1 mapping

MRI enabled us to measure the exact concentration of MH and GDS in both the intralesional and

perilesional spaces of the CCM microenvironment, both with and without FUS.

**Differential Spatial Delivery Augmentation for Varying-Sized Molecules with FUS**

One unexpected and potentially important finding that arose from our spatiotemporally

detailed T1 mapping results was that FUS differentially augments the delivery of small and large

molecules to the two pre-defined CCM tissue compartments (i.e. lesion core vs. perilesional

space). Specifically, FUS provided a greater relative benefit for (i) model small molecule drug

delivery to the perilesional space and (ii) model biologic delivery to the lesion core. This effect is

evident when using either AUC (**Figure 5C**) or maximum concentration (**Figure 5F**) as the metric

of interest. To explore the potential causes behind the observed differential spatial delivery of

varying-sized molecules with FUS, we first emphasize that FUS is known to offer varying degrees

of benefit based on the transport properties of a given molecule[34]. Noting that the increase in

permeability induced by FUS had a greater effect for MH in the perilesional space, we postulate

that the benefit of FUS for small molecule drug delivery in regions with an already disrupted BBB

(e.g. the lesion core) is less than in areas that have a more intact BBB (e.g. the perilesional

space). Conversely, for a larger molecule like GDS (17 kDa; 5 nm), crossing the disrupted BBB

in the lesion core may be less feasible due to biophysical constraints limiting the transport of a

larger molecule. FUS partially alleviates these constraints, ultimately providing more relative

benefit for larger molecules than for small molecules in the already leaky CCM core. In perilesional

regions harboring a more intact BBB, even small molecules cannot effectively cross into the brain

parenchyma. Thus, FUS yields a larger benefit for small molecule delivery in this region.

Moreover, for larger molecules, the advantage of FUS may be less pronounced in regions with a

previously intact BBB than in regions with a previously disrupted BBB, once again due to

increased biophysical transport constraints.

We also note that differences in BBB closure time, as well as clearance mechanisms within

the CCM microenvironment, for small molecules and biologics could impact the integrated

exposure of tissue to drug. Here, GDS reached its maximum concentration at ~10 minutes after

FUS (**Figure 5G and 5H**), while MH concentration was often still increasing at 20 min after FUS.

This is consistent with the hypothesis that the BBB in and around CCMs closes fairly rapidly to

larger therapeutics, which could factor into how injections are timed with respect to FUS

application. Regarding clearance, while there is evidence that FUS alters clearance mechanisms

through modification of the glymphatic system[35–37] and BBB efflux pumps[38,39], its specific influence

on the clearance of varying-sized molecules remains unclear. Our data indicate that GDS

concentrations rapidly decrease without FUS when compared to MH without FUS or GDS with

FUS, highlighting that differential clearance is also likely a significant determinant of tissue-drug

exposure.

**Potential for Clinical Impact on Therapeutic Delivery in CCM**

Here, we demonstrate that FUS enhances therapeutic delivery for molecules of different

sizes in both the CCM core and surrounding perilesional space. In the clinic, this will translate to

increased local delivery for any given standard systemic dose, thereby increasing therapeutic

index. Furthermore, enhanced on-target drug delivery reduces the risk of side effects associated

with off-target delivery. The greater benefit observed for larger molecules with FUS opens the

door for biologic delivery exploration for CCM. Indeed, our study highlights that, in the absence of

FUS, the delivery of a 5 nm model biologic drug (GDS) is minimal. There also may be rapid

clearance from both the intralesional and perilesional spaces. However, with FUS, biologic-sized

molecules are more effectively retained in both CCM compartments. These findings pave the way

for future investigation into even larger agents with promising therapeutic potential for CCM, such

as antibodies and gene therapy vectors.

Notably, FUS also offers a level of precision that can be customized for either familial or

sporadic cases of CCM. In these studies, we induce BBB opening in a substantial volume—almost

one-quarter—of the CCM brain. In contrast, our previous study showcased targeting FUS to a

smaller volume of the CCM brain[30]. For patients, FUS can be tailored to target a large volume,

which may be necessary for familial patients with multiple CCMs, or it can be focused on a singular

CCM, as would be needed for sporadic cases. Moreover, the region of delivery can also be

adapted for the mechanism of action of the delivered therapeutic. Drugs with a preventative effect

could be more widely delivered than those with specific corrective functions in the CCM

microenvironment.  Ultimately, given its ability to stabilize lesions and seamlessly integrate with

therapeutic delivery, FUS may offer a powerful platform for the treatment of CCM via image-

guided drug and gene delivery.

**Materials and Methods**

**Animals**

All animal experiments adhered to ethical guidelines and were approved by the University

of Virginia Animal Care and Use Committee. The animals were housed in accordance with

standard laboratory conditions, maintaining a temperature of 22°C and a 12-hour light/12-hour

dark cycle. The generation of the CCM murine model was established as previously detailed[31].

Briefly, *Krit1*$^{fl/null}$ or *Krit1*$^{fl/fl}$ male or females were generated under the endothelial promoter

Pdgfb$^{CreER}$. On postnatal day 5, induction of *Krit1* was initiated with a subcutaneous injection of

tamoxifen (50 µL at 2mg/mL in corn oil). Genotypes were subsequently verified using Transnetyx

(Cordova, TN). Mice were studied between 2 and 3 months old.

**MRI Acquisition**

Data for T1 maps were acquired with a set of multi-slice 2D spin echo (SE) images at

varied repetition times (TR) to generate a saturation recovery curve. 2 sets of 7 images, for a total

of 14 scans, were acquired prior to FUS and contract agent administration to obtain saturation

recovery curves with a satisfactory dynamic range. The two sets of image series were offset by

the slice thickness in the slice select plane to ensure 3D coverage of the brain. The parameters

for these scans were: TR=790, 1040, 1350, 1750, 2300, 3215, and 7000 ms, TE=6.71 ms, slice

thickness=0.6 mm, slice gap=0.6 mm, FOV=35 x 35 mm, matrix size=180 x 180, rare factor=10,

and R= 0.194 x 0.194 x 0.6 mm$^3$. After FUS and contrast agent administration, 14 SE images

were acquired with identical parameters except at a fixed TR=1040 ms. The acquisitions

alternated between slice package orientations resulting in 7 images at each slice profile geometry.

Time per acquisition was 1 minute and 28 seconds.

**Data Processing**

A saturation recovery approach was utilized to calculate $M_0$ and all T1 values (pre and

post contrast) on a voxel-by-voxel basis by fitting the data to the signal equation:

298    
$$|S| = M_0 \left(1 - e^{\frac{-TR}{T1}}\right) e^{\frac{-TE}{T2}} \qquad \text{Eqn [1]}$$

300    In equation 1, $|S|$ is the magnitude of the signal within the voxel, $M_0$ is the product of the thermal

301    equilibrium magnetization and coil sensitivity, TR is the repetition time (ms), T1 is the spin-lattice

302    relaxation (ms), TE is the echo time (ms), and T2 is the spin-spin relaxation (ms). The echo time

303    exponential is assumed to be 1 due to TE<<T2, resulting in the final form seen in equation 2.

305    
$$|S| = M_0 \left(1 - e^{\frac{-TR}{T1}}\right) \qquad \text{Eqn [2]}$$

A custom MATLAB script fit the signal magnitude data on a voxel-by-voxel basis to equation 2.

Each fitting procedure simultaneously fit the data to 8 functions: function 1 incorporated the 7 pre-

contrast variable TR scans, while functions 2-8 incorporated the singular scan at a fixed TR but

different time points. The fits were constrained to having the same $M_0$ value but allowed different

T1 values. Pre-contrast and post-contrast T1 values were then used to calculate the contrast

agent concentration on a voxel-by-voxel basis at each time point using equation 3.

314    
$$\frac{1}{T_{1\_Post}} = \frac{1}{T_{1\_Pre}} + r_1 C_1 \quad \text{Eqn [3]}$$

In equation 3, $T_{1\_Post}$ is the post-contrast value at a particular time point (ms), $T_{1\_Pre}$ is the pre-

contrast T1 value (ms), $r_1$ is the contrast agent relaxivity (L/mmol/ms), and C1 is the contrast

agent concentration (mM). At the conclusion of this process, concentration values for slice

package 1 existed for time points (minutes): 1.47, 4.40, 7.33, 10.27, 13.2, 16.13, and 19.07, while

concentration values for slice package 2 existed for time points (minutes): 2.93, 5.87, 8.80, 11.73,

14.67, 17.60, and 20.53. To obtain 3D coverage at each time point, concentration data was

calculated at the missing time points by linearly interpolating between the acquired points. This

required an assumption of 0 concentration at minute 0 for slice package 2. The 20.53-minute time

point was not used because it required data be extrapolated past minute 19.07 for slice package

1.

A second custom MATLAB script was used to calculate average concentrations with

manually drawn regions of interest (ROIs) on the concentration maps. To ensure the iron rich

intralesional data was not skewed by susceptibility artifacts, a data exclusion method was

developed. Briefly, a ROI of healthy brain tissue on the contralateral hemisphere was used to

calculate an average residuals value for the fit. If any residuals value for the voxels within the

lesion core were 3 times greater than this average, they were excluded from the analysis. The

value of 3 was empirically determined. To maintain consistency within data processing, this was

also applied to all perilesional data.

**FUS Blood-Brain Barrier Opening**

The FUS procedure was conducted using the RK-300 small bore FUS device (FUS Instruments, Toronto, CA). Mice were prepared by shaving and depilating their heads before being placed in a supine position and coupled to the transducer using degassed ultrasound gel. Blood-brain barrier opening was achieved using a 1.1 MHz single-element transducer with a 10 ms burst length over a 2000 ms period. A total of 60 sonications were administered during a 2-minute sonication duration. The FUS Instruments software, operating in the "Blood-brain Barrier" mode, facilitated PCD-modulated PNP. The feedback control system parameters were set as follows: a starting pressure of 0.2 MPa, pressure increment of 0.05 MPa, maximum pressure of 0.4 MPa, 20 sonication baselines without microbubbles, area under the curve (AUC) bandwidth of 500 Hz, AUC threshold of 10 standard deviations, pressure drop of 0.95, and frequency selection of the subharmonic, first ultraharmonic, and second ultraharmonic. Optison™ (GE HealthCare) microbubbles were intravenously injected as a bolus dose of 10^5 microbubbles per gram of body weight. Prior to sonication, the distribution of microbubble diameter and concentration was assessed using a Coulter counter (Multisizer 3; Beckman Coulter, Fullerton, California). T1 mapping MRI sequences were used to guided sonication targeting. Six non-overlapping sonication targets were placed over one frontal hemisphere with placement optimized to target CCMs.

**Contrast Agent Injections**

MultiHance ® (gadobenate dimeglumine; Bracco) and GadoSpin D™ (dendritic Gd-chelate; Viscover) were injected as a bolus intravenously at a dose of 0.01 and 0.0002 mmol, respectively, diluted in saline. Injection of contrast agent was given immediately prior to MRI acquisition for FUS⁻ control studies and immediately following the initiation of FUS for FUS⁺ studies.

**Passive Cavitation Detection**

Acoustic emissions during FUS were detected with a fiber-optic hydrophone (Precision Acoustics, Dorset, UK) of 10 um diameter and 15 mm aperture center-mounted within the ultrasound transducer. Emissions data was processed with a custom MATLAB script. The area under the curve of the acoustic emissions at the subharmonic (0.5f) and ultra-harmonics (1.5f, 2.5f) after applying a 300 Hz bandwidth filter. Broadband emissions were evaluated by summing acoustic emissions following the removal of all emissions at the fundamental frequency, harmonics (2f, 3f, 4f), subharmonic (0.5f), and ultra-harmonics (1.5f, 2.5f, 3.5f).

**Statistical Analysis**

All results reported with error bars are means with standard deviation. The "n" values per group are made evident either by individual data points shown or statement of "n" value in figure or figure legend. Statistical significance was assessed at $p < 0.05$ for all experiments and were calculated using GraphPad Prism 9 (San Diego, USA). Statistical tests are provided in the figure legends.

**Author Contributions**

DGF, MH, and RJP conceptualized the study. DGF and MH conducted the FUS experiments with the aid of CMG in animal preparation. MRI sequences and analysis were optimized by MH and GWM. MRI data was acquired and processed by MH and analyzed by DGF and MH. KAS generated experimental animals. DGF and MH designed the figures and wrote the manuscript. GWM, PT, and RJP edited the manuscript. All authors approved the manuscript.

**Acknowledgements**

This work was supported by funding from NIH R01CA279134, R01EB030409, R01EB030744, and R21NS118278 to RJP; NIH R21NS116431 and grants from Focused Ultrasound Foundation,

Be Brave for Life Foundation, and Alliance to Cure Cavernous Malformation to PT; NIH

R01CA226899 to GWM; and American Heart Association 830909 to DGF. MRI was performed in

the University of Virginia Molecular Imaging Core Laboratory, with support for the 9.4T Bruker

scanner from NIH S10OD025024. We thank Dr. Kevin Whitehead for kindly providing the mouse

strains used in this study. We are also grateful to Jeremy Gatesman of the University of Virginia

Center for Comparative Medicine for assistance with catheterization procedures.

**References**

1.  Snellings, D. A. *et al.* Cerebral Cavernous Malformation: From Mechanism to Therapy.

*Circ. Res.* **129**, 195–215 (2021).

2.  Denier, C. *et al.* Clinical features of cerebral cavernous malformations patients with

KRIT1 mutations. *Ann. Neurol.* **55**, 213–220 (2004).

3.  Gault, J., Sain, S., Hu, L. J. & Awad, I. A. Spectrum of genotype and clinical

manifestations in cerebral cavernous malformations. *Neurosurgery* **59**, 1278–1284

(2006).

4.  Gianfrancesco, F. *et al.* Highly variable penetrance in subjects affected with cavernous

cerebral angiomas (CCM) carrying novel CCM1 and CCM2 mutations. *Am. J. Med.*

*Genet. B. Neuropsychiatr. Genet.* **144B**, 691–695 (2007).

5.  Awad, I. A. & Polster, S. P. Cavernous angiomas: deconstructing a neurosurgical

disease: JNSPG 75th Anniversary Invited Review Article. *J. Neurosurg.* **131**, 1–13

(2019).

6.  Amin-Hanjani, S., Ogilvy, C. S., Ojemann, R. G. & Crowell, R. M. Risks of surgical

management for cavernous malformations of the nervous system. *Neurosurgery* **42**,

1220–1228 (1998).

7.  Hong, T. *et al.* Somatic MAP3K3 and PIK3CA mutations in sporadic cerebral and spinal

cord cavernous malformations. *Brain* **144**, 2648–2658 (2021).

8.  Zhou, Z. *et al.* Cerebral cavernous malformations arise from endothelial gain of MEKK3-

KLF2/4 signalling. *Nature* **532**, 122–126 (2016).

9.  Polster, S. P. *et al.* Atorvastatin Treatment of Cavernous Angiomas with Symptomatic

Hemorrhage Exploratory Proof of Concept (AT CASH EPOC) Trial. *Clin. Neurosurg.* **85**,

843–853 (2019).

10. Hong, C. C. *et al.* Cerebral cavernous malformations are driven by ADAMTS5 proteolysis

of versican. *J. Exp. Med.* **217**, (2020).

11.   Tang, A. T. *et al.* Endothelial TLR4 and the microbiome drive cerebral cavernous

malformations. *Nature* **545**, 305–310 (2017).

12.   Li, W. *et al.* Propranolol inhibits cavernous vascular malformations by β1 adrenergic

receptor antagonism in animal models. *J. Clin. Invest.* **131**, (2021).

13.   Shi, C. *et al.* B-Cell Depletion Reduces the Maturation of Cerebral Cavernous

Malformations in Murine Models. *J. Neuroimmune Pharmacol.* **11**, 369–377 (2016).

14.   Gibson, C. C. *et al.* Strategy for identifying repurposed drugs for the treatment of cerebral

cavernous malformation. *Circulation* **131**, 289–299 (2015).

15.   Shenkar, R. *et al.* Rho kinase inhibition blunts lesion development and hemorrhage in

murine models of aggressive Pdcd10/Ccm3 disease. *Stroke* **50**, 738–744 (2019).

16.   McKerracher, L. *et al.* A Brain-Targeted Orally Available ROCK2 Inhibitor Benefits Mild

and Aggressive Cavernous Angioma Disease. *Transl. Stroke Res.* **11**, 365–376 (2020).

17.   Shenkar, R. *et al.* RhoA kinase inhibition with fasudil versus simvastatin in murine models

of cerebral cavernous malformations. *Stroke* **48**, 187–194 (2017).

18.   Lopez-Ramirez, M. A. *et al.* Cerebral cavernous malformations form an anticoagulant

vascular domain in humans and mice. *Blood* **133**, 193–204 (2019).

19.   Detter, M. R. *et al.* Novel Murine Models of Cerebral Cavernous Malformations.

*Angiogenesis* **23**, 651–666 (2020).

20.   Cardoso, C. *et al.* Novel Chronic Mouse Model of Cerebral Cavernous Malformations.

*Stroke* **51**, 1272–1278 (2020).

21.   Mikati, A. G. *et al.* Dynamic Permeability and Quantitative Susceptibility. *Stroke* **45**, 598–

601 (2014).

22.   Girard, R. *et al.* Vascular permeability and iron deposition biomarkers in longitudinal

follow-up of cerebral cavernous malformations. *J. Neurosurg.* **127**, 102–110 (2016).

23.   Mikati, A. G. *et al.* Vascular permeability in cerebral cavernous malformations. *J. Cereb.*

*Blood Flow Metab.* **35**, 1632–1639 (2015).

24.    Yadla, S. *et al.* Cerebral cavernous malformations as a disease of vascular permeability:

from bench to bedside with caution. *Neurosurg. Focus* **29**, 1–7 (2010).

25.    Gorick, C. M. *et al.* Applications of focused ultrasound-mediated blood-brain barrier

opening. *Adv. Drug Deliv. Rev.* **191**, 114583 (2022).

26.    Fisher, D. G. & Price, R. J. Recent advances in the use of focused ultrasound for

magnetic resonance image-guided therapeutic nanoparticle delivery to the central

nervous system. *Front. Pharmacol.* **10**, 1348 (2019).

27.    Timbie, K. F., Mead, B. P. & Price, R. J. Drug and gene delivery across the blood-brain

barrier with focused ultrasound. *J. Control. Release* **219**, 61–75 (2015).

28.    Shang, X., Wang, P., Liu, Y., Zhang, Z. & Xue, Y. Mechanism of low-frequency

ultrasound in opening blood-tumor barrier by tight junction. *J. Mol. Neurosci.* **43**, 364–369

(2011).

29.    Sheikov, N. *et al.* Brain arterioles show more active vesicular transport of blood-borne

tracer molecules than capillaries and venules after focused ultrasound-evoked opening of

the blood-brain barrier. *Ultrasound Med. Biol.* **32**, 1399–1409 (2006).

30.    Fisher, D. G. *et al.* Focused Ultrasound Blood-Brain Barrier Opening Arrests the Growth

and Formation of Cerebral Cavernous Malformations. *bioRxiv* 2024.01.31.577810 (2024)

doi:10.1101/2024.08.27.609060.

31.    Fisher, D. G. *et al.* Magnetic Resonance Imaging of Mouse Cerebral Cavernomas Reveal

Differential Lesion Progression and Variable Permeability to Gadolinium. *Arterioscler.*

*Thromb. Vasc. Biol.* **43**, 958–970 (2023).

32.    Hart, B. L., Taheri, S., Rosenberg, G. A. & Morrison, L. A. Dynamic Contrast-Enhanced

MRI Evaluation of Cerebral Cavernous Malformations. *Transl. Stroke Res.* **4**, 500–506

(2013).

33.    Plummer, N. W. *et al.* Loss of p53 Sensitizes Mice with a Mutation in Ccm1 (KRIT1) to

Development of Cerebral Vascular Malformations. *Am. J. Pathol.* **165**, 1509–1518 (2004).

34.  Marty, B. *et al.* Dynamic study of blood-brain barrier closure after its disruption using

ultrasound: A quantitative analysis. *J. Cereb. Blood Flow Metab.* **32**, 1948–1958 (2012).

35.  Lee, Y. *et al.* Improvement of glymphatic–lymphatic drainage of beta-amyloid by focused

ultrasound in Alzheimer's disease model. *Sci. Rep.* **10**, 16144 (2020).

36.  Meng, Y. *et al.* Glymphatics Visualization after Focused Ultrasound-Induced Blood–Brain

Barrier Opening in Humans. *Ann. Neurol.* **86**, 975–980 (2019).

37.  Han, M., Seo, H., Choi, H., Lee, E. H. & Park, J. Localized Modification of Water Molecule

Transport After Focused Ultrasound-Induced Blood–Brain Barrier Disruption in Rat Brain.

*Front. Neurosci.* **15**, (2021).

38.  Aryal, M. *et al.* Effects on P-Glycoprotein Expression after Blood-Brain Barrier Disruption

Using Focused Ultrasound and Microbubbles. *PLoS One* **12**, e0166061 (2017).

39.  Cho, H. S. *et al.* Localized Down-regulation of P-glycoprotein by Focused Ultrasound and

Microbubbles induced Blood-Brain Barrier Disruption in Rat Brain. *Sci. Rep.* **6**, 1–10

(2016).