The role of phosphorylation and dephosphorylation in the regulation of Rubisco activase

Nikita Bhatnagar[a], , 0000-0001-7832-1347

Sarah S. Chung[a,b], sohyeon2@illinois.edu, 0009-0005-6096-5343

John Hodge[a,b], jghodge@illinois.edu, 0000-0003-0793-1871

Ryan A Boyd[a,b], raboyd2@illinois.edu, 0000-0003-4009-7700

Sang Yeol Kim[b]

Mia Sands[b], xin7@illinois.edu, 0000-0002-4925-9772

Andrew D. B. Leakey[a,b,c], leakey@illinois.edu, 0000-0001-6251-024X

Donald R. Ort[a,b,c], d-ort@illinois.edu, 0000-0002-5435-4387

Steven Burgess[a,b], sjb287@illinois.edu, 0000-0003-2353-7794

Affiliations

[a]Carl R. Woese Institute for Genomic Biology, University of Illinois Urbana-Champaign

[b]Department of Plant Biology, University of Illinois Urbana-Champaign

[c]Department of Crop Sciences, University of Illinois Urbana-Champaign

Corresponding Author: Steven J. Burgess

The author responsible for distribution of materials integral to the findings presented in this article in

accordance with the policy described in the Instructions for Authors

(https://academic.oup.com/plphys/pages/General-Instructions) is (insert author name).

Running head:

Role of phosphorylation/dephosphorylation the regulation of of Rubisco activase

SUMMARY

Rubisco activase is an ATP-dependent chaperone that facilitates dissociation of inhibitory sugar

phosphates from the catalytic sites of ribulose-1,5-bisphosphate carboxylase/oxygenase during

photosynthesis. In Arabidopsis, Rubisco activase is negatively regulated by dark-dependent

phosphorylation of threonine 78. Here we identify chloroplast localized *SHEWENELLA-LIKE PROTEIN*

PHOSPHATASE 1 (SLP1) as the major phosphatase responsible for light-dependent dephosphorylation of

Rubisco activase. The prevalence of threonine 78 in Rubisco activase was investigated across sequences

from 91 plant species, finding 29 (~32%) species shared a threonine in the same position. Analysis of

seven C3 species with an antibody raised against a threonine 78 phospho-peptide demonstrated that this

position is phosphorylated in multiple genera. However, light-dependent dephosphorylation of threonine

78 was observed only in Arabidopsis. Further, phosphorylation of threonine 78 could not be detected in

any of the four C4 grass species examined. The results suggest that despite conservation of threonine 78

in Rubisco activase from a wide range of species, a regulatory role for phosphorylation at this site is more

limited. This provides a case study for how variation in post-translational regulation can amplify

functional divergence across the phylogeny of plants beyond what is explained by sequence variation in a

metabolically important protein.

INTRODUCTION

Rubisco catalyzes the addition of CO_2 to a molecule of ribulose-1,5-bisphosphate (RuBP) forming two molecules of 3-phosoglycerate (3-PGA) and in many circumstances is the rate limiting step of CO_2 assimilation (Stitt *et al.*, 1991; Hudson *et al.*, 1992; Stitt & Schulze, 1994; Poolman *et al.*, 2000; Zhu *et al.*, 2007). To perform this reaction the enzyme must first be activated through attachment of CO_2 to a catalytic lysine in order to bind Mg^{2+} (Lorimer and Miziorko, 1980). However, the enzyme is susceptible to inhibition by attachment of RuBP prior to activation, or binding of one of several inhibitory sugar phosphates to the active site (Orr et al., 2023). Inhibitors can be released by the action of Rubisco activase (Rca) (Somerville et al., 1982; Salvucci et al., 1986; Jiang et al., 1994), which uses energy from ATP hydrolysis to drive conformational changes in Rubisco, freeing the active site (Neuwald et al., 1999; Portis & Jr., 2003; Flecken et al., 2020). Therefore Rca is critical for regulation of Rubisco activity and carbon assimilation (Parry et al., 2013; Wijewardene et al., 2021; Qu et al., 2023; Orr et al., 2023).

Rca itself is subject to regulation, which provides a link between carbon assimilation and electron transport. Most of our understanding about this process comes from Arabidopsis, where there are two isoforms of Rca which differ in size. The longer 'α' isoform possesses a 20-30 amino acid C-terminal extension (To et al., 1999) which contains two cysteine residues (Cys-451 and Cys-470). The cysteines are regulated by thioredoxin f in response to changes in the ATP/ADP arising from variation in rates of electron transport and ATP synthesis (Zhang & Portis, 1999b; Zhang et al., 2002). In the dark, the cysteines are oxidized to form a disulfide bridge (oxidized state, E_m ~344 mV pH 7.9)(Hutchison et al., 2000), increasing the sensitivity of Rca to inhibition by ADP (Portis et al., 2007), downregulating Rca activity and CO_2 fixation. However, redox regulation of Rca does not appear to be essential. Solanaceae species, such as tobacco, are reported to only express Rca-β, which lacks the C-terminal extension, (Nagarajan & Gill, 2018), and the sensitivity of Rca-β isoforms to inhibition by ADP varies (Carmo-Silva & Salvucci, 2013). Therefore, it is apparent that additional regulatory mechanisms are also important to control Rca activity.

Rca *α* and β isoforms, which differ in ADP sensitivity, can be produced by both alternate splicing (Werneke et al., 1989; Rundle & Zielinski, 1991; To et al., 1999) or from transcription of independent genes (Rundle & Zielinski, 1991; Salvucci et al., 2003). In addition, the relative expression level and catalytic properties of Rca-α and Rca-β differ across species (Rundle & Zielinski, 1991; Crafts-Brandner et al., 1997; Carmo-Silva & Salvucci, 2013; Shivhare & Mueller-Cajar, 2017; Degen et al., 2021) and can be impacted by environmental factors such as temperature (Shivhare & Mueller-Cajar, 2017; Scafaro et al., 2019; Kim et al., 2020, 2021; Degen et al., 2021) to modulate CO_2 fixation under variable conditions.

Post-translational modification of Rca has also been proposed to provide an additional layer of regulation (Amaral et al., 2024). In Arabidopsis, Threonine 78 (Thr78) is phosphorylated in the dark and

under low light intensities, but not under high light, suggesting a regulatory role (Reiland *et al.*, 2009;

Boex☐Fontvieille *et al.*, 2014; Kim *et al.*, 2016). Functional support for this hypothesis came from

several findings: (1) phosphorylation at Thr78 (Thr78-Rca) in Arabidopsis is reported to cause a

significant decrease in plant biomass, due to a decreased Rubisco activation state (Kim *et al.*, 2019); (2)

transgenic Arabidopsis plants expressing phospho-null mutants of Rca (T78A) had enhanced

photosynthetic efficiency and plant growth; and (3) substitution of Thr78-to-Ser (T78S) resulted in hyper-

phosphorylation, leading to reduced carbon assimilation and slower growth (Kim *et al.*, 2019). cpCK2

was shown to be the major kinase responsible for dark phosphorylation of AtRca, with loss-of-function

mutant lines showing a reduction in Rca phosphorylation (Kim *et al.*, 2016). However, the phosphatase

responsible for dephosphorylating Rca in the light has not been identified, and it is not clear whether this

proposed mechanism is relevant outside of Arabidopsis. To address these issues, we sought to define the

mechanisms controlling Thr78 phosphorylation, and prevalence of this post-translational modification in

photosynthetic organisms. This was achieved by considering the following five questions, (1) what is the

enzyme response for dephosphorylation of Thr78? (2) at what stage of flowering plant evolution did

Thr78 emerge? (3) how conserved is Thr78 among photosynthetic angiosperms? (4) to what extent is

Thr78 phosphorylated in different species? (5) and is light-dependent de-phosphorylation of Thr78 a

general mechanism of controlling of RCA activity?

MATERIAL AND METHODS

Determination of Rca isoform and photosynthetic species containing Thr78-Rca

To analyze the conservation of Thr78 in Rca protein across plant species, the sequence of

Arabidopsis Rca (AtRca: AT2G39730) was retrieved from the *Arabidopsis thaliana* TAIR10 genome

assembly (Lamesch *et al.*, 2012) on Phytozome v13 (Goodstein *et al.*, 2012). Syntelogs to AtRca (i.e.,

shared positional homology along chromosomes) were subsequently identified in monocots (*Setaria*

viridis Rca: Sevir.8G261433) and asterids (*Mimulus guttatus* Rca: Migut.F00758), which have homology

to this sequence using either synteny mapping features on CoGe (Lyons & Freeling, 2008; Lyons *et al.*,

2008). Setaria Rca was chosen as an example of a gene that has been previously characterized (Kim et al.,

2021) while Mimulus shares comparable synteny to Arabidopsis in the genomic region surrounding Rca

enabling identification of syntelogs in the asterids. Working from these three representatives, additional

syntelogues were locally identified based on their closest known Rca sister lineage within the Angiosperm

phylogeny across other accessible plant genomes on these databases. *Amborella trichopoda* was

leveraged as an outgroup for flowering plants in this case study given that this is known to be the one of

the earliest diverging lineages for flowering plants (Soltis *et al.*, 2011). Additionally, homologues were

surveyed, which contained threonine at position 78 phosphorylation site even if they didn't precisely fit

this stringent criterion of synteny using pBLAST on NCBI (https://blast.ncbi.nlm.nih.gov/Blast.cgi) for

key agronomic model systems. This ultimately resulted in 217 isoform CDS sequences being identified.

Alternative splicing of the same locus in some lineages meant that these CDS sequences represented 153

unique genes from across 91 species and 50 plant families. The longer, Rca-α, sequence was

preferentially retained as a taxonomic representative for each alternative splicing gene when available for

phylogenetic analysis. The CDS sequence of these genes were visualized in Seaview (Gouy *et al.*, 2010)

and codon-aligned using MUSCLE (Edgar, 2004). Obvious pseudogenes that were missing large portions

of their AAA+ ATPase domain such as the rice sequence (LOC_Os11g47980) were removed prior to

phylogenetic analysis of this DNA alignment. A maximum likelihood tree was subsequently assembled

using IQTree (Nguyen *et al.*, 2015) with the Amborella Rca sequence being specified as an outgroup and

1000 bootstraps being run to generate statistical support for nodes along the tree with the automatic

substitution model selection determining a SYM+R8 model was the best-fit according to Bayesian

Information Criterion. The resulting treefile was then visualized either in Figtree

(http://tree.bio.ed.ac.uk/software/figtree/) (Figure S3) or in R using the Phytools and Ape libraries

(Paradis & Schliep, 2019; Revell, 2024). Ancestral state reconstructions were performed using the Ape

function *ace* particularly for the original position 78 amino acid identity between syntelogous genes, and

the manner of α/ß origin (either by alternative splicing or distinct genes) for each plant family (Figure

S4).

130   Plant growth and stress treatment

Arabidopsis Col-0 and camelina seeds were sterilized in 1 % (v/v) bleach for 5 min, rinsed with 70 %

EtOH, and dried on filter paper. The sterilized seeds were sown on MS (Murashige and Skoog basal salt

mixture powder, Phyto Technology Laboratories, M524) agar (0.8 %, w/v, Sigma-Aldrich, A1296) with 1

% (w/v) sucrose. The plates were stored at 4 °C for 2 days for cold stratification before transfer to a

growth chamber. Seedlings were transferred into pots (5 cm diameter) filled with soil (Berger; PRO-MIX

BX All Purpose Growing Mix) before the third true leaf emerged, which was approximately 5 to 7 days

after germination (DAG). The other plants (brassica, soybean, green bean, and Medicago) were directly

sown in soil after sterilization. Arabidopsis homozygous mutants of SLP1 (SALK_129247C) were

ordered from ABRC, and homozygous lines were confirmed by genomic DNA PCR using LB_6313R

(TCAAACAGGATTTTCGCCTGCT), SLP1_LP (GCCTGATCATCAAGTGACCTC) and SLP1_RP

(CGGGAACCATGTCTGACTTAG) xfs designed using iSect tool by SALK

(http://signal.salk.edu/tdnaprimers.2.html) (Alonso *et al.*, 2003). All plant species were grown at 22 °C,

16 h light and 8 h dark, 200 µmol m^{-2} s^{-1} PPFD with 2100 K 400W high pressure sodium bulbs

(Ceramalux ALTO; C400S51).

Protein extraction and immunoblotting assay

Total proteins were extracted from three frozen leaf discs (1 cm in diameter) using a Qiagen Tissuelyzer for tissue grinding. The ground tissue was suspended in 0.5 mL of extraction buffer containing 62.5 mM Tris-HCl (pH 6.8), 2% SDS, 10% glycerol, 0.35 M 2-mercaptoethanol, and 0.001% protease inhibitor cocktail (Sigma-Aldrich; P9599). The suspension was heated at 95°C for 5 minutes to denature proteins and inactivate proteases, followed by centrifugation at 13,500 rpm for 30 minutes at 4°C. The supernatant was collected, and 10 µL aliquots were separated on 10% SDS-PAGE gels. Proteins from the gels were transferred electrophoretically to 0.2 µm nitrocellulose membranes (Bio-Rad) and blocked with PBS solution with 5% (w/v) non-fat milk powder (Sigma) for 1h at room temperature. For immunodetection, primary antibodies were diluted in Intercept Blocking Buffer (LI-COR; 927-70001); anti-Rca (Agrisera; AS10 700) at 1:5000 and anti-pT78 (GenScript) at 1:4000, and incubated overnight at 4°C. Modification-specific antibody was generated against the phosphopeptide antigen RGLAYDpT78DDQQDC (Arabidopsis anti-pT78). For each peptide antigen, the Cys residue at the N- or C-terminus was added for coupling to keyhole limpet hemocyanin (KLH). All custom antibodies were produced by GenScript (Piscataway, NJ, USA). Secondary detection used a fluorescent goat anti-rabbit antibody (Invitrogen; A32735) at a 1:10,000 dilution in PBS-T, incubated for 1 h at room temperature, and visualized with a LI-COR Odyssey Infrared Imaging System. All immunoblotting experiments were conducted at least twice to ensure reproducibility of results.

Protein phosphorylation assay with phosphatase inhibitor

WT Col-0 plants were grown as described above and whole rosettes were sampled and dipped in the protein phosphatase inhibitor buffer: 1% (v/v) solution of phosphatase inhibitor cocktail 2 (Sigma: P5726) and phosphatase inhibitor cocktail 3 (Sigma: P0044), overnight under dark. For dark treatment samples, the protein extraction was done using the overnight incubated samples and for light treatment samples the rosettes were exposed to 150 µmol m^{-2} s^{-1} PPFD light for 2 h and then total protein was extracted.

Physiological analysis

For light induction analysis, the leaves of 5-week WT and *slp1* mutants were dark adapted overnight before measurement in a portable infra-red gas exchange system (LI-6800, LICOR Environmental, Lincoln, Nebraska, USA) connected to a multiphase flash fluorometer (6800-01A) with a 2 cm^2 chamber. During low light incubation the plants were managed under dark conditions to make sure that Rca remained phosphorylated in both WT and *slp1* mutants. The chamber settings were 1:1 red:blue

ratio, leaf temperature of 22 °C, CO_2 level of 420 µmol mol^{-1}, and 50% relative humidity. The

measurement started with a low light of 20 µmol m^{-2} s^{-1} photosynthetic photon flux density (PPFD) for 20

min, then the PPFD was increased to high light conditions of 1000 µmol m^{-2} s^{-1} for 60 min, measurements

were logged every 10 s. For CO_2 response analysis, plants were measured at the same high light

conditions described above after finishing the light induction measurement. The CO_2 level was adjusted

to produce a plot of net CO_2 assimilation against intercellular CO_2 concentration (A-C_i curve). The CO_2

levels were changed every 90 – 120 s in the following order: 420, 300, 200, 100, 50, 10, 420, 420, 420,

600, 800, 1200, 1800 µmol mol^{-1}. The relative comparisons and significant differences between *slp1* and

WT were determined using students t-test (n = 4 to 6).

Statistical Analysis

Comparisons of rosette size and assimilation rate between *slp1* and WT were done using a two-tailed

Students' t-test using data from four to six biological replicates (individual plants) in Microsoft Excel

v16.88.

RESULTS

SHEWENELLA-LIKE PROTEIN PHOSPHATASE 1 (SLP1) dephosphorylates Rca in-vivo

Light-dependent dephosphorylation of AtRca suggested that an unknown protein phosphatase is

operating. To test this hypothesis, the phosphorylation state of Rca was assessed under light and dark

conditions in the presence and absence of a protein phosphatase inhibitor cocktail (PPIC) (Fig. 1A and B).

Under dark conditions, Rca phosphorylation increased compared to the sample lacking PPIC (Fig. 1B)

which is consistent with the presence of a protein phosphatase. Dephosphorylation of Rca occurred in the

light (Fig. 1B), even in the presence of the PPIC, which may be because the level of dephosphorylation

exceeded the ability of the PPIC to inhibit activity. To identify the gene responsible for

dephosphorylation of Rca, all known chloroplast localized protein phosphatases in Arabidopsis (White-

Gloria *et al.*, 2018) were screened for reported associations with Rca in the literature and co-expression

databases. Out of 9 chloroplast localized protein phosphatases, AtSLP1 (AT1G07010) showed a potential

association based on co-expression (Fig. 1C). We hypothesized that in the absence of *AtSLP1*, AtRca will

not be dephosphorylated in the light. To test this, a homozygous T-DNA insertion mutant of *slp1* was

identified (Figure S1) and assessed for the ability to dephosphorylate Rca in the dark and light. *slp1*

mutants showed phosphorylation of Rca in the dark, as also observed in the case of Col-0, but in the

mutant phosphorylation was observed to persist in the light (Fig. 1D).

We hypothesized that lack of Rca dephosphorylation in *slp1* mutants would lead to decreased

photosynthesis and a growth penalty. Analysis of 5-week-old plants found *slp1* mutants have a significant

reduction in rosette size relative to Col-0 under normal light conditions (Fig. 1E and F). To assess the
impact of SLP1 on steady state photosynthesis, A/Ci (net CO_2 assimilation vs internal CO_2 concentration)
curves were performed on Col-0 and *slp1* mutants, finding no difference between lines (p=0.05; Figure
S2). As there is a tight correlation between Rubisco activation state and the rate of induction of
photosynthesis during shade to sun transitions, the impact of SLP1 on the rate of photosynthetic induction
after a step-increase in photosynthetic photon flux density from 20 to 1000 µmol m^{-2} s^{-1} was assessed,
again finding no significant difference compared to Col-0 (p=0.05; Fig. 1G).

Thr 78 in Rca is common

To study the conservation of Thr78 (Arabidopsis numbering) in Rca, 1893 protein sequences
from 479 plants species were assessed. Among these, 203 species retained Thr78 (Table S1). Next, a
more stringent synteny informed ortholog phylogeny of Rca-α and Rca-β isoforms was generated across
angiosperms to address the questions: at what stage of flowering plant evolution did the putative Thr78
phosphosite emerge, and how well conserved is it among photosynthetic angiosperms? Leveraging
maximum likelihood gene family phylogeny ancestral states, Thr78 distribution was estimated for each
family, including deeper nodes within the tree such as the branch point of the *Amborellaceae* (Fig. 2). The
presence of Thr78 was observed in most plant families (Fig. 2), while among the *Poaceae*, that includes
major crop species, the presence of Isoleucine (Ile) was found to dominate at position 78 compared to
Threonine (Fig. 2). The difference in the abundance of both amino acids, Thr and Ile at this site is similar
at the evolutionary scale, and an estimate of the ancestral state was ambiguous. Interestingly, the *α* and *β*
copies appeared polyphyletic, with multiple origins of the two isoforms arising from alternate splicing or
gene duplication across the phylogeny (Fig. S4, and S5). Within the Brassicaceae *Arabidopsis thaliana*
and its immediate relatives all appear to have a single copy of Rca with alpha and beta forms being the
result of alternative splicing of a single transcript.

By contrast in the Fabaceae, *Glycine max* has four gene copies of Rca (two each of *α* and *β*) with
the Papillionoid legumes more generally having two copies of Rca. There is no equivalent duplication
event in *Cercis canadensis* that would indicate a Fabaceae-wide pattern. Therefore, this pattern of two
Papillionoid copies is consistent with the whole-genome doubling (WGD) event preceding the
diversification of this subfamily, and by undergoing an additional allopolyploid event, *Glycine max* has
two additional copies for each of these ancestral homeologues (Gill, *et al.*, 2009; Koenen, *et al.*, 2021).
Within Poaceae, *Setaria viridis* and the grasses more generally displayed two mirrored clades, which
reflect the known taxonomic sorting of this family. The grass family has undergone three WGD events (τ,
σ, and ρ), two of which are shared with monocots, with the p WGD unique to the Poaceae (Paterson *et al.*,
2004; Tang *et al.*, 2010; McKain *et al.*, 2016). Although superficially resembling the *ρ* WGD event, these

grass copies are physically proximal to their sister copies in the adjacent clade suggesting that this may

instead be an ancient tandem duplication event.

Light-dependent dephosphorylation of Thr78 in Rca is unique to Arabidopsis

Previous reports showed that both AtRca isoforms phosphorylate in the dark and are

dephosphorylated as plants are exposed to light. To test whether Thr78 phosphorylation status is light-

sensitive in species other than Arabidopsis, we screened three C3 plants from the *Brassicaceae* for dark-

dependent phosphorylation of Thr78, using a pT78 antibody (Kim et al., 2019) (Fig. 3A). As in

Arabidopsis, Rca-α and Rca-β isoforms arise from alternate splicing in *Brassicaceae* and the Thr78 site is

conserved in all proteins (Fig. 3B). However, unlike Arabidopsis, Rca in *Camelina sativa* (*Cs*), *Brassica*

oleraceae (*Bo*) and *Brassica napus* (*Bn*) phosphorylated in dark but lacked light-induced

dephosphorylation (Fig. 3A). These data suggest that although Thr78 is phosphorylated in multiple

species, light-dependent regulation may be specific to Arabidopsis.

To broaden the analysis, we performed the dephosphorylation assay on several legumes: *Glycine*

max (*Gm*), *Phaseolus vulgaris* (*Pv*), and *Medicago truncatula* (*Mt*). The Rca isoforms in these species are

derived from individual genes of *Rca-α* and *Rca-β*, unlike the situation in the *Brassicaceae*. The Rca-α

isoforms retain Thr78, whereas the β isoforms have Ile78 in *Gm* and *Mt,* and Val78 in *Pv* (Fig. 3B).

Publicly available data suggest the transcription level of Rca-α is low relative to β in both *Gm* and *Pv,*

and moderately expressed in *Mt* (Table S2). The light/dark assay on legume plants showed that both the

isoforms were expressed, and phosphorylated in the dark, but as with the Brassicaceae, did not

dephosphorylate in the light (Fig. 3A). The reason why the antibody was able to detect phosphorylation of

β forms, which lack Thr78, is unclear. It is possible that there is relaxed selectively, but we consider these

results with caution.

To study the impact of phosphorylation in C4 grass plants, we analyzed Rca phosphorylation in

four species of C4 grass: *Sorghum bicolor* (*Sb*), *Miscanthus* (*Ms*), *Saccharum officinarum* (*Su*), and

Setaria viridis (*Sv*) (Fig. 4A). Each of these species possess two separate genes for the alternative Rca

isoforms, with Rca-α containing Thr78, but with it lacking from Rca-β (Fig. 4B, C). In accordance with

previous reports (Kim *et al.*, 2021), only the Rca-β isoform could be detected under normal growth

conditions (Fig. 4A and C). An unspecific cross-reaction was present in C4 grass samples probed with

pT78 antibody, but the detected protein was too large to be Rca-β, subsequently no phosphorylation of

Rca-β was detected (Fig. 4A). To test whether a C4 Rca-α isoform could be phosphorylated, sorghum

plants were exposed to heat induce Rca-α isoform expression (Fig. 4C). Plants accumulated SbRca-α

after 24 h heat treatment, but no bands of the correct size could be detected using the pT78 antibody (Fig.

4C). To rule out the possibility that a lack of detectable Thr78 phosphorylation was due to high protein

phosphatase activity, the assay was repeated in the presence of a protein phosphatase inhibitor cocktail

(PPIC), but again protein phosphorylation was not observed under any condition (Fig. 5). This result

suggests that SbRca-α T78 is not phosphorylated by cpCK2 as it is in Arabidopsis even though the gene

for cpCK2 is present in sorghum.

DISCUSSION

Rca is subject to multiple levels of regulation to adjust carbon fixation in response to

environmental conditions (Amaral *et al.*, 2024). Differences in gene expression, biochemical properties of

isoforms, redox status, and post-translational modification of Rca have been topics of investigation

(Boex□Fontvieille *et al.*, 2014; Kim *et al.*, 2016, 2019). Dark dependent phosphorylation of Thr78 is

proposed to downregulate Rca activity leading to reduced assimilation (Kim et al. 2019). cpCK2 is

described as the major kinase responsible for phosphorylation of AtRca Thr78 based on dramatically

reduced phosphorylation in *cpcpk2* T-DNA insertion lines (Kim *et al.*, 2016). Here we identified

Shewanella-like protein phosphatase 1 (SLP1) as the second component of this regulatory mechanism,

acting to dephosphorylate AtRca Thr78 in the light (Fig. 1). SLP1 is a highly conserved protein

phosphatase reported to be blue light responsive in Arabidopsis (Depuydt & Vandepoele, 2021) and

belongs to a protein family consisting of two members, of which, SLP1 is the only chloroplast localized

isoform (Uhrig & Moorhead, 2011). A role for mitochondrial targeted SLP2 has been described in

negative regulation of gibberellic acid related seed germination (Uhrig *et al.*, 2017), but the physiological

role and targets of SLP1 have not previously been reported.

If phosphorylation of Rca has a dominant impact on activity, plants lacking SLP1 would be

predicted to see an increase Thr78 phosphorylation in the light, leading to lower Rubisco activity, and

lower A_{net}, an impact on photosynthesis, with no difference observed in assimilation rates between WT

and knockout lines during light induction or CO_2 response (Fig. 1G and Fig. S2). These findings are in

keeping with observations by Kim *et al.* (2016), who reported no change in assimilation rates when Thr78

phosphorylation state was reduced in *cpck2* mutants. These results were attributed to redundancy in Rca

regulation between protein phosphorylation and control by thioredoxin (Kim *et al.*, 2016) and the same

rationale may apply here. Support for a functional role for phosphorylation of Thr78 in Arabidopsis has

only been shown in Rca variant lines lacking the redox sensitive C-terminal extension (Kim *et al.*, 2019).

Our analysis, conducted on lines containing the C-terminal extension, are therefore consistent with the

hypothesis that Rca possesses multiple redundant regulatory mechanisms in Arabidopsis, which operate

to regulate activity in response to variable light environments. The impact of SLP1 on photosynthesis

may only be apparent in the absence of redox regulation.

Despite the absence of impact on A_{net}, the *slp1* mutant showed reduced growth (Fig. 1). Given
that the *cpck2* mutant, which also showed altered phosphorylation of Rca, did not have a growth
phenotype, we attribute this to an unrelated impact on chloroplast physiology. As a largely
uncharacterized protein, the specificity and targets of SLP1 are unknown. Phosphatases have traditionally
been considered promiscuous, with specificity dictated by regulatory subunits (Virshup and Shenolikar,
2009). SLP1 is not known to have regulatory subunits, it is therefore conceivable there are other targets
within the chloroplast that are impacted in the mutant leading to the observed growth phenotype.

Substitution of Thr78 with Serine lead to a relative increase in phosphorylation at the site, in
Arabidopsis resulted in smaller plants with decreased photosynthesis (Kim *et al.*, 2019). Altering a single
amino acid, Thr78, was proposed as a means of generating plants with better photosynthesis. However,
the prevalence of Thr78 across species, the frequency of post-translational modification and functionality
were unclear. To begin to address these questions we expanded previous searches of genomic databases,
which assessed the evolution of Rca copy number across the green lineage (Nagarajan & Gill, 2018; Kim
et al., 2019), to use a synteny based approach. Nagarajan and Gill (2018) placed the emergence of the
Rca-α CTE and alternate splicing mechanisms in the charophytes, before the divergence of land plants
and there is only one reported example of a species only having evidence for expression of Rca α
(Nagarajan & Gill, 2018). Here, despite α and β forms (bearing or lacking a c-terminus respectively)
occurring within each model system, these α and β forms appear to represent repeated cases of
evolutionary convergence on these two isoforms (Fig S5) through distinct means within each clade:
subfunctionalization (Birchler & Yang, 2022), following tandem duplication in the grasses as previously
described (Nagarajan & Gill, 2018), WGD in the legumes, or alternative splicing of a single coding
sequence into alternative isoforms in Brassicaceae. This may suggest that Rca isoforms can be quite
plastic in the way they arise within distinct plant families where α and β forms may not necessarily reflect
true homology within Rubisco activase isoforms.

Previous assessment of a smaller number of Rca sequences failed to identify any with Thr78
replaced by Serine, leading to speculation that the configuration with the attendant hyper phosphorylation
observed in Arabidopsis has been selected against (Kim *et al.*, 2019). In the initial set of Rca sequences
identified by BLAST search, we identified gene models from at least seven species (Table S1) which do
have Ser78, although these were not considered orthologs in the synteny analysis. Of the 25 species found
to contain Rca sequences with Thr78, it remains to be seen how many undergo phosphorylation. Our data
are consistent with it occurring in at least the Brassicaceae, while phosphorylation of an equivalent Thr
residue has been reported in tomato, suggesting this modification may occur more widely (Marques et al.,
2021).

We found no evidence for phosphorylation of Thr78 in monocot species, including Sorghum,

Setaria, sugarcane and Miscanthus. The peptide fragment used for antibody generation is highly

conserved, with 100% identity between Arabidopsis Rca and Sorghum Rca-α "RGLAYDTSDDQQDC"

and should therefore have been able to detect phosphorylation if present under the conditions tested.

There are several potential explanations for this finding, including an absence of the kinase, differences in

localization or expression profiles of the kinase, or regulation of the kinase. In Arabidopsis, Rca is

phosphorylated by cpCK2 (Kim et al., 2016). A homologous gene has been characterized in rice

(LOC_Os03g55490) (Lu et al., 2015) and syntenic sequences can be found in C4 grasses including

Sorghum (Sobic.001G07970) (Carvalho et al., 2020). As Sorghum operates two-celled C4 photosynthesis

for an interaction to occur between Rca and cpCK2 both must be expressed in the same cell type.

Transcriptomics suggests that cpCK2 is expressed in both mesophyll and bundle sheath cells in maize

(Chang et al., 2012), Setaria (John et al., 2014) and Sorghum (Döring et al., 2016), albeit with ~2-fold

mesophyll enrichment in maize and Setaria whereas Rca is localized in the bundle sheath. Even so, it

appears that the orthologous cpCK2 is present in the same compartment as Rca (i.e., the bundle sheath

cell chloroplast). However, in the monocot species tested Rca-β, which lacks Thr78, is the predominate

isoform under normal conditions and Rca-α, which possess Thr78, is induced during stress (Shivhare &

Mueller-Cajar, 2017; Scafaro et al., 2019; Kim et al., 2020, 2021). Whether C4 monocot cpCK2 retains

the ability to phosphorylate Thr78 in Rca-α, or if the bundle sheath stromal environment is permissive for

phosphorylation remains unclear (Kim et al., 2016). CK2 exists as a multigene family, composed of

catalytic α subunits and regulatory beta subunits (Salinas et al., 2006; Schweer et al., 2010), which form

hetero-tetramers that can differ in substrate specificity (Domańska et al., 2005). Intriguingly there appears

to be a lack of regulatory β subunits in the chloroplast (Salinas et al., 2006), so differences in regulatory

subunits impacting specificity appears unlikely, although multiple chloroplast CK2 containing complexes

have been detected (Baginsky et al., 1997) suggesting the protein does not act alone. However, further

research is required to characterize the regulation of monocot cpCK2 and whether it can phosphorylate

Rca.

Conservation of phosphorylation of Thr78 in some C3 plants is not evidence of a conserved

regulatory role. Indeed, in the species examined, strong evidence for light-dependent dephosphorylation

of Thr78 was only found for Arabidopsis, with phosphorylation of Rca detected in both the light and dark

in other C3 species (Fig. 3). These data may suggest a limited scope for phosphorylation as a means of

regulation, with most species depending on other mechanisms, such as changes in redox state, catalytic

properties or expression. It may be that alternate post-translational modifications of Rca have greater

importance in other species and several different ones have been described (Amaral et al., 2024).

Interestingly, there are several reports of phosphorylation of S79 in Arabidopsis (Willems et al., 2019),

and this modification was also observed in a Brachypodium, Chlamydomonas and maize (Willems et al.,
2019, 2024). Further phosphoproteomics of maize during a diurnal cycle suggests that rather than Thr78,
Ser79 undergoes dark-dependent phosphorylation (Gao et al., 2023). However, the prevalence of S79
phosphorylation is also likely to vary, as seen by a failure to detected in protein samples from *Glycine*
max (soybean-S86) (Soleh et al., 2017) .Therefore, the prevalence and function of Ser79 phosphorylation
is unclear, and whether it can play a similar role to Thr78 warrants further investigation.

Taken together our results suggest that Thr78 can be found in Rca sequences across the green
lineage, although the prevalence varies, and degree to which it undergoes phosphorylation is inconsistent.
Our data provides support for a mechanism whereby Thr78 is phosphorylated in the dark by cpCK2 and
dephosphorylated in the light by SLP1 in Arabidopsis. The lack of impact on photosynthesis when cpCK2
or SLP1 are knocked out suggest that phosphorylation of Thr78 acts redundantly with redox regulation.
More widely, it is likely that regulation of Rca can occur through multiple mechanisms, including redox,
sensitivity to ADP and gene expression and the relative importance of each varies.
394

Acknowledgments
We would like to thank Toby Kellogg and Shawn Abrahams for providing feedback on the phylogenic
analyses. This work was funded by the DOE Center for Advanced Bioenergy and Bioproducts Innovation
(U.S. Department of Energy, Office of Science, Biological and Environmental Research Program under
Award Number DE-SC0018420). Any opinions, findings, and conclusions or recommendations expressed
in this publication are those of the author(s) and do not necessarily reflect the views of the U.S.
Department of Energy.

Competing interests
The authors report no conflict of interest
405

406 **Authors contributions**
407 Conceptualization (SK, NB, SJB, DO), Investigation (NB, SC, MS, JH, SK), Formal Analysis (NB),
408 Project administration (NB, SJB, DO), Supervision (AL, SJB, DO), Visualization (NB), Writing - original
409 draft (NB, JH, SC, DO, SJB), Writing – review and editing (NB, JH, DO, AL, SJB).
410

FIGURE LEGENDS

Figure 1: SLP1 dephosphorylates Rca in Arabidopsis. Protein phosphatase assay with PPIC (+P) and without PPIC (-P) isoforms, comparing total abundance (A) to phosphorylation (B) of Rca by immunoblot. (C) Putative protein-protein interactions between Rca and chloroplast localized protein phosphatases. (D) Immunoblot analysis of light dependent dephosphorylation of Rca in WT (Col-0) and *slp1* mutants. (E) Image of 5-week-old rosettes comparing WT (Col-0) and slp1 plants. (F) Comparison of rosette size between WT (Col-0) and slp1 plants (n=20). (G) Photosynthetic induction curves of WT (Col-0) and *slp1* mutant plants (n=5 to 6).

Figure 2: Phylogenetic representation of the amino acid present at the phosphorylation site (peptide position 78) within extant Rca gene models. Ancestral states for families, and when available deeper orders of Angiosperms, are denoted at their corresponding nodes as pie charts. The nodes depicting the presence of Thr78 is colored green and the ones with Ile78 is colored navy blue. Less abundant amino acids at the peptide position 78 are depicted as varying shades of blue and purple. The detailed confirmation of bootstrap values is mentioned in Figure S3.

Figure 3: Light dependent dephosphorylation of Thr78 in Rca is specific to Arabidopsis. (A) Immunoblot analysis comparing protein phosphorylation and total abundance of Rca under light (L) and dark (D) conditions (At: *Arabidopsis thaliana*, Cs: *Camelina sativa*, Bn: *Brassica napus*, Bo: *Brassica oleraceae*, Gm: *Glycine max*, Pv: *Phaseolus vulgaris*, Mt: *Medicago truncatula*). The upper panel shows the total Rca protein extracted from plant samples and the middle panel shows the phosphorylated Rca protein. α and β denote the 2 isoforms of Rca protein. CBB (Coomassie Brilliant Blue) depicts the loading control. (B) Sequence alignment depicting the conservation of Thr78 amino acid in C3 species. The amino acid lacking color codes depicts the change in the Thr78 site to Isoleucine (*I*) in *Glycine max* and *Medicago truncatula*, and Valine (*V*) in Phaseolus vulgaris. The letter "A" and "B" at the end of the species labelling denotes the Rca-α and Rca-β isoforms. (C) The phylogenetic tree depicts the division of C3 Brassicaceae family and legumes based on Rca isoforms arising from protein splicing in Brassicaceae as compared to different genes in legumes and presence and absence of Thr78 amino acid in the Rca-β isoform. The blue boxes depict the presence of Thr78 amino acid whereas the red depicts the lack of Thr78. The yellow box depicts the experimental confirmation of protein phosphorylation at Thr78 phosphosite.

Figure 4: Phosphorylation of Rca is not observed in selected C4 moncots. (A) Protein phosphorylation assay depicting four C4 plant samples in light and dark conditions (At: *Arabidopsis thaliana*, Sv: *Setaria viridis*, Sb: *Sorghum bicolor*, Su: *Saccharum officinarum*, Ms: *Miscanthus*). The upper panel shows the accumulation of Rca protein quantified with Anti-Rca antibody and middle panel shows the

phosphorylated Rca protein quantified with Anti-pT78 antibody. α and β denote the 2 isoforms of Rca

protein. CBB denotes the loading control. (B) Sequence alignment of motif comparing Arabidopsis and

select C4 plants. (C) Immunoblot analysis comparing Rca phosphorylation under heat for 24 h in WT

sorghum. The numbers denote the time in hours after the heat stress. CBB denotes the loading control.

Figure 5: Sorghum lacks phosphorylation of Thr78 with or without PPIC treatment. (A) Immunoblot

analysis comparing sorghum WT samples collected after heat stress at 42^{o}C under light and dark

treatment. The control samples were collected at 28^{o}C. The + and – sign denotes presence and absence of

treatment, respectively. The left panel shows Rca protein accumulation and right panel depicts analysis

using anti-pT78 antibody.

Figure S1: The genomic DNA (gDNA) PCR with primer combinations of LB, LP and RP showed that

LB primer T-DNA insertion was confirmed in homozygous lines whereas WT showed no bands (upper

panel). The SLP1 gene primers showed gDNA bands in the WT and not in the *slp1* mutants (lower panel).

Figure S2: The *ACi* curve showed that both WT (Col-0) and slp1 mutants overlap suggesting that both

the genotypes have no difference in the photosynthetic efficiency. Error bars denote standard error. Green

dots are *slp1* mutants and black dots are WT (Col-0).

Figure S3: The conservative phylogeny structure of Rca orthologs where all nodes with bootstrap support

(blue numbers) greater than 80% being retained whereas other lower support nodes were collapsed into

polytomies.

Figure S4: Rca α (white) and β (black) isoforms on extant gene models as well as alternatively spliced

gene models which produce both Rca α and β (grey). Ancestral states for families, and when available

deeper orders of Angiosperms, are denoted at their corresponding nodes as pie charts.

Figure S5: The likely pattern that could result in the polytomies of α and β isoforms ancestry states that

Angiosperms initially beginning as Rca α lineages out of which β isoforms repeatedly arise

independently following either subfunctionalization of duplicate gene copies or alternative splicing of α

gene models to β isoforms as well. Within certain lineages where α isoforms are lost all the Rca copies

become canalized into a β isoform state thereafter.

Table S1: List of Rca protein sequences identified by BLAST search.

Table S2: Relative expression and residue at site equivalent to AtRca T78 for Rca isoforms from *G. max*,

M. truncatula and *P. vulgaris*.

REFERENCES

Alonso JM, Stepanova AN, Leisse TJ, Kim CJ, Chen H, Shinn P, Stevenson DK, Zimmerman J, Barajas P, Cheuk R, *et al.* **2003**. Genome-wide insertional mutagenesis of Arabidopsis thaliana. *Science (New York, N.Y.)* **301**: 653–7.

Amaral J, Lobo AKM, Carmo☐Silva E. 2024. Regulation of Rubisco activity in crops. *New Phytologist* **241**: 35–51.

Baginsky S, Tiller K, Link G. 1997. Transcription factor phosphorylation by a protein kinase associated with chloroplast RNA polymerase from mustard (Sinapis alba). *Plant Molecular Biology* **34**: 181–189.

Birchler JA, Yang H. 2022. The multiple fates of gene duplications: Deletion, hypofunctionalization, subfunctionalization, neofunctionalization, dosage balance constraints, and neutral variation. *The Plant Cell* **34**: 2466–2474.

Boex☐Fontvieille E, Daventure M, Jossier M, Hodges M, Zivy M, Tcherkez G. 2014. Phosphorylation pattern of <scp>R</scp>ubisco activase in *<scp>A</scp> rabidopsis* leaves. *Plant Biology* **16**: 550–557.

Carmo-Silva AE, Salvucci ME. 2013. The Regulatory Properties of Rubisco Activase Differ among Species and Affect Photosynthetic Induction during Light Transitions . *Plant Physiology* **161**: 1645–1655.

Carvalho DS, Nishimwe A V., Schnable JC. 2020. IsoSeq transcriptome assembly of C $_3$ panicoid grasses provides tools to study evolutionary change in the Panicoideae. *Plant Direct* **4**.

Chang Y-M, Liu W-Y, Shih AC-C, Shen M-N, Lu C-H, Lu M-YJ, Yang H-W, Wang T-Y, Chen SC-C, Chen SM, *et al.* **2012**. Characterizing Regulatory and Functional Differentiation between Maize Mesophyll and Bundle Sheath Cells by Transcriptomic Analysis . *Plant Physiology* **160**: 165–177.

Crafts-Brandner SJ, van de Loo FJ, Salvucci ME. 1997. The Two Forms of Ribulose-1,5-Bisphosphate Carboxylase/Oxygenase Activase Differ in Sensitivity to Elevated Temperature. *Plant Physiology* **114**: 439–444.

Crafts-Brandner SJ, Salvucci ME. 2002. Sensitivity of Photosynthesis in a C4 Plant, Maize, to Heat Stress. *Plant Physiology* **129**: 1773–1780.

Degen GE, Orr DJ, Carmo☐Silva E. 2021. Heat☐induced changes in the abundance of wheat Rubisco activase isoforms. *New Phytologist* **229**: 1298–1311.

Depuydt T, Vandepoele K. 2021. Multi☐omics network☐based functional annotation of unknown Arabidopsis genes. *The Plant Journal* **108**: 1193–1212.

Domańska K, Zieliński R, Kubiński K, Sajnaga E, Masłyk M, Bretner M, Szyszka R. 2005. Different properties of four molecular forms of protein kinase CK2 from Saccharomyces cerevisiae. *Acta biochimica Polonica* **52**: 947–51.

Döring F, Streubel M, Bräutigam A, Gowik U. 2016. Most photorespiratory genes are preferentially expressed in the bundle sheath cells of the C $_4$ grass *Sorghum bicolor*. *Journal of Experimental Botany* **67**: 3053–3064.

Flecken M, Wang H, Popilka L, Hartl FU, Bracher A, Hayer-Hartl M. **2020**. Dual Functions of a
Rubisco Activase in Metabolic Repair and Recruitment to Carboxysomes. *Cell* **183**: 457-473.e20.

Gao Z, Yang X, Mei Y, Zhang J, Chao Q, Wang B. **2023**. A dynamic phosphoproteomic analysis
provides insight into the <scp>C4</scp> plant maize (*Zea mays* L.) response to natural diurnal changes.
The Plant Journal **113**: 291–307.

**Goodstein DM, Shu S, Howson R, Neupane R, Hayes RD, Fazo J, Mitros T, Dirks W, Hellsten U,
Putnam N,** *et al.* **2012**. Phytozome: a comparative platform for green plant genomics. *Nucleic Acids
Research* **40**: D1178–D1186.

Gouy M, Guindon S, Gascuel O. **2010**. SeaView Version 4: A Multiplatform Graphical User Interface
for Sequence Alignment and Phylogenetic Tree Building. *Molecular Biology and Evolution* **27**: 221–224.

Hudson GS, Evans JR, von Caemmerer S, Arvidsson YBC, Andrews TJ. **1992**. Reduction of
Ribulose-1,5-Bisphosphate Carboxylase/Oxygenase Content by Antisense RNA Reduces Photosynthesis
in Transgenic Tobacco Plants. *Plant Physiology* **98**: 294–302.

Jiang C, Quick WP, Alred R, Kliebenstein D, Rodermel SR. **1994**. Antisense RNA inhibition of
Rubisco activase expression. *The Plant Journal* **5**: 787–798.

John CR, Smith-Unna RD, Woodfield H, Covshoff S, Hibberd JM. **2014**. Evolutionary Convergence
of Cell-Specific Gene Expression in Independent Lineages of C4 Grasses   . *Plant Physiology* **165**: 62–
75.

Kim SY, Bender KW, Walker BJ, Zielinski RE, Spalding MH, Ort DR, Huber SC. **2016**. The plastid
casein kinase 2 phosphorylates rubisco activase at the Thr-78 site but is not essential for regulation of
rubisco activation state. *Frontiers in Plant Science* **7**.

**Kim SY, Harvey CM, Giese J, Lassowskat I, Singh V, Singh V, Cavanagh AP, Cavanagh AP,
Spalding MH, Finkemeier I,** *et al.* **2019**. In vivo evidence for a regulatory role of phosphorylation of
Arabidopsis Rubisco activase at the Thr78 site. *Proceedings of the National Academy of Sciences of the
United States of America*.

Kim SY, Slattery RA, Ort DR. **2021**. A role for differential Rubisco activase isoform expression in C $_4$
bioenergy grasses at high temperature. *GCB Bioenergy* **13**: 211–223.

Kim SY, Stessman DJ, Wright DA, Spalding MH, Huber SC, Ort DR. **2020**. Arabidopsis plants
expressing only the redox-regulated Rca-α isoform have constrained photosynthesis and plant growth.
Plant Journal **103**: 2250–2262.

**Lamesch P, Berardini TZ, Li D, Swarbreck D, Wilks C, Sasidharan R, Muller R, Dreher K,
Alexander DL, Garcia-Hernandez M,** *et al.* **2012**. The Arabidopsis Information Resource (TAIR):
improved gene annotation and new tools. *Nucleic Acids Research* **40**: D1202–D1210.

Lu Q, Ding S, Reiland S, Rödiger A, Roschitzki B, Xue P, Gruissem W, Lu C, Baginsky S. **2015**.
Identification and characterization of chloroplast casein kinase II from Oryza sativa (rice). *Journal of
Experimental Botany* **66**: 175–187.

575   **Lyons E, Freeling M**. **2008**. How to usefully compare homologous plant genes and chromosomes as
576   DNA sequences. *The Plant Journal* **53**: 661–673.
577
578   **Lyons E, Pedersen B, Kane J, Alam M, Ming R, Tang H, Wang X, Bowers J, Paterson A, Lisch D,**
579   *et al.* **2008**. Finding and Comparing Syntenic Regions among Arabidopsis and the Outgroups Papaya,
580   Poplar, and Grape: CoGe with Rosids. *Plant Physiology* **148**: 1772–1781.
581
582   **Marques DN, Stolze SC, Harzen A, Nogueira ML, Batagin-Piotto KD, Piotto FA, Mason C,**
583   **Azevedo RA, Nakagami H**. **2021**. Comparative phosphoproteomic analysis of tomato genotypes with
584   contrasting cadmium tolerance. *Plant Cell Reports* **40**: 2001–2008.
585
586   **McKain MR, Tang H, McNeal JR, Ayyampalayam S, Davis JI, dePamphilis CW, Givnish TJ, Pires**
587   **JC, Stevenson DWm, Leebens-Mack JH**. **2016**. A phylogenomic assessment of ancient polyploidy and
588   genome evolution across the Poales. *Genome Biology and Evolution*: evw060.
589
590   **Nagarajan R, Gill KS**. **2018**. Evolution of Rubisco activase gene in plants. *Plant Molecular Biology* **96**:
591   69–87.
592
593   **Neuwald AF, Aravind L, Spouge JL, Koonin E V. 1999**. AAA $^{+}\square$: A Class of Chaperone-Like
594   ATPases Associated with the Assembly, Operation, and Disassembly of Protein Complexes. *Genome*
595   *Research* **9**: 27–43.
596
597   **Nguyen L-T, Schmidt HA, von Haeseler A, Minh BQ**. **2015**. IQ-TREE: A Fast and Effective
598   Stochastic Algorithm for Estimating Maximum-Likelihood Phylogenies. *Molecular Biology and*
599   *Evolution* **32**: 268–274.
600
601   **Orr DJ, Robijns AKJ, Baker CR, Niyogi KK, Carmo-Silva E**. **2023**. Dynamics of Rubisco regulation
602   by sugar phosphate derivatives and their phosphatases. *Journal of Experimental Botany* **74**: 581–590.
603
604   **Parry MAJ, Andralojc PJ, Scales JC, Salvucci ME, Carmo-Silva AE, Alonso H, Whitney SM**. **2013**.
605   Rubisco activity and regulation as targets for crop improvement. *Journal of Experimental Botany* **64**:
606   717–730.
607
608   **Paterson AH, Bowers JE, Chapman BA**. **2004**. Ancient polyploidization predating divergence of the
609   cereals, and its consequences for comparative genomics. *Proceedings of the National Academy of*
610   *Sciences* **101**: 9903–9908.
611
612   **Poolman MG, Fell DA, Thomas S**. **2000**. Modelling photosynthesis and its control. *Journal of*
613   *Experimental Botany* **51**: 319–328.
614
615   **Portis AR, Jr. 2003**. Rubisco activase - Rubisco's catalytic chaperone. *Photosynthesis Research* **75**: 11–
616   27.
617
618   **Portis AR, Li C, Wang D, Salvucci ME**. **2007**. Regulation of Rubisco activase and its interaction with
619   Rubisco. *Journal of Experimental Botany* **59**: 1597–1604.
620
621   **Qu Y, Mueller-Cajar O, Yamori W**. **2023**. Improving plant heat tolerance through modification of
622   Rubisco activase in C3 plants to secure crop yield and food security in a future warming world. *Journal of*
623   *Experimental Botany* **74**: 591–599.
624

625 **Reiland S, Messerli G, Baerenfaller K, Gerrits B, Endler A, Grossmann J, Gruissem W, Baginsky**
626 **S**. **2009**. Large-Scale Arabidopsis Phosphoproteome Profiling Reveals Novel Chloroplast Kinase
627 Substrates and Phosphorylation Networks . *Plant Physiology* **150**: 889–903.
628
629 **Ristic Z, Momčilović I, Bukovnik U, Prasad PVV, Fu J, DeRidder BP, Elthon TE, Mladenov N**.
630 **2009**. Rubisco activase and wheat productivity under heat-stress conditions. *Journal of Experimental*
631 *Botany* **60**: 4003–4014.
632
633 **Robinson SP, Portis AR**. **1989**. Adenosine triphosphate hydrolysis by purified rubisco activase. *Archives*
634 *of Biochemistry and Biophysics* **268**: 93–99.
635
636 **Rundle SJ, Zielinski RE**. **1991**. Alterations in barley ribulose-1,5-bisphosphate carboxylase/oxygenase
637 activase gene expression during development and in response to illumination. *Journal of Biological*
638 *Chemistry* **266**: 14802–14807.
639
640 **Salinas P, Fuentes D, Vidal E, Jordana X, Echeverria M, Holuigue L**. **2006**. An Extensive Survey of
641 CK2 α and β Subunits in Arabidopsis: Multiple Isoforms Exhibit Differential Subcellular Localization.
642 *Plant and Cell Physiology* **47**: 1295–1308.
643
644 **Salvucci ME, van de Loo FJ, Stecher D**. **2003**. Two isoforms of Rubisco activase in cotton, the
645 products of separate genes not alternative splicing. *Planta* **216**: 736–744.
646
647 **Salvucci ME, Portis AR, Ogren WL**. **1985**. A soluble chloroplast protein catalyzes
648 ribulosebisphosphate carboxylase/oxygenase activation in vivo. *Photosynthesis Research* **7**: 193–201.
649
650 **Salvucci ME, Portis AR, Ogren WL**. **1986**. Light and CO_2 Response of Ribulose-1,5-Bisphosphate
651 Carboxylase/Oxygenase Activation in *Arabidopsis* Leaves. *Plant Physiology* **80**: 655–659.
652
653 **Salvucci ME, Werneke JM, Ogren WL, Portis AR**. **1987**. Purification and Species Distribution of
654 Rubisco Activase. *Plant Physiology* **84**: 930–936.
655
656 **Samol I, Shapiguzov A, Ingelsson B, Fucile G, Crèvecoeur M, Vener A V., Rochaix JD,**
657 **Goldschmidt-Clermont M**. **2012**. Identification of a photosystem II phosphatase involved in light
658 acclimation in Arabidopsis. *Plant Cell* **24**.
659
660 **Sanchez de Jimenez E, Medrano L, Martinez-Barajas E**. **1995**. Rubisco Activase, a Possible New
661 Member of the Molecular Chaperone Family. *Biochemistry* **34**: 2826–2831.
662
663 **Scafaro AP, Bautsoens N, den Boer B, Van Rie J, Gallé A**. **2019**. A Conserved Sequence from Heat-
664 Adapted Species Improves Rubisco Activase Thermostability in Wheat. *Plant Physiology* **181**: 43–54.
665
666 **Schweer J, Türkeri H, Link B, Link G**. **2010**. AtSIG6, a plastid sigma factor from Arabidopsis, reveals
667 functional impact of cpCK2 phosphorylation. *The Plant Journal* **62**: 192–202.
668
669 **Shivhare D, Mueller-Cajar O**. **2017**. In Vitro Characterization of Thermostable CAM Rubisco Activase
670 Reveals a Rubisco Interacting Surface Loop. *Plant Physiology* **174**: 1505–1516.
671
672 **Soleh MA, Tanaka Y, Kim SY, Huber SC, Sakoda K, Shiraiwa T**. **2017**. Identification of large
673 variation in the photosynthetic induction response among 37 soybean [Glycine max (L.) Merr.] genotypes
674 that is not correlated with steady-state photosynthetic capacity. *Photosynthesis Research* **131**: 305–315.
675

676 **Soltis DE, Smith SA, Cellinese N, Wurdack KJ, Tank DC, Brockington SF, Refulio☐Rodriguez NF,**
677 **Walker JB, Moore MJ, Carlsward BS,** *et al.* **2011**. Angiosperm phylogeny: 17 genes, 640 taxa.
678 *American Journal of Botany* **98**: 704–730.
679
680 **Somerville CR, Portis AR, Ogren WL**. **1982**. A Mutant of *Arabidopsis thaliana* Which Lacks
681 Activation of RuBP Carboxylase *In Vivo*. *Plant Physiology* **70**: 381–387.
682
683 **Stitt M, Quick WP, Schurr U, Schulze E-D, Rodermel SR, Bogorad L**. **1991**. Decreased ribulose-1,5-
684 bisphosphate carboxylase-oxygenase in transgenic tobacco transformed with☐?antisense? rbcS. *Planta*
685 **183**.
686
687 **Stitt M, Schulze D**. **1994**. Does Rubisco control the rate of photosynthesis and plant growth? An exercise
688 in molecular ecophysiology. *Plant, Cell & Environment* **17**: 465–487.
689
690 **Stotz M, Mueller-Cajar O, Ciniawsky S, Wendler P, Hartl FU, Bracher A, Hayer-Hartl M**. **2011**.
691 Structure of green-type Rubisco activase from tobacco. *Nature structural & molecular biology* **18**: 1366–
692 70.
693
694 **Tang H, Bowers JE, Wang X, Paterson AH**. **2010**. Angiosperm genome comparisons reveal early
695 polyploidy in the monocot lineage. *Proceedings of the National Academy of Sciences* **107**: 472–477.
696
697 **To K-Y, Suen D-F, Chen S-CG**. **1999**. Molecular characterization of ribulose-1,5-bisphosphate
698 carboxylase/oxygenase activase in rice leaves. *Planta* **209**: 66–76.
699
700 **Uhrig RG, Labandera A-M, Tang L-Y, Sieben NA, Goudreault M, Yeung E, Gingras A-C, Samuel**
701 **MA, Moorhead GBG**. **2017**. Activation of Mitochondrial Protein Phosphatase SLP2 by MIA40
702 Regulates Seed Germination. *Plant Physiology* **173**: 956–969.
703
704 **Uhrig RG, Moorhead GB**. **2011**. Two Ancient Bacterial-like PPP Family Phosphatases from
705 Arabidopsis Are Highly Conserved Plant Proteins That Possess Unique Properties . *Plant Physiology*
706 **157**: 1778–1792.
707
708 **Werneke JM, Chatfield JM, Ogren WL**. **1989**. Alternative mRNA splicing generates the two
709 ribulosebisphosphate carboxylase/oxygenase activase polypeptides in spinach and Arabidopsis. *The Plant*
710 *Cell* **1**: 815–825.
711
712 **White-Gloria C, Johnson JJ, Marritt K, Kataya A, Vahab A, Moorhead GB**. **2018**. Protein Kinases
713 and Phosphatases of the Plastid and Their Potential Role in Starch Metabolism. *Frontiers in Plant Science*
714 **9**.
715
716 **Wijewardene I, Shen G, Zhang H**. **2021**. Enhancing crop yield by using Rubisco activase to improve
717 photosynthesis under elevated temperatures. *Stress Biology* **1**: 2.
718
719 **Willems P, Horne A, Van Parys T, Goormachtig S, De Smet I, Botzki A, Van Breusegem F, Gevaert**
720 **K**. **2019**. The Plant <scp>PTM</scp> Viewer, a central resource for exploring plant protein
721 modifications. *The Plant Journal* **99**: 752–762.
722
723 **Willems P, Sterck L, Dard A, Huang J, De Smet I, Gevaert K, Van Breusegem F**. **2024**. The Plant
724 PTM Viewer 2.0: in-depth exploration of plant protein modification landscapes. *Journal of Experimental*
725 *Botany*.
726

727 **Yin Z, Meng F, Song H, Wang X, Xu X, Yu D**. **2010**. Expression Quantitative Trait Loci Analysis of
728 Two Genes Encoding Rubisco Activase in Soybean . *Plant Physiology* **152**: 1625–1637.
729
730 **Zhang N, Kallis RP, Ewy RG, Portis AR**. **2002**. Light modulation of Rubisco in *Arabidopsis* requires a
731 capacity for redox regulation of the larger Rubisco activase isoform. *Proceedings of the National*
732 *Academy of Sciences* **99**: 3330–3334.
733
734 **Zhang N, Portis AR**. **1999a**. Mechanism of light regulation of Rubisco: A specific role for the larger
735 Rubisco activase isoform involving reductive activation by thioredoxin-f. *Proceedings of the National*
736 *Academy of Sciences* **96**: 9438–9443.
737
738 **Zhang N, Portis AR**. **1999b**. Mechanism of light regulation of Rubisco: A specific role for the larger
739 Rubisco activase isoform involving reductive activation by thioredoxin-f. *Proceedings of the National*
740 *Academy of Sciences* **96**: 9438–9443.
741
742 **Zhu X-G, de Sturler E, Long SP**. **2007**. Optimizing the Distribution of Resources between Enzymes of
743 Carbon Metabolism Can Dramatically Increase Photosynthetic Rate: A Numerical Simulation Using an
744 Evolutionary Algorithm. *Plant Physiology* **145**: 513–526.
745

XSDDQ Phosphorylation Site Amino Acid Legend

A
D
E
F
I
M
Q
T
V

Tandem duplications

A

B

BrassicanapusA	GLAYDTSDDQQDITRGKGMVD	94
BrassicanapusB	GLAYDTSDDQQDITRGKGMVD	94
BrassicaoleraceaA	GLAYDTSDDQQDITRGKGMVD	94
BrassicaoleraceaB	GLAYDTSDDQQDITRGKGMVD	94
ArabidopsisthalianaA	GLAYDTSDDQQDITRGKGMVD	93
ArabidopsisthalianaB	GLAYDTSDDQQDITRGKGMVD	93
CamelinasativaA	GLAYDTSDDQQDITRGKGLVD	93
CamelinasativaB	GLAYDTSDDQQDITRGKGLVD	93
GlycinemaxA	GLAYDTSDDQQDITRGKGMVD	93
PhaseolusvulgarisA	GLAYDTSDDQQDITRGKGMVD	93
GlycinemaxB	GLAYDISDDQQDITRGKGLVD	99
PhaseolusvulgarisB	GLAYDVSDDQQDITRGKGLVD	97
MedicagotruncatulaA	GLAYDTSDDQQDITRGKGMVD	94
MedicagotruncatulaB	GLAYDISDDQQDITRGKGMVD	94

+	+	−	−	+	+	−	−	Temperature: 28°C
−	−	+	+	−	−	+	+	Temperature: 42°C
+	−	+	−	+	−	+	−	Light
−	+	−	+	−	+	−	+	Dark
+	+	+	+	−	−	−	−	Water
−	−	−	−	+	+	+	+	Protein phosphatase inhibitor

α
β

Anti-Rca

α
β

Anti-pT78

CBB