Phylogenetic diversity of light dependent phosphorylation of Thr78 in Rca

Nikita Bhatnagar[a], 0000-0001-7832-1347

Sarah S. Chung[a,b], sohyeon2@illinois.edu, 0009-0005-6096-5343

John Hodge[a,b], jghodge@illinois.edu, 0000-0003-0793-1871

Ryan A Boyd[a,b], raboyd2@illinois.edu, 0000-0003-4009-7700

Sang Yeol Kim[b]

Mia Sands[b], xin7@illinois.edu, 0000-0002-4925-9772

Andrew D. B. Leakey[a,b,c], leakey@illinois.edu, 0000-0001-6251-024X

Donald R. Ort[a,b,c], d-ort@illinois.edu, 0000-0002-5435-4387

Steven Burgess[a,b], sjb287@illinois.edu, 0000-0003-2353-7794

Affiliations

[a]Carl R. Woese Institute for Genomic Biology, University of Illinois Urbana-Champaign

[b]Department of Plant Biology, University of Illinois Urbana-Champaign

[c]Department of Crop Sciences, University of Illinois Urbana-Champaign

Corresponding Author: Steven J. Burgess

The author responsible for distribution of materials integral to the findings presented in this article in

accordance with the policy described in the Instructions for Authors

(https://academic.oup.com/plphys/pages/General-Instructions) is (insert author name).

Running head:

Role of phosphorylation/dephosphorylation the regulation of of Rubisco activase

SUMMARY

Rubisco activase is an ATP-dependent chaperone that facilitates dissociation of inhibitory sugar

phosphates from the catalytic sites of ribulose-1,5-bisphosphate carboxylase/oxygenase during

photosynthesis. In Arabidopsis, Rubisco activase is negatively regulated by dark-dependent

phosphorylation of threonine 78. The prevalence of threonine 78 in Rubisco activase was investigated

across sequences from 91 plant species, finding 29 (~32%) species shared a threonine in the same

position. Analysis of seven C3 species with an antibody raised against a threonine 78 phospho-peptide

demonstrated that this position is phosphorylated in multiple genera. However, light-dependent

dephosphorylation of threonine 78 was observed only in Arabidopsis. Further, phosphorylation of

threonine 78 could not be detected in any of the four C4 grass species examined. The results suggest that

despite conservation of threonine 78 in Rubisco activase from a wide range of species, a regulatory role
for phosphorylation at this site is more limited. This provides a case study for how variation in post-
translational regulation can amplify functional divergence across the phylogeny of plants beyond what is
explained by sequence variation in a metabolically important protein.
39
40

INTRODUCTION

Rubisco catalyzes the addition of CO_2 to a molecule of ribulose-1,5-bisphosphate (RuBP) forming two molecules of 3-phosphoglycerate (3-PGA) and in many circumstances is the rate limiting step of CO_2 assimilation (Stitt *et al.*, 1991; Hudson *et al.*, 1992; Stitt & Schulze, 1994; Poolman *et al.*, 2000; Zhu *et al.*, 2007). To perform this reaction the enzyme must first be activated through attachment of CO_2 to a catalytic lysine in order to bind Mg^{2+} (Lorimer and Miziorko, 1980). However, the enzyme is susceptible to inhibition by attachment of RuBP prior to activation, or binding of one of several inhibitory sugar phosphates to the active site (Orr et al., 2023). Inhibitors can be released by the action of Rubisco activase (Rca) (Somerville et al., 1982; Salvucci et al., 1986; Jiang et al., 1994), which uses energy from ATP hydrolysis to drive conformational changes in Rubisco, freeing the active site (Neuwald et al., 1999; Portis & Jr., 2003; Flecken et al., 2020). Therefore, Rca is critical for regulation of Rubisco activity and carbon assimilation (Parry et al., 2013; Wijewardene et al., 2021; Qu et al., 2023; Orr et al., 2023).

Rca itself is subject to regulation, which provides a link between carbon assimilation and electron transport. Most of our understanding about this process comes from Arabidopsis, where there are two isoforms of Rca which differ in size. The longer 'α' isoform possesses a 20-30 amino acid C-terminal extension (To et al., 1999) which contains two cysteine residues (Cys-451 and Cys-470). The cysteines are regulated by thioredoxin f in response to changes in the ATP/ADP arising from variation in rates of electron transport and ATP synthesis (Zhang & Portis, 1999b; Zhang et al., 2002). In the dark, the cysteines are oxidized to form a disulfide bridge (oxidized state, E_m ~344 mV pH 7.9) (Hutchison et al., 2000), increasing the sensitivity of Rca to inhibition by ADP (Portis et al., 2007), downregulating Rca activity and CO_2 fixation. However, redox regulation of Rca does not appear to be essential. Solanaceae species, such as tobacco, are reported to only express Rca-β, which lacks the C-terminal extension, (Nagarajan & Gill, 2018), and the sensitivity of Rca-β isoforms to inhibition by ADP varies (Carmo-Silva & Salvucci, 2013). Therefore, it is apparent that additional regulatory mechanisms are also important to control Rca activity.

Rca α and β isoforms, which differ in ADP sensitivity, can be produced by both alternate splicing (Werneke et al., 1989; Rundle & Zielinski, 1991; To et al., 1999) or from transcription of independent genes (Rundle & Zielinski, 1991; Salvucci et al., 2003). In addition, the relative expression level and catalytic properties of Rca-α and Rca-β differ across species (Rundle & Zielinski, 1991; Crafts-Brandner et al., 1997; Carmo-Silva & Salvucci, 2013; Shivhare & Mueller-Cajar, 2017; Degen et al., 2021) and can be impacted by environmental factors such as temperature (Shivhare & Mueller-Cajar, 2017; Scafaro et al., 2019; Kim et al., 2020, 2021; Degen et al., 2021) to modulate CO_2 fixation under variable conditions.

Post-translational modification of Rca has also been proposed to provide an additional layer of regulation (Amaral et al., 2024). In Arabidopsis, Threonine 78 (Thr78) is phosphorylated in the dark and

under low light intensities, but not under high light, suggesting a regulatory role (Reiland *et al.*, 2009;
Boex☐Fontvieille *et al.*, 2014; Kim *et al.*, 2016). Functional support for this hypothesis came from
several findings: (1) phosphorylation at Thr78 (Thr78-Rca) in Arabidopsis is reported to cause a
significant decrease in plant biomass, due to a decreased Rubisco activation state (Kim *et al.*, 2019); (2)
transgenic Arabidopsis plants expressing phospho-null mutants of Rca (T78A) had enhanced
photosynthetic efficiency and plant growth; and (3) substitution of Thr78-to-Ser (T78S) resulted in hyper-
phosphorylation, leading to reduced carbon assimilation and slower growth (Kim *et al.*, 2019). cpCK2
was shown to be the major kinase responsible for dark phosphorylation of AtRca, with loss-of-function
mutant lines showing a reduction in Rca phosphorylation (Kim *et al.*, 2016). However, it is not clear
whether this proposed mechanism is relevant outside of Arabidopsis. To address this issues, we sought to
define the prevalence of this post-translational modification in photosynthetic organisms. This was
achieved by considering the following four questions, (1) at what stage of flowering plant evolution did
Thr78 emerge? (2) how conserved is Thr78 among photosynthetic angiosperms? (3) to what extent is
Thr78 phosphorylated in different species? (4) and is light-dependent de-phosphorylation of Thr78 a
general mechanism of controlling of RCA activity?
90
MATERIAL AND METHODS
Determination of Rca isoform and photosynthetic species containing Thr78-Rca
To analyze the conservation of Thr78 in Rca protein across plant species, the sequence of
Arabidopsis Rca (AtRca: AT2G39730) was retrieved from the *Arabidopsis thaliana* TAIR10 genome
assembly (Lamesch *et al.*, 2012) on Phytozome v13 (Goodstein *et al.*, 2012). Syntelogs to AtRca (i.e.,
shared positional homology along chromosomes) were subsequently identified in monocots (*Setaria*
viridis Rca: Sevir.8G261433) and asterids (*Mimulus guttatus* Rca: Migut.F00758), which have homology
to this sequence using either synteny mapping features on CoGe (Lyons & Freeling, 2008; Lyons *et al.*,
2008). Setaria Rca was chosen as an example of a gene that has been previously characterized (Kim et al.,
2021) while Mimulus shares comparable synteny to Arabidopsis in the genomic region surrounding Rca
enabling identification of syntelogs in the asterids. Working from these three representatives, additional
syntelogues were locally identified based on their closest known Rca sister lineage within the Angiosperm
phylogeny across other accessible plant genomes on these databases. *Amborella trichopoda* was
leveraged as an outgroup for flowering plants in this case study given that this is known to be the one of
the earliest diverging lineages for flowering plants (Soltis *et al.*, 2011). Additionally, homologues were
surveyed, which contained threonine at position 78 phosphorylation site even if they didn't precisely fit
this stringent criterion of synteny using pBLAST on NCBI (https://blast.ncbi.nlm.nih.gov/Blast.cgi) for
key agronomic model systems. This ultimately resulted in 217 isoform CDS sequences being identified.

Alternative splicing of the same locus in some lineages meant that these CDS sequences represented 153

unique genes from across 91 species and 50 plant families. The longer, Rca-α, sequence was

preferentially retained as a taxonomic representative for each alternative splicing gene when available for

phylogenetic analysis. The CDS sequence of these genes were visualized in Seaview (Gouy *et al.*, 2010)

and codon-aligned using MUSCLE (Edgar, 2004). Obvious pseudogenes that were missing large portions

of their AAA+ ATPase domain such as the rice sequence (LOC_Os11g47980) were removed prior to

phylogenetic analysis of this DNA alignment. A maximum likelihood tree was subsequently assembled

using IQTree (Nguyen *et al.*, 2015) with the Amborella Rca sequence being specified as an outgroup and

1000 bootstraps being run to generate statistical support for nodes along the tree with the automatic

substitution model selection determining a SYM+R8 model was the best-fit according to Bayesian

Information Criterion. The resulting treefile was then visualized either in Figtree

(http://tree.bio.ed.ac.uk/software/figtree/) or in R using the Phytools and Ape libraries (Paradis & Schliep,

2019; Revell, 2024). Ancestral state reconstructions were performed using the Ape function *ace*

particularly for the original position 78 amino acid identity between syntelogous genes, and the manner of

α/ß origin (either by alternative splicing or distinct genes) for each plant family (Figure S3).

Plant growth and stress treatment

Arabidopsis Col-0 and camelina seeds were sterilized in 1 % (v/v) bleach for 5 min, rinsed with 70 %

EtOH, and dried on filter paper. The sterilized seeds were sown on MS (Murashige and Skoog basal salt

mixture powder, Phyto Technology Laboratories, M524) agar (0.8 %, w/v, Sigma-Aldrich, A1296) with 1

% (w/v) sucrose. The plates were stored at 4 °C for 2 days for cold stratification before transfer to a

growth chamber. Seedlings were transferred into pots (5 cm diameter) filled with soil (Berger; PRO-MIX

BX All Purpose Growing Mix) before the third true leaf emerged, which was approximately 5 to 7 days

after germination (DAG). The other plants (brassica, soybean, green bean, and Medicago) were directly

sown in soil after sterilization. All plant species were grown at 22 °C, 16 h light and 8 h dark, 200 µmol

m^{-2} s^{-1} PPFD with 2100 K 400W high pressure sodium bulbs (Ceramalux ALTO; C400S51).

Protein extraction and immunoblotting assay

Total proteins were extracted from three frozen leaf discs (1 cm in diameter) using a Qiagen

Tissuelyzer for tissue grinding. The ground tissue was suspended in 0.5 mL of extraction buffer

containing 62.5 mM Tris-HCl (pH 6.8), 2% SDS, 10% glycerol, 0.35 M 2-mercaptoethanol, and 0.001%

protease inhibitor cocktail (Sigma-Aldrich; P9599). The suspension was heated at 95°C for 5 minutes to

denature proteins and inactivate proteases, followed by centrifugation at 13,500 rpm for 30 minutes at

4°C. The supernatant was collected, and 10 µL aliquots were separated on 10% SDS-PAGE gels. Proteins

from the gels were transferred electrophoretically to 0.2 µm nitrocellulose membranes (Bio-Rad) and

blocked with PBS solution with 5% (w/v) non-fat milk powder (Sigma) for 1h at room temperature. For

immunodetection, primary antibodies were diluted in Intercept Blocking Buffer (LI-COR; 927-70001);

anti-Rca (Agrisera; AS10 700) at 1:5000 and anti-pT78 (GenScript) at 1:4000, and incubated overnight at

4°C. Modification-specific antibody was generated against the phosphopeptide antigen

RGLAYDpT78DDQQDC (Arabidopsis anti-pT78). For each peptide antigen, the Cys residue at the N- or

C-terminus was added for coupling to keyhole limpet hemocyanin (KLH). All custom antibodies were

produced by GenScript (Piscataway, NJ, USA). Secondary detection used a fluorescent goat anti-rabbit

antibody (Invitrogen; A32735) at a 1:10,000 dilution in PBS-T, incubated for 1 h at room temperature,

and visualized with a LI-COR Odyssey Infrared Imaging System. All immunoblotting experiments were

conducted at least twice to ensure reproducibility of results.

Protein phosphorylation assay with phosphatase inhibitor

WT Col-0 plants were grown as described above and whole rosettes were sampled and dipped in

the protein phosphatase inhibitor buffer: 1% (v/v) solution of phosphatase inhibitor cocktail 2 (Sigma:

P5726) and phosphatase inhibitor cocktail 3 (Sigma: P0044), overnight under dark. For dark treatment

samples, the protein extraction was done using the overnight incubated samples and for light treatment

samples the rosettes were exposed to 150 µmol m^{-2} s^{-1} PPFD light for 2 h and then total protein was

extracted.

RESULTS

Thr 78 in Rca is common

To study the conservation of Thr78 (Arabidopsis numbering) in Rca, 1893 protein sequences

from 479 plants species were assessed. Among these, 203 species retained Thr78 (Table S1). Next, a

more stringent synteny informed ortholog phylogeny of Rca-α and Rca-β isoforms was generated across

angiosperms to address the questions: at what stage of flowering plant evolution did the putative Thr78

phosphosite emerge, and how well conserved is it among photosynthetic angiosperms? Leveraging

maximum likelihood gene family phylogeny ancestral states, Thr78 distribution was estimated for each

family, including deeper nodes within the tree such as the branch point of the *Amborellaceae* (Fig. 1). The

presence of Thr78 was observed in most plant families (Fig. 1), while among the *Poaceae*, that includes

major crop species, the presence of Isoleucine (Ile) was found to dominate at position 78 compared to

Threonine (Fig. 1). The difference in the abundance of both amino acids, Thr and Ile at this site is similar

at the evolutionary scale, and an estimate of the ancestral state was ambiguous. Interestingly, the *α* and *β*

copies appeared polyphyletic, with multiple origins of the two isoforms arising from alternate splicing or

gene duplication across the phylogeny (Fig. S1-S3). Within the Brassicaceae *Arabidopsis thaliana* and its

immediate relatives all appear to have a single copy of Rca with alpha and beta forms being the result of

alternative splicing of a single transcript.

By contrast in the Fabaceae, *Glycine max* has four gene copies of Rca (two each of α and β) with

the Papillionoid legumes more generally having two copies of Rca. There is no equivalent duplication

event in *Cercis canadensis* that would indicate a Fabaceae-wide pattern. Therefore, this pattern of two

Papillionoid copies is consistent with the whole-genome doubling (WGD) event preceding the

diversification of this subfamily, and by undergoing an additional allopolyploid event, *Glycine max* has

two additional copies for each of these ancestral homeologues (Gill, *et al.*, 2009; Koenen, *et al.*, 2021).

Within Poaceae, *Setaria viridis* and the grasses more generally displayed two mirrored clades, which

reflect the known taxonomic sorting of this family. The grass family has undergone three WGD events (τ,

σ, and ρ), two of which are shared with monocots, with the p WGD unique to the Poaceae (Paterson *et al.*,

2004; Tang *et al.*, 2010; McKain *et al.*, 2016). Although superficially resembling the ρ WGD event, these

grass copies are physically proximal to their sister copies in the adjacent clade suggesting that this may

instead be an ancient tandem duplication event.

Light-dependent dephosphorylation of Thr78 in Rca is unique to Arabidopsis

Previous reports showed that both AtRca isoforms phosphorylate in the dark and are

dephosphorylated as plants are exposed to light. To test whether Thr78 phosphorylation status is light-

sensitive in species other than Arabidopsis, we screened three C3 plants from the *Brassicaceae* for dark-

dependent phosphorylation of Thr78, using a pT78 antibody (Kim et al., 2019) (Fig. 2A). As in

Arabidopsis, Rca-α and Rca-β isoforms arise from alternate splicing in *Brassicaceae* and the Thr78 site is

conserved in all proteins (Fig. 2B). However, unlike Arabidopsis, Rca in *Camelina sativa* (*Cs*), *Brassica*

oleraceae (*Bo*) and *Brassica napus* (*Bn*) phosphorylated in dark but lacked light-induced

dephosphorylation (Fig. 3A). These data suggest that although Thr78 is phosphorylated in multiple

species, light-dependent regulation may be specific to Arabidopsis.

To broaden the analysis, we performed the dephosphorylation assay on several legumes: *Glycine*

max (*Gm*), *Phaseolus vulgaris* (*Pv*), and *Medicago truncatula* (*Mt*). The Rca isoforms in these species are

derived from individual genes of *Rca-α* and *Rca-β*, unlike the situation in the *Brassicaceae*. The Rca-α

isoforms retain Thr78, whereas the β isoforms have Ile78 in *Gm* and *Mt,* and Val78 in *Pv* (Fig. 2B).

Publicly available data suggest the transcription level of Rca-α is low relative to β in both *Gm* and *Pv,*

and moderately expressed in *Mt* (Table S2). The light/dark assay on legume plants showed that both the

isoforms were expressed, and phosphorylated in the dark, but as with the Brassicaceae, did not

dephosphorylate in the light (Fig. 2A). The reason why the antibody was able to detect phosphorylation of

β forms, which lack Thr78, is unclear. It is possible that there is relaxed selectively, but we consider these
results with caution.

To study the impact of phosphorylation in C4 grass plants, we analyzed Rca phosphorylation in
four species of C4 grass: *Sorghum bicolor* (*Sb*), *Miscanthus* (*Ms*), *Saccharum officinarum* (*Su*), and
Setaria viridis (*Sv*) (Fig. 3A). Each of these species possess two separate genes for the alternative Rca
isoforms, with Rca-α containing Thr78, but with it lacking from Rca-β (Fig. 3B, C). In accordance with
previous reports (Kim *et al.*, 2021), only the Rca-β isoform could be detected under normal growth
conditions (Fig. 3A and C). An unspecific cross-reaction was present in C4 grass samples probed with
pT78 antibody, but the detected protein was too large to be Rca-β, subsequently no phosphorylation of
Rca-β was detected (Fig. 3A). To test whether a C4 Rca-α isoform could be phosphorylated, sorghum
plants were exposed to heat induce Rca-α isoform expression (Fig. 3C). Plants accumulated SbRca-α
after 24 h heat treatment, but no bands of the correct size could be detected using the pT78 antibody (Fig.
3C). To rule out the possibility that a lack of detectable Thr78 phosphorylation was due to high protein
phosphatase activity, the assay was repeated in the presence of a protein phosphatase inhibitor cocktail
(PPIC), but again protein phosphorylation was not observed under any condition (Fig. 4). This result
suggests that SbRca-α T78 is not phosphorylated by cpCK2 as it is in Arabidopsis even though the gene
for cpCK2 is present in sorghum.

DISCUSSION

Rca is subject to multiple levels of regulation to adjust carbon fixation in response to
environmental conditions (Amaral *et al.*, 2024), and differences in gene expression, biochemical
properties of isoforms, redox status, and post-translational modification of Rca have been topics of
investigation (Boex☐Fontvieille *et al.*, 2014; Kim *et al.*, 2016, 2019). Dark dependent phosphorylation of
Thr78 is proposed to downregulate Rca activity leading to reduced assimilation (Kim et al. 2019). cpCK2
is described as the major kinase responsible for phosphorylation of AtRca Thr78 based on dramatically
reduced phosphorylation in *cpcpk2* T-DNA insertion lines (Kim *et al.*, 2016). Substitution of Thr78 with
Serine lead to a relative increase in phosphorylation at the site, in Arabidopsis resulted in smaller plants
with decreased photosynthesis (Kim *et al.*, 2019). Altering a single amino acid, Thr78, was proposed as a
means of generating plants with better photosynthesis. However, the prevalence of Thr78 across species,
the frequency of post-translational modification and functionality were unclear. To begin to address these
questions we expanded previous searches of genomic databases, which assessed the evolution of Rca
copy number across the green lineage (Nagarajan & Gill, 2018; Kim *et al.*, 2019), to use a synteny based
approach. Nagarajan and Gill (2018) placed the emergence of the Rca-α CTE and alternate splicing

mechanisms in the charophytes, before the divergence of land plants and there is only one reported
example of a species only having evidence for expression of Rca α (Nagarajan & Gill, 2018). Here,
despite α and β forms (bearing or lacking a c-terminus respectively) occurring within each model system,
these α and β forms appear to represent repeated cases of evolutionary convergence on these two
isoforms (Fig S5) through distinct means within each clade: subfunctionalization (Birchler & Yang,
2022), following tandem duplication in the grasses as previously described (Nagarajan & Gill, 2018),
WGD in the legumes, or alternative splicing of a single coding sequence into alternative isoforms in
Brassicaceae. This may suggest that Rca isoforms can be quite plastic in the way they arise within distinct
plant families where α and β forms may not necessarily reflect true homology within Rubisco activase
isoforms.

Previous assessment of a smaller number of Rca sequences failed to identify any with Thr78
replaced by Serine, leading to speculation that the configuration with the attendant hyper phosphorylation
observed in Arabidopsis has been selected against (Kim *et al.*, 2019). In the initial set of Rca sequences
identified by BLAST search, we identified gene models from at least seven species (Table S1) which do
have Ser78, although these were not considered orthologs in the synteny analysis. Of the 25 species found
to contain Rca sequences with Thr78, it remains to be seen how many undergo phosphorylation. Our data
are consistent with it occurring in at least the Brassicaceae, while phosphorylation of an equivalent Thr
residue has been reported in tomato, suggesting this modification may occur more widely (Marques et al.,
2021).

We found no evidence for phosphorylation of Thr78 in monocot species, including Sorghum,
Setaria, sugarcane and Miscanthus. The peptide fragment used for antibody generation is highly
conserved, with 100% identity between Arabidopsis Rca and Sorghum Rca-α "RGLAYDTSDDQQDC"
and should therefore have been able to detect phosphorylation if present under the conditions tested.
There are several potential explanations for this finding, including an absence of the kinase, differences in
localization or expression profiles of the kinase, or regulation of the kinase. In Arabidopsis, Rca is
phosphorylated by cpCK2 (Kim et al., 2016). A homologous gene has been characterized in rice
(LOC_Os03g55490) (Lu et al., 2015) and syntenic sequences can be found in C4 grasses including
Sorghum (Sobic.001G07970) (Carvalho et al., 2020). As Sorghum operates two-celled C4 photosynthesis
for an interaction to occur between Rca and cpCK2 both must be expressed in the same cell type.
Transcriptomics suggests that cpCK2 is expressed in both mesophyll and bundle sheath cells in maize
(Chang et al., 2012), Setaria (John et al., 2014) and Sorghum (Döring et al., 2016), albeit with ~2-fold
mesophyll enrichment in maize and Setaria whereas Rca is localized in the bundle sheath. Even so, it
appears that the orthologous cpCK2 is present in the same compartment as Rca (i.e., the bundle sheath
cell chloroplast). However, in the monocot species tested Rca-β, which lacks Thr78, is the predominate

isoform under normal conditions and Rca-α, which possess Thr78, is induced during stress (Shivhare &

Mueller-Cajar, 2017; Scafaro et al., 2019; Kim et al., 2020, 2021). Whether C4 monocot cpCK2 retains

the ability to phosphorylate Thr78 in Rca-α, or if the bundle sheath stromal environment is permissive for

phosphorylation remains unclear (Kim et al., 2016). CK2 exists as a multigene family, composed of

catalytic α subunits and regulatory beta subunits (Salinas et al., 2006; Schweer et al., 2010), which form

hetero-tetramers that can differ in substrate specificity (Domańska et al., 2005). Intriguingly there appears

to be a lack of regulatory β subunits in the chloroplast (Salinas et al., 2006), so differences in regulatory

subunits impacting specificity appears unlikely, although multiple chloroplast CK2 containing complexes

have been detected (Baginsky et al., 1997) suggesting the protein does not act alone. However, further

research is required to characterize the regulation of monocot cpCK2 and whether it can phosphorylate

Rca.

Conservation of phosphorylation of Thr78 in some C3 plants is not evidence of a conserved

regulatory role. Indeed, in the species examined, strong evidence for light-dependent dephosphorylation

of Thr78 was only found for Arabidopsis, with phosphorylation of Rca detected in both the light and dark

in other C3 species (Fig. 3). These data may suggest a limited scope for phosphorylation as a means of

regulation, with most species depending on other mechanisms, such as changes in redox state, catalytic

properties or expression. It may be that alternate post-translational modifications of Rca have greater

importance in other species and several different ones have been described (Amaral et al., 2024).

Taken together our results suggest that Thr78 can be found in Rca sequences across the green

lineage, although the prevalence varies, and degree to which it undergoes phosphorylation is inconsistent.

It is likely that regulation of Rca can occur through multiple mechanisms, including redox, sensitivity to

ADP and gene expression and the relative importance of each varies.

Acknowledgments

We would like to thank Toby Kellogg and Shawn Abrahams for providing feedback on the phylogenic

analyses. This work was funded by the DOE Center for Advanced Bioenergy and Bioproducts Innovation

(U.S. Department of Energy, Office of Science, Biological and Environmental Research Program under

Award Number DE-SC0018420). Any opinions, findings, and conclusions or recommendations expressed

in this publication are those of the author(s) and do not necessarily reflect the views of the U.S.

Department of Energy.

Competing interests

The authors report no conflict of interest

Authors contributions

Conceptualization (SK, NB, SJB, DO), Investigation (NB, SC, MS, JH, SK), Formal Analysis (NB),

Project administration (NB, SJB, DO), Supervision (AL, SJB, DO), Visualization (NB), Writing - original

draft (NB, JH, SC, DO, SJB), Writing – review and editing (NB, JH, DO, AL, SJB).

FIGURE LEGENDS

Figure 1: Phylogenetic representation of the amino acid present at the phosphorylation site (peptide position 78) within extant Rca gene models. Ancestral states for families, and when available deeper orders of Angiosperms, are denoted at their corresponding nodes as pie charts. The nodes depicting the presence of Thr78 is colored green and the ones with Ile78 is colored navy blue. Less abundant amino acids at the peptide position 78 are depicted as varying shades of blue and purple. The detailed confirmation of bootstrap values is mentioned in Figure S3.

Figure 2: Light dependent dephosphorylation of Thr78 in Rca is specific to Arabidopsis. (A) Immunoblot analysis comparing protein phosphorylation and total abundance of Rca under light (L) and dark (D) conditions (At: *Arabidopsis thaliana*, Cs: *Camelina sativa*, Bn: *Brassica napus*, Bo: *Brassica oleraceae*, Gm: *Glycine max*, Pv: *Phaseolus vulgaris*, Mt: *Medicago truncatula*). The upper panel shows the total Rca protein extracted from plant samples and the middle panel shows the phosphorylated Rca protein. α and β denote the 2 isoforms of Rca protein. CBB (Coomassie Brilliant Blue) depicts the loading control. (B) Sequence alignment depicting the conservation of Thr78 amino acid in C3 species. The amino acid lacking color codes depicts the change in the Thr78 site to Isoleucine (*I*) in *Glycine max* and *Medicago truncatula*, and Valine (*V*) in Phaseolus vulgaris. The letter "A" and "B" at the end of the species labelling denotes the Rca-α and Rca-β isoforms. (C) The phylogenetic tree depicts the division of C3 Brassicaceae family and legumes based on Rca isoforms arising from protein splicing in Brassicaceae as compared to different genes in legumes and presence and absence of Thr78 amino acid in the Rca-β isoform. The blue boxes depict the presence of Thr78 amino acid whereas the red depicts the lack of Thr78. The yellow box depicts the experimental confirmation of protein phosphorylation at Thr78 phosphosite.

Figure 3: Phosphorylation of Rca is not observed in selected C4 moncots. (A) Protein phosphorylation assay depicting four C4 plant samples in light and dark conditions (At: *Arabidopsis thaliana*, Sv: *Setaria viridis*, Sb: *Sorghum bicolor*, Su: *Saccharum officinarum*, Ms: *Miscanthus*). The upper panel shows the accumulation of Rca protein quantified with Anti-Rca antibody and middle panel shows the phosphorylated Rca protein quantified with Anti-pT78 antibody. α and β denote the 2 isoforms of Rca protein. CBB denotes the loading control. (B) Sequence alignment of motif comparing Arabidopsis and select C4 plants. (C) Immunoblot analysis comparing Rca phosphorylation under heat for 24 h in WT sorghum. The numbers denote the time in hours after the heat stress. CBB denotes the loading control.

Figure 4: Sorghum lacks phosphorylation of Thr78 with or without PPIC treatment. (A) Immunoblot analysis comparing sorghum WT samples collected after heat stress at 42^{o}C under light and dark treatment. The control samples were collected at 28^{o}C. The + and – sign denotes presence and absence of

treatment, respectively. The left panel shows Rca protein accumulation and right panel depicts analysis

using anti-pT78 antibody.

Figure S1: The conservative phylogeny structure of Rca orthologs where all nodes with bootstrap support

(blue numbers) greater than 80% being retained whereas other lower support nodes were collapsed into

polytomies.

Figure S2: Rca α (white) and β (black) isoforms on extant gene models as well as alternatively spliced

gene models which produce both Rca α and β (grey). Ancestral states for families, and when available

deeper orders of Angiosperms, are denoted at their corresponding nodes as pie charts.

Figure S3: The likely pattern that could result in the polytomies of α and β isoforms ancestry states that

Angiosperms initially beginning as Rca α lineages out of which β isoforms repeatedly arise

independently following either subfunctionalization of duplicate gene copies or alternative splicing of α

gene models to β isoforms as well. Within certain lineages where α isoforms are lost all the Rca copies

become canalized into a β isoform state thereafter.

Table S1: List of Rca protein sequences identified by BLAST search.

Table S2: Relative expression and residue at site equivalent to AtRca T78 for Rca isoforms from *G. max*,

M. truncatula and *P. vulgaris.*

REFERENCES

Alonso JM, Stepanova AN, Leisse TJ, Kim CJ, Chen H, Shinn P, Stevenson DK, Zimmerman J, Barajas P, Cheuk R, *et al.* 2003. Genome-wide insertional mutagenesis of Arabidopsis thaliana. *Science (New York, N.Y.)* **301**: 653–7.

Amaral J, Lobo AKM, Carmo☐Silva E. 2024. Regulation of Rubisco activity in crops. *New Phytologist* **241**: 35–51.

Baginsky S, Tiller K, Link G. 1997. Transcription factor phosphorylation by a protein kinase associated with chloroplast RNA polymerase from mustard (Sinapis alba). *Plant Molecular Biology* **34**: 181–189.

Birchler JA, Yang H. 2022. The multiple fates of gene duplications: Deletion, hypofunctionalization, subfunctionalization, neofunctionalization, dosage balance constraints, and neutral variation. *The Plant Cell* **34**: 2466–2474.

Boex☐Fontvieille E, Daventure M, Jossier M, Hodges M, Zivy M, Tcherkez G. 2014. Phosphorylation pattern of <scp>R</scp>ubisco activase in *<scp>A</scp> rabidopsis* leaves. *Plant Biology* **16**: 550–557.

Carmo-Silva AE, Salvucci ME. 2013. The Regulatory Properties of Rubisco Activase Differ among Species and Affect Photosynthetic Induction during Light Transitions . *Plant Physiology* **161**: 1645–1655.

Carvalho DS, Nishimwe A V., Schnable JC. 2020. IsoSeq transcriptome assembly of C$_3$ panicoid grasses provides tools to study evolutionary change in the Panicoideae. *Plant Direct* **4**.

Chang Y-M, Liu W-Y, Shih AC-C, Shen M-N, Lu C-H, Lu M-YJ, Yang H-W, Wang T-Y, Chen SC-C, Chen SM, *et al.* 2012. Characterizing Regulatory and Functional Differentiation between Maize Mesophyll and Bundle Sheath Cells by Transcriptomic Analysis . *Plant Physiology* **160**: 165–177.

Crafts-Brandner SJ, van de Loo FJ, Salvucci ME. 1997. The Two Forms of Ribulose-1,5-Bisphosphate Carboxylase/Oxygenase Activase Differ in Sensitivity to Elevated Temperature. *Plant Physiology* **114**: 439–444.

Crafts-Brandner SJ, Salvucci ME. 2002. Sensitivity of Photosynthesis in a C4 Plant, Maize, to Heat Stress. *Plant Physiology* **129**: 1773–1780.

Degen GE, Orr DJ, Carmo☐Silva E. 2021. Heat☐induced changes in the abundance of wheat Rubisco activase isoforms. *New Phytologist* **229**: 1298–1311.

Depuydt T, Vandepoele K. 2021. Multi☐omics network☐based functional annotation of unknown Arabidopsis genes. *The Plant Journal* **108**: 1193–1212.

Domańska K, Zieliński R, Kubiński K, Sajnaga E, Masłyk M, Bretner M, Szyszka R. 2005. Different properties of four molecular forms of protein kinase CK2 from Saccharomyces cerevisiae. *Acta biochimica Polonica* **52**: 947–51.

Döring F, Streubel M, Bräutigam A, Gowik U. 2016. Most photorespiratory genes are preferentially expressed in the bundle sheath cells of the C$_4$ grass *Sorghum bicolor*. *Journal of Experimental Botany* **67**: 3053–3064.

418
419 **Flecken M, Wang H, Popilka L, Hartl FU, Bracher A, Hayer-Hartl M**. **2020**. Dual Functions of a
420 Rubisco Activase in Metabolic Repair and Recruitment to Carboxysomes. *Cell* **183**: 457-473.e20.
421
422 **Gao Z, Yang X, Mei Y, Zhang J, Chao Q, Wang B**. **2023**. A dynamic phosphoproteomic analysis
423 provides insight into the <scp>C4</scp> plant maize (*Zea mays* L.) response to natural diurnal changes.
424 *The Plant Journal* **113**: 291–307.
425
426 **Goodstein DM, Shu S, Howson R, Neupane R, Hayes RD, Fazo J, Mitros T, Dirks W, Hellsten U,**
427 **Putnam N,** *et al.* **2012**. Phytozome: a comparative platform for green plant genomics. *Nucleic Acids*
428 *Research* **40**: D1178–D1186.
429
430 **Gouy M, Guindon S, Gascuel O**. **2010**. SeaView Version 4: A Multiplatform Graphical User Interface
431 for Sequence Alignment and Phylogenetic Tree Building. *Molecular Biology and Evolution* **27**: 221–224.
432
433 **Hudson GS, Evans JR, von Caemmerer S, Arvidsson YBC, Andrews TJ**. **1992**. Reduction of
434 Ribulose-1,5-Bisphosphate Carboxylase/Oxygenase Content by Antisense RNA Reduces Photosynthesis
435 in Transgenic Tobacco Plants. *Plant Physiology* **98**: 294–302.
436
437 **Jiang C, Quick WP, Alred R, Kliebenstein D, Rodermel SR**. **1994**. Antisense RNA inhibition of
438 Rubisco activase expression. *The Plant Journal* **5**: 787–798.
439
440 **John CR, Smith-Unna RD, Woodfield H, Covshoff S, Hibberd JM**. **2014**. Evolutionary Convergence
441 of Cell-Specific Gene Expression in Independent Lineages of C4 Grasses   . *Plant Physiology* **165**: 62–
442 75.
443
444 **Kim SY, Bender KW, Walker BJ, Zielinski RE, Spalding MH, Ort DR, Huber SC**. **2016**. The plastid
445 casein kinase 2 phosphorylates rubisco activase at the Thr-78 site but is not essential for regulation of
446 rubisco activation state. *Frontiers in Plant Science* **7**.
447
448 **Kim SY, Harvey CM, Giese J, Lassowskat I, Singh V, Singh V, Cavanagh AP, Cavanagh AP,**
449 **Spalding MH, Finkemeier I,** *et al.* **2019**. In vivo evidence for a regulatory role of phosphorylation of
450 Arabidopsis Rubisco activase at the Thr78 site. *Proceedings of the National Academy of Sciences of the*
451 *United States of America*.
452
453 **Kim SY, Slattery RA, Ort DR**. **2021**. A role for differential Rubisco activase isoform expression in C $_4$
454 bioenergy grasses at high temperature. *GCB Bioenergy* **13**: 211–223.
455
456 **Kim SY, Stessman DJ, Wright DA, Spalding MH, Huber SC, Ort DR**. **2020**. Arabidopsis plants
457 expressing only the redox-regulated Rca-α isoform have constrained photosynthesis and plant growth.
458 *Plant Journal* **103**: 2250–2262.
459
460 **Lamesch P, Berardini TZ, Li D, Swarbreck D, Wilks C, Sasidharan R, Muller R, Dreher K,**
461 **Alexander DL, Garcia-Hernandez M,** *et al.* **2012**. The Arabidopsis Information Resource (TAIR):
462 improved gene annotation and new tools. *Nucleic Acids Research* **40**: D1202–D1210.
463
464 **Lu Q, Ding S, Reiland S, Rödiger A, Roschitzki B, Xue P, Gruissem W, Lu C, Baginsky S**. **2015**.
465 Identification and characterization of chloroplast casein kinase II from Oryza sativa (rice). *Journal of*
466 *Experimental Botany* **66**: 175–187.
467

Lyons E, Freeling M. **2008**. How to usefully compare homologous plant genes and chromosomes as
DNA sequences. *The Plant Journal* **53**: 661–673.

Lyons E, Pedersen B, Kane J, Alam M, Ming R, Tang H, Wang X, Bowers J, Paterson A, Lisch D,
et al. **2008**. Finding and Comparing Syntenic Regions among Arabidopsis and the Outgroups Papaya,
Poplar, and Grape: CoGe with Rosids. *Plant Physiology* **148**: 1772–1781.

Marques DN, Stolze SC, Harzen A, Nogueira ML, Batagin-Piotto KD, Piotto FA, Mason C,
Azevedo RA, Nakagami H. **2021**. Comparative phosphoproteomic analysis of tomato genotypes with
contrasting cadmium tolerance. *Plant Cell Reports* **40**: 2001–2008.

McKain MR, Tang H, McNeal JR, Ayyampalayam S, Davis JI, dePamphilis CW, Givnish TJ, Pires
JC, Stevenson DWm, Leebens-Mack JH. **2016**. A phylogenomic assessment of ancient polyploidy and
genome evolution across the Poales. *Genome Biology and Evolution*: evw060.

Nagarajan R, Gill KS. **2018**. Evolution of Rubisco activase gene in plants. *Plant Molecular Biology* **96**:
69–87.

Neuwald AF, Aravind L, Spouge JL, Koonin E V. 1999. AAA $^{+}\square$: A Class of Chaperone-Like
ATPases Associated with the Assembly, Operation, and Disassembly of Protein Complexes. *Genome*
Research **9**: 27–43.

Nguyen L-T, Schmidt HA, von Haeseler A, Minh BQ. **2015**. IQ-TREE: A Fast and Effective
Stochastic Algorithm for Estimating Maximum-Likelihood Phylogenies. *Molecular Biology and*
Evolution **32**: 268–274.

Orr DJ, Robijns AKJ, Baker CR, Niyogi KK, Carmo-Silva E. **2023**. Dynamics of Rubisco regulation
by sugar phosphate derivatives and their phosphatases. *Journal of Experimental Botany* **74**: 581–590.

Parry MAJ, Andralojc PJ, Scales JC, Salvucci ME, Carmo-Silva AE, Alonso H, Whitney SM. **2013**.
Rubisco activity and regulation as targets for crop improvement. *Journal of Experimental Botany* **64**:
717–730.

Paterson AH, Bowers JE, Chapman BA. **2004**. Ancient polyploidization predating divergence of the
cereals, and its consequences for comparative genomics. *Proceedings of the National Academy of*
Sciences **101**: 9903–9908.

Poolman MG, Fell DA, Thomas S. **2000**. Modelling photosynthesis and its control. *Journal of*
Experimental Botany **51**: 319–328.

Portis AR, Jr. 2003. Rubisco activase - Rubisco's catalytic chaperone. *Photosynthesis Research* **75**: 11–
27.

Portis AR, Li C, Wang D, Salvucci ME. **2007**. Regulation of Rubisco activase and its interaction with
Rubisco. *Journal of Experimental Botany* **59**: 1597–1604.

Qu Y, Mueller-Cajar O, Yamori W. **2023**. Improving plant heat tolerance through modification of
Rubisco activase in C3 plants to secure crop yield and food security in a future warming world. *Journal of*
Experimental Botany **74**: 591–599.
517

518 **Reiland S, Messerli G, Baerenfaller K, Gerrits B, Endler A, Grossmann J, Gruissem W, Baginsky**
519 **S**. **2009**. Large-Scale Arabidopsis Phosphoproteome Profiling Reveals Novel Chloroplast Kinase
520 Substrates and Phosphorylation Networks . *Plant Physiology* **150**: 889–903.
521
522 **Ristic Z, Momčilović I, Bukovnik U, Prasad PVV, Fu J, DeRidder BP, Elthon TE, Mladenov N**.
523 **2009**. Rubisco activase and wheat productivity under heat-stress conditions. *Journal of Experimental*
524 *Botany* **60**: 4003–4014.
525
526 **Robinson SP, Portis AR**. **1989**. Adenosine triphosphate hydrolysis by purified rubisco activase. *Archives*
527 *of Biochemistry and Biophysics* **268**: 93–99.
528
529 **Rundle SJ, Zielinski RE**. **1991**. Alterations in barley ribulose-1,5-bisphosphate carboxylase/oxygenase
530 activase gene expression during development and in response to illumination. *Journal of Biological*
531 *Chemistry* **266**: 14802–14807.
532
533 **Salinas P, Fuentes D, Vidal E, Jordana X, Echeverria M, Holuigue L**. **2006**. An Extensive Survey of
534 CK2 α and β Subunits in Arabidopsis: Multiple Isoforms Exhibit Differential Subcellular Localization.
535 *Plant and Cell Physiology* **47**: 1295–1308.
536
537 **Salvucci ME, van de Loo FJ, Stecher D**. **2003**. Two isoforms of Rubisco activase in cotton, the
538 products of separate genes not alternative splicing. *Planta* **216**: 736–744.
539
540 **Salvucci ME, Portis AR, Ogren WL**. **1985**. A soluble chloroplast protein catalyzes
541 ribulosebisphosphate carboxylase/oxygenase activation in vivo. *Photosynthesis Research* **7**: 193–201.
542
543 **Salvucci ME, Portis AR, Ogren WL**. **1986**. Light and CO_2 Response of Ribulose-1,5-Bisphosphate
544 Carboxylase/Oxygenase Activation in *Arabidopsis* Leaves. *Plant Physiology* **80**: 655–659.
545
546 **Salvucci ME, Werneke JM, Ogren WL, Portis AR**. **1987**. Purification and Species Distribution of
547 Rubisco Activase. *Plant Physiology* **84**: 930–936.
548
549 **Samol I, Shapiguzov A, Ingelsson B, Fucile G, Crèvecoeur M, Vener A V., Rochaix JD,**
550 **Goldschmidt-Clermont M**. **2012**. Identification of a photosystem II phosphatase involved in light
551 acclimation in Arabidopsis. *Plant Cell* **24**.
552
553 **Sanchez de Jimenez E, Medrano L, Martinez-Barajas E**. **1995**. Rubisco Activase, a Possible New
554 Member of the Molecular Chaperone Family. *Biochemistry* **34**: 2826–2831.
555
556 **Scafaro AP, Bautsoens N, den Boer B, Van Rie J, Gallé A**. **2019**. A Conserved Sequence from Heat-
557 Adapted Species Improves Rubisco Activase Thermostability in Wheat. *Plant Physiology* **181**: 43–54.
558
559 **Schweer J, Türkeri H, Link B, Link G**. **2010**. AtSIG6, a plastid sigma factor from Arabidopsis, reveals
560 functional impact of cpCK2 phosphorylation. *The Plant Journal* **62**: 192–202.
561
562 **Shivhare D, Mueller-Cajar O**. **2017**. In Vitro Characterization of Thermostable CAM Rubisco Activase
563 Reveals a Rubisco Interacting Surface Loop. *Plant Physiology* **174**: 1505–1516.
564
565 **Soleh MA, Tanaka Y, Kim SY, Huber SC, Sakoda K, Shiraiwa T**. **2017**. Identification of large
566 variation in the photosynthetic induction response among 37 soybean [Glycine max (L.) Merr.] genotypes
567 that is not correlated with steady-state photosynthetic capacity. *Photosynthesis Research* **131**: 305–315.
568

Soltis DE, Smith SA, Cellinese N, Wurdack KJ, Tank DC, Brockington SF, Refulio Rodriguez NF, Walker JB, Moore MJ, Carlsward BS, _et al._ 2011. Angiosperm phylogeny: 17 genes, 640 taxa. _American Journal of Botany_ **98**: 704–730.

Somerville CR, Portis AR, Ogren WL. **1982**. A Mutant of _Arabidopsis thaliana_ Which Lacks Activation of RuBP Carboxylase _In Vivo_. _Plant Physiology_ **70**: 381–387.

Stitt M, Quick WP, Schurr U, Schulze E-D, Rodermel SR, Bogorad L. **1991**. Decreased ribulose-1,5-bisphosphate carboxylase-oxygenase in transgenic tobacco transformed with ?antisense? rbcS. _Planta_ **183**.

Stitt M, Schulze D. **1994**. Does Rubisco control the rate of photosynthesis and plant growth? An exercise in molecular ecophysiology. _Plant, Cell & Environment_ **17**: 465–487.

Stotz M, Mueller-Cajar O, Ciniawsky S, Wendler P, Hartl FU, Bracher A, Hayer-Hartl M. **2011**. Structure of green-type Rubisco activase from tobacco. _Nature structural & molecular biology_ **18**: 1366–70.

Tang H, Bowers JE, Wang X, Paterson AH. **2010**. Angiosperm genome comparisons reveal early polyploidy in the monocot lineage. _Proceedings of the National Academy of Sciences_ **107**: 472–477.

To K-Y, Suen D-F, Chen S-CG. **1999**. Molecular characterization of ribulose-1,5-bisphosphate carboxylase/oxygenase activase in rice leaves. _Planta_ **209**: 66–76.

Uhrig RG, Labandera A-M, Tang L-Y, Sieben NA, Goudreault M, Yeung E, Gingras A-C, Samuel MA, Moorhead GBG. **2017**. Activation of Mitochondrial Protein Phosphatase SLP2 by MIA40 Regulates Seed Germination. _Plant Physiology_ **173**: 956–969.

Uhrig RG, Moorhead GB. **2011**. Two Ancient Bacterial-like PPP Family Phosphatases from Arabidopsis Are Highly Conserved Plant Proteins That Possess Unique Properties . _Plant Physiology_ **157**: 1778–1792.

Werneke JM, Chatfield JM, Ogren WL. **1989**. Alternative mRNA splicing generates the two ribulosebisphosphate carboxylase/oxygenase activase polypeptides in spinach and Arabidopsis. _The Plant Cell_ **1**: 815–825.

White-Gloria C, Johnson JJ, Marritt K, Kataya A, Vahab A, Moorhead GB. **2018**. Protein Kinases and Phosphatases of the Plastid and Their Potential Role in Starch Metabolism. _Frontiers in Plant Science_ **9**.

Wijewardene I, Shen G, Zhang H. **2021**. Enhancing crop yield by using Rubisco activase to improve photosynthesis under elevated temperatures. _Stress Biology_ **1**: 2.

Willems P, Horne A, Van Parys T, Goormachtig S, De Smet I, Botzki A, Van Breusegem F, Gevaert K. **2019**. The Plant <scp>PTM</scp> Viewer, a central resource for exploring plant protein modifications. _The Plant Journal_ **99**: 752–762.

Willems P, Sterck L, Dard A, Huang J, De Smet I, Gevaert K, Van Breusegem F. **2024**. The Plant PTM Viewer 2.0: in-depth exploration of plant protein modification landscapes. _Journal of Experimental Botany_.

620 **Yin Z, Meng F, Song H, Wang X, Xu X, Yu D**. **2010**. Expression Quantitative Trait Loci Analysis of
621 Two Genes Encoding Rubisco Activase in Soybean . *Plant Physiology* **152**: 1625–1637.
622
623 **Zhang N, Kallis RP, Ewy RG, Portis AR**. **2002**. Light modulation of Rubisco in *Arabidopsis* requires a
624 capacity for redox regulation of the larger Rubisco activase isoform. *Proceedings of the National*
625 *Academy of Sciences* **99**: 3330–3334.
626
627 **Zhang N, Portis AR**. **1999a**. Mechanism of light regulation of Rubisco: A specific role for the larger
628 Rubisco activase isoform involving reductive activation by thioredoxin-f. *Proceedings of the National*
629 *Academy of Sciences* **96**: 9438–9443.
630
631 **Zhang N, Portis AR**. **1999b**. Mechanism of light regulation of Rubisco: A specific role for the larger
632 Rubisco activase isoform involving reductive activation by thioredoxin-f. *Proceedings of the National*
633 *Academy of Sciences* **96**: 9438–9443.
634
635 **Zhu X-G, de Sturler E, Long SP**. **2007**. Optimizing the Distribution of Resources between Enzymes of
636 Carbon Metabolism Can Dramatically Increase Photosynthetic Rate: A Numerical Simulation Using an
637 Evolutionary Algorithm. *Plant Physiology* **145**: 513–526.
638

XSDDQ Phosphorylation Site Amino Acid Legend

A
D
E
F
I
M
Q
T
V

Tandem duplications