The role of phosphorylation and dephosphorylation in the regulation of Rubisco

activase

Nikita Bhatnagar[a,b], , 0000-0001-7832-1347

Sarah S. Chung[a,b], sohyeon2@illinois.edu, 0009-0005-6096-5343

John Hodge[a,b], jghodge@illinois.edu, 0000-0003-0793-1871

Ryan A Boyd[b], raboyd2@illinois.edu, 0000-0003-4009-7700

Sang Yeol Kim[c]

Mia Sands[d], xin7@illinois.edu, 0000-0002-4925-9772

Andrew Leakey[a,b,,e,f], leakey@illinois.edu, 0000-0001-6251-024X

Donald R. Ort[a,b,,e,f], d-ort@illinois.edu, 0000-0002-5435-4387

Steven Burgess[a,b,,e], sjb287@illinois.edu, 0000-0003-2353-7794

Affiliations

[a]The Center for Advanced Bioenergy and Bioproducts Innovation (CABBI), University of

Illinois Urbana-Champaign

[b]Carl R. Woese Institute for Genomic Biology, University of Illinois Urbana-Champaign,

Urbana, IL 61801

[c]INARI Agriculture Inc, West Lafayette, IN, USA

[d]Department of Bioengineering, University of Illinois, Urbana-Champaign, Urbana, IL 61801,

USA

[d]Department of Plant Biology, University of Illinois Urbana-Champaign, Urbana IL 61801

[e]Department of Crop Sciences, University of Illinois Urbana-Champaign, Urbana IL 61801

Corresponding Author

26 Steven J. Burgess

28 Keywords: Rubisco activase (Rca), protein phosphorylation, post-translational modification,

29 photosynthesis

SUMMARY

Rubisco activase (Rca) is an ATP-dependent chaperone that facilitates dissociation of inhibitory sugar phosphates from the catalytic sites of ribulose-1,5-bisphosphate carboxylase/oxygenase (Rubisco) during photosynthesis. In Arabidopsis, Rca is negatively regulated by dark-dependent phosphorylation of Thr78. Here we identify chloroplast localized *SHEWENELLA-LIKE PROTEIN PHOSPHATASE 1 (SLP1)* as the major phosphatase responsible for light-dependent Rca dephosphorylation. The prevalence of Thr78 in Rca was investigated across sequences from 91 plant species, finding 29 (~32%) species shared a threonine in the same position. Analysis of seven C3 species with an antibody raised against a Thr78 phospho-peptide demonstrated that this position is phosphorylated in multiple genera. However, light-dependent dephosphorylation of Thr78 was observed only in Arabidopsis. Further, phosphorylation of Thr78 could not be detected in any of the four C4 grass species examined. The results suggest that despite conservation of Thr78 in Rca from a wide range of species, a regulatory role for phosphorylation at this site may be limited. This provides a case study for how variation in post-translational regulation can amplify functional divergence across the phylogeny of plants beyond what is explained by sequence variation in a metabolically important protein.

INTRODUCTION

Photosynthesis provides energy for all forms of life involving light-driven fixation of carbon dioxide and water into carbohydrates while producing oxygen. The rate and efficiency of this system depends upon several factors. Under saturating light, or low CO_2 concentrations, the activity of Ribulose-1,5-bis-phosphate carboxylase/oxygenase (Rubisco) is commonly limiting (Stitt *et al.*, 1991; Hudson *et al.*, 1992; Stitt & Schulze, 1994; Poolman *et al.*, 2000; Zhu *et al.*, 2007). In turn, Rubisco activation state is regulated by Rubisco activase (Rca), a member of the AAA+ ATPase superfamily (Somerville et al., 1982; Salvucci et al., 1986; Jiang et al., 1994). During CO_2 fixation, five carbon sugar phosphates can bind to Rubisco active sites preventing catalysis or blocking activation, by preventing carbamylation of Lys-21 (Orr et al., 2023). Rca uses energy from ATP hydrolysis to drive conformational changes in Rubisco, facilitating the release of the inhibitory sugar phosphates from the Rubisco active site (Neuwald et al., 1999; Portis & Jr., 2003; Flecken et al., 2020). Thus, Rca is critical for regulation of Rca activity and carbon assimilation. This also makes it a target to improve photosynthetic efficiency in crop plants (Parry et al., 2013; Wijewardene et al., 2021; Qu et al., 2023; Orr et al., 2023).

Most species produce two forms of Rca, α and β which can form mixed complexes (Salvucci et al., 1985; Zhang & Portis, 1999a), with the longer α isoform possessing a 20-30 amino acid C-terminal extension (To et al., 1999). Isoforms can arise from alternate splicing(Werneke et al., 1989; Rundle & Zielinski, 1991; To et al., 1999), or from transcription of independent genes (Rundle & Zielinski, 1991; Salvucci et al., 2003). The relative expression level, thermostability and catalytic properties of Rca-α and Rca-β differ across species (Rundle & Zielinski, 1991; Crafts-Brandner et al., 1997; Carmo-Silva & Salvucci, 2013; Shivhare & Mueller-Cajar, 2017; Degen et al., 2021) and can be impacted by environmental factors (Shivhare & Mueller-Cajar, 2017; Scafaro et al., 2019; Kim et al., 2020, 2021; Degen et al., 2021). Some species have almost equal abundance of both isoforms (e.g., Arabidopsis), whereas in others Rca-β is more highly expressed (Salvucci et al., 1987). In several C4 grasses such as sorghum, only the Rca-β isoform is abundant under normal growing conditions, while Rca-α upregulated under stress (Sanchez de Jimenez et al., 1995; Crafts-Brandner & Salvucci, 2002; Ristic et al., 2009; Yin et al., 2010). In contrast in wheat one of two β isoforms is induced at elevated temperature (Degen et al., 2021).

Some forms of Rca are sensitive to inhibition by ADP (Robinson & Portis, 1989). Most
of our understanding about this process comes from Arabidopsis, where the C-terminal extension
of Rca-α contains two cysteine residues (Cys-451 and Cys-470), which are regulated by
thioredoxin f in response to ATP/ADP ratio (Zhang & Portis, 1999b; Zhang et al., 2002). In an
oxidized state, the two cysteine residues form a disulfide bridge (oxidized state, E_m ~344 mV pH
7.9), and C-terminal domain then interacts with the nucleotide binding pocket, increasing the
sensitivity of Rca to inhibition by ADP (Portis et al., 2007). This downregulates Rca activity, in
accordance with the redox state of the chloroplast stroma. In contrast, the activities of AtRca-β
are unaffected by thioredoxin (Zhang et al., 2002). Thus Rca-α provides a mechanism to fine-
tune Rubisco activation state in dynamic light environments. However, redox regulation of Rca
does not appear to be essential. The sensitivity of Rca-β isoforms to inhibition by ADP varies
(Carmo-Silva & Salvucci, 2013). Further, studies with transgenic Arabidopsis mutants
expressing only Rca-α showed lower Rubisco activation and impaired growth (Zhang et al.,
2002; Kim et al., 2020), while plants only expressing Rca-β had higher biomass accumulation
under fluctuating light, where the impact of the rate of Rubisco reactivation on net carbon gain is
amplified relative to non-fluctuating light (Carmo-Silva & Salvucci, 2013). Solanaceae species,
such as tobacco, are reported to only express Rca-β under normal conditions (Nagarajan & Gill,
2018). Instead of redox regulation, activity of tobacco Rca-β is sensitive to the ratio of
ADP/ATP, the precise mechanism behind this sensitivity is unclear, but it has been demonstrated
to be caused by a substitution in one of 17 amino acid positions which vary between Arabidopsis
and tobacco sequences (Carmo-Silva & Salvucci, 2013). ADP sensitivity alone may therefore
explain control of Rca in species or mutants lacking an α isoform (Carmo-Silva & Salvucci,
2013).

Post-translational modification of Rca may add a supplemental layer of regulation
(Amaral et al., 2024). Phosphorylation of Threonine 78 (Thr78) in Arabidopsis Rca was
identified during a phospho-proteomic screen of dark-harvested leaf proteins (Reiland *et al.*,
2009). Subsequent research showed Thr78 was phosphorylated in the dark and under low light
intensities, but not under high light, suggesting a regulatory role (Reiland *et al.*, 2009;
Boex☐Fontvieille *et al.*, 2014; Kim *et al.*, 2016). Functional support for this hypothesis came
from several findings:(1) phosphorylation at Thr78 (Thr78-Rca) in Arabidopsis caused a
significant decrease in plant biomass that was linked to a decreased Rubisco activation state(Kim

et al., 2019); (2) transgenic Arabidopsis expressing phospho-null mutants of Rca (T78A) had
enhanced photosynthetic efficiency and plant growth; and (3) substitution of Thr78-to-Ser
(T78S) resulted in hyper-phosphorylation, and the transgenic plants expressing T78S-Rca
showed reduced carbon assimilation rate and slower plant growth under continuous and
fluctuating-light regimes (Kim *et al.*, 2019).

cpCK2 was shown to be the major kinase responsible for dark phosphorylation of AtRca,
with T-DNA insertion, loss-of-function mutant lines showing a reduction in Rca phosphorylation
(Kim *et al.*, 2016) but the phosphatase that is active in the light has not been identified. In
addition to the PPH/TAP3 and PBCP phosphatases that act in opposition to STN7 and STN8 to
regulate the kinetics of state transitions (Samol *et al.*, 2012), several other phosphatases
belonging to phosphoprotein phosphatase (PPP), protein phosphatase 2C (PP2C), and
phosphotransferase gene families have been reported to be localized in chloroplast based on the
presence of Chloroplast Transit Peptide (CTP)(White-Gloria *et al.*, 2018), but their function
remains unclear (White-Gloria *et al.*, 2018).

Here, we sought to define the mechanisms controlling Thr78 phosphorylation, and
prevalence of this PTM in photosynthetic organisms. This was achieved by considering the
following five questions, (1) what is the enzyme response for dephosphorylation of Thr78? (2) at
what stage of flowering plant evolution did Thr78 emerge? (3) how conserved is Thr78 among
photosynthetic angiosperms? (4) to what extent is Thr78 phosphorylated in different species? (4)
and is light-dependent de-phosphorylation of Thr78 a general mechanism of controlling of RCA
activity?

MATERIAL AND METHODS
Determination of Rca isoform and photosynthetic species containing Thr78-Rca
To analyze the conservation of Thr78 in Rca protein across plant species, the sequence of
Arabidopsis Rca (AtRca: AT2G39730) was retrieved from the Arabidopsis thaliana TAIR10
genome assembly (Lamesch *et al.*, 2012) on Phytozome v13 (Goodstein *et al.*, 2012). Syntelogs
to AtRca (i.e., shared positional homology along chromosomes) were subsequently identified in
monocots (*Setaria viridis* Rca: Sevir.8G261433) and asterids (*Mimulus guttatus* Rca:
Migut.F00758), which have homology to this sequence using either synteny mapping features on
CoGe (Lyons & Freeling, 2008; Lyons *et al.*, 2008). Setaria Rca was chosen as an example of a

gene that has been previously characterized (Kim et al. 2021), while Mimulus shares comparable
synteny to Arabidopsis in the genomic region surrounding Rca enabling identification of
syntelogs in the asterids. Working from these three representatives, additional syntelogues were
locally identified based on their closest known Rca sister lineage within the Angiosperm
phylogeny across other accessible plant genomes on these databases. *Amborella trichopoda* was
leveraged as an outgroup for flowering plants in this case study given that this is known to be the
one of the earliest diverging lineages for flowering plants (Soltis *et al.*, 2011). Additionally,
homologues were surveyed, which contained threonine at position 78 phosphorylation site even
if they didn't precisely fit this stringent criterion of synteny using pBLAST on NCBI
(https://blast.ncbi.nlm.nih.gov/Blast.cgi) for key agronomic model systems. This ultimately
resulted in 217 isoform CDS sequences being identified. Alternative splicing of the same locus
in some lineages meant that these CDS sequences represented 153 unique genes from across 91
species and 50 plant families. The longer, Rca-α, sequence was preferentially retained as a
taxonomic representative for each alternative splicing gene when available for phylogenetic
analysis. The CDS sequence of these genes were visualized in Seaview (Gouy *et al.*, 2010) and
codon-aligned using MUSCLE (Edgar, 2004). Obvious pseudogenes that were missing large
portions of their AAA+ ATPase domain such as the rice sequence (LOC_Os11g47980) were
removed prior to phylogenetic analysis of this DNA alignment. A maximum likelihood tree was
subsequently assembled using IQTree (Nguyen *et al.*, 2015) with the Amborella Rca sequence
being specified as an outgroup and 1000 bootstraps being run to generate statistical support for
nodes along the tree with the automatic substitution model selection determining a SYM+R8
model was the best-fit according to Bayesian Information Criterion. The resulting treefile were
then visualized either in Figtree (http://tree.bio.ed.ac.uk/software/figtree/) (Figure S3) or in R
using the Phytools and Ape libraries (Paradis & Schliep, 2019; Revell, 2024). Ancestral state
reconstructions were performed using the Ape function *ace* particularly for the original position
78 amino acid identity between syntelogous genes, and the manner of α/ß origin (either by
alternative splicing or distinct genes) for each plant family (Figure S4).
169
Plant growth and stress treatment
Arabidopsis Col-0 and camelina seeds were sterilized in 1 % (v/v) bleach for 5 min, rinsed with
70 % EtOH, and dried on filter paper. The sterilized seeds were sown on MS (Murashige and

Skoog basal salt mixture powder, Phyto Technology Laboratories, M524) agar (0.8 %, w/v,
Sigma-Aldrich, A1296) with 1 % (w/v) sucrose. The plates were stored at 4 °C for 2 days for
cold stratification before transfer to a growth chamber. Seedlings were transferred into pots (5
cm diameter) filled with soil (Berger; PRO-MIX BX All Purpose Growing Mix) before the third
true leaf emerged, which was approximately 5 to 7 days after germination (DAG). The other
plants (brassica, soybean, green bean, and Medicago) were directly sown in soil after
sterilization. Arabidopsis homozygous mutants of SLP1 (SALK_129247C) were ordered from
ABRC, and homozygous lines were confirmed by genomic DNA PCR using LB_6313R
(TCAAACAGGATTTTCGCCTGCT), SLP1_LP (GCCTGATCATCAAGTGACCTC) and
SLP1_RP (CGGGAACCATGTCTGACTTAG) xfs designed using iSect tool by SALK
(http://signal.salk.edu/tdnaprimers.2.html) (Alonso *et al.*, 2003). All plant species were grown at
22 °C, 16 h light and 8 h dark, 200 µmol m^{-2} s^{-1} PPFD with 2100 K 400W high pressure sodium
bulbs (Ceramalux ALTO; C400S51).

Protein extraction and immunoblotting assay

Total proteins were extracted from three frozen leaf discs (1 cm in diameter) using a
Qiagen Tissuelyzer for tissue grinding. The ground tissue was suspended in 0.5 mL of extraction
buffer containing 62.5 mM Tris-HCl (pH 6.8), 2% SDS, 10% glycerol, 0.35 M 2-
mercaptoethanol, and 0.001% protease inhibitor cocktail (Sigma-Aldrich; P9599). The
suspension was heated at 95°C for 5 minutes to denature proteins and inactivate proteases,
followed by centrifugation at 13,500 rpm for 30 minutes at 4°C. The supernatant was collected,
and 10 µL aliquots were separated on 10% SDS-PAGE gels. Proteins from the gels were
transferred electrophoretically to 0.2 µm nitrocellulose membranes (Bio-Rad) and blocked with
PBS solution with 5% (w/v) non-fat milk powder (Sigma) for 1h at room temperature. For
immunodetection, primary antibodies were diluted in Intercept Blocking Buffer (LI-COR; 927-
70001); anti-Rca (Agrisera; AS10 700) at 1:5000 and anti-pT78 (GenScript) at 1:4000, and
incubated overnight at 4°C. Modification-specific antibody was generated against the
phosphopeptide antigen RGLAYDpT78DDQQDC (Arabidopsis anti-pT78). For each peptide
antigen, the Cys residue at the N- or C-terminus was added for coupling to keyhole limpet
hemocyanin (KLH). All custom antibodies were produced by GenScript (Piscataway, NJ, USA).
Secondary detection used a fluorescent goat anti-rabbit antibody (Invitrogen; A32735) at a

1:10,000 dilution in PBS-T, incubated for 1 h at room temperature, and visualized with a LI-COR Odyssey Infrared Imaging System. All immunoblotting experiments were conducted at least twice to ensure reproducibility of results.

Protein phosphorylation assay with phosphatase inhibitor

WT Col-0 plants were grown as described above and whole rosettes were sampled and dipped in the protein phosphatase inhibitor buffer: 1% (v/v) solution of phosphatase inhibitor cocktail 2 (Sigma: P5726) and phosphatase inhibitor cocktail 3 (Sigma: P0044), overnight under dark. For dark treatment samples, the protein extraction was done using the overnight incubated samples and for light treatment samples the rosettes were exposed to 150 µmol m^{-2} s^{-1} PPFD light for 2 h and then total protein was extracted.

Physiological analysis

For light induction analysis, the leaves of 5-week WT and *slp1* mutants were dark adapted overnight before measurement in a portable infra-red gas exchange system (LI-6800, LICOR Environmental, Lincoln, Nebraska, USA) connected to a multiphase flash fluorometer (6800-01A) with a 2 cm^2 chamber. During low light incubation the plants were managed under dark conditions to make sure that Rca remained phosphorylated in both WT and *slp1* mutants. The chamber settings were 1:1 red:blue ratio, leaf temperature of 22 °C, CO_2 level of 420 µmol mol^{-1}, and 50% relative humidity. The measurement started with a low light of 20 µmol m^{-2} s^{-1} photosynthetic photon flux density (PPFD) for 20 min, then the PPFD was increased to high light conditions of 1000 µmol m^{-2} s^{-1} for 60 min, measurements were logged every 10 s. For CO_2 response analysis, plants were measured at the same high light conditions described above after finishing the light induction measurement. The CO_2 level was adjusted to produce a plot of net CO_2 assimilation against intercellular CO_2 concentration (A-C_i curve). The CO_2 levels were changed every 90 – 120 s in the following order: 420, 300, 200, 100, 50, 10, 420, 420, 420, 600, 800, 1200, 1800 µmol mol^{-1}. The relative comparisons and significant differences between *slp1* and WT were determined using students t-test (n = 4 to 6).

Statistical Analysis

Comparisons of rosette size and assimilation rate between *slp1* and WT were done using a two-
tailed Students' t-test using data from four to six biological replicates (individual plants) in
Microsoft Excel v16.88.

RESULTS

SHEWENELLA-LIKE PROTEIN PHOSPHATASE 1 (SLP1) dephosphorylates Rca in-vivo

Light-dependent dephosphorylation of AtRca suggested that an unknown protein
phosphatase is operating. To test this hypothesis, the phosphorylation state of Rca was assessed
under light and dark conditions in the presence and absence of a protein phosphatase inhibitor
cocktail (PPIC) (Fig. 1A and B). Under dark conditions, Rca phosphorylation increased
compared to the sample lacking PPIC (Fig. 1B) which is consistent with the presence of a protein
phosphatase. Dephosphorylation of Rca occurred in the light (Fig. 1B), even in the presence of
the PPIC, which may be because the level of dephosphorylation exceeded the ability of the PPIC
to inhibit activity. To identify the gene responsible for dephosphorylation of Rca, all known
chloroplast localized protein phosphatases in Arabidopsis (White-Gloria *et al.*, 2018) were
screened for reported associations with Rca in the literature and co-expression databases. Out of
9 chloroplast localized protein phosphatases, AtSLP1 (AT1G07010) showed a potential
association based on co-expression (Fig. 1C). We hypothesized that in the absence of *AtSLP1*,
AtRca will not be dephosphorylated in the light. To test this, a homozygous T-DNA insertion
mutant of *slp1* was identified (Figure S1) and assessed for the ability to dephosphorylate Rca in
the dark and light. *slp1* mutants showed phosphorylation of Rca in the dark, as also observed in
the case of Col-0, but in the mutant phosphorylation was also observed in the light (Fig. 1D).

We hypothesized that lack of Rca dephosphorylation in *slp1* mutants would lead to
decreased photosynthesis and a growth penalty. Analysis of 5-week-old plants found *slp1*
mutants have a significant reduction in rosette size relative to Col-0 under normal light
conditions (Fig. 1E and F). To assess the impact of SLP1 on steady state photosynthesis, A/Ci
(net CO_2 assimilation vs internal CO_2 concentration) curves were performed on Col-0 and *slp1*
mutants, finding no difference between lines (p=0.05; Figure S2). As there is a tight correlation
between Rubisco activation state and the rate of induction of photosynthesis during shade to sun
transitions, the impact of SLP1 on the rate of photosynthetic induction after a step-increase in

photosynthetic photon flux density from 20 to 1000 μmol m^{-2} s^{-1} was assessed, again finding no

significant difference compared to Col-0 (p=0.05; Fig. 1G).

Thr 78 in Rca is common

To study the conservation of Thr78 (Arabidopsis numbering) in Rca, 1893 protein

sequences from 479 plants species were assessed. Among these, 203 species retained Thr78

(Table S1). Next, a more stringent synteny informed ortholog phylogeny of Rca-α and Rca-β

isoforms was generated across angiosperms to address the questions: at what stage of flowering

plant evolution did the putative Thr78 phosphosite emerge, and how well conserved is it among

photosynthetic angiosperms? Leveraging maximum likelihood gene family phylogeny ancestral

states, Thr78 distribution was estimated for each family, including deeper nodes within the tree

such as the branch point of the *Amborellaceae* (Fig. 2). The presence of Thr78 was observed in

most plant families (Fig. 2), while among the *Poaceae*, that includes major crop species, the

presence of Isoleucine (Ile) was found to dominate at position 78 compared to Threonine (Fig.

2). The difference in the abundance of both amino acids, Thr and Ile at this site is similar at the

evolutionary scale, and an estimate of the ancestral state was ambiguous. Interestingly, the *α* and

β copies appeared polyphyletic, with multiple origins of the two isoforms arising from alternate

splicing or gene duplication across the phylogeny (Fig. S4, and S5). Within the Brassicaceae

Arabidopsis thaliana and its immediate relatives all appear to have a single copy of Rca with

alpha and beta forms being the result of alternative splicing of a single transcript.

By contrast in the Fabaceae, *Glycine max* has four gene copies of Rca (two each of *α* and

β) with the Papillionoid legumes more generally having two copies of Rca. There is no

equivalent duplication event in *Cercis canadensis* that would indicate a Fabaceae-wide pattern.

Therefore, this pattern of two Papillionoid copies is consistent with the whole-genome doubling

(WGD) event preceding the diversification of this subfamily, and by undergoing an additional

allopolyploid event, *Glycine max* has two additional copies for each of these ancestral

homeologues (Gill, *et al.*, 2009; Koenen, *et al.*, 2021). Within Poaceae, *Setaria viridis* and the

grasses more generally displayed two mirrored clades, which reflect the known taxonomic

sorting of this family. The grass family has undergone three WGD events (τ, σ, and ρ), two of

which are shared with monocots, with the p WGD unique to the Poaceae (Paterson *et al.*, 2004;

Tang *et al.*, 2010; McKain *et al.*, 2016). Although superficially resembling the *ρ* WGD event,

these grass copies are physically proximal to their sister copies in the adjacent clade suggesting
that this may instead be an ancient tandem duplication event.

Light-dependent dephosphorylation of Thr78 in Rca is unique to Arabidopsis

Previous reports showed that both AtRca isoforms phosphorylate in the dark and are
dephosphorylated as plants are exposed to light. To test whether Thr78 phosphorylation status is
light-sensitive in species other than Arabidopsis, we screened three C3 plants from the
Brassicaceae for dark-dependent phosphorylation of Thr78, using a pT78 antibody (Kim et al.,
2019) (Fig. 3A). As in Arabidopsis, Rca-α and Rca-β isoforms arise from alternate splicing in
Brassicaceae and the Thr78 site is conserved in all proteins (Fig. 3B). However, unlike
Arabidopsis, Rca in *Camelina sativa* (*Cs*), *Brassica oleraceae* (*Bo*) and *Brassica napus* (*Bn*)
phosphorylated in dark but lacked light-induced dephosphorylation (Fig. 3A). These data suggest
that although Thr78 is phosphorylated in multiple species, light-dependent regulation may be
specific to Arabidopsis.

To broaden the analysis, we performed the dephosphorylation assay on several legumes:
Glycine max (*Gm*), *Phaseolus vulgaris* (*Pv*), and *Medicago truncatula* (*Mt*). The Rca isoforms in
these species are derived from individual genes of *Rca-α* and *Rca-β*, unlike the situation in the
Brassicaceae. The Rca-α isoforms retain Thr78, whereas the β isoforms have Ile78 in *Gm* and
Mt, and Val78 in *Pv* (Fig. 3B). Publicly available data suggest the transcription level of Rca-α is
low relative to β in both *Gm* and *Pv,* and moderately expressed in *Mt* (Table S2). The light/dark
assay on legume plants showed that both the isoforms were expressed, and phosphorylated in the
dark, but as with the Brassicaceae, did not dephosphorylate in the light (Fig. 3A). The reason
why the antibody was able to detect phosphorylation of β forms, which lack Thr78, is unclear. It
is possible that there is relaxed selectively, but we consider these results with caution.

To study the impact of phosphorylation in C4 plants, we analyzed Rca phosphorylation in
four species of C4 grass: *Sorghum bicolor* (*Sb*), *Miscanthus* (*Ms*), *Saccharum officinarum* (*Su*),
and *Setaria viridis* (*Sv*) (Fig. 4A). Each of these species possess two separate genes for the
alternative Rca isoforms, with Rca-α containing Thr78, but with it lacking from Rca-β (Fig. 4B,
C). In accordance with previous reports (Kim *et al.*, 2021), only the Rca-β isoform could be
detected under normal growth conditions (Fig. 4A and C). An unspecific cross-reaction was
present in C4 grass samples probed with pT78, but the detected protein was too large to be Rca-

β, subsequently no phosphorylation of Rca-β was detected (Fig. 4A). To test whether a C4 Rca-α isoform could be phosphorylated, sorghum plants were exposed to heat induce Rca-α isoform expression (Fig. 4C). Plants accumulated SbRca-α after 24 h heat treatment, but no bands of the correct size could be detected using the pT78 antibody (Fig. 4C). To rule out the possibility that a lack of detectable Thr78 phosphorylation was due to high protein phosphatase activity, the assay was repeated in the presence of a protein phosphatase inhibitor cocktail (PPIC), but again protein phosphorylation was not observed under any condition (Fig. 5). This result suggests that SbRca-α T78 is not phosphorylated by cpCK2 as it is in Arabidopsis even though the gene for cpCK2 is present in sorghum.

DISCUSSION

Rca is subject to multiple levels of regulation to adjust carbon fixation in response to environmental conditions (Amaral *et al.*, 2024). Differences in gene expression, biochemical properties of isoforms, redox status, and post-translational modification have been topics of investigation (Boex⬚Fontvieille *et al.*, 2014; Kim *et al.*, 2016, 2019). Dark dependent phosphorylation of Thr78 is proposed to downregulate Rca activity leading to reduced assimilation (Kim et al. 2019). cpCK2 is described as the major kinase responsible for phosphorylation of AtRca Thr78 based on dramatically reduced phosphorylation in *cpcpk2* T-DNA insertion lines (Kim *et al.*, 2016). Here we identified Shewanella-like protein phosphatase 1 (SLP1) as the second component of this regulatory mechanism, acting to dephosphorylate AtRca Thr78 in the light (Fig. 1). SLP1 is a highly conserved protein phosphatase reported to be blue light responsive in Arabidopsis (Depuydt & Vandepoele, 2021) and belongs to a protein family consisting of two members, of which, SLP1 is the only chloroplast localized isoform (Uhrig & Moorhead, 2011). A role for mitochondrial targeted SLP2 has been described in negative regulation of gibberellic acid related seed germination (Uhrig *et al.*, 2017), but the physiological role and targets of SLP1 have not previously been explored.

If phosphorylation of Rca has a dominant impact on activity, plants lacking SLP1 would be predicted to see an increase Thr78 phosphorylation in the light, leading to lower Rubisco activity, and lower Anet. an impact on photosynthesis, with no difference observed in assimilation rates between WT and knockout lines during light induction or CO_2 response (Fig. 1G and Fig. S2). These findings are in keeping with observations by Kim *et al.* (2016), who

reported no change in assimilation rates when Thr78 phosphorylation state was reduced in *cpck2*
mutants. These results were attributed to redundancy in Rca regulation between protein
phosphorylation and control by thioredoxin (Kim *et al.*, 2016) and the same rationale may apply
here. Support for a functional role for phosphorylation of Thr78 in Arabidopsis has only been
shown in Rca variant lines lacking the redox sensitive C-terminal extension (Kim *et al.*, 2019).
Our analysis, conducted on lines containing the C-terminal extension, are therefore consistent
with the hypothesis that Rca possesses multiple redundant mechanisms in Arabidopsis, which
operate to regulate activity in response to variable light environments. The impact of SLP1 on
photosynthesis may only be apparent in the absence of redox regulation.

Despite the absence of impact on Anet, the *slp1* mutant showed reduced growth (Fig. 1).
Given that the *cpck2* mutant, which also showed altered phosphorylation of Rca, did not have a
growth phenotype, we attribute this to an unrelated impact on chloroplast physiology. As a
largely uncharacterized protein, the specificity and targets of SLP1 are unknown. Phosphatases
have traditionally been considered promiscuous, with specificity dictated by regulatory subunits
(ref). SLP1 is not known to have regulatory subunits, it is therefore conceivable there are other
targets within the chloroplast that are impacted in the mutant leading to the observed growth
phenotype.

Substitution of Thr78 with Serine lead to a relative increase in phosphorylation at the site,
in Arabidopsis resulted in smaller plants with decreased photosynthesis (Kim *et al.*, 2019).
Altering a single amino acid, Thr78, was proposed as a means of generating plants with better
photosynthesis. However, the prevalence of Thr78 across species, the frequency of post-
translational modification and functionality were unclear. To begin to address these questions we
expanded previous searches of genomic databases, which assessed the evolution of Rca copy
number across the green lineage (Nagarajan & Gill, 2018; Kim *et al.*, 2019), to use a synteny
based approach. Nagarajan and Gill (2018) placed the emergence of the Rca-α CTE and alternate
splicing mechanisms in the charophytes, before the divergence of land plants and there is only
one reported example of a species only having evidence for expression of Rca α (Nagarajan &
Gill, 2018). Here, despite α and β forms (bearing or lacking a c-terminus respectively) occurring
within each model system, these α and β forms appear to represent repeated cases of
evolutionary convergence on these two isoforms (Figure S5) through distinct means within each
clade: subfunctionalization (Birchler & Yang, 2022), following tandem duplication in the grasses

as previously described (Nagarajan & Gill, 2018), WGD in the legumes, or alternative splicing
of a single coding sequence into alternative isoforms in Brassicaceae. This may suggest that Rca
isoforms can be quite plastic in the way they arise within distinct plant families where α and β
forms may not necessarily reflect true homology within Rubisco activase isoforms.

Previous assessment of a smaller number of Rca sequences failed to identify any with
Thr78 replaced by Serine, leading to speculation that the configuration with the attendant hyper
phosphorylation observed in Arabidopsis has been selected against (Kim *et al.*, 2019). In the
initial set of Rca sequences identified by BLAST search, we identified gene models from at least
seven species (Table S3) which do have Ser78, although these were not considered orthologs in
the synteny analysis. Of the 25 species found to contain Rca sequences with Thr78, it remains to
be seen how many undergo phosphorylation. Our data are consistent with it occurring in at least
the Brassicaceae, while phosphorylation of an equivalent Thr residue has been reported in
tomato, suggesting this modification may occur more widely (Marques et al., 2021).

We found no evidence for phosphorylation of Thr78 in monocot species, including
Sorghum, Setaria, sugarcane and Miscanthus. The peptide fragment used for antibody generation
is highly conserved, with 100% identity between Arabidopsis Rca and Sorghum Rca-α
"RGLAYDTSDDQQDC" and should therefore have been able to detect phosphorylation if
present under the conditions tested. There are several potential explanations for this finding,
including an absence of the kinase, differences in localization or expression profiles of the
kinase, or regulation of the kinase. In Arabidopsis, Rca is phosphorylated by cpCK2 (Kim et al.,
2016). A homologous gene has been characterized in rice (LOC_Os03g55490) (Lu et al.,
2015)and syntenic sequences can be found in C4 grasses including Sorghum (Sobic.001G07970)
(Carvalho et al., 2020). As Sorghum operates two-celled C4 photosynthesis for an interaction to
occur between Rca and cpCK2 both must be expressed in the same cell type. Transcriptomics
suggests that cpCK2 is expressed in both mesophyll and bundle sheath cells in maize (Chang et
al., 2012), Setaria (John et al., 2014)and Sorghum (Döring et al., 2016), albeit with ~2-fold
mesophyll enrichment in maize and Setaria whereas Rca is localized in the bundle sheath. Even
so, it appears that the orthologous cpCK2 is present in the same compartment as Rca (i.e., the
bundle sheath cell chloroplast). However, in the monocot species tested Rca-β, which lacks
Thr78, is the predominate isoform under normal conditions and Rca-α, which possess Thr78, is
induced during stress (Shivhare & Mueller-Cajar, 2017; Scafaro et al., 2019; Kim et al., 2020,

2021). Whether C4 monocot cpCK2 retains the ability to phosphorylate Thr78 in Rca-α, or if the
bundle sheath stromal environment is permissive for phosphorylation remains unclear (Kim et
al., 2016). CK2 exists as a multigene family, composed of catalytic α subunits and regulatory
beta subunits (Salinas et al., 2006; Schweer et al., 2010), which form hetero-tetramers that can
differ in substrate specificity (Domańska et al., 2005). Intriguingly there appears to be a lack of
regulatory β subunits in the chloroplast (Salinas et al., 2006), so differences in regulatory
subunits impacting specificity appears unlikely, although multiple chloroplast CK2 containing
complexes have been detected (Baginsky et al., 1997) suggesting the protein does not act alone.
However, further research is required to characterize the regulation of monocot cpCK2 and
whether it can phosphorylate Rca.

Conservation of phosphorylation of Thr78 in some C3 plants is not evidence of a
conserved regulatory role. Indeed, in the species examined, strong evidence for light-dependent
dephosphorylation of Thr78 was only found for Arabidopsis, with phosphorylation of Rca
detected in both the light and dark in other C3 species (Fig. 3). These data may suggest a limited
scope for phosphorylation as a means of regulation, with most species depending on other
mechanisms, such as changes in redox state, catalytic properties or expression. It may be that
alternate post-translational modifications of Rca have greater importance in other species and
several different ones have been described (Amaral et al., 2024). Interestingly, there are several
reports of phosphorylation of S79 in Arabidopsis (Willems et al., 2019), and this modification
was also observed in a Brachypodium, Chlamydomonas and maize (Willems et al., 2019, 2024).
Further phosphoproteomics of maize during a diurnal cycle suggests that rather than Thr78,
Ser79 undergoes dark-dependent phosphorylation (Gao et al., 2023). However, the prevalence of
S79 phosphorylation is also likely to vary, as seen by a failure to detected in protein samples
from *Glycine max* (soybean-S86) (Soleh et al., 2017) .Therefore, the prevalence and function of
Ser79 phosphorylation is unclear, and whether it can play a similar role to Thr78 warrants further
investigation.

Taken together our results suggest that Thr78 can be found in Rca sequences across the
green lineage, although the prevalence varies, and degree to which it undergoes phosphorylation
is inconsistent. Our data provides support for a mechanism whereby Thr78 is phosphorylated in
the dark by cpCK2 and dephosphorylated in the light by SLP1 in Arabidopsis. The lack of
impact on photosynthesis when cpCK2 or SLP1 are knocked out suggest that phosphorylation of

Thr78 acts redundantly with redox regulation. More widely, it is likely that regulation of Rca can

occur through multiple mechanisms, including redox, sensitivity to ADP and gene expression

and the relative importance of each varies.

Acknowledgments

We would like to thank Toby Kellogg and Shawn Abrahams for providing feedback on the

phylogenic analyses. This work was funded by the DOE Center for Advanced Bioenergy and

Bioproducts Innovation (U.S. Department of Energy, Office of Science, Biological and

Environmental Research Program under Award Number DE-SC0018420). Any opinions,

findings, and conclusions or recommendations expressed in this publication are those of the

author(s) and do not necessarily reflect the views of the U.S. Department of Energy.

Competing interests

The authors report no conflict of interest

Authors contributions

Conceptualization (SK, NB, SJB, DO), Investigation (NB, SC, MS, JH, SK), Formal Analysis

(NB), Project administration (NB, SJB, DO), Supervision (AL, SJB, DO), Visualization (NB),

Writing - original draft (NB, JH, SC, DO, SJB), Writing – review and editing (NB, JH, DO, AL,

SJB).

FIGURE LEGENDS

Figure 1: SLP1 dephosphorylates Rca in Arabidopsis. Protein phosphatase assay with PPIC (+P) and without PPIC (-P) isoforms, comparing total abundance (A) to phosphorylation (B) of Rca by immunoblot. (C) Putative protein-protein interactions between Rca and chloroplast localized protein phosphatases. (D) Immunoblot analysis of light dependent dephosphorylation of Rca in WT (Col-0) and *slp1* mutants. (E) Image of 5-week-old rosettes comparing WT (Col-0) and slp1 plants. (F) Comparison of rosette size between WT (Col-0) and slp1 plants (n=20). (G) Photosynthetic induction curves of WT (Col-0) and *slp1* mutant plants (n=5 to 6).

Figure 2: Phylogenetic representation of the amino acid present at the phosphorylation site (peptide position 78) within extant Rca gene models. Ancestral states for families, and when available deeper orders of Angiosperms, are denoted at their corresponding nodes as pie charts. The nodes depicting the presence of Thr78 is colored green and the ones with Ile78 is colored navy blue. Less abundant amino acids at the peptide position 78 are depicted as varying shades of blue and purple. The detailed confirmation of bootstrap values is mentioned in Figure S3.

Figure 3: Light dependent dephosphorylation of Thr78 in Rca is specific to Arabidopsis. (A) Immunoblot analysis comparing protein phosphorylation and total abundance of Rca under light (L) and dark (D) conditions (At: *Arabidopsis thaliana*, Cs: *Camelina sativa*, Bn: *Brassica napus*, Bo: *Brassica oleraceae*, Gm: *Glycine max*, Pv: *Phaseolus vulgaris*, Mt: *Medicago truncatula*). The upper panel shows the total Rca protein extracted from plant samples and the middle panel shows the phosphorylated Rca protein. α and β denote the 2 isoforms of Rca protein. CBB (Coomassie Brilliant Blue) depicts the loading control. (B) Sequence alignment depicting the conservation of Thr78 amino acid in C3 species. The amino acid lacking color codes depicts the change in the Thr78 site to Isoleucine (*I*) in *Glycine max* and *Medicago truncatula*, and Valine (*V*) in Phaseolus vulgaris. The letter "A" and "B" at the end of the species labelling denotes the Rca-α and Rca-β isoforms. (C) The phylogenetic tree depicts the division of C3 Brassicaceae family and legumes based on Rca isoforms arising from protein splicing in Brassicaceae as compared to different genes in legumes and presence and absence of Thr78 amino acid in the Rca-β isoform. The blue boxes depict the presence of Thr78 amino acid whereas the red depicts the lack of Thr78. The yellow box depicts the experimental confirmation of protein phosphorylation at Thr78 phosphosite.

Figure 4: Phosphorylation of Rca is not observed in selected C4 moncots. (A) Protein phosphorylation assay depicting four C4 plant samples in light and dark conditions (At: *Arabidopsis thaliana*, Sv: *Setaria viridis*, Sb: *Sorghum bicolor*, Su: *Saccharum officinarum*, Ms: *Miscanthus*). The upper panel shows the accumulation of Rca protein quantified with Anti-Rca antibody and middle panel shows the phosphorylated Rca protein quantified with Anti-pT78 antibody. α and β denote the 2 isoforms of Rca protein. CBB denotes the loading control. (B) Sequence alignment of motif comparing Arabidopsis and select C4 plants. (C) Immunoblot analysis comparing Rca phosphorylation under heat for 24 h in WT sorghum. The numbers denote the time in hours after the heat stress. CBB denotes the loading control.

Figure 5: Sorghum lacks phosphorylation of Thr78 with or without PPIC treatment. (A) Immunoblot analysis comparing sorghum WT samples collected after heat stress at 42^{o}C under light and dark treatment. The control samples were collected at 28^{o}C. The + and – sign denotes presence and absence of treatment, respectively. The left panel shows Rca protein accumulation and right panel depicts analysis using anti-pT78 antibody.

Figure S1: The genomic DNA (gDNA) PCR with primer combinations of LB, LP and RP showed that LB primer T-DNA insertion was confirmed in homozygous lines whereas WT showed no bands (upper panel). The SLP1 gene primers showed gDNA bands in the WT and not in the *slp1* mutants (lower panel).

Figure S2: The *ACi* curve showed that both WT (Col-0) and slp1 mutants overlap suggesting that both the genotypes have no difference in the photosynthetic efficiency. Error bars denote standard error. Green dots are *slp1* mutants and black dots are WT (Col-0).

Figure S3: The conservative phylogeny structure of Rca orthologs where all nodes with bootstrap support (blue numbers) greater than 80% being retained whereas other lower support nodes were collapsed into polytomies.

Figure S4: Rca α (white) and β (black) isoforms on extant gene models as well as alternatively spliced gene models which produce both Rca α and β (grey). Ancestral states for families, and when available deeper orders of Angiosperms, are denoted at their corresponding nodes as pie charts.

Figure S5: The likely pattern that could result in the polytomies of *α* and *β* isoforms ancestry states that Angiosperms initially beginning as Rca *α* lineages out of which *β* isoforms repeatedly arise independently following either subfunctionalization of duplicate gene copies or alternative

533 splicing of α gene models to β isoforms as well. Within certain lineages where α isoforms are

534 lost all the Rca copies become canalized into a β isoform state thereafter.

REFERENCES

Alonso JM, Stepanova AN, Leisse TJ, Kim CJ, Chen H, Shinn P, Stevenson DK, Zimmerman J, Barajas P, Cheuk R, *et al.* **2003**. Genome-wide insertional mutagenesis of Arabidopsis thaliana. *Science (New York, N.Y.)* **301**: 653–7.

Amaral J, Lobo AKM, Carmo□Silva E. 2024. Regulation of Rubisco activity in crops. *New Phytologist* **241**: 35–51.

Baginsky S, Tiller K, Link G. 1997. Transcription factor phosphorylation by a protein kinase associated with chloroplast RNA polymerase from mustard (Sinapis alba). *Plant Molecular Biology* **34**: 181–189.

Birchler JA, Yang H. 2022. The multiple fates of gene duplications: Deletion, hypofunctionalization, subfunctionalization, neofunctionalization, dosage balance constraints, and neutral variation. *The Plant Cell* **34**: 2466–2474.

Boex□Fontvieille E, Daventure M, Jossier M, Hodges M, Zivy M, Tcherkez G. 2014. Phosphorylation pattern of <scp>R</scp>ubisco activase in *<scp>A</scp> rabidopsis* leaves. *Plant Biology* **16**: 550–557.

Carmo-Silva AE, Salvucci ME. 2013. The Regulatory Properties of Rubisco Activase Differ among Species and Affect Photosynthetic Induction during Light Transitions . *Plant Physiology* **161**: 1645–1655.

Carvalho DS, Nishimwe A V., Schnable JC. 2020. IsoSeq transcriptome assembly of C $_3$ panicoid grasses provides tools to study evolutionary change in the Panicoideae. *Plant Direct* **4**.

Chang Y-M, Liu W-Y, Shih AC-C, Shen M-N, Lu C-H, Lu M-YJ, Yang H-W, Wang T-Y, Chen SC-C, Chen SM, *et al.* **2012**. Characterizing Regulatory and Functional Differentiation between Maize Mesophyll and Bundle Sheath Cells by Transcriptomic Analysis . *Plant Physiology* **160**: 165–177.

Crafts-Brandner SJ, van de Loo FJ, Salvucci ME. 1997. The Two Forms of Ribulose-1,5-Bisphosphate Carboxylase/Oxygenase Activase Differ in Sensitivity to Elevated Temperature. *Plant Physiology* **114**: 439–444.

Crafts-Brandner SJ, Salvucci ME. 2002. Sensitivity of Photosynthesis in a C4 Plant, Maize, to Heat Stress. *Plant Physiology* **129**: 1773–1780.

Degen GE, Orr DJ, Carmo□Silva E. 2021. Heat□induced changes in the abundance of wheat Rubisco activase isoforms. *New Phytologist* **229**: 1298–1311.

Depuydt T, Vandepoele K. 2021. Multi□omics network□based functional annotation of unknown Arabidopsis genes. *The Plant Journal* **108**: 1193–1212.

Domańska K, Zieliński R, Kubiński K, Sajnaga E, Masłyk M, Bretner M, Szyszka R. **2005**.
Different properties of four molecular forms of protein kinase CK2 from Saccharomyces
cerevisiae. *Acta biochimica Polonica* **52**: 947–51.

Döring F, Streubel M, Bräutigam A, Gowik U. **2016**. Most photorespiratory genes are
preferentially expressed in the bundle sheath cells of the C 4 grass *Sorghum bicolor*. *Journal of*
Experimental Botany **67**: 3053–3064.

Flecken M, Wang H, Popilka L, Hartl FU, Bracher A, Hayer-Hartl M. **2020**. Dual Functions
of a Rubisco Activase in Metabolic Repair and Recruitment to Carboxysomes. *Cell* **183**: 457-
473.e20.

Gao Z, Yang X, Mei Y, Zhang J, Chao Q, Wang B. **2023**. A dynamic phosphoproteomic
analysis provides insight into the <scp>C4</scp> plant maize (*Zea mays* L.) response to natural
diurnal changes. *The Plant Journal* **113**: 291–307.

Goodstein DM, Shu S, Howson R, Neupane R, Hayes RD, Fazo J, Mitros T, Dirks W,
Hellsten U, Putnam N, *et al.* 2012. Phytozome: a comparative platform for green plant
genomics. *Nucleic Acids Research* **40**: D1178–D1186.

Gouy M, Guindon S, Gascuel O. **2010**. SeaView Version 4: A Multiplatform Graphical User
Interface for Sequence Alignment and Phylogenetic Tree Building. *Molecular Biology and*
Evolution **27**: 221–224.

Hudson GS, Evans JR, von Caemmerer S, Arvidsson YBC, Andrews TJ. **1992**. Reduction of
Ribulose-1,5-Bisphosphate Carboxylase/Oxygenase Content by Antisense RNA Reduces
Photosynthesis in Transgenic Tobacco Plants. *Plant Physiology* **98**: 294–302.

Jiang C, Quick WP, Alred R, Kliebenstein D, Rodermel SR. **1994**. Antisense RNA inhibition
of Rubisco activase expression. *The Plant Journal* **5**: 787–798.

John CR, Smith-Unna RD, Woodfield H, Covshoff S, Hibberd JM. **2014**. Evolutionary
Convergence of Cell-Specific Gene Expression in Independent Lineages of C4 Grasses   . *Plant*
Physiology **165**: 62–75.

Kim SY, Bender KW, Walker BJ, Zielinski RE, Spalding MH, Ort DR, Huber SC. **2016**.
The plastid casein kinase 2 phosphorylates rubisco activase at the Thr-78 site but is not essential
for regulation of rubisco activation state. *Frontiers in Plant Science* **7**.

Kim SY, Harvey CM, Giese J, Lassowskat I, Singh V, Singh V, Cavanagh AP, Cavanagh
AP, Spalding MH, Finkemeier I, *et al.* 2019. In vivo evidence for a regulatory role of
phosphorylation of Arabidopsis Rubisco activase at the Thr78 site. *Proceedings of the National*
Academy of Sciences of the United States of America.

Kim SY, Slattery RA, Ort DR. **2021**. A role for differential Rubisco activase isoform
expression in C 4 bioenergy grasses at high temperature. *GCB Bioenergy* **13**: 211–223.

Kim SY, Stessman DJ, Wright DA, Spalding MH, Huber SC, Ort DR. **2020**. Arabidopsis
plants expressing only the redox-regulated Rca-α isoform have constrained photosynthesis and
plant growth. *Plant Journal* **103**: 2250–2262.
632

633 **Lamesch P, Berardini TZ, Li D, Swarbreck D, Wilks C, Sasidharan R, Muller R, Dreher**
634 **K, Alexander DL, Garcia-Hernandez M, *et al.* 2012**. The Arabidopsis Information Resource
635 (TAIR): improved gene annotation and new tools. *Nucleic Acids Research* **40**: D1202–D1210.
636

637 **Lu Q, Ding S, Reiland S, Rödiger A, Roschitzki B, Xue P, Gruissem W, Lu C, Baginsky S**.
638 **2015**. Identification and characterization of chloroplast casein kinase II from Oryza sativa (rice).
639 *Journal of Experimental Botany* **66**: 175–187.
640

641 **Lyons E, Freeling M**. **2008**. How to usefully compare homologous plant genes and
642 chromosomes as DNA sequences. *The Plant Journal* **53**: 661–673.
643

644 **Lyons E, Pedersen B, Kane J, Alam M, Ming R, Tang H, Wang X, Bowers J, Paterson A,**
645 **Lisch D, *et al.* 2008**. Finding and Comparing Syntenic Regions among Arabidopsis and the
646 Outgroups Papaya, Poplar, and Grape: CoGe with Rosids. *Plant Physiology* **148**: 1772–1781.
647

648 **Marques DN, Stolze SC, Harzen A, Nogueira ML, Batagin-Piotto KD, Piotto FA, Mason C,**
649 **Azevedo RA, Nakagami H**. **2021**. Comparative phosphoproteomic analysis of tomato genotypes
650 with contrasting cadmium tolerance. *Plant Cell Reports* **40**: 2001–2008.
651

652 **McKain MR, Tang H, McNeal JR, Ayyampalayam S, Davis JI, dePamphilis CW, Givnish**
653 **TJ, Pires JC, Stevenson DWm, Leebens-Mack JH**. **2016**. A phylogenomic assessment of
654 ancient polyploidy and genome evolution across the Poales. *Genome Biology and Evolution*:
655 evw060.
656

657 **Nagarajan R, Gill KS**. **2018**. Evolution of Rubisco activase gene in plants. *Plant Molecular*
658 *Biology* **96**: 69–87.
659

660 **Neuwald AF, Aravind L, Spouge JL, Koonin E V. 1999**. AAA $^{+}\square$: A Class of Chaperone-Like
661 ATPases Associated with the Assembly, Operation, and Disassembly of Protein Complexes.
662 *Genome Research* **9**: 27–43.
663

664 **Nguyen L-T, Schmidt HA, von Haeseler A, Minh BQ**. **2015**. IQ-TREE: A Fast and Effective
665 Stochastic Algorithm for Estimating Maximum-Likelihood Phylogenies. *Molecular Biology and*
666 *Evolution* **32**: 268–274.
667

668 **Orr DJ, Robijns AKJ, Baker CR, Niyogi KK, Carmo-Silva E**. **2023**. Dynamics of Rubisco
669 regulation by sugar phosphate derivatives and their phosphatases. *Journal of Experimental*
670 *Botany* **74**: 581–590.
671

Parry MAJ, Andralojc PJ, Scales JC, Salvucci ME, Carmo-Silva AE, Alonso H, Whitney
SM. **2013**. Rubisco activity and regulation as targets for crop improvement. *Journal of*
Experimental Botany **64**: 717–730.

Paterson AH, Bowers JE, Chapman BA. **2004**. Ancient polyploidization predating divergence
of the cereals, and its consequences for comparative genomics. *Proceedings of the National*
Academy of Sciences **101**: 9903–9908.

Poolman MG, Fell DA, Thomas S. **2000**. Modelling photosynthesis and its control. *Journal of*
Experimental Botany **51**: 319–328.

Portis AR, Jr. 2003. Rubisco activase - Rubisco's catalytic chaperone. *Photosynthesis Research*
75: 11–27.

Portis AR, Li C, Wang D, Salvucci ME. **2007**. Regulation of Rubisco activase and its
interaction with Rubisco. *Journal of Experimental Botany* **59**: 1597–1604.

Qu Y, Mueller-Cajar O, Yamori W. **2023**. Improving plant heat tolerance through
modification of Rubisco activase in C3 plants to secure crop yield and food security in a future
warming world. *Journal of Experimental Botany* **74**: 591–599.

Reiland S, Messerli G, Baerenfaller K, Gerrits B, Endler A, Grossmann J, Gruissem W,
Baginsky S. **2009**. Large-Scale Arabidopsis Phosphoproteome Profiling Reveals Novel
Chloroplast Kinase Substrates and Phosphorylation Networks . *Plant Physiology* **150**: 889–903.

Ristic Z, Momčilović I, Bukovnik U, Prasad PVV, Fu J, DeRidder BP, Elthon TE,
Mladenov N. **2009**. Rubisco activase and wheat productivity under heat-stress conditions.
Journal of Experimental Botany **60**: 4003–4014.

Robinson SP, Portis AR. **1989**. Adenosine triphosphate hydrolysis by purified rubisco activase.
Archives of Biochemistry and Biophysics **268**: 93–99.

Rundle SJ, Zielinski RE. **1991**. Alterations in barley ribulose-1,5-bisphosphate
carboxylase/oxygenase activase gene expression during development and in response to
illumination. *Journal of Biological Chemistry* **266**: 14802–14807.

Salinas P, Fuentes D, Vidal E, Jordana X, Echeverria M, Holuigue L. **2006**. An Extensive
Survey of CK2 α and β Subunits in Arabidopsis: Multiple Isoforms Exhibit Differential
Subcellular Localization. *Plant and Cell Physiology* **47**: 1295–1308.

Salvucci ME, van de Loo FJ, Stecher D. **2003**. Two isoforms of Rubisco activase in cotton, the
products of separate genes not alternative splicing. *Planta* **216**: 736–744.

Salvucci ME, Portis AR, Ogren WL. **1985**. A soluble chloroplast protein catalyzes
ribulosebisphosphate carboxylase/oxygenase activation in vivo. *Photosynthesis Research* **7**: 193–
201.

718
719 **Salvucci ME, Portis AR, Ogren WL**. **1986**. Light and CO $_2$ Response of Ribulose-1,5-
720 Bisphosphate Carboxylase/Oxygenase Activation in *Arabidopsis* Leaves. *Plant Physiology* **80**:
721 655–659.
722
723 **Salvucci ME, Werneke JM, Ogren WL, Portis AR**. **1987**. Purification and Species
724 Distribution of Rubisco Activase. *Plant Physiology* **84**: 930–936.
725
726 **Samol I, Shapiguzov A, Ingelsson B, Fucile G, Crèvecoeur M, Vener A V., Rochaix JD,
727 Goldschmidt-Clermont M**. **2012**. Identification of a photosystem II phosphatase involved in
728 light acclimation in Arabidopsis. *Plant Cell* **24**.
729
730 **Sanchez de Jimenez E, Medrano L, Martinez-Barajas E**. **1995**. Rubisco Activase, a Possible
731 New Member of the Molecular Chaperone Family. *Biochemistry* **34**: 2826–2831.
732
733 **Scafaro AP, Bautsoens N, den Boer B, Van Rie J, Gallé A**. **2019**. A Conserved Sequence from
734 Heat-Adapted Species Improves Rubisco Activase Thermostability in Wheat. *Plant Physiology*
735 **181**: 43–54.
736
737 **Schweer J, Türkeri H, Link B, Link G**. **2010**. AtSIG6, a plastid sigma factor from
738 Arabidopsis, reveals functional impact of cpCK2 phosphorylation. *The Plant Journal* **62**: 192–
739 202.
740
741 **Shivhare D, Mueller-Cajar O**. **2017**. In Vitro Characterization of Thermostable CAM Rubisco
742 Activase Reveals a Rubisco Interacting Surface Loop. *Plant Physiology* **174**: 1505–1516.
743
744 **Soleh MA, Tanaka Y, Kim SY, Huber SC, Sakoda K, Shiraiwa T**. **2017**. Identification of
745 large variation in the photosynthetic induction response among 37 soybean [Glycine max (L.)
746 Merr.] genotypes that is not correlated with steady-state photosynthetic capacity. *Photosynthesis*
747 *Research* **131**: 305–315.
748
749 **Soltis DE, Smith SA, Cellinese N, Wurdack KJ, Tank DC, Brockington SF,
750 Refulio□Rodriguez NF, Walker JB, Moore MJ, Carlsward BS, *et al.* 2011**. Angiosperm
751 phylogeny: 17 genes, 640 taxa. *American Journal of Botany* **98**: 704–730.
752
753 **Somerville CR, Portis AR, Ogren WL**. **1982**. A Mutant of *Arabidopsis thaliana* Which Lacks
754 Activation of RuBP Carboxylase *In Vivo*. *Plant Physiology* **70**: 381–387.
755
756 **Stitt M, Quick WP, Schurr U, Schulze E-D, Rodermel SR, Bogorad L**. **1991**. Decreased
757 ribulose-1,5-bisphosphate carboxylase-oxygenase in transgenic tobacco transformed
758 with□?antisense? rbcS. *Planta* **183**.
759
760 **Stitt M, Schulze D**. **1994**. Does Rubisco control the rate of photosynthesis and plant growth? An
761 exercise in molecular ecophysiology. *Plant, Cell & Environment* **17**: 465–487.
762

Stotz M, Mueller-Cajar O, Ciniawsky S, Wendler P, Hartl FU, Bracher A, Hayer-Hartl M. **2011**. Structure of green-type Rubisco activase from tobacco. *Nature structural & molecular biology* **18**: 1366–70.

Tang H, Bowers JE, Wang X, Paterson AH. **2010**. Angiosperm genome comparisons reveal early polyploidy in the monocot lineage. *Proceedings of the National Academy of Sciences* **107**: 472–477.

To K-Y, Suen D-F, Chen S-CG. **1999**. Molecular characterization of ribulose-1,5-bisphosphate carboxylase/oxygenase activase in rice leaves. *Planta* **209**: 66–76.

Uhrig RG, Labandera A-M, Tang L-Y, Sieben NA, Goudreault M, Yeung E, Gingras A-C, Samuel MA, Moorhead GBG. **2017**. Activation of Mitochondrial Protein Phosphatase SLP2 by MIA40 Regulates Seed Germination. *Plant Physiology* **173**: 956–969.

Uhrig RG, Moorhead GB. **2011**. Two Ancient Bacterial-like PPP Family Phosphatases from Arabidopsis Are Highly Conserved Plant Proteins That Possess Unique Properties . *Plant Physiology* **157**: 1778–1792.

Werneke JM, Chatfield JM, Ogren WL. **1989**. Alternative mRNA splicing generates the two ribulosebisphosphate carboxylase/oxygenase activase polypeptides in spinach and Arabidopsis. *The Plant Cell* **1**: 815–825.

White-Gloria C, Johnson JJ, Marritt K, Kataya A, Vahab A, Moorhead GB. **2018**. Protein Kinases and Phosphatases of the Plastid and Their Potential Role in Starch Metabolism. *Frontiers in Plant Science* **9**.

Wijewardene I, Shen G, Zhang H. **2021**. Enhancing crop yield by using Rubisco activase to improve photosynthesis under elevated temperatures. *Stress Biology* **1**: 2.

Willems P, Horne A, Van Parys T, Goormachtig S, De Smet I, Botzki A, Van Breusegem F, Gevaert K. **2019**. The Plant <scp>PTM</scp> Viewer, a central resource for exploring plant protein modifications. *The Plant Journal* **99**: 752–762.

Willems P, Sterck L, Dard A, Huang J, De Smet I, Gevaert K, Van Breusegem F. **2024**. The Plant PTM Viewer 2.0: in-depth exploration of plant protein modification landscapes. *Journal of Experimental Botany*.

Yin Z, Meng F, Song H, Wang X, Xu X, Yu D. **2010**. Expression Quantitative Trait Loci Analysis of Two Genes Encoding Rubisco Activase in Soybean . *Plant Physiology* **152**: 1625–1637.

Zhang N, Kallis RP, Ewy RG, Portis AR. **2002**. Light modulation of Rubisco in *Arabidopsis* requires a capacity for redox regulation of the larger Rubisco activase isoform. *Proceedings of the National Academy of Sciences* **99**: 3330–3334.

809   **Zhang N, Portis AR**. **1999a**. Mechanism of light regulation of Rubisco: A specific role for the
810   larger Rubisco activase isoform involving reductive activation by thioredoxin-f. *Proceedings of*
811   *the National Academy of Sciences* **96**: 9438–9443.
812
813   **Zhang N, Portis AR**. **1999b**. Mechanism of light regulation of Rubisco: A specific role for the
814   larger Rubisco activase isoform involving reductive activation by thioredoxin-f. *Proceedings of*
815   *the National Academy of Sciences* **96**: 9438–9443.
816
817   **Zhu X-G, de Sturler E, Long SP**. **2007**. Optimizing the Distribution of Resources between
818   Enzymes of Carbon Metabolism Can Dramatically Increase Photosynthetic Rate: A Numerical
819   Simulation Using an Evolutionary Algorithm. *Plant Physiology* **145**: 513–526.
820

XSDDQ Phosphorylation
Site Amino Acid Legend

A
D
E
F
I
M
Q
T
V
Tandem
duplications

A

B

A

At　Sv　Sb　Su　Ms

L D　L D　L D　L D　L D

kDa

55 —

43 —

Anti-Rca

55 —

43 —

Anti-pT78

55 —

43 —

CBB

B

	D R W R G L A Y D □ S D D Q Q D I T R G K G	
Arabidopsisthaliana B	D R W R G L A Y D T S D D Q Q D I T R G K G	90
Arabidopsisthaliana A	D R W R G L A Y D T S D D Q Q D I T R G K G	90
Setariaviridis B	D R W R G L V N D I S D D Q Q D I T R G K G	83
Zeamays B	D R W R G L A Y D I S D D Q Q D I T R G K G	86
Sorghumbicolor B	D R W R G L A Y D V S D D Q Q D I T R G K G	93
Misin B	D R W R G L A F D I S D D Q Q D I T R G K G	88
Setariaviridis A	D R W R G L A H D I S D D Q Q D I T R G K G	85
Sorghumbicolor A	D R W R G L A Y D T S D D Q Q D I T R G K G	98
Zeamays A	D R W R G L A L D T S D D Q Q D I T R G K G	84
Misin A	D R W R G L A Y D V S D D Q Q D I T R G K G	96

C

Sb-WT-Heat

0　0.5　1　1.5　2　24hr

kDa

43 —

Anti-Rca

43 —

Anti-pT78

43 —

CBB