Title: Are human super-predators always super-scary? A meta-analysis of wild animal

behavioural responses to human interactions

Authors:

Shawn Dsouza[1*]; Email: shawn27.dsouza@pm.me

Kartik Shanker[1]; Email: kshanker@iisc.ac.in

Maria Thaker[1]; Email: mthaker@iisc.ac.in

Affiliations: [1] Centre for Ecological Sciences, Indian Institute of Science, Bengaluru, India.

Author Contributions: The study was conceptualised by SD with substantial input from MT

and KS. Data collection and analysis was conducted by SD. The first draft of the manuscript and

subsequent revisions were written with equal contribution by all authors.

Data Accessibility Statement: Data and code used in this manuscript are available at

https://doi.org/10.5281/zenodo.15099679.

Keywords: HIREC, risk disturbance, non-consumptive, predator – prey interactions, fear

Abstract: 200; **Main Text:** 6246; **References:** 88; **Figures:** 5; **Tables:** 0; **Text boxes:** 0;

*Corresponding Author:

Shawn Dsouza

Mob No.: +91 9821158693

Email: shawn27.dsouza@pm.me

Address: Centre for Ecological Sciences, Indian Institute of Science, Bangalore 560012.

Abstract

Humans interact with wild animals through lethal activities (e.g., hunting, fishing) as well as non-lethal activities (e.g., tourism, recreation). While it is known that these different interactions alter animal behaviour, their relative impacts are not well understood. We conducted a systematic review and meta-analysis of the past 30 years of research, focusing on changes in foraging, vigilance, and movement behaviours in wild animals. Our results show that lethal human activities cause significant behavioural changes, with targeted species increasing vigilance and reducing foraging in affected areas. Active non-lethal activities elicited similar but generally weaker patterns, with many species showing little to no change in the magnitude of behavioural response. In contrast, passive non-lethal interactions, like those involving roads and development, produced highly variable responses depending on the species. These findings illustrate the importance of differentiating between types of human activity when considering their effects on wildlife. Notably, the most pronounced behavioural changes occurred where humans posed a direct lethal threat. Gaps in the literature, such as uneven data within and across species, and limited information on the history or context of human interactions currently limit our ability to better predict when and why animals change their behaviour in response to humans.

Introduction

Humans have shifted natural environments in dramatic ways, due in part to their unique ecology which has enabled them to occupy every ecoregion of the world and play multiple functional roles (Darimont et al. 2023). As predators, humans are more deadly than most other predators in both marine and terrestrial systems because they can target multiple trophic levels simultaneously and extract animals at a faster rate than natural predators (Darimont et al. 2015). Humans also occupy natural spaces for a variety of seemingly benign activities such as resource extraction, tourism, and hiking. Human development over the past two centuries has led to the alteration of environments at a rate that has not been witnessed in the history of the planet (Estes et al. 2011; Sih 2013). Wild animals therefore have no choice but to respond to human activities in the natural world.

Human social organisation and use of tools enables them to target multiple trophic levels at high efficiency (Treves & Naughton-Treves 1999). Thus, humans have been referred to as generalist super-predators (Clinchy et al. 2016), taking a disproportionately large number of animals from ecosystems. In fact, the rate of mortality of prey species from human predators, such as hunters and fishers, are much greater in magnitude than the sum of kill by all natural predators in both terrestrial and marine ecosystems (Darimont et al. 2015). Thus, humans create truly risky conditions for animals that they intentionally kill in the wild (Oriol-Cotterill et al. 2015).

In addition to hunting and fishing, humans also interact with wild animals through non–lethal activities (Boyle & Samson 1985), which can be further distinguished as active or passive.

Active non-lethal interactions include tourism, walking in parks, hiking in nature reserves, scuba

diving and snorkelling. Studies have shown that these activities can negatively influence animals

even though the humans in their habitats are not actively killing them (e.g. Brown et al. 2012;

Loehr et al. 2005; Montero-Quintana et al. 2020; Newey 2007). Passive non-lethal interactions

are even more pervasive, as they include engineered alterations to environments. Roads and

settlements, in particular, have the greatest potential to influence the behaviour of animals (e.g.

Cappa et al. 2017; Mehlhoop et al. 2022, etc). Some types of "passive" human interactions may

also cause significant mortality, such as through collisions on glass (Loss et al. 2015) and the

high occurrence of road-kill globally (Benson et al. 2015). Although some studies suggest that

non-lethal human activities can negatively influence animal behaviours (Boyle & Samson 1985;

selected examples: Papouchis et al. 2001; Davis et al. 1997; Westekemper et al. 2018; Ladle et

al. 2019; Nevin & Gilbert 2005), other studies show that animals habituate quickly to human

encounters and may even benefit from these encounters (see Smith et al. 2024, Gaynor et al.

2025). As such, non-lethal interactions may have positive or negative effects on wild animals

(see Smith et al. 2024). Thus, it remains unclear whether non-lethal interactions of humans on

animals are as strong as the lethal interactions in eliciting fear and changing behaviours.

Given that humans interact with animals in the wild with both lethal and non-lethal intent, the

individual and combined effects of these interactions on animal behaviours can potentially be

very high. In a recent narrative review, Smith et al. (2024) outlined a framework describing the

pathways by which lethal and non-lethal human interactions can change behavioural and

physiological responses of animals, with consequences for population growth and abundance.

They conclude that perceived risk of humans may induce phenotypic changes in wild animals,

but as expected, the evidence is mixed (Smith et al. 2024). What is still missing is a quantitative

assessment of the relative magnitude of effect for different types of human activities, which

would illustrate when, why and how wild animals respond to human-induced risk.

In this paper, we synthesise and review the current evidence for human-induced behavioural

changes in wild animals. We categorised studies based on whether wild animals were exposed to

interactions with humans of lethal intent (hunting, fishing), active and non-lethal intent

(ecotourism, hiking), and passive and non-lethal intent (roads and human settlements) (Figure 1

and 2). One of the clearest and most effective responses that prey use to minimise risk is by

shifting behaviour (Lima and Dill 1990). Decades of research has shown that animals change

their behaviours in response to natural predators, following a trade-off between energy gain and

safety that directly affect fitness (Brown & Kotler 2004; Sih 1984). Thus, we focused on changes

in foraging behaviours (bite rates and foraging time), vigilance (vigilance time and proportion)

and movement (displacement, home range size, and movement rate) of animals in response to

human interactions. Immediate responses, such as flight or escape, were excluded from this

meta-analysis as they occur at short time scales and may not have long-term fitness

consequences in and of themselves. Physiological responses, such as changes in hormone levels

(Sherrif et al. 2020; Smith et al. 2024), were also not included here, as these physiological

changes can influence multiple behavioural pathways and are variable across taxa (Sheriff and

Thaler 2014).

We predict that, in areas where humans have lethal intent towards animals, these animals will

strongly invest in anti-predator behavioural strategies, spending more time being vigilant, less

time foraging, and increasing their movement rate or displacement (Figure 1). We expect a lower

intensity of anti-predator responses to active and passive non-lethal interactions with humans, as

wild animals may be able to perceive such interactions as non-threatening or even beneficial

based on cues given out by humans in the habitat (Wirsing et al. 2021, Figure 1).

From the systematic review of published studies, we derived effect sizes that describe the

magnitude of behavioural change across these categories of human interactions. We then

conducted a meta-analysis to determine whether active or passive non-lethal interactions with

humans influence wild animal behaviours to the same extent as lethal human interactions. We

also tested the potential influence of body size and trophic level on the magnitude of behavioural

effects based on the hypothesis that these factors influence which animals are targeted by the

lethal actions of humans (i.e., they are hunted, trapped or fished). Finally, we highlight where

gaps in the literature have yielded skewed conclusions about human impacts on the behaviour of

wild animals and suggest avenues for future research in this field.

Methods

To determine the strength of behavioural change (i.e. effect sizes) caused by human activities,

we conducted a systematic literature survey that included studies where foraging, vigilance or

movement behaviours were measured in wild animals in response to potentially lethal human

interactions (hunting, including poaching and trapping, and fishing), non-lethal but active

interactions (walking in parks, hiking in nature reserves, off-road bicycling and animal

watching), and non-lethal but passive interactions (roads and human settlements) (Figure 2). We

focus on only wild animal behaviour that is indicative of long-term foraging - safety trade-offs

(Lima & Dill, 1990), thus we excluded studies of escape behaviour (typically measured as flight

initiation distance) and studies that measure animal physiology. We also excluded studies where

human interactions were induced to simulate a natural predator approach, if there was no

information about the underlying context of human interaction (i.e., lethal and non-lethal intent).

Search and literature database

We conducted a scoping search using the Web of Science database and their advanced search

function. Our initial search term was "(risk OR fear) AND (human OR anthropogenic) AND

(behavio*) AND ALL=(predat*)". The search terms were applied across all fields. The initial

search returned 2319 results on 12th January 2023. We downloaded the data and conducted a

literature mapping analysis to identify additional search terms and updated our search string to

"(human OR anthropo*) AND (risk OR fear OR NCE OR "trait mediated effect" OR

nonconsumptive* effect) AND (behavio*) AND (predat*)". Using our updated search string, we

conducted searches on the SCOPUS and WOS database as well as OpenThesis for grey literature

on 14th February 2023. We then used Rayyan, a literature management and screening software

to deduplicate our database (Foo et al. 2021; Ouzzani et al. 2016), which resulted in 7562

abstracts. This review was not registered.

Screening

We piloted our initial screening protocol on 100 randomly selected papers using Rayyan

(Ouzzani et al. 2016). We decided inclusion and exclusion criteria by evaluating titles, abstracts,

and keywords to determine the population studied, the exposure or intervention to the

population, the control if any and the outcome (see Appendix A for detailed inclusion criteria).

The initial screening was conducted by three independent researchers, who discussed conflicting

decisions on inclusion or exclusion of abstracts until a consensus was reached. We applied the

finalised screening protocol to all abstracts in our database and were left with 436 papers to

further evaluate. Papers chosen for inclusion during our initial screening were full-text screened.

We included 71 studies at this stage (See appendix A for our complete screening protocol).

Backward and forward search

We identified key studies and reviews during our screening for backward and forward search.

The key studies (n = 14) included major reviews, studies that first addressed the effects of human

interactions on animal behaviour and concept papers (see Appendix B for a full list). Backward

search identified papers cited by these studies and reviews, and forward search identified papers

that cited these key papers. We used a citation chaining tool called ResearchRabbit to execute the

search on 12th March 2024. We added the results to our existing database and deduplicated it to

create the final paper database for the review (Foo et al. 2021). After screening the additional

abstracts, we screened full-texts of 40 papers and 14 were added. Our final sample included 44

studies for our meta-analysis and an additional 41 for review (See Appendix B for complete list).

We have included a PRISMA (EcoEvo) flowchart and check-list in Appendix A and F,

respectively (O'Dea et al. 2021; Page et al. 2021).

Data collection

We collected data on the behavioural responses, study species, trophic level (Consumer, Primary

Predator and Secondary Predator), location of the study, treatments, controls, and sample size.

Trophic level, dietary guild and body mass of the study species from each study was curated

from Mammalbase (Lintulaakso 2021), Animaltraits database (Herberstein et al. 2022),

Handbook of the Birds of the World (Del Hoyo et al. 1992) and Fishbase (Froese & Pauly 2010).

We included omnivores as either Primary or Secondary predators based on their diets (e.g. bear

species were included as secondary predators, see Appendix A).

For the animal responses, we extracted data on the mean effects of foraging, vigilance, and

movement behaviours and associated variance measures (Figure 3). Studies that quantify

foraging and vigilance behaviours used focal individual sampling and scan sampling. The

measured behaviours were time spent vigilant, time spent foraging, frequency of vigilance

behaviour in a group, rate of individual vigilance behaviour, and foraging (bite) rates. We

included movement studies that measured movement rates (speed), displacement (distance

travelled), and home range size using GPS tags, collars and transmitters. Home range size was

included as it is a function of displacement and speed in a given sampling period. Where results

were not clearly reported we either used raw data to calculate mean effects and variance

measures or used a plot digitiser to extract data from figures when raw data were not available.

Studies in which data extraction was not possible were excluded from our meta-analysis and

used only for qualitative review. We did not include papers that only reported test statistics in

our meta-analysis (Mikolajewicz & Komarova 2019, See Appendix B for a complete list of

studies).

There were several types of study approaches that measured animal responses to human

activities. The majority of studies were observational or natural experiments. Spatial contrasts

included designs in which behaviours were compared across protected and unprotected areas for

lethal interactions (n = 8), areas with and without tourists for active interactions (n = 6) and areas

close and far away from roads or settlements for passive interactions (n = 5). Studies also

employed auditory stimuli (n = 5) such as road noise or human speech in areas with hunting

pressure or high traffic to test for the effect of lethal interactions and passive interactions

respectively. Other experimental studies employed direct human disturbance to simulate active

interactions (n = 6) and lethal interactions in areas with hunting (n = 3). Four studies used

temporal contrast in human activities such as hunting closures and tourist seasons to determine

the effects of active (n = 1) and lethal (n = 3) interactions. Finally, ten studies did not implement

or utilise a contrast for their comparison and instead determined correlations between human

activities for active interactions and distance from road for passive interaction on behaviour.

Analysis

We divided the studies into categories based on their study designs as experimental or

observational. We converted proportional and percentage data (means and measures of

uncertainty) to absolute values (number of individuals or time in seconds), and we converted all

measures of uncertainty to variance (using sample size to convert SE to SD) before analysis.

Furthermore, we calculated standardised mean difference for studies that reported an explicit

treatment and control (negative or positive), as $d = \dfrac{x_{treatment} - x_{control}}{s_{pooled}}$, where x_i is the mean of

treatment   and   control   outcomes   respectively   and

$s_{pooled} = \sqrt{\dfrac{s^2_{control}(n_{control}-1) + s^2_{treatmen}(n_{treatment}-1)}{n_{treatment} + n_{control} - 2}}$, here s^2_i is the sample variance and n_i is the

sample size.

All the correlational studies reported unstandardised regression coefficients (b), and thus, we

calculated   the   standardised   mean   differences   from b as $d = \dfrac{b}{s_{pooled}}$,   where

$s_{pooled} = \sqrt{\dfrac{s^2_y(N-1) - b^2\left(\dfrac{n_{treatment}\, n_{control}}{n_{treatmen} + n_{control}}\right)}{N-2}}$ here N is the total sample size (Lipsey and Wilson

2001). Where generalised linear models were used or where data was transformed prior to fitting

a linear model, we back transformed the coefficients to the original scale. We used a

control:treatment contrast, with treatment being those groups that were exposed to lethal or non-

lethal interactions and control being groups not exposed to interactions. The model coefficients

of studies which used a control:treatment contrast were multiplied by -1 to maintain consistency

across studies. We assigned each study a unique identifier (study ID); when we extracted

multiple data points from studies, we assigned a unique identifier to each data point (data ID).

We first fit an intercept only multilevel meta-analytic (MLMA) model using the metafor package

(Viechtbauer 2010) with species, study, and data id as random effects to determine the degree of

heterogeneity across studies after accounting for known sources of variation as follows:

244
$$y_{ij} = \mu + u_{ij} + s_i + d_j + \eta_{ij}$$

where, y_{ij} is the effect from j^{th} data point of the i^{th} study, μ is the overall mean across studies, u_{ij}

is the i^{th} estimate within the j^{th} study, s_i is the random effect due to the i^{th} study, d_j is the random

effect due to the j^{th} data point and η_{ij} is the error of the j^{th} estimate from the i^{th} study. The random

effect framework assumes that each study has its own true effect size which is derived from a

population of true effect sizes. Since this meta-analysis involves a comparison across taxa in

different habitats and geographies, the random effect framework is appropriate (Nakagawa et al.

2022; Noble et al. 2022). We used the Open Tree of Life API via the `rotl` package to build a

phylogenetic tree of species included in our meta-analysis (Michonneau et al. 2016;

OpenTreeOfLife et al. 2019). We also fit a phylogenetic model with relatedness as random

effect.

255
$$y_{ijk} = \mu + u_{ij} + s_i + d_j + p_k + \eta_{ij}$$

where, p_k is random effect due to the k^{th} species. Phylogeny did not capture significant variation

across all outcomes (Chamberlain et al. 2012; Cinar et al. 2022). We thus used a model with only

study identity, and data identity as random effects. We used the Q statistic to determine leftover

variation. Furthermore, we used both I^2 and H^2 to determine heterogeneity across studies (Harrer

et al. 2021). We used the final MLMA to determine whether there was an overall significant

effect across studies. Finally, we used the MLMA to perform a multiple regression to determine

the effect of type of human interactions and animal size on the observed effects. We have

described our analysis of sensitivity, robustness and bias of our dataset in Appendix D. We

report the mean and 95% confidence intervals of the strength of each behavioural response we

have considered in our meta-analysis; foraging, vigilance, and movement, to each type of human

interaction we have considered in our meta-analysis; lethal, active non-lethal and passive non-

lethal, as measured by standardised mean difference of responses to treatment and responses to

control. Additionally, we present regression analyses examining the effect of the different human

interaction types on these behaviours, including T statistics and significance levels.

All data preparation and analysis were conducted in R – 4.0.1 (R Core Team 2024). Data and

code are available at Dsouza *et al.* (2025).

Results

After screening, we included 85 studies in our systematic review, a majority of which were

published in the past two decades (Appendix A). Of these, 44 studies reported enough

information to be included in the meta-analysis, covering 38 species. These studies were widely

distributed across geographies and habitats ranging from 78ºN to 43ºS and 169ºE to 123ºW

(Figure 2). Across outcomes, 24 studies reported measures of foraging behaviours, which

included bite rate and time spent foraging, 29 studies focused on vigilance behaviours, which

included time spent vigilant and vigilance rate, and 9 studies measured movement behaviour,

which included home range size, displacement, and rate of movement (Figure 3). Eighteen

studies reported data from more than one major category of behaviour. The majority of these

papers (n = 36) studied primary consumers, such as *Cervus elaphus* and *Capreolus capreolus*

(Figure 4, Appendix A). Only 7 studies included secondary and tertiary consumers (Figure 4),

thus statistical comparisons across trophic levels were not conducted.

Changes in behaviours across types of human interactions

Across studies, the overall effect of human interactions on foraging behaviours was negative

(SMD = -0.178, 95% CI = -0.738 – 0.382, T_{23} = -0.624, p = 0.533) and on vigilance was positive

(SMD = 0.641, 95% CI = 0.292 – 0.991, T_{27} = 3.593, p = 0.0003). However, the effect of human

interactions on movement behaviour across studies was negligible (SMD = 0.191, 95% CI =

-0.219 – 0.594, T_6 = -0.172, p = 0.868, Appendix E). Upon fitting the MLMA model, there was

still significant heterogeneity among studies that was not accounted for by random effects

(Appendix E).

Foraging behaviour of animals decreased significantly in response to lethal human interactions

(SMD = -1.193, 95% CI = -2.279 – -0.108, Z_5 = -2.154, p = 0.031), and to a lesser extent in

response to passive non-lethal interactions (SMD = -0.255, 95% CI = -1.392 – 0.882, Z_9= -0.439,

p = 0.66, Figure 4). Active non-lethal interactions had an overall significant positive effect on

foraging (SMD = 0.331, 95% CI = -0.115 – 0.778, Z_{20} = 1.455, p = 0.145, Figure 4). Vigilance

behaviour increased significantly in response to lethal interactions (SMD = 1.1, 95% CI = 0.014–

2.186, Z_{12} = 2.991, p = 0.003) and active non-lethal interactions (SMD = 0.829, 95% CI =

0.549– 1.11, Z_{24} = 5.798, p = 0). However, vigilance behaviour was significantly reduced in

passive non-lethal interactions (SMD = -1.339, 95% CI= -2.042 – -0.054, Z_6 = -2.042, p =

0.041). There was no significant overall effect of human interactions on movement (Figure 5).

Overall, when we consider all studies in this analysis, foraging behaviours were significantly

more suppressed by lethal human activities than by active non-lethal human interactions (β =

-0.955 ± 0.51, T_{34} = -1.874, p = 0.07, Figure 5). There was no significant difference in the

magnitude of effect on foraging behaviours between active and passive non-lethal human

interactions. The magnitude of behavioural effects on vigilance behaviour due to lethal human

interactions was significantly higher than that of active non-lethal interactions ($\beta = 1.347 \pm$

0.553, $T_{46} = 2.433$, p $= 0.019$, Figure 5). There were no significant differences in the effect of

human interactions on movement behaviours across the three types of interactions (Appendix E,

Figure 5).

The body size of the animal did not influence the effect of human activities on foraging ($\beta = 0 \pm$

0, $T_{31} = -0.31$, $p = 0.75$), movement ($\beta = 0.005 \pm 0.006$, $T_{16} = 0.948$, $p = 0.37$), or vigilance

behaviour ($\beta = 0 \pm 0$, $T_{41} = 0.089$, $p = 0.92$).

Discussion

The term human "super-predator" was coined to highlight the efficiency, lethality, and scale at

which human predators operate (Clinchy et al. 2016; Darimont et al. 2015). Recent frameworks,

such as those proposed by Smith et al. (2024), have begun to assess the ecological impacts of

humans as super-predators. Building on this, our study offers a first step toward quantifying

these effects by conducting a meta-analysis of wild animal behavioural responses to human

interactions, differentiating between those with lethal and non-lethal intent. Our analysis,

spanning a broad geographic and taxonomic range (Figure 2), reveals that lethal interactions

exert the strongest influence on overall wild animal behaviour, followed by active non-lethal

interactions. In contrast, passive interactions, such as those with roads and human developments,

had a suppressive effect on wild animal vigilance behaviours but did not show consistent effects

on foraging or movement (Figure 5). Importantly, we found significant variation in wild animal

responses to human activity across interaction types (Figure 4), underscoring the strong context-

dependence of these effects across species. Below, we discuss the findings of our meta-analysis

within the broader context of human-animal interactions.

Are humans always super-scary when present?

Humans have hunted other animals for the majority of their evolutionary history (Treves &

Naughton-Treves 1999). This long history of interaction includes hunting, fishing, and trapping

which result in the direct removal of wild animals. Darimont et al.'s (2015) analysis revealed that

humans have killed a larger biomass of prey species than most other predators in their respective

ecosystems. Given the efficiency and scale at which humans kill animals, potential prey should

respond to human hunters very strongly (Clinchy et al. 2016; Crawford et al. 2022). Our meta-

analysis showed that, across taxa and geographical locations, wild animals significantly altered

both their foraging and vigilance behaviours in areas where lethal interactions with humans

occur. In line with the foraging – vigilance trade off (Brown & Kotler 2004), we found that

animals reduced foraging and significantly increased vigilance in response to human predators,

akin to the way they would respond to other predators in their environment.

There was, however, some variability in the strength of animal responses to humans with lethal

intent (Figure 4). Moose (*Alces alces*), elk (Cervus elaphus) and fallow deer (*Dama dama*)

showed the strongest reductions in foraging behaviour, elk showed the strongest increase in

vigilance, and roe deer (*Capreolus capreolus*) the strongest reduction in movement. The

magnitude of behavioural change in impala (*Aepyceros melampus*) depended on the study.

Matson et al. (2005) recorded much lower vigilance for impala than Crosmary et al. (2011), who

measured animals approaching water—a context where need and risk intersect. This pattern

highlights how for some species, internal state modulates risk sensitivity, as supported by

broader research on state-dependent anti-predator behaviour (Wirsing et al. 2021; Zanette et al.

2020).

Similar to lethal interactions with humans, active interactions with humans that are of non-lethal

intent also elicited an overall positive effect on wild animal vigilance behaviour (Figure 5). The

aggregate effect also indicated an increase in foraging in the presence of non-lethal humans;

however, this effect was neither large nor significant (Figure 5). According to the risk –

disturbance hypothesis, animals are expected to perceive even non-lethal human disturbance

similarly to the threat of predation (Frid & Dill 2002). Overall, we found that wild animals

responded to non–lethal active human interactions to a lesser extent than lethal disturbance

(Figure 5). Variation in the strength of response to non-lethal active interactions was also lower

than that seen for lethal interactions, indicating a more consistently muted response to human

presence across taxa.

Human altered environments have variable effects

Passive human interactions, such as roads and human settlements, did not significantly influence

wild animal foraging or movement behaviour. However, vigilance behaviours were significantly

lower in response to human altered environments (Figure 5). While this may, in part, reflect the

paucity of studies that directly assess these effects, we nevertheless found substantial variation in

behavioural responses to passive human interactions across studies (Figure 4). Unlike active

lethal interactions, which typically produced reduced or non-significant effects, responses to

passive interactions were often large in magnitude, and highly variable.

Some wild animals appear to benefit from passive human interactions. The human–shield hy-

pothesis proposes that human presence can alter interspecific interactions by deterring predators

or dominant competitors and creating safe areas for prey or subordinate competitors (sensu

Berger 2007, Moller 2008; Proudman et al. 2021; Shannon et al. 2014; Abrahms et al. 2025).

Roads, for example, remove cover typically used by ambush predators, paradoxically making

areas adjacent to roads "safer" for certain prey species (Berger 2007; Moll et al. 2017). From our

meta-analysis, evidence of a positive effect of passive human interactions can be seen in the in-

creased foraging and reduced vigilance behaviours of prairie dogs (*Cynomys ludovicianus*),

white-footed mice (*Peromyscus leucopus*), and Verreaux's sifakas (*Propithecus verreauxi*)

(Shannon et al. 2014; Giordano et al. 2022; Chen-Kraus et al. 2022; Figure 4). It should be noted,

though, that roads can also be ecological traps, offering new foraging opportunities while in-

creasing mortality risks (Smith et al. 2021).

Human-altered environments may not always confer advantages. In some cases, human develop-

ment increases predation risk by opening habitats and making it more difficult for prey to min-

imise detection and escape (Smith et al. 2016). Human infrastructure also poses direct threats to

wildlife via vehicle collisions, poisoning, or conflict with livestock interests (Harris et al. 2018;

van der Kolk et al. 2024; Loss et al. 2015). Consistent with these risks, several studies have doc-

umented reductions in foraging and heightened vigilance of elk (*Cervus elaphus*) near roads and

other passive disturbances (Ciuti et al. 2012; Shannon et al. 2014; Jayakody et al. 2008). Forag-

ing responses in pronghorn (*Antilocapra americana*) were mixed, potentially attributable to dif-

ferences in study design. For instance, Gavin et al. (2006) employed distance from roads as a

continuous control variable, while Shannon et al. (2014) used categorical comparisons between

disturbed and undisturbed road sites. Ultimately, whether animals perceive human altered envi-

ronments as refuges or threats varies greatly by species and ecological context (Wirsing et al.

2021; Smith et al. 2021).

Comparing lethal and non-lethal human interactions within species

The biggest challenge in understanding the relative impacts of lethal and non-lethal human inter-

actions on wild animals is the fact that only three studies tested these effects in the same species.

An especially informative example from our meta-analysis is elk (*Cervus elaphus*), the only

species in our dataset for which foraging and vigilance behaviours were assessed under both

lethal and non-lethal human disturbance conditions (Appendix A: Table A1; Figure 4). Elk con-

sistently reduced foraging and increase vigilance, regardless of interaction type, and in some

contexts, displayed a pronounced increase in vigilance even without a concomitant change in for-

aging (Jayakody et al. 2008). Additionally, lethal interactions with humans induced a marked

increase in elk movement rates (Ciuti et al. 2012; Shannon et al. 2014). Such strong anti-predator

behavioural responses towards humans, regardless of whether they have lethal or non-lethal in-

tent, may reflect a long history of hunting pressure on elk (Ciuti et al. 2012).

In addition to elk, several other species in our dataset—including willow grouse, brown bear,

whale shark, roe deer, fallow deer, and badgers—have documented histories of lethal or

antagonistic interactions with humans (See appendix A: Table A1). Yet, the nature and strength

of their behavioural responses were highly variable (Figure 4). These examples underscore that a

mere history of lethal encounters is insufficient to predict how animals will respond to humans in

their environment. The presence and magnitude of behavioural responses to humans may depend

on a number of additional factors such as intensity of hunting or persecution, nature of human

activities in the habitat, duration since cessation of hunting or persecution and current forms of

human interaction (Smith et al. 2021, 2024).

The criteria of our meta-analysis precluded us from including some other interesting examples of

when lethal and non-lethal interactions with humans combine to influence animal behaviour. For

example, moose (*Alces alces*) in Norway avoid human infrastructure in hunting areas, suggesting

a synergistic effect of different human disturbance types (Mehlhoop et al. 2002). Such changes in

behaviour may have compounding negative effects on wild animals. Roe deer and wild boar

change habitat use in response to hunting pressure, which in turn makes them more vulnerable to

death from vehicle collisions (Zuberogoitia et al. 2014). These multiple-exposure scenarios high-

light a critical area for future research that our meta-analysis could not fully address.

Behavioural trade-offs within individuals

About half of the studies in our meta-analysis tested the effects of human interactions on

foraging and vigilance simultaneously (n = 18). This allows us to comment on direct foraging–

vigilance trade-offs (Brown and Kotler, 1991). Qualitatively, there is good consensus among

these studies that animals reduce foraging and proportionally increase vigilance in response to

both lethal and active non-lethal interactions with humans. Similarly, there is consensus that

animals increase foraging and reduce vigilance in response human altered habitats (Appendix A:

Table A3).

While our meta-analytic approach enables unit-less comparison across studies (Noble et al.

2022), this does not change the fact that different behavioural metrics may be indicative of

different underlying processes, and different study designs may yield different conclusions. For

example, movement and space use in response to passive human disturbances seem to be

governed by distinct behavioural processes. Wild boar (*Sus scrofa*) in urbanised environments

occupy smaller home ranges than those in undisturbed forests, yet move greater distances within

these ranges, reflecting both reduced spatial requirements and increased activity (Podgorski et al.

2013; Zanette et al. 2020). Another notable example is the contrasting movement responses of

roe deer (*Capreolus capreolus*) to lethal human presence (Figure 4). However, this discrepancy

is plausibly attributable to methodological differences between studies. Specifically, Grignolio et

al. 2011 used a temporal control to test the effect of hunting, whereas Picardi et al. (2009)

employed a spatial control. These mixed responses indicate how different behavioural metrics

and designs can result in different responses to human disturbance within the same species.

Perception of risk and the role of habituation

Contrary to expectation, some animals showed little change in behaviour in response to lethal

interactions with humans. Despite being hunted, puma (*Puma concolor*, Smith et al. 2017),

badgers (*Meles meles*, Clinchy et al. 2016), and impala (*Aepyceros melampus*, Matson et al.

2005), which were included in our meta-analysis, did not change either foraging or vigilance

behaviour or both (Figure 4). These species may be under-reacting to humans in their

environment, and their lack of a response may reflect the nature of how these animals are killed

and the ability for animals to identify and assess human predator cues (Smith et al. 2021). For

example, animals may not respond the same way to humans with guns and traps as they provide

different cues. On the other hand, some animals seem to overreact, such as those that

significantly change their behaviours in response to non-lethal human interactions, leading

potentially to high energy investment or chronic stress (e.g. Blumstein 2003; Westekemper et al.

2018; Larson et al. 2016; Nunes et al. 2019 review: Larson et al. 2016).

Animals may also habituate to human encounters or human-induced changes in the environment,

which may explain the subdued or non-significant responses to active non-lethal human interac-

tions. In our meta-analysis, blackbirds (*Turdus merula*), thinhorn sheep (*Ovis dalli*), whale

sharks (*Rhincodon typus*), semipalmated plovers (*Charadrius semipalmatus*) and western sand-

pipers (*Calidris mauri)* showed no change in either foraging or vigilance in response to active

human activity in their environment (Fernandez-Juricic et al., 2001, Loehr et al. 2005, Montero-

quintana et al., 2020, Murchison et al., 2016, respectively). Developmental stage may also medi-

ate responsiveness: for instance, while adult thinhorn sheep increase vigilance in response to

non-lethal disturbance, juveniles do not; and only juveniles show flexibility in their foraging be-

haviour (Loehr et al. 2005). Similar patterns occur in hooded cranes (*Grus monacha*), which vary

vigilance but not foraging responses across age classes (Li et al. 2015). These observations sug-

gest habituation, wherein repeated exposure to non-lethal humans results in dampened anti-

predator responses over time.

A recent meta-analysis of camera trapping data from 21 countries during the COVID-19 pan-

demic also suggests that habituation may explain significant variation in animal responses to

non-lethal human presence (Burton et al., 2024). For example, Uchida & Blumstein (2021) found

that yellow-bellied marmots (*Marmota flaviventer*) in highly disturbed forests grew habituated to

humans after repeated exposure and reduced their flight initiation distances. Similar examples

have been cited from tourist interactions with animals in the African savannah (Knight 2009),

killer whale and Hector's dolphin interactions with whale watchers (Bejder et al. 1999; Williams

et al. 2002) and penguin interactions with researchers (Shelton et al. 2004).

As with non-lethal active human interactions, not all taxa included in this meta-analysis re-

sponded to passive human interactions: chipmunks (*Tamias striatus*) and Gaspésie caribou

(*Rangifer tarandus caribou*), for example, show limited behavioural shifts to passive human

presence for reasons that are unclear (Giordano et al. 2022; Lesmerises et al., 2017). The hetero-

geneity in risk perception and habituation contributes to the overall variation observed in be-

havioural responses across taxa and highlights the need for further research encompassing a

broader diversity of species and behavioural metrics (Figure 1).

Context dependent variability in behavioural responses to humans

The results of our meta-analysis reveal the importance of context in how animals respond to

humans. Ecological and biological factors, such as trophic level, body size, and group size can

influence the perception of risk by animals. Unfortunately, there were too few studies in the

meta-analysis that enabled statistical evaluation of these effects. These are, however, important

enough to warrant some discussion.

When humans kill top predators, they create new risk that these animals have not previously

experienced. Predators may or may not be equipped to respond to new threats leading to many

extirpations (e.g. wolves in Yellowstone (Laundré et al. 2001), bears in Alaska (Merkle et al.

2013), puma in California and Florida (Nickel et al. 2021). Predators may be overly cautious

around humans, leading to greater missed opportunities for fitness-enhancing activities. For

example, Suraci et al. (2019) showed that human presence induced puma (*Puma concolor*) to

abandon their kill sites. Based on camera trap data, Burton et al., (2024) showed that carnivores

avoided humans to a greater extent than wild animals belonging to other foraging guilds. Chronic

exposure to humans may thus lead to reduced fitness for risk-averse individuals. Alternatively,

predators may not be cautious enough around humans leading to greater direct mortality (Smith

et al. 2021). The severity and direction of the response may be proportional to the extent and

duration of persecution by humans. The majority of studies we found in our search focused on

primary consumers and not predators (Appendix A), making it difficult to assess trophic effects.

This gap remains an important area for future research.

Like predators, large animals may be disproportionately targeted by humans for economic

reasons and as trophies. Larger and healthy animals are rarely targeted by natural predators and

thus are less likely to invest in anti-predator behaviour, which is similar to apex predators.

Within trophic levels large herbivorous animals have been shown to increase activity in the

presence of humans (Burton et al., 2024). We found no evidence for a consistent effect of body

size on wild animal behavioural responses to human interactions (Appendix E). However when

we look at the responses of individual species across studies (Figure 4), behavioural response to

lethal interactions were largest for large herbivores such as deer (*Dama dama*), elk (*Cervus*

elaphus) and moose (*Alces alces*) (Deer: Pecorella et al. 2016; Elk: Ciuti et al., 2012, Proffitt et

al., 2009, Shannon et al., 2014, Jayakody et al., 2008; Moose: Bhardwaj et al., 2022).

Finally, most species in our dataset are group living to some degree, while only three species

(predators) are considered solitary. Group living typically reduces predation risk through the

dilution and many-eyes effects (Foster & Treherne 1981; Pulliam 1973). Whether grouping

similarly alters responses to humans compared to natural predators remains unclear and could

not be evaluated here.

Conclusion

Here, we have presented an extensive synthesis of the available literature on the effect of human

interactions on animal behaviour. Although our study does not bridge the gap between behaviour

and the resulting demographic consequences for populations (see Smith et al. 2024), it is

important to note that the demographic effects of non-consumptive effects (NCEs) remain

contentious (Sheriff et al. 2020). Future research should aim to more explicitly link human

interactions, behavioural changes, fitness consequences, and demographic outcomes.

Our general conclusions and suggestions for future research are below:

1. Both lethal and non–lethal interactions with humans elicit changes in wild animal

behaviour, but lethal interactions have a more consistent and larger effect.

2. There is considerable variation in the response of animals to the different types of human

interactions that has yet to be disentangled. Factors such as frequency of interactions,

history of hunting, and taxon identity may modulate the response of animals to human

disturbance.

3. Studies that directly compare the effects of lethal interactions and non-lethal within the

same species are necessary to determine whether and when animals can infer the intent of

humans.

4. Future research may benefit from contrasting humans as lethal predators with natural

predators in the same space and across trophic levels, given the potential for

compounding effects on animals and ecosystems.

581 **Acknowledgements**

582 Funding for this research was provided by the Prime Ministers Research Fellowship to SD with

583 some support from DBT-Wellcome Trust India Alliance grant (IA/I/19/2/504639) to MT. We

584 would like to thank Dr. Nitya Mohanty and Dr. Daniel Noble for their invaluable inputs during

585 the drafting of this manuscript.

586 **References**

Abrahms, B., Rafiq, K., Nisi, A.C., Jordan, N.R., Wilson, A.M., Loveridge, A.J., et al. (2025). Intraguild competition mediates human avoidance in an endangered African large carnivore. Proceedings of the Royal Society B: Biological Sciences, 292, 20242835.

Bejder, L., Dawson, S.M. & Harraway, J.A. (1999). Responses by Hector's dolphins to boats and swimmers in Porpoise bay, New Zealand. *Mar. Mammal Sci.*, 15, 738–750.

Benson, J., Mahoney, P. & Patterson, B. (2015). Spatiotemporal variation in selection of roads influences mortality risk for canids in an unprotected landscape. *OIKOS*, 124, 1664–1673.

Berger, J. (2007). Fear, human shields and the redistribution of prey and predators in protected areas. *Biol. Lett.*, 3, 620–623.

Bhardwaj, M., Lodnert, D., Olsson, M., Winsvold, A., Eilertsen, S.M., Kjellander, P., *et al.* (2022). Inducing fear using acoustic stimuli—A behavioural experiment on moose (Alces alces) in Sweden. *Ecol. Evol.*, 12.

Blumstein, D. (2003). Flight-initiation distance in birds is dependent on intruder starting distance. *J. Wildl. Manag.*, 67, 852–857.

Boyle, S.A. & Samson, F.B. (1985). Effects of Nonconsumptive Recreation on Wildlife: A Review. *Wildl. Soc. Bull. 1973-2006*, 13, 110–116.

Brown, J.S. & Kotler, B.P. (2004). Hazardous duty pay and the foraging cost of predation: Foraging cost of predation. *Ecol. Lett.*, 7, 999–1014.

Burton, A.C., Beirne, C., Gaynor, K.M., Sun, C., Granados, A., Allen, M.L., *et al.* (2024). Mammal responses to global changes in human activity vary by trophic group and landscape. *Nat Ecol Evol*, 8, 924–935.

Cappa, F., Giannoni, S. & Borghi, C. (2017). Effects of roads on the behaviour of the largest South American artiodactyl (Lama guanicoe) in an Argentine reserve. *Anim. Behav.*, 131, 131–136.

Chamberlain, S.A., Hovick, S.M., Dibble, C.J., Rasmussen, N.L., Van Allen, B.G., Maitner, B.S., *et al.* (2012). Does phylogeny matter? Assessing the impact of phylogenetic information in ecological meta-analysis. *Ecol. Lett.*, 15, 627–636.

Cinar, O., Nakagawa, S. & Viechtbauer, W. (2022). Phylogenetic multilevel meta-analysis: A simulation study on the importance of modelling the phylogeny. *Methods Ecol. Evol.*, 13, 383–395.

Clinchy, M., Zanette, L.Y., Roberts, D., Suraci, J.P., Buesching, C.D., Newman, C., *et al.* (2016). Fear of the human "super predator" far exceeds the fear of large carnivores in a model mesocarnivore. *Behav. Ecol.*, arw117.

Crawford, D.A., Conner, L.M., Clinchy, M., Zanette, L.Y. & Cherry, M.J. (2022). Prey tells, large herbivores fear the human 'super predator.' *Oecologia*, 198, 91–98.

Crosmary, W., Valeix, M., Fritz, H., Madzikanda, H. & Cote, S. (2012). African ungulates and their drinking problems: Hunting and predation risks constrain access to water. *Anim. Behav.*, 83, 145–153.

Darimont, C.T., Fox, C.H., Bryan, H.M. & Reimchen, T.E. (2015). The unique ecology of human predators. *Science*, 349, 858–860.

Darimont, C.T., Cooke, R., Bourbonnais, M.L., Bryan, H.M., Carlson, S.M., Estes, J.A., et al. (2023). Humanity's diverse predatory niche and its ecological consequences. Commun Biol, 6, 609.

Davis, D., Banks, S., Birtles, A., Valentine, P. & Cuthill, M. (1997). Whale sharks in Ningaloo Marine Park: Managing tourism in an Australian marine protected area. *Tour. Manag.*, 18, 259–271.

Del Hoyo, J., Elliott, A. & Sargatal, J. (1992). *Handbook of the birds of the world*. Lynx edicions Barcelona.

Dill, L.M., Heithaus, M.R. & Walters, C.J. (2003). Behaviourally mediated indirect interactions in marine communities and their conservation implications. *Ecology*, 84, 1151–1157.

Dsouza, Shawn, Kartik Shanker, and Maria Thaker. 2025. "Are Super-Predators Also Super-Scary? A Systematic Review and Meta-Analysis of Animal Responses to Human Interactions. [Dataset]" Zenodo. https://doi.org/10.5281/zenodo.15099679.

Fernández-Juricic, Miguel Clavijo, María Dolores Jiménez, Eloy Asensio, & Elena Lucas. (2001). Bird tolerance to human disturbance in urban parks of Madrid (Spain): Management implications.

Estes, J.A., Terborgh, J., Brashares, J.S., Power, M.E., Berger, J., Bond, W.J., *et al.* (2011). Trophic Downgrading of Planet Earth. *Science*, 333, 301–306.

Foo, Y.Z., O'Dea, R.E., Koricheva, J., Nakagawa, S. & Lagisz, M. (2021). A practical guide to question formation, systematic searching and study screening for literature reviews in ecology and evolution. *Methods Ecol. Evol.*, 12, 1705–1720.

Foster, W.A. & Treherne, J.E. (1981). Evidence for the dilution effect in the selfish herd from fish predation on a marine insect. *Nature*, 293, 466–467.

Frid, A. & Dill, L. (2002). Human-caused disturbance stimuli as a form of predation risk. *Conserv. Ecol.*, 6.

Froese, R. & Pauly, D. (2010). FishBase.

Gaynor, K.M., Hojnowski, C.E., Carter, N.H. & Brashares, J.S. (2018). The influence of human disturbance on wildlife nocturnality. *Science*, 360, 1232–1235.

Gaynor, K.M., Wooster, E.I.F., Martinig, A.R., Green, J.R., Chhen, A., Cuadros, S., et al. (2025). The Human Shield Hypothesis: Does Predator Avoidance of Humans Create Refuges for Prey? Ecology Letters, 28, e70138.

Grignolio, S., Merli, E., Bongi, P., Ciuti, S. & Apollonio, M. (2011). Effects of hunting with hounds on a non-target species living on the edge of a protected area. *Biol. Conserv.*, 144, 641–649.

Harrer, M., Cuijpers, P., Furukawa, T.A. & Ebert, D.D. (2021). *Doing Meta-Analysis With R: A Hands-On Guide*. 1st edn. Chapman & Hall/CRC Press, Boca Raton, FL and London.

Harris, C., Thomas, L., Falcone, E., Hildebrand, J., Houser, D., Kvadsheim, P., *et al.* (2018). Marine mammals and sonar: Dose-response studies, the risk-disturbance hypothesis and the role of exposure context. *J. Appl. Ecol.*, 55, 396–404.

Herberstein, M.E., McLean, D.J., Lowe, E., Wolff, J.O., Khan, M.K., Smith, K., *et al.* (2022). AnimalTraits - a curated animal trait database for body mass, metabolic rate and brain size. *Sci. Data*, 9, 265.

Jayakody, S., Sibbald, A., Gordon, I. & Lambin, X. (2008). Red deer Cervus elephus vigilance behaviour differs with habitat and type of human disturbance. *Wildl. Biol.*, 14, 81–91.

Knight, J. (2009). Making Wildlife Viewable: Habituation and Attraction. *Soc. Anim.*, 17, 167–184.

van der Kolk, H.-J., Smit, C.J., Allen, A.M., Ens, B.J. & van de Pol, M. (2024). Frequency-dependent tolerance to aircraft disturbance drastically alters predicted impact on shorebirds. *Ecol. Lett.*, 27, e14452.

Ladle, A., Avgar, T., Wheatley, M., Stenhouse, G., Nielsen, S. & Boyce, M. (2019). Grizzly bear response to spatio-temporal variability in human recreational activity. *J. Appl. Ecol.*, 56, 375–386.

Larson, C.L., Reed, S.E., Merenlender, A.M. & Crooks, K.R. (2016). Effects of Recreation on Animals Revealed as Widespread through a Global Systematic Review. *PLOS ONE*, 11, e0167259.

Laundré, J.W., Hernández, L. & Altendorf, K.B. (2001). Wolves, elk, and bison: reestablishing the "landscape of fear" in Yellowstone National Park, U.S.A. *Can. J. Zool.*, 79, 1401–1409.

Lima, S.L. & Dill, L.M. (1990). Behavioral decisions made under the risk of predation: a review and prospectus. *Can. J. Zool.*, 68, 619–640.

Lintulaakso, K. (2021). MammalBase — Database of traits, measurements and diets of the species in class Mammalia.

Loehr, J., Kovanen, M., Carey, J., Hogmander, H., Jurasz, C., Karkkainen, S., *et al.* (2005). Gender- and age-class-specific reactions to human disturbance in a sexually dimorphic ungulate. *Can. J. Zool.*, 83, 1602–1607.

Loss, S.R., Will, T. & Marra, P.P. (2015). Direct Mortality of Birds from Anthropogenic Causes. *Annu. Rev. Ecol. Evol. Syst.*, 46, 99–120.

Matson, T., Goldizen, A. & Putland, D. (2005). Factors affecting the vigilance and flight behaviour of impalas. *SOUTH AFRICAN JOURNAL OF WILDLIFE RESEARCH*, 35, 1-11

Mehlhoop, A., Van Moorter, B., Rolandsen, C., Hagen, D., Granhus, A., Eriksen, R., *et al.* (2022). Moose in our neighborhood: Does perceived hunting risk have cascading effects on tree performance in vicinity of roads and houses? *Ecol. Evol.*, 12.

Merkle, J., Robinson, H., Krausman, P. & Alaback, P. (2013). Food availability and foraging near human developments by black bears. *J. Mammal.*, 94, 378–385.

Michonneau, F., Brown, J.W. & Winter, D.J. (2016). rotl: an R package to interact with the Open Tree of Life data. *Methods Ecol. Evol.*, 7, 1476–1481.

Mikolajewicz, N. & Komarova, S.V. (2019). Meta-Analytic Methodology for Basic Research: A Practical Guide. *Front. Physiol.*, 10, 203.

Moller, A. (2008). Flight distance of urban birds, predation, and selection for urban life. *Behav. Ecol. Sociobiol.*, 63, 63–75.

Montero-Quintana, A., Vazquez-Haikin, J., Merkling, T., Blanchard, P. & Osorio-Beristain, M. (2020). Ecotourism impacts on the behaviour of whale sharks: an experimental approach. *ORYX*, 54, 270–275.

Murchison, C., Zharikov, Y. & Nol, E. (2016). Human Activity and Habitat Characteristics Influence Shorebird Habitat Use and Behavior at a Vancouver Island Migratory Stopover Site. *ENVIRONMENTAL MANAGEMENT*, 58, 386–398.

Nakagawa, S., Lagisz, M., Jennions, M.D., Koricheva, J., Noble, D.W.A., Parker, T.H., *et al.* (2022). Methods for testing publication bias in ecological and evolutionary meta-analyses. *Methods Ecol. Evol.*, 13, 4–21.

Nevin, O. & Gilbert, B. (2005). Perceived risk, displacement and refuging in brown bears: positive impacts of ecotourism? *Biol. Conserv.*, 121, 611–622.

Newey, P. (2007). Foraging behaviour of the Common Myna (Acridotheres tristis) in relation to vigilance and group size. *EMU*, 107, 315–320.

Nickel, B., Suraci, J., Nisi, A. & Wilmers, C. (2021). Energetics and fear of humans constrain the spatial ecology of pumas. *Proc. Natl. Acad. Sci. U. S. A.*, 118.

Noble, D.W.A., Pottier, P., Lagisz, M., Burke, S., Drobniak, S.M., O'Dea, R.E., *et al.* (2022). Meta-analytic approaches and effect sizes to account for 'nuisance heterogeneity' in comparative physiology. *J. Exp. Biol.*, 225, jeb243225.

Nunes, J., Blumstein, D., Giglio, V., Barros, F. & Quimbayo, J. (2019). Reef fish antipredator behaviour in remote islands does not reflect patterns seen in coastal areas. *Ethol. Ecol. Evol.*, 31, 557–567.

O'Dea, R.E., Lagisz, M., Jennions, M.D., Koricheva, J., Noble, D.W.A., Parker, T.H., *et al.* (2021). Preferred reporting items for systematic reviews and meta-analyses in ecology and evolutionary biology: a PRISMA extension. *Biol. Rev.*, 96, 1695–1722.

OpenTreeOfLife, Redelings, B., Reyes, L.L.S., Cranston, K.A., Allman, J., Holder, M.T., *et al.* (2019). Open tree of life synthetic tree.

Oriol-Cotterill, A., Valeix, M., Frank, L., Riginos, C. & Macdonald, D. (2015). Landscapes of Coexistence for terrestrial carnivores: the ecological consequences of being downgraded from ultimate to penultimate predator by humans. *OIKOS*, 124, 1263–1273.

Ouzzani, M., Hammady, H., Fedorowicz, Z. & Elmagarmid, A. (2016). Rayyan—a web and mobile app for systematic reviews. *Syst. Rev.*, 5, 210.

Page, M.J., McKenzie, J.E., Bossuyt, P.M., Boutron, I., Hoffmann, T.C., Mulrow, C.D., *et al.* (2021). The PRISMA 2020 statement: an updated guideline for reporting systematic reviews. *BMJ*, 372, n71.

Papouchis, Francis J. Singer, & William B. Sloan. (2001). Responses of Desert Bighorn Sheep to Increased Human Recreation. *J. Wildl. Manag.*

Pecorella, I., Ferretti, F., Sforzi, A. & Macchi, E. (2016). Effects of culling on vigilance behaviour and endogenous stress response of female fallow deer. *Wildl. Res.*, 43, 189–196.

Picardi, S., Basille, M., Peters, W., Ponciano, J., Boitani, L. & Cagnacci, F. (2019). Movement responses of roe deer to hunting risk. *J. Wildl. Manag.*, 83, 43–51.

Podgorski, T., Bas, G., Jedrzejewska, B., Sonnichsen, L., Sniezko, S., Jedrzejewski, W., *et al.* (2013). Spatiotemporal behavioral plasticity of wild boar (Sus scrofa) under contrasting conditions of human pressure: primeval forest and metropolitan area. *JOURNAL OF MAMMALOGY*, 94, 109–119.

Proudman, N., Churski, M., Bubnicki, J., Nilsson, J. & Kuijper, D. (2021). Red deer allocate vigilance differently in response to spatio-temporal patterns of risk from human hunters and wolves. *Wildl. Res.*, 48, 163–174.

Pulliam, H.R. (1973). On the advantages of flocking. *J Theor Biol*, 38, 419–422.

R Core Team. (2024). *R: a language and environment for statistical computing* (manual). R Foundation for Statistical Computing, Vienna, Austria.

Shannon, G., Cordes, L., Hardy, A., Angeloni, L. & Crooks, K. (2014). Behavioral Responses Associated with a Human-Mediated Predator Shelter. *PLOS ONE*, 9.

Shelton, E., Higham, J. & Seddon, P. (2004). Habituation, penguin research and ecotourism: Some thoughts from left field. *N. Z. J. Zool.*, 31.

Sheriff, M. J., & Thaler, J. S. (2014). Ecophysiological effects of predation risk; An integration across disciplines. Oecologia, 176, 607–611. https://doi.org/10.1007/s0044 2-014-3105-5

Sheriff, M.J., Peacor, S.D., Hawlena, D. & Thaker, M. (2020). Non-consumptive predator effects on prey population size: A dearth of evidence. *J. Anim. Ecol.*, 89, 1302–1316.

Sih, A. (1984). The Behavioral Response Race Between Predator and Prey. *Am. Nat.*, 123, 143–150.

Sih, A. (2013). Understanding variation in behavioural responses to human-induced rapid environmental change: a conceptual overview. *Anim. Behav.*, Special Issue: Behavioural Plasticity and Evolution, 85, 1077–1088.

Smith, J., Suraci, J., Clinchy, M., Crawford, A., Roberts, D., Zanette, L., *et al.* (2017). Fear of the human "super predator" reduces feeding time in large carnivores. *Proc. R. Soc. B-Biol. Sci.*, 284.

Smith, J.A., Gaynor, K.M. & Suraci, J.P. (2021). Mismatch Between Risk and Response May Amplify Lethal and Non-lethal Effects of Humans on Wild Animal Populations. *Front. Ecol. Evol.*, 9, 604973.

Smith, J.A., McDaniels, M.E., Peacor, S.D., Bolas, E.C., Cherry, M.J., Dorn, N.J., *et al.* (2024). Population and community consequences of perceived risk from humans in wildlife. *Ecol. Lett.*, 27, e14456.

Smith, J.A., Wang, Y. & Wilmers, C.C. (2016). Spatial characteristics of residential development shift large carnivore prey habits. *J. Wildl. Manag.*, 80, 1040–1048.

Suraci, J., Smith, J., Clinchy, M., Zanette, L. & Wilmers, C. (2019). Humans, but not their dogs, displace pumas from their kills: An experimental approach. *Sci. Rep.*, 9.

Treves, A. & Naughton-Treves, L. (1999). Risk and opportunity for humans coexisting with large carnivores. *J. Hum. Evol.*, 36, 275–282.

Uchida, K. & Blumstein, D. (2021). Habituation or sensitization? Long-term responses of yellow-bellied marmots to human disturbance. *Behav. Ecol.*, 32, 668–678.

Viechtbauer, W. (2010). Conducting meta-analyses in R with the metafor package. *Journal of Statistical Software*, 36, 1–48.

Westekemper, Horst Reinecke, Johannes Signer, et al. (2018). Stay on trails - effects of human recreation on the spatiotemporal behaviour of red deer Cervus elaphus in a German National Park. Wildl. Biol.

Williams, R., Trites, A.W. & Bain, D.E. (2002). Behavioural responses of killer whales (Orcinus orca) to whale-watching boats: opportunistic observations and experimental approaches. *J. Zool.*, 256, 255–270.

Wirsing, A.J., Heithaus, M.R., Brown, J.S., Kotler, B.P. & Schmitz, O.J. (2021). The context dependence of non-consumptive predator effects. *Ecol. Lett.*, 24, 113–129.

Zanette, L. & Clinchy, M. (2020). Ecology and Neurobiology of Fear in Free-Living Wildlife. *ANNUAL REVIEW OF ECOLOGY, EVOLUTION, AND SYSTEMATICS, VOL 51, 2020*, 51, 297–318.

Zuberogoitia, I., del Real, J., Torres, J.J., Rodríguez, L., Alonso, M. & Zabala, J. (2014). Ungulate vehicle collisions in a peri-urban environment: consequences of transportation infrastructures planned assuming the absence of ungulates. *PLoS One*, 9, e107713.

Figures and tables

Figure 1: The different scenarios in which humans interact with wild animals and the

hypothesised consequences on animal behaviour. We expect a gradient of response with the

strongest behavioural changes of animals in areas where interactions with humans are of lethal

intent (i.e. hunting and fishing), followed by non-lethal but active interactions (i.e. ecotourism)

and non-lethal but passive interactions (i.e. roads). Values below 0 represent a reduction in the

behaviour while those above 0 represent an increase relative to animals in areas without that

human activity.

Figure 2: Geographical distribution of studies included in the systematic review.

Figure 3: Distribution of studies included in the meta-analysis. Shown are the various human ac-

tivities that were considered under lethal, active non-lethal and passive non-lethal as well as the

specific behavioural measurements of wild animals within the broad categories of foraging, vigi-

lance and movement. Number of studies at each node are indicated in brackets.

Figure 4: The effect of human interactions (lethal, active interaction, passive interaction) across

taxa and trophic levels. Negative values on the x-axis indicate a reduction in that behaviour,

whereas positive values indicate an increase in that behaviour, relative to control conditions

where no human activity is present.

Figure 5: Summary effect of human interactions on foraging, movement, and vigilance

behaviour across studies included in this meta-analysis. For animal movement, active non-lethal

interactions were excluded due to the low sample size.