Auditory perception and neural representation of temporal features are altered by age but not by cochlear synaptopathy

Friederike Steenken [1,2,3¶], Rainer Beutelmann [1,2,3¶], Henning Oetjen [1,2,3], Christine Köppl [1,2,3], Georg M. Klump [1,2,3*]

1 Cluster of Excellence "Hearing4all", Carl von Ossietzky University Oldenburg, 26111 Oldenburg, Germany

2 Research Centre Neurosensory Science, Carl von Ossietzky University Oldenburg, 26111 Oldenburg, Germany

3 Department of Neuroscience, School of Medicine and Health Science, Carl von Ossietzky University Oldenburg, 26111 Oldenburg, Germany

¶ Shared first author

* Corresponding author

Email: georg.klump@uol.de

Running Title: TFS processing and perception

Author contributions: G.M.K. and C.K. designed research; F.S. and H.O. performed research; R.B. contributed analytic tools; F.S., R.B., H.O., and G.M.K. analyzed data; F.S., R.B., H.O., C.K., and G.M.K. wrote the paper.

Keywords: temporal fine structure, temporal envelope, auditory nerve, perception, gerbil, age-related hearing loss, synapse loss

Abstract

Age-related hearing loss is a complex phenomenon. The earliest-onset degenerative event is the gradual loss of neural connections between cochlea and auditory brainstem. To probe for perceptual deficits that might arise from this loss, cochlear synaptopathy was induced pharmacologically in young-adult gerbils, which were then tested in a challenging listening task for the perception of temporal fine structure. Treated gerbils behaved no differently than normal-hearing, young-adult animals. In contrast, old gerbils, which typically express many cochlear and central-neural pathologies, showed impaired perception. To probe for the underlying mechanisms, single-unit responses were obtained from the auditory nerve to the same test stimuli. Responses from old gerbils showed no impairment in temporal locking to the stimulus fine structure. However, responses were significantly more driven by slower temporal fluctuations of the stimulus envelope, suggesting that the central auditory system may be unable to extract the relevant information for discrimination from such altered inputs.

Introduction

Many elderly people have difficulties understanding speech in noisy environments, despite having normal tone-detection thresholds in quiet. To explain this condition of compromised supra-threshold perception, for example of speech in background noise (e.g., Füllgrabe et al. 2015, Gómez-Álvarez et al. 2023, Lopez-Poveda 2014, Moore 2014, Parthasarathy et al. 2020), despite an absence of detectable hearing deficits in the pure-tone audiogram in quiet, is currently an important challenge in auditory neuroscience. It has been suggested to be associated with a reduced number of synapses between auditory-nerve (AN) fibers and inner hair cells (IHCs), which is commonly referred to as cochlear synaptopathy (e.g., Bharadwaj et al. 2014, Kujawa and Liberman 2009, Liberman 2017, Liberman and Kujawa 2017). Synaptopathy is one of the aging processes in the cochlea, and, in addition, can be caused by excessive noise exposure, both accumulating during a human's life time (e.g., Liberman et al. 2016, Wu et al. 2019, but see Johannesen et al. 2019). Animal experiments demonstrate that exposure to loud sounds can produce such synaptopathy, even without permanent loss in hearing sensitivity (e.g., Kujawa and Liberman 2009). Furthermore, the number of functional synapses decreases with increasing age (e.g., gerbil: Bovee et al. 2024, Gleich et al. 2016, Steenken et al. 2021; mouse: Jeng et al. 2020, Parthasaraty and Kujawa 2018, Sergeyenko et al. 2013; rat: Cai et al. 2018; human: Viana et al. 2015).

Synaptopathy is postulated to be causal for compromised processing of temporal stimulus features (e.g., Parthasarathy and Kujawa 2018), especially of the temporal fine structure (TFS) of sounds (reviewed in Moore 2019), resulting in difficulties of speech comprehension in background noise. Specific psychophysical tests that evaluate the performance to

distinguish between stimuli that differ in TFS have been developed. These tasks are then expected to be sensitive to synaptopathy (for review see Moore 2014). The most commonly used test is the TFS1 test (Moore and Sek 2009) which probes the discrimination of harmonic and inharmonic tone complexes that strongly differ in TFS but only marginally in the envelope of the waveform. The previously found link between diminished perception of stimuli differing in TFS (Moore et al. 2006) and sensory-neural hearing loss in human patients has also been found in TFS1 test results (Mathew et al. 2015). Furthermore, a correlation between age and TFS1 sensitivity has been observed (e.g., Eipert et al. 2019, Füllgrabe et al. 2015, Moore et al. 2012).

Whereas post-mortem studies of human subjects have provided evidence for age-related synapse loss (Viana et al. 2015, Wu et al. 2019), it has been impossible to relate this loss to deficits in supra-threshold sound perception of living subjects. Animal studies have suggested a close correlation between the number of surviving synapses and supra-threshold amplitude growth of auditory brainstem responses (ABRs) or compound action potentials (CAPs; e.g., Bourien et al. 2014, Bramhall et al. 2018, Buran et al. 2010, Kujawa and Liberman 2009, Lin et al. 2011, Sergeyenko et al. 2013, Shi et al. 2016). A number of attempts have been made to infer the synapse loss in humans from non-invasive physiological measures, such as ABR, and psychophysical measures of the processing of sounds. However, even studies involving large samples of human subjects have not produced strong evidence that synaptopathy in humans has functional consequences for supra-threshold temporal perception (e.g., Bramhall et al. 2019, Carcagno and Plack 2022, Guest et al. 2018). Attempts to relate life-time noise exposure of human subjects to their supra-threshold perceptual deficits have also not been conclusive (e.g., Prendergast et al. 2017, 2019, Füllgrabe et al. 2020). Controlled animal studies are one strategy for testing the influence of synapse loss on discrimination of complex sounds that differ in TFS. In those, synaptopathy can be experimentally manipulated and the degree of cochlear dysfunction can be related to neural responses and performance in behavioral tests, using the same stimuli. With this approach we can test the hypothesis that synaptopathy negatively affects the discrimination of stimuli that differ in their TFS.

Animal studies relating behavioral measures of performance in response to a variety of stimuli to synaptopathy have so far also provided mixed results (reviewed in Henry 2022). An investigation by Henry and Abrams (2021), which experimentally induced synaptopathy in budgerigars by kainic acid infusions at the round window, failed to demonstrate a relationship between the amplitude of the CAP (taken as a measure of cochlear synaptopathy) and tone-detection thresholds in noise. Studies in Macaque monkeys that produced a noise-induced hearing loss (NIHL) by acoustic trauma demonstrated differences between normal-hearing and acoustically traumatized monkeys for a) tone detection in noise (Burton et al. 2020), b)

masking release for tone detection in modulated maskers compared to unmodulated maskers, and c) spatial release from masking for tone detection in noise (Mackey et al. 2021). However, these studies used high trauma levels and long exposure durations. Thus, effects were not limited to synapse loss at the IHC but included both outer hair cell (OHC) and IHC damage. Studies in chinchilla AN fibers, involving normal-hearing subjects and subjects with NIHL, did reveal differences in the neural envelope representation but failed to demonstrate differences in the neural representation of TFS that match the psychophysical effects observed in human subjects (e.g., Henry and Heinz 2012, Kale and Heinz 2010, Kale et al. 2014).

To clarify the relation between synaptopathy, AN fiber responses, and behavior, we embarked on a study in Mongolian gerbils (*Meriones unguiculatus*) that combined behavioral measurements of sensitivity in the TFS1 test and the representation of harmonic and inharmonic complex tones with the same fundamental frequency (f_0) by AN fibers. In addition, synaptopathy of many of the experimental subjects was characterized by directly counting the numbers of functional synapses on IHCs, and these measures were related to sensitivity in the TFS1 test. Because compromised performance in the TFS1 test has been observed in elderly humans, we compared the perception of young-adult and old gerbils. In order to experimentally induce synaptopathy in young-adult gerbils and thus dissociate the effects of synaptopathy from effects of age, we treated young-adult gerbils with an infusion of ouabain at the round window. This treatment has been shown to reduce the number of synapses on gerbil IHCs (Bourien et al. 2014). We observed a lower perceptual sensitivity of old compared to young-adult gerbils in the TFS1 test. However, contrary to our hypothesis, ouabain treatment did not affect the performance in the TFS1 task, and the responses of AN fibers that survived ouabain treatment did not differ from those of gerbils that were not treated.

Results

Synapse loss was similar in old gerbils and gerbils treated with a high dosage of ouabain.

Synapses were counted at the cochlear location corresponding to 2 kHz in 54 gerbils, divided into the five gerbil groups, as listed in Table 1, row 4. Young-adult, untreated gerbils had, on average, 23 (SE ±0.5) synapses per IHC (Fig. 1). Synapse numbers were significantly different between gerbil groups (univariate ANOVA, [$F_{(4,49)} = 5.496$, $p = 0.001$]). Compared to untreated young adults, the average number of synapses per IHC in gerbils treated with 70 µM ouabain, and in old gerbils, was significantly reduced, to 19 (SE ±1.4) and 19 (SE ±0.7), respectively (Bonferroni corrected pairwise comparisons: young-adult vs. 70 µM

[p = 0.011]; young-adult vs. old [p = 0.002]). No significant difference between ears treated

with a high dose of ouabain and old ears was found, indicating that both animal groups had

similar degrees of synaptopathy in the region of interest. In surgery-only ears and ears

treated with 40 µM ouabain, average synapse numbers (surgery only: 23 (SE ±0.5), 40 µM

ouabain: 22 (SE ±0.7)) were not significantly different from untreated young-adult ears,

suggesting no effect of the surgical procedure itself and no effect of the lower ouabain dose.

These results were also consistent across more basal cochlear locations (up to 8 kHz

equivalent frequency, see Supplemental Materials). Based on this, we pooled the single-unit

data from surgery-only and 40 µM ouabain treated gerbils into a "sham" group. The gerbils

treated with 70 µM ouabain are henceforth referred to simply as "ouabain". There were no

significant differences in synapse numbers between gerbils tested behaviorally and the other

gerbils from the corresponding groups not tested behaviorally (2-way ANOVA: factor

inclusion in behavior [F(1,44) = 0.330, p = 0.568] and no interactions between inclusion and

gerbil groups [F(4,44) = 1.179, p = 0.333]).

Figure 1: Synapse number was reduced after ouabain treatment and in old age.

Number of functional synapses per IHC, at the cochlear position corresponding to 2 kHz, for

young-adult gerbils (red, N = 19), surgery-only-treated gerbils (orange, N = 3), gerbils treated

with a low dose (40 µM) of ouabain (green, N = 8), gerbils treated with a high dose (70 µM)

of ouabain (blue, N = 10), and old gerbils (gray, N = 14). Box plots display the median (center

line), mean (white circled dot), 25th and 75th percentiles (upper and lower edges of the

*boxes), and maximum and minimum (whiskers). Significance levels are * p = 0.011 and ***

p = 0.0022.

Threshold sensitivity and frequency tuning were not affected by the synapse loss.

Induced synaptopathy, with a similar dose of ouabain, and causing a similar extent of synapse loss as in the present study, was previously found to not impair threshold sensitivity (assessed via ABR) or OHC function (assessed via distortion-product otoacoustic emissions; Bourien et al. 2014). Here, we assessed CAP thresholds and single-unit AN tuning to test for normal, basic cochlear function in our young-adult, ouabain-treated gerbils (see Table 1 in Methods for sample sizes of AN fibers). CAP thresholds in response to chirp stimuli were obtained from all ouabain- and sham-treated animals before surgery, and again at the start of terminal neurophysiology, and in most cases, for both ears. None of our treatment groups showed a significant threshold difference between the two measurements (Mixed-model ANOVA, [$F_{(2,35)} = 0.222$, $p = 0.802$]; mean difference < 5 dB). Neural bandwidths of single AN fibers clearly varied with BF (Fig. 2), as expected, but did not differ significantly between untreated young adults and gerbils treated with 70 μM ouabain (Mixed-model ANOVA, [BF effect: $F_{(1,76)} = 71.67$, $p = 1.41 \times 10^{-12}$], [treatment effect: $F_{(1,76)} = 0.228$, $p = 0.634$]). Thus, the ouabain treatment did neither affect basic cochlear sensitivity nor frequency selectivity.

In our old gerbils, CAP thresholds showed the typical, variable sensitivity loss observed previously for individuals of comparable age (Heeringa 2024). The mean threshold elevation was 36 dB, compared to young adults (T-test, $p = 1.77 \times 10^{-5}$, n(young-adult cochleae) = 38, n(old cochleae) = 15). It was also shown previously that old gerbils still show normal, sharp frequency selectivity of their tuning curve tips, i.e., near threshold (Heeringa et al. 2020, Hellstrom and Schmiedt 1996). This was confirmed for our sample of old gerbils (Fig. 2; Mixed-model ANOVA, [BF effect: $F_{(1,90)} = 101.58$, $p = 1.96 \times 10^{-16}$], [age effect: $F_{(1,90)} = 2.28$, $p = 0.135$]) and is consistent with only mild OHC pathologies in aging gerbils (Tarnowski et al. 1991, Adams and Schulte 1997, Steenken et al. 2024).

Figure 2: *Bandwidths of neural response curves were not affected by ouabain treatment or age. Half-maximal bandwidth of response curves of single AN fibers, obtained with pure tones at a level <= 20 dB above the individual fibers' rate threshold, as a function of BF. Color code for the different treatment groups is the same as in Fig. 1.*

Average rate did not carry sufficient information about inharmonic frequency shift.

Spike responses to TFS1 stimuli of 94 AN fibers from 26 animals were recorded (for details see Table 1 in Materials and Methods). BFs ranged from 470 Hz to 11,528 Hz, with a median of 3,995 Hz. The SRs of these fibers ranged from 0 sp/s to 144 sp/s with a median of 60 sp/s. Average stimulated rates (4.8 sp/s - 223 sp/s, median 135 sp/s) and driven rates

(i.e., average rate minus SR, 4.8 sp/s - 215 sp/s, median 67.6 sp/s) were calculated from the

spike counts across the stimulus duration of 375 ms, excluding on- and offset ramps.

Most importantly, the inharmonic frequency shift Δf% - as a covariate in the ANOVA - had no

significant effect on mean driven rate $[F(1,943) = 0.018, p = 0.8933]$ (Fig. 3, overlapping

colored and black distribution curves in each panel). The overall distribution of driven rate

also did not change substantially with Δf%. Thus, the inharmonic frequency shift had no

effect on the mean driven rate, and AN spike rate was not a potential cue for the gerbil in the

behavioral task.

Beyond this robust, basic result, there were differences in discharge rates between the

animal groups and stimulus conditions that might reflect different sampling biases of single

units in the different experimental groups. These details are reported in the Supplemental

Material. We acknowledge that our single-unit sample was not sufficient, i.e., not all

subpopulations of AN fibers were sufficiently sampled for all animal treatment groups, to

conclusively resolve every aspect that may be interesting to explore. The power of our

approach lies in the direct linkage of several levels of investigation – cochlear synaptic

morphology, single-unit representation and behavioral performance – and, in the main

manuscript, we focus on the core question of synaptopathy and its relation to TFS

perception.

Figure 3: Mean driven spike rates did not differ between frequency shifts. *Histograms of*

average driven spike rates during presentation of the TFS1 stimuli (black and colored) and

average spontaneous rate, without acoustic stimulation (gray). Subject groups are arranged
in columns and stimulus conditions in rows. Colors code the frequency shift Δf in percent of
f_0, with the responses to harmonic stimuli shown in black and responses to the inharmonic
stimuli grading from yellow (least inharmonic) to red (most inharmonic). The legends apply to
panels above them. In each panel, the maximal number of fibers (with 40 stimulus repetitions
per fiber) that the histograms are based on is indicated; note that not all fibers were tested
with each frequency shift Δf. Clipped spontaneous rate bars are marked with an arrow and
the maximal relative proportion beyond the axis limit.

Neural representation of TFS was not degraded in AN fibers of old or synaptopathic gerbils.

The same set of AN fibers was probed for the neural responses to the fine structure and
envelope of TFS1 stimuli (for details see Table 1) using the TFS log-z-ratio (see section
"Neural recordings - Data Analysis"). A high TFS log-z-ratio indicates strong phase locking to
the stimulus TFS. The main findings are listed below; for more details see Supplemental
Material. Importantly, the inharmonic frequency shift, Δf%, as a covariate in the ANOVA,
affected the TFS log-z-ratio significantly and strongly [main effect: $F_{(1,943)} = 104.403$,
$p = 2.569×10^{-23}$], suggesting that AN fibers followed the fine structure of the stimulus and
thus represented the frequency shift in their spiking pattern (Fig. 4A-J, L). Δf% significantly
interacted with stimulus condition and BF-class [interaction: $F_{(2,943)} = 20.292$,
$p = 2.353×10^{-9}$; $F_{(1,943)} = 31.050$, $p = 3.282×10^{-8}$, respectively]. In response to
400/1600 Hz, the shifts affected the temporal representation most (i.e., higher Δf% resulted
in higher TFS log-z-ratios). Additionally, TFS log-z-ratios of low-BF fibers were influenced
more by the inharmonic shifts than those of high-BF fibers (again, higher Δf% result in higher
TFS log-z-ratios).
TFS representation by the TFS log-z-ratio metric was not significantly different between AN
fibers from different gerbil groups (p = 0.154). Also, no interaction between group and Δf%
occurred. Together, this indicates that aged fibers (Fig. 4D, H, L) and fibers that survived the
treatment with ouabain (Fig. 4C, G) had responses that did not differ from young-adult (Fig.
4A, E, I) and sham-treated fibers (Fig. 4B, F, J). As expected, TFS log-z-ratios of AN fibers
responding to the stimulus condition with higher f_c were lower, compared to those responding
to stimuli with lower f_c (200/1600 Hz: 0.987, 400/1600 Hz: 1.111, 400/3200 Hz: 0.394; [main
effect: $F_{(2,943)} = 7.495$, $p = 5.894×10^{-4}$]). This reflects the known decrease in phase-locking
with increasing stimulus frequency (Versteegh et al. 2011).

Representation of f0 was enhanced in AN fibers of old gerbils.

As a metric for the relative representation of TFS vs. stimulus envelope, the ENV/TFS log-z-ratio was used (see section "Neural recordings - Data Analysis"). This metric reflects the relative emphasis in the AN fibers' responses of either TFS (negative ENV/TFS log-z-ratio) or envelope locking (positive ENV/TFS log-z-ratio).

Here, over all animal groups, Δf% had a marginally significant effect on the ENV/TFS log-z-ratio [main effect: $F(1,943) = 4.278$, $p = 0.039$] and stimulus condition had a highly significant effect on the ENV/TFS log-z-ratio [main effect: $F(2,943) = 43.929$, $p = 5.744 \times 10^{-19}$]. Posthoc comparisons revealed that, for 400/3200 Hz, envelope representation (mean ENV/TFS log-z-ratio = 0.263) was stronger than fine structure representation, indicating stronger phase-locking of AN fibers to low-frequency stimulus components. Furthermore, stimulus condition showed a significant interaction with gerbil groups [interaction: $F(6,943) = 11.439$, $p = 2.242 \times 10^{-12}$]. In old gerbils, the strength of phase-locking was nearly equal to the envelope (Fig. 4D, left inset; Fig. 5D) and to the TFS (Fig. 4D, right inset; Fig. 5D'), in response to 200/1600 Hz.

Figure 4: Frequency shift representation in the neural responses differed between groups and stimulus conditions. *Vector strength frequency spectra of responses to harmonic and inharmonic stimuli, averaged across repetitions and fibers. Subject groups are arranged in columns and stimulus conditions in rows. Colors code the frequency shift Δf in percent of f0, with the same color code as in Fig. 3. Dashed lines indicate the limits of the stimulus bandpass filters. The inset panels show enlarged examples of the spectra ranging*

around the envelope frequency (f₀, 200 or 400 Hz) and the fine structure (center) frequency
(TFS peak frequency; 1000, 800, or 2000 Hz). The thin gray lines mark the y-axis scaling of
the inset panels. In each panel, the maximal number of fibers (with 40 stimulus repetitions
per fiber) that the spectra are based on is indicated; note that not all fibers were tested with
each frequency shift Δf.

In response to 400/3200 Hz, fibers from old gerbils also showed better envelope locking (Fig.
4L, Fig. 5L). Since TFS representation was unchanged in old animals (see above), this result
reflects enhanced envelope coding by AN fibers obtained in old gerbils, compared to all other
animal groups (Fig. 4A-D, I-L, left insets). In contrast, AN fibers of young-adult gerbils,
irrespective of treatment, tended to respond strongly to the fine structure of the stimulus,
especially for conditions with lower f_c of 1600 Hz (i.e., showed VS peaks at the respective,
shifted frequencies; Fig. 4A-H, right insets; Fig. 5A'-C', E'-G').

Over all animal groups, low-SR fibers had, on average, positive ENV/TFS log-z-ratios, that is,
phase-locked more strongly to the stimulus envelope, whereas high-SR fibers showed, on
average, negative ENV/TFS log-z-ratios, that is, phase-locked more strongly to the TFS
[main effect: $F(1,943) = 29.138$, $p = 8.533 \times 10^{-8}$].

Figure 5: Frequency shift representation differed between envelope and TFS
frequency ranges. *Frequency shift of the z-value peak versus stimulus frequency shift Δf.*

Subject groups are arranged in columns and stimulus conditions in double rows. Odd rows
show the data at f_0 and even rows at $f_{maxpeak}$ (the frequency of maximal average response, cf.
Fig. 11). Each circle represents four dimensions: the stimulus frequency shift Δf (position on
the x-axis), the frequency shift of the z-value peak (position on the y-axis), the average
vector strength (z-score) across all respective fibers (color saturation, see legend at the
bottom), and the percentage of fibers with a frequency shift of the z-value peak at the
corresponding stimulus frequency shift Δf (circle radius). The largest circles correspond to
100% of fibers (e.g., in panel C), medium sized circles to values around 50% (e.g., panel J).
Colors code the frequency shift Δf in percent of f_0, with the same color code as in Fig. 3. In
each panel, the maximal number of fibers (with 40 stimulus repetitions per fiber) that the z-
value peaks are based on is indicated; note that not all fibers were tested with each
frequency shift Δf.

Age, but not synapse loss, affected behavioral discrimination of TFS1 stimuli.

To investigate the effect of synapse loss on performance in the TFS1 task, the behavior of
young-adult gerbils, without and with cochlear synaptopathy, and old gerbils was compared
for the different stimulus conditions defined by the combinations of f_0 and f_c. The sensitivity of
the four groups of gerbils in the different stimulus conditions is illustrated in Fig. 6, in relation
to the frequency shift Δf%. The sensitivity of young-adult gerbils of the different treatment
groups was quite similar, whereas the old gerbils had considerably lower sensitivity. A GLMM
ANOVA with the sensitivity *d'* as dependent variable, condition and treatment group as
factors, and frequency shift Δf% as a covariate (only trials with Δf% > 5 included, for which
data from all treatment groups were obtained), revealed significant main effects of treatment
group   [$F(3,387) = 8.18$, $p = 2.725 \times 10^{-5}$],   stimulus   condition   [$F(2,387) = 22.71$,
$p = 4.736 \times 10^{-10}$, and Δf% [$F(1,387) = 20.457$, $p = 8.121 \times 10^{-6}$]. No significant interactions
were observed. There was a general increase in sensitivity with increasing Δf%. Old gerbils
showed lower sensitivities than most young-adult gerbils, irrespective of their treatment (all
$p < 0.002$, Bonferroni corrected). There was no significant difference in sensitivity between
the treatment groups of young gerbils. To separately investigate the effects of f_0 and
harmonic number, *N*, on sensitivity, we conducted planned comparisons in two additional
GLMM ANOVAs, each containing only the data of two conditions, with the same dependent
variable and factors as the initial ANOVA. The planned comparison with the data of stimulus
conditions 200/1600 Hz and 400/1600 Hz was used to investigate the effect of f_0 (which
inherently includes a change in harmonic number, N) on sensitivity. This ANOVA revealed an
effect of f_0, with a higher sensitivity for increasing f_0 [$F(1,257) = 45.001$, $p = 1.248 \times 10^{-10}$].
Furthermore, an effect of Δf% [$F(1,257) = 13.539$, $p = 2.847 \times 10^{-4}$] was observed. Similar to

the previous analysis, the main effect of gerbil group [$F_{(3,257)}$ = 4.878, p = 2.570×10⁻³] reflected the low sensitivity of the old gerbils compared to all other groups. The second planned comparison with the data of stimulus conditions 400/3200 Hz and 400/1600 Hz, with equal f_0, investigated the dependence of sensitivity on f_c. An effect of f_c [$F_{(1,256)}$ = 12.656, p = 4.459×10⁻⁴] was shown, with decreasing sensitivity for increasing f_c. As in the previous comparison, this ANOVA also revealed increased sensitivity with increasing Δf% [$F_{(1,256)}$ = 13.484, p = 2.929×10⁻⁴], whereas the main effect of gerbil group [$F_{(3,256)}$ = 4.692, p = 3.299×10⁻³] again reflected the lower sensitivity of the old group compared to all other groups. No interactions were observed in either planned comparison.

The frequency-shift threshold reflects Δf% for a criterion of *d'* = 1 and was calculated for each gerbil and condition. If no threshold could be derived because all *d'* values were below 1, a threshold of 100% f_0 was assumed for further statistical analysis. On average, old gerbils were only able to discriminate inharmonically shifted TFS1 stimuli for the condition 400/1600 Hz (see *d'* values > 1 in Fig. 6D, H, L). All young-adult gerbil treatment groups achieved average sensitivities above threshold for all stimulus conditions.

Figure 6: The gerbils' discrimination performance improved with larger frequency shifts. *Sensitivity index d' for behavioral discrimination, as a function of stimulus frequency shift Δf in percent of f₀. Reference lines at d' = 1 indicate the assumed threshold value for meaningful discrimination performance. Subject groups are arranged in columns and stimulus conditions in rows. Colors code the subject group. Circled black dots show the mean across subjects, boxes show the median and interquartile ranges, whiskers show the*

extrema, except for outliers, which are displayed as colored dots. Data points are considered

outliers, if they lie beyond 1.5 times the distance between median and upper or lower

quartile, respectively. For each panel, the maximal number of subjects in the respective

group is indicated. Unfilled boxes mark conditions completed by less than half of those

subjects in the respective group.

Frequency-shift thresholds for the three different stimulus conditions are shown in Fig. 7. A

GLMM ANOVA was conducted with Δf% thresholds as the dependent variable and condition

and treatment group as factors. There were main effects for stimulus condition

$[F_{(2,60)} = 8.99$, $p < 0.0005]$ and treatment group $[F_{(3,60)} = 6.84$, $p < 0.0005]$, with no

interaction between both factors. Average thresholds were not significantly different between

the 200/1600 Hz and 400/3200 Hz conditions, while thresholds were lower in the

400/1600 Hz than in the 200/1600 Hz condition ($p < 0.0005$). The pairwise comparison

revealed that old gerbils had significantly higher thresholds than gerbils of all other animal

groups (all $p < 0.005$).

 Figure 7: Old gerbils were typically unable to perceive frequency shifts whereas

 ouabain treatment did not impair discrimination. Behavioral discrimination thresholds

 based on the data shown in Fig. 6. Thresholds were defined as the lowest (linearly

 interpolated) stimulus frequency shift Δf in percent of f_0 at which the sensitivity index d'

 crossed d' = 1. The threshold was set to 100% if this criterion was never met. Colors code

 the subject group. Circled black dots show the mean across subjects, boxes show the

 median and interquartile ranges, whiskers show the extrema, except for outliers, which are

 displayed as colored dots. Data points are considered outliers, if they lie beyond 1.5 times

 the distance between median and upper or lower quartile, respectively. The y-axes at the

 *right show the same frequency shifts in Hz. Significance levels are ** p < 0.002 for all three*

comparisons, young-adult vs. old p = 1.53×10^{-3}, sham vs. old p = 1.11×10^{-3}, ouabain high vs.
old p = 0.382×10^{-3}.

No differences were observed between the young-adult gerbil groups with different
treatments. To evaluate the effect of f0 on the threshold, a planned comparison with the data
of stimulus conditions 200/1600 Hz and 400/1600 Hz (i.e., harmonic numbers 8 and 4,
respectively) was conducted in an additional ANOVA, with frequency-shift thresholds as
dependent variable, and condition and gerbil group as factors. A main effect of condition was
revealed, indicating that f0 (respectively harmonic number N at 1600 Hz fc) did affect the
threshold [$F(1,40) = 15.219$, $p = 3.577×10-4$]. There was no significant effect of gerbil group;
all groups showed similarly low thresholds for the 400/1600 Hz condition. A second
additional ANOVA with the data of stimulus conditions 400/3200 Hz and 400/1600 Hz (i.e.,
harmonic numbers 8 and 4, respectively) investigated the dependency of threshold on fc
(respectively harmonic number N). An effect of gerbil group [$F(3,40) = 10.460$, $p = 3.272×10-$
5] and condition [$F(1,40) = 16.213$, $p = 2.455×10-4$] was observed. The pairwise comparison
revealed lower thresholds with decreasing fc (respectively harmonic number N) and a
significant difference between untreated and old gerbils (all $p ≤ 0.001$), whereas the young-
adult gerbil treatment groups did not differ.

It has been suggested that human subjects compensate difficulties in speech perception in a
noisy background due to deficits in TFS processing by increasing their listening effort
(Parthasarathy et al. 2020). We investigated the possibility that the ouabain-treated young
animals may have compensated for their loss of synapses by increasing their listening effort,
predicted to result in longer response latencies. To this end, we analyzed the distributions of
response latencies for a range of frequency shifts between 2.65 % and 31.88 % applying a
mixed model ANOVA with response latency as the dependent variable and group (ouabain-
treated young animals vs. untreated young-adult animals), stimulus conditions (200/1600,
400/1600, and 400/3200) and frequency shift as factors. Response latency was significantly
affected by the extent of the frequency shift [$F(4,3451) = 7.516$, $p = 5.027×10^{-6}$] and by the
stimulus condition [$F(2,3451) = 3.621$, $p = 2.687×10^{-2}$], but not by the group
[$F(1,3451) = 15.219$, $p = 3.719×10^{-1}$]. There were no significant interactions. Thus, we
conclude that there is no evidence that ouabain-treated young animals compensated for a
putative perceptual deficit by increasing the listening effort.

In contrast, a similar analysis comparing untreated young-adult and old animals revealed
significantly longer response times in old animals compared to young-adult animals
(difference 66 ms, [$F(1,2251) = 13.156$, $p = 2.930×10-4$]). Despite the longer response
latencies in old animals, suggesting increased listening effort, the performance of the old
animals did not reach that of young-adult animals.

In summary, the behavioral data showed that the old gerbils differed from all other treatment
groups of young-adult gerbils. This result suggests that age considerably affected both
perceptual sensitivity and thresholds while synapse numbers had little effect.

Discussion

The ability to process temporal fine structure (TFS) of sounds is a key factor for speech
perception. Its age-related decline has been held responsible for compromised speech
comprehension in the elderly, especially in challenging noisy acoustic backgrounds
(Füllgrabe et al. 2015, Moore 2014, 2019). It has been suggested that age-related or noise-
induced synaptopathy results in the observed perceptual deficits regarding the TFS of
sounds (e.g., Moore 2014, 2019). However, so far a direct test of this hypothesis is lacking.
Here, we present data that investigate the performance of the Mongolian gerbil in the TFS1
test (Moore and Sek 2009), a psychoacoustic paradigm that has been applied in human
subjects to evaluate the ability to discriminate between stimuli based on the TFS. We
determined the TFS1 test performance in young-adult gerbils and compared this to the
performance of old gerbils (≥ 36 months of age, corresponding to humans ≥ 60 years of age,
Castaño-González et al. 2024), demonstrating an age-related decline in TFS sensitivity
similar to that found in human subjects (Füllgrabe et al. 2015, Moore et al. 2012). However,
young-adult gerbils with experimentally induced synaptopathy (by applying ouabain), to a
degree corresponding to the synaptopathy in old gerbils (Steenken et al. 2021, this study),
showed no decline in behavioral TFS sensitivity. This result suggests that reduced synapse
numbers alone cannot explain the perceptual deficits for TFS in old subjects. Auditory-nerve
(AN) fiber responses of young-adult, young-adult ouabain-treated, and old gerbils, elicited by
the stimuli presented in the TFS1 test, indicate that the representation of TFS by AN fibers is
not affected by ouabain treatment or age. In old gerbils that show compromised TFS1 test
performance behaviorally, however, the representation of the signal envelope by AN fibers
was enhanced compared to the other groups. We propose that this enhancement of the
signal envelope cue, which does not differentiate the stimuli in the TFS1 test, may cause the
perceptual deficits in old gerbils by overriding the usability of the TFS cues. Thus, cue
representation may be associated with the perceptual deficits, but not reduced synapse
numbers, as originally proposed.

Gerbil as a model for human temporal-fine-structure perception.

428 In the TFS1 test, subjects must discriminate an inharmonic (I) complex from a harmonic (H)
429 complex, in which the frequency difference between the components determines the period
430 of the envelope. The I complex is created by shifting all components of the H complex by a
431 fixed frequency. This shift results in pairs of I and H complexes with similar envelopes but

different TFS. Thus, TFS is the only acoustic cue which allows to discriminate the stimuli. Excitation-pattern differences due to the shift of the frequency components in the I complex compared to the H complex can be eliminated in the TFS1 test by appropriate spectral filtering (e.g., Marmel et al. 2015). In addition, the gerbil has larger auditory filter bandwidths than human subjects (Kittel et al. 2002). Since we used a similar filter bandwidth for the gerbil as that applied in studies involving humans (e.g., Eipert et al. 2019, Marmel et al. 2015, Moore and Sek 2009), cochlear excitation pattern differences are unlikely to have provided usable cues for discrimination by the gerbils in the present study, at least for conditions with a harmonic number N = 8 (200/1600 Hz, 400/3200 Hz). However, it appears possible that for the stimulus condition 400/1600 Hz (harmonic number N = 4), the gerbils' discrimination may have relied on excitation pattern differences (see model results in Fig. 8). The modeled cochlear excitation patterns also suggest that the noise masker in which the I and H complexes were presented ensured that the stimuli did not produce usable distortion products for the gerbils. Thus, as in the study with human subjects, the only usable cue for the discrimination for conditions with a harmonic number N = 8 was the TFS.

Figure 8: Gerbil excitation patterns for harmonic (thick black lines in panels A, C, E) and inharmonic stimuli close to the frequency shift at the TFS1 threshold (thick dark green lines in panels A, C, E) and their differences (panels B, D, F) for the three stimulus conditions. Thin black and green lines in panels A, C, E show the physical sound spectra for the corresponding harmonic and inharmonic stimuli, respectively. Red lines in panels A, C, E show the excitation pattern elicited by the pink noise masker. Since the level of excitation produced by the pink noise is less than 30 dB below that produced by the complex tones, distortion products will be masked. Excitation patterns were calculated with the assumption that the gerbil auditory filter bandwidth is 1.8 times the human auditory filter bandwidth (see

Kittel et al. 2002). The thick green lines in panels B, D, F show the differences between the excitation pattern elicited by the harmonic reference and the inharmonic stimuli close to the frequency shift at the TFS1 threshold. The thin yellow and orange lines show these differences for the different frequency shifts of the inharmonic stimuli (color codes as in Fig. 5). The grey-shaded areas in panels B, D, F indicate the 5%/95% percentiles of the amplitude statistics in the harmonic reference signals. The blue dashed lines in panels B, D, F show the gerbil threshold for detecting a change in the intensity of a 70 dB 1-kHz tone (Sinnott et al. 1992). Only for the condition 400/1600 Hz does the change in the level of the excitation pattern due to the frequency shift exceed the threshold for detecting the intensity difference. For this condition, the gerbils could solve the task, at their TFS1 threshold, by comparing the difference in excitation elicited by the harmonic and frequency-shifted inharmonic stimulus. For the other conditions, this change is less than the intensity difference limen. Thus, for these two conditions with harmonic number N of 8 the gerbils cannot rely on differences in the excitation patterns but must solve the task by comparing the temporal fine structure.

Figure 9 provides an overview of gerbil TFS1 thresholds in comparison to human thresholds. Human thresholds were selected from studies involving stimuli with spectral components and fundamental frequencies (f_0) similar to those presented to the gerbils in the present study, allowing a comparison of gerbil and human performance. There is a broad overlap between gerbil and human TFS1 thresholds. Both, in humans and gerbils, old subjects showed higher (i.e., worse) TFS1 thresholds than young subjects (for human studies see Eipert et al. 2019, Füllgrabe et al. 2015, Moore et al. 2012). This similarity indicates that the gerbil is a good model for investigating the mechanisms underlying TFS perception that may also apply to humans. Behavioral TFS1 thresholds in young-adult gerbils with ouabain-induced synaptopathy, similar to that typical for old gerbils, were similar to TFS1 thresholds in untreated young-adult gerbils. This result suggests that increased behavioral thresholds in old subjects were not due to the reduced number of synapses. Thus, in the gerbil, the hypothesis that age-related deterioration of TFS perception is a direct consequence of synaptopathy can be rejected.

The observed extent of age-related or noise-induced loss of type-I afferent synapses on IHC varies widely between species and studies. For example, in ageing CBA/CaJ mice, mean losses of between 20 and 50% of afferent synapses (depending on cochlear location and precise age) were reported (Sergeyenko et al. 2013, Kobrina et al. 2020). Humans showed more pronounced losses of peripheral axons, of 40–100%, again depending on cochlear location, precise age, and noise history (Wu et al. 2019, 2021). The age-related and induced synapse losses in our gerbils were in a more moderate range, around 20% (Steenken et al. 2021, this study). Thus, it is possible that a more severe, induced synaptopathy would have

resulted in behavioral deficits in young-adult gerbils. However, in the absence of additional noise or pharmacologically induced damage, our study provides strong evidence for other factors causing temporal processing problems with advancing age. Our 3-year-old gerbils are approximately comparable to a 60-year-old human (Castano-Gonzalez et al. 2024) with beginning but not yet clinically relevant hearing loss (Hamann et al. 2002).

Figure 9: Frequency shift detection thresholds of gerbils and humans for TFS1 test stimuli in relation to the center frequency f_c of the harmonic complex. Filled symbols represent gerbil data from the present study. Human data are from published material with the limitation that the f_c was in the range from 500 Hz to 4500 Hz and the fundamental f_0 of the harmonic complex was in the range of 100 Hz to 400 Hz to make the range of parameter values similar to those of the present study (for references see legend, normal hearing NH, hearing loss HL). The higher and lower thresholds shown for the gerbil data reflect thresholds at f_c of 1600 Hz for fundamentals f_0 of 200 Hz and 400 Hz, respectively.

Auditory-nerve fiber responses do not explain differences in behavioral sensitivity.

Since the use of excitation patterns by the gerbil is unlikely due to its larger auditory-filter bandwidths for stimulus conditions of 200/1600 Hz and 400/3200 Hz (Fig. 8), we explored the temporal responses of AN fibers for alternative explanations of compromised TFS perception. TFS representation requires the ability of AN fibers to phase-lock (reviewed in Joris et al. 2004, Rose et al. 1967). Here we showed that, throughout all tested gerbil groups, the H and I complexes were neurally distinguishable based on phase-locking to the stimulus fine structure (Fig. 4, 5). In contrast, no distinction was possible based on the average firing rate of AN fibers (Fig. 3). Neural phase-locking reached highest z-values for stimuli with f_c of

1600 Hz and were lower for 3200 Hz (Fig. 5), which reflects the known decrease in phase-
locking of AN fibers for stimuli above 1.5 kHz (Versteegh et al. 2011). Several studies in
animal models of cochlear dysfunction showed that phase locking is not compromised in
single AN fibers, including quiet-aged gerbils (Heeringa et al. 2020), noise-exposed cats
(Miller et al. 1997), chinchillas (Henry and Heinz 2012, Kale et al. 2013), and kanamycin-
treated guinea pigs (Harrison and Evans 1979). Consistent with these findings, the surviving
AN fibers in old gerbils did not show compromised TFS representation of I and H complexes
compared to young-adult gerbils (Fig. 4, 5).

Old gerbils showed significant synapse loss compared to untreated young-adult gerbils at the
cochlear location corresponding to 2 kHz (Fig. 1, Fig. S1 for all frequencies), consistent with
what was shown previously (Steenken et al. 2021). Ouabain, a cardiac glycoside that inhibits
Na^+-K^+-ATPase pumps of AN fibers (O'Brien et al. 1994), applied to the round window,
results in a dose-dependent loss of AN fibers in gerbils (Bourien et al. 2014, Lang et al. 2005,
Schmiedt et al. 2002). In our hands, a low dose (40 µM) of ouabain had no effect at the
cochlear location equivalent to 2 kHz. In contrast, a higher dose of 70 µM resulted in a loss
that was similar to that observed in old gerbils (Fig. 1). The surviving fibers of ouabain-
treated gerbils showed no deterioration of frequency tuning and TFS representation
compared to untreated or sham-treated young-adult gerbils (Fig. 2, Fig. 4A,C,E,G); however,
the behavioral outcomes of this study clearly showed that only old gerbils had deteriorated
discrimination abilities between H and I complexes (Fig. 7). We thus showed conclusively
that neither synapse loss, to an extent typical for old gerbils, nor impaired TFS
representation, at the level of single AN fibers, could explain the deteriorated perception in
old gerbils.

Enhancement of confounding/distracting AN cues and changed
central processing may explain the behavioral deficit in old
subjects.

Surviving AN fibers in old gerbils did not show compromised fine-structure representation
(Fig. 4, 5). However, old gerbils exhibited deteriorated performance in the behavioral test in
response to TFS1 test stimuli (Fig. 6, 7). Despite the overall robust phase-locking ability,
other pathologies are known to develop with advancing age.

First, the age-related atrophy of the stria vascularis within the lateral wall of cochleae
(Gratton and Schulte 1995) results in a decreased endocochlear potential (Schulte and
Schmiedt 1992), which in turn increases the threshold (i.e., decreases the sensitivity) of AN
fibers (Schmiedt et al. 1996). Envelope representation, measured as synchronization to
amplitude-modulated stimuli, is non-monotonic with sound level, with a pronounced peak
near the threshold of the fiber (Heeringa et al. 2023, Joris and Yin 1992, Kale and Heinz

2010). Thus, the typical sensitivity loss in old subjects will cause a difference in neural
representations between young-adult and old subjects, even if the test stimulus levels are
kept constant. Alternatively, stimulus levels can be raised for old subjects, to an equivalent
sensation level that compensates for their sensitivity loss. Consistent with that prediction, AN
fibers of old gerbils show enhanced representation of the fundamental frequency of speech
vowels, compared to fibers from young adults, when responding to fixed-level stimuli
(Heeringa et al. 2023), but do not show enhanced envelope representation in response to
frozen noise presented at similar sensation level (Heeringa et al. 2020). Old gerbils of the
same age as in the present study invariably show hearing loss, although to variable extents
(Hellstrom and Schmiedt 1990, Steenken et al. 2021, this study). Thus, since TFS1 stimuli in
the current study were presented at a fixed absolute level of 68 dB SPL, fibers of old animals
were stimulated closer to their individual thresholds than fibers of young adults, which is
expected to increase their phase locking to f_0, as observed. Our findings match those of Kale
and Heinz (2010) for AN fibers of chinchillas with noise-induced hearing loss. After noise
exposure, envelope coding was enhanced, in approximate proportion to the fiber's sensitivity
loss, while the representation of TFS remained unchanged (Kale and Heinz 2010). Together,
these AN data suggest that both age-related and noise-induced hearing loss result in a
relative deficit of TFS representation that may ultimately contribute to perceptual problems.

A decreased endocochlear potential may also affect the AN fibers' frequency selectivity,
indirectly via compromised OHC function (Ruggero and Rich 1991). In chinchillas with noise-
induced hearing loss, deteriorated frequency tuning of AN fibers and tail hypersensitivity led
to abnormal temporal coding, such that a mismatch emerged between the fibers'
characteristic frequency and the dominant frequency represented in their TFS responses
(Henry et al. 2016), including responses to TFS1 stimuli (Kale et al. 2013). However, this is
an additional effect, unique to noise damage, and not likely to apply to our aged gerbils. In
old gerbils, the frequency selectivity of the AN fibers' sensitive tuning-curve tips is not
significantly affected (Heeringa et al. 2020, Hellstrom and Schmied 1996, this study) and
there is no evidence for tail hypersensitivity (Hellstrom and Schmied 1996), consistent with
only mild OHC pathologies in aging gerbils (Tarnowski et al. 1991, Adams and Schulte 1997,
Steenken et al. 2024). Furthermore, in chinchillas treated with furosemide, a drug that
reversibly reduces the endocochlear potential, the frequency tuning curves of AN fibers were
affected in their tip-to-tail ratios, however, much less drastically than after noise damage
(Henry et al. 2019). Together this suggests that AN fibers in aged animals, without
confounding noise-induced hearing loss, experience only mild impairments of frequency
tuning that did not cause the changes to temporal coding observed in our old gerbils.

Second, and moving beyond cochlear pathologies, age-related changes in GABAergic
processing in the central auditory system do occur (Caspary et al. 1995). Such changes were

also specifically shown for the gerbil (Gleich et al. 2014, Kessler et al. 2020) and were shown to affect temporal processing (Gleich et al. 2003). The gerbils used by Kessler and colleagues (2020) were derived from the same population as the gerbils used for the present study and had the same age range for old animals. Therefore, it can be assumed that the gerbils in the current study showed a change in GABAergic processing in the brain that could affect processing of such supra-threshold stimuli as being used in the TFS1 test.

Third, olivocochlear efferent activity is a potential modulator of OHC gain (by medial olivocochlear neurons, MOC) and afferent activity (by lateral olivocochlear neurons, LOC). Beyond this general observation it is, however, difficult to speculate about its specific role in the TFS1 test, as almost nothing is known about efferent activity under naturalistic conditions in a behaving animal (reviewed by Lauer et al. 2022). We note that efferent activity is believed to be reduced under general anesthesia (reviewed by Guinan 2011) and possibly abnormal in other ways, considering the potential top-down inputs to the efferent neurons from extensive brain networks (reviewed by Schofield 2011, Romero and Trussell 2022). Thus, it is reasonable to assume a reduced efferent influence in our AN data, compared to the behavioral test situation. As for changes with aging, we tentatively assume more comparable efferent influences in young-adult and old gerbils. It was recently shown that, despite age-related losses in both MOC and LOC cochlear innervation, this basically reflected the loss of efferent target structures (OHC and type-I afferents), with the surviving cochlear circuitry remaining largely normal (Steenken et al. 2024). The main difference was an increased proportion of OHCs without any efferent innervation, predominantly in low-frequency cochlear regions (Steenken et al. 2024). Such OHCs are thus not under efferent control, and they are more numerous (about 10 – 30%) in old gerbils.

Conclusion

We demonstrated that the gerbils' ability to behaviorally discriminate TFS1 stimuli is neither affected by synapse loss nor by alterations to TFS representation by their AN fibers. However, the elevated neural response thresholds in old gerbils enhanced the neural envelope representation. This enhancement may have overridden the still intact TFS representation.

Acknowledgments

This work was supported by the DFG priority program "PP 1608", the DFG Cluster of Excellence EXC 1077/1 "Hearing4all" (Project ID 390895286). We thank Laurel Carney for helpful comments on a previous version of this manuscript. Susanne Groß for assistance with the behavioral experiments. Amarins N. Heeringa for providing analysis scripts for AN

data and analyzing electrophysiological data. Nadine Thiele for generous assistance with the
gerbil care and for implementing the blinding process. Furthermore, many thanks go to Sonja
Standfest for the great assistance with gerbil care. Sonny Bovee is acknowledged for
processing cochleae, Julia Forst and Carina Lützow for assistance with the spike detection
and immunostainings, and Julia Winter for assistance with the immunostainings. We
acknowledge the Fluorescence Microscopy Service Unit, Carl von Ossietzky University of
Oldenburg, for the use of the imaging facilities.

Materials and Methods

Experimental model and subject details

Sixty adult, agouti-colored Mongolian gerbils (*Meriones unguiculatus*) of both sexes (32 M,
28 F) were studied (Table 1). The gerbils originated from Charles River Laboratories and
were bred in an in-house facility of the University of Oldenburg. Groups of gerbils were
tested, differing in age and ouabain treatment: untreated young-adult, surgery-only young-
adult, ouabain-treated young-adult, and old gerbils. Young-adult gerbils were between 3 and
19 months of age and old gerbils were 36 months or older. All animals were housed
individually or in pairs in Type IV cages, which were provided with nesting material. During
behavioral testing, access to water was not limited, but animals were food-deprived, such
that the weight of young-adult and old gerbils was 65 - 75 g and 70 - 85 g, respectively, to
motivate the animals to perform for a food reward. All protocols and procedures were
approved by the Niedersächsisches Landesamt für Verbraucherschutz und
Lebensmittelsicherheit (LAVES), Germany, permit AZ 33.19-42502-04-15/1990. All
procedures were performed in compliance with the NIH Guide on Methods and Welfare
Consideration in Behavioral Research with Animals (National Institute of Mental Health,
2002).

Table 1: Numbers of gerbils of different treatment and age groups within the behavioral,
electrophysiological, and histological part of this study. Note that numbers in category 2-
4 do not sum up to total numbers, since many gerbils were used in all three study parts.
For the neural data, the core data for this study are the AN fibers which were tested with
TFS1 stimuli. However, some additional fibers, in some animal groups also additional
animals, were included for the analysis of neural tuning; these numbers are provided
separately. Furthermore, both untreated gerbil groups were supplemented with data
from eight young-adult gerbils and twelve old gerbils for which synapse counts were
available, from Steenken et al. (2021).

		Young	Surgery	Ouabain	Ouabain	Old

		adult	only	40 µM	70 µM	
1	**Total** numbers of animals in this study	21	4	8	10	17
2	Animals in **behavioral** study	4	4	5	8	3
3	Animals with **neural tuning** data (# AN fibers)	5 (27)	3 (17)	6 (38)	7 (21)	6 (13)
	neural TFS1 resp. (# AN fibers)	5 (37)	2 (10)	3 (8)	6 (14)	6 (23)
4	Animals with **synapse counts**	19	3	8	10	14

Young-adult gerbils were on average 9.9 ± 5.4 months of age (N = 21) and old gerbils were

38.8 ± 2.5 months old (N = 17) when electrophysiological recordings were conducted and

cochleae were harvested. Treated young-adult gerbils were either 15.1 ± 3.6 (treated with a

low concentration of ouabain, N = 8) or 12.7 ± 4.2 (high concentration of ouabain, N = 10). All

surgery-only gerbils were 17.5 ± 1 months old (N = 4). Gerbil groups had significantly

different ages (univariate ANOVA [$F(4,55) = 129.449$; $p = 2.677 \times 10^{-26}$]). Bonferroni corrected

post-hoc tests revealed that old gerbils were significantly different from all other young-adult

gerbil groups with respect to age, while all young-adult gerbil groups, irrespective of

treatment, did not differ.

Methods details

Stimulus paradigm

The stimuli were similar to those used for the TFS1 test in humans (for a detailed description

see Moore and Sek 2009). In short, the reference stimulus was a harmonic tone (H)

complex. The test stimulus that was discriminated from the reference by the gerbils in the

behavioral tests or by analyses of AN-fiber responses was an inharmonic tone (I) complex

created by shifting all frequency components upwards. This procedure creates H and I

complexes with the same envelope periodicity. In the following, the frequency shift, Δf, will be

expressed by the percentage of the fundamental frequency, f_0 (Weber fraction).

The f_0 was either 200 or 400 Hz. H and I complexes were generated with components from f_0

up to 9 kHz. H and I complexes were passed through the same 3^{rd}- or 5^{th}-order Butterworth

infinite-impulse-response band-pass filter (for AN recording and behavioral experiments,

respectively), with a center frequency, f_c, of 1600 or 3200 Hz, to reduce possible cues related

to differences in the excitation pattern on the basilar membrane (Moore and Moore 2003).

The 3-dB down points of the filter were separated by 7 f_0, which resulted in a flat spectral

region of 5 f_0. The amplitude at the filter skirts decreased by at least 30 dB per octave

(except for the 3rd-order filter with f_c of 1600 Hz and f_0 of 400 Hz, for which the slope was

18 dB/oct).

The phase of each harmonic was randomly chosen for each test trial, but was matched for

the H and I complexes within a trial to ensure similar envelopes for both stimuli (see Fig. 10

A-D). A pink noise with frequencies between 100 and 11,050 Hz was added to the signal to

mask distortion products possibly emerging in the inner ear, and signal components outside

the filter passband. The masker power spectral density was 13 dB SPL at 1 kHz, equivalent

to 37 dB SPL in within the gerbil's auditory equivalent rectangular bandwidth centered on

1 kHz, to mask distortion products of H and I stimuli in the gerbil (Faulstich and Kössl 1999,

Bourien et al. 2014). The level of each component in the tone complex was 60 dB SPL

resulting in an overall level of 68 dB SPL. The duration of each H or I complex stimulus was

0.4 s. In the behavioral experiment, stimulus onset and offset were ramped with a 25-ms

Hann window. Stimuli presented to AN fibers were not ramped but still filtered. By analyzing

the steady-state response only, any on- and offset effects were excluded.

Figure 10: Examples illustrating stimulus waveforms and typical auditory-nerve

responses. Stimulus waveforms (A-D), peri-stimulus-time histograms (PSTHs, E-H), and

average vector strength as a function of frequency (I-L). Columns 1 (A,E,I) and 2 (B,F,J)

show data from a high (H)SR fiber, columns 3 (C,G,K) and 4 (D,H,L) show data from a low

(L)SR fiber. Both fibers are from a young-adult, untreated animal and are matched in best

frequency. The stimulus condition was $f_0 = 400$ Hz and $f_c = 1600$ Hz. A and C show examples

of the harmonic stimulus (Δf = 0%, black) and its envelope (gray), B and D show examples of

a strongly inharmonic stimulus (Δf = 8%, red), its envelope (gray), and the harmonic stimulus

(black) as a reference.The corresponding neural responses are displayed in the same colors

in the second and third row, respectively. Arrow markers are added (A-D) to highlight the

differences in fine structure between harmonic and inharmonic stimuli. The PSTHs (E-H)

show the instantaneous spike rate, averaged across 40 repetitions of identical stimuli. Vector

strength (I to L) is shown as a function of frequency, averaged across 40 stimulus

presentations. The inset panels show enlarged examples of the spectra in the envelope

frequency range (400 Hz) and the fine structure frequency range (800 Hz).

In both the behavioral experiments and in AN-fiber recordings, three combinations of f_0 and f_c

were used: $f_c = 1600$ Hz, and $f_0 = 200$ Hz; $f_c = 1600$ Hz, and $f_0 = 400$ Hz, and $f_c = 3200$ Hz

and $f_0 = 400$ Hz. The combination of $f_0 = 200$ Hz and $f_c = 3200$ Hz was not used because a

pilot behavioral test showed that two young untreated gerbils were not able to discriminate I

and H even for the largest value of Δf.

The harmonic rank, *N*, of a condition describes the rank of the center frequency in the

harmonic tone complex. The highest value for Δf was 42.5% of the fundamental f_0. Lower

values for Δf were computed by dividing 42.5% by 2n, with n being 1 or increasing integers.

The smallest Δf presented in the behavioral tests was 0.33% f_0. In old gerbils tested

behaviorally, additional test stimuli with Δf = 31.87% f_0 and 15.94% f_0 were presented to

ensure that old gerbils were presented with a sufficient number of trials allowing a salient

percept for discrimination.

In recordings from gerbil AN fibers, only subsets of the stimuli used in behavior were

presented to save time and minimize the number of incomplete datasets due to deterioration

or loss of single-unit isolation. The frequency shifts were further grouped into either "high"

(10.6, 15.9, 21.25, 31.8, and 42.5 Hz) or "low" (0.66, 1.3, and 5.31 Hz) and the stimuli were

presented in three different blocks per stimulus condition. The first block comprised the H

and two I complex stimuli, one with a high and one with a low shift. The second block

comprised I complex stimuli, two high shifts and another low shift. The third block always

comprised the remaining shifts (1.3, 10.6, and 42.5 Hz). The order of stimuli within the blocks

was random between fibers. A silent period of 0.6 s occurred between stimuli. This

organization of stimuli ensured that the H complex was always presented and could be

compared to one low and one high shift. The second and third block of stimuli were recorded

with declining success.

The periodic envelopes of H and I complexes created in the way described above are similar

and offer no cues for discrimination, whereas the TFS of H and I complexes is not identical

(Fig. 10A-D). The TFS of I complexes changes across sequential periods while the TFS of

the H complex does not; only this difference can be used for discrimination. The small

random change in the TFS and envelope of both H and I due to the addition of pink noise

offers no discrimination cues. The bandpass filtering of the tone complexes served to
minimize cochlear excitation pattern cues.

Experimentally induced synaptopathy using ouabain

For experiments including ouabain treatment, gerbils were anesthetized with ketamine
(Ketamin 10%, bela-pharm GmbH, Vechta, Germany; 135 mg/kg body weight) and xylazine
(Xylazin 2%, Ceva Tiergesundheit GmbH, Düsseldorf, Germany; 6 mg/kg body weight)
mixed with 0.9% saline injected intra-peritoneal. Maintenance doses of one third of the initial
dose were given hourly or as needed. Depth of anesthesia was monitored by withdrawal
reflex to toe pinches and via an ECG with needle electrodes in the front- and contralateral
hind leg, continuously displayed on an oscilloscope (SDS 1102CNL, SIGLENT Technologies,
Hamburg, Germany). Body temperature was continuously controlled with a homeothermic
blanket (Harvard Apparatus, Holliston, MA, USA). Experiments were carried out in a sound-
attenuating booth. The gerbil's head was fixed within a bite-bar (David Kopf Instruments,
Tujunga, CA, USA). Pure, moisturized oxygen (1.5 l/min) was delivered from a tube pointing
towards the nose.

Twenty-two gerbils (11M, 11F, aged 3 – 8 months at the beginning of treatment) received
either ouabain or artificial perilymph solution. Twenty-four hours before the oubain or artificial
perilymph injection, gerbils were treated with antibiotics (oral; Baytril (0.5%, Bayer Animal
Health GmbH, Monheim am Rhein, Germany; 0.3 ml per gerbil/day) and a non-steroidal
antiphlogistic/ analgetic agent (oral; Meloxidyl (0.5 mg/ml, CevaTiergesundheit GmbH; 0.3 ml
per gerbil/day; or Novalgin; oral; 500 mg/ml, Sanofi-Aventis Deutschland GmbH, Frankfurt,
Germany, 0.21 ml per gerbil/day). The antibiotic treatment was continued for nine more days,
the antiphlogistic/ analgetic agent for at least three days. Additionally, gerbils received an
antiemetic (oral; Emeprid, Ceva Tiergesundheit GmbH; 0.21 ml per gerbil/day) and, if
needed, also sterofundin (4 ml s.c.; B. Braun SE, Melsungen, Germany) or amynin (4 ml s.c.;
Boehringer Ingelheim Pharma GmbH Co KG, Ingelheim am Rhein, Germany) as infusion
solutions if gerbils were dehydrated, all post-operatively.

Ouabain treatment was carried out according to Bourien et al. (2014). After gerbils were
anesthetized, they received a sub-cutaneous infusion with sterofundin (4 ml). The location
where the headholder met the skin of the face, was treated with 2% xylocaine ointment
(Lidocainhydrochloride, Aspen Pharma Trading Limited, Dublin, Ireland). The skin covering
the bulla was disinfected with 70% ethanol. A small incision was placed behind the ear to
access the bulla. A hole was drilled into the bone with an angled blade and subsequently
widened with forceps to reach the round window. A syringe needle tip was placed onto the
round window to apply one of the following solutions onto the round window membrane:
Eight gerbils were treated with 40 µM ouabain (Ouabain octahydrate, Sigma-Aldrich, Saint
Louis, MO, USA), ten with 70 µM ouabain, dissolved in artificial perilymph

solution (137 mM NaCl, 5 mM KCl, 2 mM CaCl2, 1 mM MgCl2, 1 mM NaHCO3, and 11 mM
glucose; pH 7.4). Four gerbils received only artificial perilymph solution (surgery-only
treatment). In 15 gerbils, the solution was continuously delivered by a pump (150 µl for 30
Min). In seven gerbils, the round window niche was bathed with the solution by manually
operating the syringe and after ten minutes, the solution was aspirated and reapplied. This
procedure was repeated three times. Surgery was concluded by suturing the skin incision
over the open, middle ear. Gerbils were binaurally treated (n = 19), except those that were
destined for electrophysiology only, which were monaurally treated (n = 3).

Note that the experimenter conducting the electrophysiology and histology was blinded to the
type of treatment: the solution that was transferred to the round window was aliquoted by
another person. Only after the synapse counts were completed, the treatment was revealed.

790   Counting inner-hair-cell synapses

791   Immunohistochemistry

Processing of cochleae was carried out as described in detail in Steenken et al. (2022). After
concluding the single-unit recordings, gerbils were euthanized with an overdose of
pentobarbital (i.p.; Narcoren, Merial GmbH, Hallbergmoos, Germany, 480 mg/kg body
weigh). Twenty gerbils were cardiovascularly perfused with a mixture of phosphate-buffered
saline (PBS; 137 mM NaCl, 10 mM phosphate and 2.7 mM KCl, pH 7.4) followed by 4%
paraformaldehyde in PBS. To prevent blood coagulation, in 15 of these gerbils, heparin
(ratiopharm, 25,000 IE/5 ml; 0.2 ml/100 ml) was added to PBS. In 34 gerbils, the bulla was
rapidly exposed after euthanasia and small openings were carefully created at the apex and
base of the bony cochlear walls. Cochleae were then fixed in 4% paraformaldehyde in PBS
for two days on a shaker at 8 °C. After fixation, all cochleae were decalcified using 0.5 M
ethylenediaminetetraacetic acid for two days. Subsequently, cochleae were treated with 1%
Triton X-100-PBS for 1 hour to permeabilize the tissue and were then rinsed with 0.2% Triton
X-100-PBS. To block unspecific binding sites, cochleae were then incubated in 3% bovine
serum albumin (BSA, + 0.2% Triton X-100-PBS) blocking solution for 1 hour at room
temperature.

Antibodies to label hair cells (anti-MyosinVIIa, IgG polyclonal rabbit; Proteus Biosciences,
Ramona, CA, USA; diluted 1:400), presynaptic ribbons (anti-CtBP2 (C-terminal binding
protein) antibody, IgG1 monoclonal mouse; BD Biosciences, Eysins, Switzerland; diluted
1:400), and postsynapstic receptor patches (anti-GluR2a antibody, IgG2a monoclonal
mouse; Millipore, Burlington, MA, USA; diluted 1:200) in blocking solution were used.
Cochleae were incubated in this mixture at 37 °C overnight. Next, cochleae were again
rinsed in 0.2% Triton X-100-PBS.

Secondary antibodies were chosen to match the hosts of the primary antibodies: goat anti-
mouse (IgG1)-AF488 (Molecular Probes Inc., Eugene, OR, USA; diluted 1:1000), goat anti-

mouse (IgG2a)-AF568 (Invitrogen, Carlsbad, CA, USA; diluted 1:500), and donkey anti-rabbit

(IgG)-AF647 (Molecular Probes Inc.; diluted 1:1000). Secondary antibodies were diluted in

blocking solution and cochleae were incubated in this solution at 37 °C overnight, and

subsequently rinsed in PBS.

After the immunostaining, a subset of the cochleae from untreated animals older than 12

months received treatment with an auto-fluorescence quencher (n = 13; TrueBlack Lipofuscin

Autofluorescence Quencher, 20x in DMF, Biotum, Hayward, CA, USA). These cochleae were

cut in half and incubated in a 5% true black-70% ethanol-mixture for 1 minute and then

rinsed in PBS. Finally, all cochleae were micro-dissected under a stereomicroscope and the

resulting 5 - 11 organ-of-Corti pieces were mounted on slides using Vectashield Mounting

Medium (Vector Laboratories, Burlingame, CA, USA, H-1000).

Image acquisition

Overview images of cochlear pieces were acquired with a light microsope (Nikon 90i with

NIS Elements software, Version 4.30, Nikon, Minato, Tokio, Japan) using a 4x objective.

After measuring the length of the cochlea as a line along the row of IHCs with the NIS

software, the cochlear location at 3.8 mm from the apex, corresponding to 2 kHz (Müller

1996), was chosen, because 2 kHz was part of all stimulus conditions (data for additional

locations corresponding to 4, 8, and 16 kHz are provided in the Supplemental Materials).

Images were taken at this position with a confocal microscope (Leica Microsystem CMS

GmbH, Wetzlar, Germany, Leica TCS SP8 system) using an oil-immersion objective (40x,

numerical aperture 1.3). Lasers (488-nm and 522-nm optically pumped semiconductor laser),

or a 638-nm diode, respectively, excited the different fluorescence tags, and released

photons were counted with a hybrid detector. Image stacks had voxel dimensions of 0.05 -

0.0998 μm (XY) and 0.3 μm (Z). All confocal stacks were deconvolved (Huygens Essentials,

Version 15.10, SVI, Hilversum, Netherlands) with default settings (maximum iteration: 80;

signal to noise ratio: 10; quality threshold: 0.01), using a theoretical point-spread function.

Synapses were counted using ImageJ (FIJI, Schindelin et al. 2012). For each cochlear

position, the number of functional synapses was evaluated for 5 IHCs by manually counting

the co-localized ribbons and glutamate patches. Contrast and brightness were manually

adjusted for each stack.

Evoked potential recordings

Hearing sensitivity of all gerbils was evaluated prior to and during AN recordings using ABRs

and CAPs. For ABRs, a platinum needle electrode was placed subcutaneously near the

ipsilateral mastoid. For CAPs, a custom-made silver-wire ball electrode was placed at the

round window niche after opening the bulla. For both measurements, the reference electrode

was placed in the neck, and a ground electrode was placed on one hindleg. The stimulus

presentation was controlled via a custom-written MATLAB software (Mathworks, Natick, MA,
USA) and delivered through a RME Hammerfall DSP Multiface II sound card (RME Audio,
Haimhausen, Germany). The stimuli were amplified (HB7, TDT Inc.) and presented to the
gerbil's ear via earphones (IE 800, Sennheiser, Wedemark, Germany) attached to an ear-
bar.

Evoked potentials were amplified (x10^3) and band-pass filtered (ABR: 0.3 - 3 kHz; CAP:
0.005 - 10 kHz) using an ISO-80 pre-amplifier (World Precision Instruments, Sarasota, FL,
USA), and sampled using the digital signal processor (Hammerfall DSP Multiface II, RME
Audio; 48 kHz sampling rate) controlled by custom MATLAB software (Mathworks, Natick,
MA, USA).

Prior to ouabain treatment (CAP), and again prior to AN recordings (ABR and CAP),
responses to chirps were recorded at a range of levels, from below threshold up to 20 dB
above threshold, in steps of 5 dB, and averaged over 300 repetitions. During single-unit
recording sessions, ABR thresholds were re-checked periodically and visually evaluated, and
the experiment was terminated if threshold deteriorated by more than 30 dB. For post-hoc
analysis, CAP amplitudes (N1 – P1) were determined with the use of a custom-written
Matlab script. Threshold was defined as the first level step that exceeded a 2 µV CAP.

Single-unit recordings

Stimulus presentation

Stimuli for single-unit recordings were generated with custom-written MATLAB-scripts
controlling a digital signal processor (RX6, TDT Inc., Alachua, FL, USA) and an attenuator
(PA5, TDT Inc.).The signal was then routed through a headphone buffer (HB7, TDT Inc.) and
presented via earphones (IE 800, Sennheiser, Wedemark, Germany) attached to an ear-bar.
Stimuli were calibrated at the start of each experiment using a probe-tube microphone (ER7-
C, Etymotic Research Inc., Elk Grove Village, IL, USA) attached to the ear-bar and a
microphone amplifier (MA3, TDT Inc., 40 dB amplification).

Single-unit recordings

Single-unit data was obtained in 27 gerbils (see Table 1). After the gerbil was anesthetized, a
single dose of non-steroidal antiphlogistic agent (meloxicam, 0.2 mg/kg, Boehringer
Ingelheim Pharma GmbH Co KG) was administered subcutaneously. The head of the gerbil
was oriented with a bite bar and firmly fixed to the setup via a screw glued with dental
cement to the exposed skull. The pinna of one ear was removed to place an ear-bar directly
at the bony edge of the ear canal. The ear-bar was sealed with petroleum jelly to the bone to
form a closed sound system. To prevent pressure buildup in the middle ear, a small hole was
drilled into the dorsal bulla with an angled blade.

The AN was accessed via a dorsal approach. The skin covering the occipital bone and the

bone itself were removed and the lateral part of the cerebellum was aspirated. The brainstem

was left intact and was gently pushed medially using tiny saline soaked cotton balls to

access the proximal part of the eighth cranial nerve. A glass electrode (GB120F-10, Science

Products GmbH, Hofheim am Taunus, Germany; pulled using a P-2000, Sutter Instruments

Co., Novato, CA, USA), filled with 3 M KCl-solution and with a resistance of ~10 - 30 MΩ,

was placed above the AN under visual control. The electrode was advanced through the AN

via a remote-controlled piezo motor (Burleigh 6000 ULN inchworm motor controller and 6005

ULN handset, Burleigh, Inc., Fishers, NY, USA). Recordings were amplified (WPI 767, World

Precision Instruments Inc., Sarasota, FL, USA), filtered for line-frequency noise (50 Hz, Hum

Bug, Quest Scientific Instruments Inc., North Vancouver, BC, Canada), and digitized

(sampling rate: 48828 Hz; RX6, TDT Inc., Alachua, FL, USA). Additionally, the analog signal

was guided to an audio monitor (MS2, TDT Inc.) and displayed on an oscilloscope (SDS

1102CNL, SIGLENT Technologies, Hamburg, Germany).

Search stimuli (broadband noise, 1 - 9 kHz at 50 dB SPL) were played while the electrode

was advanced into the AN. When a unit was isolated, its best frequency (BF) was first

assessed audio-visually using 50-ms duration tones. Next, a response curve was obtained

by presenting pure tone bursts (50-ms duration, including 5-ms raised-cosine rise/fall times,

0 - 30 dB above threshold, 3 repetitions) at a range of frequencies within approximately

±1 octave around the audio-visually determined BF. Subsequently, tone bursts at BF were

presented over a range of levels (10 - 79 dB SPL, 50-ms duration including 5-ms raised-

cosine rise/fall times, 10 repetitions) to derive a rate-level function. At randomly inserted

trials, no stimulus was presented to determine the spontaneous rate (SR). Subsequently,

TFS1 stimuli were presented at a fixed level of 60 dB SPL.

Spike detection from the recorded signal was conducted off-line with a custom-written

MATLAB script (Steenken et al. 2022b). In short, the bandpass-filtered (300 – 3000 Hz)

recordings were manually screened for threshold-crossing events. The threshold could be

adjusted on a trial-by-trial basis, to compensate for variations in spike amplitude. The time of

each spike's peak amplitude was saved as the spike time. Spike trains were excluded if: 1.)

more than 10% of the spikes for a given stimulus condition were judged to be below the set

threshold criterion , or 2.) any inter-spike intervals < 0.6 ms occurred, indicating inadequate

unit isolation (Heil et al. 2007), or 3.) a pre-potential was present in the spike waveform,

suggesting a recording from the ventral cochlear nucleus (Keine and Rubsamen 2015), or 4.)

the rate-level function was non-monotonic, also suggesting a potential cochlear nucleus

recording (e.g., Davis et al. 1996, Joris et al. 1994, Sinex et al. 2001).

Neural recordings - Data Analysis

BF was defined as the frequency that evoked the highest mean discharge rate in the recorded response curve. Threshold at BF was determined from the rate-level curve, as the lowest stimulus level where the firing rate was higher than [mean spontaneous rate +1.2 times standard deviation of spontaneous rate] and higher than 15 spikes/s. The bandwidth of the response curve, at 50% of the maximal driven rate, was taken as a measure of neural tuning, but only if the response curve had been obtained at ≤ 20 dB above threshold.

A number of measures were derived from the AN responses to TFS1 stimuli: a) Peri-stimulus time histograms (PSTHs), b) rate histograms, and c) vector strengths (VS) corresponding to certain periodicities in the stimuli. Only the 375-ms long steady-state part of the responses, excluding a 12.5 ms onset and offset fringe, respectively, was analyzed.

First, peri-stimulus time histograms (PSTHs) were calculated for each animal/fiber/condition/Δf combination, pooling over repeated stimulus presentations within each combination. Spike times were collected into histograms using bin widths of 1/10 of the f_0 period (0.5 ms for 200 Hz and 0.25 ms for 400 Hz). The histogram counts were converted into instantaneous spike rates by dividing by the bin width and by the number of stimulus repetitions.

Second, average rate and SR histograms were calculated for each group/condition/Δf combination, pooling over repeated stimulus presentations, animals, and fibers within each combination. Average rates were computed over the 375-ms steady-state portion of the response. SR was calculated from trials without stimulation. Driven rate was calculated as the difference between stimulated absolute rate and SR. Average driven rates and SRs were collected into histograms using bin widths of 2.5 spikes per second and 5 spikes per second, respectively. The average-driven-rate histogram counts were divided by the number of spike trains to convert the counts to relative frequency and smoothed with a moving-average window of 3 bin lengths.

Third, VS frequency spectra were calculated for each group/condition/Δf combination, pooling over repeated stimulus presentations, animals, and fibers within each combination. For this, all-order inter-spike intervals (ISIs) were calculated from each spike train, and collected into histograms using a bin width of 50 μs, corresponding to a sampling rate of 20 kHz. The bin positions ranged from -375 ms to +375 ms. A zero-centered gaussian window with a standard deviation of 200 ms was multiplied with the histogram to suppress fringe effects and spurious side lobes in the spectra. The histogram counts were divided by the total number of ISIs. A subsequent fast Fourier transform of the histograms resulted in the ISI power spectrum, and the square root of the spectral magnitudes was identical to the VS at each bin center frequency. The spectral resolution was 1.33 Hz.

958   VS single-peak data was calculated for each animal/fiber/condition/Δf combination, pooling
959   over repeated stimulus presentations within each combination. In contrast to the spectra, the
960   peak VS values were calculated individually for each relevant frequency f:

961
$$V(f) = \left| \frac{1}{N} \sum_{i=1}^{N} e^{-i2\pi f t_i} \right| ,$$

where t_i are the spike times, and N is the number of spikes in each spike train. For
significance analysis of the VS, the Rayleigh statistic z (henceforth referred to as z-value)
was calculated: $z = N_{avg} \cdot V_{avg}^2(f)$, where N_{avg} is the average number of spikes per spike train
and V_{avg} is the average VS across stimulus presentations. The Rayleigh z statistic has useful
properties for statistical analyses because VS values resulting from higher numbers of spikes
have larger weights, and logarithms of ratios of z-values are approximately normally
distributed, as opposed to VS ratios or differences.

As a metric for TFS representation, the log ratio between two z-values was derived: First,
from responses to the H complex, the z-value, z_0, at the frequency component that the fibers
on average responded to maximally, $f_{maxpeak}$, was obtained. Second, this value was divided
by the z-value, $z_{\Delta f}$, at the shifted frequency of the I complex, also in the maximal average
response range (filled circles in Fig. 11). This ratio was used as the dependent variable in the
first ANOVA model and reflects high temporal resolution; large positive values correspond to
stronger phase-locked responses to the TFS of the stimulus. This metric was termed the TFS
log-z-ratio.

Another log ratio between z-values was used as a metric for the balance between envelope
and TFS representation (crosses in Fig. 11): The (harmonic, unshifted) z_0 value at the
fundamental frequency, f_0, was divided by the (inharmonic, shifted) $z_{\Delta f}$ value in the maximal
average response range, $f_{maxpeak}$. Large positive log ratios indicate a stronger response to the
stimulus envelope, whereas large negative log ratios indicate a stronger response to the
stimulus TFS. This metric was termed ENV/TFS log-z-ratio.

Figure 11: Schematic representation of statistical contrasts. *Schematic of the frequency*
spectrum of a harmonic (black solid lines) and inharmonic (red dashed lines) TFS1 stimulus
in panel A and schematic representation of the phase locking spectrum in a neuronal
response in panel B. Black arrows and symbols point to the harmonic, unshifted frequencies,
termed "z_0" and red arrows and symbols point to the inharmonic, shifted frequencies, termed
"z_{df}". The annotated brackets below the frequency axis mark representative regions of the
fundamental frequency, f_0, and of the maximal average response, $f_{maxpeak}$. Filled connected
circles show the log-z-ratio used for assessing the TFS representation, connected crosses
show the log-z-ratio used for assessing the balance between f_0/ENV and TFS representation.
The representation of Δf shift in the responses was also tested by calculating z-values at the
different harmonic frequencies, combined with all possible Δf shifts. The actual position of the
maximum z-value in terms of Δf and the corresponding z-value is summarized in Fig. 4 for
two TFS peak frequencies (to which the Δf shifts were added): at f_0 of the corresponding
condition and at 1 kHz for the 200/1600 Hz condition, at 800 Hz for the 400/1600 Hz
condition, and at 2 kHz for the 400/3200 Hz condition, respectively. The latter TFS peak

frequencies were identified as the harmonics actually present in the acoustic stimulus, which
had the maximum z-value in the summarized spectra (Fig. 4).

For the single-unit data, several main factors were defined. The low-SR fiber class comprised
fibers with SR ≤ 18 spikes/s, and the high-SR class comprised fibers with SR > 18 spikes/s.
In addition, fibers in the low-BF class had BF ≤ 1850 Hz, and fibers in the high-BF class had
BF > 1850 Hz.

Behavioral procedure

The behavioral experiments were conducted in a single-walled sound-attenuating chamber
(Industrial Acoustics, Type 401 A, Industrial Acoustic Company GmbH, Mönchengladbach,
Germany) lined with a 15-cm thick layer of sound- absorbing foam (Pinta Acoustics Pyramide
100/50 on Pinta Acustics PLANO Type 50/0, Seyboth & Co., Regensburg, Germany). The
walls and all devices within the chamber produced no relevant sound reflections. The time for
the reverberation in the chamber to decay by 60 dB was 12 ms, indicating nearly anechoic
conditions. These conditions allowed faithful presentation of the TFS1 stimuli with the desired
acoustic cues in the free sound field.

In the center of the chamber, a 30-cm-long wire-mesh platform was located at 90-cm height,
with an elevated pedestal at its center where the gerbils waited during the measurement.
This construction minimized sound reflections. At one end of the platform, a food bowl was
located, attached via a tube to a custom-made feeder that did not obstruct the sound path.
Gerbils were rewarded for correct discrimination with a 10-mg custom-made food pellet
(Altromin International, type 1324 rodent pellets enriched with sunflower oil and spelt flour;
Altromin Spezialfutter GmbH & Co. KG, Lage, Germany). A loudspeaker (Canton Plus XS,
frequency range: 150 Hz - 21 kHz, Canton, Weilrod, Germany) was mounted 30 cm in front
of the pedestal to the side of the food bowl, at 0° elevation and azimuth relative to the gerbil's
normal head position when sitting on the pedestal. Out of the gerbil's reach, a system of two
custom-made light barriers was installed to detect the gerbil's position and facing direction on
the pedestal. An infrared camera (Conrad 150001 C-MOS camera module, Conrad
Electronics, Hirschau, Germany) was positioned above the platform to observe the gerbil's
movements on the platform under invisible infrared light. The stimulus presentation,
registration of light barriers switching, and feeder were controlled by custom-written software
on a Linux-based PC with an RME sound-card (Hammerfall DSP Multiface II, sampling
frequency 48 kHz). The signal output from the sound card was manually attenuated (Texio
type RA-902A, TEXIO Technology Corporation, Kanagawa, Japan), amplified (Rotel type
RMB 1506, Rotel Tokyo, Japan), and presented by the loudspeaker. Before testing started
on each day, the system was calibrated (±1 dB) with a sound level meter (Brüel and Kjaer
type 2238 Mediator, Hottinger Brüel & Kjær, Virum, Denmark) positioned on the elevated
pedestal at about the position of the gerbil's head.

The behavioral experiment used an operant Go/NoGo paradigm. Gerbils were trained to wait

on the pedestal facing in the direction of the loudspeaker. During the whole session, the H

complex was played every 1.3 s as a reference stimulus. As soon as the gerbil jumped onto

the pedestal facing the loudspeaker, a trial was started with a random waiting time between 2

and 7 s. When the waiting time had elapsed, a target stimulus, the I complex, was played

instead of the reference H complex. The target stimuli could either be equal to the reference

H complex (sham trial) for estimating the false-alarm rate or an I complex (test trial) for

testing discrimination ability. Leaving the pedestal before the waiting time elapsed resulted in

the restart of the trial with a new waiting time. Leaving the pedestal within 1.3 s after the

onset of the target stimulus of a test trial, as indicated by the light barriers, was registered as

a "hit" and resulted in a food reward. Staying on the pedestal in a test trial was counted as a

"miss" with no food reward. The animal had to leave the pedestal and jump back onto it in the

correct sequence to start a new trial after a "miss" occurred. Leaving the pedestal within 1.3 s

after the start of the target stimulus in a sham trial was registered as a "false alarm", while

staying on the pedestal during a sham trial was registered as a "correct rejection". No food

reward was provided in sham trials. To keep the gerbils motivated during the session, a

salient test trial not included in the analysis was inserted after each correct rejection.

Each session contained 10 blocks with 9 trials each (six test trials, three sham trials). All

stimuli within a given block were from the same condition (200/1600 Hz, 400/1600 Hz, or

400/3200 Hz). The first block of a session was a warmup block with $\Delta f = 42.5\%$ that was not

included in the analysis. To interleave simple and more difficult conditions, blocks of the

different conditions were pseudo-randomly distributed throughout the session. The least

salient condition (200/1600 Hz) did not occur within two consecutive blocks. Additionally, the

sham and test trials within each block were pseudo-randomly distributed with the constraint

that no sham trials followed one another within and across blocks. Data collection started

when the gerbil achieved a false alarm rate below 20% in three consecutive sessions with hit

rates for the three highest Δf values that would allow determining a discrimination threshold.

Sessions were included in the analysis if the gerbil completed all 90 trials and the false alarm

rate did not exceed 20%. Data collection continued until the sample size for each condition

and Δf value was at least 20 trials. The sensitivity for each Δf value was defined by applying

the z-transform, $d' = z(\text{hit rate}) - z(\text{false-alarm rate})$. Individual discrimination thresholds for

each animal and condition were then calculated from sensitivity, d', as a function of

inharmonic frequency shift, $\Delta f\%$. A threshold at $d' = 1$ was obtained by linearly interpolating

between the first $\Delta f\%$ shift where $d' > 1$ and the previous $\Delta f\%$ shift with $d' \leq 1$. If all d' values

were below 1, a linear fit was estimated through all d' values as a function of $\Delta f\%$ shifts and

threshold was extrapolated at $d' = 1$. If all d' values were larger than 1, the threshold was set

to half of the minimal $\Delta f\%$ shift measured in the specific experiment. If none of the above was

possible and either all $d' < 1$, the fitted slope was not positive, or the extrapolated threshold

larger than 100% shift (equivalent to $\Delta f = f_0$ at threshold), the final threshold was limited to

100% (5 out of 72 thresholds).

Statistical analysis

All analyses of variance were calculated with IBM SPSS Statistics for Windows, Version 29.0

(IBM Corp, Armonk, NY). The family-wise error rate of all multiple post-hoc comparisons was

controlled by Bonferroni correction of the significance levels.

Supporting information

Data availability

The data that are presented in the figures of the present manuscript and were statistically

analyzed are uplodaded to a Zenodo repository (https://doi.org/10.5281/zenodo.15546625).

References

Adams, J. C., & Schulte, B. A. (1997). Histopathologic observations of the aging gerbil
cochlea. Hearing Research, 104, 101-111. https://doi.org/10.1016/S0378-
5955(96)00184-0
Bharadwaj, H. M., Verhulst, S., Shaheen, L., Liberman, M. C., & Shinn-Cunningham, B. G.
(2014). Cochlear neuropathy and the coding of supra-threshold sound. *Frontiers in
Systems Neuroscience*, *8*, 26. https://doi.org/10.3389/fnsys.2014.00026
Bourien, J., Tang, Y., Batrel, C., Huet, A., Lenoir, M., Ladrech, S., Desmadryl, G., Nouvian,
R., Puel, J.-L., & Wang, J. (2014). Contribution of auditory nerve fibers to compound
action potential of the auditory nerve. *Journal of Neurophysiology*, *112*(5), 1025–1039.
https://doi.org/10.1152/jn.00738.2013
Bovee, S., Klump, G. M., Pyott, S. J., Sielaff, C., & Köppl, C. (2024). Cochlear ribbon synapses
in aged gerbils. *International Journal of Molecular Sciences*, *25*(5).
https://doi.org/10.3390/ijms25052738
Bramhall, N., Beach, E. F., Epp, B., Le Prell, C. G., Lopez-Poveda, E. A., Plack, C. J., Schaette,
R., Verhulst, S., & Canlon, B. (2019). The search for noise-induced cochlear
synaptopathy in humans: Mission impossible? *Hearing Research*, *377*, 88–103.
https://doi.org/https://doi.org/10.1016/j.heares.2019.02.016
Bramhall, N. F., McMillan, G. P., Kujawa, S. G., & Konrad-Martin, D. (2018). Use of non-
invasive measures to predict cochlear synapse counts. *Hearing Research*, *370*, 113–119.
https://doi.org/https://doi.org/10.1016/j.heares.2018.10.006
Buran, B. N., Strenzke, N., Neef, A., Gundelfinger, E. D., Moser, T., & Liberman, M. C.
(2010). Onset coding is degraded in auditory nerve fibers from mutant mice lacking
synaptic ribbons. *The Journal of Neuroscience*, *30*(22), 7587.
https://doi.org/10.1523/JNEUROSCI.0389-10.2010
Burton, J. A., Mackey, C. A., MacDonald, K. S., Hackett, T. A., & Ramachandran, R. (2020).
Changes in audiometric threshold and frequency selectivity correlate with cochlear
histopathology in macaque monkeys with permanent noise-induced hearing loss.
Hearing Research, *398*, 108082.
https://doi.org/https://doi.org/10.1016/j.heares.2020.108082
Cai, R., Montgomery, S. C., Graves, K. A., Caspary, D. M., & Cox, B. C. (2018). The FBN rat
model of aging: investigation of ABR waveforms and ribbon synapse changes.

Neurobiology of Aging, *62*, 53–63.
https://doi.org/https://doi.org/10.1016/j.neurobiolaging.2017.09.034

Carcagno, S., & Plack, C. J. (2022). Relations between speech-reception, psychophysical temporal processing, and subcortical electrophysiological measures of auditory function in humans. *Hearing Research*, *417*, 108456.
https://doi.org/https://doi.org/10.1016/j.heares.2022.108456

Caspary, D. M., Milbrandt, J. C., & Helfert, R. H. (1995). Central auditory aging: GABA changes in the inferior colliculus. *Experimental Gerontology*, *30*(3–4), 349–360.
https://doi.org/10.1016/0531-5565(94)00052-5

Castaño-González, K., Köppl, C., & Pyott, S.J. (2024). The crucial role of diverse animal models to investigate cochlear aging and hearing loss. *Hearing Research 445*: 108989.
https://doi.org/10.1016/j.heares.2024.108989

Davis, K. A., Ding, J., Benson, T. E., & Voigt, H. F. (1996). Response properties of units in the dorsal cochlear nucleus of unanesthetized decerebrate gerbil. *Journal of Neurophysiology*, *75*(4), 1411–1431. https://doi.org/10.1152/jn.1996.75.4.1411

Eipert, L., Selle, A., & Klump, G. M. (2019). Uncertainty in location, level and fundamental frequency results in informational masking in a vowel discrimination task for young and elderly subjects. *Hearing Research*, *377*, 142–152.
https://doi.org/https://doi.org/10.1016/j.heares.2019.03.015

Faulstich, M., & Kössl, M. (1999). Neuronal response to cochlear distortion products in the anteroventral cochlear nucleus of the gerbil. *The Journal of the Acoustical Society of America*, *105*(1), 491–502. https://doi.org/10.1121/1.424586

Füllgrabe, C., Moody, M., & Moore, B. C. J. (2020). No evidence for a link between noise exposure and auditory temporal processing for young adults with normal audiograms. *The Journal of the Acoustical Society of America*, *147*(6), EL465–EL470.
https://doi.org/10.1121/10.0001346

Füllgrabe, C., Moore, B. C. J., & Stone, M. A. (2015). Age-group differences in speech identification despite matched audiometrically normal hearing: contributions from auditory temporal processing and cognition. *Frontiers in Aging Neuroscience*, *6*, 347.
https://doi.org/10.3389/fnagi.2014.00347

Gleich, O., Hamann, I., Klump, G. M., Kittel, M., & Strutz, J. (2003). Boosting GABA improves impaired auditory temporal resolution in the gerbil. In *Neuroreport* (Vol. 14, Issue 14, pp. 1877–1880). https://doi.org/10.1097/00001756-200310060-00024

Gleich, O., Netz, J., & Strutz, J. (2014). Comparing the inferior colliculus of young and old gerbils (Meriones unguiculatus) with an emphasis on GABA. *Experimental Gerontology*, 57, 155-162. https://doi.org/10.1016/j.exger.2014.05.018

Gleich, O., Semmler, P., & Strutz, J. (2016). Behavioral auditory thresholds and loss of ribbon synapses at inner hair cells in aged gerbils. *Experimental Gerontology*, *84*, 61–70. https://doi.org/10.1016/j.exger.2016.08.011

Gómez-Álvarez, M., Johannesen, P. T., Coelho-de-Sousa, S. L., Klump, G. M., & Lopez-Poveda, E. A. (2023). The relative contribution of cochlear synaptopathy and reduced inhibition to age-related hearing impairment for people with normal audiograms. *Trends in Hearing*, *27*, 23312165231213190.
https://doi.org/10.1177/23312165231213191

Gratton, M. A., & Schulte, B. A. (1995). Alterations in microvasculature are associated with atrophy of the stria vascularis in quiet-aged gerbils. *Hearing Research*, *82*(1), 44–52.
https://doi.org/https://doi.org/10.1016/0378-5955(94)00161-I

Guest, H., Munro, K. J., Prendergast, G., Millman, R. E., & Plack, C. J. (2018). Impaired speech perception in noise with a normal audiogram: No evidence for cochlear synaptopathy and no relation to lifetime noise exposure. *Hearing Research*, *364*, 142–151. https://doi.org/https://doi.org/10.1016/j.heares.2018.03.008

Guinan, J. J. (2011). Physiology of the medial and lateral olivocochlear systems. In D. Ryugo, R. R. Fay, & A. N. Popper (Eds.), Auditory and Vestibular Efferents (pp. 39-81). Springer Science+Business Media, LLC. https://doi.org/10.1007/978-1-4419-7070-1_3

Hamann, I., Gleich, O., Klump, G. M., Kittel, M. C., Boettcher, F. A., Schmiedt, R. A., & Strutz, R. (2002). Behavioral and evoked-potential thresholds in young and old Mongolian gerbils (Meriones unguiculatus). Hearing Research, 171(1-2), 82-95.

https://doi.org/10.1016/S0378-5955(02)00454-9

Harrison, R. V, & Evans, E. F. (1979). Some aspects of temporal coding by single cochlear
fibres from regions of cochlear hair cell degeneration in the guinea pig. *Archives of Oto-*
Rhino-Laryngology, *224*(1), 71–78. https://doi.org/10.1007/BF00455226

Heeringa AN. (2024). Single-unit data for sensory neuroscience: Responses from the
auditory nerve of young-adult and aging gerbils. *Scientific Data*, 11(1):411. doi:
10.1038/s41597-024-03259-3

Heeringa, A. N., Jüchter, C., Beutelmann, R., Klump, G. M., & Köppl, C. (2023). Altered
neural encoding of vowels in noise does not affect behavioral vowel discrimination in
gerbils with age-related hearing loss. *Frontiers in Neuroscience*, *17*.
https://www.frontiersin.org/journals/neuroscience/articles/10.3389/fnins.2023.12389
41

Heeringa, A. N., Zhang, L., Ashida, G., Beutelmann, R., Steenken, F., & Koeppl, C. (2020).
Temporal coding of single auditory nerve fibers is not degraded in aging gerbils. *The*
Journal of Neuroscience, *40*(2), 343–354. https://doi.org/10.1523/JNEUROSCI.2784-
18.2019

Heil, P., Neubauer, H., Irvine, D. R. F., & Brown, M. (2007). Spontaneous activity of
auditory-nerve fibers: insights into stochastic processes at ribbon synapses. *The Journal*
of Neuroscience, *27*(31), 8457–8474.

Hellstrom, L. I., & Schmiedt, R. A. (1990). Compound action potential input/output
functions in young and quiet-aged gerbils. *Hearing Research*, *50*(1), 163–174.
https://doi.org/https://doi.org/10.1016/0378-5955(90)90042-N

Hellstrom, L. I., & Schmiedt, R. A. (1996). Measures of tuning and suppression in single-fiber
and whole-nerve responses in young and quiet-aged gerbils. *The Journal of the*
Acoustical Society of America, *100*(5), 3275–3285. https://doi.org/10.1121/1.417211

Henry, K. S. (2022). Animal models of hidden hearing loss: Does auditory-nerve-fiber loss
cause real-world listening difficulties? *Molecular and Cellular Neuroscience*, *118*,
103692. https://doi.org/https://doi.org/10.1016/j.mcn.2021.103692

Henry, K. S., & Abrams, K. S. (2021). Normal tone-in-noise sensitivity in trained budgerigars
despite substantial auditory-nerve injury: No evidence of hidden hearing loss. *The*
Journal of Neuroscience, *41*(1), 118 LP – 129.
https://doi.org/10.1523/JNEUROSCI.2104-20.2020

Henry, K. S., & Heinz, M. G. (2012). Diminished temporal coding with sensorineural hearing
loss emerges in background noise. *Nature Neuroscience*, *15*(10), 1362–1364.
https://doi.org/10.1038/nn.3216

Henry, K. S., Kale, S., & Heinz, M. G. (2016). Distorted Tonotopic Coding of Temporal
Envelope and Fine Structure with Noise-Induced Hearing Loss. J Neurosci, 36(7), 2227-
2237. https://doi.org/10.1523/JNEUROSCI.3944-15.2016

Henry, K. S., Sayles, M., Hickox, A. E., & Heinz, M. G. (2019). Divergent auditory nerve
encoding deficits between two common etiologies of sensorineural hearing loss. *The*
Journal of Neuroscience, *39*(35), 6879 LP – 6887.
https://doi.org/10.1523/JNEUROSCI.0038-19.2019

Jeng, J., Ceriani, F., Olt, J., Brown, S. D. M., Holley, M. C., Bowl, M. R., Johnson, S. L., &
Marcotti, W. (2020). Pathophysiological changes in inner hair cell ribbon synapses in
the ageing mammalian cochlea. *Journal of Physiology*, *598*(19), 4339–4355.
https://doi.org/10.1113/JP280018

Johannesen, P. T., Buzo, B. C., & Lopez-Poveda, E. A. (2019). Evidence for age-related
cochlear synaptopathy in humans unconnected to speech-in-noise intelligibility deficits.
Hearing Research, *374*, 35–48.
https://doi.org/https://doi.org/10.1016/j.heares.2019.01.017

Joris, P. X., Schreiner, C. E., & Rees, A. (2004). Neural processing of amplitude-modulated
sounds. *Physiological Reviews*, *84*(2), 541–577.
https://doi.org/10.1152/physrev.00029.2003

Joris, P. X., Smith, P. H., & Yin, T. C. T. (1994). Enhancement of neural synchronization in
the anteroventral cochlear nucleus. II. Responses in the tuning curve tail. *Journal of*
Neurophysiology, *71*(3), 1037–1051. https://doi.org/10.1152/jn.1994.71.3.1022

Joris, P. X., & Yin, T. C. T. (1992). Responses to amplitude-modulated tones in the auditory

nerve of the cat. *The Journal of the Acoustical Society of America*, *91*(1), 215–232.
https://doi.org/10.1121/1.402757
Kale, S., & Heinz, M. G. (2010). Envelope coding in auditory nerve fibers following noise-
induced hearing loss. *Journal of the Association for Research in Otolaryngology :*
JARO, *11*(4), 657–673. https://doi.org/10.1007/s10162-010-0223-6
Kale, S., Micheyl, C., & Heinz, M. G. (2013). Effects of sensorineural hearing loss on temporal
coding of harmonic and inharmonic tone complexes in the auditory nerve. *Advances in*
Experimental Medicine and Biology, *787*, 109–118. https://doi.org/10.1007/978-1-
4614-1590-9_13
Kale, S., Micheyl, C., & Heinz, M. G. (2014). Implications of within-fiber temporal coding for
perceptual studies of F0 discrimination and discrimination of harmonic and
inharmonic tone complexes. *Journal of the Association for Research in*
Otolaryngology : JARO, *15*(3), 465–482. https://doi.org/10.1007/s10162-014-0451-2
Keine, C., & Ruebsamen, R. (2015). Inhibition shapes acoustic responsiveness in spherical
bushy cells. *The Journal of Neuroscience 35*(22), 8579–8592.
https://doi.org/10.1523/JNEUROSCI.0133-15.2015
Kessler, M., Mamach, M., Beutelmann, R., Lukacevic, M., Eilert, S., Bascuñana, P., Fasel, A.,
Bengel, F. M., Bankstahl, J. P., Ross, T. L., Klump, G. M., & Berding, G. (2020).
GABA(A) receptors in the Mongolian gerbil: a PET study using [(18)F]flumazenil to
determine receptor binding in young and old animals. *Molecular Imaging and Biology*,
22(2), 335–347. https://doi.org/10.1007/s11307-019-01371-0
Kittel, M., Wagner, E., & Klump, G. M. (2002). An estimate of the auditory-filter bandwidth
in the Mongolian gerbil. *Hearing Research*, *164*(1), 69–76.
https://doi.org/https://doi.org/10.1016/S0378-5955(01)00411-7
Kobrina, A., Schrode, K. M., Screven, L. A., Javaid, H., Weinberg, M. M., Brown, G., Board,
R., Villavisanis, D. F., Dent, M. L., & Lauer, A. M. (2020). Linking anatomical and
physiological markers of auditory system degeneration with behavioral hearing
assessments in a mouse (Mus musculus) model of age-related hearing loss. Neurobiol
Aging, 96, 87-103. https://doi.org/10.1016/j.neurobiolaging.2020.08.012
Kujawa, S. G., & Liberman, M. C. (2009). Adding insult to injury: Cochlear nerve
degeneration after "temporary" noise-induced hearing loss. *The Journal of*
Neuroscience, *29*(45), 14077–14085. https://doi.org/10.1523/JNEUROSCI.2845-
09.2009
Lang, H., Schulte, B. A., & Schmiedt, R. A. (2005). Ouabain induces apoptotic cell death in
type I spiral ganglion neurons, but not type II neurons. *Journal of the Association for*
Research in Otolaryngology : JARO, *6*(1), 63–74. https://doi.org/10.1007/s10162-004-
5021-6
Lauer, A. M., Jimenez, S. V., & Delano, P. H. (2022). Olivocochlear efferent effects on
perception and behavior. Hearing Research, 419.
https://doi.org/10.1016/j.heares.2021.108207
Liberman, M. C. (2017). Noise-induced and age-related hearing loss: new perspectives and
potential therapies [version 1; peer review: 4 approved]. *F1000Research*, *6*(927).
https://doi.org/10.12688/f1000research.11310.1
Liberman, M. C., Epstein, M. J., Cleveland, S. S., Wang, H., & Maison, S. F. (2016). Toward a
differential diagnosis of hidden hearing loss in humans. *PLOS ONE*, *11*(9), e0162726-.
https://doi.org/10.1371/journal.pone.0162726
Liberman, M. C., & Kujawa, S. G. (2017). Cochlear synaptopathy in acquired sensorineural
hearing loss: Manifestations and mechanisms. *Hearing Research*, *349*, 138–147.
https://doi.org/10.1016/j.heares.2017.01.003
Lin, H. W., Furman, A. C., Kujawa, S. G., & Liberman, M. C. (2011). Primary Neural
Degeneration in the Guinea Pig Cochlea After Reversible Noise-Induced Threshold
Shift. *Journal of the Association for Research in Otolaryngology*, *12*(5), 605–616.
https://doi.org/10.1007/s10162-011-0277-0
Lopez-Poveda, E. A. (2014). Why do I hear but not understand? Stochastic undersampling as
a model of degraded neural encoding of speech. *Frontiers in Neuroscience, 8*, 348.
https://doi.org/10.3389/fnins.2014.00348
Mackey, C. A., McCrate, J., MacDonald, K. S., Feller, J., Liberman, L., Liberman, M. C.,

Hackett, T. A., & Ramachandran, R. (2021). Correlations between cochlear
pathophysiology and behavioral measures of temporal and spatial processing in noise
exposed macaques. *Hearing Research*, *401*, 108156.
https://doi.org/https://doi.org/10.1016/j.heares.2020.108156
Marmel, F., Plack, C. J., Hopkins, K., Carlyon, R. P., Gockel, H. E., & Moore, B. C. J. (2015).
The role of excitation-pattern cues in the detection of frequency shifts in bandpass-
filtered complex tones. *The Journal of the Acoustical Society of America*, *137*(5), 2687–
2697. https://doi.org/10.1121/1.4919315
Mathew, A. K., Purdy, S. C., Welch, D., Pontoppidan, N. H., & Rønne, F. M. (2016).
Electrophysiological and behavioural processing of complex acoustic cues. *Clinical*
Neurophysiology, *127*(1), 779–789. https://doi.org/10.1016/j.clinph.2015.04.002
Miller, R. L., Schilling, J. R., Franck, K. R., & Young, E. D. (1997). Effects of acoustic trauma
on the representation of the vowel /ɛ/ in cat auditory nerve fibers. *The Journal of the*
Acoustical Society of America, *101*(6), 3602–3616. https://doi.org/10.1121/1.418321
Moore, B. C., Glasberg, B. R., & Hopkins, K. (2006). Frequency discrimination of complex
tones by hearing-impaired subjects: Evidence for loss of ability to use temporal fine
structure. *Hearing Research*, *222*(1–2), 16–27.
https://doi.org/10.1016/j.heares.2006.08.007
Moore, B. C. J. (2014). Auditory processing of temporal fine structure: effects of age and
hearing loss. In *Auditory processing of temporal fine structure*. World Scientific
Publishing Co. Pte. Ltd.
Moore, B. C. J. (2019). The roles of temporal envelope and fine structure information in
auditory perception. *Acoustical Science and Technology*, *40*(2), 61–83.
https://doi.org/10.1250/ast.40.61
Moore, B. C. J., Glasberg, B. R., Stoev, M., Fuellgrabe, C., & Hopkins, K. (2012). The
influence of age and high-frequency hearing loss on sensitivity to temporal fine
structure at low frequencies (L). *The Journal of the Acoustical Society of America*,
131(2, 1), 1003–1006. https://doi.org/10.1121/1.3672808
Moore, B. C. J., & Sek, A. (2009). Development of a fast method for determining sensitivity
to temporal fine structure. *International Journal of Audiology*, *48*(4), 161–171.
https://doi.org/10.1080/14992020802475235
Moore, G. A., & Moore, B. C. J. (2003). Perception of the low pitch of frequency-shifted
complexes. *The Journal of the Acoustical Society of America*, *113*(2), 977–985.
https://doi.org/10.1121/1.1536631
Müller, M. (1996). The cochlear place-frequency map of the adult and developing mongolian
gerbil. *Hearing Research*, *94*(1–2), 148–156. https://doi.org/10.1016/0378-
5955(95)00230-8
National Institute of Mental Health. (2002). *Methods and Welfare Considerations in*
Behavioral Research with Animals: Report of a National Institutes of Health
Workshop. (A. Morrison, H. Evans, N. Ator, & R. Nakamura, Eds.). NIH Publication No.
02-5083.
O'Brien, W. J., Lingrel, J. B., & Wallick, E. T. (1994). Ouabain binding kinetics of the rat
alpha two and alpha three isoforms of the sodium-potassium adenosine triphosphate.
Archives of Biochemistry and Biophysics 310: 32–39.
https://doi.org/10.1006/abbi.1994.1136
Parthasarathy, A., Hancock, K. E., Bennett, K., DeGruttola, V., & Polley, D. B. (2020).
Bottom-up and top-down neural signatures of disordered multi-talker speech
perception in adults with normal hearing. *ELife*, *9*, e51419.
https://doi.org/10.7554/eLife.51419
Parthasarathy, A., & Kujawa, S. G. (2018). Synaptopathy in the aging cochlea: characterizing
early-neural deficits in auditory temporal envelope processing. *The Journal of*
Neuroscience, *38*(32), 7108–7119. https://doi.org/10.1523/JNEUROSCI.3240-17.2018
Prendergast, G., Couth, S., Millman, R. E., Guest, H., Kluk, K., Munro, K. J., & Plack, C. J.
(2019). Effects of age and noise exposure on proxy measures of cochlear synaptopathy.
Trends in Hearing, *23*, 2331216519877301. https://doi.org/10.1177/2331216519877301
Prendergast, G., Millman, R. E., Guest, H., Munro, K. J., Kluk, K., Dewey, R. S., Hall, D. A.,
Heinz, M. G., & Plack, C. J. (2017). Effects of noise exposure on young adults with

normal audiograms II: Behavioral measures. *Hearing Research*, *356*, 74–86. https://doi.org/https://doi.org/10.1016/j.heares.2017.10.007

Romero, G. E., & Trussell, L. O. (2022). Central circuitry and function of the cochlear efferent systems. Hear Res, 425, 108516. https://doi.org/10.1016/j.heares.2022.108516

Rose, J. E., Brugge, J. F., Anderson, D. J., & Hind, J. E. (1967). Phase-locked response to low-frequency tones in single auditory nerve fibers of the squirrel monkey. *Journal of Neurophysiology*, *30*(4), 769–793. https://doi.org/10.1152/jn.1967.30.4.769

Ruggero, M. A., & Rich, N. C. (1991). Furosemide alters organ of corti mechanics: Evidence for feedback of outer hair cells upon the basilar membrane. The Journal of Neuroscience, 11(4), 1057-1067.

Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T., Preibisch, S., Rueden, C., Saalfeld, S., Schmid, B., Tinevez, J.-Y., White, D. J., Hartenstein, V., Eliceiri, K., Tomancak, P., & Cardona, A. (2012). Fiji: an open-source platform for biological-image analysis. *Nature Methods*, *9*(7), 676–682. https://doi.org/10.1038/nmeth.2019

Schmiedt, R. A., Mills, J. H., & Boettcher, F. A. (1996). Age-related loss of activity of auditory-nerve fibers. *Journal of Neurophysiology*, *76*(4), 2799–2803. https://doi.org/10.1152/jn.1996.76.4.2799

Schmiedt, R. A., Okamura, H., Lang, H., & Schulte, B. A. (2002). Ouabain application to the round window of the gerbil cochlea: A model of auditory neuropathy and apoptosis. *Journal of the Association for Research in Otolaryngology : JARO.*, *3*(3), 223–233. https://doi.org/10.1007/s101620020017

Schofield, B. R. (2011). Central descending auditory pathways. In D. Ryugo, R. R. Fay, & A. N. Popper (Eds.), Auditory and Vestibular Efferents (pp. 261-290). Springer Science+Business Media, LLC. https://doi.org/10.1007/978-1-4419-7070-1_9

Schulte, B. A., & Schmiedt, R. A. (1992). Lateral wall Na,K-ATPase and endocochlear potentials decline with age in quiet-reared gerbils. *Hearing Research*, *61*(1–2), 35–46. https://doi.org/10.1016/0378-5955(92)90034-k

Sergeyenko, Y., Lall, K., Liberman, M. C., & Kujawa, S. G. (2013). Age-related cochlear synaptopathy: An early-onset contributor to auditory functional decline. *The Journal of Neuroscience*, *33*(34), 13686–13694. https://doi.org/10.1523/JNEUROSCI.1783-13.2013

Shi, L., Chang, Y., Li, X., Aiken, S. J., Liu, L., & Wang, J. (2016). Coding deficits in noise-induced hidden hearing loss may stem from incomplete repair of ribbon synapses in the cochlea. *Frontiers in Neuroscience, 10*. https://www.frontiersin.org/journals/neuroscience/articles/10.3389/fnins.2016.00231

Sinex, D. G., López, D. E., & Warr, W. B. (2001). Electrophysiological responses of cochlear root neurons. *Hearing Research*, *158*(1–2), 28–38. https://doi.org/10.1016/S0378-5955(01)00293-3

Sinnott JM, Brown CH, Brown FE. (1992). Frequency and intensity discrimination in Mongolian gerbils, African monkeys and humans. *Hearing Research*, 59(2):205-12. doi: 10.1016/0378-5955(92)90117-6.

Steenken, F., Bovee, S., & Köppl, C. (2022a). Immunolabeling and Counting Ribbon Synapses in Young Adult and Aged Gerbil Cochleae. *JoVE, 182*, e63874. https://doi.org/doi:10.3791/63874

Steenken, F., Heeringa, A. N., Beutelmann, R., Zhang, L., Bovee, S., Klump, G. M., & Köppl, C. (2021). Age-related decline in cochlear ribbon synapses and its relation to different metrics of auditory-nerve activity. *Neurobiology of Aging*. https://doi.org/https://doi.org/10.1016/j.neurobiolaging.2021.08.019

Steenken F, Oetjen H, Beutelmann R, Carney LH, Koeppl C, Klump GM. (2022b). Neural processing and perception of Schroeder-phase harmonic tone complexes in the gerbil: Relating single-unit neurophysiology to behavior. *European Journal of Neuroscience, 56*(3):4060-4085. doi: 10.1111/ejn.15744.

Steenken, F., Pektaş, A., & Köppl, C. (2024). Age-related changes in olivocochlear efferent innervation in gerbils. Frontiers in Synaptic Neuroscience, 16. https://doi.org/10.3389/fnsyn.2024.1422330

Tarnowski, B. I., Schmiedt, R. A., Hellstrom, L. I., Lee, F. S., & Adams, J. C. (1991). Age-

related changes in cochleas of mongolian gerbils. Hearing Research, 54, 123-134. https://doi.org/10.1016/0378-5955(91)90142-V

Versteegh, C., Meenderink, S. W. F., & Heijden, M. van der. (2011). Response characteristics in the apex of the gerbil cochlea studied through auditory nerve recordings. *Journal of the Association for Research in Otolaryngology*, *12*(3), 301–316. https://doi.org/10.1007/s10162-010-0255-y

Viana, L. M., O'Malley, J. T., Burgess, B. J., Jones, D. D., Oliveira, C. A. C. P., Santos, F., Merchant, S. N., Liberman, L. D., & Liberman, M. C. (2015). Cochlear neuropathy in human presbycusis: Confocal analysis of hidden hearing loss in post-mortem tissue. *Hearing Research*, *327*, 78–88. https://doi.org/10.1016/j.heares.2015.04.014

Wu, P. Z., Liberman, L. D., Bennett, K., de Gruttola, V., O'Malley, J. T., & Liberman, M. C. (2019). Primary neural degeneration in the human cochlea: Evidence for hidden hearing loss in the aging ear. *Neuroscience*, *407*, 8–20. https://doi.org/10.1016/j.neuroscience.2018.07.053

Wu, P. Z., O'Malley, J. T., de Gruttola, V., & Liberman, M. C. (2021). Primary Neural Degeneration in Noise-Exposed Human Cochleas: Correlations with Outer Hair Cell Loss and Word-Discrimination Scores. J Neurosci, 41(20), 4439-4447. https://doi.org/10.1523/JNEUROSCI.3238-20.2021

Supplemental Information

This supplemental information summarizes extended statistical results, which are not essential for the support of the findings in the main manuscript.

Synapse counts
supplemental to "*Synapse loss was similar in old gerbils and gerbils treated with a high dosage of ouabain*"

	2 kHz	4 kHz	8 kHz	16 kHz
young-adult	23 (+/-0,5, N=19)	21 (+/-0,4, N=18)	19 (+/-0,5, N=18)	20 (+/-0,5, N=18)
sham	23 (+/-0,5, N=3)	18 (+/-1, N=3)	16 (+/-0,4, N=3)	21 (+/-0,4, N=3)
ouabain low	22 (+/-0,7, N=8)	19 (+/-0,6, N=8)	16 (+/-0,3, N=8)	19 (+/-0,6, N=8)
ouabain high	19 (+/-1,4, N=10)	17 (+/-1,4, N=10)	12 (+/-1,7, N=10)	13 (+/-2,5, N=11)
old	19 (+/-0,7, N=14)	17 (+/-0,8, N=12)	15 (+/-0,6, N=12)	16 (+/-0,6, N=13)

Mean number of functional synapses per IHC, at the cochlear positions corresponding to 2, 4, 8, and 16 kHz, with standard errors of the mean and number of ears.

The mean number of functional synapses per IHC was significantly different between gerbil groups for cochlear locations corresponding to all four frequencies.
(univariate ANOVAs: 2 kHz: F = 5.496, p = 9.782×10^{-4}; 4 kHz: F = 4.995, p = 1.988×10^{-3};
8 kHz: F = 7.933, p = 5.974×10^{-5}; 16 kHz: F = 6.176, p = 4.305×10^{-4})
Post-hoc tests revealed significant differences (at the 5% level, with Bonferroni correction) between the young-adult and the ouabain-high groups for all equivalent frequencies, between the young-adult and the old group for 2, 4, and 8 kHz, and between the ouabain-low and the ouabain-high group for 16 kHz.

Fig. S1: Synapse number was reduced after ouabain treatment and in old age. Number of functional synapses per IHC, at the cochlear positions corresponding to 2, 4, 8, and 16 kHz, for young-adult gerbils (red), sham-treated gerbils (surgery only, orange), gerbils treated with a low dose (40µM) of ouabain (green), gerbils treated with a high dose (70µM) of ouabain (blue), and old gerbils (gray). Box plots display the median (center line), mean (white circled dot), 25th and 75th percentiles (upper and lower edges of the boxes), and maximum and minimum (whiskers).

Fig. S2: Median spontaneous rates did not differ between animal groups. Individual spontaneous rates for all auditory-nerve fibers reported in this study (dots), group medians (box center lines), quartiles (box boundaries), minima, and maxima (whiskers). The dashed black line indicates the boundary between low- and high-spontaneous-rate fibers (18 spikes/s). Counts of included fibers within each combination of group and spontaneous rate class are indicated at to the corresponding side of the box plots. A Kruskal-Wallis test confirmed that the group medians are not significantly different at the 5% level.

Fig. S3: Median best frequency was lower in the young-adult sample. Individual best frequencies for all auditory-nerve fibers reported in this study (dots), and box plots in the same style as in Fig. S1. The dashed black line indicates the boundary between low and high best frequency class (1.85 kHz), as used in the results statistics. Counts of included fibers within each combination of group and best frequency class are indicated at to the corresponding side of the box plots. A Kruskal-Wallis test confirmed that the group medians are significantly different at the 5% level, with a Bonferroni-corrected post-hoc test showing that only the young-adult group median differs significantly from all other groups.

Below, we list the full ANOVA results, including the principal effects that are reported in the main manuscript. The factors used in the ANOVAs reported here had the following factor levels:

- **group:** young-adult, sham, ouabain-high, old
- **condition:** 200/1600, 400/1600, 400/3200 (f_0/f_c in Hz)
- **BF class:** low: BF ≤ 1850 Hz, high: BF > 1850 Hz
- **SR class:** low: SR ≤ 18 spikes/s, high: SR > 18 spikes/s
- **Δf%:** as a covariate ranging from 0 to 21.25%

All significances are reported at the 5% level, with Bonferroni correction for post-hoc tests and planned comparisons.

Effects on mean driven rate
supplemental to "*Average rate does not carry sufficient information about inharmonic frequency shift*"

Note that driven rate (mean rate - spontaneous rate) was used here. The differences found below are consistent with known differences between low-SR (higher driven rates) and high-SR fibers (lower driven rates; Steenken et al., 2021), and with expected differences across BF classes (low-BF fibers less driven by stimuli with f_c = 3200 Hz and high-BF fibers less driven by stimuli with f_c = 1600 Hz). Differences between groups are also consistent with the sample distributions shown in Figs. S1 and S2.

- **Significant main effects:**
    - **group:** $F(3,943)=10.017$, $p=1.676\times10^{-6}$
        - Mean driven rate was lower in sham group than in any other group.
    - **BF class:** $F(1,943)=6.297$, $p=0.0123$
        - Mean driven rate was lower for high BF class (BF > 1850 Hz).
- **Significant two-way interactions:**
    - **condition × group:** $F(6,943)=2.525$, $p=0.0198$
        - Planned comparisons between groups within each condition, and between conditions within each group:
          In the 400/3200 condition, the mean driven rate of the sham group was significantly lower than in any other group. The mean driven rate in the sham group was significantly lower than in the young-adult group for each condition.
    - **condition × BF class:** $F(2,943)=10.562$, $p=2.909\times10^{-5}$
        - Planned comparisons between conditions within each BF class, and between BF classes within each condition:
          In the x/1600 conditions the mean driven rate in the high-BF class was significantly lower than in the low-BF class. No significant differences between conditions within each BF class.
    - **group × BF class:** $F(3,943)=10.189$, $p=1.316\times10^{-6}$
        - Planned comparisons between groups within each BF class, and between BF classes within each group:
          In the low-BF class, the mean driven rate in the sham group was significantly lower than in all other groups. For the ouabain-high and old groups, the mean driven rate in the high-BF class was lower than in the low-BF class.

- o **group × SR class:** $F_{(2,943)}=8.148$, $p=3.103\times10^{-4}$
  - – Planned comparisons between groups within each SR class, and between SR classes within each group:
    In the young-adult group, the mean driven rate in the high-SR class was lower than in the low-SR class. In the high-SR class, the mean driven rate in the sham group was lower than in any other group. (No samples for the sham group in the low-SR class).
- **Non-significant main effects and interactions:**
  - o *Δf%: $F_{(1,943)}=0.018$, $p=0.8933$ (reported in main manuscript)*
  - o **condition:** $F_{(2,943)}=2.187$, $p=0.1129$
  - o **SR class:** $F_{(1,943)}=1.884$, $p=0.1702$
  - o **group × Δf%:** $F_{(3,943)}=0.394$, $p=0.7575$
  - o **BF class × Δf%:** $F_{(1,943)}=0.052$, $p=0.8195$
  - o **condition × Δf%:** $F_{(2,943)}=0.081$, $p=0.9219$
  - o **SR class × Δf%:** $F_{(1,943)}=0.002$, $p=0.9657$
  - o **condition × SR class:** $F_{(2,943)}=0.698$, $p=0.4977$
  - o **BF class × SR class:** $F_{(1,943)}=0.388$, $p=0.5336$

Effects on TFS representation (TFS log-z-ratio)

supplemental to "*Neural representation of TFS was not degraded in AN fibers of old or synaptopathic gerbils*"

A high TFS log-z-ratio indicates strong phase locking to the stimulus TFS.

- **Significant main effects:**
  - $\Delta f\%$: $F(1,943)=104.403$, $p=2.569\times10^{-23}$ (reported in main manuscript)
  - **condition:** $F(2,943)=7.495$, $p=5.894\times10^{-4}$ (reported in main manuscript)
    – TFS log-z-ratio was significantly lower in the 400/3200 condition, compared to both the 200/1600 and 400/1600 conditions
- **Significant two-way interactions:**
  - **condition × $\Delta f\%$:** $F(2,943)=20.292$, $p=2.353\times10^{-9}$ (reported in main manuscript)
    – Based on the parameter estimates of the ANOVA model:
    The slope of the TFS log-z-ratio vs. $\Delta f\%$ was significantly different from 0, for the 200/1600 and 400/1600 condition. It was significantly higher in the 400/1600 condition, compared to the 200/1600 condition.
  - **BF class × $\Delta f\%$:** $F(1,943)=31.05$, $p=3.282\times10^{-8}$ (reported in main manuscript)
    – Based on the parameter estimates of the ANOVA model:
    The slope of the TFS log-z-ratio vs. $\Delta f\%$ was significantly higher in the low-BF class than in the high-BF class.
  - **condition × group:** $F(6,943)=7.583$, $p=5.808\times10^{-8}$
    – planned comparisons between groups within each condition, and between conditions within each group:
    In the 200/1600 condition, TFS log-z-ratio in the ouabain-high group was significantly higher than in the old group. For all groups except the old group, TFS log-z-ratio in the 400/1600 condition was significantly higher than in the 400/3200 condition. In the sham group, the TFS log-z-ratio in the 400/1600 condition was significantly higher than in the 200/1600 condition. In the ouabain-high group, the TFS log-z-ratio in the 200/1600 condition was significantly higher than in the 400/3200 condition.
  - **condition × BF class:** $F(2,943)=5.774$, $p=3.200\times10^{-3}$
    – Planned comparisons between BF classes within each condition, and between conditions within each BF class:
    In the high-BF class, TFS log-z-ratio in the 400/3200 condition was significantly lower than in any other condition. In the 400/3200 condition, the TFS log-z-ratio in the high-BF class was significantly lower than in the low-BF class.

- o **group × BF class:** $F(3,943)=4.228$, $p=5.600\times10^{-3}$
  - – Planned comparisons between groups within each BF class, and between BF classes within each group:
    In the high-BF class, the TFS log-z-ratio in the ouabain-high group was significantly higher than in the old group.
- o **BF class × SR class:** $F(1,943)=4.918$, $p=0.0268$
  - – In the high-SR class, TFS log-z-ratio in the low-BF class was significantly higher than in the high-BF class.

- **Non-significant main effects and interactions:**
  - o *group: $F(3,943)=1.757$, $p=0.1537$ (reported in main manuscript)*
  - o *group × Δf%: $F(3,943)=1.193$, $p=0.3112$ (reported in main manuscript)*
  - o **BF class:** $F(1,943)=0.103$, $p=0.7479$
  - o **SR class:** $F(1,943)=0.985$, $p=0.3212$
  - o **condition × SR class:** $F(2,943)=2.302$, $p=0.1006$
  - o **group × SR class:** $F(2,943)=0.556$, $p=0.5736$
  - o **SR class × Δf%:** $F(1,943)=1.203$, $p=0.2730$

Effects on ENV/TFS representation balance (ENV/TFS log-z-ratio)

supplemental to "*Representation of f_0 was enhanced in AN fibers of old gerbils*"

The ENV/TFS log-z-ratio reflects the relative emphasis in the AN fibers' responses on either TFS (negative ENV/TFS log-z-ratio) or envelope locking (positive ENV/TFS log-z-ratio).

- **Significant main effects:**
  - ***condition:*** *F(2,943)=43.929, p=5.744×10^{-19} (reported in main manuscript)*
    - The 200/1600 and 400/1600 conditions had negative ENV/TFS log-z-ratios, the 400/3200 condition had a positive ENV/TFS log-z-ratio. All were significantly different from each other, with the 200/1600 condition showing the most negative value.
  - ***SR class:*** *F(1,943)=29.138, p=8.533×10^{-08} (reported in main manuscript)*
    - In the high-SR class, the ENV/TFS log-z-ratio was negative and significantly different from zero. In the low-SR class, it was not significantly different from zero.
  - Δ***f%:*** *F(1,943)=4.278, p=0.0389 (reported in main manuscript)*
- **Significant two-way interactions:**
  - ***condition × group:*** *F(6,943)=11.439, p=2.242×10^{-12} (reported in main manuscript)*
    - Planned comparisons between groups within each condition, and between conditions within each group:
    In addition to the main condition effect: In the 200/1600 condition, the ENV/TFS log-z-ratio was more negative in the sham group than both in the young-adult group, and in the old group. In the 400/3200 condition, the ENV/TFS log-z-ratio was more positive in the old group than in the young-adult group. In the sham group, the ENV/TFS log-z-ratio was more negative in the 200/1600 condition than in the 400/1600 condition.
  - **group × BF class:** F(3,943)=11.354, p=2.551×10^{-7}
    - Planned comparisons between groups within each BF class, and between BF classes within each group:
    In the high-BF class, the ENV/TFS log-z-ratio was significantly more negative in the sham and ouabain-high groups than in the young-adult and old groups.
  - **BF class × SR class:** F(1,943)=5.234, p=0.0224
    - In the order from most negative to most positive ENV/TFS log-z-ratio: low-BF/high-SR, high-BF/high-SR, high-BF/low-SR, low-BF/low-SR. Both high-SR values were significantly below zero, and different from each other; both low-SR values were not significantly different from zero, and not different from each other.

- **Non-significant main effects and interactions:**
  - **group:** $F(3,943)=0.838$, $p=0.4731$
  - **BF class:** $F(1,943)=0.439$, $p=0.5076$
  - **group × Δf%:** $F(3,943)=0.236$, $p=0.8711$
  - **condition × Δf%:** $F(2,943)=1.673$, $p=0.1883$
  - **BF class × Δf%:** $F(1,943)=0.005$, $p=0.9410$
  - **SR class × Δf%:** $F(1,943)=0.58$, $p=0.4464$
  - **condition × BF class:** $F(2,943)=0.414$, $p=0.6611$
  - **condition × SR class:** $F(2,943)=0.522$, $p=0.5936$
  - **group × SR class:** $F(2,943)=2.442$, $p=0.0875$