Auditory perception and neural representation of temporal fine structure are impaired by age but not by cochlear synaptopathy

Friederike Steenken [1,2,3¶], Rainer Beutelmann [1,2,3¶], Henning Oetjen [1,2,3], Christine Köppl [1,2,3], Georg M. Klump [1,2,3*]

1 Cluster of Excellence "Hearing4all", Carl von Ossietzky University Oldenburg, 26111 Oldenburg, Germany

2 Research Centre Neurosensory Science, Carl von Ossietzky University Oldenburg, 26111 Oldenburg, Germany

3 Department of Neuroscience, School of Medicine and Health Science, Carl von Ossietzky University Oldenburg, 26111 Oldenburg, Germany

¶ Shared first author

* Corresponding author
Email: georg.klump@uol.de

Running Title: TFS processing and perception

Author contributions: G.M.K. and C.K. designed research; F.S. and H.O. performed research; R.B. contributed analytic tools; F.S., R.B., H.O., and G.M.K. analyzed data; F.S., R.B., H.O., C.K., and G.M.K. wrote the paper.

Keywords: temporal fine structure, temporal envelope, auditory nerve, perception, gerbil, age-related hearing loss, synapse loss

1 **Abstract**

Age-related hearing loss is a complex phenomenon. The earliest-onset degenerative event is
the gradual loss of neural connections between cochlea and auditory brainstem. To probe for
perceptual deficits that might arise from this loss, cochlear synaptopathy was induced
pharmacologically in young-adult gerbils which were then tested in a challenging listening task
for the perception of temporal fine structure. Treated gerbils behaved no differently than
normal-hearing, young-adult animals. In contrast, old gerbils, which typically express many
cochlear and central-neural pathologies, showed impaired perception. To probe for the
underlying mechanisms, single-unit responses were obtained from the auditory nerve to the
same test stimuli. Responses from old gerbils showed no impairment in temporal locking to
the stimulus fine structure. However, responses were significantly more driven by slower
temporal fluctuations of the stimulus envelope, suggesting that the central auditory system
may be unable to extract the relevant information for discrimination from such altered inputs.

14
15

Introduction

Many elderly people have difficulties understanding speech in noisy environments, despite having normal tone-detection thresholds in quiet. This condition of compromised supra-threshold perception, for example of speech in background noise (e.g., Füllgrabe et al. 2015, Gómez-Álvarez et al. 2023, Lopez-Poveda 2014, Moore 2014, Parthasarathy et al. 2020), despite an absence of detectable hearing deficits in the pure-tone audiogram in quiet, is referred to as hidden hearing loss (HHL, Schaette and McAlpine 2011). It has been suggested that HHL is associated with a reduced number of synapses between auditory-nerve fibers and inner hair cells, which is commonly referred to as cochlear synaptopathy (e.g., Bharadwaj et al. 2014, Kujawa and Liberman 2009, Liberman 2017, Liberman and Kujawa 2017). Synaptopathy is one of the aging processes in the cochlea, and, in addition, can be caused by excessive noise exposure, both accumulating during a human's life time (e.g., Liberman et al. 2016, Wu et al. 2019, but see Johannesen et al. 2019). Animal experiments demonstrate that exposure to loud sounds can produce such synaptopathy, even without permanent loss in hearing sensitivity (e.g., Kujawa and Liberman 2009). Furthermore, the number of functional synapses decreases with increasing age (e.g., gerbil: Bovee et al. 2024, Gleich et al. 2016, Steenken et al. 2021; mouse: Jeng et al. 2020, Parthasaraty and Kujawa 2018, Sergeyenko et al. 2013; rat: Cai et al. 2018; human: Viana et al. 2015).

Synaptopathy is postulated to be causal for compromised processing of temporal stimulus features (e.g., Parthasarathy and Kujawa 2018), especially of the temporal fine structure (TFS) of sounds (reviewed in Moore 2019), resulting in difficulties of speech comprehension in background noise. Specific psychophysical tests that evaluate the performance to distinguish between stimuli that differ in TFS have been developed. These tasks are then expected to be sensitive to synaptopathy (for review see Moore 2014). The most commonly used test is the TFS1 test (Moore and Sek 2009) which probes the discrimination of harmonic and inharmonic tone complexes that strongly differ in TFS but only marginally in the envelope of the waveform. The previously found link between diminished perception of stimuli differing in TFS (Moore et al. 2006) and SNHL in human patients has also been found in TFS1 test results (Mathew et al. 2015). Furthermore, a correlation between age and TFS1 sensitivity has been observed (e.g., Eipert et al. 2019, Füllgrabe et al. 2015, Moore et al. 2012).

Whereas post-mortem studies of human subjects have provided evidence for age-related synapse loss (Viana et al. 2015, Wu et al. 2019), it has been impossible to relate this loss to HHL; that is, deficits in supra-threshold sound perception of living subjects. Animal studies have suggested a close correlation between the number of surviving synapses and supra-threshold amplitude growth of auditory brainstem responses (ABRs) or compound action potentials (e.g., Bourien et al. 2014, Bramhall et al. 2018, Buran et al. 2010, Kujawa and Liberman 2009, Lin et al. 2011, Sergeyenko et al., 2013, Shi et al. 2016). A number of attempts

have been made to infer the synapse loss in humans from non-invasive physiological measures, such as ABR, and psychophysical measures of the processing of sounds. However, even studies involving large samples of human subjects have not produced strong evidence that synaptopathy in humans has functional consequences for supra-threshold temporal perception (e.g., Bramhall et al. 2019, Carcagno and Plack 2022, Guest et al. 2018). Attempts to relate life-time noise exposure of human subjects to their supra-threshold perceptual deficits have also not been conclusive (e.g., Prendergast et al. 2017, 2019, Füllgrabe et al. 2020). Controlled animal studies are one strategy for testing the influence of synapse loss on discrimination of complex sounds that differ in TFS. In those, synaptopathy can be experimentally manipulated and the degree of cochlear dysfunction can be related to neural responses and performance in behavioral tests, using the same stimuli. With this approach we can test the hypothesis whether synaptopathy negatively affects the discrimination of stimuli that differ in their TFS.

Animal studies relating behavioral measures of performance in response to a variety of stimuli to synaptopathy have so far also provided mixed results (reviewed in Henry 2022). An investigation by Henry and Abrams (2021), which experimentally induced synaptopathy in budgerigars by kainic acid infusions at the round window, failed to demonstrate a relationship between the amplitude of the compound action potential (taken as a measure of cochlear synaptopathy) and tone-detection thresholds in noise. Studies in Macaque monkeys that produced a noise-induced hearing loss (NIHL) by acoustic trauma demonstrate differences between normal-hearing and acoustically traumatized monkeys for a) tone detection in noise (Burton et al. 2020), b) masking release for tone detection in modulated maskers compared to unmodulated maskers, and c) spatial release from masking for tone detection in noise (Mackey et al. 2021). However, these studies used high trauma levels and long exposure durations. Thus, effects were not limited to synapse loss at the inner hair cell but included both outer- and inner-hair-cell damage. Studies in chinchilla auditory-nerve (AN) fibers, involving normal-hearing subjects and subjects with NIHL, did reveal differences in the neural envelope representation but failed to demonstrate differences in the neural representation of TFS that match the psychophysical effects observed in human subjects (e.g., Henry and Heinz 2012, Kale and Heinz 2010, Kale et al. 2014).

To clarify the relation between synaptopathy, AN fiber responses, and behavior, we embarked on a study in Mongolian gerbils (*Meriones unguiculatus*) that combined behavioral measurements of sensitivity in the TFS1 test and the representation of harmonic and inharmonic complex tones with the same fundamental frequency (f_0) by AN fibers. In addition, synaptopathy of many of the experimental subjects was characterized by directly counting the numbers of functional synapses on inner hair cells, and these measures were related to sensitivity in the TFS1 test. Because compromised performance in the TFS1 test has been

observed in elderly humans, we compared the perception of young-adult and old gerbils. In order to experimentally induce synaptopathy in young-adult gerbils and, thus dissociate the effects of synaptopathy from effects of age, we treated young-adult gerbils with an infusion of ouabain at the round window. This treatment has been shown to reduce the number of synapses on gerbil inner hair cells (Bourien et al. 2014). We observed a lower perceptual sensitivity of old compared to young-adult gerbils in the TFS1 test. However, contrary to our hypothesis, ouabain treatment did not affect the performance in the TFS1 task, and the responses of AN fibers that survived ouabain treatment did not differ from gerbils that were not treated.

Results

Synapse loss was similar in old gerbils and gerbils treated with a high dosage of ouabain

Synapses were counted at the cochlear location corresponding to 2 kHz in 54 gerbils, divided into the five gerbil groups, as listed in Table 1, row 4. Young-adult, untreated gerbils had, on average, 23 (SE ±0.5) synapses per inner hair cell (Fig. 1). Synapse numbers were significantly different between gerbil groups (univariate ANOVA: F = 5.496, p = 0.001). Compared to untreated young adults, the average number of synapses per inner hair cell in gerbils treated with 70 µM ouabain and in old gerbils was significantly reduced, to 19 (SE ±1.4) and 19 (SE ±0.7), respectively (Bonferroni corrected pairwise comparisons: young-adult vs. 70 µM [p = 0.011]; young-adult vs. old [p = 0.002]). No significant difference between ears treated with a high dose of ouabain and old ears was found, indicating that both animal groups had similar degrees of synaptopathy in the region of interest. In surgery-only ears and ears treated with 40 µM ouabain, average synapse numbers (surgery only: 23 (SE ±0.5), 40 µM ouabain: 22 (SE ±0.7)) were not significantly different compared to untreated young-adult ears, suggesting no effect of the surgical procedure itself and no effect of the lower ouabain dose. These results were also consistent across more basal cochlear locations (up to 8 kHz equivalent frequency, see Supplemental Materials). Based on this, we pooled the single-unit data from surgery-only and 40 µM ouabain treated gerbils into a "sham" group. The gerbils treated with 70 µM ouabain are henceforth referred to simply as "ouabain".

Figure 1: Synapse number was reduced after ouabain treatment and in old age. *Number*
of functional synapses per IHC, at the cochlear positions corresponding to 2 kHz, for young-
adult gerbils (red, N=19), sham-treated gerbils (surgery only, orange, N=3), gerbils treated with
a low dose (40µM) of ouabain (green, N=8), gerbils treated with a high dose (70µM) of ouabain
(blue, N=10), and old gerbils (gray, N=14). Box plots display the median (center line), mean
(white circled dot), 25th and 75th percentiles (upper and lower edges of the boxes), and
maximum and minimum (whiskers).
129

Average rate does not carry sufficient information about inharmonic frequency shift

130
131

Spike responses to TFS1 stimuli of 94 AN fibers from 26 animals were recorded (for details
see Table 1 in Materials and Methods). BFs ranged from 470 Hz to 11,528 Hz, with a median
of 3,995 Hz. The SRs of these fibers ranged from 0 sp/s to 144 sp/s with a median of 60 sp/s.
Average stimulated rates (4.8 sp/s - 223 sp/s, median 135 sp/s) and driven rates (i.e., average
rate minus SR, 4.8 sp/s - 215 sp/s, median 67.6 sp/s) were calculated from the spike counts
across the stimulus duration of 375 ms, excluding on- and offset ramps.
Most importantly, the inharmonic frequency shift $\Delta f\%$ - as a covariate in the ANOVA - had no
significant effect on mean driven rate [$F(1;943) = 0.018$, $p = 0.8933$] (Fig. 2, overlapping
colored and black distribution curves in each panel). The overall distribution of driven rate also
did not change substantially with $\Delta f\%$. Thus, the inharmonic frequency shift had no effect on
the mean driven rate, and AN spike rate was not a potential cue for the gerbil in the behavioral
task.

Beyond this robust, basic result, there were differences in discharge rates between the animal groups and stimulus conditions that might reflect different sampling biases of single units in the different experimental groups (see Supplemental Material for details).

Figure 2: Mean driven spike rates did not differ between frequency shifts. *Histograms of average driven spike rates during presentation of the TFS1 stimuli (black and colored) and average spontaneous rate, without acoustic stimulation (gray). Subject groups are arranged in columns and stimulus conditions in rows. Colors code the frequency shift Δf in percent of f₀, with the responses to harmonic stimuli shown in black and responses to the inharmonic stimuli grading from yellow (least inharmonic) to red (most inharmonic). The legends apply to panels above them. In each panel, the maximal number of fibers (with 40 stimulus repetitions per fiber) that the histograms is based on is indicated; note that not all fibers were tested with each frequency shift Δf. Clipped spontaneous rate bars are marked with an arrow and the maximal relative frequency beyond the axis limit.*

162 Neural representation of TFS was not degraded in AN fibers of old

163 or synaptopathic gerbils.

The same set of AN fibers was probed for the neural responses to the fine structure and
envelope of TFS1 stimuli (for details see Table 1) using the TFS log-z-ratio (see section "Neural
recordings - Data Analysis"). A high TFS log-z-ratio indicates strong phase locking to the
stimulus TFS. The main findings are listed below; for more details see Supplemental Material.
Importantly, the inharmonic frequency shift, Δf%, as a covariate in the ANOVA, affected the
TFS log-z-ratio significantly and strongly [main effect: $F(1;943) = 104.403$, $p = 2.569\times10^{-23}$],
suggesting that AN fibers followed the fine structure of the stimulus and thus represented the
frequency shift in their spiking pattern (Fig. 3A-J, L). Δf% significantly interacted with stimulus
condition and BF-class [interaction: $F(2;943) = 20.292$, $p = 2.353\times10^{-9}$; $F(1;943) = 31.050$,
$p = 3.282\times10^{-8}$, respectively]. In response to 400/1600 Hz, the shifts affected the temporal
representation most (i.e., higher Δf% resulted in higher TFS log-z-ratios). Additionally, TFS log-
z-ratios of low-BF fibers were influenced more by the inharmonic shifts than those of high-BF
fibers (again, higher Δf% result in higher TFS log-z-ratios).

TFS representation by the TFS log-z-ratio metric was not significantly different between AN
fibers from different gerbil groups ($p = 0.154$). Also, no interaction between group and Δf%
occurred. Together, this indicates that aged fibers (Fig. 3D, H, L) and fibers that survived the
treatment with ouabain (Fig. 3C, G) had responses that did not differ from young-adult (Fig.
3A, E, I) and sham-treated fibers (Fig. 3B, F, J). As expected, TFS log-z-ratios of AN fibers
responding to the stimulus condition with higher f_c were lower, compared to those responding
to stimuli with lower f_c (200/1600 Hz: 0.987, 400/1600 Hz: 1.111, 400/3200 Hz: 0.394; [main
effect: $F(2;943) = 7.495$, $p = 5.894\times10^{-4}$]). This reflects the known decrease in phase-locking
with increasing stimulus frequency (Versteegh et al., 2011).

186 Representation of f_0 was enhanced in AN fibers of old gerbils

As a metric for the relative representation of TFS vs. stimulus envelope, the ENV/TFS log-z-
ratio was used (see section "Neural recordings - Data Analysis"). This metric reflects the
relative emphasis in the AN fibers' responses of either TFS (negative ENV/TFS log-z-ratio) or
envelope locking (positive ENV/TFS log-z-ratio).

Here, over all animal groups, Δf% had a marginally significant effect on the ENV/TFS log-z-
ratio [main effect: $F(1;943) = 4.278$, $p = 0.039$] and stimulus condition had a highly significant
effect on the ENV/TFS log-z-ratio [main effect: $F(2;943) = 43.929$, $p = 5.744\times10^{-19}$]. Posthoc
comparisons revealed that, for 400/3200 Hz, envelope representation (mean ENV/TFS log-z-
ratio = 0.263) was stronger than fine structure representation, indicating stronger phase-
locking of AN fibers to low-frequency stimulus components. Furthermore, stimulus condition

Figure 3: Frequency shift representation in the neural responses differed between groups and stimulus conditions. _Vector strength frequency spectra of responses to harmonic and inharmonic stimuli, averaged across repetitions and fibers. Subject groups are arranged in columns and stimulus conditions in rows. Colors code the frequency shift Δf in percent of f_0, with the same color code as in Fig. 4. Dashed lines indicate the limits of the stimulus bandpass filters. The inset panels show enlarged examples of the spectra ranging around the envelope frequency (f_0, 200 or 400 Hz) and the fine structure (center) frequency (TFS peak frequency; 1000, 800, or 2000 Hz). The thin gray lines mark the y-axis scaling of the inset panels. In each panel, the maximal number of fibers (with 40 stimulus repetitions per fiber) that the spectra are based on is indicated; note that not all fibers were tested with each frequency shift Δf._

showed a significant interaction with gerbil groups [interaction: $F(6;943) = 11.439$, $p = 2.242 \times 10^{-12}$]. In old gerbils, the strength of phase-locking was nearly equal to the envelope (Fig. 3D, left inset; Fig. 4D) and to the TFS (Fig. 3D, right inset; Fig. 4D'), in response to 200/1600 Hz. In response to 400/3200 Hz, fibers from old gerbils also showed better envelope locking (Fig. 3L, Fig. 4L). Since TFS representation was unchanged in old animals (see above), this result reflects enhanced envelope coding by AN fibers obtained in old gerbils, compared to all other animal groups (Fig. 3A-D, I-L, left insets). In contrast, AN fibers of young-adult gerbils, irrespective of treatment, tended to respond strongly to the fine structure of the

stimulus, especially for conditions with lower f_c of 1600 Hz (i.e., showed VS peaks at the respective, shifted frequencies; Fig. 3A-H, right insets; Fig. 4A'-C', E'-G').

Over all animal groups, low-SR fibers had, on average, positive ENV/TFS log-z-ratios, that is, phase-locked more strongly to the stimulus envelope, whereas high-SR fibers showed, on average, negative ENV/TFS log-z-ratios, that is, phase-locked more strongly to the TFS [main effect: $F(1;943) = 29.138$, $p = 8.533×10^{-8}$].

***Figure 4: Frequency shift representation differed between envelope and TFS frequency ranges.** Frequency shift of the z-value peak versus stimulus frequency shift Δf. Subject groups are arranged in columns and stimulus conditions in double rows. Odd rows show the data at f_0 and even rows at f_{max}. Each circle represents four dimensions: the stimulus frequency shift Δf (position on the x-axis), the frequency shift of the z-value peak (position on the y-axis), the average vector strength (z-score) across all respective fibers (color saturation, see legend at the bottom), and the percentage of fibers with a frequency shift of the z-value peak at the corresponding stimulus frequency shift Δf (circle radius). The largest circles correspond to 100% of fibers (e.g., in panel C), medium sized circles to values around 50% (e.g., panel J). Colors code the frequency shift Δf, with the harmonic stimulus in black and the inharmonic stimuli fading from yellow (least inharmonic) to red (most inharmonic). In each panel, the maximal number of fibers (with 40 stimulus repetitions per fiber) that the z-value peaks are based on is indicated; note that not all fibers were tested with each frequency shift Δf.*

Age, but not synapse loss, affected behavioral discrimination of TFS1 stimuli

To investigate the effect of synapse loss on performance in the TFS1 task, the behavior of young-adult gerbils, without and with cochlear synaptopathy, and old gerbils was compared for the different stimulus conditions defined by the combinations of f_0 and f_c. The sensitivity of the four groups of gerbils in the different stimulus conditions is illustrated in Fig. 5, in relation to the frequency shift $\Delta f\%$. The sensitivity of young-adult gerbils of the different treatment groups was quite similar, whereas the old gerbils had considerably lower sensitivity. A GLMM ANOVA with the sensitivity *d'* as dependent variable, condition and treatment group as factors, and frequency shift $\Delta f\%$ as a covariate (only trials with $\Delta f\% > 5$ included, for which data from all treatment groups were obtained), revealed significant main effects of treatment group [$F(3,387) = 8.18$, $p = 2.725 \times 10^{-5}$], stimulus condition [$F(2,387) = 22.71$, $p = 4.736 \times 10^{-10}$, and $\Delta f\%$ [$F(1,387) = 20.457$, $p = 8.121 \times 10^{-6}$]. No significant interactions were observed. There was a general increase in sensitivity with increasing $\Delta f\%$. Old gerbils showed lower sensitivities than most young-adult gerbils, irrespective of their treatment (all $p < 0.002$, Bonferroni corrected). There was no significant difference in sensitivity between the treatment groups of young gerbils. To separately investigate the effects of f_0 and harmonic number, *N*, on sensitivity, we conducted planned comparisons in two additional GLMM ANOVAs, each containing only the data of two conditions, with the same dependent variable and factors as the initial ANOVA. The planned comparison with the data of stimulus conditions 200/1600 Hz and 400/1600 Hz was used to investigate the effect of f_0 (which inherently includes a change in harmonic number, N) on sensitivity. This ANOVA revealed an effect of f_0, with a higher sensitivity for increasing f_0 [$F(1,257) = 45.001$, $p = 1.248 \times 10^{-10}$]. Furthermore, an effect of $\Delta f\%$ [$F(1, 257) = 13.539$, $p = 2.847 \times 10^{-4}$] was observed. Similar to the previous analysis, the main effect of gerbil group [$F(3, 257) = 4.878$, $p = 2.570 \times 10^{-3}$] reflected the low sensitivity of the old gerbils compared to all other groups. The second planned comparison with the data of stimulus conditions 400/3200 Hz and 400/1600 Hz, with equal f_0, investigated the dependence of sensitivity on f_c. An effect of f_c [$F(1,256) = 12.656$, $p = 4.459 \times 10^{-4}$] was shown, with decreasing sensitivity for increasing f_c. As in the previous comparison, this ANOVA also revealed increased sensitivity with increasing $\Delta f\%$ [$F(1,256) = 13.484$, $p = 2.929 \times 10^{-4}$], whereas the main effect of gerbil group [$F(3,256) = 4.692$, $p = 3.299 \times 10^{-3}$] again reflected the lower sensitivity of the old group compared to all other groups. No interactions were observed in either planned comparison.

The frequency-shift threshold reflects $\Delta f\%$ for a criterion of *d'* = 1 and was calculated for each gerbil and condition. If no threshold could be derived because all *d'* values were below 1, a threshold of 100% f_0 was assumed for further statistical analysis. On average, old gerbils were only able to discriminate inharmonically shifted TFS1 stimuli for the condition 400/1600 Hz (see

d' values > 1 in Fig. 5D, H, L). All young-adult gerbil treatment groups achieved average
sensitivities above threshold for all stimulus conditions.

Figure 5: The gerbils' discrimination performance improved with larger frequency
shifts*. Sensitivity index d' for behavioral discrimination, as a function of stimulus frequency*
shift Δf in percent of f$_0$. Reference lines at d' = 1 indicate the assumed threshold value for
meaningful discrimination performance. Subject groups are arranged in columns and stimulus
conditions in rows. Colors code the subject group. Circled black dots show the mean across
subjects, boxes show the median and interquartile ranges, whiskers show the extrema, except
for outliers, which are displayed as colored dots. Data points are considered outliers, if they lie
beyond 1.5 times the distance between median and upper or lower quartile, respectively. For
each panel, the maximal number of subjects in the respective group is indicated. Unfilled boxes
mark conditions completed by less than half of those subjects in the respective group.

Frequency-shift thresholds for the three different stimulus conditions are shown in Fig. 6. A
GLMM ANOVA was conducted with Δf% thresholds as the dependent variable and condition
and treatment group as factors. There were main effects for stimulus condition [$F_{(2,60)}$ = 8.99,
p < 0.0005] and treatment group [$F_{(3,60)}$ = 6.84, p<0.0005], with no interaction between both
factors. Average thresholds were not significantly different between the 200/1600 Hz and
400/3200 Hz conditions, while thresholds were lower in the 400/1600 Hz than in the
200/1600 Hz condition (p < 0.0005). The pairwise comparison revealed that old gerbils had
significantly higher thresholds than gerbils of all other animal groups (all p < 0.005).

Figure 6: Old gerbils were typically unable to perceive frequency shifts whereas ouabain
treatment did not impair discrimination. *Behavioral discrimination thresholds based on the*
data shown in Fig. 5. Thresholds were defined as the lowest (linearly interpolated) stimulus
frequency shift Δf in percent of f_0, at which the sensitivity index d' crossed d'=1. The threshold
was set to 100% if this criterion was never met. Colors code the subject group. Circled black
dots show the mean across subjects, boxes show the median and interquartile ranges,
whiskers show the extrema, except for outliers, which are displayed as colored dots. Data
points are considered outliers, if they lie beyond 1.5 times the distance between median and
upper or lower quartile, respectively. The y-axes at the right show the same frequency shifts
in Hz.

No differences were observed between the young-adult gerbil groups with different treatments.
To evaluate the effect of f_0 on the threshold, a planned comparison with the data of stimulus
conditions 200/1600 Hz and 400/1600 Hz (i.e., harmonic numbers 8 and 4, respectively) was
conducted in an additional ANOVA, with frequency-shift thresholds as dependent variable, and
condition and gerbil group as factors. A main effect of condition was revealed, indicating that
f_0 (respectively harmonic number N at 1600 Hz f_c) did affect the threshold [$F(1,40) = 15.219$,
$p = 3.577 \times 10^{-4}$]. There was no significant effect of gerbil group; all groups showed similarly low
thresholds for the 400/1600 Hz condition. A second additional ANOVA with the data of stimulus
conditions 400/3200 Hz and 400/1600 Hz (i.e., harmonic numbers 8 and 4, respectively)
investigated the dependency of threshold on f_c (respectively harmonic number N). An effect of
gerbil group [$F(3,40) = 10.460$, $p = 3.272 \times 10^{-5}$] and condition [$F(1,40) = 16.213$, $p = 2.455 \times 10^{-4}$]
was observed. The pairwise comparison revealed lower thresholds with decreasing f_c
(respectively harmonic number N) and a significant difference between untreated and old
gerbils (all $p \leq 0.001$), whereas the young-adult gerbil treatment groups did not differ.
In summary, the behavioral data showed that the old gerbils differed from all other treatment
groups of young-adult gerbils. This result suggests that age considerably affected both
perceptual sensitivity and thresholds while synapse numbers had little effect.

Discussion

The ability to process temporal fine structure (TFS) of sounds is a key factor for speech
perception and its age-related decline has been held responsible for compromised speech
comprehension in the elderly, especially in challenging noisy acoustic backgrounds (Füllgrabe
et al. 2015, Moore 2014, 2019). It has been suggested that age-related or noise-induced
synaptopathy results in the observed perceptual deficits regarding the TFS of sounds (e.g.,
Moore 2014, 2019). However, so far a direct test of this hypothesis is lacking. Here, we present
data that investigate the performance of the Mongolian gerbil in the TFS1 test (Moore and Sek
2009), a psychoacoustic paradigm that has been applied in human subjects to evaluate the
ability to discriminate between stimuli based on the TFS. We determined the TFS1 test
performance in young-adult gerbils and compare this to the performance of old gerbils (≥ 36
months of age, corresponding to humans ≥ 60 years of age, Castaño-González et al. 2024),
demonstrating an age-related decline in TFS sensitivity similar to that found in human subjects
(Füllgrabe et al. 2015, Moore et al. 2012). However, young-adult gerbils with experimentally
induced synaptopathy (by applying ouabain), to a degree corresponding to the synaptopathy
in old gerbils (Steenken et al. 2021, this study), showed no decline in TFS sensitivity. This
result suggests that reduced synapse numbers alone cannot explain the perceptual deficits for
TFS in old subjects. Auditory-nerve (AN) fiber responses of young-adult, young-adult ouabain-
treated, and old gerbils, elicited by the stimuli presented in the TFS1 test, indicate that the

representation of TFS in AN fibers is not affected by ouabain treatment or age. In old gerbils that show compromised TFS1 test performance behaviorally, however, the representation of the signal envelope by AN fibers was enhanced compared to the other groups. We propose that this enhancement of the signal envelope cue, which does not differentiate the stimuli in the TFS1 test, may cause the perceptual deficits in old gerbils by overriding the usability of the TFS cues. Thus, cue representation may cause the perceptual deficits, but not reduced synapse numbers, as originally proposed.

Gerbil as a model for human temporal-fine-structure perception

In the TFS1 test, subjects must discriminate an inharmonic (I) complex from a harmonic (H) complex, in which the frequency difference between the components determines the period of the envelope. The I complex is created by shifting all components of the H complex by a fixed frequency. This shift results in pairs of I and H complexes with similar envelopes but different TFS. Thus, TFS is the only acoustic cue which allows to discriminate the stimuli. Excitation-pattern differences due to the shift of the frequency components in the I complex compared to the H complex can be eliminated in the TFS1 test by appropriate spectral filtering (e.g., Marmel et al. 2015). In addition, the gerbil has larger auditory filter bandwidths than human subjects (Kittel et al. 2002). Since we used a similar filter bandwidth for the gerbil as that applied in studies involving humans (e.g., Eipert et al. 2019, Marmel et al. 2015, Moore and Sek 2009), cochlear excitation pattern differences are unlikely to have provided usable cues for discrimination by the gerbils in the present study. Furthermore, the noise masker in which the I and H complexes were presented ensured that the stimuli did not produce usable distortion products for the gerbils. Thus, as in the study with human subjects, the only usable cue for the discrimination was the TFS.

Figure 7 provides an overview of gerbil TFS1 thresholds in comparison to human thresholds. Human thresholds were selected from studies involving stimuli with spectral components and fundamental frequencies (f_0) similar to those presented to the gerbils in the present study, allowing a comparison of gerbil and human performance. There is a broad overlap between gerbil and human TFS1 thresholds. Both, in humans and gerbils, old subjects showed higher (i.e., worse) TFS1 thresholds than young subjects (for human studies see Eipert et al. 2019, Füllgrabe et al. 2015, Moore et al. 2012). This similarity indicates that the gerbil is a good model for investigating the mechanisms underlying TFS perception that may also apply to humans. Behavioral TFS1 thresholds in young-adult gerbils with ouabain-induced synaptopathy, similar to that typical for old gerbils, were similar to TFS1 thresholds in untreated young-adult gerbils. This result suggests that increased thresholds in old subjects were not due to the reduced number of synapses. Thus, in the gerbil, the hypothesis that age-related deterioration of TFS perception is a direct consequence of synaptopathy can be rejected.

Figure 7: Frequency shift detection thresholds of gerbils and humans for TFS1 test stimuli in relation to the center frequency f_c of the harmonic complex. *Filled symbols represent gerbil data from the present study. Human data are from published material with the limitation that the f_c was in the range from 500 Hz to 4500 Hz and the fundamental f_0 of the harmonic complex was in the range of 100 Hz to 400 Hz to make the range of parameter values similar to those of the present study (for references see legend, normal hearing NH, hearing loss HL).*

Auditory-nerve fiber responses do not explain differences in behavioral sensitivity

Since the use of excitation patterns by the gerbil is unlikely due to its larger auditory-filter bandwidths, we explored the temporal responses of AN fibers for alternative explanations of compromised TFS perception. TFS representation requires the ability of AN fibers to phase-lock (reviewed in Joris et al. 2004, Rose et al. 1967). Here we showed that, throughout all tested gerbil groups, the H and I complexes were neurally distinguishable based on phase-locking to the stimulus fine structure (Fig. 3, 4). In contrast, no distinction was possible based on the average firing rate of AN fibers (Fig. 2). Neural phase-locking reached highest z-values for stimuli with f_c of 1600 Hz and were lower for 3200 Hz (Fig. 4), which reflects the known decrease in phase-locking of AN fibers for stimuli above 1.5 kHz (Versteegh et al. 2011). Several studies in animal models of cochlear dysfunction showed that phase locking is not compromised in single AN fibers, including quiet-aged gerbils (Heeringa et al. 2020), noise-exposed cats (Miller et al. 1997), chinchillas (Henry and Heinz 2012, Kale et al. 2013), and kanamycin-treated guinea pigs (Harrison and Evans 1979). Consistent with these findings, the

surviving fibers in old, synaptopathic gerbils did not show compromised temporal fine structure
representation of I and H complexes compared to young-adult gerbils (Fig. 3, 4).
Old gerbils showed significant synapse loss compared to untreated young-adult gerbils at the
cochlear location corresponding to 2 kHz (Fig. 1), consistent with what was shown previously
(Steenken et al. 2021). Ouabain, a cardiac glycoside that inhibits Na^+-K^+-ATPase pumps of
AN fibers (O'Brien et al. 1994), applied to the round window, results in a dose-dependent loss
of AN fibers in gerbils (Bourien et al. 2014, Lang et al. 2005, Schmiedt et al. 2002). In our
hands, a low dose (40 µM) of ouabain had no effect at the cochlear location equivalent to
2 kHz. In contrast, a higher dose of 70 µM resulted in a loss that was similar to that observed
in old gerbils (Fig. 1). The surviving fibers of ouabain-treated gerbils showed no deterioration
of TFS representation compared to untreated or sham-treated young-adult gerbils (Fig.
3A,C,E,G); however, the behavioral outcomes of this study clearly showed that only old gerbils
had deteriorated discrimination abilities between H and I complexes. We thus showed
conclusively that neither synapse loss, to an extent typical for old gerbils, nor impaired TFS
representation, at the level of single AN fibers, could explain the deteriorated perception in old
gerbils.

Enhancement of confounding/distracting AN cues and changed central processing may explain the behavioral deficit in old subjects

Surviving AN fibers in old gerbils did not show compromised fine-structure representation (Fig.
3, 4). However, old gerbils exhibited deteriorated performance in the behavioral test in
response to TFS1 test stimuli (Fig. 5, 6). Despite the overall robust phase-locking ability, other
pathologies are known to develop with advancing age.
First, the age-related atrophy of the stria vascularis within the lateral wall of cochleae (Gratton
and Schulte 1995) results in a decreased endocochlear potential (Schulte and Schmiedt 1992),
which in turn increases the threshold (i.e., decreases the sensitivity) of AN fibers (Schmiedt et
al. 1996). Envelope representation, measured as synchronization to amplitude-modulated
stimuli, is non-monotonic with sound level, with a pronounced peak near the threshold of the
fiber (Heeringa et al. 2023, Joris and Yin 1992). Thus, the typical sensitivity loss in old subjects
will cause a difference in neural representations between young-adult and old subjects, even
if the test stimulus levels are kept constant. Alternatively, stimulus levels can be raised for old
subjects, to an equivalent sensation level that compensates for their sensitivity loss. Consistent
with that prediction, AN fibers of old gerbils show enhanced representation of the fundamental
frequency of speech vowels, compared to fibers from young adults, when responding to fixed-
level stimuli (Heeringa et al. 2023), but do not show enhanced envelope representation in
response to frozen noise presented at similar sensation level (Heeringa et al. 2020). Old gerbils

of the same age as in the present study invariably show hearing loss, although to variable extents (Hellstrom and Schmiedt 1990, Steenken et al. 2021). Thus, since TFS1 stimuli in the current study were presented at a fixed absolute level of 68 dB SPL, fibers of old animals were stimulated closer to their individual thresholds than fibers of young adults, which is expected to increase their phase locking to f_0, as observed. This increase in phase-locking to f_0 may have confounded their perception. In old gerbils, the frequency selectivity of AN fibers is not significantly affected (Heeringa et al. 2020, Hellstrom and Schmied 1996), unlike that in noise-induced hearing loss which can cause additional changes to temporal coding. Chinchillas with noise-induced hearing loss showed compromised temporal coding in response to TFS1 stimuli (Kale et al. 2013), which is explained by a mismatch between the AN fiber's characteristic frequency and the frequency of the stimulus component. In AN fibers from noise-damaged animals, this mismatch arises as the consequence of changes to their frequency tuning (Henry et al. 2019). In chinchillas treated with furosemide, a drug that reversibly reduces the endocochlear potential, the AN fibers' frequency tuning curves remained unaffected, whereas their sensitivity is decreased (Henry et al., 2019). The same can be seen in aged gerbils (Heeringa et al. 2020, Hellstrom and Schmied 1996), and, thus, a distorted tonotopy caused by changed frequency tuning is unlikely.

Second, age-related changes in GABAergic processing in the central auditory system have been found (Caspary et al. 1995), including the gerbil (Gleich et al. 2014, Kessler et al. 2020) and shown to affect temporal processing (Gleich et al. 2003). The gerbils used by Kessler and colleagues (2020) were derived from the same population as the gerbils used for the present study and had the same age range for old animals. Therefore, it can be assumed that the gerbils in the current study showed a change in GABAergic processing in the brain that could affect processing of such supra-threshold stimuli as being used in the TFS1 test.

Conclusion

We demonstrated that the gerbils' ability to discriminate TFS1 stimuli is neither affected by synapse loss nor by alterations to TFS representation by their AN fibers. However, the elevated neural response thresholds in old gerbils enhanced the neural envelope representation. This enhancement may have overridden the still intact TFS representation.

Acknowledgments

This work was supported by the DFG priority program "PP 1608", the DFG Cluster of Excellence EXC 1077/1 "Hearing4all" (Project ID 390895286). We thank Laurel Carney for helpful comments on a previous version of this manuscript. Susanne Groß for assistance with the behavioral experiments. Amarins N. Heeringa for providing analysis scripts for AN data.

Nadine Thiele for generous assistance with the gerbil care and for implementing the blinding

process. Furthermore, many thanks go to Sonja Standfest for the great assistance with gerbil

care. Sonny Bovee is acknowledged for processing cochleae, Julia Forst and Carina Lützow

for assistance with the spike detection and immunostainings, and Julia Winter for assistance

with the immunostainings. We acknowledge the Fluorescence Microscopy Service Unit, Carl

von Ossietzky University of Oldenburg, for the use of the imaging facilities.

Materials and Methods

Experimental model and subject details

Sixty adult, agouti-colored Mongolian gerbils (*Meriones unguiculatus*) of both sexes (32 M, 28 F) were studied (Table 1). The gerbils originated from Charles River Laboratories and were bred in an in-house facility of the University of Oldenburg. Groups of gerbils were tested, differing in age and ouabain treatment: untreated young-adult, surgery-only young-adult, ouabain-treated young-adult, and old gerbils. Young-adult gerbils were between 3 and 19 months of age and old gerbils were 36 months or older. All animals were housed individually or in pairs in Type IV cages, which were provided with nesting material. During behavioral testing, access to water was not limited, but animals were food-deprived, such that the weight of young-adult and old gerbils was 65 - 75 g and 70 - 85 g, respectively, to motivate the animals to perform for a food reward. All protocols and procedures were approved by the Niedersächsisches Landesamt für Verbraucherschutz und Lebensmittelsicherheit (LAVES), Germany, permit AZ 33.19-42502-04-15/1990. All procedures were performed in compliance with the NIH Guide on Methods and Welfare Consideration in Behavioral Research with Animals (National Institute of Mental Health, 2002).

Table 1: Numbers of gerbils of different treatment and age groups within the behavioral, electrophysiological, and histological part of this study. Note that numbers in category 2-4 do not sum up to total numbers, since many gerbils were used in all three study parts. Furthermore, both untreated gerbil groups were supplemented with data from eight young-adult gerbils and twelve old gerbils for which synapse counts were available, from Steenken et al. (2021).

		Young adult	Surgery only	Ouabain 40 µM	Ouabain 70 µM	Old
1	**Total** numbers of animals in this study	21	4	8	10	17
2	Animals in **behavioral** study	4	4	5	8	3
3	Animals with **neural** responses to **TFS1 stimuli** (number of fibers)	5 (37)	2 (10)	3 (8)	6 (14)	6 (23)
4	Animals from which **cochleae were harvested**	19	3	8	10	14

Young-adult gerbils were on average 10.3 ± 5.7 months of age (N = 21) and old gerbils were

38.4 ± 2.5 months old (N = 17) when electrophysiological recordings were conducted and

cochleae were harvested. Treated young-adult gerbils were either 15.1 ± 3.6 (treated with a

low concentration of ouabain, N = 8) or 12.7 ± 4.2 (high concentration of ouabain, N = 10). All

surgery-only gerbils were 18 months old (N = 4). Gerbil groups had significantly different ages

(univariate ANOVA: F=99.343; $p < 0.0005$). Bonferroni corrected post-hoc tests revealed that

old gerbils were significantly different from all other young-adult gerbil groups with respect to

age, while all young-adult gerbil groups, irrespective of treatment, did not differ.

Methods details

Stimulus paradigm

The stimuli were similar to those used for the TFS1 test in humans (for a detailed description

see Moore and Sek 2009). In short, the reference stimulus was a harmonic tone (H) complex.

The test stimulus that was discriminated from the reference by the gerbils in the behavioral

tests or by analyses of AN-fiber responses was an inharmonic tone (I) complex created by

shifting all frequency components upwards. This procedure creates H and I complexes with

the same envelope periodicity. In the following, the frequency shift, Δf, will be expressed by

the percentage of the fundamental frequency, f_0 (Weber fraction).

The f_0 was either 200 or 400 Hz. H and I complexes were generated with components from f_0

up to 9 kHz. H and I complexes were passed through the same 3^{rd}- or 5^{th}-order Butterworth

infinite-impulse-response band-pass filter (for AN recording and behavioral experiments,

respectively), with a center frequency, f_c, of 1600 or 3200 Hz, to reduce possible cues related

to differences in the excitation pattern on the basilar membrane (Moore and Moore 2003). The

3-dB down points of the filter were separated by 7 f_0, which resulted in a flat spectral region of

5 f_0. The amplitude at the filter skirts decreased by at least 30 dB per octave (except for the

3^{rd}-order filter with f_c of 1600 Hz and f_0 of 400 Hz, for which the slope was 18 dB/oct).

The phase of each harmonic was randomly chosen for each test trial, but was matched for the

H and I complexes within a trial to ensure similar envelopes for both stimuli (see Fig. 1 A-D).

A pink noise with frequencies between 100 and 11,050 Hz was added to the signal to mask

distortion products possibly emerging in the inner ear, and signal components outside the filter

passband. The masker level was 20 dB SPL at 1 kHz, to mask distortion products of H and I

stimuli in the gerbil (Faulstich and Kössl, 1999). The level of each component in the tone

complex was 60 dB SPL resulting in an overall level of 68 dB SPL. The duration of each H or

I complex stimulus was 0.4 s. In the behavioral experiment, stimulus onset and offset were

ramped with a 25-ms Hann window. Stimuli presented to AN fibers were not ramped but still

filtered. By analyzing the steady-state response only, any on- and offset effects were excluded.

Figure 8: Examples illustrating stimulus waveforms and typical auditory-nerve responses. *Stimulus waveforms (A-D), peri-stimulus-time histograms (PSTHs, E-H), and average vector strength as a function of frequency (I-L). Columns 1 (A,E,I) and 2 (B,F,J) show data from a HSR fiber, columns 3 (C,G,K) and 4 (D,H,L) show data from a LSR fiber. Both fibers are from a young-adult, untreated animal and are matched in best frequency. The stimulus condition was f_0=400 Hz and f_c=1600 Hz. A and C show examples of the harmonic stimulus (Δf=0%, black) and its envelope (gray), B and D show examples of a strongly inharmonic stimulus (Δf=8%, red), its envelope (gray), and the harmonic stimulus (black) as a reference.The corresponding neural responses are displayed in the same colors in the second and third row, respectively. Arrow markers are added (A-D) to highlight the differences in fine structure between harmonic and inharmonic stimuli. The PSTHs (E-H) show the instantaneous spike rate, averaged across 40 repetitions of identical stimuli. Vector strength (I to L) is shown as a function of frequency, averaged across 40 stimulus presentations. The inset panels show enlarged examples of the spectra in the envelope frequency range (400 Hz) and the fine structure frequency range (800 Hz).*

In both the behavioral experiments and in AN-fiber recordings, three combinations of f_0 and f_c

were used: f_c = 1600 Hz, and f_0 = 200 Hz; f_c = 1600 Hz, and f_0 = 400 Hz, and f_c = 3200 Hz and

f_0 = 400 Hz. The combination of f_0 = 200 Hz and f_c = 3200 Hz was not used because a pilot

behavioral test showed that two young untreated gerbils were not able to discriminate I and H

even for the largest value of Δf.

The harmonic rank, *N*, of a condition describes the rank of the center frequency in the harmonic

tone complex. The highest value for Δf was 42.5% of the fundamental f_0. Lower values for Δf

were computed by dividing 42.5% by 2n, with n being 1 or increasing integers. The smallest

Δf presented in the behavioral tests was 0.33% f_0. In old gerbils tested behaviorally, additional

test stimuli with Δf = 31.87% f_0 and 15.94% f_0 were presented to ensure that old gerbils were

presented with a sufficient number of trials allowing a salient percept for discrimination.

In recordings from gerbil AN fibers, only subsets of the stimuli used in behavior were presented

to save time and minimize the number of incomplete datasets due to deterioration or loss of

single-unit isolation. The frequency shifts were further grouped into either "high" (10.6, 15.9,

21.25, 31.8, and 42.5 Hz) or "low" (0.66, 1.3, and 5.31 Hz) and the stimuli were presented in

three different blocks per stimulus condition. The first block comprised the H and two I complex

stimuli, one with a high and one with a low shift. The second block comprised I complex stimuli,

two high shifts and another low shift. The third block always comprised the remaining shifts

(1.3, 10.6, and 42.5 Hz). The order of stimuli within the blocks was random between fibers. A

silent period of 0.6 s occurred between stimuli. This organization of stimuli ensured that the H

complex was always presented and could be compared to one low and one high shift. The

second and third block of stimuli were recorded with declining success.

The periodic envelopes of H and I complexes created in the way described above are similar

and offer no cues for discrimination, whereas the TFS of H and I complexes is not identical

(Fig. 8A-D). The TFS of I complexes changes across sequential periods while the TFS of the

H complex does not; only this difference can be used for discrimination. The small random

change in the TFS and envelope of both H and I due to the addition of pink noise offers no

discrimination cues. The bandpass filtering of the tone complexes served to minimize cochlear

excitation pattern cues.

Experimentally introduced synaptopathy using ouabain

For experiments including ouabain treatment, gerbils were anesthetized with ketamine

(Ketamin 10%, bela-pharm GmbH, Vechta, Germany; 135 mg/kg body weight) and xylazine

(Xylazin 2%, Ceva Tiergesundheit GmbH, Düsseldorf, Germany; 6 mg/kg body weight) mixed

with 0.9% saline injected intra-peritoneal. Maintenance doses of one third of the initial dose

were given hourly or as needed. Depth of anesthesia was monitored by withdrawal reflex to

toe pinches and via an ECG with needle electrodes in the front- and contralateral hind leg,

continuously displayed on an oscilloscope (SDS 1102CNL, SIGLENT Technologies, Hamburg,

Germany). Body temperature was continuously controlled with a homeothermic blanket
(Harvard Apparatus, Holliston, MA, USA). Experiments were carried out in a sound-attenuating
booth. The gerbil's head was fixed within a bite-bar (David Kopf Instruments, Tujunga, CA,
USA). Pure, moisturized oxygen (1.5 l/min) was delivered from a tube pointing towards the
nose.

Twenty-two gerbils (11M, 11F, aged 3 – 8 months at the beginning of treatment) received
either ouabain or artificial perilymph solution. Twenty-four hours before the oubain or artificial
perilymph injection, gerbils were treated with antibiotics (oral; Baytril (0.5%, Bayer Animal
Health GmbH, Monheim am Rhein, Germany; 0.3 ml per gerbil/day) and a non-steroidal
antiphlogistic/ analgetic agent (oral; Meloxidyl (0.5 mg/ml, CevaTiergesundheit GmbH; 0.3 ml
per gerbil/day; or Novalgin; oral; 500 mg/ml, Sanofi-Aventis Deutschland GmbH, Frankfurt,
Germany, 0.21 ml per gerbil/day). The antibiotic treatment was continued for nine more days,
the antiphlogistic/ analgetic agent for at least three days. Additionally, gerbils received an
antiemetic (oral; Emeprid, Ceva Tiergesundheit GmbH; 0.21 ml per gerbil/day) and, if needed,
also sterofundin (4 ml s.c.; B. Braun SE, Melsungen, Germany) or amynin (4 ml s.c.;
Boehringer Ingelheim Pharma GmbH Co KG, Ingelheim am Rhein, Germany) as infusion
solutions if gerbils were dehydrated, all post-operatively.

Ouabain treatment was carried out according to Bourien et al. (2014). After gerbils were
anesthetized, they received a sub-cutaneous infusion with sterofundin (4 ml). The location
where the headholder met the skin of the face, was treated with 2% xylocaine ointment
(Lidocainhydrochloride, Aspen Pharma Trading Limited, Dublin, Ireland). The skin covering the
bulla was disinfected with 70% ethanol. A small incision was placed behind the ear to access
the bulla. A hole was drilled into the bone with an angled blade and subsequently widened with
forceps to reach the round window. A syringe needle tip was placed onto the round window to
apply one of the following solutions onto the round window membrane: Eight gerbils were
treated with 40 µM ouabain (Ouabain octahydrate, Sigma-Aldrich, Saint Louis, MO, USA), ten
with 70 µM ouabain, dissolved in artificial perilymph solution (137 mM NaCl, 5 mM KCl, 2 mM
CaCl2, 1 mM MgCl2, 1 mM NaHCO3, and 11 mM glucose; pH 7.4). Four gerbils received only
artificial perilymph solution (surgery-only treatment). In 15 gerbils, the solution was
continuously delivered by a pump (150 µl for 30 Min). In seven gerbils, the round window niche
was bathed with the solution by manually operating the syringe and after ten minutes, the
solution was aspirated and reapplied. This procedure was repeated three times. Surgery was
concluded by suturing the skin incision over the open, middle ear. Gerbils were binaurally
treated (n = 19), except those that were destined for electrophysiology only, which were
monaurally treated (n = 3).

Note that the experimenter conducting the electrophysiology and histology was blinded to the
type of treatment: the solution that was transferred to the round window was aliquoted by
another person. Only after the synapse counts were completed, the treatment was revealed.

Counting inner-hair-cell synapses

Immunohistochemistry

Processing of cochleae was carried out as described in detail in Steenken et al. (2022). After
concluding the single-unit recordings, gerbils were euthanized with an overdose of
pentobarbital (i.p.; Narcoren, Merial GmbH, Hallbergmoos, Germany, 480 mg/kg body weigh).
Twenty gerbils were cardiovascularly perfused with a mixture of phosphate-buffered saline
(PBS; 137 mM NaCl, 10 mM phosphate and 2.7 mM KCl, pH 7.4) followed by 4%
paraformaldehyde in PBS. To prevent blood coagulation, in 15 of these gerbils, heparin
(ratiopharm, 25,000 IE/5 ml; 0.2 ml/100 ml) was added to PBS. In 34 gerbils, the bulla was
rapidly exposed after euthanasia and small openings were carefully created at the apex and
base of the bony cochlear walls. Cochleae were then fixed in 4% paraformaldehyde in PBS for
two days on a shaker at 8 °C. After fixation, all cochleae were decalcified using 0.5 M
ethylenediaminetetraacetic acid for two days. Subsequently, cochleae were treated with 1%
Triton X-100-PBS for 1 hour to permeabilize the tissue and were then rinsed with 0.2% Triton
X-100-PBS. To block unspecific binding sites, cochleae were then incubated in 3% bovine
serum albumin (BSA, + 0.2% Triton X-100-PBS) blocking solution for 1 hour at room
temperature.

Antibodies to label hair cells (anti-MyosinVIIa, IgG polyclonal rabbit; Proteus Biosciences,
Ramona, CA, USA; diluted 1:400), presynaptic ribbons (anti-CtBP2 (C-terminal binding
protein) antibody, IgG1 monoclonal mouse; BD Biosciences, Eysins, Switzerland; diluted
1:400), and postsynapstic receptor patches (anti-GluR2a antibody, IgG2a monoclonal mouse;
Millipore, Burlington, MA, USA; diluted 1:200) in blocking solution were used. Cochleae were
incubated in this mixture at 37 °C overnight. Next, cochleae were again rinsed in 0.2% Triton
X-100-PBS.

Secondary antibodies were chosen to match the hosts of the primary antibodies: goat anti-
mouse (IgG1)-AF488 (Molecular Probes Inc., Eugene, OR, USA; diluted 1:1000), goat anti-
mouse (IgG2a)-AF568 (Invitrogen, Carlsbad, CA, USA; diluted 1:500), and donkey anti-rabbit
(IgG)-AF647 (Molecular Probes Inc.; diluted 1:1000). Secondary antibodies were diluted in
blocking solution and cochleae were incubated in this solution at 37 °C overnight, and
subsequently rinsed in PBS.

After the immunostaining, a subset of the cochleae from untreated animals older than 12
months received treatment with an auto-fluorescence quencher (n = 13; TrueBlack Lipofuscin
Autofluorescence Quencher, 20x in DMF, Biotum, Hayward, CA, USA). These cochleae were
cut in half and incubated in a 5% true black-70% ethanol-mixture for 1 minute and then rinsed

in PBS. Finally, all cochleae were micro-dissected under a stereomicroscope and the resulting

5 - 11 organ-of-Corti pieces were mounted on slides using Vectashield Mounting Medium

(Vector Laboratories, Burlingame, CA, USA, H-1000).

Image acquisition

Overview images of cochlear pieces were acquired with a light microsope (Nikon 90i with NIS

Elements software, Version 4.30, Nikon, Minato, Tokio, Japan) using a 4x objective. After

measuring the length of the cochlea as a line along the row of inner hair cells with the NIS

software, the cochlear location at 3.8 mm from the apex, corresponding to 2 kHz (Müller,

1996), was chosen, because 2 kHz was part of all stimulus conditions (data for additional

locations corresponding to 4, 8, and 16 kHz are provided in the Supplemental Materials).

Images were taken at this position with a confocal microscope (Leica Microsystem CMS

GmbH, Wetzlar, Germany, Leica TCS SP8 system) using an oil-immersion objective (40x,

numerical aperture 1.3). Lasers (488-nm and 522-nm optically pumped semiconductor laser),

or a 638-nm diode, respectively, excited the different fluorescence tags, and released photons

were counted with a hybrid detector. Image stacks had voxel dimensions of 0.05 - 0.0998 µm

(XY) and 0.3 µm (Z). All confocal stacks were deconvolved (Huygens Essentials, Version

15.10, SVI, Hilversum, Netherlands) with default settings (maximum iteration: 80; signal to

noise ratio: 10; quality threshold: 0.01), using a theoretical point-spread function.

Synapses were counted using ImageJ (FIJI, Schindelin et al. 2012). For each cochlear

position, the number of functional synapses was evaluated for 5 inner hair cells by manually

counting the co-localized ribbons and glutamate patches. Contrast and brightness were

manually adjusted for each stack.

ABR recordings

Hearing sensitivity of all gerbils was evaluated prior to and during AN recordings using ABRs.

The recording electrode was placed near the ipsilateral mastoid, the reference electrode was

placed in the neck, and a ground electrode was placed on one hindleg. The stimulus

presentation was controlled via a custom-written MATLAB software (Mathworks, Natick, MA,

USA) and delivered through a RME Hammerfall DSP Multiface II sound card (RME Audio,

Haimhausen, Germany). The stimuli were amplified (HB7, TDT Inc.) and presented to the

gerbil's ear via earphones (IE 800, Sennheiser, Wedemark, Germany) attached to an ear-bar.

ABRs were amplified (x10^3) and band-pass filtered (0.3 - 3 kHz) using an ISO-80 pre-

amplifier (World Precision Instruments, Sarasota, FL, USA), and sampled using the digital

signal processor (Hammerfall DSP Multiface II, RME Audio; 48 kHz sampling rate) controlled

by custom MATLAB software (Mathworks, Natick, MA, USA).

Prior to AN recordings, responses to chirps were recorded at a range of levels, from below

threshold up to 20 dB above threshold in steps of 5 dB, repeated 300 times. Thresholds from

averaged ABR waveforms were detected visually during the experiment. During single-unit recording sessions, ABR thresholds were re-checked periodically, and the experiment was terminated if threshold deteriorated by more than 30 dB.

Single-unit recordings

Stimulus presentation

Stimuli for single-unit recordings were generated with custom-written MATLAB-scripts controlling a digital signal processor (RX6, TDT Inc., Alachua, FL, USA) and an attenuator (PA5, TDT Inc.).The signal was then routed through a headphone buffer (HB7, TDT Inc.) and presented via earphones (IE 800, Sennheiser, Wedemark, Germany) attached to an ear-bar. Stimuli were calibrated at the start of each experiment using a probe-tube microphone (ER7-C, Etymotic Research Inc., Elk Grove Village, IL, USA) attached to the ear-bar and a microphone amplifier (MA3, TDT Inc., 40 dB amplification).

Single-unit recordings

Single-unit data was obtained in 22 gerbils (see Table 1). After the gerbil was anesthetized, a single dose of non-steroidal antiphlogistic agent (meloxicam, 0.2 mg/kg, Boehringer Ingelheim Pharma GmbH Co KG) was administered subcutaneously. The head of the gerbil was oriented with a bite bar and firmly fixed to the setup via a screw glued with dental cement to the exposed skull. The pinna of one ear was removed to place an ear-bar directly at the bony edge of the ear canal. The ear-bar was sealed with petroleum jelly to the bone to form a closed sound system. To prevent pressure buildup in the middle ear, a small hole was drilled into the dorsal bulla with an angled blade.

The AN was accessed via a dorsal approach. The skin covering the occipital bone and the bone itself were removed and the lateral part of the cerebellum was aspirated. The brainstem was left intact and was gently pushed medially using tiny saline soaked cotton balls to access the proximal part of the eighth cranial nerve. A glass electrode (GB120F-10, Science Products GmbH, Hofheim am Taunus, Germany; pulled using a P-2000, Sutter Instruments Co., Novato, CA, USA), filled with 3 M KCl-solution and with a resistance of ~10 - 30 MΩ, was placed above the AN under visual control. The electrode was advanced through the AN via a remote-controlled piezo motor (Burleigh 6000 ULN inchworm motor controller and 6005 ULN handset, Burleigh, Inc., Fishers, NY, USA). Recordings were amplified (WPI 767, World Precision Instruments Inc., Sarasota, FL, USA), filtered for line-frequency noise (50 Hz, Hum Bug, Quest Scientific Instruments Inc., North Vancouver, BC, Canada), and digitized (sampling rate: 48828 Hz; RX6, TDT Inc., Alachua, FL, USA). Additionally, the analog signal was guided to an audio monitor (MS2, TDT Inc.) and displayed on an oscilloscope (SDS 1102CNL, SIGLENT Technologies, Hamburg, Germany).

Search stimuli (broadband noise, 1 - 9 kHz at 50 dB SPL) were played while the electrode was advanced into the AN. When a unit was isolated, its frequency response range was first assessed audio-visually using 50-ms duration tones. Next, the best frequency (BF) of each fiber was assessed by presenting pure tone bursts (50-ms duration, including 5-ms raised-cosine rise/fall times, 0 - 30 dB above threshold, 3 repetitions) at a range of frequencies within approximately ±1 octave around the audio-visually determined best response frequency. Subsequently, tone bursts at BF were presented over a range of levels (10 - 79 dB SPL, 50-ms duration including 5-ms raised-cosine rise/fall times, 10 repetitions) to derive a rate-level function. At randomly inserted trials, no stimulus was presented to determine the spontaneous rate (SR). Subsequently, TFS1 stimuli were presented at a fixed level of 60 dB SPL.

Spike detection from the recorded signal was conducted off-line with a custom-written MATLAB script (Steenken et al., 2022b). In short, the bandpass-filtered (300 – 3000 Hz) recordings were manually screened for threshold-crossing events. The threshold could be adjusted on a trial-by-trial basis, to compensate for variations in spike amplitude. The time of each spike's peak amplitude was saved as the spike time. Spike trains were excluded if: 1.) more than 10% of the spikes for a given stimulus condition were judged to be below the set threshold criterion , or 2.) any inter-spike intervals < 0.6 ms occurred, indicating inadequate unit isolation (Heil et al. 2007), or 3.) a pre-potential was present in the spike waveform, suggesting a recording from the ventral cochlear nucleus (Keine and Rubsamen 2015), or 4.) the rate-level function was non-monotonic, also suggesting a potential cochlear nucleus recording (e.g., Davis et al. 1996, Joris et al. 1994, Sinex et al. 2001).

Neural recordings - Data Analysis

A number of measures were derived from the AN spike times: a) Peri-stimulus time histograms (PSTHs), b) rate histograms, and c) vector strengths (VS) corresponding to certain periodicities in the stimuli. Only the 375-ms long steady-state part of the responses, excluding a 12.5 ms onset and offset fringe, respectively, was analyzed.

First, peri-stimulus time histograms (PSTHs) were calculated for each animal/fiber/condition/Δf combination, pooling over repeated stimulus presentations within each combination. Spike times were collected into histograms using bin widths of 1/10 of the f_0 period (0.5 ms for 200 Hz and 0.25 ms for 400 Hz). The histogram counts were converted into instantaneous spike rates by dividing by the bin width and by the number of stimulus repetitions.

Second, average rate and SR histograms were calculated for each group/condition/Δf combination, pooling over repeated stimulus presentations, animals, and fibers within each combination. Average rates were computed over the 375-ms steady-state portion of the response. SR was calculated from trials without stimulation. Driven rate was calculated as the difference between stimulated absolute rate and SR. Average driven rates and SRs were collected into histograms using bin widths of 2.5 spikes per second and 5 spikes per second,

respectively. The average-driven-rate histogram counts were divided by the number of spike trains to convert the counts to relative frequency and smoothed with a moving-average window of 3 bin lengths.

Third, VS frequency spectra were calculated for each group/condition/Δf combination, pooling over repeated stimulus presentations, animals, and fibers within each combination. For this, all-order inter-spike intervals (ISIs) were calculated from each spike train, and collected into histograms using a bin width of 50 µs, corresponding to a sampling rate of 20 kHz. The bin positions ranged from -375 ms to +375 ms. A zero-centered gaussian window with a standard deviation of 200 ms was multiplied with the histogram to suppress fringe effects and spurious side lobes in the spectra. The histogram counts were divided by the total number of ISIs. A subsequent fast Fourier transform of the histograms resulted in the ISI power spectrum, and the square root of the spectral magnitudes was identical to the VS at each bin center frequency. The spectral resolution was 1.33 Hz.

VS single-peak data was calculated for each animal/fiber/condition/Δf combination, pooling over repeated stimulus presentations within each combination. In contrast to the spectra, the peak VS values were calculated individually for each relevant frequency f:

$$V(f) = \left| \frac{1}{N} \sum_{i=1}^{N} e^{-i2\pi f t_i} \right| ,$$

where t_i are the spike times, and N is the number of spikes in each spike train. For significance analysis of the VS, the Rayleigh statistic z (henceforth referred to as z-value) was calculated: $z = N_{avg} \cdot V_{avg}^2(f)$, where N_{avg} is the average number of spikes per spike train and V_{avg} is the average VS across stimulus presentations. The Rayleigh z statistic has useful properties for statistical analyses because VS values resulting from higher numbers of spikes have larger weights, and logarithms of ratios of z-values are approximately normally distributed, as opposed to VS ratios or differences.

As a metric for TFS representation, the log ratio between two z-values was derived: First, from responses to the H complex, the z-value, z_0, at the frequency component that the fibers on average responded to maximally, $f_{maxpeak}$, was obtained. Second, this value was divided by the z-value, $z_{\Delta f}$, at the shifted frequency of the I complex, also in the maximal average response range (filled circles in Fig. 9). This ratio was used as the dependent variable in the first ANOVA model and reflects high temporal resolution; large positive values correspond to stronger phase-locked responses to the TFS of the stimulus. This metric was termed the TFS log-z-ratio.

Another log ratio between z-values was used as a metric for the balance between envelope and TFS representation (crosses in Fig. 9): The (harmonic, unshifted) z_0 value at the fundamental frequency, f_0, was divided by the (inharmonic, shifted) $z_{\Delta f}$ value in the maximal average response range, $f_{maxpeak}$. Large positive log ratios indicate a stronger response to the

stimulus envelope, whereas large negative log ratios indicate a stronger response to the stimulus TFS. This metric was termed ENV/TFS log-z-ratio.

Figure 9: Schematic representation of statistical contrasts. Schematic of the frequency spectrum of a harmonic (black solid lines) and inharmonic (red dashed lines) TFS1 stimulus in panel A and schematic representation of the phase locking spectrum in a neuronal response in panel B. Black arrows and symbols point to the harmonic, unshifted frequencies, termed "z_0" and red arrows and symbols point to the inharmonic, shifted frequencies, termed "z_{df}". The annotated brackets below the frequency axis mark representative regions of the fundamental frequency, f_0, and of the maximal average response, fmaxpeak. Filled connected circles show the log-z-ratio used for assessing the TFS representation, connected crosses show the log-z-ratio used for assessing the balance between f_0/ENV and TFS representation.

The representation of Δf shift in the responses was also tested by calculating z-values at the

different harmonic frequencies, combined with all possible Δf shifts.. The actual position of the

maximum z-value in terms of Δf and the corresponding z-value is summarized in Fig. 4 for two

TFS peak frequencies (to which the Δf shifts were added): at f_0 of the corresponding condition

and at 1 kHz for the 200/1600 Hz condition, at 800 Hz for the 400/1600 Hz condition, and at

2 kHz for the 400/3200 Hz condition, respectively. The latter TFS peak frequencies were

identified as the harmonics actually present in the acoustic stimulus, which had the maximum

z-value in the summarized spectra (Fig. 3).

For the single-unit data, several main factors were defined. The low-SR fiber class comprised

fibers with SR ≤ 18 spikes/s, and the high-SR class comprised fibers with SR > 18 spikes/s. In

addition, fibers in the low-BF class had BF ≤ 1850 Hz, and fibers in the high-BF class had BF

> 1850 Hz.

Behavioral procedure

The behavioral experiments were conducted in a single-walled sound-attenuating chamber

(Industrial Acoustics, Type 401 A, Industrial Acoustic Company GmbH, Mönchengladbach,

Germany) lined with a 15-cm thick layer of sound- absorbing foam (Pinta Acoustics Pyramide

100/50 on Pinta Acustics PLANO Type 50/0, Seyboth

& Co., Regensburg, Germany). The walls and all devices within the chamber produced no

relevant sound reflections. The time for the reverberation in the chamber to decay by 60 dB

was 12 ms, indicating nearly anechoic conditions. These conditions allowed faithful

presentation of the TFS1 stimuli with the desired acoustic cues in the free sound field.

In the center of the chamber, a 30-cm-long wire-mesh platform was located at 90-cm height,

with an elevated pedestal at its center where the gerbils waited during the measurement. This

construction minimized sound reflections. At one end of the platform, a food bowl was located,

attached via a tube to a custom-made feeder that did not obstruct the sound path. Gerbils were

rewarded for correct discrimination with a 10-mg custom-made food pellet (Altromin

International, type 1324 rodent pellets enriched with sunflower oil and spelt flour; Altromin

Spezialfutter GmbH & Co. KG, Lage, Germany). A loudspeaker (Canton Plus XS, frequency

range: 150 Hz - 21 kHz, Canton, Weilrod, Germany) was mounted 30 cm in front of the

pedestal to the side of the food bowl, at 0° elevation and azimuth relative to the gerbil's normal

head position when sitting on the pedestal. Out of the gerbil's reach, a system of two custom-

made light barriers was installed to detect the gerbil's position and facing direction on the

pedestal. An infrared camera (Conrad 150001 C-MOS camera module, Conrad Electronics,

Hirschau, Germany) was positioned above the platform to observe the gerbil's movements on

the platform under invisible infrared light. The stimulus presentation, registration of light

barriers switching, and feeder were controlled by custom-written software on a Linux-based

PC with an RME sound-card (Hammerfall DSP Multiface II, sampling frequency 48 kHz). The

signal output from the sound card was manually attenuated (Texio type RA-902A, TEXIO

Technology Corporation, Kanagawa, Japan), amplified (Rotel type RMB 1506, Rotel Tokyo,

Japan), and presented by the loudspeaker. Before testing started on each day, the system

was calibrated (±1 dB) with a sound level meter (Brüel and Kjaer type 2238 Mediator, Hottinger

Brüel & Kjær, Virum, Denmark) positioned on the elevated pedestal at about the position of

the gerbil's head.

The behavioral experiment used an operant Go/NoGo paradigm. Gerbils were trained to wait

on the pedestal facing in the direction of the loudspeaker. During the whole session, the H

complex was played every 1.3 s as a reference stimulus. As soon as the gerbil jumped onto

the pedestal facing the loudspeaker, a trial was started with a random waiting time between 2

and 7 s. When the waiting time had elapsed, a target stimulus, the I complex, was played

instead of the reference H complex. The target stimuli could either be equal to the reference

H complex (sham trial) for estimating the false-alarm rate or an I complex (test trial) for testing

discrimination ability. Leaving the pedestal before the waiting time elapsed resulted in the

restart of the trial with a new waiting time. Leaving the pedestal within 1.3 s after the onset of

the target stimulus of a test trial, as indicated by the light barriers, was registered as a "hit" and

resulted in a food reward. Staying on the pedestal in a test trial was counted as a "miss" with

no food reward. The animal had to leave the pedestal and jump back onto it in the correct

sequence to start a new trial after a "miss" occurred. Leaving the pedestal within 1.3 s after the

start of the target stimulus in a sham trial was registered as a "false alarm", while staying on

the pedestal during a sham trial was registered as a "correct rejection". No food reward was

provided in sham trials. To keep the gerbils motivated during the session, a salient test trial not

included in the analysis was inserted after each correct rejection.

Each session contained 10 blocks with 9 trials each (six test trials, three sham trials). All stimuli

within a given block were from the same condition (200/1600 Hz, 400/1600 Hz, or

400/3200 Hz). The first block of a session was a warmup block with Δf = 42.5% that was not

included in the analysis. To interleave simple and more difficult conditions, blocks of the

different conditions were pseudo-randomly distributed throughout the session. The least

salient condition (200/1600 Hz) did not occur within two consecutive blocks. Additionally, the

sham and test trials within each block were pseudo-randomly distributed with the constraint

that no sham trials followed one another within and across blocks. Data collection started when

the gerbil achieved a false alarm rate below 20% in three consecutive sessions with hit rates

for the three highest Δf values that would allow determining a discrimination threshold.

Sessions were included in the analysis if the gerbil completed all 90 trials and the false alarm

rate did not exceed 20%. Data collection continued until the sample size for each condition

and Δf value was at least 20 trials. The sensitivity for each Δf value was defined by applying

the z-transform, $d' = z(hit\ rate) - z(false\text{-}alarm\ rate)$. Individual discrimination thresholds for

each animal and condition were then calculated from sensitivity, d', as a function of inharmonic

frequency shift, $\Delta f\%$. A threshold at $d' = 1$ was obtained by linearly interpolating between the

first $\Delta f\%$ shift where $d' > 1$ and the previous $\Delta f\%$ shift with $d' <= 1$. If all d' values were below

1, a linear fit was estimated through all d' values as a function of $\Delta f\%$ shifts and threshold was

extrapolated at $d' = 1$. If all d' values were larger than 1, the threshold was set to half of the

minimal $\Delta f\%$ shift measured in the specific experiment. If none of the above was possible and

either all $d' < 1$, the fitted slope was not positive, or the extrapolated threshold larger than 100%

shift (equivalent to $\Delta f = f_0$ at threshold), the final threshold was limited to 100% (5 out of 72

thresholds).

Statistical analysis

918 All analyses of variance were calculated with IBM SPSS Statistics for Windows, Version 29.0

919 (IBM Corp, Armonk, NY). The family-wise error rate of all multiple post-hoc comparisons was

920 controlled by Bonferroni correction of the significance levels.

Supportimg information

923 At the end of the manuscript plus figures file

References

Bharadwaj, H. M., Verhulst, S., Shaheen, L., Liberman, M. C., & Shinn-Cunningham, B. G.
(2014). Cochlear neuropathy and the coding of supra-threshold sound. *Frontiers in*
Systems Neuroscience, *8*, 26. https://doi.org/10.3389/fnsys.2014.00026
Bourien, J., Tang, Y., Batrel, C., Huet, A., Lenoir, M., Ladrech, S., Desmadryl, G., Nouvian,
R., Puel, J.-L., & Wang, J. (2014). Contribution of auditory nerve fibers to compound
action potential of the auditory nerve. *Journal of Neurophysiology*, *112*(5), 1025–1039.
https://doi.org/10.1152/jn.00738.2013
Bovee, S., Klump, G. M., Pyott, S. J., Sielaff, C., & Köppl, C. (2024). Cochlear ribbon synapses
in aged gerbils. *International Journal of Molecular Sciences*, *25*(5).
https://doi.org/10.3390/ijms25052738
Bramhall, N., Beach, E. F., Epp, B., Le Prell, C. G., Lopez-Poveda, E. A., Plack, C. J., Schaette,
R., Verhulst, S., & Canlon, B. (2019). The search for noise-induced cochlear
synaptopathy in humans: Mission impossible? *Hearing Research*, *377*, 88–103.
https://doi.org/https://doi.org/10.1016/j.heares.2019.02.016
Bramhall, N. F., McMillan, G. P., Kujawa, S. G., & Konrad-Martin, D. (2018). Use of non-
invasive measures to predict cochlear synapse counts. *Hearing Research*, *370*, 113–119.
https://doi.org/https://doi.org/10.1016/j.heares.2018.10.006
Buran, B. N., Strenzke, N., Neef, A., Gundelfinger, E. D., Moser, T., & Liberman, M. C.
(2010). Onset coding is degraded in auditory nerve fibers from mutant mice lacking
synaptic ribbons. *The Journal of Neuroscience*, *30*(22), 7587.
https://doi.org/10.1523/JNEUROSCI.0389-10.2010
Burton, J. A., Mackey, C. A., MacDonald, K. S., Hackett, T. A., & Ramachandran, R. (2020).
Changes in audiometric threshold and frequency selectivity correlate with cochlear
histopathology in macaque monkeys with permanent noise-induced hearing loss.

Hearing Research, *398*, 108082.
https://doi.org/https://doi.org/10.1016/j.heares.2020.108082

Cai, R., Montgomery, S. C., Graves, K. A., Caspary, D. M., & Cox, B. C. (2018). The FBN rat model of aging: investigation of ABR waveforms and ribbon synapse changes. *Neurobiology of Aging*, *62*, 53–63.
https://doi.org/https://doi.org/10.1016/j.neurobiolaging.2017.09.034

Carcagno, S., & Plack, C. J. (2022). Relations between speech-reception, psychophysical temporal processing, and subcortical electrophysiological measures of auditory function in humans. *Hearing Research*, *417*, 108456.
https://doi.org/https://doi.org/10.1016/j.heares.2022.108456

Caspary, D. M., Milbrandt, J. C., & Helfert, R. H. (1995). Central auditory aging: GABA changes in the inferior colliculus. *Experimental Gerontology*, *30*(3–4), 349–360.
https://doi.org/10.1016/0531-5565(94)00052-5

Castaño-González, K., Köppl, C., & Pyott, S.J. (2024). The crucial role of diverse animal models to investigate cochlear aging and hearing loss. *Hearing Research 445*: 108989.
https://doi.org/10.1016/j.heares.2024.108989

Davis, K. A., Ding, J., Benson, T. E., & Voigt, H. F. (1996). Response properties of units in the dorsal cochlear nucleus of unanesthetized decerebrate gerbil. *Journal of Neurophysiology*, *75*(4), 1411–1431. https://doi.org/10.1152/jn.1996.75.4.1411

Eipert, L., Selle, A., & Klump, G. M. (2019). Uncertainty in location, level and fundamental frequency results in informational masking in a vowel discrimination task for young and elderly subjects. *Hearing Research*, *377*, 142–152.
https://doi.org/https://doi.org/10.1016/j.heares.2019.03.015

Faulstich, M., & Kössl, M. (1999). Neuronal response to cochlear distortion products in the anteroventral cochlear nucleus of the gerbil. *The Journal of the Acoustical Society of America*, *105*(1), 491–502. https://doi.org/10.1121/1.424586

Füllgrabe, C., Moody, M., & Moore, B. C. J. (2020). No evidence for a link between noise exposure and auditory temporal processing for young adults with normal audiograms. *The Journal of the Acoustical Society of America*, *147*(6), EL465–EL470.
https://doi.org/10.1121/10.0001346

Füllgrabe, C., Moore, B. C. J., & Stone, M. A. (2015). Age-group differences in speech identification despite matched audiometrically normal hearing: contributions from auditory temporal processing and cognition. *Frontiers in Aging Neuroscience*, *6*, 347.
https://doi.org/10.3389/fnagi.2014.00347

Gleich, O., Hamann, I., Klump, G. M., Kittel, M., & Strutz, J. (2003). Boosting GABA improves impaired auditory temporal resolution in the gerbil. In *Neuroreport* (Vol. 14, Issue 14, pp. 1877–1880). https://doi.org/10.1097/00001756-200310060-00024

Gleich, O., Netz, J., & Strutz, J. (2014). Comparing the inferior colliculus of young and old gerbils (Meriones unguiculatus) with an emphasis on GABA. *Experimental Gerontology*, 57, 155-162. https://doi.org/10.1016/j.exger.2014.05.018

Gleich, O., Semmler, P., & Strutz, J. (2016). Behavioral auditory thresholds and loss of ribbon synapses at inner hair cells in aged gerbils. *Experimental Gerontology*, *84*, 61–70. https://doi.org/10.1016/j.exger.2016.08.011

Gómez-Álvarez, M., Johannesen, P. T., Coelho-de-Sousa, S. L., Klump, G. M., & Lopez-Poveda, E. A. (2023). The relative contribution of cochlear synaptopathy and reduced inhibition to age-related hearing impairment for people with normal audiograms. *Trends in Hearing*, *27*, 23312165231213190.
https://doi.org/10.1177/23312165231213191

Gratton, M. A., & Schulte, B. A. (1995). Alterations in microvasculature are associated with atrophy of the stria vascularis in quiet-aged gerbils. *Hearing Research*, *82*(1), 44–52.
https://doi.org/https://doi.org/10.1016/0378-5955(94)00161-I

Guest, H., Munro, K. J., Prendergast, G., Millman, R. E., & Plack, C. J. (2018). Impaired speech perception in noise with a normal audiogram: No evidence for cochlear synaptopathy and no relation to lifetime noise exposure. *Hearing Research*, *364*, 142–151. https://doi.org/https://doi.org/10.1016/j.heares.2018.03.008

Harrison, R. V, & Evans, E. F. (1979). Some aspects of temporal coding by single cochlear fibres from regions of cochlear hair cell degeneration in the guinea pig. *Archives of Oto-*

Rhino-Laryngology, 224(1), 71–78. https://doi.org/10.1007/BF00455226

Heeringa, A. N., Jüchter, C., Beutelmann, R., Klump, G. M., & Köppl, C. (2023). Altered
neural encoding of vowels in noise does not affect behavioral vowel discrimination in
gerbils with age-related hearing loss. *Frontiers in Neuroscience, 17*.
https://www.frontiersin.org/journals/neuroscience/articles/10.3389/fnins.2023.12389
41

Heeringa, A. N., Zhang, L., Ashida, G., Beutelmann, R., Steenken, F., & Koeppl, C. (2020).
Temporal coding of single auditory nerve fibers is not degraded in aging gerbils. *The*
Journal of Neuroscience, 40(2), 343–354. https://doi.org/10.1523/JNEUROSCI.2784-
18.2019

Heil, P., Neubauer, H., Irvine, D. R. F., & Brown, M. (2007). Spontaneous activity of
auditory-nerve fibers: insights into stochastic processes at ribbon synapses. *The Journal*
of Neuroscience, 27(31), 8457–8474.

Hellstrom, L. I., & Schmiedt, R. A. (1990). Compound action potential input/output
functions in young and quiet-aged gerbils. *Hearing Research, 50*(1), 163–174.
https://doi.org/https://doi.org/10.1016/0378-5955(90)90042-N

Hellstrom, L. I., & Schmiedt, R. A. (1996). Measures of tuning and suppression in single-fiber
and whole-nerve responses in young and quiet-aged gerbils. *The Journal of the*
Acoustical Society of America, 100(5), 3275–3285. https://doi.org/10.1121/1.417211

Henry, K. S. (2022). Animal models of hidden hearing loss: Does auditory-nerve-fiber loss
cause real-world listening difficulties? *Molecular and Cellular Neuroscience, 118*,
103692. https://doi.org/https://doi.org/10.1016/j.mcn.2021.103692

Henry, K. S., & Abrams, K. S. (2021). Normal tone-in-noise sensitivity in trained budgerigars
despite substantial auditory-nerve injury: No evidence of hidden hearing loss. *The*
Journal of Neuroscience, 41(1), 118 LP – 129.
https://doi.org/10.1523/JNEUROSCI.2104-20.2020

Henry, K. S., & Heinz, M. G. (2012). Diminished temporal coding with sensorineural hearing
loss emerges in background noise. *Nature Neuroscience, 15*(10), 1362–1364.
https://doi.org/10.1038/nn.3216

Henry, K. S., Sayles, M., Hickox, A. E., & Heinz, M. G. (2019). Divergent auditory nerve
encoding deficits between two common etiologies of sensorineural hearing loss. *The*
Journal of Neuroscience, 39(35), 6879 LP – 6887.
https://doi.org/10.1523/JNEUROSCI.0038-19.2019

Jackson, H. M., & Moore, B. C. J (2014). The role of excitation-pattern, temporal-fine-
structure, and envelope cues in the discrimination of complex tones. *The Journal of the*
Acoustical Society of America, 135(3), 1356–1370.
https://dx.doi.org/10.1121/1.4864306

Jeng, J., Ceriani, F., Olt, J., Brown, S. D. M., Holley, M. C., Bowl, M. R., Johnson, S. L., &
Marcotti, W. (2020). Pathophysiological changes in inner hair cell ribbon synapses in
the ageing mammalian cochlea. *Journal of Physiology, 598*(19), 4339–4355.
https://doi.org/10.1113/JP280018

Johannesen, P. T., Buzo, B. C., & Lopez-Poveda, E. A. (2019). Evidence for age-related
cochlear synaptopathy in humans unconnected to speech-in-noise intelligibility deficits.
Hearing Research, 374, 35–48.
https://doi.org/https://doi.org/10.1016/j.heares.2019.01.017

Joris, P. X., Schreiner, C. E., & Rees, A. (2004). Neural processing of amplitude-modulated
sounds. *Physiological Reviews, 84*(2), 541–577.
https://doi.org/10.1152/physrev.00029.2003

Joris, P. X., Smith, P. H., & Yin, T. C. T. (1994). Enhancement of neural synchronization in
the anteroventral cochlear nucleus. II. Responses in the tuning curve tail. *Journal of*
Neurophysiology, 71(3), 1037–1051. https://doi.org/10.1152/jn.1994.71.3.1022

Joris, P. X., & Yin, T. C. T. (1992). Responses to amplitude-modulated tones in the auditory
nerve of the cat. *The Journal of the Acoustical Society of America, 91*(1), 215–232.
https://doi.org/10.1121/1.402757

Kale, S., & Heinz, M. G. (2010). Envelope coding in auditory nerve fibers following noise-
induced hearing loss. *Journal of the Association for Research in Otolaryngology :*
JARO, 11(4), 657–673. https://doi.org/10.1007/s10162-010-0223-6

Kale, S., Micheyl, C., & Heinz, M. G. (2013). Effects of sensorineural hearing loss on temporal coding of harmonic and inharmonic tone complexes in the auditory nerve. *Advances in Experimental Medicine and Biology*, *787*, 109–118. https://doi.org/10.1007/978-1-4614-1590-9_13

Kale, S., Micheyl, C., & Heinz, M. G. (2014). Implications of within-fiber temporal coding for perceptual studies of F0 discrimination and discrimination of harmonic and inharmonic tone complexes. *Journal of the Association for Research in Otolaryngology : JARO*, *15*(3), 465–482. https://doi.org/10.1007/s10162-014-0451-2

Keine, C., & Ruebsamen, R. (2015). Inhibition shapes acoustic responsiveness in spherical bushy cells. *The Journal of Neuroscience 35*(22), 8579–8592. https://doi.org/10.1523/JNEUROSCI.0133-15.2015

Kessler, M., Mamach, M., Beutelmann, R., Lukacevic, M., Eilert, S., Bascuñana, P., Fasel, A., Bengel, F. M., Bankstahl, J. P., Ross, T. L., Klump, G. M., & Berding, G. (2020). GABA(A) receptors in the Mongolian gerbil: a PET study using [(18)F]flumazenil to determine receptor binding in young and old animals. *Molecular Imaging and Biology*, *22*(2), 335–347. https://doi.org/10.1007/s11307-019-01371-0

Kittel, M., Wagner, E., & Klump, G. M. (2002). An estimate of the auditory-filter bandwidth in the Mongolian gerbil. *Hearing Research*, *164*(1), 69–76. https://doi.org/https://doi.org/10.1016/S0378-5955(01)00411-7

Kujawa, S. G., & Liberman, M. C. (2009). Adding insult to injury: Cochlear nerve degeneration after "temporary" noise-induced hearing loss. *The Journal of Neuroscience*, *29*(45), 14077–14085. https://doi.org/10.1523/JNEUROSCI.2845-09.2009

Lang, H., Schulte, B. A., & Schmiedt, R. A. (2005). Ouabain induces apoptotic cell death in type I spiral ganglion neurons, but not type II neurons. *Journal of the Association for Research in Otolaryngology : JARO*, *6*(1), 63–74. https://doi.org/10.1007/s10162-004-5021-6

Liberman, M. C. (2017). Noise-induced and age-related hearing loss: new perspectives and potential therapies [version 1; peer review: 4 approved]. *F1000Research*, *6*(927). https://doi.org/10.12688/f1000research.11310.1

Liberman, M. C., Epstein, M. J., Cleveland, S. S., Wang, H., & Maison, S. F. (2016). Toward a differential diagnosis of hidden hearing loss in humans. *PLOS ONE*, *11*(9), e0162726-. https://doi.org/10.1371/journal.pone.0162726

Liberman, M. C., & Kujawa, S. G. (2017). Cochlear synaptopathy in acquired sensorineural hearing loss: Manifestations and mechanisms. *Hearing Research*, *349*, 138–147. https://doi.org/10.1016/j.heares.2017.01.003

Lin, H. W., Furman, A. C., Kujawa, S. G., & Liberman, M. C. (2011). Primary Neural Degeneration in the Guinea Pig Cochlea After Reversible Noise-Induced Threshold Shift. *Journal of the Association for Research in Otolaryngology*, *12*(5), 605–616. https://doi.org/10.1007/s10162-011-0277-0

Lopez-Poveda, E. A. (2014). Why do I hear but not understand? Stochastic undersampling as a model of degraded neural encoding of speech. *Frontiers in Neuroscience*, *8*, 348. https://doi.org/10.3389/fnins.2014.00348

Mackey, C. A., McCrate, J., MacDonald, K. S., Feller, J., Liberman, L., Liberman, M. C., Hackett, T. A., & Ramachandran, R. (2021). Correlations between cochlear pathophysiology and behavioral measures of temporal and spatial processing in noise exposed macaques. *Hearing Research*, *401*, 108156. https://doi.org/https://doi.org/10.1016/j.heares.2020.108156

Marmel, F., Plack, C. J., Hopkins, K., Carlyon, R. P., Gockel, H. E., & Moore, B. C. J. (2015). The role of excitation-pattern cues in the detection of frequency shifts in bandpass-filtered complex tones. *The Journal of the Acoustical Society of America*, *137*(5), 2687–2697. https://doi.org/10.1121/1.4919315

Mathew, A. K., Purdy, S. C., Welch, D., Pontoppidan, N. H., & Rønne, F. M. (2016). Electrophysiological and behavioural processing of complex acoustic cues. *Clinical Neurophysiology*, *127*(1), 779–789. https://doi.org/10.1016/j.clinph.2015.04.002

Miller, R. L., Schilling, J. R., Franck, K. R., & Young, E. D. (1997). Effects of acoustic trauma on the representation of the vowel /ɛ/ in cat auditory nerve fibers. *The Journal of the*

Acoustical Society of America, *101*(6), 3602–3616. https://doi.org/10.1121/1.418321

Mishra, S. K., Pada, M. R., & Raj, S. (2015). Influence of musical training on sensitivity to temporal fine structure. *International Journal of Audiology 54*, 220-226. https://doi.org/10.3109/14992027.2014.969411

Moore, B. C., Glasberg, B. R., & Hopkins, K. (2006). Frequency discrimination of complex tones by hearing-impaired subjects: Evidence for loss of ability to use temporal fine structure. *Hearing Research*, *222*(1–2), 16–27. https://doi.org/10.1016/j.heares.2006.08.007

Moore, B. C. J. (2014). Auditory processing of temporal fine structure: effects of age and hearing loss. In *Auditory processing of temporal fine structure*. World Scientific Publishing Co. Pte. Ltd.

Moore, B. C. J. (2019). The roles of temporal envelope and fine structure information in auditory perception. *Acoustical Science and Technology*, *40*(2), 61–83. https://doi.org/10.1250/ast.40.61

Moore, B. C. J., Glasberg, B. R., Stoev, M., Fuellgrabe, C., & Hopkins, K. (2012). The influence of age and high-frequency hearing loss on sensitivity to temporal fine structure at low frequencies (L). *The Journal of the Acoustical Society of America*, *131*(2, 1), 1003–1006. https://doi.org/10.1121/1.3672808

Moore, B. C. J., & Sek, A. (2009). Development of a fast method for determining sensitivity to temporal fine structure. *International Journal of Audiology*, *48*(4), 161–171. https://doi.org/10.1080/14992020802475235

Moore, G. A., & Moore, B. C. J. (2003). Perception of the low pitch of frequency-shifted complexes. *The Journal of the Acoustical Society of America*, *113*(2), 977–985. https://doi.org/10.1121/1.1536631

Müller, M. (1996). The cochlear place-frequency map of the adult and developing mongolian gerbil. *Hearing Research*, *94*(1–2), 148–156. https://doi.org/10.1016/0378-5955(95)00230-8

National Institute of Mental Health. (2002). *Methods and Welfare Considerations in Behavioral Research with Animals: Report of a National Institutes of Health Workshop*. (A. Morrison, H. Evans, N. Ator, & R. Nakamura, Eds.). NIH Publication No. 02-5083.

O'Brien, W. J., Lingrel, J. B., & Wallick, E. T. (1994). Ouabain binding kinetics of the rat alpha two and alpha three isoforms of the sodium-potassium adenosine triphosphate. *Archives of Biochemistry and Biophysics 310*: 32–39. https://doi.org/10.1006/abbi.1994.1136

Parthasarathy, A., Hancock, K. E., Bennett, K., DeGruttola, V., & Polley, D. B. (2020). Bottom-up and top-down neural signatures of disordered multi-talker speech perception in adults with normal hearing. *ELife*, *9*, e51419. https://doi.org/10.7554/eLife.51419

Parthasarathy, A., & Kujawa, S. G. (2018). Synaptopathy in the aging cochlea: characterizing early-neural deficits in auditory temporal envelope processing. *The Journal of Neuroscience*, *38*(32), 7108–7119. https://doi.org/10.1523/JNEUROSCI.3240-17.2018

Prendergast, G., Couth, S., Millman, R. E., Guest, H., Kluk, K., Munro, K. J., & Plack, C. J. (2019). Effects of age and noise exposure on proxy measures of cochlear synaptopathy. *Trends in Hearing*, *23*, 2331216519877301. https://doi.org/10.1177/2331216519877301

Prendergast, G., Millman, R. E., Guest, H., Munro, K. J., Kluk, K., Dewey, R. S., Hall, D. A., Heinz, M. G., & Plack, C. J. (2017). Effects of noise exposure on young adults with normal audiograms II: Behavioral measures. *Hearing Research*, *356*, 74–86. https://doi.org/https://doi.org/10.1016/j.heares.2017.10.007

Rose, J. E., Brugge, J. F., Anderson, D. J., & Hind, J. E. (1967). Phase-locked response to low-frequency tones in single auditory nerve fibers of the squirrel monkey. *Journal of Neurophysiology*, *30*(4), 769–793. https://doi.org/10.1152/jn.1967.30.4.769

Schaette, R., & McAlpine, D. (2011). Tinnitus with a normal audiogram: Physiological evidence for hidden hearing loss and computational model. *The Journal of Neuroscienc 31*(38), 13452–13457. https://doi.org/10.1523/JNEUROSCI.2156-11.2011

Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T., Preibisch, S., Rueden, C., Saalfeld, S., Schmid, B., Tinevez, J.-Y., White, D. J., Hartenstein, V.,

Eliceiri, K., Tomancak, P., & Cardona, A. (2012). Fiji: an open-source platform for biological-image analysis. *Nature Methods*, *9*(7), 676–682. https://doi.org/10.1038/nmeth.2019

Schmiedt, R. A., Mills, J. H., & Boettcher, F. A. (1996). Age-related loss of activity of auditory-nerve fibers. *Journal of Neurophysiology*, *76*(4), 2799–2803. https://doi.org/10.1152/jn.1996.76.4.2799

Schmiedt, R. A., Okamura, H., Lang, H., & Schulte, B. A. (2002). Ouabain application to the round window of the gerbil cochlea: A model of auditory neuropathy and apoptosis. *Journal of the Association for Research in Otolaryngology : JARO.*, *3*(3), 223–233. https://doi.org/10.1007/s101620220017

Schulte, B. A., & Schmiedt, R. A. (1992). Lateral wall Na,K-ATPase and endocochlear potentials decline with age in quiet-reared gerbils. *Hearing Research*, *61*(1–2), 35–46. https://doi.org/10.1016/0378-5955(92)90034-k

Sergeyenko, Y., Lall, K., Liberman, M. C., & Kujawa, S. G. (2013). Age-related cochlear synaptopathy: An early-onset contributor to auditory functional decline. *The Journal of Neuroscience*, *33*(34), 13686–13694. https://doi.org/10.1523/JNEUROSCI.1783-13.2013

Shi, L., Chang, Y., Li, X., Aiken, S. J., Liu, L., & Wang, J. (2016). Coding deficits in noise-induced hidden hearing loss may stem from incomplete repair of ribbon synapses in the cochlea. *Frontiers in Neuroscience, 10*. https://www.frontiersin.org/journals/neuroscience/articles/10.3389/fnins.2016.00231

Sinex, D. G., López, D. E., & Warr, W. B. (2001). Electrophysiological responses of cochlear root neurons. *Hearing Research*, *158*(1–2), 28–38. https://doi.org/10.1016/S0378-5955(01)00293-3

Steenken, F., Bovee, S., & Köppl, C. (2022). Immunolabeling and Counting Ribbon Synapses in Young Adult and Aged Gerbil Cochleae. *JoVE, 182*, e63874. https://doi.org/doi:10.3791/63874

Steenken, F., Heeringa, A. N., Beutelmann, R., Zhang, L., Bovee, S., Klump, G. M., & Köppl, C. (2021). Age-related decline in cochlear ribbon synapses and its relation to different metrics of auditory-nerve activity. *Neurobiology of Aging*. https://doi.org/https://doi.org/10.1016/j.neurobiolaging.2021.08.019

Tarnowska, E., Wicher, A., & Moore, B. C. J. (2019). The effect of musicianship, contralateral noise, and ear of presentation on the detection of changes in temporal fine structure. The Journal of the Acoustical Society of America 146 (1), 1-10. https://doi.org/10.1121/1.5114820

Versteegh, C., Meenderink, S. W. F., & Heijden, M. van der. (2011). Response characteristics in the apex of the gerbil cochlea studied through auditory nerve recordings. *Journal of the Association for Research in Otolaryngology*, *12*(3), 301–316. https://doi.org/10.1007/s10162-010-0255-y

Viana, L. M., O'Malley, J. T., Burgess, B. J., Jones, D. D., Oliveira, C. A. C. P., Santos, F., Merchant, S. N., Liberman, L. D., & Liberman, M. C. (2015). Cochlear neuropathy in human presbycusis: Confocal analysis of hidden hearing loss in post-mortem tissue. *Hearing Research*, *327*, 78–88. https://doi.org/10.1016/j.heares.2015.04.014

Wu, P. Z., Liberman, L. D., Bennett, K., de Gruttola, V., O'Malley, J. T., & Liberman, M. C. (2019). Primary neural degeneration in the human cochlea: Evidence for hidden hearing loss in the aging ear. *Neuroscience*, *407*, 8–20. https://doi.org/10.1016/j.neuroscience.2018.07.053

Supplemental Information

This supplemental information summarizes extended statistical results, which are not essential for the support of the findings in the main manuscript.

Synapse counts

supplemental to "*Synapse loss was similar in old gerbils and gerbils treated with a high dosage of ouabain*"

	2 kHz	4 kHz	8 kHz	16 kHz
young-adult	23 (+/-0,5, N=19)	21 (+/-0,4, N=18)	19 (+/-0,5, N=18)	20 (+/-0,5, N=18)
sham	23 (+/-0,5, N=3)	18 (+/-1, N=3)	16 (+/-0,4, N=3)	21 (+/-0,4, N=3)
ouabain low	22 (+/-0,7, N=8)	19 (+/-0,6, N=8)	16 (+/-0,3, N=8)	19 (+/-0,6, N=8)
ouabain high	19 (+/-1,4, N=10)	17 (+/-1,4, N=10)	12 (+/-1,7, N=10)	13 (+/-2,5, N=11)
old	19 (+/-0,7, N=14)	17 (+/-0,8, N=12)	15 (+/-0,6, N=12)	16 (+/-0,6, N=13)

Mean number of functional synapses per IHC, at the cochlear positions corresponding to 2, 4, 8, and 16 kHz, with standard errors of the mean and number of ears.

The mean number of functional synapses per IHC was significantly different between gerbil groups for cochlear locations corresponding to all four frequencies.

(univariate ANOVAs: 2 kHz: F = 5.496, p = 9.782×10^{-4}; 4 kHz: F = 4.995, p = 1.988×10^{-3}; 8 kHz: F = 7.933, p = 5.974×10^{-5}; 16 kHz: F = 6.176, p = 4.305×10^{-4})

Post-hoc tests revealed significant differences (at the 5% level, with Bonferroni correction) between the young-adult and the ouabain-high groups for all equivalent frequencies, between the young-adult and the old group for 2, 4, and 8 kHz, and between the ouabain-low and the ouabain-high group for 16 kHz.

Fig. S1: Synapse number was reduced after ouabain treatment and in old age. Number of functional synapses per IHC, at the cochlear positions corresponding to 2, 4, 8, and 16 kHz, for young-adult gerbils (red), sham-treated gerbils (surgery only, orange), gerbils treated with a low dose (40µM) of ouabain (green), gerbils treated with a high dose (70µM) of ouabain (blue), and old gerbils (gray). Box plots display the median (center line), mean (white circled dot), 25th and 75th percentiles (upper and lower edges of the boxes), and maximum and minimum (whiskers).

Fig. S2: Median spontaneous rates did not differ between animal groups. Individual spontaneous rates for all auditory-nerve fibers reported in this study (dots), group medians (box center lines), quartiles (box boundaries), minima, and maxima (whiskers). The dashed black line indicates the boundary between low- and high-spontaneous-rate fibers (18 spikes/s). A Kruskal-Wallis test confirmed that the group medians are not significantly different at the 5% level.

Fig. S3: Median best frequency was lower in the young-adult sample. Individual best frequencies for all auditory-nerve fibers reported in this study (dots), and box plots in the same style as in Fig. S1. The dashed black line indicates the boundary between low and high best frequency class (1.85 kHz), as used in the results statistics. A Kruskal-Wallis test confirmed that the group medians are significantly different at the 5% level, with a Bonferroni-corrected post-hoc test showing that only the young-adult group median differs significantly from all other groups.

Below, we list the full ANOVA results, including the principal effects that are reported in the main manuscript. The factors used in the ANOVAs reported here had the following factor levels:

- **group:** young-adult, sham, ouabain-high, old
- **condition:** 200/1600, 400/1600, 400/3200 (f_0/f_c in Hz)
- **BF class:** low: BF ≤ 1850 Hz, high: BF > 1850 Hz
- **SR class:** low: SR ≤ 18 spikes/s, high: SR > 18 spikes/s
- **Δf%:** as a covariate ranging from 0 to 21.25%

All significances are reported at the 5% level, with Bonferroni correction for post-hoc tests and planned comparisons.

Effects on mean driven rate
supplemental to "*Average rate does not carry sufficient information about inharmonic frequency shift*"

Note that driven rate (mean rate - spontaneous rate) was used here. The differences found below are consistent with known differences between low-SR (higher driven rates) and high-SR fibers (lower driven rates; Steenken et al., 2021), and with expected differences across BF classes (low-BF fibers less driven by stimuli with f_c = 3200 Hz and high-BF fibers less driven by stimuli with f_c = 1600 Hz). Differences between groups are also consistent with the sample distributions shown in Figs. S1 and S2.

- **Significant main effects:**
    - **group:** $F_{(3,943)}$=10.017, p=1.676×10^{-6}
        - Mean driven rate was lower in sham group than in any other group.
    - **BF class:** $F_{(1,943)}$=6.297, p=0.0123
        - Mean driven rate was lower for high BF class (BF > 1850 Hz).
- **Significant two-way interactions:**
    - **condition × group:** $F_{(6,943)}$=2.525, p=0.0198
        - Planned comparisons between groups within each condition, and between conditions within each group:
          In the 400/3200 condition, the mean driven rate of the sham group was significantly lower than in any other group. The mean driven rate in the sham group was significantly lower than in the young-adult group for each condition.
    - **condition × BF class:** $F_{(2,943)}$=10.562, p=2.909×10^{-5}
        - Planned comparisons between conditions within each BF class, and between BF classes within each condition:
          In the x/1600 conditions the mean driven rate in the high-BF class was significantly lower than in the low-BF class. No significant differences between conditions within each BF class.
    - **group × BF class:** $F_{(3,943)}$=10.189, p=1.316×10^{-6}
        - Planned comparisons between groups within each BF class, and between BF classes within each group:
          In the low-BF class, the mean driven rate in the sham group was significantly lower than in all other groups. For the ouabain-high and old groups, the mean driven rate in the high-BF class was lower than in the low-BF class.

- o **group × SR class:** $F_{(2,943)}=8.148$, $p=3.103\times10^{-4}$
  - – Planned comparisons between groups within each SR class, and between SR classes within each group:
    In the young-adult group, the mean driven rate in the high-SR class was lower than in the low-SR class. In the high-SR class, the mean driven rate in the sham group was lower than in any other group. (No samples for the sham group in the low-SR class).
- **Non-significant main effects and interactions:**
  - o *Δf%: $F_{(1,943)}=0.018$, p=0.8933 (reported in main manuscript)*
  - o **condition:** $F_{(2,943)}=2.187$, $p=0.1129$
  - o **SR class:** $F_{(1,943)}=1.884$, $p=0.1702$
  - o **group × Δf%:** $F_{(3,943)}=0.394$, $p=0.7575$
  - o **BF class × Δf%:** $F_{(1,943)}=0.052$, $p=0.8195$
  - o **condition × Δf%:** $F_{(2,943)}=0.081$, $p=0.9219$
  - o **SR class × Δf%:** $F_{(1,943)}=0.002$, $p=0.9657$
  - o **condition × SR class:** $F_{(2,943)}=0.698$, $p=0.4977$
  - o **BF class × SR class:** $F_{(1,943)}=0.388$, $p=0.5336$

Effects on TFS representation (TFS log-z-ratio)
supplemental to "*Neural representation of TFS was not degraded in AN fibers of old or synaptopathic gerbils*"

A high TFS log-z-ratio indicates strong phase locking to the stimulus TFS.

- **Significant main effects:**
  - Δ**f%:** $F(1,943)=104.403$, $p=2.569\times10^{-23}$ (reported in main manuscript)
  - **condition:** $F(2,943)=7.495$, $p=5.894\times10^{-4}$ (reported in main manuscript)
    - TFS log-z-ratio was significantly lower in the 400/3200 condition, compared to both the 200/1600 and 400/1600 conditions
- **Significant two-way interactions:**
  - **condition × Δf%:** $F(2,943)=20.292$, $p=2.353\times10^{-9}$ (reported in main manuscript)
    - Based on the parameter estimates of the ANOVA model:
      The slope of the TFS log-z-ratio vs. Δf% was significantly different from 0, for the 200/1600 and 400/1600 condition. It was significantly higher in the 400/1600 condition, compared to the 200/1600 condition.
  - **BF class × Δf%:** $F(1,943)=31.05$, $p=3.282\times10^{-8}$ (reported in main manuscript)
    - Based on the parameter estimates of the ANOVA model:
      The slope of the TFS log-z-ratio vs. Δf% was significantly higher in the low-BF class than in the high-BF class.
  - **condition × group:** $F(6,943)=7.583$, $p=5.808\times10^{-8}$
    - planned comparisons between groups within each condition, and between conditions within each group:
      In the 200/1600 condition, TFS log-z-ratio in the ouabain-high group was significantly higher than in the old group. For all groups except the old group, TFS log-z-ratio in the 400/1600 condition was significantly higher than in the 400/3200 condition. In the sham group, the TFS log-z-ratio in the 400/1600 condition was significantly higher than in the 200/1600 condition. In the ouabain-high group, the TFS log-z-ratio in the 200/1600 condition was significantly higher than in the 400/3200 condition.
  - **condition × BF class:** $F(2,943)=5.774$, $p=3.200\times10^{-3}$
    - Planned comparisons between BF classes within each condition, and between conditions within each BF class:
      In the high-BF class, TFS log-z-ratio in the 400/3200 condition was significantly lower than in any other condition. In the 400/3200 condition, the TFS log-z-ratio in the high-BF class was significantly lower than in the low-BF class.

- o **group × BF class:** $F(3,943)=4.228$, $p=5.600\times10^{-3}$
  - – Planned comparisons between groups within each BF class, and between BF classes within each group:
    In the high-BF class, the TFS log-z-ratio in the ouabain-high group was significantly higher than in the old group.
- o **BF class × SR class:** $F(1,943)=4.918$, $p=0.0268$
  - – In the high-SR class, TFS log-z-ratio in the low-BF class was significantly higher than in the high-BF class.

- **Non-significant main effects and interactions:**
  - o *group: $F(3,943)=1.757$, $p=0.1537$ (reported in main manuscript)*
  - o *group × Δf%: $F(3,943)=1.193$, $p=0.3112$ (reported in main manuscript)*
  - o **BF class:** $F(1,943)=0.103$, $p=0.7479$
  - o **SR class:** $F(1,943)=0.985$, $p=0.3212$
  - o **condition × SR class:** $F(2,943)=2.302$, $p=0.1006$
  - o **group × SR class:** $F(2,943)=0.556$, $p=0.5736$
  - o **SR class × Δf%:** $F(1,943)=1.203$, $p=0.2730$

Effects on ENV/TFS representation balance (ENV/TFS log-z-ratio)
supplemental to "*Representation of f_0 was enhanced in AN fibers of old gerbils*"

The ENV/TFS log-z-ratio reflects the relative emphasis in the AN fibers' responses on either TFS (negative ENV/TFS log-z-ratio) or envelope locking (positive ENV/TFS log-z-ratio).

- **Significant main effects:**
  - ○ *condition: F(2,943)=43.929, p=5.744×10⁻¹⁹ (reported in main manuscript)*

    — The 200/1600 and 400/1600 conditions had negative ENV/TFS log-z-ratios, the 400/3200 condition had a positive ENV/TFS log-z-ratio. All were significantly different from each other, with the 200/1600 condition showing the most negative value.
  - ○ **SR class:** *F(1,943)=29.138, p=8.533×10⁻⁰⁸ (reported in main manuscript)*

    — In the high-SR class, the ENV/TFS log-z-ratio was negative and significantly different from zero. In the low-SR class, it was not significantly different from zero.
  - ○ **Δ*f*%:** *F(1,943)=4.278, p=0.0389 (reported in main manuscript)*
- **Significant two-way interactions:**
  - ○ *condition × group: F(6,943)=11.439, p=2.242×10⁻¹² (reported in main manuscript)*

    — Planned comparisons between groups within each condition, and between conditions within each group:

    In addition to the main condition effect: In the 200/1600 condition, the ENV/TFS log-z-ratio was more negative in the sham group than both in the young-adult group, and in the old group. In the 400/3200 condition, the ENV/TFS log-z-ratio was more positive in the old group than in the young-adult group. In the sham group, the ENV/TFS log-z-ratio was more negative in the 200/1600 condition than in the 400/1600 condition.
  - ○ **group × BF class:** F(3,943)=11.354, p=2.551×10⁻⁷

    — Planned comparisons between groups within each BF class, and between BF classes within each group:

    In the high-BF class, the ENV/TFS log-z-ratio was significantly more negative in the sham and ouabain-high groups than in the young-adult and old groups.
  - ○ **BF class × SR class:** F(1,943)=5.234, p=0.0224

    — In the order from most negative to most positive ENV/TFS log-z-ratio: low-BF/high-SR, high-BF/high-SR, high-BF/low-SR, low-BF/low-SR. Both high-SR values were significantly below zero, and different from each other; both low-SR values were not significantly different from zero, and not different from each other.

- **Non-significant main effects and interactions:**
    - **group:** $F_{(3,943)}=0.838$, $p=0.4731$
    - **BF class:** $F_{(1,943)}=0.439$, $p=0.5076$
    - **group × Δf%:** $F_{(3,943)}=0.236$, $p=0.8711$
    - **condition × Δf%:** $F_{(2,943)}=1.673$, $p=0.1883$
    - **BF class × Δf%:** $F_{(1,943)}=0.005$, $p=0.9410$
    - **SR class × Δf%:** $F_{(1,943)}=0.58$, $p=0.4464$
    - **condition × BF class:** $F_{(2,943)}=0.414$, $p=0.6611$
    - **condition × SR class:** $F_{(2,943)}=0.522$, $p=0.5936$
    - **group × SR class:** $F_{(2,943)}=2.442$, $p=0.0875$