Tracing active members in microbial communities by BONCAT and click chemistry-based enrichment of newly synthesised proteins

Patrick Hellwig[1,2]*, Daniel Kautzner[3], Robert Heyer[3,4], Anna Dittrich[5], Daniel Wibberg[6,7], Tobias Busche[8,9], Anika Winkler[8,9], Udo Reichl[1,2], Dirk Benndorf[1,2,10]*

*:   corresponding author

1.   Otto-von-Guericke University Magdeburg, Bioprocess Engineering, Universitätsplatz 2, 39106 Magdeburg, Germany;

2.   Bioprocess Engineering, Max Planck Institute for Dynamics of Complex Technical Systems, Sandtorstraße 1, 39106 Magdeburg, Germany

3.   Multidimensional Omics Analyses group, Faculty of Technology, Bielefeld University, Universitätsstraße 25, 33615 Bielefeld,

4.   Multidimensional Omics Analyses group, Leibniz-Institut für Analytische Wissenschaften – ISAS – e.V., Bunsen-Kirchhoff-Straße 11, 44139 Dortmund

5.   Department of Systems Biology, Institute of Biology, Otto-von-Guericke University Magdeburg, , Universitätsplatz 2, 39106 Magdeburg, Germany

6.   Institute for Genome Research and Systems Biology, CeBiTec, Bielefeld University, Universitätsstraße 25, 33615 Bielefeld, Germany

7.   Institute of Bio- and Geosciences IBG-5, Computational Metagenomics, Forschungszentrum Jülich GmbH, 52425 Juelich, Germany

8.   Center for Biotechnology - CeBiTec, Bielefeld University, Universitätsstraße 27, D-33615 Bielefeld, Germany.

9.   Medical School East Westphalia-Lippe, Bielefeld University, Universitätsstraße 27, D-33615 Bielefeld, Germany.

10. Microbiology, Anhalt University of Applied Sciences, Bernburger Straße 55, 06354 Köthen, Germany

ORCID ID:

•   P. Hellwig: 0000-0003-3280-9042

•   A. Dittrich: 0000-0002-1056-4219

•   D. Wibberg: 0000-0002-1331-4311

•   T. Busche: 0000-0001-9211-8927

•   U. Reichl: 0000-0001-6538-1332

Emails:

hellwig@mpi-magdeburg.mpg.de

daniel.kautzner@uni-bielefeld.de

anna.dittrich@ovgu.de

robert.heyer@isas.de

dwibberg@cebitec.uni-bielefeld.de

tbusche@cebitec.uni-bielefeld.de

awinkler@cebitec.uni-bielefeld.de

ureichl@mpi-magdeburg.mpg.de

benndorf@mpi-magdeburg.mpg.de

Keywords: Metaproteomics; mass spectrometry; BONCAT; click chemistry; microbiome;

microbial activity; anaerobic digestion; biogas; microbial interactions; Metagenomics

Running title: Microbial Protein Synthesis Tracking

Abstract (196 words):

A comprehensive understanding of microbial community dynamics is fundamental to the

advancement of environmental microbiology, human health, and biotechnology.

Metaproteomics, i.e. the analysis of all proteins in a microbial community, provides insights

into these complex systems. Microbial adaptation and activity depend to an important extent

on newly synthesized proteins (nP), however, the distinction between nP and bulk proteins is

challenging. The application of bioorthogonal non-canonical amino acid tagging (BONCAT)

with click chemistry has demonstrated efficacy in the enrichment of nP in pure cultures.

However, the transfer of this technique to microbial communities has proven challenging and

has therefore not been used on microbial communities before. To address this, a new

workflow with efficient and specific nP enrichment was developed using a laboratory-scale

mixture of labelled *E. coli* and unlabelled yeast. This workflow was successfully applied to an

anaerobic microbial community with initially low BONCAT efficiency. A substrate shift from

glucose to ethanol selectively enriched nP with minimal background. The identification of

bifunctional alcohol dehydrogenase and a syntrophic interaction between an ethanol-utilizing

bacterium and two methanogens (hydrogenotrophic and acetoclastic) demonstrates the

potential of metaproteomics targeting nP to trace microbial activity in complex microbial

communities.

Abbreviations:

AA	amino acids
AD	anaerobic digestion
ADH	alcohol dehydrogenase
AHA	4-azido-l-homoalanine
BONCAT	bioorthogonal non-canonical amino acid tagging
BSA	bovine serum albumin
CC	click chemistry
DBCO	dibenzocyclooctyne
DTT	dithiothreitol
FASP	filter-aided sample preparation
HPG	homopropargylglycine
HPLC	high-pressure liquid chromatography
IAA	iodoacetamide
KEGG	Kyoto Encyclopedia of Genes and Genomes
LBR	laboratory biogas reactor
LC	liquid chromatography
MAG	metagenome assembled genome
MC	microbial community
MS	mass spectrometry
Mgf	mascot generic file
ncAA	non-canonical amino acids
nP	newly synthesized proteins
PASEF	parallel accumulation–serial fragmentation

PBS	phosphate-buffered saline
PEG	polyethylene glycol
rpm	rounds per minute
RT	room temperature
SDS	sodium dodecyl sulfate
SIP	stable isotope labeling
SOP	standard operation procedure
SS	disulfide
TYGS	Type (Strain) Genome Server

56 **Graphical Abstract:**

57
58

Introduction

Microbial communities (MC) include single-celled eukaryotes, archaea, bacteria, and viruses

[1–3]. These communities are found in various environments, including soil and water, and

within living organisms such as animals and humans. Additionally, MC are used for industrial

processes such as wastewater treatment or energy production e.g. in anaerobic digestion (AD)

[4]. The understanding of MC is still very limited due to their inherent complexity. Enhancing

our understanding will help to cure diseases, optimize biotechnological processes, and even

act against climate change [5–7]. Due to its immense importance in the analysis and

characterization of MC gained prominence in the last decades.

Metagenomics is currently the most widely used method to characterise MCs. It involves the

simultaneous genomic analysis of all members of an MC and provides insight into the

taxonomic composition and metabolic potential of an MC [8]. However, only in combination

with other meta-omics techniques such as metaproteomics, which analyses all proteins of an

MC, it is possible to fully decipher the functions of the MC [9].

Microbial adaptation to changing environments or process parameters and the resulting

changed microbial activity occurs primarily through the expression of newly synthesized

proteins (nP), which can be hardly distinguished from already present bulk proteins.

Consequently, the analysis of the nP would provide a novel dimension of metaproteomics,

thereby facilitating a deeper comprehension of MC activities.

The standard method of tracking nP is stable isotope labeling (SIP). In protein-SIP stable

isotopes, such as ^{13}C, ^{15}N, and ^{2}H (deuterium), are introduced into the cultivation media.

These isotopes are incorporated into amino acids and the modified amino acids will be

incorporated in nP [10, 11]. This allows a distinction between labeled nP from non-labeled

proteins. Alternatively, advances in bioorthogonal chemistry have enabled another way of

identification of nP. One technique is bioorthogonal non-canonical amino acid tagging

(BONCAT), which allows the selective labelling of nP [12]. In BONCAT, analog non-

canonical amino acids (ncAA) are added to cells or MCs. During protein synthesis, the tRNA-

amino acyl synthetase of an amino acid mistakenly selects the ncAA instead of the canonical

amino acid. 4-Azido-L-homoalanine (AHA) and L-Homopropargylglycine (HPG). Both are

the ncAAs replacing methionine [13]. Note, incorporation of AHA and HPG does not depend

on genetic modification of tRNAs in the uptaking cells. Studies in *E. coli* showed that the

likelihood of incorporation is 1:390 for AHA and 1:500 for HPG [13]. The incorporation rate

of ncAA into the amino acid chain of a protein correlates negatively with the concentration of

the targeted amino acid in the medium [14]. This amino acid should therefore be avoided.

However, Ignacio et al. (2023) [15] have recently developed a method called THRONCAT, in

which ncAA incorporation is not affected by external canonical amino acid concentrations,

which could significantly expand the range of applications for ncAA labeling in the future.

The incorporation of ncAAs into nP results in a mass shift that can be detected by mass

spectrometry (MS). Nevertheless, the low proportion of labelled nP in relation to the bulk

proteins in MC is so far insufficient to permit the reliable identification of nP through

metaproteomics.

As a remedy, BONCAT can be combined with click chemistry (CC) in the form of azide-

alkyne cycloaddition [16, 17]. Thus, a fluorescence or biotin marker can be attached to the

AHA (azide group) or HPG (alkyne group) [18]. BONCAT and CC have previously been

applied to visualize and enrich nP from eukaryotic cells, such as neurons [14], HeLa cells

[19], or HEK cells [20] and from viruses and their host cells [21]. However, BONCAT

combined with CC has been used mainly in cell culture experiments with only one cell type.

Hatzenpichler et al. [22] showed that the application of BONCAT in MC is possible and

discussed the advantages and disadvantages of BONCAT and its application to MC in great

detail [23]. Additionally, Reichart et al. [24] showed that sorting active ncAA-labelled

microbes from non-active microbes with flow cytometry is possible. BONCAT and CC also

show promise for bacteriophages in MC [25, 26].

In the case of metaproteomics, the detection of nP is of paramount importance for the

understanding of MC. However, the currently available enrichment methods for nP are

designed for pure cultures with high ncAA-labeling. In contrast, low enrichment efficiencies

of nP and an enormous, unspecific binding of bulk proteins are achieved when using more

complex and less ncAA-labelled MC samples. Therefore, there is a clear need for the

development of reliable, specific, and reproducible methods for the detection of nP in MC *via*

MS. The aim of this work was thus to develop a workflow that allows the enrichment of nP

from growing MC for the detection *via* MS. The main focus was on the enrichment of AHA-

labelled nP since the conditions of BONCAT have to be individually adapted to each

experimental condition and MC in particular to the dynamics of protein translation [22, 27].

Therefore, we developed a test system for nP enrichment. It should demonstrate the

specificity of nP enrichment and be easy to reproduce in any laboratory. Based on these

restrictions, a test system consisting of *E. coli* and *Saccharomyces cerevisiae* (yeast) was

chosen. *E. coli* was labelled with AHA after a substrate shift from glucose to lactose that

induces metabolic changes and thus expression of nP involved in lactose metabolism. Next,

unlabelled yeast cells were added. The established workflow allowed the enrichment of *E.*

coli nPs involved in lactose metabolism without contamination by yeast proteins.

As a proof of concept, the workflow was applied to enrich nP from an anaerobic MC derived

from a laboratory-scale biogas reactor (LBR). Given the slow growth of this anaerobic MC,

the efficiency of ncAA-labeling was expected to be suboptimal. However, if nP can be

enriched from this MC and reliably detected in MS, the developed workflow should be

applicable to other slowly growing MC. The LBR used for this study was operated with

glucose as the primary substrate. The objective was to shift the substrate to ethanol so that the

nP associated with the consumption of ethanol as a secondary substrate would be labelled

with AHA and could subsequently be enriched with the established workflow. It was

anticipated that successful nP enrichment would allow to trace the degradation pathway of

ethanol under anaerobic conditions. It may even be possible to gain new insights into this

poorly described metabolic pathway [28].

Material and Methods

In the following we briefly describe the workflow established (Figure 1). A detailed

description of the methods, including a step-by-step standard operation procedure (SOP), can

be found in Supplementary Note 1. All cultivation experiments were performed in triplicates.

AHA-labeling of *E. coli* nP

The carbon source of an E. coli (DSM 5911) culture grown aerobically at 37°C in M9 media

was switched from glucose to lactose while the cells were incubated with 100 μM AHA

(sterile filtrated). The same substrate shift was done for control cells without adding AHA.

After the harvest, the two *E. coli* cultures were separately mixed with an overnight yeast

culture. The mixture of *E. coli* and yeast is named "test system" in the following (see SOP 1).

AHA-labeling of nP from MC of LBR

50 mL of cell suspensions from an LBR MC (1 L total volume, 40°C, 75 rpm, pH 7.4 – 7.5,

Supplementary Table 1) was sampled using a 50 mL syringe. The cell suspensions were then

transferred to nitrogen-flushed 100 mL serum bottles, diluted 1:1 with 50 mL anaerobic

medium without glucose (Supplementary Table 2), and incubated for one hour. Following

this, 1 mL of 10% (vol./vol.) ethanol solution was added. Additionally, for BONCAT, 1 mL of

20 mM AHA was introduced, while controls received 1 mL of water. An extra control was

prepared with 3.8 mM glucose instead of ethanol. The cultures were sealed and incubated for

24 hours at 40°C with gentle stirring until harvest.

Cell disruption with a ball mill

Samples from the test system and the LBR MC were suspended in 5 mL of 20 mM Tris/HCl-

MgCl buffer (pH 7.5) and treated with 1 g/mL of silica beads and 0.5 µL/mL of Cyanase[TM]

Nuclease solution. The cells were lysed in a ball mill, incubated at 37°C for 15 minutes, and

then centrifuged. The resulting supernatant was transferred to a new reaction tube(see SOP 1).

<u>Protein extraction with acetone (for pure cultures)</u>

The proteins of the test system were precipitated with the five-fold volume of ice-cold acetone

at -20°C for at least 1 h followed by centrifugation. The supernatant was discarded, and the

protein pellets were dried under a fume hood at room temperature (RT). The dry protein

pellets were resuspended in incubation buffer I. The protein samples were centrifuged, and the

supernatants were transferred to new reaction tubes (see SOP 1).

<u>Protein extraction with methanol and chloroform (for environmental samples)</u>

The MC proteins were extracted using a mixture of methanol, chloroform, and water,

followed by centrifugation [29]. After carefully removing the top phase, methanol was added,

and the mixture was vortexed and centrifuged again. The resulting protein pellets were dried

and resuspended in incubation buffer II. After centrifugation, the supernatant was transferred

to new tubes.

<u>Protein quantification</u>

Amido black assay was used to quantify the protein concentration of each sample, as

described earlier [30].

<u>Click chemistry with fluorophores and SDS-PAGE with fluorescent scan</u>

AHA incorporation was assessed using 10 µg protein from each sample. For this purpose, the

dye Dibenzocyclooctyne (DBCO) Cyanin 5.5 was attached using CC following SOP 2 in

Supplementary Note 1. The fluorophore-tagged proteins were separated in a 1.5 mm 10 %

SDS-PAGE and scanned twice: once with a LI-COR Odyssey Classic fluorescence scanner at

700 nm and a second time after Coomassie staining [30, 31]. (see SOP 5).

Click chemistry with biotin linker

Based on the fluorescence signal (previous paragraph), 100 μg protein of the test system and

200 μg protein of the MC were used for the attachment of DBCO-SS-Biotin via CC (see SOP

2).

Enrichment of AHA-labelled proteins

The nP of the test system and the MC were enriched using Dynabeads™ MyOne™

Streptavidin C1 beads and the protocol described in SOP 3. The beads were blocked with

bovine serum albumin (BSA), amino acids (AA), or left blocked.

FASP digestion

The tryptic digestion was performed using FASP [30] with few modifications (see SOP 4).

The proteins contained in the dithiothreitol (DTT) elution phase and in the elution phase of

the SDS boiled beads were precipitated with acetone. 25 μg protein of each sample before CC

("not enriched samples") was precipitated with acetone (control without enrichment). The

resulting protein pellets were dissolved in 200 μL 8 M urea buffer and transferred to a filter

unit (Centrifugal Filter Unit, 10 kDa). In FASP digestion, a ratio of 1:100 (trypsin quantity to

protein quantity) was used. For the enriched nP fractions, 100 ng trypsin was used. After

extraction and concentration via vacuum centrifuge, the eluted peptides were transferred to

HPLC vials for MS measurement.

LC-MS/MS measurements

The peptides of each sample were measured using a timsTOF™ pro mass spectrometer

(Bruker Daltonik GmbH, Bremen, Germany) coupled online to an UltiMate® 3000 nano

splitless reversed-phase nanoHPLC (Thermo Fisher Scientific, Dreieich, Germany) in

PASEF® mode.

Illumina library preparation, MiSeq sequencing, and metagenome assembly for protein

database

Three samples of the LBR used in this study were collected in different weeks for

metagenomic sequencing to generate a protein database for the metaproteomic analysis in this

study. The samples from the LBR were sequenced independently using PCR-free libraries

prepared with the Illumina TruSeq® DNA PCR-free kit [32]. The libraries were subjected to

quality control and sequenced on the MiSeq platform (2×300 bp paired-end, v3 chemistry).

After sequencing, data were stripped of adapters and low-quality reads [33], followed by

assembly with Megahit (v1.1.1) [34] and gene prediction with Prodigal v.2.6.0 [35].

Metagenomic binning [36, 37] was performed with Bowtie 2 [38] and MetaBAT2 (76

Metagenome assembled genome (MAGs)) [39], while completeness and contamination were

assessed with BUSCO (v5.7.0) [40]). The translated amino acid sequences of the predicted

genes were generated and the replicates were merged for a protein database generation with

Contig informations. For the most abundant MAGs, the MAG sequence was uploaded to the

Type (Strain) Genome Server (TYGS) to evaluate the taxonomy [41, 42].

<u>Protein identification using Mascot and the MetaProteomeAnalyzer</u>

With Compass Data Analysis software (version 5.1.0.177, Bruker Corporation, Bremen,

Germany) the raw files were converted into Mascot Generic Files (mgf) and mgf were

searched with Mascot (version 2.6) for the peptide spectrum matches. For the test system, a

defined UniProtKB/SwissProt database (08/05/2022) containing *E. coli* K12

(taxonomy_id:83333) and yeast (taxonomy_id:4932) proteins was used. A metagenome from

the LBR was used for the MC samples (see Illumina library preparation, MiSeq sequencing,

and metagenome assembly). Metaproteins were generated from the MC samples using

MetaProteomeAnalyzer software (version 3.0 [43]).

<u>Further analysis of the protein data</u>

The protein data were uploaded to Prophane (version 6.2.6) [44] for basic local alignment

search tool analysis of the functional and taxonomic annotation. For all unknown KO

numbers after Functional Ontology Assignments for Metagenomes annotation [45],

KofamKOALA [46] was used to identify possible missing KO numbers (Supplementary Table

3 and Supplementary Note 1).

Results and Discussion

<u>Development of a nP enrichment workflow with the test system</u>

Prior to developing the enrichment protocol, the *E. coli* (AHA-labelled) and yeast (unlabelled)

test system was refined by optimising the AHA labelling of *E. coli,* measured by detecting the

proportion of fluorescent *E. coli* cells generated through click chemistry. (Supplementary

Figure 1). Attempts were made to enrich nP of *E. coli* from the test system with known

enrichment protocols (e.g., [20, 21, 47]) but these were unsuccessful (data not shown). The

major issues were that MS of tryptic digests was either unable to detect any proteins, or

detected similar proportions of yeast proteins (unspecific background) and *E. coli* proteins.

The results highlighted the need for a test system to monitor nP enrichment, and the

importance of implementing negative and positive controls to regulate non-specific binding in

each experimental procedure.

A first successful enrichment of nP after click chemistry was obtained with DBCO-SS-PEG3-

biotin and the use of Dynabeads™ MyOne™ Streptavidin C1 beads (Supplementary Figure

2). The nP were eluted under mild conditions from streptavidin beads by the addition of DTT

through the reductive cleavage of the disulfide bridge in the DBCO-SS-PEG3-biotin.

Therefore, DBCO-SS-PEG3-biotin appeared to be an optimal choice for our nP enrichment.

However, many unspecific yeast proteins (background) were co-eluted with the nP

(Supplementary Figure 2).

Therefore, washing was optimized. An adapted washing strategy [20] and the use of

phosphate-buffered saline (PBS) buffer with increased NaCl concentration (237 mM) reduced

the amount of background yeast proteins. However, it strongly reduced the reproducibility of

nP elution (Figure 2 A). For higher reproducibility of enrichment, the loss of nP during elution

should be minimized.

Blocking the beads with 1 mg/mL BSA (e.g. [48, 49]), before loading the samples, reduced

the amount of yeast protein background further but unfortunately also increased the variability

in nP elution (Figure 2 A). Additionally, the elution phase was found to be contaminated with

BSA proteins. Blocking the beads seemed to be important for the removal of non-specific

protein, but BSA was not optimal.

Inspired by Nicora et al. (2013) [50], the beads were blocked with an AA blocking solution

containing 16.67 µg/mL of leucine (aliphatic hydrophobic), tryptophan (aromatic

hydrophobic), histidine (polar positive), glutamine (polar neutral), glutamic acid (polar

negative) and glycine (nonpolar neutral) each. These AAs should not cause steric effects and

should prevent any unspecific interactions with the beads. This method ("AA method")

maintained a low background and exhibited minimal variability and loss in nP elution (Figure

2 A), making it the most promising approach.

<u>Examining the AA method using the test system</u>

In total, 1,558 ± 53 proteins (885 ± 25 *E. coli*, 590 ± 29 yeast) were identified in the AHA-

labelled but non-enriched test system ("not enriched sample") (Figure 2 B). Over a third of

these proteins were yeast proteins. After nP enrichment, 607 ± 16 proteins (555 ± 17 *E. coli*,

21 ± 2 yeast) were identified, reducing the amount of identified yeast proteins by over 96%

while retaining 63% of the *E. coli* proteins (Figure 2 B). Additionally, the AA method resulted

in a minimal background in the unlabelled control test system (without AHA-labelled

proteins) with 10 ± 3 proteins (6 ± 3 *E. coli*, 1 ± 1 yeast).

Furthermore, the AA method allowed the detection of 48 proteins that were absent in the not

enriched samples (Supplementary Table 2). Of these, 29 were *E. coli* proteins that related to

cell metabolism, movement, transport, and protein biosynthesis. Additionally, 19 very low-

abundant yeast proteins were detected, likely co-eluting with AHA-labelled *E. coli* nP. This

co-elution did not occur in the AA method control, which confirmed the specificity of the

elution of labelled nP by DTT. Boiling of the beads with SDS-PAGE sample buffer released

no additional proteins showing that DTT was very efficient in elution (Supplementary Figures

3 and 4). Consequently, the selected elution with DTT is not only specific but also highly

efficient.

Due to the substrate shift from glucose to lactose, the enriched *E. coli* nP should include

proteins responsible for lactose degradation encoded by the *lac operon,* namely β-

galactosidase (*lacZ*), galactoside O-acetyltransferase (*lacA*), and β-galactoside permease

(*lacY*) [51, 52]. Indeed, the AA method successfully enriched β-galactosidase and galactoside

O-acetyltransferase by a relative foldchange of 3.7. However, β-galactoside permease was not

detected, neither in the enriched sample nor in the not enriched sample. As it is an integral

membrane protein it was probably lost during protein extraction [53].

Lactose consists of galactose and glucose. Therefore, the pathway for galactose metabolism

proteins should also be enriched with the AA method. The assignment of the detected nP to

the KEGG pathway for galactose metabolism showed that all enzymes for galactose

degradation were identified and at least 2-fold enriched, including β-galactosidase (EC

3.2.1.23; 217 ± 6 spectra), aldose 1-epimerase (EC 5.1.3.3; 14 ± 1 spectra), galactokinase (EC

2.7.1.6; 50 ± 5 spectra), galactose-1-phosphate uridylyltransferase (EC 2. 7.7.12; 13 ± 3

spectra), UTP-glucose-1-phosphate uridylyltransferase (EC 2.7.7.9; 3 ± 1 spectra), and UDP-

glucose 4-epimerase (EC 5.1.3.2; 34 ± 2 spectra) (Figure 3). Glycolytic enzymes were also

identified; however, they did not exhibit a 2-fold increase in the enriched nP compared to the

not enriched sample. This might be explained by the fact, that cells were already grown on

glucose as substrate before the substrate shift (Figure 3). Furthermore, all dehydrogenases

involved in the mixed acid fermentation of *E. coli* [54] were enriched after the substrate shift

by the AA method(Supplementary Table 2). The substrate switch was carried out by

centrifugation, which presumably led to anaerobic conditions for a short time and thus

facilitated the formation of these enzymes. This further emphasizes the significance of nP as a

conduit of knowledge regarding the activities of microorganisms.

<u>Enrichment of nP in anaerobic MC of an LBR using the AA method</u>

The next objective was to demonstrate that the established workflow is also effective for

weaker AHA-labelled and more complex MC. Therefore, the nP of an anaerobic MC from an

LBR were labelled with AHA while the substrate was switched from glucose to ethanol.

Initial quality control of successful AHA incorporation by SDS-PAGE revealed a weak

fluorescent signal that represents nP tagged with DBCO-Cyanin 5.5 by CC (Supplementary

Figure 5). The lower incorporation of AHA in the MC compared to a pure test sample

reinforces the necessity for specific and effective enrichment of nP to reliably detect nP in

MCs with MS. Unfortunately, in one out of three biological replicates of the experiment, no

fluorescence was detectable (Supplementary Figure 5). Therefore, this replicate was excluded

from further analysis. The other two replicates were processed with the AA method and

subsequent metaproteome analysis.

To achieve more accurate identification of proteins within the LBR MC, a metagenome was

sequenced from the LBR, and a corresponding protein database was generated. 76 MAGs

were binned.

In total, 2063 (replicate 1) and 2065 (replicate 2) proteins were identified in the not enriched

sample of the AHA-labelled MC ("Not enriched sample AHA"), while 525 (replicate 1) and

356 (replicate 2) proteins were identified after the enrichment of nP ("Elution AHA").

Application of the AA method to the non-labelled control ("Elution no AHA") showed a low

non-specific protein background of 7 (replicate 1) and 25 (replicate 2) proteins (Figure 4 A).

Notably, with the AA method, 26 proteins were identified (both replicates), which were absent

in the not enriched sample (Figure 4 B). Consequently, the specific enrichment of nP should

provide a better insight into the potential conversion of ethanol to methane in this LBR than

the proteome analysed without enrichment.

Ethanol-associated taxa based on nP enrichment

Initially, the enrichment of the nP was considered for the 20 most abundant MAGs of the AA

method and without the AA method each based on their protein abundance divided by the

MAG length (29 MAGs in total, Supplementary Table 6). The ratio of the MAGs between

the AA method and without the AA method was calculated based on their protein abundance.

The results of the two replicates demonstrate a consistent pattern of variation in protein

abundance among the MAGs under investigation. Bin_27 (family *Propionibacteriaceae*),

Bin_24 (class *Methanomicrobia*), Bin_41 (class *Clostridia*), and Bin 53 (species *Candidatus*

Methanoculleus thermohydrogenitrophicus) are of particular interest due to their high protein

abundance in all samples. The protein abundance of Bin_27 has been depleted by the AA

method, whereas the proteins of the other three MAGs are similarly abundant in all samples.

While certain MAGs, such as Bin_74 (family *Desulfobacteraceae*), Bin_62 (unknown

Bacteria), Bin_46 (unknown *Bacteria*), Bin_11 (family *Synergistaceae*), and Bin_20 (family

Alcaligenaceae), do not stand out due to their high protein abundance, they do exhibit strong

protein enrichment with the AA method. In contrast, certain MAGs, including Bin_7 (family

Pseudomonadaceae),   Bin_23   (family   *Methanobacteriaceae*),   Bin_69   (family

Flavobacteriaceae), Bin_71 (class *Clostridia*), Bin_75 (class *Actinomycetia*), and Bin_67

(family *Pseudomonadaceae*), exhibited a depletion pattern in their protein abundance after

using the AA method (Figure 5).

Upon closer examination of the protein functions from the enriched MAGs, a notable

abundance of stress proteins (up to 100% in some MAGs) was observed. Although ethanol

can have a positive effect on methanogenesis in AD [55], the addition of ethanol in

combination with the transfer of the MC from continuous LBR to serum bottles and mixing

with a medium may have induced the expression of these stress proteins [56, 57].

Consequently, stress proteins were excluded from further analyses to ensure a metabolic

enrichment of MAGs (Figure 5). In the case of the high-abundant MAGs, the protein

abundance after the removal of stress proteins has only decreased slightly from Bin_41.

However, in the case of the low-abundance MAGs, there have been some significant changes.

MAGs such as Bin_74 (family *Desulfobacteraceae*), Bin_46 (unknown *Bacteria*), and Bin_20

(genus *Castellaniella*) showed more than 10-fold protein enrichment using the AA method. In

contrast, proteins from MAGs such as Bin_13 (family: *Solibacteraceae*), Bin_34 (class

Betaproteobacteria), Bin_25 (species *Candidatus Brevifilum fermentans*) and Bin_62

(unknown *Bacteria*) could no longer be identified after the stress-induced proteins were

removed from the analysis. Therefore, the addition of ethanol resulted in increased expression

of metabolic proteins in Bin_74, Bin_46, and Bin_20, whereas MAGs such as Bin_13,

Bin_34, Bin_25, and Bin_62 primarily synthesized stress proteins.

Ethanol-associated functions based on nP enrichment

Upon closer examination of Bin_46, Bin_74, and Bin_20, it was discovered that Bin_74 and

Bin_20 expressed gluconeogenesis proteins. Too few proteins could be identified from

Bin_46 to make conclusions about its metabolism. A review of the individual genes revealed

that each of the three MAGs possesses at least two genes that encode for a potential alcohol

dehydrogenase (ADH). Additionally, *Castellaniella* (Bin_20) is known to utilize acetyl-CoA

from acetate, ethanol, or pyruvate *via* the glyoxylate cycle [58–62]. The incubation period for

these three strains in the presence of ethanol was likely insufficient for the strains to multiply

sufficiently to allow for the detection of these proteins. Without enrichment with the AA

method, they would thus not have been the focus of analyses. In follow-up studies, the

incubation times with ethanol could be extended so that also the metabolisms can be analysed

[63].

The analysis of the Functional Ontology Assignments for Metagenomes in comparison to a

sample with glucose (as in the LBR) showed that there were clear differences due to the

substrate shift (Supplementary Figure 6). A clustered heatmap of the Functional Ontology

Assignments for Metagenomes and most abundant MAGs showed that Bin_27, Bin_24,

Bin_41, and Bin_53 in particular account for most of the functions of the microbial

community of both replicates (Figure 6 and Supplementary Figure 7).

A detailed analysis of proteomes of all samples identified 29 different ADHs, with 7 ADHs

(Bin_41, 2x Bin_27, Bin_66, Bin_24, Bin_70, and Bin_53) as nP. Out of these 7 ADHs, 3

poorly classified ADHs (FHCFJDLK_121438 (Bin_41), FHCFJDLK_86872 (Bin_53), and

FHCFJDLK_294919 (Bin_70, only in replicate 1)) were increased ($\geq$ 2.5-fold) after

enrichment (Supplementary Table 3 and 4). An analysis with Alphafold [64, 65] and

KofamKOALA revealed that all three poorly classified ADHs are responsible for ethanol

degradation (see Supplementary Table 4). However, Bin_70 was a very low abundant MAG

(0.13%). This MAG belongs to the genus *Tepidiphilus*, which is known as a secondary

fermenter and syntrophic acetate oxidizer [66, 67]. The abundance of this MAG is however

insufficient to definitively ascertain its metabolic processes in this MC. Based on the proteins,

Bin_41 appears to be an acetate producer. All proteins for acetate formation from ethanol

were found in the proteome of Bin_41 [68], except an aldehyde dehydrogenase, which is

necessary for the further degradation of acetaldehyde [69–71]. It is therefore assumed that the

ADH of Bin_41 is bifunctional, converting ethanol into acetaldehyde (EC: 1.1.1.1) and then

directly to acetyl-CoA (EC: 1.2.1.10, Supplementary Figure 8) [72–75]. Bifunctionality poses

a challenge in the annotation of KO numbers, as normally only one KO number is assigned to

one protein. Note, that it was only through enrichment of nP that we became aware of the

protein and were able to consider the possibility of bifunctionality. Furthermore, NADH

dehydrogenases, which are crucial for replenishing depleted NAD+ levels during anaerobic

ethanol oxidation, were enriched from Bin_41 [76]. The removed electrons are probably

transferred to a membrane-bound cytochrome-coupled formate dehydrogenase, which utilizes

free CO_2 and protons for formate formation at iron-containing 2Fe-2S clusters [77].

Additionally, ATP appeared to be generated through substrate chain phosphorylation [77, 78] (

Supplementary Table 4). However, a major portion of stored energy is probably consumed for

the reduction of formate due to coupled proton influx [79].

Bin_53 (species *Candidatus Methanoculleus thermohydrogenitrophicus*) is closely related to

Methanoculleus, which is known to use ethanol and secondary alcohols as electron donors for

methanogenesis [80]. Ethanol is converted into acetate by a reaction that is coupled with

NADP+ reduction, which is then recovered by NADPH oxidoreductase (EC 1.5.1.40) [81, 82]

delivering reduction equivalents to hydrogenotrophic methanogenesis (Supplementary Figure

9). Since ethanol is not completely oxidized to CO_2, an alternative source of CO_2 is required

for hydrogenotrophic methanogenesis [83, 84]. A formate dehydrogenase (EC 1.17.1.9) was

identified that converts formate into CO_2, which can then be used for hydrogenotrophic

methanogenesis and feeds released reduction equivalents into hydrogenotrophic

methanogenesis. *Methanoculleus* is also known to use formate [84]. A syntropic interaction

between Bin_53 and Bin_41 is plausible with Bin_41 delivering formate to Bin_53 [85–87].

Furthermore, Bin_53 also should be able to consume H_2 which could result in the inhibition

of Bin_41 when accumulating in the medium [88–90]. But syntrophic transfer of H_2 probably

did not play a major role in the LBR MC since no hydrogenase capable of releasing H_2 was

detected in Bin_41, (Figure 7). The phenomenon of syntrophic ethanol oxidation between

fermenting bacteria and methanogenic archaea observed here has been repeatedly observed in

anaerobic digesters [28, 91–93].

In Bin_24, all proteins necessary for acetoclastic methanogenesis were identified, indicating

its potential to convert the acetate produced in Bin_41 and Bin_53 into methane and CO_2

(Supplementary Figure 10 and Figure 7). Therefore, based on the enriched nP, a syntrophic

relationship between these three MAGs is highly probable. In this scenario, ethanol is

converted to acetate by Bin_53 and Bin_41, which transfer formate to each other, and Bin_24

consumes the acetate. However, other sources of H_2, formate, acetate, and CO_2, e.g.

originating from fermentative degradation of microbial biomass [94], and additional partners

involved in ethanol oxidation (e.g. by minor abundant bins) cannot be excluded

(Supplementary Figures 11 and 12).

<u>Further Improvements and important considerations for implementing the workflow</u>

A successful BONCAT experimentis the precondition for this workflow. Our experiments

demonstrated the sufficient efficacy of BONCAT in slow-growing anaerobic MC. We advise

reading Hatzenpichler et al [22, 23, 95] and Landor et al. [27] as they address pivotal aspects

of a successful BONCAT experiment and the potential cytotoxicity of BONCAT. Following a

successful BONCAT, our workflow can be readily implemented in any laboratory (SOPs in

Supplementary Note 1). In the case of suboptimal nP yields despite a successful BONCAT

experiment, increasing the bead volume is recommended. The sensitivity and reproducibility

of quantitative measurements could be further improved by data-independent acquisition mass

spectrometry measurements, such as DIA-PASEF$^®$ [96]. In addition, fluorescence-activated

cell sorting of ncAA-labelled cells from MC [24] could generate corresponding metagenomes

for improved protein identification. Alternatively, the workflow could be adapted to HPG,

pyrrolysine, or β-ethynylserine, although this has not yet been tested but the same biotin

linker is also available for copper-based click chemistry [15, 97].

Conclusion

The combination of BONCAT and the direct enrichment of nP is an effective approach for

investigating MC using MS. Enriched nP reflects metabolic or adaptive processes, making it

easier to get new insights into the MC by highlighting relevant proteins. The optimized nP

enrichment workflow was demonstrated to be effective by a test system of *E. coli* and yeast

and may serve as a reference for future nP enrichments. The application of the workflow on a

slow-growing anaerobic MC enabled the description of syntrophic interaction in an MC

consuming ethanol as substrate.

Acknowledgment

We want to thank Dr.-Ing. Ute Thron from GETEC Green Energy GmbH for providing the

fermentation substrate for the inoculum of the LBR.

Funding

The bioinformatics support of the BMBF-funded project "Bielefeld-Gießen Center for

Microbial Bioinformatics" (BiGi) (BMBF grant FKZ 031A533) within the German Network

for Bioinformatics Infrastructure (de.NBI) is gratefully acknowledged. Funded by the

Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) – 446327964).

Figures

Figure 1: Workflow of the enrichment of newly synthesized proteins. **1.:** nP were labelled with AHA. **2.:** Free cysteines of the proteins were blocked

with IAA (not shown). The labelled nP were tagged with DBCO-S-S-PEG₃-biotin within 1 h. **3.:** The biotin-tagged nP were bound to magnetic

streptavidin beads (blocked with amino acids in advance). All proteins not bound to the beads were removed by harsh washing with urea, sodium

dodecyl sulfate (SDS), and acetonitrile (ACN). The bound proteins were eluted from the streptavidin beads with DTT. The specifically enriched nP

were transferred to a new tube, tryptic digested, and further analysed with LC-MS/MS.

Figure 2: Number of identified proteins of the test system using the new enrichment method

for nP. **A:** Comparison of the number of identified proteins from the test system between the

tested methods for nP enrichment. All proteins identified in the elution with AHA were

compared between the three enrichment methods: blocking with amino acids, blocking with

BSA, and no blocking. **B:** The different fractions of nP enrichment with the AA method were

compared. The number of proteins identified in the not enriched and AHA labelled sample

("Not enriched sample"), in the elution fraction of AHA labelled sample ("Elution AHA"),

and in the elution fraction without AHA labelling ("Elution No AHA") are shown. All proteins

with at least two spectra were considered. The proteins were grouped into the taxonomies *E.*

coli and yeast. Proteins with shared peptides were grouped as "Shared peptides". The bars

represent the average of each group, and the error bars represent the standard derivation of

three technical replicates. The raw data can be found in Supplementary Table 2.

GALACTOSE METABOLISM

Figure 3: KEGG map of all identified proteins from galactose metabolism in *E. coli* in the test system (map00052). **Orange** indicates proteins

enriched at least 2-fold with the developed AA method. **Blue** shows the proteins with no significant difference in abundance between the AA method

and not enriched sample. **Purple** represents all proteins uniquely identified in the not enriched sample. Boxes with several colours show similar

proteins with different abundances for this step. **Pink arrows** highlight the conversion pathway of galactose to α-D-glucose-1-phosphate. Blank

boxes were not identified in any sample. Sample data were normalized by dividing the number of spectra of each protein by the sum of the spectra

for each sample. The abundance of each protein was averaged over three replicates, and the relative abundance ratio between the not enriched

sample and elution was calculated for comparison [98] .

Figure 4: Number of identified metaproteins from AD using the AA method. The proteins were grouped based on shared peptides and the number of

the resulting metaproteins was plotted. Each sample was prepared in biological duplicates, which are indicated in the plot by red and blue bars. The

samples are the not enriched but AHA labelled sample ("Not enriched sample"), the enriched nPs of the AHA labelled sample in the elution fraction

("Elution AHA"), and in the elution fraction of the control without AHA labelling ("Elution No AHA"). **B:** The Venn diagram illustrates the

505 complete set of proteins identified in the respective replicates of the non-enriched ("Not enriched") and nP-enriched ("Enriched") samples [99]. The

506 raw data can be found in Supplementary Table 3.

Figure 5: Bubble plot of the MAGs with the highest protein abundance identified using the AA method. The protein abundance of the 20 most abundant MAGs identified by the AA method and the 20 most abundant MAGs identified in the non-enriched sample were compared in 2 replicates

(a total of 29 different MAGs). The fold changes were calculated based on the ratio of the relative spectral abundance of the MAG, which was

normalised by the MAG length in the samples. The taxonomies were identified using the TYGS. The left bubble plot of each replicate shows the

MAG abundance based on all identified proteins and the right without stress proteins. "C" in brackets shows the completeness of the bins. "+" in the

right bubble plot indicates that the MAG could no longer be identified. "*" are MAGs where the species are known: Bin_68 (*Candidatus*

Syntrophosphaera thermopropionivorans), Bin_67 (*Halopseudomonas salegens*), Bin_53 (*Candidatus Methanoculleus thermohydrogenitrophicus*),

Bin_5 (*Acidilutibacter cellobiosedens*), Bin_25 (*Candidatus Brevifilum fermentans*) and Bin_22 (*Fermentimonas caenicola*) (also Supplementary

Table   6).

Figure 6: Alternated biological processes in the MC due to ethanol addition The functional

ontology assignments for metagenomes of all enriched nP and MAGs with the AA method

were analysed using a heatmap created with R Studio(2023.06.0 Build 421) and pheatmap

(1.0.12). Previously, the protein abundances were normalised by dividing the spectral count of

each protein by the total spectral abundance of the sample. Afterwards, a pivot table was

generated, to sum up the relative abundances of the proteins based on the functional

assignment and the MAG. The functional ontology assignments were normalized using a z-

score per row (overall MAGs). also Supplementary Table 5.

Figure 7: Suggestion for an anaerobic syntrophic ethanol oxidation based on nP enrichment.

Intracellular metabolites are represented by unfilled circles, while extracellular metabolites

are represented by filled circles. Metabolic pathways are designated by the colours of the

arrows, which indicate the direction of metabolite reactions. Enzymes identified with the AA

method are denoted by boxes on the arrows. The boxes with thicker lines indicate that the

enzymes have been enriched at least twofold by the AA method. Exceptions include enzymes

with EC numbers 1.5.98.2 and 3.5.4.27 that were not identified using the AA method or in the

not enriched sample. The inspiration for this illustration is from Zhao et al. (2019) [100]

Fig.7.

Supplementary

Supplementary note 1: SOPs and detailed method description

Supplementary note 2: additional figures and tables

Supplementary table 1: lab scale biogas reactor information

Supplementary table 2: protein data testsystem

Supplementary table 3: metaprotein data ethanol

Supplementary table 4: identified proteins for syntrophic interactions

Supplementary table 5: biolog processes

Supplementary table 6: input bubble plot

Supplementary table 7: raw data protein quantification

Declarations

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

Data Availability Statement:

Proteome data were stored on PRIDE with the accession number: PXD047252, Username:
reviewer_pxd047252@ebi.ac.uk, Password: 0kWF7k9n

The metagenome database for protein identification is stored on ENA under the accession
number: PRJEB70937

Conflicts of Interest: Not applicable.

Author contributions (CRediT):

• Conceptualization: P.H., D.B.

• Project administration: P.H., D.B.

• Formal analysis: P.H., D.K., D.W.

• Funding acquisition: U.R., R.H.

• Investigation: P.H., D.K., D.W.

• Methodology: P.H., D.K., D.W., T.B., A.W.

• Project administration: D.B., U.R.

• Software: -

• Resources: D.B., U.R., R.H.

• Supervision: D.B., U.R.

• Validation: PH., D.K., D.W., T.B., A.W.

• Visualization: P.H., D.K.

References

1. Finlay BJ, Clarke KJ. Ubiquitous dispersal of microbial species. Nature. 1999;400:828.

doi:10.1038/23616.

2. Burman E, Bengtsson-Palme J. Microbial Community Interactions Are Sensitive to Small

Changes in Temperature. Front Microbiol. 2021;12:672910.

doi:10.3389/fmicb.2021.672910.

3. Suttle CA. The significance of viruses to mortality in aquatic microbial communities.

Microb Ecol. 1994;28:237–43. doi:10.1007/BF00166813.

4. Heyer R, Hellwig P, Maus I, Walke D, Schlüter A, Hassa J, et al. Breakdown of hardly

degradable carbohydrates (lignocellulose) in a two-stage anaerobic digestion plant is

favored in the main fermenter. Water Res. 2024;250:121020.

doi:10.1016/j.watres.2023.121020.

5. Waldrop MP, Firestone MK. Response of microbial community composition and function

to soil climate change. Microb Ecol. 2006;52:716–24. doi:10.1007/s00248-006-9103-3.

6. Reid G, Younes JA, van der Mei HC, Gloor GB, Knight R, Busscher HJ. Microbiota

restoration: natural and supplemented recovery of human microbial communities. Nat

Rev Microbiol. 2011;9:27–38. doi:10.1038/nrmicro2473.

7. Vrieze J de, Verstraete W. Perspectives for microbial community composition in

anaerobic digestion: from abundance and activity to connectivity. Environ Microbiol.

2016;18:2797–809. doi:10.1111/1462-2920.13437.

8. Tringe SG, Mering C von, Kobayashi A, Salamov AA, Chen K, Chang HW, et al.

Comparative metagenomics of microbial communities. Science. 2005;308:554–7.

doi:10.1126/science.1107851.

9. Wilmes P, Bond PL. Metaproteomics: studying functional gene expression in microbial

ecosystems. Trends Microbiol. 2006;14:92–7. doi:10.1016/j.tim.2005.12.006.

10. Biévre PD, Taylor PDP. Table of the isotopic compositions of the elements. International

Journal of Mass Spectrometry and Ion Processes. 1993;123:149–66. doi:10.1016/0168-

1176(93)87009-H.

11. Becker GW. Stable isotopic labeling of proteins for quantitative proteomic applications.

Briefings in Functional Genomics. 2008;7:371–82. doi:10.1093/bfgp/eln047.

12. Dieterich DC, Lee JJ, Link AJ, Graumann J, Tirrell DA, Schuman EM. Labeling,

detection and identification of newly synthesized proteomes with bioorthogonal non-

canonical amino-acid tagging. Nat Protoc. 2007;2:532–40. doi:10.1038/nprot.2007.52.

13. Kiick KL, Saxon E, Tirrell DA, Bertozzi CR. Incorporation of azides into recombinant

proteins for chemoselective modification by the Staudinger ligation. Proc Natl Acad Sci U

S A. 2002;99:19–24. doi:10.1073/pnas.012583299.

14. Dieterich DC, Link AJ, Graumann J, Tirrell DA, Schuman EM. Selective identification of

newly synthesized proteins in mammalian cells using bioorthogonal noncanonical amino

acid tagging (BONCAT). Proc Natl Acad Sci U S A. 2006;103:9482–7.

doi:10.1073/pnas.0601637103.

15. Ignacio BJ, Dijkstra J, Mora N, Slot EFJ, van Weijsten MJ, Storkebaum E, et al.

THRONCAT: metabolic labeling of newly synthesized proteins using a bioorthogonal

threonine analog. Nat Commun. 2023;14:3367. doi:10.1038/s41467-023-39063-7.

16. Kolb HC, Finn MG, Sharpless KB. Click Chemistry: Diverse Chemical Function from a

Few Good Reactions. Angew. Chem. Int. Ed. 2001;40:2004–21. doi:10.1002/1521-

3773(20010601)40:11<2004::AID-ANIE2004>3.0.CO;2-5.

17. Johnson JA, Lu YY, van Deventer JA, Tirrell DA. Residue-specific incorporation of non-

canonical amino acids into proteins: recent developments and applications. Curr Opin

Chem Biol. 2010;14:774–80. doi:10.1016/j.cbpa.2010.09.013.

18. van Kasteren S, Rozen DE. Using click chemistry to study microbial ecology and

evolution. ISME Commun. 2023;3:9. doi:10.1038/s43705-022-00205-5.

19. Bagert JD, Xie YJ, Sweredoski MJ, Qi Y, Hess S, Schuman EM, Tirrell DA. Quantitative,

time-resolved proteomic analysis by combining bioorthogonal noncanonical amino acid

tagging and pulsed stable isotope labeling by amino acids in cell culture. Mol Cell

Proteomics. 2014;13:1352–8. doi:10.1074/mcp.M113.031914.

20. Zhang G, Bowling H, Hom N, Kirshenbaum K, Klann E, Chao MV, Neubert TA. In-depth

quantitative proteomic analysis of de novo protein synthesis induced by brain-derived

neurotrophic factor. J Proteome Res. 2014;13:5707–14. doi:10.1021/pr5006982.

21. Franco M, D'haeseleer PM, Branda SS, Liou MJ, Haider Y, Segelke BW, El-Etr SH.

Proteomic Profiling of Burkholderia thailandensis During Host Infection Using Bio-

Orthogonal Noncanonical Amino Acid Tagging (BONCAT). Front Cell Infect Microbiol.

2018;8:370. doi:10.3389/fcimb.2018.00370.

22. Hatzenpichler R, Scheller S, Tavormina PL, Babin BM, Tirrell DA, Orphan VJ. In situ

visualization of newly synthesized proteins in environmental microbes using amino acid

tagging and click chemistry. Environ Microbiol. 2014;16:2568–90. doi:10.1111/1462-

2920.12436.

23. Hatzenpichler R, Krukenberg V, Spietz RL, Jay ZJ. Next-generation physiology

approaches to study microbiome function at single cell level. Nature Reviews

Microbiology. 2020;18:241–56. doi:10.1038/s41579-020-0323-1.

24. Reichart NJ, Jay ZJ, Krukenberg V, Parker AE, Spietz RL, Hatzenpichler R. Activity-

based cell sorting reveals responses of uncultured archaea and bacteria to substrate

amendment. ISME J. 2020;14:2851–61. doi:10.1038/s41396-020-00749-1.

25. Pasulka AL, Thamatrakoln K, Kopf SH, Guan Y, Poulos B, Moradian A, et al.

Interrogating marine virus-host interactions and elemental transfer with BONCAT and

nanoSIMS-based methods. Environ Microbiol. 2018;20:671–92. doi:10.1111/1462-

2920.13996.

26. Hellwig P, Dittrich A, Heyer R, Reichl U, Benndorf D. Detection, isolation and
characterisation of phage-host complexes using BONCAT and click chemistry; 2024.

27. Landor LAI, Bratbak G, Larsen A, Tjendra J, Våge S. Differential toxicity of
bioorthogonal non-canonical amino acids (BONCAT) in Escherichia coli. J Microbiol
Methods. 2023;206:106679. doi:10.1016/j.mimet.2023.106679.

28. Feng D, Guo X, Lin R, Xia A, Huang Y, Liao Q, et al. How can ethanol enhance direct
interspecies electron transfer in anaerobic digestion? Biotechnol Adv. 2021;52:107812.
doi:10.1016/j.biotechadv.2021.107812.

29. Wessel D, Flügge UI. A method for the quantitative recovery of protein in dilute solution
in the presence of detergents and lipids. Anal Biochem. 1984;138:141–3.
doi:10.1016/0003-2697(84)90782-6.

30. Heyer R, Schallert K, Büdel A, Zoun R, Dorl S, Behne A, et al. A Robust and Universal
Metaproteomics Workflow for Research Studies and Routine Diagnostics Within 24 h
Using Phenol Extraction, FASP Digest, and the MetaProteomeAnalyzer. Front Microbiol.
2019;10:1883. doi:10.3389/fmicb.2019.01883.

31. Laemmli UK. Cleavage of structural proteins during the assembly of the head of
bacteriophage T4. Nature. 1970;227:680–5. doi:10.1038/227680a0.

32. Maus I, Tubbesing T, Wibberg D, Heyer R, Hassa J, Tomazetto G, et al. The Role of
Petrimonas mucosa ING2-E5AT in Mesophilic Biogas Reactor Systems as Deduced from
Multiomics Analyses. Microorganisms 2020. doi:10.3390/microorganisms8122024.

33. Wibberg D, Andersson L, Tzelepis G, Rupp O, Blom J, Jelonek L, et al. Genome analysis
of the sugar beet pathogen Rhizoctonia solani AG2-2IIIB revealed high numbers in
secreted proteins and cell wall degrading enzymes. BMC Genomics. 2016;17:245.
doi:10.1186/s12864-016-2561-1.

34. Li D, Liu C-M, Luo R, Sadakane K, Lam T-W. MEGAHIT: an ultra-fast single-node

solution for large and complex metagenomics assembly via succinct de Bruijn graph.

Bioinformatics. 2015;31:1674–6. doi:10.1093/bioinformatics/btv033.

35. Hyatt D, Chen G-L, Locascio PF, Land ML, Larimer FW, Hauser LJ. Prodigal:

prokaryotic gene recognition and translation initiation site identification. BMC

Bioinformatics. 2010;11:119. doi:10.1186/1471-2105-11-119.

36. Nguyen TH, Zucker J-D. Enhancing Metagenome-based Disease Prediction by

Unsupervised Binning Approaches. In: 2019 11th International Conference on Knowledge

and Systems Engineering (KSE); 10/24/2019 - 10/26/2019; Da Nang, Vietnam: IEEE;

2019. p. 1–5. doi:10.1109/KSE.2019.8919295.

37. Berckx F, van Nguyen T, Bandong CM, Lin H-H, Yamanaka T, Katayama S, et al. A tale

of two lineages: how the strains of the earliest divergent symbiotic Frankia clade spread

over the world. BMC Genomics. 2022;23:602. doi:10.1186/s12864-022-08838-5.

38. Langmead B, Salzberg SL. Fast gapped-read alignment with Bowtie 2. Nat Methods.

2012;9:357–9. doi:10.1038/nmeth.1923.

39. Kang DD, Li F, Kirton E, Thomas A, Egan R, An H, Wang Z. MetaBAT 2: an adaptive

binning algorithm for robust and efficient genome reconstruction from metagenome

assemblies. PeerJ. 2019;7:e7359. doi:10.7717/peerj.7359.

40. Manni M, Berkeley MR, Seppey M, Zdobnov EM. BUSCO: Assessing Genomic Data

Quality and Beyond. Curr Protoc. 2021;1:e323. doi:10.1002/cpz1.323.

41. Meier-Kolthoff JP, Göker M. TYGS is an automated high-throughput platform for state-

of-the-art genome-based taxonomy. Nat Commun. 2019;10:2182. doi:10.1038/s41467-

019-10210-3.

42. Meier-Kolthoff JP, Carbasse JS, Peinado-Olarte RL, Göker M. TYGS and LPSN: a

database tandem for fast and reliable genome-based classification and nomenclature of

prokaryotes. Nucleic Acids Res. 2022;50:D801-D807. doi:10.1093/nar/gkab902.

43. Muth T, Behne A, Heyer R, Kohrs F, Benndorf D, Hoffmann M, et al. The

MetaProteomeAnalyzer: a powerful open-source software suite for metaproteomics data

analysis and interpretation. J Proteome Res. 2015;14:1557–65. doi:10.1021/pr501246w.

44. Schiebenhoefer H, Schallert K, Renard BY, Trappe K, Schmid E, Benndorf D, et al. A

complete and flexible workflow for metaproteomics data analysis based on

MetaProteomeAnalyzer and Prophane. Nat Protoc. 2020;15:3212–39.

doi:10.1038/s41596-020-0368-7.

45. Prestat E, David MM, Hultman J, Taş N, Lamendella R, Dvornik J, et al. FOAM

(Functional Ontology Assignments for Metagenomes): a Hidden Markov Model (HMM)

database with environmental focus. Nucleic Acids Res. 2014;42:e145-e145.

doi:10.1093/nar/gku702.

46. Aramaki T, Blanc-Mathieu R, Endo H, Ohkubo K, Kanehisa M, Goto S, Ogata H.

KofamKOALA: KEGG Ortholog assignment based on profile HMM and adaptive score

threshold. Bioinformatics. 2020;36:2251–2. doi:10.1093/bioinformatics/btz859.

47. Somasekharan SP, Stoynov N, Rotblat B, Leprivier G, Galpin JD, Ahern CA, et al.

Identification and quantification of newly synthesized proteins translationally regulated

by YB-1 using a novel Click–SILAC approach. Journal of Proteomics. 2012;77:e1-e10.

doi:10.1016/j.jprot.2012.08.019.

48. Kovalchuk SI, Anikanov NA, Ivanova OM, Ziganshin RH, Govorun VM. Bovine serum

albumin as a universal suppressor of non-specific peptide binding in vials prior to nano-

chromatography coupled mass-spectrometry analysis. Anal Chim Acta. 2015;893:57–64.

doi:10.1016/j.aca.2015.08.027.

49. Suelter CH, DeLuca M. How to prevent losses of protein by adsorption to glass and

plastic. Anal Biochem. 1983;135:112–9. doi:10.1016/0003-2697(83)90738-8.

50. Nicora CD, Anderson BJ, Callister SJ, Norbeck AD, Purvine SO, Jansson JK, et al.

Amino acid treatment enhances protein recovery from sediment and soils for

metaproteomic studies. Proteomics. 2013;13:2776–85. doi:10.1002/pmic.201300003.

51. Kennell D, Riezman H. Transcription and translation initiation frequencies of the

Escherichia coli lac operon. J Mol Biol. 1977;114:1–21. doi:10.1016/0022-

2836(77)90279-0.

52. Roderick SL. The lac operon galactoside acetyltransferase. C R Biol. 2005;328:568–75.

doi:10.1016/j.crvi.2005.03.005.

53. Zhao Y, Zhang W, Kho Y, Zhao Y. Proteomic analysis of integral plasma membrane

proteins. Anal Chem. 2004;76:1817–23. doi:10.1021/ac0354037.

54. Clark DP. The fermentation pathways of Escherichia coli. FEMS Microbiology Letters.

1989;63:223–34. doi:10.1111/j.1574-6968.1989.tb03398.x.

55. Jin Du, Qidong Yin, Mengqi Gu, Guangxue Wu. New insights into the effect of ethanol

and volatile fatty acids proportions on methanogenic activities and pathways.

Environmental Research. 2021;194:110644. doi:10.1016/j.envres.2020.110644.

56. Cao H, Du Wei, Yang Y, Shang Y, Li G, Zhou Y, et al. Systems-level understanding of

ethanol-induced stresses and adaptation in E. coli. Sci Rep. 2017;7:44150.

doi:10.1038/srep44150.

57. Guan N, Li J, Shin H-D, Du G, Chen J, Liu L. Microbial response to environmental

stresses: from fundamental mechanisms to practical applications. Appl Microbiol

Biotechnol. 2017;101:3991–4008. doi:10.1007/s00253-017-8264-y.

58. Kornberg HL, Krebs HA. Synthesis of cell constituents from C2-units by a modified

tricarboxylic acid cycle. Nature. 1957;179:988–91. doi:10.1038/179988a0.

59. Kornberg HL, Madsen NB. Synthesis of C4-dicarboxylic acids from acetate by a

glyoxylate bypass of the tricarboxylic acid cycle. Biochim Biophys Acta. 1957;24:651–3.

doi:10.1016/0006-3002(57)90268-8.

60. Dunn MF, Ramírez-Trujillo JA, Hernández-Lucas I. Major roles of isocitrate lyase and

malate synthase in bacterial and fungal pathogenesis. Microbiology (Reading).

2009;155:3166–75. doi:10.1099/mic.0.030858-0.

61. Foss S, Heyen U, Harder J. Alcaligenes defragrans sp. nov., description of four strains

isolated on alkenoic monoterpenes ((+)-menthene, alpha-pinene, 2-carene, and alpha-

phellandrene) and nitrate. Syst Appl Microbiol. 1998;21:237–44. doi:10.1016/S0723-

2020(98)80028-3.

62. Kämpfer P, Denger K, Cook AM, Lee S-T, Jäckel U, Denner EBM, Busse H-J.

Castellaniella gen. nov., to accommodate the phylogenetic lineage of Alcaligenes

defragrans, and proposal of Castellaniella defragrans gen. nov., comb. nov. and

Castellaniella denitrificans sp. nov. Int J Syst Evol Microbiol. 2006;56:815–9.

doi:10.1099/ijs.0.63989-0.

63. Koch S, Kohrs F, Lahmann P, Bissinger T, Wendschuh S, Benndorf D, et al. RedCom: A

strategy for reduced metabolic modeling of complex microbial communities and its

application for analyzing experimental datasets from anaerobic digestion. PLoS Comput

Biol. 2019;15:e1006759. doi:10.1371/journal.pcbi.1006759.

64. Jumper J, Evans R, Pritzel A, Green T, Figurnov M, Ronneberger O, et al. Highly

accurate protein structure prediction with AlphaFold. Nature. 2021;596:583–9.

doi:10.1038/s41586-021-03819-2.

65. Varadi M, Anyango S, Deshpande M, Nair S, Natassia C, Yordanova G, et al. AlphaFold

Protein Structure Database: massively expanding the structural coverage of protein-

sequence space with high-accuracy models. Nucleic Acids Res. 2022;50:D439-D444.

doi:10.1093/nar/gkab1061.

66. Tang Y-Q, Matsui T, Morimura S, Wu X-L, Kida K. Effect of temperature on microbial

community of a glucose-degrading methanogenic consortium under hyperthermophilic

chemostat cultivation. J Biosci Bioeng. 2008;106:180–7. doi:10.1263/jbb.106.180.

67. Poddar A, Lepcha RT, Das SK. Taxonomic study of the genus Tepidiphilus: transfer of

Petrobacter succinatimandens to the genus Tepidiphilus as Tepidiphilus succinatimandens

comb. nov., emended description of the genus Tepidiphilus and description of

Tepidiphilus thermophilus sp. nov., isolated from a terrestrial hot spring. Int J Syst Evol

Microbiol. 2014;64:228–35. doi:10.1099/ijs.0.056424-0.

68. Walter KA, Bennett GN, Papoutsakis ET. Molecular characterization of two Clostridium

acetobutylicum ATCC 824 butanol dehydrogenase isozyme genes. J Bacteriol.

1992;174:7149–58. doi:10.1128/jb.174.22.7149-7158.1992.

69. Kessler D, Leibrecht I, Knappe J. Pyruvate-formate-lyase-deactivase and acetyl-CoA

reductase activities of Escherichia coli reside on a polymeric protein particle encoded by

adhE. FEBS Lett. 1991;281:59–63. doi:10.1016/0014-5793(91)80358-a.

70. Paysan-Lafosse T, Blum M, Chuguransky S, Grego T, Pinto BL, Salazar GA, et al.

InterPro in 2022. Nucleic Acids Res. 2023;51:D418-D427. doi:10.1093/nar/gkac993.

71. Jones P, Binns D, Chang H-Y, Fraser M, Li W, McAnulla C, et al. InterProScan 5:

genome-scale protein function classification. Bioinformatics. 2014;30:1236–40.

doi:10.1093/bioinformatics/btu031.

72. Burton RM, Stadtman ER. The oxidation of acetaldehyde to acetyl coenzyme A. J Biol

Chem. 1953;202:873–90.

73. Fleischmann A, Darsow M, Degtyarenko K, Fleischmann W, Boyce S, Axelsen KB, et al.

IntEnz, the integrated relational enzyme database. Nucleic Acids Res. 2004;32:D434-7.

doi:10.1093/nar/gkh119.

74. Gómez-Manzo S, Escamilla JE, González-Valdez A, López-Velázquez G, Vanoye-Carlo

A, Marcial-Quino J, et al. The oxidative fermentation of ethanol in Gluconacetobacter

diazotrophicus is a two-step pathway catalyzed by a single enzyme: alcohol-aldehyde

Dehydrogenase (ADHa). Int J Mol Sci. 2015;16:1293–311. doi:10.3390/ijms16011293.

75. Hitschler L, Nissen LS, Kuntz M, Basen M. Alcohol dehydrogenases AdhE and AdhB

with broad substrate ranges are important enzymes for organic acid reduction in

Thermoanaerobacter sp. strain X514. Biotechnol Biofuels. 2021;14:187.

doi:10.1186/s13068-021-02038-1.

76. Ramos AR, Grein F, Oliveira GP, Venceslau SS, Keller KL, Wall JD, Pereira IAC. The

FlxABCD-HdrABC proteins correspond to a novel NADH dehydrogenase/heterodisulfide

reductase widespread in anaerobic bacteria and involved in ethanol metabolism in

Desulfovibrio vulgaris Hildenborough. Environ Microbiol. 2015;17:2288–305.

doi:10.1111/1462-2920.12689.

77. Wawrik B, Marks CR, Davidova IA, McInerney MJ, Pruitt S, Duncan KE, et al.

Methanogenic paraffin degradation proceeds via alkane addition to fumarate by

'Smithella' spp. mediated by a syntrophic coupling with hydrogenotrophic methanogens.

Environ Microbiol. 2016;18:2604–19. doi:10.1111/1462-2920.13374.

78. Zhang B, Lingga C, Bowman C, Hackmann TJ. A New Pathway for Forming Acetate and

Synthesizing ATP during Fermentation in Bacteria. Appl Environ Microbiol.

2021;87:e0295920. doi:10.1128/AEM.02959-20.

79. Schuchmann K, Chowdhury NP, Müller V. Complex Multimeric FeFe Hydrogenases:

Biochemistry, Physiology and New Opportunities for the Hydrogen Economy. Front

Microbiol. 2018;9:2911. doi:10.3389/fmicb.2018.02911.

80. Frimmer U, Widdel F. Oxidation of ethanol by methanogenic bacteria. Arch. Microbiol.

1989;152:479–83. doi:10.1007/BF00446933.

81. Song J, Wang J, Wang X, Zhao H, Hu T, Feng Z, et al. Improving the Acetic Acid

Fermentation of Acetobacter pasteurianus by Enhancing the Energy Metabolism. Front

Bioeng Biotechnol. 2022;10:815614. doi:10.3389/fbioe.2022.815614.

82. Berk H, Thauer RK. Function of coenzyme F420-dependent NADP reductase in

methanogenic archaea containing an NADP-dependent alcohol dehydrogenase. Arch.

Microbiol. 1997;168:396–402. doi:10.1007/s002030050514.

83. Maus I, Wibberg D, Stantscheff R, Eikmeyer F-G, Seffner A, Boelter J, et al. Complete

genome sequence of the hydrogenotrophic, methanogenic archaeon Methanoculleus

bourgensis strain MS2(T), Isolated from a sewage sludge digester. J Bacteriol.

2012;194:5487–8. doi:10.1128/jb.01292-12.

84. Kougias PG, Campanaro S, Treu L, Zhu X, Angelidaki I. A novel archaeal species

belonging to Methanoculleus genus identified via de-novo assembly and metagenomic

binning process in biogas reactors. Anaerobe. 2017;46:23–32.

doi:10.1016/j.anaerobe.2017.02.009.

85. Thiele JH, Zeikus JG. Control of Interspecies Electron Flow during Anaerobic Digestion:

Significance of Formate Transfer versus Hydrogen Transfer during Syntrophic

Methanogenesis in Flocs. Appl Environ Microbiol. 1988;54:20–9.

doi:10.1128/aem.54.1.20-29.1988.

86. Manzoor S, Schnürer A, Bongcam-Rudloff E, Müller B. Complete genome sequence of

Methanoculleus bourgensis strain MAB1, the syntrophic partner of mesophilic acetate-

oxidising bacteria (SAOB). Stand Genomic Sci. 2016;11:80. doi:10.1186/s40793-016-

0199-x.

87. Singh A, Schnürer A, Westerholm M. Enrichment and description of novel bacteria

performing syntrophic propionate oxidation at high ammonia level. Environ Microbiol.

2021;23:1620–37. doi:10.1111/1462-2920.15388.

88. Reddy CA, Bryant MP, Wolin MJ. Characteristics of S organism isolated from

Methanobacillus omelianskii. J Bacteriol. 1972;109:539–45. doi:10.1128/jb.109.2.539-

545.1972.

89. Bryant MP, Wolin EA, Wolin MJ, Wolfe RS. Methanobacillus omelianskii, a symbiotic

association of two species of bacteria. Arch Mikrobiol. 1967;59:20–31.

doi:10.1007/BF00406313.

90. Morris BEL, Henneberger R, Huber H, Moissl-Eichinger C. Microbial syntrophy:

interaction for the common good. FEMS Microbiol Rev. 2013;37:384–406.

doi:10.1111/1574-6976.12019.

91. Seitz H-J, Schink B, Pfennig N, Conrad R. Energetics of syntrophic ethanol oxidation in

defined chemostat cocultures. Arch. Microbiol. 1990;155:82–8.

doi:10.1007/BF00291279.

92. Summers ZM, Fogarty HE, Leang C, Franks AE, Malvankar NS, Lovley DR. Direct

exchange of electrons within aggregates of an evolved syntrophic coculture of anaerobic

bacteria. Science. 2010;330:1413–5. doi:10.1126/science.1196526.

93. Keller A, Schink B, Müller N. Alternative Pathways of Acetogenic Ethanol and Methanol

Degradation in the Thermophilic Anaerobe Thermacetogenium phaeum. Front Microbiol.

2019;10:423. doi:10.3389/fmicb.2019.00423.

94. Khesali Aghtaei H, Püttker S, Maus I, Heyer R, Huang L, Sczyrba A, et al. Adaptation of

a microbial community to demand-oriented biological methanation. Biotechnol Biofuels

Bioprod. 2022;15:125. doi:10.1186/s13068-022-02207-w.

95. Hatzenpichler R, Orphan VJ. Detection of Protein-Synthesizing Microorganisms in the

Environment via Bioorthogonal Noncanonical Amino Acid Tagging (BONCAT). In:

McGenity TJ, Timmis KN, Nogales B, editors. Hydrocarbon and Lipid Microbiology

Protocols. Berlin, Heidelberg: Springer Berlin Heidelberg; 2016. p. 145–157.

doi:10.1007/8623_2015_61.

96. Gómez-Varela D, Xian F, Grundtner S, Sondermann JR, Carta G, Schmidt M. Increasing

taxonomic and functional characterization of host-microbiome interactions by DIA-

PASEF metaproteomics. Front Microbiol. 2023;14:1258703.
doi:10.3389/fmicb.2023.1258703.

97. Fekner T, Li X, Lee MM, Chan MK. A pyrrolysine analogue for protein click chemistry.
Angew. Chem. Int. Ed. 2009;48:1633–5. doi:10.1002/anie.200805420.

98. Kanehisa M, Furumichi M, Sato Y, Kawashima M, Ishiguro-Watanabe M. KEGG for
taxonomy-based analysis of pathways and genomes. Nucleic Acids Res. 2023;51:D587-
D592. doi:10.1093/nar/gkac963.

99. Oliveros JC. Venny: An interactive tool for comparing lists with Venn's diagrams.; 2015.

100.   Zhao Z, Wang J, Li Y, Zhu T, Yu Q, Wang T, et al. Why do DIETers like drinking:
Metagenomic analysis for methane and energy metabolism during anaerobic digestion
with ethanol. Water Res. 2020;171:115425. doi:10.1016/j.watres.2019.115425.