1   The gut microbiome of honey-producing wasps has converged on that of social bees

Jo-anne C. Holley[1,2]*, Alexia N. Martin[1,3], Anna T. Pham[1], Jennifer Schlauch[1,4], Nancy A. Moran[2]

*Corresponding author: Jo-anne C. Holley

jo.holley@utexas.edu

1 Freshman Research Initiative, College of Natural Sciences, University of Texas at Austin, Austin, TX,

78712

2 Department of Integrative Biology, University of Texas at Austin, Austin, TX, 78712

3 Department of Entomology and Nematology, University of California Davis, CA, 95616

4 Department of Ecology and Evolutionary Biology, University of California Irvine, CA, 92697

Competing interests: The authors declare that they have no competing interests.

Key words: microbiome, honey wasps, paper wasps, honey bees, bumble bees, diet switch

ABSTRACT

Honey-feeding social bees, including honey bees and bumble bees, possess distinctive gut microbiomes

that provide benefits to hosts, such as defense against pathogens and parasites. Microbiome members

are transmitted through social interactions within colonies. The Mexican honey wasp (*Brachygastra*

mellifica) represents an independent origin of honey-storing within a group of social Hymenoptera. Honey

wasps feed on and store honey, but they prey on other insects as a protein source, and do not consume

pollen as do bees. We surveyed the gut microbiomes of Mexican honey wasps across sites within Texas

using 16S rRNA community profiling, and we estimated microbiome size using qPCR. For comparison, we

also surveyed non-honey feeding wasps from six families, collected in the same region. We found that

honey wasps have large microbiomes dominated by characteristic bacterial species, in contrast to other

wasps, which had smaller, more variable microbiomes dominated by environmental bacteria. All honey

wasps contained strains of *Bifidobacterium* and *Bombilactobacillus* that were closely related to symbionts

of bumble bees and other bees, suggesting their acquisition via host-switching. Some individuals also

harbored a close relative of *Candidatus* Schmidhempelia bombi (Orbaceae), an uncultured bumble bee

symbiont, again suggesting host-switching. The most prevalent species was an uncultured *Lactobacillus*,

related to *L. acetotolerans*, and potentially representing an independent acquisition of environmental

Lactobacillus. The switch to honey feeding, combined with a highly social life history, appears to have

facilitated the establishment of a microbiome resembling those of honey bee and bumble bees.

IMPORTANCE

Honey-feeding social insects such as honey bees and bumble bees have conserved gut

microbiomes that are transmitted among nestmates. The bacteria consist of host-associated species that

benefit the bee hosts by providing defense against pathogens and parasites, and potentially by

contributing to digestion. The microbiomes of wasps are less studied, although some species have bacteria

found in solitary bees. Here, we looked at a group of wasps that have a diet much like that of honey bees,

predicting that a diet switch from nectar to stored honey available year-round enabled them to acquire a

microbiome like that of honey-feeding bees. Honey wasps (*Brachygastra* spp.) are the only group of wasps

that store honey. Their large nests are made from paper, and colonies survive for several years. We found

that Mexican honey wasps have a conserved gut microbiome with similarities to those of other honey-

feeding bees. Several microbiome members are most closely related to bacteria in bumble bees,

suggesting their acquisition via host switching. The most common member was a previously unknown

Lactobacillus. Solitary wasps and other paper wasps that do not make honey have smaller microbiomes

dominated by environmental bacteria, suggesting that honey feeding has shaped the gut bacterial

communities of honey wasps.

INTRODUCTION

Gut microbial communities often provide numerous benefits to insect hosts (1). They can

protect against pathogens or parasites (2–4), enhance or become essential for physiological

development (5), metabolize indigestible substrates consumed by the host (6, 7) or neutralize toxic

compounds (8, 9). Depending on the host species, microbiome members can be highly host-specific or

can consist of bacteria widespread in the environment. Well-studied examples of host-restricted

microbiomes are found in the honey-feeding bees, which belong to a clade consisting of honey bees

(genus *Apis*), bumble bees (genus *Bombus*) and stingless bees (tribe Meliponini). These social bees are

associated with a microbiome that is relatively consistent among different species, reflecting a shared

evolutionary history (10). Most species harbor representatives of five bacterial lineages that appear to

have colonized a common ancestor 80 million years ago and to have subsequently diversified with hosts,

though a few bee lineages have lost ancestral associations (11). For example, in the best studied host,

the western honey bee (*A. mellifera*), adult guts are dominated by these five bacterial groups

(sometimes called the "core" microbiome), with a number of other "accessory" bacteria present more

erratically. These bacteria have been implicated in digestion and detoxification of the nectar, honey, and

pollen diet of their hosts (8, 12–14). Conversely, solitary bees (such as non-corbiculate bees) have more

variable gut microbiomes containing bacteria commonly associated with many bee lineages, flowers

(nectar and pollen), and other environmental sources (15–19). The gut microbiomes can vary with

pathogens (20) and urbanization (21, 22), although some of these bacteria may also protect against

pathogens (23).

Dietary shifts are frequently associated with a corresponding shift in gut microbial communities

(24, 25). A new diet may facilitate or necessitate the acquisition of new symbionts including symbionts

that can contribute to nutritional processing. Cockroach gut microbiomes reflect the transition to wood

feeding: omnivorous, litter-feeding cockroaches have similar gut bacteria regardless of diet composition

(26); however, the switch to wood-feeding in *Cryptocercus* cockroaches was accompanied by the

acquisition of lignocellulose-degrading protists (27). These symbionts are maintained in the wood-

feeding lower termites but were later lost in the soil and litter-feeding higher termites, which possess

polysaccharide-degrading fungal and bacterial symbionts that resemble the microbiomes of litter-

feeding cockroaches (7, 28–30). Conversely, on a shorter evolutionary timeline, the vulture bee, *Trigona*

necrophaga, adopted an obligately necrophagous diet rather than relying on pollen and honey-feeding

as in other stingless bees. This diet shift is linked to a higher abundance of environmentally acquired

acidophiles and more variable microbiomes compared to related honey-feeding bees (31).

Here, we investigate the gut microbiome of a honey-feeding wasp, representing a convergent

independent evolution of honey feeding in a group unrelated to honey bees. Three species of honey

wasps (*Brachygastra* spp.*,* Vespidae paper wasps) are known to uniquely store and feed on honey rather

than nectar (32, 33). This diet shift is rare among wasps; only *Polistes annularis* is known to store honey

to overwinter (34), although *Mischocyttarus* spp. store honey short-term (35). Mexican honey wasps,

Brachygastra mellifica, share other life history traits with honey bees: colonies contain thousands of

individuals, can live for several years, and new nests are established by swarming (33, 36). To address

how this dietary shift may have affected the microbiome composition of the Mexican honey wasp, we

characterized the gut microbiomes of wasps across their range in Texas (Fig. 1). We sequenced the V4

region of the 16S rRNA gene to determine the microbiome composition of 64 wasps from 13 nests.

Honey wasp microbiomes were compared to those of *Polistes* paper wasps and several other solitary

wasps. Our findings indicate that Mexican honey wasps acquired a unique microbial community

composed of bacteria associated with honey-feeding bees plus a novel species of *Lactobacillus*. This

composition differs greatly from that observed for other sympatric wasps, including paper wasp

relatives and solitary wasps, both of which have smaller and more variable gut microbiomes derived

from environmental sources.

RESULTS

Microbiomes of honey wasps

We sampled 64 Mexican honey wasps consisting of three to six workers sampled from 13 nests,

and five workers collected from flowers in Austin (Fig. 1). For comparison, 47 other wasps were

sampled, but only 26 of these wasp samples (13 *Polistes* paper wasps and 13 solitary wasps) had

adequate read counts to be included in the analysis (Table S1-S3, explanation of process in

Supplementary materials and methods). As controls, we included blank samples and two honey bee

samples. The blanks had no detectable product, and the honey bee samples were dominated by known

honey bee gut symbionts (Fig. 2, Table S3).

Honey wasps from all nests possessed *Bifidobacterium* and *Lactobacillus* (Fig. 2), both of which

are absent from the *Polistes* and solitary wasps. *Lactobacillus* ASV2 was present in four solitary wasps,

while eight other *Lactobacillus* ASVs (2, 7, 8, 24, 44, 56, 66, 67) and two *Bifidobacterium* ASVs were

confined to honey wasps. Honey wasps also possessed *Convivina* (ASV16 and ASV25), *Fructilactobacillus*

(ASV13), and *Bombilactobacillus* (ASV19) throughout the Texas range, but these were not present in all

individuals (Fig. 2B). Honey wasps from some San Antonio nests had bacteria related to *Ca.*

Schmidhempelia bombi (ASV14 and ASV28), an undescribed Neisseriaceae (ASV27), and *Asaia* (ASV34).

Based on the "core" microbiome analysis, some ASVs of *Lactobacillus*, *Bifidobacterium*,

Bombilactobacilus, *Fructilactobacillus*, and *Convivina* occur in most honey wasp individuals and in high

abundance (Fig. S1). A NMDS ordination of the Bray-Curtis dissimilarity index shows that honey wasps

cluster together apart from the other wasps and honey bees studied (Fig. 3A, Table S4, PERMANOVA, F =

9.334, p = 0.001). Pairwise comparisons further support this finding (Table S5 honey wasps:other wasps,

F = 15.366, adjusted p = 0.003; honey wasps:honey bees, F = 4.458, adjusted p = 0.006). Beta dispersions

significantly differed between the groups (Table S6, betadisper, F = 16.472, p = 0.001).

Microbiomes of *Polistes*, solitary wasps

The microbiomes of *Polistes* and solitary wasps sometimes contained *Bifidobacterium* or

Lactobacillus, but these ASVs differed from those found in honey wasps (Fig. 2B). *Polistes* and solitary

wasps shared Enterobacterales ASVs (ASV4 and ASV6) with honey wasps, and they more often

contained *Lactococcus*, *Gluconobacter*, *Gilliamella*, *Acinetobacter*, *Pseudomonas,* and *Atopobiaceae* (Fig.

2). Core microbiome analysis showed that *Polistes* were typified by the presence of Enterobacterales,

Lactococcus, *Acinetobacter*, and *Exiguobacterium* (Fig. S1). No bacteria were detectable in more than

40% of *Polistes* individuals sampled, indicating that *Polistes* microbial communities are often absent or

tiny. Several *Polistes* ASVs were shared with the 13 solitary wasps (Enterobacterales ASV4 and

Acinetobacter ASV31), but *Gilliamella* (ASV 32) and another Enterobacterales (ASV6) were more

prevalent in *Polistes* samples. Overall, the prevalence of bacterial ASVs was lower in the individuals of

other wasp species compared to the honey wasps.

Absolutes sizes of honey wasp microbiomes

Based on qPCR of a region of the bacterial 16S rRNA gene, honey wasps have substantial gut

communities, averaging greater than 10^8 gene copies per bee individual, with relatively little variability

in community size (Fig. 4). In contrast, the quantity of gut bacteria in *Polistes* and in solitary wasps was

lower and highly variable, ranging from below the limit of detection to 10^{10} copies of the bacterial 16S

rRNA gene (Fig. 4, Table S7). *Polistes* wasps had a median of 3.1×10^5 copies of 16S rRNA gene, ranging

from 10^3 to 10^7 copies, and solitary wasps had a similar wide range of bacterial load (Table S8, Dunn's

test, $z = -1.662$, adjusted $p = 0.290$).

Microbiome diversity among honey wasp nests

Honey wasp microbiome composition varied among sample locations and varied significantly

among nests (Fig. 3B, Table S4-S5, PERMANOVA, $F = 3.198$, $p = 0.001$; betadisper, $F = 0.973$, $p = 0.498$).

No nests had distinctly different microbiomes based on pairwise comparisons of honey wasps from all

13 nests (Table S6, adjusted for multiple comparisons). The primary members of the microbiome,

Bifidobacterium and *Lactobacillus*, were present in all nests, but there were multiple ASVs for each that

were differentially distributed among nests. Honey wasp *Bifidobacterium* was represented by three

ASVs. Indicator species analysis revealed that ASV1 was significantly associated with five nests while

ASV3 was associated with four different nests (Table S9). *Bifidobacterium* ASV1 and ASV3 rarely

cooccurred in the same host*,* but ASV3 typically occurred with *Bifidobacterium* ASV5 (Fig. 2B).

Lactobacillus ASV2 was present in all nests, but the indicator species analysis associated ASV2 with five

particular nests (Table S9, indicator value = 0.548, $p = 0.022$). Other *Lactobacillus* ASVs were associated

with a subset of nests. For example, *Lactobacillus* ASV24 was associated with San Antonio nest 6 and

ASV44 with the Port Lavaca nest (Table S9; ASV24: indicator value = 0.814, p = 0.001; ASV44: indicator

value = 0.861, p = 0.001). Some bacteria had a limited association with only a few nests, for example,

San Antonio nests 1, 2, and 3 were characterized by the presence of *Ca.* Schmidhempelia (Table S9;

ASV14: indicator value = 0.634, p = 0.007; ASV28: indicator value = 0.649, p = 0.004). Additionally,

Wolbachia and *Lactococcus* were present in the single nests collected from Kingsville and Port Lavaca,

while the Austin nest showed high abundance of *Fructilactobacillus* (Fig. 2B, Table S3).

Laboratory isolation of honey wasp bacteria

170 We isolated pure cultures of *Ca.* Schmidhempelia (wjB12) and several strains of *Bifidobacterium*

171 (efB7, efB4, wjB1) from two San Antonio nests (SA1 and SA7) grown on heart infusion agar (HIA) with 5%

172 sheep's blood under 5% CO_2 at 35°C. A *Fructilactobacillus vespulae* isolate (sjhB6) and additional

173 *Bifidobacterium* isolate (sjhB46) were also cultured in MRS broth under anaerobic conditions at 31°C.

Phylogenetic relationships of key honey wasp bacteria

We constructed phylogenies of three bacterial groups prevalent in honey wasp microbiomes

belonging to Orbaceae, *Bifidobacteria*, and Lactobacillaceae. Within Orbaceae, the honey wasps

possessed two ASVs and a bacterial isolate (strain wjB12) that form a well-supported clade together

with *Candidatus* Schmidhempelia bombi, an uncultured Orbaceae known only from bumble bees (Fig.

5A).

Three *Bifidobacterium* ASVs and numerous *Bifidobacterium* isolates from honey wasps also form

a clade with multiple species of *Bifidobacterium* isolated from several species European bumble bees (*B.*

commune, *B. bohemicum*, *B. bombi*) (Fig. 5B). This clade is not closely related to the *Bifidobacterium*

'asteroides' clade associated with honey bees, bumble bees, carpenter bees, and *Osmia* mason bees.

Honey wasps also possess a variety of bacteria in the family Lactobacillaceae. Eight ASVs

represent an undescribed species of *Lactobacillus* that is closely related to the clade of *Lactobacillus*

associated with honey bees, bumble bees, and *Osmia* bees (formerly referred to as "Firm-5") (Fig. 5C). It

is also closely related to *Lactobacillus acetotolerans*, an acetic-acid tolerant bacterium isolated from

vinegar. This bacterium remains uncultured, and the 16S rRNA gene amplicon variation suggests there is

strain diversity within the honey wasps with ASV2 found in 92% of honey wasps sampled.

Bombilactobacillus (formerly "Firm-4") was sequenced from multiple honey wasps but remains

uncultured (Fig. 5C). Previously the genus was known only from honey bees, bumble bees, and

carpenter bees. The sequences from honey wasps were close to *Bombilactobacillus apium* isolated from

the Asian honey bee, *Apis cerana*. Additionally, honey wasps possess the vespid wasp-associated

Fructilactobacillus vespulae, a *Lactiplantilactobacillus*, and another Lactobacillaceae (Fig. 5C).

DISCUSSION

Honey wasps (genus *Brachygastra*) represent an independent evolution of honey storage and

feeding in a eusocial member of the Hymenoptera, and have also acquired a gut microbiome very

similar to that of social bees. Honey wasps are not closely related to honey-feeding bee lineages, yet

most members of the honey wasp microbiome are related to bacteria known from bumble and honey

bees (Fig. 6). These include *Bifidobacterium* and *Ca.* Schmidhempelia (both related to species solely

known from bumble bees), *Bombilactobacillus*, and *Convivina*, though these were not consistently found

in workers from all nests (Fig. 2A). Practically all honey wasps in this study possessed a novel

Lactobacillus in high abundance that is related to other bee-associated *Lactobacillus* species but not

known from bumble bees and honey bees. *Fructilactobacillus vespulae* was also present in some

colonies, showing they also possess bacteria associated with paper wasps from the subfamily Vespinae.

The high microbial abundance (Fig. 4) is comparable to that observed for honey bees, averaging slightly

lower, which may reflect the smaller body size of honey wasps (10) (Fig. 1A). The honey wasp

microbiome differs from that of other wasps (Fig. 3A); the *Polistes* paper wasps and solitary wasps had

microbiomes with lower abundance variable composition, and dominated by environmentally acquired

bacteria; they lacked all core members of the honey wasp microbiome. Some environmental bacteria

were shared among honey wasps and *Polistes* and solitary wasps, including Enterobacteriacae,

Pseudomonas, *Lactococcus*, and some others found in environmental sources such as nectar and in

other pollinating Hymenoptera (19, 37–39).

Honey wasps have a diverse consortium of Lactobacillaceae that is distinct from that of honey

bees or the other wasps included in this study. All honey wasps host an uncultured *Lactobacillus* which

belongs to a clade of honey bee-associated *Lactobacillus* (formally Firm-5); multiple ASVs were detected

indicating strain diversity in this taxon (Fig. 5C). *Bombilactobacillus* ASVs in honey wasps are closely

related to *B. apium* isolated from *Apis cerana* (40). Honey wasps also consistently possess three related

Bifidobacterium that belong to the same clade as *Bifidobacterium commune*, *Bifidobacterium bombi*,

and *Bifidobacterium bohemicum* (Fig. 5B), all isolated from European bumble bees (41, 42). These three

Bifidobacterium ASVs were found in high abundance in all honey wasp nests. However, individual wasps

had either ASV1 or ASV3 in high abundance, not both, suggesting competition or another

incompatibility. Several nests of honey wasps in San Antonio had *Candidatus* Schmidhempelia bombi,

known from metagenomic samples of *Bombus impatiens* (Fig. 5A) (10, 43). *Ca.* Schmidhempelia is

abundant in the colonization phase of *B. impatiens* but decreases as bees age (44). These findings

suggest that the core species in guts of Mexican honey wasps were acquired from social bees some time

in the past. Honey wasps and bumble bees may have co-occurred in North America for at least several

million years based on dated phylogenies of the Vespids (45) and *Bombus* (46) providing ample time for

the lineages to interact. Notably, however, several honey wasp bacteria are more similar to honey bee

and bumble bee species located in Asia and Europe. This could be attributed to long-term patterns of

symbiont acquisition and loss or an incomplete knowledge of symbiont distributions.

Also present in the honey wasps were bacteria commonly associated with other bees. These

include *Lactococcus, Convivina*, and some *Lactobacillaceae* such as *Lactiplantibacillus*. Three honey

wasps collected individually from Austin had high abundances of a bacterium that may be a member of

Candidatus Stammerula. The short ASV aligns with 16S sequences in the NCBI database with 99%

sequence identity to an uncultured symbiont of tephritid flies (47, 48).

This study adds to others indicating that vespid wasps have some reliable associations with gut

bacteria. The subfamily Vespinae contains four genera, and two have been the focus of microbiome

research. *Vespula pensylvanica*, *Vespula vulgaris* and *Vespa velutina* are all associated with several

bacteria that were found in the Mexican honey wasp. *Fructilactobacillus vespulae* was found in most

honey wasp nests and highly abundant in one nest. This bacterium was isolated from *Vespula vulgaris* in

China, and sequenced from *Vespula pensylvanica* in the US (37) and *Vespa velutina* in Belgium, showing

an association with vespid wasps on three continents. *Convivina was* detected in about 40% of the

honey wasps representing all sampled nests, but not in other wasps. *Convivina intestini* was isolated

from *Bombus terrestris* (49) and *Convivina praedatoris* was found in all individuals in a single *V. velutina*

nest collected in Belgium (outside of the Korean native range of *V. velutina*) (50). Similarly,

Lactiplantibacillus was found in about 43% of the honey wasps, and is another possible member of the

V. velutina microbiome. A study of five *Vespa* species in the South Korean native range did not find

Convivina and *Lactiplantibacillus* among the top 10 bacterial genera (51), so these associations may be

site-specific.

Polistes wasps did not share many bacteria with honey wasps and were associated with

Enterobacteriaceae and *Gilliamella*, *Acinetobacter*, *Gluconobacter,* and *Lactobacillus* (Fig. 2A). These

bacteria were found in the four species of *Polistes* inconsistently and are typically associated with flower

nectar and nectar-feeding solitary bees (15, 52, 53), and they have been found in *Vespa* spp. (51) and

Vespula pensylvanica (37). *Gilliamella* sp. (ASV 32) was found in the mutilid, crabronid, sphecid, and

several species of *Polistes*. *Gilliamella* is one of the core bacteria in honey bees and bumble bees (39)

but some strains have been detected in native bee species and in nectar (52). The amount of gut

bacteria based on 16S rRNA gene copies is lower and more variable in *Polistes* and solitary wasps

compared to honey wasps. It is unknown why this is, but could be due to conditions of the wasp gut

which may make it hard for microbes to colonize or because not all animals require a gut

microbiome (1, 54).

Flowers are an important hub for the transmission of microbes (15), and bacteria are present in

the nectar and pollen (16). The shared food source of nectar may have enabled honey wasps to acquire

bacteria from honey-feeding bees. *Polistes* workers also feed on nectar and share it with larvae, as do

many solitary wasps, explaining how they acquire some bacteria. However, many adult bees have

microbiomes similar to that of their pollen provisions (18, 19, 21, 22, 52, 55), whereas *Polistes* derive

protein from insect prey. Overall, it appears that the studied *Polistes* and solitary wasp species contain

variable environmentally acquired bacteria with considerable variation among individuals.

We note that determining microbiome composition from PCR amplicons of extracted DNA

cannot distinguish between DNA from bacteria colonizing the guts, DNA from bacteria that are simply

present in the food, and extracellular DNA from dead bacteria. The possibility that profiles represent

these other sources is especially great for microbiomes consisting of environmental bacteria, such as

those in the *Polistes* and solitary wasps. Since many bees and wasps can visit the same flowers, where

they may contaminate nectar, small numbers of live or dead cells from other host species may be

ingested in food and stored food provisions.

Bumble bees or honey bees appear to have contributed some bacterial lineages to honey wasps,

but numerous others were not transferred. Mexican honey wasps do not possess members of the

Bifidobacterium "asteroides", *Snodgrassella,* or *Gilliamella* clades, which are found in bumble bees and

honey bees (10, 56). Honey wasps also lack the "Firm-5" *Lactobacillus* species associated with honey

bees but possess an undescribed relative. *Snodgrassella* and *Gilliamella* form a biofilm along the ileum

of honey bees and bumble bees (57, 58). Their absence is not surprising given that they have been lost

by *Melipona* stingless bees (11) and that honey wasps do not have a defined ileum gut structure (Fig.

1A). Furthermore, multiple *Gilliamella spp.* synthesize pectin-degrading enzymes for the breakdown of

cell walls including pollen (12, 13). Mexican honey wasps source protein from insect prey, so pollen-

degrading bacteria would be unnecessary. Also missing are the "accessory" bacteria in the honey bee

microbiome which include species in the genera *Apilactobacillus*, *Bombella*, *Arsenophonus*, *Bartonella*,

Parasaccharabacter, *Apibacter*, and *Frischella perrara* (39, 59).

Social and gregarious insects can facilitate the transmission of symbionts among group members

by direct contact (60). Social spiders, for example, acquire their bacteria from co-nesting females via

regurgitation resulting in homogenized microbiomes among members of the same nest (61). Honey

wasp nests can house up to 20,000 individuals, share food sources through trophallaxis, and establish

new nests by swarming (33, 36), providing frequent opportunities for symbiont transfer among

nestmates and to new nests. Sociality, however, is not essential for maintaining a conserved, host-

restricted microbiome. Brood provisions also serve as a microbial transmission route for solitary bees

such as *Osmia* Mason bees (18, 62), and *Xylocopa* Carpenter bees have conserved microbiomes with

only moderate social interactions within communal nests (38, 63, 64). The gut microbiomes of many

solitary bees are acquired from environmental sources (pollen and nectar) and brood provisions and as

such are heterogeneous (19). *Polistes* paper wasps are eusocial and have a means of bacterial

transmission among nestmates and generations, yet their gut microbiomes are composed of

environmental bacteria found in nectar and in solitary bees. It appears that solitary wasps acquire

bacteria from environmental sources based on the wasps sequenced in this study.

Mexican honey wasps occur from central Texas south to Panama, so our collection range

represents a small part of the *B. mellifica* range. Furthermore, the Mexican honey wasp was restricted to

the USA/Mexico border until around 1995 (33) and recently expanded approximately 500 kilometers

north over 3 years. We collected 13 nests across the Texan range and found significant variation in

microbiome composition between nests. Individual worker microbiomes are more similar to those of

nestmates. Differences between nests are due to the presence and abundance of particular ASVs,

indicating strain variation among the core microbiome members, such as the uncultured *Lactobacillus*,

with eight ASVs that form a clade (fig. 5c), and the *Bifidobacterium* with 4 ASVs that varied among hosts.

All individuals in all nests had a microbiome dominated by *Lactobacillus* and *Bifidobacterium*

with the exception of the Austin nest, where three of the five individuals were dominated by

Fructilactobacillus vespulae. Additionally, five individual honey wasps were collected from flowers (not

nests) in Austin. One had the characteristic microbiome profile, and the four other wasps had higher

abundance of Enterobacteriaceae, *Ca*. Stammerula, and *Conviva*. These two Austin collection sites sit

close to the northernmost extent of the Mexican honey wasp range, suggesting a change in microbial

diversity associated with edge populations, though our sampling is not sufficient for conclusions. Two

species of English butterflies have microbiomes with decreased diversity and a shift in composition in

the northern part of their range. (65) Similarly, the microbiomes of range-expanding plants were more

similar to each other than to those of conspecifics (66). Further research into honey wasps throughout

Mexico and Central America would reveal whether the core microbiome is stable throughout its range.

The transition to long-term honey-feeding and storage in a wasp is only known in three species

of *Brachygastra* honey wasps (*B. mellifica*, *B. azteca*, and *B. lecheguana*) (32, 33). This diet switch may

have facilitated the acquisition of bee-associated bacteria as honey wasps feed on a carbohydrate-rich

diet similar to that of honey-feeding bees. However, the carbohydrate profile of honey wasp honey

differs from that of honey bee honey, which may influence the composition of gut bacteria. Honey bee

honey is predominantly glucose and fructose whereas, in a study based on seven nests, honey wasp

honey contains less glucose and fructose (23% and 30% respectively), more sucrose (~9.8%), with 28.6%

of the sugars not identified (33).

Honey wasps routinely collect honeydew from aphids and psyllids (33) which contains large

amounts of melezitose and trehalose (67). Honey bees can also collect honeydew when nectar is not

available (68), studies in Greece showed that honey can have up to 14% melezitose when honey bees

forage directly on honeydew from aphids on fir trees (67, 69). Gut bacteria of honey bees can

metabolize a variety of sugars, including some monosaccharides toxic to honey bees (8). *Bifidobacterium*

spp. have extensive abilities to degrade plant-derived polysaccharides (70), and bee-associated

Bifidobacterium can degrade and use hemicelluloses (13). Honey wasps likely do not ingest significant

amounts of polysaccharides, since they do not eat pollen, so metabolic capabilities of their microbiomes

may be very different from those of honey bees and bumble bees, which rely on pollen. Investigation of

the functional capabilities of the *Bifidobacterium* strains isolated from honey wasps. will offer insights

into their metabolic capabilities and potential benefits to hosts.

Overall, the Mexican honey wasp gut microbiome differs greatly from microbiomes of related

wasps, in both size and composition. It is an amalgamation of bacteria that appear to have originated

primarily from bumble bees or honeybees, and from vespid wasps. The switch from feeding on nectar

and insect prey to a stable, year-long supply of stored honey may have facilitated (or been facilitated by)

the acquisition of bacteria associated with honey-feeding bees.

METHODS

345   METHODS

Sample Collection

Sample Collection

Honey wasps from 13 nests, individual honey wasps, *Polistes* wasps, solitary wasps, and honey bees

were collected across sites in central and south Texas (Fig. 1, Table S1). Guts were dissected from

specimens using sterile technique and stored in 100% ethanol at -80°C. Additional honey wasp guts

were preserved in 20% glycerol at -80°C for bacterial isolation.

DNA Extractions, Sequencing, and qPCR

DNA was extracted from whole guts using the CTAB/phenol-chloroform extraction method (71). The full

length or V4 region of the 16S rRNA gene was PCR amplified and sequenced on an Illumina MiSeq

platform at the Genome Sequencing and Analysis Facility (University of Texas at Austin). Quantitative

PCR was performed on the DNA extracts to quantify total 16S rRNA gene copies as an estimate of

bacterial abundance.

Bacterial Isolation and Identification

The preserved honey wasp gut homogenates from two nests (SA1 and SA7) and an individual collected

in Austin were grown on HIA + 5% sheep's blood or in MRS broth at 35°C in 5% CO_2 or 31°C in anaerobic

conditions for 1-3 days. To identify bacterial isolates, the 16S rRNA gene was PCR amplified, and

amplicons were sequenced at the DNA Sequencing Facility (University of Texas at Austin).

Data Analysis and Phylogenetic Reconstruction

Illumina sequencing files were processed in R using the dada2 pipeline. Taxonomy was assigned using

the SILVA 138.1 database (72) and manually confirmed using NCBI BLASTn. Read counts were checked

after each filtering step, resulting in the removal of 21 *Polistes* and solitary wasps from the dataset (see

Supplementary methods, Table S2). The remaining samples were rarefied to 90% of the minimum

sampling depth (Fig. S2, Table S3). The core microbiomes of honey wasps, *Polistes*, and solitary wasps

were identified using the microbiome package. Indicator species analysis was also performed using the

indicspecies package to determine if any ASVs were significantly associated with each honey wasp nest.

Statistical differences between gut bacterial communities across honey wasp nests and insect types

were analyzed using PERMANOVA (*adonis2*), and assumptions were assessed using beta dispersion

analysis. Post-hoc pairwise comparisons with Bonferroni adjustment were performed to directly

compare bacterial communities across nests and insect types.

The 16S rRNA gene phylogenies were reconstructed for Lactobacillaceae, Orbaceae, and

Bifidobacterium, which were commonly amplified ASVs in the honey wasps and isolated bacteria. Full

length 16S rRNA gene sequences were obtained from GenBank. These were aligned using MAFFT with

ASV sequences and full length 16S rRNA from the isolated bacteria. Maximum likelihood phylogenies

were reconstructed using IQtree with auto-generated substitution models and 100 standard bootstrap

alignments.

DATA AVAILABILITY

Raw 16*S* rRNA gene amplicon reads are deposited in the National Center for Biotechnology

Information Sequence Read Archive under accession PRJNA932820. ASV sequences, taxonomic

assignments, read numbers, metadata can be found in the supplement. Scripts for the analysis can be

found on GitHub (https://github.com/annatpham/honey-wasp-microbiome), and insect specimens are

deposited at the University of Texas Insect Collection (UTIC).

ACKNOWLEDGEMENTS

We thank G. Rankin, H. Koch, S. Bard, D. Fall, S. Biles, J. Hare, M. Reed, B. Freitag, B. Fehrenkamp

and for specimen acquisition and collection. Thanks also to A. Santillana for images, A. Wild for

specimen ID, and E. Frederick and K. Chow for technical assistance. This work was supported by USDA-

NIFA grant (2017-06473), NSF award (NSF 2103208), and the Texas EcoLab program, and

undergraduates were supported by the Freshman Research Initiative (UT Austin).

CITATIONS

1. Moran NA, Ochman H, Hammer TJ. 2019. Evolutionary and ecological consequences of gut microbial communities. Annu Rev Ecol Evol Syst 50:451–475.

2. Koch H, Schmid-Hempel P. 2012. Gut microbiota instead of host genotype drive the specificity in the interaction of a natural host-parasite system. Ecol Lett 15:1095–1103.

3. Lang H, Duan H, Wang J, Zhang W, Guo J, Zhang X, Hu X, Zheng H. 2022. Specific strains of honeybee gut *Lactobacillus* stimulate host immune system to protect against pathogenic Hafnia alvei. Microbiol Spectr 10:e0189621.

4. Steele MI, Motta EVS, Gattu T, Martinez D, Moran NA. 2021. The gut microbiota protects bees from invasion by a bacterial pathogen. Microbiol Spectr 9:e00394–21.

5. Shukla SP, Plata C, Reichelt M, Steiger S, Heckel DG, Kaltenpoth M, Vilcinskas A, Vogel H. 2018. Microbiome-assisted carrion preservation aids larval development in a burying beetle. Proc Natl Acad Sci U S A 115:11274–11279.

6. Salem H, Bauer E, Kirsch R, Berasategui A, Cripps M, Weiss B, Koga R, Fukumori K, Vogel H, Fukatsu T, Kaltenpoth M. 2017. Drastic genome reduction in an herbivore's pectinolytic symbiont. Cell 171:1520–1531.e13.

7. Arora J, Kinjo Y, Šobotník J, Buček A, Clitheroe C, Stiblik P, Roisin Y, Žifčáková L, Park YC, Kim KY, Sillam-Dussès D, Hervé V, Lo N, Tokuda G, Brune A, Bourguignon T. 2022. The functional evolution of termite gut microbiota. Microbiome 10:78.

8. Zheng H, Nishida A, Kwong WK, Koch H, Engel P, Steele MI, Moran NA. 2016. Metabolism of toxic sugars by strains of the bee gut symbiont *Gilliamella apicola*. mBio 7:10.1128/mbio.01326–16.

9. Hammer TJ, Bowers MD. 2015. Gut microbes may facilitate insect herbivory of chemically

defended plants. Oecologia 179:1–14.

10. Kwong WK, Medina LA, Koch H, Sing K-W, Soh EJY, Ascher JS, Jaffé R, Moran NA. 2017.

Dynamic microbiome evolution in social bees. Sci Adv 3:e1600513.

11. Cerqueira AES, Hammer TJ, Moran NA, Santana WC, Kasuya MCM, da Silva CC. 2021.

Extinction of anciently associated gut bacterial symbionts in a clade of stingless bees. ISME

J 15:2813–2816.

12. Engel P, Martinson VG, Moran NA. 2012. Functional diversity within the simple gut

microbiota of the honey bee. Proc Natl Acad Sci U S A 109:11002–11007.

13. Zheng H, Perreau J, Powell JE, Han B, Zhang Z, Kwong WK, Tringe SG, Moran NA. 2019.

Division of labor in honey bee gut microbiota for plant polysaccharide digestion. Proc Natl

Acad Sci U S A 116:25909–25916.

14. Li Y, Leonard SP, Powell JE, Moran NA. 2022. Species divergence in gut-restricted

bacteria of social bees. Proc Natl Acad Sci U S A 119:e2115013119.

15. Keller A, McFrederick QS, Dharampal P, Steffan S, Danforth BN, Leonhardt SD. 2021.

(More than) Hitchhikers through the network: the shared microbiome of bees and flowers.

Curr Opin Insect Sci 44:8–15.

16. Vannette RL. 2020. The floral microbiome: Plant, pollinator, and microbial perspectives.

Annu Rev Ecol Evol Syst 51:363–386.

17. Dew RM, McFrederick QS, Rehan SM. 2020. Diverse diets with consistent core microbiome

in wild bee pollen provisions. Insects 11:499.

18. Keller A, Grimmer G, Steffan-Dewenter I. 2013. Diverse microbiota identified in whole intact

nest chambers of the red mason bee *Osmia bicornis* (Linnaeus 1758). PLoS One

8:e78296.

19. Voulgari-Kokota A, McFrederick QS, Steffan-Dewenter I, Keller A. 2019. Drivers, diversity,

and functions of the solitary-bee microbiota. Trends Microbiol 27:1034–1044.

20. Fernandez De Landa G, Alberoni D, Baffoni L, Fernandez De Landa M, Revainera PD,

Porrini LP, Brasesco C, Quintana S, Zumpano F, Eguaras MJ, Maggi MD, Di Gioia D. 2023.

The gut microbiome of solitary bees is mainly affected by pathogen assemblage and

partially by land use. Environ Microbiome 18:38.

21. McFrederick QS, Rehan SM. 2019. Wild bee pollen usage and microbial communities co-

vary cross landscapes. Microb Ecol 77:513–522.

22. Nguyen PN, Rehan SM. 2023. Wild bee and pollen microbiomes across an urban–rural

divide. FEMS Microbiol Ecol 99:fiad158.

23. McFrederick QS, Mueller UG, James RR. 2014. Interactions between fungi and bacteria

influence microbial community structure in the *Megachile rotundata* larval gut. Proc Biol Sci

281:20132653.

24. Groussin M, Mazel F, Sanders JG, Smillie CS, Lavergne S, Thuiller W, Alm EJ. 2017.

Unraveling the processes shaping mammalian gut microbiomes over evolutionary time. Nat

Commun 8:1–12.

25. Sudakaran S, Kost C, Kaltenpoth M. 2017. Symbiont acquisition and replacement as a

source of ecological innovation. Trends Microbiol 25:375–390.

26. Lampert N, Mikaelyan A, Brune A. 2019. Diet is not the primary driver of bacterial

community structure in the gut of litter-feeding cockroaches. BMC Microbiol 19:238.

27. Ohkuma M, Noda S, Hongoh Y, Nalepa CA, Inoue T. 2009. Inheritance and diversification

of symbiotic trichonymphid flagellates from a common ancestor of termites and the

cockroach *Cryptocercus*. Proc Biol Sci 276:239–245.

28. Bucek A, Šobotník J, He S, Shi M, McMahon DP, Holmes EC, Roisin Y, Lo N, Bourguignon

T. 2019. Evolution of termite symbiosis informed by transcriptome-based phylogenies. Curr

Biol 29:3728–3734.e4.

29. Mikaelyan A, Dietrich C, Köhler T, Poulsen M, Sillam-Dussès D, Brune A. 2015. Diet is the

primary determinant of bacterial community structure in the guts of higher termites. Mol

Ecol 24:5284–5295.

30. Dietrich C, Köhler T, Brune A. 2014. The cockroach origin of the termite gut microbiota:

patterns in bacterial community structure reflect major evolutionary events. Appl Environ

Microbiol 80:2261–2269.

31. Figueroa Laura L., Maccaro Jessica J., Krichilsky Erin, Yanega Douglas, McFrederick

Quinn S. 2021. Why did the bee eat the chicken? Symbiont gain, loss, and retention in the

vulture bee microbiome. mBio 12:e02317–21.

32. Richards O. 1978. The social wasps of the Americas excluding the Vespinae. British

Museum, London.

33. Sugden EA, McAllen RL. 1994. Observations on foraging, population and nest biology of

the Mexican honey wasp, *Brachygastra mellifica* (Say) in Texas [Vespidae: Polybiinae]. J

Kans Entomol Soc 67:141–155.

34. Strassmann JE. 1979. Honey caches help female paper wasps (*Polistes annularis*) survive

Texas winters. Science 204:207–209.

35. Barbosa BC, Detoni M, Maciel TT, Prezoto F. 2017. Resource storage in the neotropical

social wasp *Mischocyttarus socialis* (Saussure, 1854) (Vespidae: Polistini).

Sociobiology 64:356–358.

36. Hastings MD, Queller DC, Eischen F, Strassmann JE. 1998. Kin selection, relatedness, and

worker control of reproduction in a large-colony epiponine wasp, *Brachygastra mellifica*.

Behav Ecol 9:573–581.

37. Rothman JA, Loope KJ, McFrederick QS, Wilson Rankin EE. 2021. Microbiome of the wasp

Vespula pensylvanica in native and invasive populations, and associations with Moku virus.

PLoS One 16:e0255463.

38. Holley JC, Jackson MN, Pham AT, Hatcher SC, Moran NA. 2022. Carpenter Bees

(*Xylocopa*) Harbor a Distinctive Gut Microbiome Related to That of Honey Bees and

Bumble Bees. Appl Environ Microbiol 88:e00203–22.

39. Kwong WK, Moran NA. 2016. Gut microbial communities of social bees. Nat Rev Microbiol

14:374–384.

40. Kang JP, Huo Y, Hoang VA, Yang DU, Yang DC, Kang SC. 2021. *Bombilactobacillus*

apium sp. nov., isolated from the gut of honeybee (*Apis cerana*). Arch Microbiol 203:2193–

2198.

41. Praet J, Meeus I, Cnockaert M, Aerts M, Smagghe G, Vandamme P. 2015. *Bifidobacterium*

commune sp. nov. isolated from the bumble bee gut. Antonie Van Leeuwenhoek 107:1307–

1313.

42. Killer J, Kopecný J, Mrázek J, Rada V, Benada O, Koppová I, Havlík J, Straka J. 2009.

Bifidobacterium bombi sp. nov., from the bumblebee digestive tract. Int J Syst Evol

Microbiol 59:2020–2024.

43. Martinson VG, Magoc T, Koch H, Salzberg SL, Moran NA. 2014. Genomic features of a

bumble bee symbiont reflect its host environment. Appl Environ Microbiol 80:3793–3803.

44. Hammer TJ, Easton-Calabria A, Moran NA. 2023. Microbiome assembly and maintenance

across the lifespan of bumble bee workers. Mol Ecol 32:724–740.

45. Menezes RST, Lloyd MW, Brady SG. 2020. Phylogenomics indicates Amazonia as the

major source of Neotropical swarm-founding social wasp diversity. Proc Biol Sci

287:20200480.

46. Hines HM. 2008. Historical biogeography, divergence times, and diversification patterns of

bumble bees (Hymenoptera: Apidae: Bombus). Syst Biol 57:58–75.

47. Mazzon L, Piscedda A, Simonato M, Martinez-Sañudo I, Squartini A, Girolami V. 2008.

Presence of specific symbiotic bacteria in flies of the subfamily Tephritinae (Diptera

Tephritidae) and their phylogenetic relationships: proposal of "*Candidatus* Stammerula

tephritidis." Int J Syst Evol Microbiol 58:1277–1287.

48. Viale E, Martinez-Sañudo I, Brown JM, Simonato M, Girolami V, Squartini A, Bressan A,

Faccoli M, Mazzon L. 2015. Pattern of association between endemic Hawaiian fruit flies

(Diptera, Tephritidae) and their symbiotic bacteria: Evidence of cospeciation events and

proposal of "*Candidatus* Stammerula trupaneae." Mol Phylogenet Evol 90:67–79.

49. Praet J, Meeus I, Cnockaert M, Houf K, Smagghe G, Vandamme P. 2015. Novel lactic acid

bacteria isolated from the bumble bee gut: *Convivina intestini* gen. nov., sp. nov.,

Lactobacillus bombicola sp. nov., and *Weissella bombi* sp. nov. Antonie Van Leeuwenhoek

50. Hettiarachchi A, Cnockaert M, Joossens M, Laureys D, De Clippeleer J, Vereecken NJ,

Michez D, Smagghe G, de Graaf DC, Vandamme P. 2023. *Convivina* is a specialised core

gut symbiont of the invasive hornet *Vespa velutina*. Insect Mol Biol

https://doi.org/10.1111/imb.12847.

51. Do Y, Park WB, Park JK, Kim CJ, Choi MB. 2023. Gut bacterial diversity in *Vespa velutina*

and implications for potential adaptation in South Korea. Pest Manag Sci 79:5180–5185.

52. McFrederick QS, Thomas JM, Neff JL, Vuong HQ, Russell KA, Hale AR, Mueller UG. 2017.

Flowers and wild Megachilid bees share microbes. Microb Ecol 73:188–200.

53. Crotti E, Rizzi A, Chouaia B, Ricci I, Favia G, Alma A, Sacchi L, Bourtzis K, Mandrioli M,

Cherif A, Bandi C, Daffonchio D. 2010. Acetic acid bacteria, newly emerging symbionts of

insects. Appl Environ Microbiol 76:6963–6970.

54. Hammer TJ, Sanders JG, Fierer N. 2019. Not all animals need a microbiome. FEMS

Microbiol Lett 366:fnz117.

55. Graystock P, Rehan SM, McFrederick QS. 2017. Hunting for healthy microbiomes:

determining the core microbiomes of *Ceratina*, *Megalopta*, and *Apis* bees and how they

associate with microbes in bee collected pollen. Conserv Genet 18:701–711.

56. Chen J, Wang J, Zheng H. 2021. Characterization of *Bifidobacterium apousia* sp. nov.,

Bifidobacterium choladohabitans sp. nov., and *Bifidobacterium polysaccharolyticum* sp.

nov., three novel species of the genus *Bifidobacterium* from honey bee gut. Syst Appl

Microbiol 44:126247.

57. Martinson VG, Moy J, Moran NA. 2012. Establishment of characteristic gut bacteria during

development of the honeybee worker. Appl Environ Microbiol 78:2830–2840.

58. Hammer TJ, Le E, Martin AN, Moran NA. 2021. The gut microbiota of bumblebees.

Insectes Soc 68:287–301.

59. Villabona N, Moran N, Hammer T, Reyes A. 2023. Conserved, yet disruption-prone, gut

microbiomes in neotropical bumblebees. mSphere 8:e0013923.

60. Onchuru TO, Javier Martinez A, Ingham CS, Kaltenpoth M. 2018. Transmission of

mutualistic bacteria in social and gregarious insects. Curr Opin Insect Sci 28:50–58.

61. Rose C, Lund MB, Søgård AM, Busck MM, Bechsgaard JS, Schramm A, Bilde T. 2023.

Social transmission of bacterial symbionts homogenizes the microbiome within and across

generations of group-living spiders. ISME Commun 3:1–10.

62. Kueneman JG, Gillung J, Van Dyke MT, Fordyce RF, Danforth BN. 2022. Solitary bee

larvae modify bacterial diversity of pollen provisions in the stem-nesting bee, *Osmia*

cornifrons (Megachilidae). Front Microbiol 13:1057626.

63. Handy MY, Sbardellati DL, Yu M, Saleh NW, Ostwald MM, Vannette RL. 2023. Incipiently

social carpenter bees (*Xylocopa*) host distinctive gut bacterial communities and display

geographical structure as revealed by full-length PacBio 16S rRNA sequencing. Mol Ecol

32:1530–1543.

64. Christensen SM, Srinivas SN, McFrederick QS, Danforth BN, Buchmann SL, Vannette RL.

2024. Symbiotic bacteria and fungi proliferate in diapause and may enhance overwintering

survival in a solitary bee. ISME J 18:wrae089.

65. Montoya D, de Jong M. 2023. Warming-induced range expansion effects on the diversity

and composition of the gut microbiome: a case study with two butterflies. bioRxiv.

66. Ramirez KS, Snoek LB, Koorem K, Geisen S, Bloem LJ, Ten Hooven F, Kostenko O,

Krigas N, Manrubia M, Caković D, van Raaij D, Tsiafouli MA, Vreš B, Čelik T, Weser C,

Wilschut RA, van der Putten WH. 2019. Range-expansion effects on the belowground plant

microbiome. Nat Ecol Evol 3:604–611.

67. Fischer MK, Shingleton AW. 2001. Host plant and ants influence the honeydew sugar

composition of aphids. Funct Ecol 15:544–550.

68. Meiners JM, Griswold TL, Harris DJ, Ernest SKM. 2017. Bees without flowers: Before peak

bloom, diverse native bees find insect-produced honeydew sugars. Am Nat 190:281–291.

69. Santas LA. 1983. Insects producing honeydew exploited by bees in Greece. Apidologie

14:93–103.

70. Milani C, Lugli GA, Duranti S, Turroni F, Mancabelli L, Ferrario C, Mangifesta M, Hevia A,

Viappiani A, Scholz M, Arioli S, Sanchez B, Lane J, Ward DV, Hickey R, Mora D, Segata N,

Margolles A, van Sinderen D, Ventura M. 2015. Bifidobacteria exhibit social behavior

through carbohydrate resource sharing in the gut. Sci Rep 5:1–14.

71. Powell JE, Martinson VG, Urban-Mead K, Moran NA. 2014. Routes of acquisition of the gut

microbiota of the honey bee *Apis mellifera*. Appl Environ Microbiol 80:7378–7387.

72. Quast C, Pruesse E, Yilmaz P, Gerken J, Schweer T, Yarza P, Peplies J, Glöckner FO.

2013. The SILVA ribosomal RNA gene database project: improved data processing and

web-based tools. Nucleic Acids Res 41:D590–6.

FIGURE LEGENDS

Figure 1: Specimen images and collecting locations. (A) Mexican honey wasp, *Brachygastra mellifica*, (B) common paper wasp, *Polistes exclamans*, (C) honey bee, *Apis mellifera*. Specimens have their guts removed to show the internal structure. The bottom section of the images were cut and rotated to orientate the midguts to the right and sting to the left. (D) Map of Texas showing the collecting sites of honey wasps, *Polistes* paper wasps and solitary wasps, and honey bees.

Figure 2: Relative abundance of (A) the top 16 bacteria genera and (B) the top 50 amplicon sequence variants (ASVs) from honey wasps, other wasps, and honey bees. Honey wasps are grouped by nest, with the exception of five individuals (Ind). Nests are named based on their location and organized on the plot from north (top) to south Texas (bottom). Other wasps are grouped by family (Vesp = Vespidae, Mut = Mutillidae, Thynn = Thynnidae, Crab = Crabronidae, Sphec = Sphecidae, Ich = Ichneumonidae). Honey wasps typically possess a characteristic microbiome that distinguishes them from other wasp species.

Figure 3: Nonmetric multidimensional scaling (NMDS) ordination of Bray-Curtis dissimilarity index. (A) Honey wasp gut microbial communities compared to those of other wasps and honey bees (stress = 0.19). (B) Honey wasp nest gut microbial communities (stress = 0.24).

Figure 4: Gut bacterial abundance of honey wasps, *Polistes* paper wasps, and solitary wasps. Dashed line is the median. The gray shaded area is the region below the detection threshold, and the yellow shaded area represents the range of 16S rRNA gene copies for *Apis mellifera* from Kwong et al. (10). Honey wasps (n = 10) were from nests in San Antonio (SA1 and SA3), *Polistes* samples represent five species (n = 19), and solitary wasps represent 7 families (n = 30, including four solitary eumenine vespid wasps). Honey wasp gut microbial communities significantly differed from those of paper wasps and solitary wasps in all pairwise comparisons (honey wasps:paper wasps, z = -3.417, adjusted p = 0.002; honey wasps:solitary wasps, z = -4.990, adjusted p = 1.8×10^{-6}).

Figure 5: Phylogenies of the 16S rRNA gene of the major bacterial taxa from the honey wasps. (A) Orbaceae, (B) Bifidobacteriaceae, and (C) Lactobacillaceae. Phylogenies were inferred by maximum likelihood, bootstrap support indicated by black circles. Colors represent bacterial genera or clades associated with bees, host source listed in parentheses.

Figure 6: Diagram showing the highly abundant bacteria found in the Mexican honey wasp (slate blue) that have host-switched from bumble bees (brown), honey bees (cream), and vespid bees (light blue). Bacterial names in black were found in honey wasps and names in gray are host-restricted bacteria that were not found in honey wasps. Bacteria commonly associated with solitary bees (gray) and the nectar in wild flowers (gray) were not found in honey wasps. The *Lactobacillus** present in honey wasps is likely a new species unknown from other insects.

Bumble bee

Bifidobacterium
Convivina
Ca. Schmidhempelia

Honey bee

Bombilactibacillus
Bifidobacterium
Gilliamella
Lactobacillus
Snodgrassella
Frischella

Bifidobacterium
Bombilactobacillus
Convivina
Ca. Schmidhempelia
Fructobacillus
*Lactobacillus**

Wild flower

Acinetobacter
Glucnobacter
Pseudomonas
Rosenbergiella
Saccharibacter

Honey wasp

Vespid wasp

Fructolactobacillus
Convivina
Lactococcus
Leucnostoc

Solitary bee

Apilactobacillus
Acinetobacter
Arsenophonus