NeuroSCAN: Exploring **Neuro**development via **S**patiotemporal **C**ollation of **A**natomical **N**etworks

Authors: Noelle L. Koonce[1], Sarah E. Emerson[1], Dhananjay Bhaskar[2], Manik Kuchroo[2], Mark W. Moyle[1,10], Pura Arroyo-Morales[1], Nabor Vázquez Martínez[1], Smita Krishnaswamy[2, 3, 4, 5], William Mohler[6*], Daniel Colón-Ramos[1,7,8,9*]

1. Department of Neuroscience and Department of Cell Biology, Wu Tsai Institute, Yale University, New Haven, CT, USA
2. Department of Genetics, Yale University School of Medicine, New Haven, CT, USA
3. Department of Computer Science, Yale University, New Haven, CT, USA
4. Program for Computational Biology and Bioinformatics, Yale University, New Haven, CT, USA
5. Program for Applied Mathematics, Yale University, New Haven, CT, USA
6. Department of Genetics and Genome Sciences and Center for Cell Analysis and Modeling, University of Connecticut Health Center, Farmington, CT, USA
7. MBL Fellow, Marine Biological Laboratory, Woods Hole, MA, USA
8. Wu Tsai Institute, Yale University; New Haven, CT 06510, USA
9. Instituto de Neurobiología, Recinto de Ciencias Médicas, Universidad de Puerto Rico; San Juan 00901, Puerto Rico
10. Department of Biology, Brigham Young University-Idaho, Rexburg, ID, USA

*Corresponding authors emails: daniel.colon-ramos@yale.edu and wmohler@neuron.uchc.edu

Summary

Volume electron microscopy (vEM) datasets such as those generated for connectome studies allow nanoscale quantifications and comparisons of the cell biological features underpinning circuit architectures. Quantifications of cell biological relationships in the connectome result in rich multidimensional datasets that benefit from data science approaches, including dimensionality reduction and integrated graphical representations of neuronal relationships. We developed NeuroSCAN, an online open-source platform that bridges sophisticated graph analytics from data science approaches with the underlying cell biological features in the connectome. We apply NeuroSCAN to

a complete published record of *C. elegans* brain neuropils and demonstrate how these integrated representations of neuronal relationships facilitate comparisons across connectomes, catalyzing new insights on the structure-function of the circuits and their changes during development. NeuroSCAN is designed for intuitive examination and comparisons across connectomes, enabling synthesis of knowledge from high-level abstractions of neuronal relationships derived from data science techniques to the detailed identification of the cell biological features underpinning these abstractions.

Keywords:

Connectome; Contactome; *C. elegans*; Nerve ring; Neuropil; Electron microscopy; Neurodevelopment

Introduction

Neural circuit structure supports function. The underlying image data that yields anatomical connectomes (or wiring diagrams) are obtained from volume electron microscopy (vEM) data (Collinson et al. 2023). These last decades have seen an increase in the generation of these high quality EM datasets, and since the first connectome was published for *C. elegans* (White et al. 1986), over 30 EM-based anatomical connectomes have been generated for whole organisms or tissue regions (Burns et al. 2013; S. Y. Takemura et al. 2013; Wanner, Genoud, and Friedrich 2016; S. Takemura, Nern, et al. 2017; Maclachlan et al. 2018; Marc et al. 2018; Pipkin et al. 2018; Svara et al. 2022; Randel et al. 2014; S. Takemura et al. 2015; S. Takemura, Aso, et al. 2017; Motta et al. 2019; Zheng et al. 2018; Scheffer et al. 2020; Shapson-Coe et al. 2021; Azevedo 2024; The MICrONS Consortium et al. 2023; Vishwanathan et al. 2017; Yim et al. 2024; Phelps et al. 2021; Ryan, Lu, and Meinertzhagen 2016). The expansion in available anatomical connectomes has resulted from recent advancements in: 1) data generation (via semi automation of EM data acquisition (Xu et al. 2017; Eberle and Zeidler 2018; Zheng et al. 2018; Phelps et al. 2021); and 2) alignments, segmentations and reconstructions

(including recent implementation of AI-driven methods) (Berger, Seung, and Lichtman
2018; Galili, Jefferis, and Costa 2022). As these developing methodologies continue to
improve, they will continue to facilitate the generation of additional connectomes of whole
brains and organisms.

The increasing availability of ssEM datasets, including the first series of
developmental connectomes published for *C. elegans* (Witvliet et al. 2021) has
highlighted the need for new tools to enable intuitive examination and comparisons across
connectomes to promote novel discoveries (Kasthuri et al. 2015; Lichtman, Pfister, and
Shavit 2014; Barabási et al. 2023). It has also underscored the fact that ssEM datasets
contain a wealth of untapped information that has yet to be fully examined, represented
and integrated for more comprehensive analyses (Perez et al. 2014; Brittin et al. 2021).
For example, ssEM datasets enable nanoscale explorations of the underlying cell
biological features that govern the properties of neural circuit architectures (Rivlin et al.
2024; Brittin et al. 2021; Moyle et al. 2021; Witvliet et al. 2021). Yet most of these cell
biological features (cell morphologies, contact profiles, organelle positions and shapes,
etc) are not currently represented in most anatomical connectomes. Quantification of cell
biological data result in high-dimensional datasets that require new approaches for their
analyses and representations. The advances in ssEM data generation and the resulting
need for new methodologies in data science and integrated representations of neuronal
relationships (e.g. from neuronal positions to neuropil structures) is akin to how advances
in genetic sequencing required new methodologies in bioinformatics and new, integrated
representations of genomic data (e.g. from gene sequence to gene structure) (Swanson
and Lichtman 2016). Addressing this gap holds the promise of integrating new knowledge
from the fields of cell biology, neurodevelopment, physiology and systems neuroscience
towards explaining how nervous system structure underpins its function.

Most representations of anatomical connectomes have focused on defining
neuronal relationships at the level of the chemical synapse (NemaNode; WormWiring;
EleganSign; FlyWire) (Witvliet et al. 2021; Cook et al. 2019; Fenyves et al. 2020;
Dorkenwald et al. 2023). While the existence of chemical synapses between neuron pairs
is an important feature of neuronal communication, these representations do not capture
other neuroanatomical features that also underlie neuron structure and function, including

contact sites from adjacent (or nearby) neurons. Recent work in *C. elegans* examined

neuronal relationships by quantifying neuron-neuron contact sites to build contact profiles,

or contactomes (Moyle et al. 2021; Brittin et al. 2021; Witvliet et al. 2021). Examination of

the contactome with data science approaches uncovered structural principles that were

not evident from interrogating the synaptic connectome alone (Moyle et al. 2021; Brittin

et al. 2021). These included the existence of higher-order structural motifs and the

stratification of neurons (Moyle et al. 2021), whose hierarchical assembly during

development is guided by centrally located pioneer neurons (Rapti et al. 2017). Moreover,

integrating neuronal adjacencies (contactome) with synaptic profiles (connectome)

allowed for a deeper understanding of the functional segregation of neurons within the

stratified neuropil structures (Brittin et al. 2021; Moyle et al. 2021). Key to achieve this

were data science approaches such as Diffusion Condensation (DC) and C-PHATE

(Brugnone et al. 2019; Moon et al. 2019), which resulted in reduced dimensionality of the

neuronal relationships, revealing architectural motifs across various scales of granularity,

from individual neurons within circuits, to individual circuits within the neuropil. These

techniques produced abstract graphs that enabled exploration of these computationally

identified groups (Moyle et al. 2021). DC/C-PHATE graphs are powerful tools, but they

have yet to be integrated to connectomics datasets as to enable explorations of the

underlying cell biological features. This limits their effectiveness for hypothesis generation

and comparative analyses across connectomes.

To address this, we generated NeuroSCAN, a tool for exploring neuroarchitectures

across ssEM datasets via novel representations of the connectome, contactome, and

anatomical networks. NeuroSCAN is an online, open-source platform that facilitates

comparisons of neuronal features and relationships across ssEM data to catalyze new

insights of the relationships that underpin architectural and functional motifs of the nerve

ring neuropil. NeuroSCAN builds on recent publications in whole-brain EM datasets,

integrating the latest set of developmental connectomes (Witvliet et al. 2021) and

employing data science tools (Brugnone et al. 2019; Moon et al. 2019) to examine

neuronal relationships based on contact profiles. We demonstrate how these integrated

representations of neuronal relationships facilitate comparisons across these

connectomes, catalyzing new insights on their structure-function and changes during

development. NeuroSCAN achieves this by addressing three challenges in current neuronal representations: 1) accessibility of specific neuronal cell biological features (i. e. synapses and contacts), 2) integration of features for examining neuronal relationships across anatomical scales, and 3) spatiotemporal comparisons of these features across developmental datasets.

These challenges were addressed by 1) creating representations of contact sites and establishing the ability to visualize subsets of synaptic sites; 2) enabling synchronous visualization of neuron morphologies, contacts and synapses and integrating these cell biological features with algorithmically-generated graphical representations of neuronal relationships; and 3) enabling simultaneous exploration of these relational representations across developmental connectomes. NeuroSCAN was designed as a suite of tools that facilitates future incorporation of additional datasets and representations with the goal of enabling integrated data exploration beyond the available *C. elegans* connectomes. The NeuroSCAN-based approaches used here for *C. elegans* could be applicable to other systems as new EM-based datasets and reconstructions become available.

Results

2.1. Comparing contactome-based relationships using C-PHATE.

The adult hermaphrodite *C. elegans* nerve ring is a neuropil of 181 neurons of known identities, morphologies, contact profiles, and synaptic partners (White et al. 1986). Even for this relatively small neuropil, representations of a single feature type, such as neuronal contact profiles, constitute over 100,000 data points of multidimensional information: cell identity, region of contact, presence of synapses, etc. Analysis of this multidimensional information requires approaches that can both capture higher order patterns of organization while enabling researchers to access the underlying cell biological features resulting in these relationships. We implemented Diffusion Condensation (DC), a clustering algorithm that iteratively group neurons based on the quantitative similarities of their 'contact' or 'adjacency' profiles (Brugnone et al. 2019; Moyle et. al 2021). Briefly, DC makes use of pair-wise quantifications of adjacent neuron

contacts to move neurons with similar adjacency profiles closer together by applying a
diffusion filter in a multidimensional manifold. This diffusion filter effectively removes
variability between neurons at each iteration. As iterations proceed, individual neurons
(and eventually groups of neurons) are clustered together based on how close their
diffused contact profiles are to one another in the manifold (Brugnone et al. 2019). In this
way, DC uncovers hierarchical neuronal relationships in the contactome (Moyle et al.
2021).

To ensure accurate comparisons of DC across available EM datasets (Witvliet et
al. 2021; White et al. 1986), we first empirically set minimum-distance adjacency
thresholds (measured in pixels; Table S1) to build adjacency profiles (see Methods,
schematized in Figure 1A-C). We then quantified the lengths of the physical adjacencies
(or contacts) between neuron pairs (Brittin et al. 2021) and built an adjacency matrix for
each of the seven selected *C. elegans* contactome datasets (Figure 1C; Methods). To
visualize and compare the results from DC, we used C-PHATE (Moon et al. 2019; Moyle
et. al 2021), a 3-D visualization tool that builds a hierarchical visual representation of the
DC agglomeration procedure (Figure 1 D-E). In C-PHATE visualizations, the DC output
is mapped in 3-D space with spheres. Initially, all individual neurons in the neuropil
dataset are at the periphery of the C-PHATE graph (left hand side in schematic in Figure
1D, edges of graph in Figure 1E). Neurons are iteratively condensed based on the
similarity of their adjacency profiles (schematized in Figure 1D). In the last iteration of DC,
there is a single point at the center of the C-PHATE graph which contains the entire
neuropil (Figure 1E, red dot). C-PHATE representations enable visualization and
comparisons of contactomes across datasets, and explorations of neuronal relationships,
from individual neuron interactions to circuit-circuit bundling (Figure 1F and Figure 2).

By Larval stage 1 (L1), neuronal differentiation has concluded and 90% of the
neurons in the neuropil (161 neurons out of the 181 neurons) have entered the nerve ring
and adopted characteristic morphologies and positions (Sun and Hobert 2023). Although
the organism grows ~5 fold from L1 to the adult, contacts in the nerve ring are also largely
established by L1 and preserved during postembryonic growth (Witvliet et al. 2021). In
agreement with this, when we used DC and C-PHATE to examine contactomes from
these datasets we consistently identified four main superclusters– Stratum 1, Stratum 2,

Stratum 3, and Stratum 4 (Figure 2B-F). These findings are consistent with previous
studies on the Larval Stage 4 (L4) and adult contactomes (Moyle et al. 2021), and further
suggest that neurons establish core relationships during embryogenesis and maintain
them into adulthood. Moreover, aligning the neuronal morphologies of strata members
reveals a persistent layered organization to the nerve ring neuropil (Figure 2 G-K), and
exploring the functional identities of the neurons in each stratum suggests that there is
spatial segregation of sensory information and motor outputs (see (Moyle et al. 2021) and
tables S3-S6). Our findings are in agreement with previous reports that the organization
of the nerve ring is largely established in embryogenesis, and then maintained during
postembryonic growth (Witvliet et al. 2021). Our findings also demonstrate the utility of
DC and C-PHATE analyses in extracting, visualizing and comparing the structure of the
neuropil architecture across contactomes.

Because DC and C-PHATE allow for the examination of relationships at varying
levels of granularity, they also facilitate the interrogation of the architectural motifs that
underly distinct neural strata. A more detailed examination of clusters reveals that while
the overall strata are preserved, the underlying neuronal configurations undergo changes
during postembryonic growth (Figure 2 B-F, Figure 3 and Tables S3-S6). Three general
features were extracted from these analyses: 1) individual neurons renegotiate their
positions in the context of the identified C-PHATE clusters in different developmental
contactomes, suggesting developmental changes; 2) the degree of these changes varied
across the distinct strata; and 3) the degree of these changes mapped onto known
features of each stratum, such as plasticity. For example, Stratum 1, which contains
shallow reflex circuits, displayed the fewest changes among the developmental
connectomes (Figure 3 B-F; Table S3). On the other hand, Strata containing circuits
associated with behavioral plasticity (Stratum 3 and Stratum 4), displayed the largest
changes across postembryonic development (Figure 3 H-L; Table S5 - S6).

To examine the changes in DC/C-PHATE during postembryonic development, we
made the C-PHATE plots fully interactive. This enables users to hover over and identify
members of each intermediate cluster, to highlight specific cell trajectories via pseudo-
coloring, and compare specific neuronal relationship dynamics across development
within a multiview window of distinct C-PHATE plots (Figure 1 E-F, Figure S6, Video S1).

Because C-PHATE graphs ultimately represent cells of known identities, we reasoned that interactive mapping of the C-PHATE cluster objects to their component cellular identities and anatomies could yield greater insights on neurodevelopmental changes, linking the algorithmic abstractions of the relationships with the cell biological features and their changes across development (Figure 4).

To examine our hypothesis and determine the utility of C-PHATE for discovery, we inspected specific regions where the distribution, or 'shape' of superclusters changed across the set of developmental contactomes. When comparing C-PHATE graphs representing distinct contactomes, we accounted for changes in the iterations at which "merge events" (or co-clustering of neurons) occurred. The logic in considering the 'iterations of the merge events' is because variations in contact profiles influence changes in iterations of merge events. Based on these criteria, we focused on a region displaying changes in Strata 3 and 4 and using the interactive C-PHATE graphs (Figure 4 A-E), we determined the identities of neurons that changed clustering patterns across the developmental contactomes (Figure 4F and G). Specifically, we focused on two interneurons, named AIML and PVQL, which we observed undergo a change in their cluster assignment from Stratum 4 (at L1) to Stratum 3 (at Larva stage 4, L4; Figure 4A and D). We pseudo-colored the trajectories of the AIML and PVQL neurons in C-PHATE to explore the changes in their merge events throughout the developmental stages (Figure 4F and G; Figure S1; Table S7). Comparing L1, L2 (Larval Stage 2) and L3 (Larval Stage 3) datasets, we observe the AIML and PVQL neurons merge at iterations 16, 14 and 22 (respectively). The increasing numbers of iterations across the L1, L2 and L3 datasets suggests the relative contact profiles of AIML and PVQL diverge across these contactomes (Figure 4F and G; Figure S1; Table S7). Yet, between the L4 and Adult datasets, we observe the PVQL and the AIML neurons merge at iterations 20 in the L4 and iteration 14 in the Adult (Figure S1; Table S7). The decrease in the number of iterations required for the merge event suggests that the relative contact relationships of AIML and PVQL eventually converge between L4 and adult animals. Comparison of the identities of the neurons that co-cluster with AIML and PVQL similarly suggests that the contact relationships varied across developmental stages (Figure 4F and G; Figure S1, Table S7).

2.2. Visualizing contact profiles in individual cells.

DC/C-PHATE changes should result from changes in adjacency or contact profiles. To link the observed changes in the C-PHATE graphs with the cell-biological changes in contact profiles, we generated a tool that would simultaneously enable: 1) 3D visualization of the cell-cell contact sites onto individual neuronal morphologies; 2) examination and comparisons of these contact profiles throughout development for the available contactomes; and 3) integration with DC/C-PHATE to link C-PHATE cluster objects to the 3-D morphologies of the algorithmically clustered cells. With these capabilities integrated, we could simultaneously view the contactome from two complementary perspectives – at an abstract systems level via DC/C-PHATE and at a cell biological level via 3D contact modeling – to perceive the architectural themes that underlie similar network patterns.

To create this tool, we generated 3D models of the area of physical contact between adjacent neuron pairs (Table S1-S2; Methods; Figure 5; Figure S2). Visualizing contacts from all adjacent neurons builds a multi-colored skeleton of the neuron morphology mapped onto the boundaries of this neuron (Figure 5A and C). Because the identities of the neurons are known and linked to the 3D contact models, we built text pop ups that define the contact partners for each site (Figure 5C). Furthermore, since neuron names are consistent across EM datasets, we can link and compare contact sites across development (Figure 5D). Additionally, we can analyze the representations of contact sites in the context of DC/C-PHATE clustering profiles (Figure 5B), 3D models of neuronal morphologies (Figure 1F), and 3D models of synaptic sites for any neuron(s) across development (Figure 7).

We used the integrated tools of DC/C-PHATE and 3D representations of the contact profiles to examine the potential cell biological changes leading to the DC/C-PHATE clustering changes observed for the AIML neuron during development. With these tools, we observed changes in the identities of the contacts made in the dorsal region of the AIML neurite (Figure 5D; Figure S3). Specifically, in the L2 stage (as compared to L1), we observed a decrease in the contacts from PVQL and an increase in

contacts from the AVF neurons. This change persists to the adult stage (Figure 5D; Figure
S3).

To then determine the possible source of these developmental changes in
contacts, we visualized 3D models of the segmented morphologies for these neurons
across L1 to adulthood (Figure 6). We find that AIM and PVQ neurons maintain similar
morphologies throughout development (Figure 6C), while AVF neurons undergo
substantial neurite outgrowth onto new regions of contact between AIM and PVQ (Figure
6 B-D). Specifically, the data revealed that although the AVF neurons terminally
differentiate in the embryo, they do not grow into the nerve ring until the L2 stage, and
continue to grow until the Adult stage (Figure 6B-D). The AVF neurons grow in between
the AIM and PVQ neurons (Figure 6D), altering their contact profiles, which likely
contributes to the observed changes in the C-PHATE graphs (although we note that
DC/C-PHATE representations systematically cluster neurons based on relative
similarities across contact profiles, not solely by scoring changes in specific contacts
within any given pair; Figure 4F and G; Figure 5B and D; Video S2). We also observe that
both AVFL and AVFR grow into the nerve ring alongside AIML, later continuing to grow
around to reach AIMR, and that these relationships were also reflected in the C-PHATE
graphs in terms of the clustering profiles throughout development; Figure 4G; Figure S1).

We then examined if the developmental changes in contact profiles result in
changes in circuitry. We examined this by layering on synaptic information. Despite
dwindling AIM-PVQ contacts, AIM and PVQ neurons maintained their synaptic
relationship throughout development, with synaptic sites observed primarily at the base
of AIM neurons, a region of persistent contact with PVQ (Figure 7A-B). We observed that
increases in contacts between AIM and AVF neurons resulted in additional *en passant*
synapses at the new points of contact, beginning at the L2 stage and continuing to
adulthood (Figure 7A-B). We also observed that AVF forms synapses with the adjacent
PVQ neurons (Figure 7; Figure S4).

In summary, by integrating, representing and comparing datasets using the new
C-PHATE tools and contact profiles in NeuroSCAN, we identified developmental changes
in the relationships of AIM, AVF and PVQ. This case-study highlights the utility of
combining cell biological representations (such as morphologies, contacts and synapses)

with coarse-grained systems-level representations (like DC/C-PHATE) of vEM datasets

to uncover developmental changes which could be further explored experimentally.

Therefore, NeuroSCAN serves as a powerful platform for generating hypotheses for

empirical testing, which can lead to insights into the dynamics of circuit development.

2.3 NeuroSCAN: Facilitating multi-layered interrogation of neuronal relationships

in the *C. elegans* nerve ring throughout larval development

NeuroSCAN is built as a web-based client-server system designed to enable the

sharing of anatomical connectomics data with an emphasis on facilitating the analyses of

neuropil relationships across hierarchies and scales. To achieve this, we integrated tools

of neuroanatomical investigation from the available *C. elegans* nerve ring connectomes

and contactomes with a collection of 3-D modeled elements (morphologies, contacts and

synapses and C-PHATE) representing different aspects of neuronal architecture and

relationships (Figure 8). NeuroSCAN differs from other available web-based tools in this

area with the integration of C-PHATE graphs that enable exploration of hierarchical

organizations of stratified fascicles, the availability of new tools to examine the

contactome, and the integration of these data with existing connectome and

morphological datasets across developmental stages.

NeuroSCAN has eight key user-driven features: (1) C-PHATE, with the ability to

highlight clusters containing neurons of interest (Figure S6; Video S1), (2) reconstructions

of neuronal morphologies (Figure S10; Video S3), (3) reconstructions of neuronal

morphologies of C-PHATE cluster members with a right-click on C-PHATE clusters

(Video S1), (4) 3-D renderings of neuronal contacts to visualize the spatial distribution of

contact profiles (Figure S5; Video S4), (5) 3-D representations of synaptic sites with the

option to visualize subsets of those sites (Figure S7; Video S4), (6) the ability to perform

side-by-side comparisons across development (Figure S11; Video S3), (7) the option to

pseudo color each object to highlight points of interest (Figure S11; Video S3), and (8)

each item is an individual object with the ability to be further customized by the user

(Figure S11-S12).

The NeuroSCAN website architecture and data structure were designed to

integrate these key user-driven features via a modular platform and linked datasets. The

architecture uses Geppetto, an open-source platform designed for neuroscience
applications, modularity, and large datasets (Cantarelli et al. 2018). Briefly, the
architecture is effectively separated into two applications, a frontend React/JavaScript
bundle that is delivered to the client, rendering the neuron data and assets, and a NodeJS
application that exposes a JSON API, serving the neuron data and assets based on user
interactions (Figure S13). The backend uses a Postgres Database to store underlying
data (Figure S14), a Persistent Storage Volume that houses and serves static assets,
and a variable number of Virtual Machines to run the frontend and backend application
code, scaling as needed to accommodate traffic. The User Interface is a React application
that allows users to filter, sort, and search through the Neurons so that they can be added
to an interactive canvas (Figure S13). When users add Neurons to a viewer, a .gltf file is
loaded in for a given model (Synapses, Neurons, Contacts) at the selected developmental
stage (Figure S13), which can then be manipulated in the 3D environment or layered with
other meshes as needed. NeuroSCAN can be used on common web-browsers (e.g.
Google Chrome, Safari) and mobile devices.

The underlying data model makes use of tables representing Synapses, Neurons,
Contacts and Developmental Stages. Relationships between these models are
represented by foreign keys (Figure S14). Source data is defined in a file-tree structure
containing various assets (such as .gltf files representing various entities), as well as
CSV's which store relationships across entities. The directory structure outlines a vertical
hierarchy, starting at the developmental stages, then branching downwards onto neuron
and synapse data. A Python script is invoked to traverse the directory tree and parse the
files, writing to the database accordingly. This configuration enables: 1) verification of the
ingested data and 2) quick search times through the datasets to identify related items.
Code is version-controlled in GitHub (https://github.com/colonramoslab/NeuroSCAN) and
deployed through a CI/CD pipeline when updates are committed to the main branch
(Figure S13).

2.4 NeuroSCAN: practical considerations

We offer seven practical considerations for users. First, NeuroSCAN is available
on mobile platforms as a quick and convenient way to look up neuron morphologies and

relationships. Second, since contact sites offer the ability to explore the surrounding
neurons and the position(s) of contact between adjacent neurons, NeuroSCAN is
designed to enable studies of adjacent neurons (e.g. phenotypes that result in site-
specific ectopic synapses; neuron morphology changes that may affect specific
surrounding neurons; developmental events requiring communication between neurons,
etc.). Third, C-PHATE can be used to identify neurons with similar contact profiles.
Because contact profiles are associated with circuits identities (Moyle et al. 2021),
exploration of neuronal relationships via C-PHATE can be used to identify new
relationships between specific neurons and circuits. Fourth, visualization of subsets of
synaptic and contact sites allows direct comparisons to light microscopy approaches such
as cell-specific labeling of synapses or GFP-Reconstitution across synaptic partners
(Feinberg et al. 2008). Fifth, because the color and transparency of each 3-D model can
be customized, users can further integrate NeuroSCAN outputs of additional atlases (for
gene expression, neurotransmitter and receptor expression, functional connectivity,
etc.(Packer et al. 2019; Taylor et al. 2021; Wang et al. 2023; Fenyves et al. 2020; Randi
et al. 2023) and directly use the NeuroSCAN outputs to create figures and comparisons
(as done for this paper). Sixth, although synaptic sites with BWM (body wall muscles) are
included in NeuroSCAN, the current data model limits the ability to search for these non-
neuronal cells. Users can search for neurons with synapses to BWM to find this datatype.
Seventh, to enable direct comparisons between our data representations and the primary
EM data, the original annotations have been preserved and can be accessed by users
via the sister app, CytoSHOW (CytoSHOW.org). As the data continues to be curated, the
modular design of NeuroSCAN and its companionship with CytoSHOW enables
integration of future annotations.

Discussion

NeuroSCAN is an integrative tool for analyzing detailed, web-based
representations of neuronal connectomes and contactomes throughout post-embryonic
development in *C. elegans*. Connectomes and contactomes are derived from volume
electron microscopy (vEM) micrographs of neuropil regions (Witvliet et al. 2021; White et

al. 1986). These EM micrographs are information-rich and have the potential to reveal
architectural motifs across scales, from the nanoarchitecture of the neuron to the
neuroanatomy of each circuit in the brain. Cell biological features, such as contact profiles
and synaptic positions, can be rigorously quantified and systematically represented as
graphs capturing multidimensional relationships. These representations require
methodologies from data science that enable dimensionality reduction and comparisons
of the architecture across scales. Yet to derive new intuitions about the spatiotemporal
events leading to the architecture that shapes its function, it is necessary to integrate and
compare these various representations, bridging knowledge from the cell biological
events to the systems-level network relationships. NeuroSCAN is designed to achieve
this integration, enabling synthesis of knowledge ranging from the abstractions of
neuronal relationships in C-PHATE to the cell biological features underpinning these
abstractions. We provide a case study to illustrate how integration of analyses performed
in NeuroSCAN can result in new insights. First, we demonstrated the discovery process
with C-PHATE representations to identify neurons that undergo changes in their
contactome during development. Second, we developed 3-D representations of contact
sites to analyze the local neuronal regions that were identified via DC/C-PHATE analysis.
Third, we visualized and compared these representations across development to zero-in
on cell biological changes in neuronal morphologies and synaptic positions across neuron
classes. Our case study demonstrates the utility NeuroSCAN to facilitate exploration of
neuronal relationships, leading to new insights on structural features of the connectome
and hypotheses for empirical testing.
427

3.1 Comparisons of NeuroSCAN to other connectomics atlases
NeuroSCAN is one of several efforts centered around interpreting the *C. elegans*
EM datasets. Other open-source tools for data exploration in *C. elegans* include efforts
to capture neuron morphologies and synaptic information (including integration of new
connectomes across larval development), to map neurotransmitter and receptor
expression, and to record whole brain functional connectivity across genotypes (Witvliet
et al. 2021; Altun, Z.F. et al. 2002; Cook et al. 2019; Fenyves et al. 2020; Randi et al.
2023). NeuroSCAN was inspired by tools like NemaNode and WormWiring (Witvliet et al.

2021; Cook et al. 2019), which enable 3-D visualizations of neuronal morphologies and
synaptic sites with synaptic subsets restricted to pre or postsynaptic sites. In NeuroSCAN
we sought to generate and integrate information beyond the synaptic connectome to
include local neuronal regions (contactome) and neuronal morphologies across available
developmental vEM datasets. Contactomes represent features that have been largely
overlooked in connectomic datasets, and which capture circuit structures not evident by
inspecting solely synaptic relationships (Brittin et al. 2018). NeuroSCAN extends existing
representations to also offer user-driven experience with choice over the visualization of
specific synaptic sites, the option to search for synaptic partners, and the ability to
customize the color of each synaptic representation (Figure 7). NeuroSCAN
representations complement resource databases like WormAtlas, which hosts digitized
electron micrographs and schematics of neuron morphologies with aggregated
information on each neuron (Altun, Z.F. et al. 2002). As such, NeuroSCAN extends an
existing suite of open-source resources to facilitate community wide exploration of vEM
datasets.
451
3.2 NeuroSCAN design and future directions
NeuroSCAN code and development was intentional in its design as an open-
source resource that is modular and allows integration of additional features and data
structures (Cantarelli et al. 2018). It is a hypothesis-generating tool that can be equally
used by educators seeking to teach neuroanatomical principles, and researchers seeking
to identify changes across connectome datasets. NeuroSCAN could be integrated into
emerging datasets, including developmental time-courses of cell-specific transcriptomic
data that would enable further insights on the molecular events underpinning neuronal
development and function– from synaptogenic processes to the logic of neurotransmitter
use (Packer et al. 2019; Taylor et al. 2021; Fenyves et al. 2020) and how it sustains
functional connectivity (Randi et al. 2023). Future iterations of NeuroSCAN could also
include positions and relationships of neurons to non-neuronal cell types, as well as the
relative networks of segmented and quantified organelles within cells. NeuroSCAN could
be used to compare new datasets from genetic variants, from animals trained under
specific conditions or from additional developmental datasets across embryogenesis. As

such, the pipeline and design of NeuroSCAN can serve as a sandbox to examine the
value of the integration of datasets in exploring representations of neuronal relationships
across connectomes.

NeuroSCAN is one of several efforts centered around interpreting the *C. elegans*
vEM datasets, and forms part of a longer tradition that has leveraged the pioneering
datasets generated for *C. elegans* connectomes towards exploring structure-function
relationships in the nervous system. While the smaller scale of the *C. elegans* neuropil
allowed us to rigorously vet the utility of these approaches, we suggest that these same
methods would be beneficial in comparative studies in neuropils of other species,
including those with less stereotypically formed connectomes. We suggest that contact
profiles, along with neuron morphologies and synaptic partners, can act as 'fingerprints'
for individual neurons and neuron classes. These 'fingerprints' can be aligned across
animals of the same species to create identities for neurons. Frameworks for systematic
connectomics analysis in tractable model systems such as *C. elegans* are critical in laying
a foundation for future analyses in other organisms with up to a billion-fold increase in
neurons (Toga et al. 2012). Therefore, we envision these collective efforts akin to the
foundational work from *C. elegans* in pioneering genomic analysis and annotations ahead
of the Human Genome Project (Stein et al. 2001; Collins and Fink 1995). We believe that
further integration of datasets in platforms like NeuroSCAN would be key in determining
the representations and features necessary for the interpretation and analyses of other
connectomes.
488

Acknowledgements:

We are grateful for current and former members of the Colón-Ramos lab for their
guidance and suggestions, in particular, Agustín Almoril-Porras and Malcolm Díaz García
for assisting with data formatting, Patricia Chanabá-López and Andrea Cuentas-Condori
for feedback on the NeuroSCAN website, Mayra Blakey for administrative roles in

managing contracts for funding distribution, and Ben Clark and Milind Singh for feedback

on the paper. We also thank Stephen Larson, Dario Del Piano and Zoran Sinnema

(MetaCell) and Jamie Emerson (Bilte Co.) for website software development, method

reporting and hosting services. We thank Brandi Mattson for editing early paper drafts.

We acknowledge Ryan Christensen and Hari Shroff (Janelia Research Campus) and

Patrick La Riviere (University of Chicago) for helpful discussions and guidance for the

NeuroSCAN website. We thank the Research Center for Minority Institutions program,

the Marine Biological Laboratories (MBL), and the Instituto de Neurobiología de la

Universidad de Puerto Rico for providing meeting and brainstorming platforms. D.A.C-R.

acknowledges the Whitman Fellows program at MBL for providing funding and space for

discussions valuable to this work. Research in D.A.C-R. and W.A.M. labs was supported

by NIH grant R24-OD016474. This work was also funded by the NIH/NINDS grant R35

NS132156-01, DP1 NS111778 and R01 NS076558–2.

Authorship Contributions

N.L.K. Conceptualization; Data curation; Investigation; Methodology; Project
Administration; Validation; Visualization; Writing- Original Draft
S.E.E. Conceptualization; Data curation; Formal Analysis; Investigation; Project
Administration; Software; Visualization; Writing- Original Draft
D.B. Formal Analysis; Software; Writing - Original Draft
M.W.M. Conceptualization; Formal Analysis; Investigation; Project Administration;
Software; Writing- Review & Editing
P.A.-M. Data curation; Writing- Review & Editing
N.V.M. Data curation; Investigation; Writing - Review & Editing
S.K. Resources; Supervision

W.A.M. Conceptualization; Data curation; Formal Analysis; Funding Acquisition;
Methodology; Resources; Software; Supervision; Validation; Writing- Review & editing;
Corresponding Author
D.A.C.-R. Conceptualization; Funding Acquisition; Resources; Supervision;
Visualization; Writing - Review & editing; Corresponding Author
525
526

Competing Interests

528
529 Authors do not declare any competing interests.
530

Declaration of generative AI and AI-assisted technologies in the writing process.

533 During the preparation of this work the author(s) used ChatGPT in order to improve
534 readability. After using this tool, the author(s) reviewed and edited the content as
535 needed and take full responsibility for the content of the published article.
536
537

Bibliography

Altun, Z.F., Herndon, L.A., Wolkow, C.A., Crocker, C., Lints, R., and Hall, D.H. (ed.s). 2002.
"WormAtlas." 2024 2002. http://www.wormatlas.org.
Azevedo, Anthony. 2024. "Connectomic Reconstruction of a Female Drosophila Ventral Nerve
Cord | Nature." June 26, 2024. https://www.nature.com/articles/s41586-024-07389-x.
Barabási, Dániel L., Ginestra Bianconi, Ed Bullmore, Mark Burgess, SueYeon Chung, Tina Eliassi-
Rad, Dileep George, et al. 2023. "Neuroscience Needs Network Science." *The Journal of*
Neuroscience: The Official Journal of the Society for Neuroscience 43 (34): 5989–95.
https://doi.org/10.1523/JNEUROSCI.1014-23.2023.
Berger, Daniel R., H. Sebastian Seung, and Jeff W. Lichtman. 2018. "VAST (Volume Annotation
and Segmentation Tool): Efficient Manual and Semi-Automatic Labeling of Large 3D
Image Stacks." *Frontiers in Neural Circuits* 12. https://doi.org/10.3389/fncir.2018.00088.
Brittin, C. A., S. J. Cook, D. H. Hall, S. W. Emmons, and N. Cohen. 2021. "A Multi-Scale Brain Map
Derived from Whole-Brain Volumetric Reconstructions." *Nature* 591 (7848): 105–10.
https://doi.org/10.1038/s41586-021-03284-x.

Brittin, C. A., Steven J. Cook, David H. Hall, Scott W. Emmons, and Netta Cohen. 2018.
"Volumetric Reconstruction of Main Caenorhabditis Elegans Neuropil at Two Different
Time Points." *bioRxiv*, 485771. https://doi.org/10.1101/485771.
Brugnone, N., A. Gonopolskiy, M. W. Moyle, M. Kuchroo, D. van Dijk, K. R. Moon, D. Colon-
Ramos, G. Wolf, M. J. Hirn, and S. Krishnaswamy. 2019. "Coarse Graining of Data via
Inhomogeneous Diffusion Condensation." *Proc IEEE Int Conf Big Data* 2019
(December):2624–33. https://doi.org/10.1109/BigData47090.2019.9006013.
Burns, Randal, William Gray Roncal, Dean Kleissas, Kunal Lillaney, Priya Manavalan, Eric
Perlman, Daniel R. Berger, et al. 2013. "The Open Connectome Project Data Cluster:
Scalable Analysis and Vision for High-Throughput Neuroscience." *Scientific and
Statistical Database Management : International Conference, SSDBM ... : Proceedings.
International Conference on Scientific and Statistical Database Management*.
https://doi.org/10.1145/2484838.2484870.
Cantarelli, Matteo, Boris Marin, Adrian Quintana, Matt Earnshaw, Robert Court, Padraig
Gleeson, Salvador Dura-Bernal, R. Angus Silver, and Giovanni Idili. 2018. "Geppetto: A
Reusable Modular Open Platform for Exploring Neuroscience Data and Models."
Philosophical Transactions of the Royal Society of London. Series B, Biological Sciences
373 (1758). https://doi.org/10.1098/rstb.2017.0380.
Collins, Francis S., and Leslie Fink. 1995. "The Human Genome Project." *Alcohol Health and
Research World* 19 (3): 190–95.
Collinson, Lucy M., Carles Bosch, Anwen Bullen, Jemima J. Burden, Raffaella Carzaniga, Cheng
Cheng, Michele C. Darrow, et al. 2023. "Volume EM: A Quiet Revolution Takes Shape."
Nature Methods 20 (6): 777–82. https://doi.org/10.1038/s41592-023-01861-8.
Colón-Ramos, Daniel A., Milica A. Margeta, and Kang Shen. 2007. "Glia Promote Local
Synaptogenesis through UNC-6 (Netrin) Signaling in C. Elegans." *Science (New York, N.Y.)*
318 (5847): 103–6. https://doi.org/10.1126/science.1143762.
Cook, Steven J., Travis A. Jarrell, Christopher A. Brittin, Yi Wang, Adam E. Bloniarz, Maksim A.
Yakovlev, Ken C. Q. Nguyen, et al. 2019. "Whole-Animal Connectomes of Both
Caenorhabditis Elegans Sexes." *Nature* 571 (7763): 63–71.
https://doi.org/10.1038/s41586-019-1352-7.
Dorkenwald, Sven, Arie Matsliah, Amy R Sterling, Philipp Schlegel, Szi-chieh Yu, Claire E.
McKellar, Albert Lin, et al. 2023. "Neuronal Wiring Diagram of an Adult Brain." *bioRxiv*,
January, 2023.06.27.546656. https://doi.org/10.1101/2023.06.27.546656.
Duncan, Leighton H., Mark W. Moyle, Lin Shao, Titas Sengupta, Richard Ikegami, Abhishek
Kumar, Min Guo, et al. 2019. "Isotropic Light-Sheet Microscopy and Automated Cell
Lineage Analyses to Catalogue Caenorhabditis Elegans Embryogenesis with Subcellular
Resolution." *JoVE*, no. 148 (June), e59533. https://doi.org/10.3791/59533.
Eberle, Anna Lena, and Dirk Zeidler. 2018. "Multi-Beam Scanning Electron Microscopy for High-
Throughput Imaging in Connectomics Research." *Frontiers in Neuroanatomy* 12
(December). https://doi.org/10.3389/fnana.2018.00112.
Feinberg, Evan H., Miri K. Vanhoven, Andres Bendesky, George Wang, Richard D. Fetter, Kang
Shen, and Cornelia I. Bargmann. 2008. "GFP Reconstitution Across Synaptic Partners

(GRASP) Defines Cell Contacts and Synapses in Living Nervous Systems." *Neuron* 57 (3):
353–63. https://doi.org/10.1016/j.neuron.2007.11.030.
Fenyves, Bánk G., Gábor S. Szilágyi, Zsolt Vassy, Csaba Sőti, and Peter Csermely. 2020. "Synaptic
Polarity and Sign-Balance Prediction Using Gene Expression Data in the Caenorhabditis
Elegans Chemical Synapse Neuronal Connectome Network." *PLOS Computational*
Biology 16 (12): e1007974. https://doi.org/10.1371/journal.pcbi.1007974.
Galili, Dana S, Gregory SXE Jefferis, and Marta Costa. 2022. "Connectomics and the Neural Basis
of Behaviour." *Current Opinion in Insect Science* 54 (December):100968.
https://doi.org/10.1016/j.cois.2022.100968.
Kasthuri, Narayanan, Kenneth Jeffrey Hayworth, Daniel Raimund Berger, Richard Lee Schalek,
José Angel Conchello, Seymour Knowles-Barley, Dongil Lee, et al. 2015. "Saturated
Reconstruction of a Volume of Neocortex." *Cell* 162 (3): 648–61.
https://doi.org/10.1016/j.cell.2015.06.054.
Lichtman, Jeff W., Hanspeter Pfister, and Nir Shavit. 2014. "The Big Data Challenges of
Connectomics." *Nature Neuroscience* 17 (11): 1448–54.
https://doi.org/10.1038/nn.3837.
Maclachlan, Catherine, Daniela A. Sahlender, Shuichi Hayashi, Zoltán Molnár, and Graham
Knott. 2018. "Block Face Scanning Electron Microscopy of Fluorescently Labeled Axons
Without Using Near Infra-Red Branding." *Frontiers in Neuroanatomy* 12.
https://doi.org/10.3389/fnana.2018.00088.
Marc, Robert E., Crystal Lynn Sigulinsky, Rebecca L. Pfeiffer, Daniel Emrich, James Russell
Anderson, and Bryan William Jones. 2018. "Heterocellular Coupling Between Amacrine
Cells and Ganglion Cells." *Frontiers in Neural Circuits* 12.
https://doi.org/10.3389/fncir.2018.00090.
Moon, Kevin R., David van Dijk, Zheng Wang, Scott Gigante, Daniel B. Burkhardt, William S.
Chen, Kristina Yim, et al. 2019. "Visualizing Structure and Transitions in High-
Dimensional Biological Data." *Nature Biotechnology* 37 (12): 1482–92.
https://doi.org/10.1038/s41587-019-0336-3.
Motta, Alessandro, Manuel Berning, Kevin M. Boergens, Benedikt Staffler, Marcel Beining, Sahil
Loomba, Philipp Hennig, Heiko Wissler, and Moritz Helmstaedter. 2019. "Dense
Connectomic Reconstruction in Layer 4 of the Somatosensory Cortex." *Science* 366
(6469): eaay3134. https://doi.org/10.1126/science.aay3134.
Moyle, M. W., K. M. Barnes, M. Kuchroo, A. Gonopolskiy, L. H. Duncan, T. Sengupta, L. Shao, et
al. 2021. "Structural and Developmental Principles of Neuropil Assembly in C. Elegans."
Nature 591 (7848): 99–104. https://doi.org/10.1038/s41586-020-03169-5.
Packer, Jonathan S., Qin Zhu, Chau Huynh, Priya Sivaramakrishnan, Elicia Preston, Hannah
Dueck, Derek Stefanik, et al. 2019. "A Lineage-Resolved Molecular Atlas of C. Elegans
Embryogenesis at Single-Cell Resolution." *Science (New York, N.Y.)* 365 (6459).
https://doi.org/10.1126/science.aax1971.
Perez, Alex J., Mojtaba Seyedhosseini, Thomas J. Deerinck, Eric A. Bushong, Satchidananda
Panda, Tolga Tasdizen, and Mark H. Ellisman. 2014. "A Workflow for the Automatic
Segmentation of Organelles in Electron Microscopy Image Stacks." *Frontiers in*
Neuroanatomy 8 (November). https://doi.org/10.3389/fnana.2014.00126.

Phelps, Jasper S., David Grant Colburn Hildebrand, Brett J. Graham, Aaron T. Kuan, Logan A.
Thomas, Tri M. Nguyen, Julia Buhmann, et al. 2021. "Reconstruction of Motor Control
Circuits in Adult Drosophila Using Automated Transmission Electron Microscopy." *Cell*
184 (3): 759-774.e18. https://doi.org/10.1016/j.cell.2020.12.013.
Pipkin, Jason E., Eric Allen Bushong, Mark H. Ellisman, and William B. Kristan Jr. 2018.
"Verifying, Challenging, and Discovering New Synapses Among Fully EM-Reconstructed
Neurons in the Leech Ganglion." *Frontiers in Neuroanatomy* 12 (November).
https://doi.org/10.3389/fnana.2018.00095.
Randel, Nadine, Albina Asadulina, Luis A. Bezares-Calderón, Csaba Verasztó, Elizabeth A.
Williams, Markus Conzelmann, Réza Shahidi, and Gáspár Jékely. 2014. "Neuronal
Connectome of a Sensory-Motor Circuit for Visual Navigation." eLife. eLife Sciences
Publications Limited. May 27, 2014. https://doi.org/10.7554/eLife.02730.
Randi, Francesco, Anuj K. Sharma, Sophie Dvali, and Andrew M. Leifer. 2023. "Neural Signal
Propagation Atlas of Caenorhabditis Elegans." *Nature* 623 (7986): 406–14.
https://doi.org/10.1038/s41586-023-06683-4.
Rapti, Georgia, Chang Li, Alan Shan, Yun Lu, and Shai Shaham. 2017. "Glia Initiate Brain
Assembly through Noncanonical Chimaerin-Furin Axon Guidance in C. Elegans." *Nature*
Neuroscience 20 (10): 1350–60. https://doi.org/10.1038/nn.4630.
Rivlin, Patricia K., Michal Januszewski, Kit D. Longden, Erika Neace, Louis K. Scheffer,
Christopher Ordish, Jody Clements, et al. 2024. "Connectomic Analysis of Mitochondria
in the Central Brain of Drosophila." bioRxiv.
https://doi.org/10.1101/2024.04.21.590464.
Ryan, Kerrianne, Zhiyuan Lu, and Ian A Meinertzhagen. 2016. "The CNS Connectome of a
Tadpole Larva of Ciona Intestinalis (L.) Highlights Sidedness in the Brain of a Chordate
Sibling." Edited by Eve Marder. *eLife* 5 (December):e16962.
https://doi.org/10.7554/eLife.16962.
Scheffer, Louis K, C Shan Xu, Michal Januszewski, Zhiyuan Lu, Shin-ya Takemura, Kenneth J
Hayworth, Gary B Huang, et al. 2020. "A Connectome and Analysis of the Adult
Drosophila Central Brain." Edited by Eve Marder, Michael B Eisen, Jason Pipkin, and
Chris Q Doe. *eLife* 9 (September):e57443. https://doi.org/10.7554/eLife.57443.
Schmid, Benjamin, Johannes Schindelin, Albert Cardona, Mark Longair, and Martin Heisenberg.
2010. "A High-Level 3D Visualization API for Java and ImageJ." *BMC Bioinformatics* 11
(1): 274. https://doi.org/10.1186/1471-2105-11-274.
Shapson-Coe, Alexander, Michał Januszewski, Daniel R. Berger, Art Pope, Yuelong Wu, Tim
Blakely, Richard L. Schalek, et al. 2021. "A Connectomic Study of a Petascale Fragment of
Human Cerebral Cortex." *bioRxiv*, January, 2021.05.29.446289.
https://doi.org/10.1101/2021.05.29.446289.
Stein, L., P. Sternberg, R. Durbin, J. Thierry-Mieg, and J. Spieth. 2001. "WormBase: Network
Access to the Genome and Biology of Caenorhabditis Elegans." *Nucleic Acids Research*
29 (1): 82–86. https://doi.org/10.1093/nar/29.1.82.
Sun, Haosheng, and Oliver Hobert. 2023. "Temporal Transitions in the Postembryonic Nervous
System of the Nematode Caenorhabditis Elegans: Recent Insights and Open Questions."

Special Issue: Temporal Patterning in the CNS 142 (June):67–80. https://doi.org/10.1016/j.semcdb.2022.05.029.

Svara, Fabian, Dominique Förster, Fumi Kubo, Michał Januszewski, Marco dal Maschio, Philipp J. Schubert, Jörgen Kornfeld, et al. 2022. "Automated Synapse-Level Reconstruction of Neural Circuits in the Larval Zebrafish Brain." *Nature Methods* 19 (11): 1357–66. https://doi.org/10.1038/s41592-022-01621-0.

Swanson, Larry W., and Jeff W. Lichtman. 2016. "From Cajal to Connectome and Beyond." *Annual Review of Neuroscience*. Annual Reviews. https://doi.org/10.1146/annurev-neuro-071714-033954.

Takemura, S. Y., A. Bharioke, Z. Lu, A. Nern, S. Vitaladevuni, P. K. Rivlin, W. T. Katz, et al. 2013. "A Visual Motion Detection Circuit Suggested by Drosophila Connectomics." *Nature* 500 (7461): 175–81. https://doi.org/10.1038/nature12450.

Takemura, Shin-ya, Yoshinori Aso, Toshihide Hige, Allan Wong, Zhiyuan Lu, C Shan Xu, Patricia K Rivlin, et al. 2017. "A Connectome of a Learning and Memory Center in the Adult Drosophila Brain." *eLife* 6 (July):e26975. https://doi.org/10.7554/eLife.26975.

Takemura, Shin-ya, Aljoscha Nern, Dmitri B. Chklovskii, Louis K. Scheffer, Gerald M. Rubin, and Ian A. Meinertzhagen. 2017. "The Comprehensive Connectome of a Neural Substrate for 'ON' Motion Detection in Drosophila." Edited by Alexander Borst. *Elife* 6 (April):e24394. https://doi.org/10.7554/eLife.24394.

Takemura, Shin-ya, C. Shan Xu, Zhiyuan Lu, Patricia K. Rivlin, Toufiq Parag, Donald J. Olbris, Stephen Plaza, et al. 2015. "Synaptic Circuits and Their Variations within Different Columns in the Visual System of Drosophila." *Proceedings of the National Academy of Sciences of the United States of America* 112 (44): 13711–16. https://doi.org/10.1073/pnas.1509820112.

Taylor, Seth R., Gabriel Santpere, Alexis Weinreb, Alec Barrett, Molly B. Reilly, Chuan Xu, Erdem Varol, et al. 2021. "Molecular Topography of an Entire Nervous System." *Cell* 184 (16): 4329-4347.e23. https://doi.org/10.1016/j.cell.2021.06.023.

The MICrONS Consortium, The MICrONS consortium, J. Alexander Bae, Mahaly Baptiste, Caitlyn A. Bishop, Agnes L. Bodor, Derrick Brittain, JoAnn Buchanan, et al. 2023. "Functional Connectomics Spanning Multiple Areas of Mouse Visual Cortex." bioRxiv. https://doi.org/10.1101/2021.07.28.454025.

Toga, Arthur W., Kristi A. Clark, Paul M. Thompson, David W. Shattuck, and John Darrell Van Horn. 2012. "Mapping the Human Connectome." *Neurosurgery* 71 (1): 1–5. https://doi.org/10.1227/NEU.0b013e318258e9ff.

Vishwanathan, Ashwin, Kayvon Daie, Alexandro D. Ramirez, Jeff W. Lichtman, Emre R. F. Aksay, and H. Sebastian Seung. 2017. "Electron Microscopic Reconstruction of Functionally Identified Cells in a Neural Integrator." *Current Biology: CB* 27 (14): 2137-2147.e3. https://doi.org/10.1016/j.cub.2017.06.028.

Wang, Chen, Berta Vidal, Surojit Sural, Curtis Loer, G. Robert Aguilar, Daniel M. Merritt, Itai Antoine Toker, Merly C. Vogt, Cyril Cros, and Oliver Hobert. 2023. "A Neurotransmitter Atlas ofC. Elegansmales and Hermaphrodites," December. https://doi.org/10.1101/2023.12.24.573258.

Wanner, Adrian A., Christel Genoud, and Rainer W. Friedrich. 2016. "3-Dimensional Electron
Microscopic Imaging of the Zebrafish Olfactory Bulb and Dense Reconstruction of
Neurons." *Scientific Data* 3 (1): 160100. https://doi.org/10.1038/sdata.2016.100.
White, John Graham, Eileen Southgate, J. N. Thomson, and Sydney Brenner. 1986. "The
Structure of the Nervous System of the Nematode *Caenorhabditis Elegans*."
Philosophical Transactions of the Royal Society of London. B, Biological Sciences 314
(1165): 1–340. https://doi.org/doi:10.1098/rstb.1986.0056.
Witvliet, D., B. Mulcahy, J. K. Mitchell, Y. Meirovitch, D. R. Berger, Y. Wu, Y. Liu, et al. 2021.
"Connectomes across Development Reveal Principles of Brain Maturation." *Nature* 596
(7871): 257–61. https://doi.org/10.1038/s41586-021-03778-8.
Xu, C Shan, Kenneth J Hayworth, Zhiyuan Lu, Patricia Grob, Ahmed M Hassan, José G García-
Cerdán, Krishna K Niyogi, Eva Nogales, Richard J Weinberg, and Harald F Hess. 2017.
"Enhanced FIB-SEM Systems for Large-Volume 3D Imaging." Edited by Jeremy Nathans.
eLife 6 (May):e25916. https://doi.org/10.7554/eLife.25916.
Yim, Hyunsoo, Daniel T. Choe, J. Alexander Bae, Myung-kyu Choi, Hae-Mook Kang, Ken C. Q.
Nguyen, Soungyub Ahn, et al. 2024. "Comparative Connectomics of Dauer Reveals
Developmental Plasticity." *Nature Communications* 15 (1): 1546.
https://doi.org/10.1038/s41467-024-45943-3.
Zheng, Z., J. S. Lauritzen, E. Perlman, C. G. Robinson, M. Nichols, D. Milkie, O. Torrens, et al.
2018. "A Complete Electron Microscopy Volume of the Brain of Adult Drosophila
Melanogaster." *Cell* 174 (3): 730-743.e22. https://doi.org/10.1016/j.cell.2018.06.019.
744

Figures

Figure 1. DC/C-PHATE representations of contactome-based relationships. DC/C-PHATE graphs enable representations of neuronal contact relationships. To build DC/C-PHATE graphs we **(A)** analyzed serial section EM datasets of the *C. elegans* nerve ring neuropil (located in the head of the animal). **(B)** Single cross section of the nerve ring (surrounding the pharynx), with segmented neurites pseudo-colored. Dark box corresponds to the zoomed-in image in (C). The cross section is from the JSH dataset digitally segmented (Brittin et al. 2021). **(C)** Zoom-in cross section with three arbitrary neurons (called A, B, C) highlighted by overlaying opaque cartoon (2-D, left image) and 3-D shapes (middle image) to represent the segmentation process and the neuronal contact sites (highlighted Yellow, Yellow dashed, Red). Contacts are quantified for all neuron pairs across the contactome (See Methods), to generate a Contact Matrix (represented here as a table, schematized for the three arbitrary neurons selected and in

which specific contact quantities are represented by a color scale and not numerical values). **(D)** Schematic of how the Diffusion Condensation algorithm (visualized with C-PHATE) works. DC/C-PHATE makes use of the contact matrix to group neurons based on similar adjacency profiles (Brugnone et al. 2019; 2019; Moyle et al. 2021), schematized here for the three neurons in (C). **(E)** Screenshot of the 3-D C-PHATE graph from a Larva stage 1 (L1; 0 hours post hatching;) contactome, with individual neurons represented as spheres at the periphery. Neurons were iteratively clustered towards the center, with the final iteration containing the nerve ring represented as a sphere in the center of the graph (Highlighted in maroon). **(F)** Integration in NeuroSCAN of the DC/C-PHATE and EM-derived 3-D neuron morphology representations allow users to point to each sphere in the graph and determine cellular or cluster identities for each iteration. Shown here and circled in Red, an arbitrarily selected cluster (in E), with the identities of the neurons belonging to that cluster (four letter codes in the column to the left of F) and the corresponding neuronal morphologies (right) of this group of neurons in the EM-reconstructed nerve ring (with individual neurons pseudo-colored ccording to their names to the left). Compass: Anterior (A), Posterior (P), Dorsal (D), Ventral (V), Left (L), Right (R).

778
779
Figure 2. Implementation of DC/C-PHATE to developmental contactomes reveal a
conserved layered organization maintained during post-embryonic growth. (A)
Cartoon of the *C. elegans* head and nerve ring (outlined with black box). Below, nerve
ring reconstruction from EM data of an L1 animal (5 hours post hatching), with all
neurons in gray. Scale bar 2 um. **(B-F)** DC/C-PHATE plots generated for available
contactomes across *C. elegans* larval development, colored by stratum identity as
described (Moyle et al. 2021). Individual neurons are located at the edges of the graph
and condense centrally. The four super-clusters identified and all iterations before are
colored accordingly. The identity of the individual neurons belonging to each stratum,
and at each larval stage, were largely preserved, and are provided in (Tables S3-S6).
Some datasets contain 5 or 6 super-clusters (colored dark purple, yellow and orange),
which are classified as groups of neurons that are differentially categorized across the
developmental connectomes. **(G-K)** Volumetric reconstruction of the *C. elegans*
neuropil (from EM serial sections for the indicated larval stages (columns)) with the
neurons colored based on their strata identity. Scale bar 2 um; Anterior (A) left, Dorsal
(D) up.
796

797
798

Figure 3. Examination of the architectural motifs underlying the distinct strata across development. Visualization of **(A-F)** Stratum 1 (Red) and **(G-L)** Strata 3 and 4 (Blue and Green) reveal motifs that are preserved (Strata 1) and change (Strata 3 and 4) across developmental contactomes (L1 to Adult, left to right, as indicated by labels on top). **(B-F)** Cropped view of Stratum 1 at each developmental stage showing a similar shape of two 'horn-like' clusters in the C-PHATE graphs. These two clusters have similar neuronal memberships, which are largely invariant across developmental contactomes (Table S3). **(H-L)** Cropped view of Strata 3 and 4 at each developmental stage highlighting differences in the organization and number of neurons contained in each of the Blue and Green strata, which is particularly distinct when comparing (H) L1 and (K) L4 (Tables S5 and S6). There is an additional supercluster (Yellow in (I-J)) at stages L2 and L3 that contains neurons of S3 and S4 identity.

Figure 4. Case study: AIML and PVQL neurons change clustering patterns across the developmental contactomes (A-E) C-PHATE plots across development, with the trajectories of AIM neurons (in purple) and the rest of the spheres colored by stratum identity (see Figure 2). **(F-G)** Zoom in of the AIM, PVQ, and AVF trajectories corresponding to Larval Stage 1 (A, dotted box) and in (G), Larval Stage 3 (C, dashed box). Note how the relationship between AIM and PVQ neurons in the C-PHATE graph varies for each of the examined contactomes across development, as seen by the iterations before co-clustering (Figure S1, Table S7).

826
827
Figure 5. Case Study: Visualization of contact profiles in individual neurons. (A)
Cartoon schematic of the head of the animal with the AIM neurons (Purple) and
Pharynx (gray), and (dotted box) a 3-D reconstruction of the AIM neuron morphology
from the L1 (0 hours post hatching) dataset. **(B)** Zoom-in of the simplified DC/C-PHATE
clustering of the AIM (Purple), PVQ (Orange) and AVF (Green) neurons for the
contactome of an L3 animal. **(C)** 3-D representation of all contacts onto the AIM neuron
morphology in an L1 animal, and colored based on contacting partner identity as
labeled (right) in the detailed inset (black box) region. **(D)** AIM-PVQ contacts (in orange)
and AIM-AVF contacts (in green), projected onto the AIM neurons (light purple) across
developmental stages and augmented for the figure (see non-augmented contacts in
Figure S3). Scale bar 2 um.
839
840

841
Figure 6. Case study: Segmented morphologies of AIM, PVQ and AVF across
larval development. (A) Cartoon schematic of the *C. elegans* head, pharynx (gray)
and examined neurons with dashed black box representing the nerve ring region. **(B)**
Schematic representation of the outgrowth path of the AVF neurons as observed by EM
(Witvliet et al. 2021). AVFL and AVFR (green) grow along the AIML neuron (purple)
onto the AIMR neurite. The distal end of the AVF neurite is highlighted with a black
arrowhead in the schematic. **(C)** Neuronal morphologies of AIM (purple), PVQ (orange),
AVF (green) across post embryonic development, as indicated, with black arrowhead
pointing to AVF outgrowth. Scale bar = 2um. Regions for insets (L1, dotted box; L2,
dashed box) correspond to (D). **(D)** Morphologies of these neurons (rotated to the
posterior view) display the AVF neurons' positions between the AIM and PVQ neurons
at the L2 stage. Indicated outgrowth between neurons continues to the Adult stage
(Video S2). Note how AVF outgrowth alters contact between PVQ and AIM (Figure 5D).

Figure 7. Case study: AIM-PVQ and AIM-AVF synaptic positions across development. (A) AIM-PVQ synaptic sites (dark orange arrowheads) and AIM-AVF synaptic sites (dark green arrowheads) in the segmented AIM neurons and reconstructed across post embryonic development from original connectomics data. Scale bar = 2um. **(B)** Schematic of the AIM, PVQ and AVF circuitry across development based on synaptic connectivity and focusing on the stage before AVF outgrowth (L1), during AVF outgrowth (L2) and Adult; arrow direction indicates pre to post synaptic connection, and arrow thickness indicates relative number of synaptic sites (finest, <5 synapses; medium, 5-10 synapses; thickest, 11-30 synapses). **(C)** Zoom in of synaptic sites (green) in the Adult connectome and embedded into the AIM neuron morphology (light purple). In NeuroSCAN, presynaptic sites are displayed as blocks and postsynaptic sites as spheres, and a scaling factor is applied to the 3-D models (Methods).

871
Figure 8. NeuroSCAN is a tool that enables integrated comparisons of neuronal
relationships across development. With NeuroSCAN, users have integrated access
to: C-PHATE plots, 3-D morphological renderings, neuronal contact sites and synaptic
representations. Through stage-specific C-PHATE renderings, users can explore
neuronal relationships from high dimensional contactome data. **(Top)** On C-PHATE
plots, schematized here, each sphere represents an individual neuron, like AVF or AIM,
or a group of neurons clustered together during algorithm iterations. **(Right)** 3D
renderings of AIM neurons (Purple), PVQ neurons (Orange), AVF neurons (Green) can
be visualized in the context of the entire nerve ring or other circuits (gray). **(Left)**
AIM:AVF contact sites (green) onto the AIM neuron (purple) with the AIM-AVF synaptic
sites (orange). Inset shows zoomed in of contacts and synapses- presynaptic sites
(blocks) postsynaptic sites (spheres). Data depicted here are from the L3 stage (27
hours post hatching).

Star Methods

Key Resources Table

Reagent or Resource	Source
Deposited data	
L1 (3), L2, L3	(Witvliet et al. 2021)
Electron Microscopy L4 (JSH), Adult (N2U)	(White et al. 1986)
Neuron Digital Segmentation L4 (JSH), Adult (N2U	(Brittin et al. 2021)
Synaptic coordinates L4 (JSH), Adult (N2U)	Cook et al. 2019
Adjacencies	(Moyle et al. 2021; Brittin et al. 2018); this paper
C-PHATE visualization	CytoSHOW.org; (Moyle et al. 2021; Moon et al. 2019); this paper
Software and Algorithms	
CytoSHOW	http://www.cytoshow.org/ https://github.com/mohler/CytoSHOW
Diffusion Condensation	(Brugnone et al. 2019), this paper
Adjacency algorithm	(Brittin et al. 2018)

RESOURCE AVAILABILITY

Lead Contact

Further information and requests can be directed to Daniel.colon-ramos@yale.edu.

Data and Code Availability

Figures in this article have been generated with NeuroSCAN (Figures 5D, Figures 6-7,

Figure S2G-I, Figure S3, Figure S4, Figure S5 A-B, Figure 8, Figures S6-S12, Videos S1-

S4) and CytoSHOW (Figures 1-4, Figure 5A and C, Figure S1, Figure S 5C). Data can

be visualized via the viewer at NeuroSCAN.net or by downloading glTF files from
NeuroSCAN and using a glTF viewer to visualize them. Additionally, the data generated
for NeuroSCAN is available in .OBJ file format (Brittin et al. 2018) (and can be visualized
from a local hard drive with CytoSHOW (http://neuroscan.cytoshow.org/). All excel files
for Diffusion Condensation iterations and adjacency quantifications can be found in
Tables S3- S13. Tutorials for NeuroSCAN are available on NeuroSCAN.net upon opening
the website, within the main menu of the website (Figure S8), and in the supplementary
materials (Figure S5-S12; Videos S1 and S3-S4). These tutorials generally cover the
process of engaging in analysis at and across specific developmental stages by filtering
the data items and adding items to viewers (Figure S10). General understanding for how
to use C-PHATE to analyze neuronal relationships can be found in Figure 1, Figure 4,
Figure S6, Video S1, and in our previous publication (Moyle et al. 2021). For additional
information on filters and in-viewer changes to the data (colors, developmental stages,
downloading data) see Figure S5, Figure S7, Figure S11, Figure S12, and Videos S3-S4.
All code for website development is available at Github
(https://github.com/colonramoslab/NeuroSCAN) and for information on website
architecture and data model see Figures S13-S14.
918
919

920   EXPERIMENTAL MODEL AND SUBJECT DETAILS
921

Volume electron microscopy (vEM) data and segmentation of neurons and synapses from
(Witvliet et al. 2021; White et al. 1986; Brittin et al. 2018; Cook et al. 2019) . We analyzed
available EM datasets that were transversely sectioned and segmented (Witvliet et al.
2021; Brittin et al. 2021; White et al. 1986). We deleted the CAN neurons in the L1-L3
datasets to keep these datasets consistent with the legacy datasets L4 and Adult (N2U),
which do not contain CAN neurons (as in Moyle et al. 2021). The L1-L3 datasets do
contain CAN neurons, however their contacts are minimal.
929
930

931   METHOD DETAILS

All 3-D object isosurfaces (Morphologies (Neurons), Contacts, Synapses, C-PHATE plots) were generated from segmented EM datasets using a modified version of the ImageJ 3D viewer plug-in (Schmid et al. 2010) implemented in CytoSHOW (cytoshow.org). This tool employs the marching cubes algorithm for polygon-generation. All 3-D objects are first exported as wavefront (.OBJ) files then converted to GL Transmission Format (.glTF) file format which does not distort the resolution but compacts the file information to enable faster loading times in the web-based 3-D viewer.

Pixel Threshold Distance for Adjacency Profiles and Contacts

We identified two challenges in compiling Electron Microscopy (EM) datasets for comparisons: 1) how to uniformly capture neuronal relationships based on areas of physical adjacency (contact) across datasets that have differences in volume depth and in x-y-z resolutions, and 2) how to standardize across datasets in which membrane boundaries had been called using a variety of methods, including contrast methods and segmentation methods (hand-drawn vs predicted via centroid node expansion by a shallow convolutional neural network) (Witvliet et al. 2021; Brittin et al. 2018; White et al. 1986). To address this, we first standardized the region of the neuropil across all developmental stages as in (Moyle et al. 2021). Briefly, all cell bodies were deleted, and we used the entry of the nerve ring neurons into the ventral cord as the posterior boundary landmark for the entire volume, focusing on the AIY Zone 2(Colón-Ramos, Margeta, and Shen 2007); slice range Table S1). Previously reported adjacency profiles used 10 pixels (or 45 nm) as the pixel threshold distance for the L4 (JSH) and Adult (N2U) datasets (Moyle et al. 2021). To account for differences in resolution (x-y axis) and in calling membrane boundaries between the L4 and Adult datasets and L1-L3 datasets, we designed a protocol to define the pixel threshold for each dataset. In short, for two cells that are in direct contact (Figure S2 D) in the manually segmented datasets (L4 and Adult), we calculated the length of overlap needed to reach from the segmented edge of one cell, across the membrane, and into the adjacent cell, when the segmented area of one cell is expanded by 45 nm (10 pixels). This results in an average overlap of 30 nm for directly contacting cells in the L4 dataset. Then, in each computationally segmented

dataset (L1-L3), we empirically tested the distance (e.g. 55 nm, 60 nm, 62 nm) required

to achieve a similar overlap of 30 nm in direct contact cells. That empirical number (in

nm) was used for adjacency calculations and rendering of contacts. The numbers were

converted from nanometers into pixels to create a pixel threshold distance for each

dataset, and these are shown in Table S1.

Once these corrections had been applied, we calculated the cell-to-cell adjacency

scores for all cell pairs in each dataset by using the measure_adjacency algorithm from

https://github.com/cabrittin/volumetric_analysis; (Brittin et al. 2018) (Tables S8-S13).

Adjacency matrices were used for Diffusion condensation (Brugnone et al. 2019).

Diffusion Condensation

Diffusion condensation (DC) is a dynamic, time-inhomogeneous process designed to

create a sequence of multiscale data representations by condensing information over

time (Brugnone et al. 2019). The primary objective of this technique is to capture and

encode meaningful abstractions from high-dimensional data, facilitating tasks such as

manifold learning, denoising, clustering, and visualization. The underlying principle of

diffusion condensation is to iteratively apply diffusion operators that adapt to the evolving

data representation, effectively summarizing the data at multiple scales.

The diffusion condensation process begins with the initialization of an initial data

representation, typically the raw high-dimensional data or a preprocessed version. This

initial representation is used to construct a diffusion operator, a matrix derived from a

similarity matrix that reflects the local geometry of the data. The similarity metric, such as

Euclidean distance or cosine similarity, plays a crucial role in defining these local

relationships.

Once the initial diffusion operator is established, the algorithm proceeds to the

diffusion step. In this step, the diffusion operator is applied to the data, smoothing it by

spreading information along the edges of the similarity graph. This operation captures the

intrinsic geometry of the data while reducing noise. The specific form of the diffusion

operator, such as the heat kernel or graph Laplacian, significantly impacts how

information is propagated during this step.

Following the diffusion step, the condensation step updates the data
representation by aggregating diffused data points if the distance between them falls
below a 'merge threshold'. This step creates a more compact and abstract representation
of the data.

These diffusion and condensation steps are iteratively repeated. At each iteration,
the diffusion operator is recomputed based on the updated diffuse data representation,
ensuring that the process adapts to the evolving structure of the data. The iterations
continue until a stopping criterion is met, such as convergence of the data representation
to a single point.

The output of the diffusion condensation process is a sequence of multiscale data
representations. Each representation in this sequence captures the data at a different
level of abstraction, with earlier representations preserving more detailed information and
later representations providing more condensed summaries. This sequence of
representations can be utilized for various tasks, including manifold learning, denoising,
clustering, and visualization. By iteratively smoothing and condensing the data, diffusion
condensation reveals the underlying structure of high-dimensional datasets. A detailed
algorithm description is provided in Box 1 and Algorithm 1.

Diffusion Condensation

Initialization: Let $\mathbf{X} = \{x_1, x_2, \ldots, x_n\}$ be the set of n data points in a high-dimensional space. Construct the affinity matrix $\mathbf{A}$, where A_{ij} measures the similarity between x_i and x_j. Typically,

$$A_{ij} = \exp\left(-\frac{\|x_i - x_j\|^2}{2\sigma^2}\right)$$

for a chosen scale parameter σ.

Diffusion Operator: Define the degree matrix $\mathbf{D}$ as a diagonal matrix where $D_{ii} = \sum_j A_{ij}$. Construct the diffusion operator

$$\mathbf{P} = \mathbf{D}^{-1}\mathbf{A}$$

which normalizes the affinity matrix.

Diffusion Step: Apply the diffusion operator to the data:

$$\mathbf{Y} = \mathbf{P}\mathbf{X}$$

This step smooths the data, capturing the intrinsic geometry.

Condensation Step: After each diffusion step, merge data points that are within a small distance, ϵ, from each other to form a condensed representation. Specifically, data points x_i and x_j are merged if

$$\|x_i - x_j\| < \epsilon.$$

This merging process produces a set of condensed cluster centers $\mathbf{C} = \{c_1, c_2, \ldots, c_k\}$, where each center represents the mean of merged data points.

Iteration: Repeat the diffusion and condensation steps, adjusting the parameter σ adaptively, until convergence or for a predefined number of iterations.

1011

1012   **Box 1:** Mathematical description of Diffusion Condensation

1013

1014

1: **Input:** Data matrix $\mathbf{X} = \{x_1, x_2, \ldots, x_n\} \in \mathbb{R}^{n \times d}$, number of iterations T, scale parameter σ, condensation threshold ϵ

2: **Output:** Condensed data matrix $\mathbf{X}_{\text{condensed}}$

3: **Initialize:** Construct affinity matrix $\mathbf{A}$, degree matrix $\mathbf{D}$, and diffusion operator $\mathbf{P}$

4: $\mathbf{A}_{ij} \leftarrow \exp\left(-\frac{\|x_i - x_j\|^2}{2\sigma^2}\right)$ for all i, j

5: $\mathbf{D} \leftarrow \text{diag}\left(\sum_j \mathbf{A}_{ij}\right)$

6: $\mathbf{P} \leftarrow \mathbf{D}^{-1}\mathbf{A}$

7: **for** iteration $= 1$ to T **do**

8:    **Diffusion Step:**
      $\mathbf{Y} \leftarrow \mathbf{PX}$

9:    **Condensation Step:**
      Merge data points x_i and x_j if $\|x_i - x_j\| < \epsilon$ to form condensed cluster centers $\mathbf{C} = \{c_1, c_2, \ldots, c_k\}$

10:    $\mathbf{X} \leftarrow \mathbf{C}$

11:    **Update:**

12:      $\mathbf{A}_{ij} \leftarrow \exp\left(-\frac{\|x_i - x_j\|^2}{2\sigma^2}\right)$ for all i, j

13:      $\mathbf{D} \leftarrow \text{diag}\left(\sum_j \mathbf{A}_{ij}\right)$

14:      $\mathbf{P} \leftarrow \mathbf{D}^{-1}\mathbf{A}$

15: **end for**

16: **Return:** $\mathbf{X}_{\text{condensed}} \leftarrow \mathbf{X}$

1015

1016   **Algorithm 1:** Pseudocode for Diffusion Condensation

1017

1018 **C-PHATE**

1019

C-PHATE is an extension of the PHATE technique (Moon et al. 2019) which is specifically

aimed at handling and visualizing high-dimensional biological data. C-PHATE is

specifically designed to handle compositional data, which are datasets where the

components represent parts of a whole and are inherently constrained. It learns the

intrinsic manifold of the data, effectively capturing non-linear relationships and structures

that are not apparent with traditional methods like PCA or t-SNE.

The C-PHATE algorithm starts by loading affinity matrices associated with specific

clusterings obtained from diffusion condensation. These matrices are normalized to

generate kernel matrices that emphasize the strength of connections within each cluster.

The algorithm then builds a connectivity matrix by integrating these kernel matrices based

on cluster assignments over multiple time points. This is achieved by first initializing the

matrix with kernel matrices along its diagonal and then filling in off-diagonal blocks with

transition probabilities that reflect how clusters transition from one time point to the next.

Next, we apply the PHATE dimensionality reduction technique to the connectivity
matrix to generate 3D embeddings of the data. These embeddings are derived from
multiple iterations of diffusion condensation, capturing the geometry of the data at various
levels of granularity. The resulting coordinates are saved for subsequent analysis.

The final step involves visualizing the PHATE results in a 3D graphics tool,
CytoSHOW (Java-based; CytoSHOW.org; https://github.com/mohler/CytoSHOW; (Moyle
et al. 2021). The results are plotted in a 3D environment, with functionality enabling
rollover labels to display information about clustered cells. This requires cross-referencing
output tables from the original data collection. CytoSHOW is an interactive tool that allows
for assigning colors and annotations to individual neurons and clusters of interest. A
detailed algorithm description is provided in Box 2 and Algorithm 2.

1: **Input:** Output of the Diffusion Condensation algoprithm, number of iterations T
2: **Output:** Low-dimensional embedding $\mathbf{Y}$
3: **Initialize:** Load affinity matrices and cluster assignments from diffusion condensation output
4: **for** $t = 1$ to T **do**
5:    Load affinity matrix $\mathbf{A}_t$ from file
6:    Compute degree matrix $\mathbf{D}_t$ where $D_{t_{ii}} = \sum_j A_{t_{ij}}$
7:    Normalize to get kernel matrix $\mathbf{K}_t = \mathbf{D}_t^{-1/2} \mathbf{A}_t \mathbf{D}_t^{-1/2}$
8: **end for**
9: $\mathbf{C}_{\text{PHATE}} \leftarrow$ zero matrix with kernel matrices $\mathbf{K}_t$ along the diagonal
10: **for** $t = 1$ to $T - 1$ **do**
11:    Compute transition probabilities matrix $\mathbf{P}_{t,t+1}$ for clusters from time t to $t+1$
12:    $\mathbf{P}_{t,t+1}[i,j] \leftarrow \frac{\text{Number of points moving from cluster } i \text{ to cluster } j}{\text{Total number of points in cluster } i \text{ at time } t}$
13:    Update off-diagonal blocks of $\mathbf{C}_{\text{PHATE}}$ based on $\mathbf{P}_{t,t+1}$
14: **end for**
15: Compute the PHATE embedding $\mathbf{Y}$ from $\mathbf{C}_{\text{PHATE}}$
16: **Return:** $\mathbf{Y}$

1044

1045   **Algorithm 2:** Pseudocode for C-PHATE

C-PHATE

Given n data points, $\mathbf{X} = \{x_1, x_2, \ldots, x_n\}$, and the diffusion condensation output, consisting of $\mathbf{C}_t = \{c_1, c_2, \ldots, c_k\}$ denoting the merged data points and $\mathbf{A}_t$ denoting the affinity matrix at iteration t.

Kernel Matrix: For each iteration, t, compute the degree matrix $\mathbf{D}$, where $D_{ii} = \sum_j A_{ij}$. Then, normalize the affinity matrix to construct the kernel matrix $\mathbf{K}_t$:

$$\mathbf{K}_t = \mathbf{D}^{-1/2}\mathbf{A}_t\mathbf{D}^{-1/2}$$

Initial Connectivity Matrix: Initialize the connectivity matrix $\mathbf{C}_{\text{PHATE}}$ with zeros. Next, populate it with the kernel matrices, $\mathbf{K}_t$, along its diagonal, reflecting self-connections within each cluster at each time point.

Update Transition Probabilities: For each pair of adjacent time points t and $t + 1$, compute a transition probability matrix to determine how points transition between clusters C_t and C_{t+1}. Each entry p_{ij} in this matrix represents the probability of moving from cluster i at time t to cluster j at time $t+1$. p_{ij} is calculated by counting the number of points moving from cluster i to cluster j and normalizing by the total number of points in cluster i at time t. This can be expressed as:

$$p_{ij} = \frac{\text{Number of points moving from } i \text{ to } j}{\text{Total number of points in cluster } i \text{ at time } t}$$

Use these transition probabilities to populate the off-diagonal blocks of $\mathbf{C}_{\text{PHATE}}$

Dimensionality Reduction: Apply the PHATE algorithm to the final connectivity matrix $\mathbf{C}_{\text{PHATE}}$ to obtain the low-dimensional embedding $\mathbf{Y}$:

$$\mathbf{Y} = \text{PHATE}(\mathbf{C}_{\text{PHATE}})$$

Visualization: Visualize low-dimensional embedding $\mathbf{Y}$ in CytoSHOW.

1046

Box 2: Mathematical description of C-PHATE

The python code for C-PHATE allows for user specification of four numerical

parameters within the command line, and we used the same set of values for all C-PHATE

plots shown in this report (100, 30, 50, 1). The first two integers define the weighting of

connectivity between the current condensation step t and previous steps t-1 (weighting =

100) or t-2 (30), respectively, during construction of the connectivity matrix. Values 100

and 30 consistently resulted in a series of plotted clustering trajectories that form a dome-

like convergence of paths, enhancing our visual perception of relative relationships and

showcasing the superclusters that constitute anatomical strata in the nerve ring neuropil
(Video S1). The reproducibility of the dome shape depends on assigning two specific
PHATE parameters (https://phate.readthedocs.io/en/stable/api.html) to non-default
values when calling PHATE, the "t" value is set to 50; the "random_state" value is set to
1.

1060
Electron Microscopy based 3-D Models
1061
1062
To make 3-D models of neuron morphologies from vEM datasets, we created Image-J
format regions of interest (ROIs) using published segmentation data (Witvliet et al. 2021;
White et al. 1986; Brittin et al. 2021). For a given cell, the stack of all sectioned ROIs was
then used to draw binary image masks as input to a customized version of the marching
cubes algorithm (Schmid et al. 2010) to build and save a 3-D isosurface. All steps of this
pipeline were executed within the ImageJ-based Java program, CytoSHOW (Duncan et
al. 2019). Slightly modified versions of this workflow were also followed for: 1) generating
cell-to-cell contact ROIs and 2) for generating 3-D representations of synaptic objects.

To align the 3-D models from the variously oriented vEM datasets, all surfaces
from a given specimen were rotated and resized to fit a consensus orientation and scale.
This was achieved by applying a rotation matrix multiplication and scaling factor to all
vertex coordinates in isosurfaces comprising each modeled dataset (Table S2). Each 3-
D object (morphology, contact or synapse) was then exported as a Wavefront file (.OBJ)
and then web-optimized by conversion to a Draco-compressed .GLTF file. Each neuron
was assigned a type-specific color that is consistent across all datasets to enable facile
visual comparison. All the original EM annotations that were used to create the
representative 3D models in NeuroSCAN have been preserved, and can be accessed via
the sister app, CytoSHOW (https://github.com/mohler/CytoSHOW; (Duncan et al. 2019)).

1081
Morphologies
1082
1083
Neuron morphologies were linked across datasets for users to visualize changes over
time. To enhance 3-D graphics performance without sacrificing gross morphologies we
employed a defined amount of data reduction when building each cell-morphology object.
NeuroSCAN can therefore display multiple (or even all) neurons of a specimen within a

single viewer. The number of vertices for a given object was decreased by reducing 10-
fold the pixel resolution of the stacked 2-D masks input into the marching cubes algorithm
of CytoSHOW.

1091

Nerve Ring

To make a simplified mesh of the overall nerve ring shape, individual neuron ROIs
were fused together into a single nerve-ring-scale-stack of image masks. This was used
for input to the marching cubes algorithm. The union of all overlapping enlarged neurite
ROIs in a vEM section was data reduced (20-fold reduced pixel resolution). This rendered
a performance-friendly outer shell of the nerve ring.

1098

Contacts

To build 3-D representations of neuron-neuron contacts, we captured the degree
of overlap when an adjacent cell outline was expanded by the specimen-specific,
empirically-defined pixel threshold distance listed in Table S1 (see Figure S2). This was
done for each cell outline. This expansion step employs a custom-written method in
CytoSHOW that increases the scale of the adjacent outlined region by the pixel threshold
distance (Table S1; Figure S2 B and E), while maintaining its congruent shape. The entire
collection of captured 2-D contact overlaps (Figure S2 C and F) for each adjacent neuron
pairs was then reconstructed as a single 3-D object (Figure S2 H). Contact patches shown
in NeuroSCAN are largely reciprocal (e.g. if there is a AIML contact from PVQL then there
will be a PVQL contact from AIML), but rarely, 2-D overlap regions may be too small to
be reliably converted to 3-D isosurfaces by the marching cubes algorithm, resulting in
absence of an expected reciprocal contact model within the collection. Contacts, like cell
morphology models, are named to be automatically linked across time-point datasets and
to facilitate user-driven visualization of changes over time.

1114

Synapses

Synaptic positions were derived from the original datasets and segmentations, which
annotate synaptic sites in the EM cross-sections (White et al. 1986; Cook et al. 2019;
Witvliet et al. 2021). To represent these coordinates in the 3-D segmented neurons, we

used Blocks (presynaptic sites), Spheres (postsynaptic sites) and Stars (electrical

synapses). The synaptic 3-D objects were placed at the annotated coordinates (White et

al. 1986; Cook et al. 2019; Witvliet et al. 2021). Additionally, the objects were scaled with

the scaling factor (Table S2). Synaptic objects were named by using standard

nomenclature across all datasets, as explained in Supplementary Figure 7.

We note that the L4 and Adult datasets and the L1-L3 datasets were prepared and

annotated by different groups (White et al. 1986; Cook et al. 2019; Witvliet et al. 2021).

Integration of these datasets reveals nanoscale disagreements in the alignment of the

boundaries and synapses. Our representations reflect the original annotations by the

authors. Because of these disagreements in annotations, the synapses are not linked

across datasets. However, all the original EM annotations that were used to create the

representative 3D models in NeuroSCAN, including the synaptic annotations, have been

preserved, and can be accessed by the users via the sister app, CytoSHOW

(CytoSHOW.org).

1133

Supplemental Information:

Supplemental Figures

1136
1137

1138
Figure S1. DC/C-PHATE clustering of AIM, PVQ and AVF across postembryonic
development. (A-E) A cropped view of the DC/C-PHATE plot colored to identify
individual neurons and clustering events in (A) Larva stage 1 (5 hours post hatching); (B)
Larva stage 2, (23 hours post hatching); (C) Larva Stage 3 (27 hours post hatching); (D)
Larva stage 4 (~36 hours post hatching) and (E) Adult (~48 hours post hatching). See
also Video S1 and Table S7.

1145
1146

1147
1148

1149
Figure S2. Projecting contact profiles onto the segmented neuronal shapes. (A-C)
Graphical representations of the strategy utilized for creating the contact profiles for each
of the adjacent neurons (purple, red, cyan) onto a cross section of the neuron of interest
(Neuron A, yellow). **(D-F)** Electron micrograph from the L4 dataset with two adjacent
neurons colored yellow and cyan. To build 3-D reconstructions of contact sites from
adjacent neurons, we analyzed segmented neurons from the electron microscopy
datasets in each slice (A, D). Each adjacent neuron is expanded in all directions to the
pixel threshold distance (specified for each dataset; Table S1; Methods; CytoSHOW.org)
(B, E). A new ROI (region of interest; purple, red, cyan in C; green in F) is created from
the overlapping areas between the neuron of interest (yellow) and the adjacent neurons
(C,F). **(G-I)** 3-D reconstruction of neuron (yellow) (G) with adjacent neuron (cyan), (H)
with contact sites captured (green) across all slices, and (I) with contact areas from the
adjacent neuron augmented (green) as seen in Figure 5 D.

1163
1164

1165
1166
1167

1168
Figure S3. AIM contact sites. Contact sites from PVQ (Orange and highlighted with
orange arrowheads) and from AVF (Green and highlighted with green arrowheads)
across developmental stages (as indicated) and projected onto the segmented AIM
neurons (transparent purple). This figure is the unmodified NeuroSCAN outputs of contact
profiles that corresponds to Figure 5D. In Figure 5D these contact profiles were
augmented. Scale bar = 2 um. See also Figure 5 and Video S4.
1175

1176
1177 **Figure S4. AVF synaptic sites.** Synaptic sites displayed onto transparent (green) AVF
1178 neurons across developmental stages. Presynaptic sites (spheres) and postsynaptic sites
1179 (Blocks) are visualized between the AVF neurons and the AIM (Purple) neurons, PVQ
1180 (Orange) neurons and other AVF (either AVFL or AVFR; opaque green) neuron; Scale
1181 bar = 2 um.
1182

1183
1184
Figure S5. Visualization of contact sites in NeuroSCAN. (A) Search for a specific
neuron (here, AIM) to filter **(B)** the list of contacts corresponding to the developmental
slider. Neuron A (AIML, here) is the neuron onto which the contacts will be mapped. The
Contacts dropdown menu sorts neurons alphabetically (here, colored according to the
contact patch color in C). **(C)** 3-D reconstruction of all AIM contacts at L3 stage. See also
Video S3-S4. In the Figure 5D, contacts are augmented.
1191
1192

1193
Figure S6. C-PHATE tutorial in NeuroSCAN. (A) Add the C-PHATE plot corresponding
to the position of the purple circle on the developmental slider (yellow box) by clicking **(B)**
the + sign. **(C)** Screenshot of C-PHATE plot at L4 (36 hours post hatching), spheres
represent individual neurons at the outer edge of the plot and DC iterations increase
towards the center where spheres represent clusters of neurons and eventually the entire
nerve ring. **(D)** Screenshot of C-PHATE plot at L4 (36 hours post hatching) with the
spheres/clusters containing the AIM neurons highlighted (Blue) by selecting the AIM
neurons within the lightbulb menu (red box). See also Video S1. NeuroSCAN features in
this figure are not shown to scale.
1203

Figure S7. Visualization of synaptic sites with NeuroSCAN. **(A)** Search for synaptic
sites for specific neuron(s) (e.g. AIM, PVQ) and choose a developmental time point with
the slider. **(B)** Synapses dropdown menu contains a list of objects representing pre- and
postsynaptic sites corresponding to all neuron names in the search bar and sorted
alphabetically. Searched neurons can be used with the synaptic filter **(C)** to select for
synapses type (electrical or chemical; Note: only use this feature for L4_36 hours post
hatching and Adult_ 48 hours post hatching) and to filter objects by synaptic specialization
(pre or post; gray dotted box)), **(D)** which will follow the filter logic (example shown for
AIM and PVQ). **(E)** To enable visualization of subsets of synapses and differentiate
between pre- and postsynaptic sites, each synapse contains object(s) representing the
postsynaptic sites(s) as spheres (Blue and Purple) and the presynaptic site as a block
(Orange). These are ordered "by synapse", with all postsynaptic objects, then the
presynaptic object. This specific example corresponds to a 3-D representation of the
PVQL (Orange, Pre)->AIAL (Blue, Post), AIML (Purple, Post) synapse. **(F-G)** All synaptic
sites contain the name of the presynaptic neuron (Orange), neuron type (chemical,
electrical or undefined), list of postsynaptic neuron(s) (Blue) and Unique identifier (Black;
Section, letter) for cases with multiple synapses between the same neurons. The 'section'
is unique to each synapse between specified neurons and at that specific developmental
stage. It is listed in order of its antero-posterior position in the neuron. Synapse names
are not linked through developmental datasets. If the synapse is polyadic, there will be
multiple postsynaptic neuron names and objects associated with a single presynaptic site.
See also Video S4.

A   Opening Page   **B**   Menu

Figure S8. Opening page view and menu. **(A)** View of opening page. **(B)** Menu for access to the 'About' window for referencing source information, the Tutorial, and the developmental Promoter database. See also Video S3.

Figure S9. The NeuroSCAN interface enables interrogation of neuronal relationships across development. (A) The left facing arrow to minimize the left panel and optimize space for the viewer windows. The interface contains four main parts: **(B-E)** Filters, **(F-J)** Results, **(K-M)** Viewer Navigation, and **(N-Q)** viewer windows. Filter Results by (C) searching for neuron names, (D) selecting a dataset with the developmental slider (in hours post- hatching), (E) and filtering synapses based on the pre- or post-synaptic partner on the neurons that are on the search bar. (F) Results drop down menus (filtered by B) for **(G)** Neuronal morphologies (shown in the viewer as purple in (O)), (H) Contacts (shown in green (O)); (I) Synapses (shown in Orange in (O)); and (J) C-PHATE (shown in (Q)), which gets filtered by the developmental slider in (D). (K) Viewer Navigation to rotate the 3-D projections in all viewers simultaneously (Play All) and which contains a drop-down menu for each viewer (L,M). The viewers are named as Viewer 1 (L, N) or CPHATE viewer (M, P) and followed by information of the developmental stage and the hours post hatching for the objects in the viewer. (O) Reconstruction of the AIM neurons

1251 with AVF contacts and synapses at L3 (27 hours post hatching; scale bar = 2 um. (Q) C-
1252 PHATE plot at L1 (0 hours post hatching). See also Video S3.
1253
1254
1255

1256
Figure S10. Select and Add objects to viewers. (A) Click "select (number) items" to
select all items in the dropdown list (green box), or **(A')** click the hexagon next to each
item (green box). **(B)** Click "Add Selected" (purple box) to add all selected items or **(B')**
click "Add to" (purple box) to add each item individually. **(C)** To add the selected item(s)
to an existing viewer of the same developmental stage or to a new viewer, choose a
viewer as indicated. **(D)** Click "Deselect (number) items" (orange box) to deselect items.
See also Video S3 and S4.
1264
1265

1266
1267
1268
Figure S11. In-viewer toolbar features. (A) In-viewer toolbar for Neurons, Contacts and
Synapses and C-PHATE (shown here, only Neurons). **(B, K)** Change the background
color of viewer from dark (white box, moon) to white (white box, sun). **(C, L)** Change the
color of any objects by selecting a desired color, transparency or color code and selecting
the object (or instance) name (here, AIML and AIMR). **(D, M)** Change developmental
stage for items in the viewer by using the in-viewer developmental slider. **(N)** Add 3-D
representations of the Nerve Ring for that developmental stage. **(E, O)** Record and
download movies for the viewer. **(F,P)** Download .gltf files and viewer screenshot (png).
(G) Rotate objects around the y-axis. **(H)** Zoom in and **(I)** zoom out, and **(J)** reset objects
to original positions in the viewer. See also Video S3.

1279
Figure S12. Viewer navigation menu. (A) Navigation bar contains a drop-down menu
for each viewer (shown here, six viewers at varied developmental stages) and a "Play all"
button for simultaneously rotating all objects in each viewer around the y-axis (Video S3).
Each viewer dropdown menu contains a dropdown menu for Neurons (green box),
Contacts and Synapses. **(B)** Viewer 6 with reconstructions of three neurons (AIML and
AIMR, purple; PVQL, orange) at Larval Stage 4 (L4), 36 hours post hatching. **(C)** Browse
and Select objects in the viewer by navigating the nested dropdown menus. **(D)** Manage
objects in viewers with options to select, group, hide, and delete objects in each viewer.
Objects can be deleted with "select" and keyboard "delete". See also Video S3.
1289
1290
1291

1292
Figure S13. NeuroSCAN architecture. (A) Source data is defined in a file tree structure
that contains various assets such as .gltf files representing various entities, as well as
CSVs storing relationships across entities (Data model in Figure S14). The directory
structure outlines a vertical hierarchy starting at the developmental stages, then
branching downwards through neuron, C-PHATE, contact and synapse data. A python
script can be invoked to traverse the directory tree and parse the files, writing to the
database accordingly. This enables verification of the ingested data and quick search
times through the datasets to identify the related items. The architecture uses Geppetto
backend and frontend (Cantarelli et al. 2018). **(B)** The backend uses a Postgres Database
to store underlying data, a Persistent Storage Volume that houses and serves static
assets, and the User Interface is a React application that filters, sorts, and searches
through the Neurons to be added to an interactive canvas. **(C)** A variable number of Virtual
Machines run the frontend and backend application code, scaling as needed to
accommodate traffic. The frontend React/Javascript bundle that is delivered to the **(D)**
client, rendering the neuron data and assets, and a NodeJS application that exposes a
JSON API, serving the neuron data and assets based on user interactions.
1309
1310

NeuroSCAN Data Model

A — Instance

Instance	
name/id	
files	
metadata	

B — Neuron

Neuron	
name (id)	str
files	List<File>
timepoints	List<int>
wormatlas	URL
location	str

C — Developmental Stage

Developmental Stage	
id	str
name	str
begin	int
end	int
order	int
timepoints	List<int>

D — C-PHATE

C-PHATE	
id	str
name	List<Neuron>
files	List<File>
structure	JSON
timepoints	List<int>

E — Contact

Contact	
id	str
neuronA	Neuron
neuronB	Neuron
files	List<File>
timepoints	List<int>
weight	int

F — Synapse

Synapse	
id	str
neuronPre	Neuron
neuronPost	List<Neuron>
files	List<File>
type	["chemical","electrical"]
timepoints	List<int>
section	str
zs	str
position	["pre", "post"]
neuronSite	str

Figure S14. NeuroSCAN data model. (A) Reference scheme for B-F; Instance refers to the category (e.g. B, Neuron; C, Developmental Stage), which contains a name or identifier (id) for each object, lists of files associated with the instance (C, Developmental Stage does not have files), and metadata to further describe each instance which is usually a string (str) or an integer (int). **(B)** The neuron name is the foundation for the Contacts, Synapses, and C-PHATE which enables integration across each of these representations and across developmental stages (timepoints) with metadata from WormAtlas (https://wormatlas.org/MoW_built0.92/MoW.html). **(C)** The Developmental Stages are named by the larval stages (L1, L2, L3, L4, Adult), and the metadata captures the list of timepoints within those developmental stages (i.e. L1, 0 hours post hatching and L1, 5 hours post hatching). **(D)** C-PHATE objects are named with a list of Neurons. **(E)** Contacts link to the Neuron names (Neuron A and Neuron B nomenclature in Figure S5) and metadata annotates the weight or the number of pixels of contact quantified in the source Electron Microscopy micrographs. **(F)** Synapses link to the Neuron names (Pre, Post, type, and section described in Figure S7).

Supplemental Videos

Video S1. Visualization of hierarchical relationships using C-PHATE plots in NeuroSCAN. The process for rendering a C-PHATE plot at the L4 stage (36 hours post hatching) with the real-time loading speed. In the viewer, 3-D visualization of a C-PHATE plot (shades of cyan), which is rotated to show the dome-shape of the plots and to orient the plot to correspond to Figures 2 and 4. The highlight functionality is used to show the spheres containing AIM (teal), then PVQ (teal). The spheres of the first iterations, containing AIM and PVQ, are identified, selected and colored magenta. The AVF neurons are highlighted in teal, and the first AIM and AVF containing clusters are identified, selected and colored yellow. The first clusters containing AIML, AVF and PVQL are identified and colored green. Neurons in the left yellow and magenta clusters are reconstructed with a right click on the sphere and "Add to new viewer" selection.

Video S2. Analysis of AIM, PVQ and AVF neuronal morphologies in developmental datasets. 3-D visualizations of AIM (Purple), PVQ (Orange) and AVF (Green) at (Left viewer) L1 (5 hours post hatching) and (Right viewer) L3 (27 hours post hatching) in NeuroSCAN. Note that at L1, AVF has not grown into the nerve ring, therefore, only AIM and PVQ are present, but by L3, the AVF neurons have grown between the AIM and PVQ neurons.

Video S3. Navigating NeuroSCAN features that enable integration of Neurons, Contacts and Synapses across developmental datasets. Upon first opening NeuroSCAN, a tutorial will launch (Figure S8). In the NeuroSCAN menu one can read about NeuroSCAN, access the tutorial, and navigate to the embryonic promoter database (Figure S8). The video shows the user searching neurons (AIM and PVQ) and adding neurons to the viewers (Figure S10). Side-by-side viewers with AIML, AIMR, and PVQL enable comparisons across developmental stages (L1, 0 hours post hatching and L4, 36 hours post hatching). Also shown in the video are the use of the in-viewer toolbar (Figure S11) and navigation menu (Figure S12) for object exploration.

Video S4. Exploring Contacts and Synapses using NeuroSCAN. Video of user navigating the tools of NeuroSCAN to examine synapses and contact profiles to yield results as in (Figures S7 and S9). AIM neurons (Transparent Purple), AIM (Purple)-PVQ synaptic sites (Orange), and AIM-PVQ contact sites (Orange) at L1 (5 hours post hatching) are added into Viewer 1. AIM neurons (Transparent Purple), AIM(Purple)-PVQ synaptic sites (Orange), and AIM-PVQ contact sites (Orange), AVF (Green)-AIM synaptic sites, and AVF-AIM contact sites (Green) at L3 (27 hours post hatching) are added into Viewer 2. Contact sites and synaptic sites are compared across developmental stages by hiding AIM neurons. All contact sites for AIM are added for L1 (5 hours post hatching) into Viewer 3.

Supplemental Tables

Table S1. Nerve ring regions, resolutions, and pixel threshold distances used to calculate adjacency matrices and to create contact sites for each dataset.

Table S2. Scaling factors and rotation corrections for 3-D representations of Neurons, Contacts and Synapses for each dataset.

Table S3. Stratum 1 (Red) Sankey diagrams of clustered neurons for each Diffusion Condensation iteration in each dataset.

Table S4. Stratum 2 (Purple) Sankey diagrams of clustered neurons for each Diffusion Condensation iteration in each dataset.

Table S5. Stratum 3 (Blue) Sankey diagrams of clustered neurons for each Diffusion Condensation iteration in each dataset.

Table S6. Stratum 4 (Green) Sankey diagrams of clustered neurons for each Diffusion Condensation iteration in each dataset.

Table S7. Sankey diagrams of AIM, PVQ and AVF containing clusters for each Diffusion Condensation iteration in each dataset.

Tables S8. L1 (0 hours post hatching) adjacency counts and searchable counter for summed adjacencies. Type the name of a "Neuron of Interest" (NOI) in the indicated cell to filter for the summed adjacency counts for each contact partner. For each partner, there are two columns: Total number of contacts (number of EM sections NOI and partner are in contact) and Total Weights (summed number of pixels NOI and partner contacts).

Tables S9. L1 (5 hours post hatching) adjacency counts and searchable counter for summed adjacencies. Type the name of a "Neuron of Interest" (NOI) in the indicated cell to filter for the summed adjacency counts for each contact partner. For each partner, there are two columns: Total number of contacts (number of EM sections NOI and partner are in contact) and Total Weights (summed number of pixels NOI and partner contacts).

Tables S10. L2 (23 hours post hatching) adjacency counts and searchable counter for summed adjacencies. Type the name of a "Neuron of Interest" (NOI) in the indicated cell to filter for the summed adjacency counts for each contact partner. For each partner, there are two columns: Total number of contacts (number of EM sections NOI and partner are in contact) and Total Weights (summed number of pixels NOI and partner contacts).

Tables S11. L3 (27 hours post hatching) adjacency counts and searchable counter for summed adjacencies. Type the name of a "Neuron of Interest" (NOI) in the indicated cell to filter for the summed adjacency counts for each contact partner. For each partner, there are two columns: Total number of contacts (number of EM sections NOI and partner are in contact) and Total Weights (summed number of pixels NOI and partner contacts).

Tables S12. L4 (36 hours post hatching) adjacency counts and searchable counter for summed adjacencies. Type the name of a "Neuron of Interest" (NOI) in the indicated cell to filter for the summed adjacency counts for each contact partner. For each partner, there are two columns: Total number of contacts (number of EM sections NOI

and partner are in contact) and Total Weights (summed number of pixels NOI and
partner contacts).
Tables S13. Adult (48 hours post hatching) adjacency counts and searchable counter
for summed adjacencies. Type the name of a "Neuron of Interest" (NOI) in the indicated
cell to filter for the summed adjacency counts for each contact partner. For each
partner, there are two columns: Total number of contacts (number of EM sections NOI
and partner are in contact) and Total Weights (summed number of pixels NOI and
partner contacts).

1430

1431

1432

1433
1434