1 **Selective autophagy of ribosomes balances a tradeoff between starvation survival and growth**
2 **resumption**
3
4 **Authors**

5 Joel Tuomaala[1,2], Devanarayanan Siva Sankar[3], Julie Perey[1], Sacha Psalmon[1,2], Nicholas Stroustrup[4], Joern
6 Dengjel[3], Benjamin D. Towbin[1,*]
7

Affiliations
[1] Institute of Cell Biology, University of Bern; Bern, Switzerland.

[2] Graduate School for Cellular and Biomedical Sciences, University of Bern; Bern, Switzerland.

[3] Department of Biology, University of Fribourg; Fribourg 1700, Switzerland.

[4] Centre for Genomic Regulation (CRG), The Barcelona Institute of Science and Technology (BIST).

*corresponding author (benjamin.towbin@unibe.ch)

Abstract

In environments with fluctuating nutrient abundance, organisms must survive periods of starvation, yet
quickly resume growth upon food encounter. A tradeoff between these objectives is well-documented in
microbes, where it is caused by the need to partition the total cellular protein content between growth-
and survival-enhancing proteins. However, the molecular mechanisms of growth-survival tradeoffs in
multicellular animals remain largely unknown. Here, we addressed this mechanism for *C. elegans* by
measuring the dynamic changes of its proteome during starvation using live imaging and proteomics. We
found that starved animals catabolize ribosomal proteins through autophagy, which provides essential
energy for survival while preserving organismal integrity. However, the resulting decline in ribosomes
delayed growth resumption upon refeeding until pre-starvation ribosome levels were restored. Genetic
inhibition of ribosomal autophagy had a dual effect: although it accelerated growth after short starvation,
it compromised survival during prolonged starvation. These findings reveal the rate of ribosomal
catabolism as a key determinant of a tradeoff between starvation survival and rapid growth resumption
whose tuning may adapt animals to different starvation durations. Our research shows how the need to
balance protein allocation between growth and survival constrains animal physiology, highlighting the
mechanistic role of proteome resource limitation in whole-organism tradeoffs.

Introduction

In nature, organisms face alternating periods of food abundance and scarcity. They must therefore effectively withstand times of starvation and rapidly resume growth when food becomes available. This dual requirement necessitates a delicate balance between investing in growth and survival [1–3].

Tradeoffs between growth and survival have been extensively characterized in microorganisms [4,5], where they are primarily governed by the partitioning of a limited proteome between growth-promoting and survival-enhancing functions [6–10]. In microbes, the growth rate follows a simple linear relation with the ribosome content called the 'growth law' [11,12]. Thus, while high expression of survival factors improves survival under starvation, it also sequesters resources from the production of ribosomes that would be needed for maximal growth [7].

Growth-survival tradeoffs are also prevalent in multicellular animals [2,13], but their underlying cellular and molecular mechanisms remain poorly understood. Do the molecular constraints of proteome partitioning [14] also explain tradeoffs occurring in animals? Or is a more nuanced and comprehensive mechanistic analysis required to understand tradeoffs in these much more complex multicellular systems? Addressing these questions can identify universal principles of organismal tradeoffs and reveal fundamental design principles of biological systems.

During starvation, eukaryotes catabolize their protein content by autophagy [15,16]. While autophagy supports survival by providing vital energy and metabolites, it can also harm survival by degrading essential cellular components [17]. In principle, this threat could be mitigated by the selective degradation of proteins that are not needed during starvation, such as ribosomes, which remain largely idle during starvation-induced growth arrest. Indeed, selective autophagy of ribosomes, called ribophagy, occurs in amino acid-starved yeast [18]. However, whether starvation also induces ribophagy in other eukaryotes and if ribophagy has a physiological role for the starvation resistance of animals remains contentious [19–24].

To resolve this controversy and explore the role of protein resource allocation in tradeoffs at a multicellular scale, we used the nematode *C. elegans* as the model system. The natural habitat of *C. elegans* is characterized by boom-bust cycles of rapid population growth and extended starvation [15,25]. An appropriate balance between survival and growth resumption is thus central to its fitness in the wild, and its genetic tractability make it well-suited to address the mechanisms involved.

Here, we monitored the dynamic changes of the proteome during starvation and refeeding of *C. elegans* larvae by proteomics and live imaging techniques. Our measurements showed unambiguously that starvation induces selective ribophagy, leading to a progressive decline in the ribosome levels. This decline in ribosomes delayed resumption of growth upon refeeding until the pre-starvation ribosome concentrations were restored. Reminiscent of microbial growth laws, the speed of growth resumption was near-linearly related to the ribosome levels at the start of refeeding, which suggests that linear growth laws also inform on physiological tradeoffs in complex multicellular animals. Indeed, by genetically manipulating the rate of ribophagy, we could predictably alter the balance between growth and survival and show that preserving ribosomes accelerates growth resumption but impairs survival during prolonged starvation.

Together, our findings reveal the rate and selectivity of ribophagy as key determinants in balancing starvation survival and growth resumption in a multicellular animal. This discovery highlights that the partitioning of limited proteome resources plays a key mechanistic role in tradeoffs at an organismal scale,

73 bridging the gap between microbial and multicellular biology in the molecular understanding of
74 fundamental tradeoffs.

Results

Ribosomal proteins decline rapidly during starvation

To measure proteome dynamics during starvation, we conducted quantitative label-free proteomics [26] of
L1 larvae of *C. elegans* after 1, 4, and 9 days of starvation (Fig. 1a). Using suitable algorithms to compare
relative amounts of different proteins within the same sample (iBAQ [27]) and relative changes of individual
proteins across different samples (LFQ [28]), we detected 2624 and 1787 proteins consistently across all
conditions and repeats. Ribosomes contributed 12.1% (s.e.m. = 0.2%) to the total detected protein mass,
as estimated by iBAQ, on day 1 of starvation and dropped by 27% (s.e.m. = 1.3%) to 9.0% (s.e.m. = 0.3%)
by day 9 of starvation. Non-ribosomal cytoplasmic proteins declined by only 12.5% (s.e.m. = 1.7%), whereas
nuclear and filamentous or membrane bound proteins increased in their relative abundance (Fig. 1d).
Consistent with previous reports [29], the total mitochondrial protein mass also declined rapidly (-23.8%,
s.e.m. = 3.0%). Notably, our experiments measured protein abundance relative to the total protein mass
and not the absolute protein amount per animal. The observed decline in ribosomal proteins is thus not a
byproduct of an overall size reduction or protein loss and shows that ribosomal proteins decline more
rapidly than the total proteome. An increase in relative protein abundance, as observed for nuclear
proteins (Fig. 1d-e), may be due to *de novo* synthesis of these proteins or due to a net decline that was
slower than that of the total proteome.

Comparisons of the relative changes of individual proteins over time, estimated by LFQ, confirmed these
conclusions. Ribosomal proteins, on average, declined more strongly than non-ribosomal cytoplasmic
proteins (Fig. 1e, Fig. S1a-b). Consistently, proteins involved in translation, including but not restricted to
ribosomes, declined more strongly than the average proteome (Fig. 1f). For LFQ-based measurements, no
significant difference between the decline of mitochondrial and cytoplasmic proteins was observed (Fig.
1e). This difference between iBAQ and LFQ can occur when, on average, the more mass-abundant
mitochondrial proteins have a faster relative decline. Indeed, abundant mitochondrial ribosomal proteins
were among the most rapidly declining mitochondrial proteins (Fig. S1a). Yolk proteins also declined
rapidly, consistent with their role in nutrient storage [30]. However, since only three yolk proteins were
detected, contributing a mere 0.1% to the total detected proteome mass, the depletion of yolk was not
statistically significant (Fig. 1e).

We next validated the mass-spectrometry-based measurements by luminescence-based detection using
HiBiT [31] (Fig. 1b) and by live imaging. For HiBiT, we endogenously tagged the ribosomal protein RPL-17 and
the yolk protein VIT-2 at their C-termini. Confirming measurements by mass-spectrometry, RPL-17 and VIT-
2 declined rapidly (-7.0%/day and -10.9%/day) normalized to total protein (Fig. 1g), showing their
preferential degradation compared to other proteins. The quantitative difference in decline rates observed
with HiBiT and mass spectrometry may be due to technical differences, or due to biological differences
among individual ribosomal proteins that are masked by the measurement noise of mass-spectrometry.

Figure 1. Rapid ribosomal protein decline during starvation. a-c. Experimental design of measurements of protein dynamics during starvation of L1 larvae by mass spectrometry (a) HiBiT (b) and live imaging (c). **d.** Mass fraction of indicated proteome subgroups on day 1 and day 9 of starvation (groups are based on GO cellular component annotation and quantified using iBAQ). Numbers inside sectors indicate mass fraction. Numbers outside sectors indicate the relative change in fraction size between days 1 and 9. **e,f.** Box plots of rates of change measured for individual proteins in indicated groups (quantified using LFQ). Stars above boxes: significance of comparison between groups (robust one-way ANOVA, adjusted post-hoc comparisons). Stars below boxes: difference to zero (adjusted Wilcox signed rank test) ***p<0.001 **p<0.01. **g.** Protein quantification by HiBit normalized to total protein at t = 0. error bars: s.e.m. from 3 biological repeats. line: linear regression (*rpl-17:HiBiT* r²=0.94, *vit-2:HiBiT* r²=0.79). shaded area: 95% CI of linear fit. **h.** Mean volumetric protein density of GFP-fluorescence over starvation time. shaded area: 95% CI. n = 146, 323 and 137 animals from 3 to 4 biological repeats for *lmn-1:gfp*, *daf-21:gfp* and *rpl-29:gfp*. **i.** Images of indicated strains after 0h and 55h starvation in agarose chambers. White: animal outline. Color scale: GFP intensity with equal contrast adjustment for the same genotypes. Animals were computationally straightened for display.

For live imaging, we created a strain with an endogenous GFP insertion in the ribosomal protein RPL-29, and used strains with integrated transgenes expressing the non-ribosomal cytoplasmic protein DAF-21:GFP[32] and the nuclear lamin LMN-1:GFP[33]. For each strain, we placed at least 100 eggs individually in

arrayed agarose-based chambers [34] that did not contain food. Upon hatching from the eggs, the starving
L1 larvae were imaged hourly by fluorescence microscopy for 55 hours, at an exposure that did not cause
significant photobleaching (Fig. S1c). We then determined the animal volume and fluorescent protein
concentration (fluorescence/volume) by computational segmentation and straightening [35]. In agreement
with proteomics data, the concentration of ribosomal RPL-29:GFP declined during starvation, nuclear LMN-
1:GFP increased in fluorescence, and cytosolic DAF-21:GFP fluorescence concentration remained
approximately constant (Fig. 1h,i & Fig. S1d-e). Taken together, the rate of decline varies among different
cytoplasmic proteins during starvation with ribosomal and translation-associated proteins being among
the most rapidly declining cytoplasmic proteins.

Preferential ribosomal protein decline is autophagy-dependent

136
To test if the decline in ribosomal proteins required autophagy, we measured RPL-29:GFP in starving L1
larvae carrying mutations in the autophagy factors *atg-18*/WIPI or *unc-51*/ULK1 [29,36]. In both autophagy
deficient mutants, the decline of RPL-29:GFP was significantly slower than in wild type, albeit not
completely abolished (Fig. 2a). The residual decline was proportional to the overall decline in animal
volume, such that ribosome density was constant during starvation of these mutants (Fig. S2a-c). The
residual decline may be due to ribosomal protein degradation unrelated to autophagy, or due to residual
autophagic activity in the mutants. Importantly, although *atg-18* and *unc-51* are known to impair starvation
resistance [29,36], both mutants remained viable throughout the measurement period of 55 hours, as
determined by their motility and a lack of death-associated swelling [37] (Fig. S2a-b,d).

To determine the subcellular fate of ribosomal proteins, we created strains in which RPL-34 and RPL-22
were endogenously tagged with mCherry. Unlike GFP, mCherry remains fluorescent under the acidic
lysosomal conditions [38–40]. Upon starvation, RPL-34:mCherry (Fig. 2b) and RPL-22:mCherry (Fig. S2e)
became spatially heterogeneous (Fig. 2b,c & Fig. S2e,f), forming fluorescent foci. These foci were weaker
or absent in *atg-18* and *unc-51* mutants and no foci occurred for RPL-29:GFP in wild type animals (Fig.
2b,c), suggesting that mCherry foci represent accumulation in the acidic lysosomal compartment [40].
Consistently, 80% of RPL-34:mCherry foci were surrounded by, or co-localized with LMP-1:GFP, the *C.
elegans* homolog of the lysosome associated membrane protein (LAMP) [41] (Fig. 2d). We conclude that the
decline of ribosomal protein during starvation is largely autophagy dependent, and that ribosomal proteins
decline more rapidly than the average cytoplasmic proteome, consistent with the occurrence of an
autophagy pathway selective for ribosomes in *C. elegans*.

Figure 2. Ribosomal protein decline is autophagy-dependent. a. Total RPL-29:GFP fluorescence in wild type and in autophagy deficient animals during starvation. Mean and 95% CI of the mean for n = 283, 299, 101 animals from 3 biological repeats for *unc-51(e1189)* and *atg-18(gk378)* and 2 biological repeats for wild type. **b.** RPL-34:mCherry and RPL-29:GFP in wild type and autophagy deficient individuals at hatch and after 48h starvation. Contrast is equal for the same genotype. Focus formation is autophagy and starvation dependent, and only occurs for acid-insensitive mCherry, but not for GFP. **c.** Box plot of relative change in coefficient of variation (CV) among pixel intensities for strains and conditions shown in (b) (113, 49, 71, and 25 individuals per strain). Significance between group means (adjusted paired sample t-test): ***p<0.001, *p<0.05 (Table S1). **d.** Co-localization of RPL-34:mCherry and lysosomal membrane associated protein LMP-1:GFP at hatching and after 24h of starvation. Arrowheads: ribosomal foci, surrounded by LMP-1:GFP.

Ribosomal protein amount after starvation predicts recovery growth rate

To explore the impact of ribosomal decline on starvation survival and growth during recovery, we starved L1 larvae expressing RPL-29:GFP for 1, 3, and 6 days before feeding them in agarose chambers. By live imaging, we measured their organismal growth and ribosome expression at a time resolution of 15 minutes throughout development (Fig. 3a). In this assay, the larval molts of *C. elegans* (M1 to M4) can be detected by a halt in volume growth prior to ecdysis [34,42] (Fig. 3b), allowing precise determination of the larval stage durations. Consistent with previous reports [43], starvation delayed the completion of the L1 stage by

approximately two hours per day of starvation (Fig. 3c). This lag time was inversely correlated (r^2=0.87)
with the total RPL-29:GFP fluorescence measured at the first time point of feeding (Fig. 3d,e). The
correlation occurred across different durations of starvation, and across individuals exposed to the same
duration of starvation (Fig. 3d). Thus, regardless of the starvation conditions experienced, the remaining
RPL-29:GFP after starvation predicted the time a given individual needed to complete the first larval stage
with high accuracy (Fig. 3d). RPL-29:GFP fluorescence also correlated with body volume, but the observed
correlation between growth and RPL-29:GFP level was not due to volume differences alone. The correlation
between animal volume and larval stage duration was weaker than that between total RPL-29:GFP and
larval stage duration ($r^2 = 0.67$ vs. $r^2 = 0.85$, Fig. S3a). Moreover, we also detected a significant correlation
(r^2=0.64) between the larval stage duration and the RPL-29:GFP density (GFP/volume) (Fig. S3b). During
the first larval stage, starved animals recovered ribosomes to near normal levels (Fig. S3c). Consistently,
starvation duration had a small or no effect on larval stage durations, growth rates, ribosome expression,
and body volumes from the second larval stage onwards (Fig. 3b, Fig. S3c-d).

188
Figure 3. Ribosome level predicts recovery speed. a. Experimental design of starvation recovery measurements. **b.**
Mean animal volume during post starvation feeding after 1, 3 and 6-day starvation for n = 67, 66 and 43 indivduals.
Squares: molts between larval stages. Shaded area: 95% CI of the mean. Means were computed by aligning individual
trajectories to the molts prioir to averaging, followed by re-scaling to the mean larval stage duration. **c.** Box plots of
L1 duration after 1, 3 and 6-day starvations for 89, 89 and 71 individuals, respectively. Significance between group

means (two sample t-test): ***p<0.001 (Table S1). **d.** RPL-29:GFP fluorescence vs. the inverse of the L1 larval stage
duration after recovery from 1, 3 and 6-day starvation for 85, 83 and 44 individuals. Lines: linear correlation with 95%
CI. Black: all starvation durations pooled (r^2=0.85). blue, gray, red: separated by starvation duration (r^2=0.62, r^2=0.76
and r^2=0.19 for 1, 3, and 6-days). **e.** Images of the individuals expressing the highest and the lowest RPL-29:GFP after
6-day starvation at indicated times of feeding. Contrast is matched for animals at the same time of feeding. White:
body outline. Color scale: GFP intensity. Animals were computationally straightened for display.

We conclude that the level of ribosomes remaining after starvation is a good predictor for the rate at which
L1 larvae resume development. Our data suggests that the need to re-synthesize degraded ribosomes, and
possibly other proteins required for growth, slows down recovery from starvation.

Hyperactivation of the mTORC1 activator RAGA-1 reduces ribosomal decline

To slow-down ribosomal protein decline experimentally during starvation, we genetically engineered the
mTORC1 activator Ras-related GTP-binding protein A (RAGA-1), the single *C. elegans* ortholog of
mammalian RagA/B [44]. RAGA-1 normally switches from a GTP to a GDP bound state upon amino acid
starvation, which represses mTORC1 activity. We introduced a mutation that stabilizes the GTP bound form,
hereafter called RAGA-1^{GTP}, that constitutively activates mTORC1 regardless of amino acid availability[45],
and ultimately upregulates biosynthetic pathways and inhibits autophagy [46] (Fig. 4a).

To test if RAGA-1^{GTP} animals retained ribosomes during starvation, we conducted quantitative label-free
proteomics in wild type and mutant animals side-by-side. We sampled on days 1, 4, and 7, during which
viability of RAGA-1^{GTP} animals was not compromised compared to wild type (Fig. 4h, Fig. S5f). The RAGA-
1^{GTP} mutation significantly dampened the decline in the mass fraction of ribosomal proteins estimated by
iBAQ (Fig. 4b-c, -32.8%, s.e.m. = 1.4% vs. -17.1%, s.e.m. = 1.8%, $\Delta_{(WT - RAGA-1-GTP)}$ = -15.7%). RAGA-1^{GTP} had a
much smaller effect on the decline of cytoplasmic and mitochondrial proteins (Δ = -6.3% for both, Fig. 4b-
d), suggesting that this mutation reduces the selectivity of ribophagy.

We note that the proteome mass fraction sizes differed between our initial measurements of wild type
alone (Fig. 1d) and when measured for comparison with RAGA-1^{GTP} (Fig. 4b-c). These differences may be
due to experimental batch effects or due to the higher number of proteins detected in the latter
experiment (2789 and 2052 proteins for iBAQ and LFQ). Importantly, both experiments consistently show
that ribosomes decline more rapidly than the average cytoplasm in wild type animals.

Consistent with changes in proteome fractions, the median rate of decline of individual ribosomal proteins
(based on LFQ) was faster in wild type than in RAGA-1^{GTP} animals (-9.7%/day vs. -6.0%/day, Δ = -3.7%/day,
Fig. 4d). This difference between RAGA-1^{GTP} and wild type was similarly pronounced for proteins associated
with translation in general (Δ = -3.2%/day, Fig. S4c), but less pronounced for mitochondrial proteins (Δ = -
1.3%/day), non-ribosomal cytosolic proteins (Δ = -1.1%/day), yolk proteins (Δ < 0.1%/day), and various
nuclear components (Fig. 4d & Fig. S4a). Thus, *raga-1* influenced the turnover of ribosomal proteins more
strongly than that of other proteins (Fig. 4b-d & Fig. S4a-b).

Figure 4. Reducing ribophagy shifts a tradeoff between starvation survival and recovery: a. Schmatic of RAGA-1 in mTORC1 signalling. **b,c**. Quantification of proteome mass fractions on day 1 and 7 of starvation as described in Fig. 1, but for wild type and RAGA-1[GTP] animals. Percentages inside sectors indicate group fraction size. Percentages outside sectors indicate proportional change in fraction size from the 1st day to the 7th day of starvation. **d.** Box plots of rates of decline for individual proteins grouped by GO cellular component-terms for wild-type (solid fill) and RAGA-1[GTP] (dotted fill). Stars above boxes: differences between group means (robust two-way ANOVA, adjusted post-hoc comparison). Stars below boxes: difference to zero (adjusted Wilcox signed rank test). (***p<0.001, **p<0.01). **e,f**. Ratio of relative growth rate ((dV/dt)/V) between RAGA-1[GTP] and wild type animals during *ad libitum* feeding after hatching (e) and after indicated duration of starvation (f) for the first 8 hours of feeding. Solid line: mean of 177 to 386 individuals across 2 to 5 biological repeats per strain and condition. Shaded area: 95% CI of the mean. **g**. Fraction of individuals that successfully recovered growth after indicated starvation duration. A total of 3467 wild type and 2621 RAGA-1[GTP] individuals were sampled across 3 biological repeats. **h.** Starvation survival in agarose chambers. 48 wildtype and 70 RAGA-1[GTP] animals. **i.** Proposed model: trade-off between starvation survival and starvation recovery is tuned by the selectivity of autophagy towards ribosomes and possibly other proteins required for translation and growth. Heatmap shows simulated relative population growth for competing strains with different growth and

survival balances (rows) under different durations of starvation (columns). The optimal compromise that maximizes
selective fitness (yellow) depends on the duration of starvation.

We confirmed these data by live imaging. Immediately after hatching, RPL-29:GFP expression was 10%
higher in RAGA-1^{GTP} animals than in wild-type animals (Fig. S4d-e), consistent with mTORC1
hyperactivation. Throughout 55 hours of starvation, RPL-29:GFP total fluorescence, fluorescence relative
to the starting levels, and volumetric density remained higher in RAGA-1^{GTP} (Fig. S4d-g). RAGA-1^{GTP} did not
significantly affect the body volume during starvation (Fig. S4h-i), consistent with a quantitative reduction
of ribophagy by RAGA-1^{GTP}, rather than a total halt of bulk autophagy (Fig. 4b-d). Measurement of protein
production rates by photo-bleaching [47] showed that RPL-29:GFP production during starvation was
negligible in mutant and wild type animals (Fig. S5a-d). The difference in the rate of decline between
mutant and wild type are therefore not due to differences in ribosomal protein synthesis, but due to
different rates of degradation. Differences in fluorescence between RAGA-1^{GTP} and wild type were also not
explained by a GFP tag associated with the RAGA-1^{GTP} mutation, since the expression of this tag was near
negligible compared to that of RPL-29:GFP (Fig. S5e). Together, we conclude that hyperactivation of
mTORC1 activity by RAGA-1^{GTP} slows down the starvation-induced decline of ribosomal proteins and
possibly of other translation associated proteins.

RAGA-1^{GTP} impairs starvation resistance and speeds up growth during recovery
Consistent with previous findings in mice [48], RAGA-1^{GTP} mutants had reduced starvation resistance with
more than 50% of the animals failing to recover growth after 5 days of starvation (Fig. 4g), and a sharp drop
of viability (based on motility) after 9 days (Fig. 4h). To test if impaired survival during long starvation was
paralleled by a faster recovery of RAGA-1^{GTP} animals after short starvation, we measured the growth rate
of wild-type and RAGA-1^{GTP} animals in agarose chambers without starvation and after 1, 2, and 3 days of
starvation. We observed no significant difference in the growth rate of RAGA-1^{GTP} and wild-type animals
that were not exposed to starvation (Fig. 4e). However, after 1 and 2 days of starvation, RAGA-1^{GTP} mutants
grew significantly faster than wild-type during the first two hours of recovery (Fig. 4f, Fig. S6a-b,i-j). This
growth advantage of RAGA-1^{GTP} was transient and disappeared later in development when wild type
animals had reconstituted their ribosomes (Fig. S6c-f). Thus, although the RAGA-1^{GTP} did not accelerate
growth *per se*, the mutation accelerated recovery from starvation, possibly by retaining higher levels of
ribosomes. No significant growth benefit of RAGA-1^{GTP} animals was observed after 3 days of starvation (Fig.
4f & Fig. S6b,i,j), possibly due to an increased frailty of this mutant after longer starvation (Fig. 4g,h) or a
smaller difference in ribosome levels between wild type and mutant after 3 days of starvation (Fig. S6e,f).

Together, these data show that hyperactivation of RAGA-1 accelerates growth during recovery after short
periods of starvation but impairs growth and survival after longer starvation periods. The balance between
the retention of ribosomal proteins, and possibly other translation factors, and their degradation to fuel
metabolism provides a parsimonious explanation for the widely observed tradeoff between growth and
survival [5,6,8,49–51] (Fig. 4i).

Best compromise between recovery speed and survival depends on starvation duration
To explore under which conditions a rapid recovery may be favorable compared to a high starvation
survival, we simulated population growth using realistic life-history parameters for *C. elegans* in a
mathematical model. In the model, starvation caused a developmental delay τ that increased linearly with
time (τ = δ * t), as experimentally observed [43]. We modelled starvation survival by Hill function with half-
point σ matching measured data (Fig. 4h). To model the tradeoff, δ and σ were proportionally coupled (δ ∝
σ). Strains with different tradeoff strategies were competed against each other until the total population

had expanded 20-fold (representing food exhaustion). These simulations showed that a longer median
survival σ at the cost of a longer delay τ were beneficial for long durations of starvation (Fig. 4i) whereas a
short delay and short survival were beneficial for short durations of starvation. The optimal compromise
between growth and survival thus depends on the duration of starvation periods, suggesting that the
ribophagy rate is an adaptive trait whose optimal value depends on the specific starvation regime.

Discussion

In a dynamic nutritional environment, the ability to survive extended starvation and to rapidly resume
growth when encountering food is crucial for organismal fitness. In the face of tradeoffs, these tasks must
be optimally balanced, in accordance with the frequency and duration of starvation periods. Evidently, if
episodes of starvation are long, it is crucial to be able to survive long periods without food. However, if the
expected duration of starvation is short, a strategy that accelerates growth resumption at the expense of
survival may be advantageous (Fig. 4i). Here, we employed quantitative assays to reveal a mechanism by
which *C. elegans* can adjust the balance between these key aspects of organismal physiology.

We showed that *C. elegans* selectively catabolize ribosomal proteins during starvation through autophagy
(Figs. 1 & 2), a process called ribophagy [18]. This strategy enhances survival by providing energy without
degrading essential cellular components but comes at the cost of delayed growth resumption upon
refeeding (Figs. 3 & 4). Notably, the synthesis of ribosomal proteins itself depends on functional ribosomes.
The loss of ribosomes therefore not only impairs the onset of growth, but also the re-synthesis of
ribosomes themselves. This combined effect may explain the observed developmental delay of up to
several days following recovery from starvation [43,52].

Unlike other pathways of selective autophagy [53], the mechanisms governing ribophagy are only beginning
to be characterized [24]. While amino acid starvation is known to induce ribophagy in yeast [18], the specific
receptor mediating this process has not been identified. In mammalian cells, the protein NUFIP1 has been
proposed as a ribophagy receptor[21], but conflicting evidence has cast doubt on its role [20]. Similarly, it
remained contentious to what extent selective ribophagy indeed contributes to the decline in ribosomes
relative to unspecific bulk autophagy [19,22]. Importantly, our study not only demonstrates that ribosomes
are degraded by autophagy during *C. elegans* starvation (Fig. 2), but also that the rate of ribophagy is faster
than the bulk degradation of other cytoplasmic proteins (Fig. 1). Moreover, genetically manipulating the
ribophagy rate directly impacted both starvation survival and the speed of growth recovery (Fig. 4). These
findings firmly establish the biological relevance of regulated ribosomal turnover in a physiological context,
providing a foundation for future investigations of its molecular control.

Quantitative studies of growth-survival tradeoffs have primarily focused on bacteria [4,5], where the growth
rate follows a linear relation with the ribosomal protein content [11]. This simple relation has enabled
accurate predictions of cellular growth dynamics when switching between conditions [10,54,55], that did not
rely on modelling the thousands of molecular reactions controlling cellular growth. Our data in *C. elegans*
reveal a similarly linear relationship between ribosome levels and the speed of growth recovery from
starvation (Fig. 3). This finding suggests that the core principles governing proteome partitioning and
growth control are portable from the microbial to the multicellular scale, at least for *C. elegans*, enabling
coarse-grained models of organismal biology.

It is important to note that the partitioning of the proteome between different functions is not restricted
to ribosomes. Indeed, we find that also other translation-related proteins are rapidly catabolized during
starvation (Fig. 1) and may also be limiting for growth recovery. Similarly, the genetic modulation of mTOR

signalling through *raga-1* did not only impact ribophagy, but also the turnover of other proteins required for growth and translation (Fig. 4). Whereas our current live imaging measurements have focused on ribosomal proteins, ribosomes evidently are not the only proteins required for growth but serve here as a representative example of growth-enhancing proteins more generally. Whether the selectivity of autophagy towards ribosomes and other translation factors is controlled by the same or by independent mechanisms presents an important mechanistic question for future studies.

In addition to the tradeoff between starvation survival and growth resumption examined in this study, the proteome resource allocation to ribosomes may also be important for other organismal tradeoffs. For instance, the striking correlation between the translation rate and lifespan [56] indicates that a similar mechanism may underlie the well-known tradeoff between growth and longevity. In this context, high production of ribosomes may accelerate growth but sequester resources from expression of survival-enhancing proteins that counteract aging. The quantitative methodology and framework established here now empower us to systematically address this hypothesis and uncover the full scope of proteome partitioning in shaping diverse organismal tradeoffs.

An important question raised by our work is how environmental conditions shape the evolution of selective ribophagy. Our theoretical analysis predicts that the best compromise between starvation survival and rapid growth resumption depends on the average duration of starvation periods. One would thus expect a rapid ribophagy rate in strains adapted to long starvation periods, while adaptation to shorter periods of starvation should favor the preservation of ribosomes, which would facilitate rapid growth resumption. Our molecular characterization of ribophagy in an organismal context opens exciting opportunities to test these predictions and understand how molecular processes shape the evolution of growth and survival along organismal tradeoffs.

Materials and Methods

C. elegans strains and maintenance

Animals were grown and maintained at 25°C using OP50-1 *E. coli* on nematode growth medium (NGM) according to standard procedures. The following strains were used in this study:

WBT100: *rpl-17(xe232)[rpl-17:HiBit]; xeSi296[peft-3::luc-gfp::unc-54 3'UTR, unc-119(+)] II* (this study), used in Fig. 1a-b,d-g, Fig. S1a-b, Fig. S7a,c-f

WBT203: *vit-2(syb3768)[vit-2::GSSG::3xflag::GSSG::HiBiT] X* (this study), used in Fig. 1g

GW8: *ygIs[lmn-1::GFP::lmn-1 unc-119(+)] X; unc-119(ed3) III* ([33]), used in Fig. 1c,h-i, Fig. S1d-e

RBW2661: *hutSi2661 [hsp-90p::eGFPT::unc-54 3'UTR + Cbr-unc-119 (+)] II* ([32]), used in Fig. 1c,h-i, Fig. S1d-e

PHX1945: *rpl-29(syb1945) [rpl-29:GFP]* (this study), used in Fig. 1c,h-i, Fig. 2a-c, Fig. 3, Fig. 4e-g, Fig. S1c-e, Fig. S2a-d,f, Fig. S3, Fig. S4d-i, Fig. 5Sf, Fig. S6a-b,e-f, Fig. S6

N2: Bristol N2 wild isolate, used in Fig. 1c,g-i, Fig. S1d-e, Fig. S5a-e

WBT257: *rpl-29(syb1945) [rpl-29:GFP] IV; unc-51(e1189) V* (this study, allele e1189 from CGC registered strain CB1189 by J. A. Hodgin lab, Oxford University), used in Fig. 2a, Fig. S2a-c,f

WBT277: *rpl-29(syb1945) [rpl-29:GFP] IV; atg-18(gk378) V* (this study, allele gk378 from [57]), used in Fig. 2a, Fig. S2a-c,f

PHX1880: *rpl-34(syb1880) [rpl-34:RPL-34:mCherry] IV* (this study), used in Fig. 2b-c, Fig. S2f

PHX1903: *rpl-22(syb1903) [rpl-22:RPL-22:mCherry] II* (this study), used in Fig. S2e

WBT230: *rpl-34(syb1880) [rpl-34:RPL-34:mCherry] IV; unc-51(e1189) V* (this study, allele e1189 from CGC
registered strain CB1189 by J. A. Hodgin lab, Oxford University), used in Fig. 2b-c

WBT234: *rpl-34(syb1880) [rpl-34:RPL-34:mCherry] IV; atg-18(gk378) V* (this study, allele gk378 from [57]),
used in Fig. 2b-c

WBT343: *unc-119(ed3)III; pwls50[lmp-1p:lmp-1::GFP+Cb-unc-119(+)]; rpl-34(syb1880) [rpl-34:RPL-*
34:mCherry] IV (this study, allele pwls50 from CGC registered strain RT258 by B. D. Grant lab, Rutgers
University), used in Fig. 2d

WBT64: *xeSi296 [Peft-3::luc::gfp::unc-54 3'UTR, unc-119(+)] raga-1(xe190) II* (this study, allele xeSi296
from [58], allele xe190 from [45]), used in Fig. 4b-d, Fig. 4Sa-c, Fig. S7b,g-i

HW1939: *xeSi296 [peft-3::luc::gfp::unc-54 3'UTR, unc-119(+)] II* ([58]), used in Fig. 4b-d, Fig. 4Sa-c, Fig.
S7b,g-i

WBT330: *raga-1(wbm40) [raga-1::AID::EmGFP] II; xeSi376ubs38[Peft-3::TIR1(*ubs38 *[F79G]):mRuby::unc-*
54 3'UTR, cb-unc-119(+)] III; rpl-29(syb1945) [rpl-29:GFP] IV (this study, allele xeSi376ubs38 from [59]), used
in Fig. 4h, Fig. S5f

WBT289: *raga-1(wbm40abt3) [raga-1(GF)::AID::EmGFP] II;* xeSi376ubs38*[Peft-3::TIR1(*ubs38
[F79G]):mRuby::unc-54 3'UTR, cb-unc-119(+)] III; rpl-29(syb1945) [rpl-29:GFP] IV (this study, allele
xeSi376ubs38 from [59]), used in Fig. 4b-h, Fig. S4d-i, Fig. S5c-f, Fig. S6

WBT231: *raga-1(wbm40abt3) [raga-1(GF)::AID::EmGFP] II* (this study), used in Fig. S5e

Egg collection and starvation cultures
Animals were grown on NGM plates at 25°C for at least 3 generations prior to starvation. For mass
spectrometry experiments, the peptone concentration in NGM was increased to 20mg/ml to reach
sufficient numbers of individuals. Mixed populations were synchronized by bleaching and the eggs were
allowed to develop into gravid adults on new plates. The adults were filtered with a cell strainer with 40µm
mesh size (Pluriselect) to remove younger animals and incubated in 35mM serotonin in S-basal in a rotating
cradle at 25°C for 1.5h. Eggs were harvested and separated from adults with a 40µm mesh, and the eggs
were filtered a 20µm mesh (Neolab) to remove hatched L1 larvae. The purified eggs were then either
loaded into agarose chambers for timelapse imaging or allowed to hatch overnight in S-basal (without
cholesterol and ethanol) in a rotating cradle at 25°C at a concentration of 10 eggs/µl. The following day,
the hatched L1 larvae were filtered with 20µm mesh to remove unhatched eggs. The L1 larvae were then
starved in S-basal (without cholesterol and ethanol) in a rotating cradle at 25°C at a density of 10 L1/µl for
the remainder of the starvation duration.

For mass spectrometry experiments, the synchronized gravid animals were bleach synchronized instead of
serotonin treated to obtain eggs. The eggs were allowed to hatch overnight in S-basal (without cholesterol
and ethanol) in an orbital shaker at 25°C at a concentration 10 eggs/µl. The hatched L1 larvae were filtered
with a 20µm mesh to remove unhatched eggs. The L1 larvae were starved in sterile S-basal cultures

(without cholesterol and ethanol) in orbital shaker at 25°C at 10 L1/µl for the remainder of the starvation
duration.

Unlabeled mass spectrometry
Starving L1 larvae were collected (10^6 animals/sample) from triplicate cultures (S-basal without cholesterol
and ethanol at 25°C in orbital shaker at 10 animals/µl) and resuspended into 700µl PBS without additives.
The pellets were flash frozen by pipetting dropwise into liquid nitrogen. Frozen beads were powdered using
cryogenic SPEX SamplePrep 6775 Freezer/Mill and stored as frozen powder at -80°C.

Powdered pellets were dissolved in 1% sodium deoxycholate buffer containing 50 mM Tris-HCl pH 8.
Samples were treated with 1 mM DTT and alkylated using 5.5 mM iodoacetamide for 10 min at RT. Samples
were in-solution digested with trypsin (Promega). Tryptic peptides were purified by STAGE tips and LC-
MS/MS measurements were performed on a QExactive plus and an Orbitrap Exploris 480 mass
spectrometer coupled to EasyLC 1200 nanoflow-HPLCs (all Thermo Scientific) using data-dependent
acquisition. MaxQuant software (version 1.6.2.10 & 2.0.1.0) [60] was used for analyzing the MS raw files
performing peak detection, peptide quantification and identification using a Uniprot *C. elegans* database,
November 2017. Carbamidomethylcysteine was set as fixed modification and oxidation of methionine was
set as variable modification. The MS/MS tolerance was set to 20 ppm and four missed cleavages were
allowed for Trypsin/P as enzyme specificity. Based on a forward-reverse database, protein and peptide
FDRs were set to 0.01, minimum peptide length was set to seven, and at least one unique peptide had to
be identified. MaxQuant results were analysed using Perseus software (version 1.6.2.3)[61]. iBAQ and LFQ
values were used for expression proteomics analyses. iBAQ values of the respective samples were median
normalized. Only proteins with valid values in all three biological replicates in respective groups were
considered for further analysis.

iBAQ and LFQ data were analyzed using R-studio (v4.3.2). Mass fractions were determined by multiplying
iBAQ quantification of each detected protein by its molecular weight. Resulting mass values were grouped
and summed based on GO terms and divided by the total iBAQ mass. Molecular weights and GO terms
were acquired from UniProtKB by list search of protein IDs reported by MaxQuant. The rate of change for
individual proteins was computed by the slope of a linear best fit through logarithmic LFQ fraction
measurements. Slopes of individual proteins were grouped based on whether the GO terms associated
with the protein matched category presented in Fig. 1d-f and 4d (e.g. "ribosome"). For figures shown in
main text, each protein was allowed to appear in only one group (see Table S2 for list of proteins in each
group). Equivalent analysis allowing for group overlap is shown in supplemental material (Fig. S7c-e).

HiBiT measurements
Starving L1 larvae were collected (3000 animals/sample) from triplicate cultures into 100µl lysis buffer (PBS
pH8, 0.25% triton, protease inhibitors (Sigma P8340)), and freeze-thawed three times in liquid nitrogen
and 37°C water bath. Samples were lysed by sonication (Bioruptor) at 4°C and diluted with 600 µl PBS prior
to HiBiT assay.

After lysis all pipetting was conducted using motorized pipettes (Sartorius) and a pipetting robot (Opentron)
to minimize experimental variation. HiBiT intensity was measured daily in three biological replicates (3
measurements x 3 wells x 3 replicates / day) in 384-well format using a HiBiT quantification kit (Promega
Nano-Glo). 5µl HiBiT reaction mixture (1:100 LgBiT protein and 1:50 HiBiT lytic substrate in HiBiT lytic
buffer) was combined with 5µl sample. Plates were shaken at room temperature for 5min prior to
measuring luminescence on a luminometer (SpectromaxL).

Protein concentration in each sample was measured using micro-BCA kit (Pierce) in three biological replicates (3 measurements x 3 wells x 3 replicates / day) in 96-well format. 140ul sample was combined with 140ul BCA reagent (25:24:1 mix of reagents MA:MB:MC) at 4°C. The plates were shaken for 5min, incubated stationary in 37°C incubator for 2h, and measured with Spectromax340 562nm in endpoint mode.

Data analysis was done in R (v4.3.2). Background signals were subtracted (BCA reagent with lysis buffer for micro-BCA, N2 worm lysate with reaction mixture for HiBiT) and HiBiT measurements were normalized with total protein BCA measurements to acquire HiBiT fraction. The fractions were normalized again by the average of the 3 biological replicates at t=0 and a linear model was fitted to the normalized values.

Live imaging in agarose-based chambers

Arrayed agarose chambers were manufactured by stamping a microscopic chamber pattern into gel of 4.5% agarose in S-basal using a PDMS template (Wunderlichips) based on Turek et al. [62], incorporating the modifications by Stojanovski et al. [34]. Animals were retained isolated in separate chambers and imaged every hour for starvation experiments, and every 15 minutes for post-starvation growth experiments. Chamber dimensions were 270x270x20μm for L1 starvation experiments in Fig. 1, Fig. 1S, Fig. 2 and Fig. 2S. Chamber depth was decreased to 10μm for L1 starvation experiments in Fig. 4S and for all L1 starvation recovery experiments in Fig. 4 to improve image focus, accuracy of segmentation, and volume estimates. Chamber dimensions were increased to 600x600x20μm for post-starvation growth experiments in Fig. 3 and Fig. S3 to accommodate animal growth at L3 and L4 stages.

For starvation experiments, the chambers were manufactured without adding cholesterol and ethanol to the agarose. 0.003% Triton was added to the eggs to reduce clumping and the eggs were pelleted. The egg pellet was gently broken up before pipetting 5μl concentrated eggs onto freshly manufactured chambers. 2-fold or older eggs inside a droplet of S-basal without cholesterol or ethanol were then individually guided to chambers using an eyelash pick.

For post-starvation growth experiments, chambers were filled with OP50-1 *E. coli* by scraping an *E. coli* lawn off a standard NGM plate and spreading it onto the chambers with a piece of 3% agarose gel prior to loading animals as described by Stojanovski et al. [34]. In Fig. 4, the agarose of the chambers contained 0.05% DMSO. Starved L1 larvae were pipetted next to the array of chambers and then moved individually to chambers using an eyelash pick. Chamber array was prevented from drying during loading by pipetting 20μl deionized water around the outer perimeter of the chamber array.

After loading animals into the chambers, the agarose chambers were inverted onto dishes with gas permeable polymer bottom of high optical quality (Ibidi) and surrounded by 3% low melting temperature agarose. The assembly was covered with thin layer of PDMS to prevent evaporation and the dish was closed and sealed with parafilm and used for imaging.

A Nikon Ti2 wide-field epifluorescence microscope equipped with a SpectraX light source (Lumencor) and a Prime BSI camera (Photometrics) was used for all live imaging experiments. A 10x 0.45NA objective was used in Fig. 3b-e, Fig. S3a-d, Fig. S4e-g, and Fig. S6. A 20x 0.75NA objective was used in Fig. 1h-I, Fig. S1c-e, Fig. 2a-c, Fig. S2a-f, and Fig. S5a-h. Effective pixel size in the sample plane was 0.65 and 0.325μm, respectively. Software auto-focus of NIS Elements Software (Nikon) was used to find the central focal plane in the GFP or mCherry channel of the ribosomal markers using single band pass filters. For all experiments, the temperature was maintained at 25 °C using an incubator encapsulating the entire microscope (life imaging services).

 Confocal imaging of RPL-22:mCherry foci
In Fig. 2d, a Nikon CREST X-light spinning disk confocal microscope was used to image L1 animals
immobilized on agarose pads by 10mM Levamisole and a coverslip. Animals were imaged at hatch or after
24h starvation. A 100x 1.45NA oil immersion objective was used to capture Z-stacks around manually set
focus plane with a single band pass filters for mCherry and GFP using 200ms exposure. Images were
deconvoluted using Huygens software.

Detection of animals and animal fluorescence in agar chambers
A Matlab script was used to process raw images as described [34]. Sobel algorithm using the edge function
of Matlab (R2022b) was used to segment the animals from raw fluorescent images. Detected outline
endpoints closest to each other were connected to close gaps in the animal outline. From the outline a
binary mask was created to separate animals from background. For L1 starvation experiments in Fig. 1 and
Fig. S1 a U-net [63] network was trained to segment the binary mask of animals from raw brightfield images.
Segmentation was performed in brightfield channel to allow for the segmentation of non-fluorescent N2
control animals. The Sobel-based masks were used as the U-net training data. For both methods animal
fluorescence was measured from the area covered by the mask in the fluorescent channel. For Fig. 1h-I,
N2 animals were imaged in parallel and their auto fluorescent signal was subtracted from all measurements
due to the comparatively weak fluorescence signal of LMN-1:GFP. For Figs. 2a and 3d, only the background
signal from agar or bacteria was subtracted, as the tissue autofluorescence was negligible compared to the
fluorescence of RPL-29:GFP (Fig. S5e).

The masks were computationally straightened using a custom made Matlab function [34]. Straightened masks
were classified as either 'egg', 'worm' or 'error' using a decision tree-based classifier that was trained on a
small subset of manually annotated masks. The classifier was an ensemble of 20 bootstrap aggregation
decision trees (TreeBagger function of Matlab) that was trained on features length, standard deviation of
width, cv of width, maximal width, median width, maximal width/median width, volume, volume/length,
and entropy of width [34]. All experiments were classified with the same classifier, except for experiments
with autophagy deficient animals, whose occasionally irregular outline necessitated training of a separate
classifier for reliable error detection. Values derived from images with errors or missing animals were
excluded from further analyses.

Computation of volume and growth rate
The volume of the animals in agarose chambers was derived from the straightened binary image masks
assuming rotational symmetry [35]. For growing animals, larval stage transitions were determined based on
the timing of the maximum of the second time derivative of the logarithm of animal volume. Using a
graphical user interface created in Matlab [34], the ecdysis timings were corrected manually based on
plateaus in animal growth curves caused prior to ecdysis. To minimize measurement error, for Fig. 3d and
Fig. S3a,b, the fluorescence and volume at t0 was estimated by fitting a linear model to the first 20
timepoints and the predicted value at t0 was plotted. Individuals for which less than 6 data points were
available in the first 20 timepoints were excluded. For Fig. 3c, the volume measurement was scaled to the
larval development by linearly interpolating the volume at 100 points per larval stage. The interpolated
values are plotted over the mean larval stage duration, such that the beginning and end of each larval stage
are aligned on the x-axis. The same scaling was used for fluorescence in Fig. S3c.

The rate of growth during starvation recovery for Fig. 4e-f was calculated in Matlab by smoothing every
individual animal's volume with the fit function's smoothing spline option using a smoothing parameter of
0.0001. Fits were manually inspected to ensure that the resulting volume interpolations are appropriate,

after which volume was interpolated at a time resolution of 1 minute. Volume growth rate by minute was
calculated as the local slope of the smoothed volume curve by using gradient function. Individuals' growth
rates were normalized to the mean growth rate of the control condition, and a mean of normalized growth
rates was determined for every replicate chamber array.

Photobleaching based detection of GFP synthesis
L1 larvae starved for 16h in agarose chambers were imaged hourly for three hours before exposure to a
477nm laser (Celesta light engine, Lumencor) for 8 seconds at 100% laser power until approximately 60%
of GFP signal had been bleached. Viability was not affected based on motility. Animals were subsequently
imaged for 20 hours, the auto-fluorescence was subtracted using N2 animals as control, and the
fluorescence signal was normalized by the first timepoint after photobleaching (Fig. S5b-c).

Data analysis: The net change in total fluorescence of GFP (g) is the difference in its production with rate
p and degradation with rate constant d, such that:

$$\dot{g} = p - d * g \qquad \text{and} \qquad \frac{\dot{g}}{g} = \frac{p}{g} - d$$

Thus, for p > 0, the relative rate of change in GFP ($\frac{\dot{g}}{g}$) would increase after photobleaching, assuming that
the rate constant d is unaffected by photobleaching. Conversely, if $\frac{\dot{g}}{g}$ is unaffected by photobleaching (as
observed in Fig. S5b,d), p must be negligible.

Starvation recovery and survival assays
The ability of animals to recover growth after starvation (Fig. 4g) was measured daily at 25 °C by placing a
total of 50-150 starving L1 larvae onto the bacterial lawn of fresh NGM plates from 3 independent cultures
and counted immediately. After 2 or 3 days of growth (for shorter and longer than 4 day-starvation,
respectively) animals that had exited L1 (based on their larger size) were considered to have resumed
development and recovered from starvation.

Starvation survival (Fig. 4h) was determined by placing a minimum of 36 eggs per strain into agarose
chambers without food and imaging every 17 min for 3 weeks at 25 °C. Worms were computationally
detected as described above. Time of death was determined as the time of permanent loss of motility that
was followed by death associated swelling (Fig. S2d).

Population Growth Model
Population growth for different tradeoff strategies was computed using numerical simulations in
Matlab(R2022a) by calculating the relative fraction of animals among different strategies after different
times of starvation during one generation. Survival s after time t of starvation was described by a Hill
function with Hill coefficient n=4 and half-point σ.

$$s(t) = \frac{1}{1 + (^t/_\sigma)^4}$$

The lag time τ was determined by a linear relation τ = δ * t, where δ is the additional lag time per time of
starvation. The tradeoff was model by the linear coupling $\sigma = k * \delta$. Parameter k was set to 56 days. 20
strains with δ being linearly spaced between 0.01 and 0.15 were computationally competed against each
other, starting with a uniformly distributed population. This parameter range corresponds to a median
survival σ spanning between 1.5 days and 22 days, and a lag times τ after 7 days of starvation between 1.7

hours and 25 hours, centered around the experimentally observed values shown in Fig. 3. Reproduction
was kept equal among all strains, starting after 40 hours plus the respective lag time τ at a constant
reproduction rate of 6 progeny per hour for 48 hours. The population was allowed to grow until it multiplied
20 times to simulate food exhaustion. The data shown in Fig. 4i is the relative frequency of each strain in
the entire population at the end of the first starvation/recovery cycle.

Statistics
A robust one-way ANOVA (Fig. 1e-f) or a robust two-way ANOVA (Fig. 4d) (t1way and t2way from R package
WRS2(v.1.1-4)) was performed to evaluate effects of genotype and protein group on the rate of change in
LFQ fraction sizes (Table S3). For group-wise comparisons, Scheffe post-hoc test for one-way ANOVA and
Games-Howel post-hoc test for two-way ANOVA was performed (Table S4). See Table S5 for testing
statistical assumptions. Where appropriate, normal one- and two-way ANOVA, Kruskal-Wallis, Tukey's HSD,
Dunn, and Tukey's HSD with Games-Howel modification tests are also reported (Table S3, Table S4). For
LFQ group difference to zero in Figures 1e-f and 4d, a Benjamini-Hochberg multiple comparison adjusted
Wilcox signed rank test was used. Bonferroni and Holm-Bonferroni adjustments are also reported (Table
S6).

For Fig. 2c, Fig. S2f, and Fig. S6i-j a Benjamini-Hochberg multiple comparison adjusted two-sample t-test
was performed (Table S1, Bonferroni and Holm-Bonferroni corrections are also reported).

References

1. Stearns, S. C. *The Evolution of Life Histories*. vol. 249 (Oxford University Press Oxford, 1992).

2. Mangel, M. & Stamps, J. Trade-offs between growth and mortality and the maintenance of individual variation in growth. *Evol. Ecol. Res.* **3**, (2001).

3. Rion, S. & Kawecki, T. J. Evolutionary biology of starvation resistance: what we have learned from Drosophila. *J. Evol. Biol.* **20**, 1655–1664 (2007).

4. Bergkessel, M., Basta, D. W. & Newman, D. K. The physiology of growth arrest: uniting molecular and environmental microbiology. *Nat. Rev. Microbiol.* **14**, 549–562 (2016).

5. Abram, F., Arcari, T., Guerreiro, D. & O'Byrne, C. P. Chapter Four - Evolutionary trade-offs between growth and survival: The delicate balance between reproductive success and longevity in bacteria. in *Advances in Microbial Physiology* (eds. Poole, R. K. & Kelly, D. J.) vol. 79 133–162 (Academic Press, 2021).

6. Shoval, O. *et al.* Evolutionary Trade-Offs, Pareto Optimality, and the Geometry of Phenotype Space. *Science* **336**, 1157–1160 (2012).

7. Yang, Y. *et al.* Temporal scaling of aging as an adaptive strategy of *Escherichia coli*. *Sci. Adv.* **5**, eaaw2069 (2019).

8. Biselli, E., Schink, S. J. & Gerland, U. Slower growth of Escherichia coli leads to longer survival in carbon starvation due to a decrease in the maintenance rate. *Mol. Syst. Biol.* **16**, e9478 (2020).

9. Basan, M. Resource allocation and metabolism: the search for governing principles. *Curr. Opin. Microbiol.* **45**, 77–83 (2018).

10. Basan, M. *et al.* A universal trade-off between growth and lag in fluctuating environments. *Nature* **584**, 470–474 (2020).

11. Scott, M. & Hwa, T. Bacterial growth laws and their applications. *Curr. Opin. Biotechnol.* **22**, 559–65 (2011).

12. Schaechter, M., MaalØe, O. & Kjeldgaard, N. O. Dependency on Medium and Temperature of Cell Size and Chemical Composition during Balanced Growth of *Salmonella typhimurium*. *Microbiology* **19**, 592–606 (1958).

13. Dmitriew, C. M. The evolution of growth trajectories: what limits growth rate? *Biol. Rev.* **86**, 97–116 (2011).

14. Liebermeister, W. *et al.* Visual account of protein investment in cellular functions. *Proc. Natl. Acad. Sci.* **111**, 8488–8493 (2014).

15. Baugh, L. R. & Hu, P. J. Starvation Responses Throughout the *Caenorhabditis elegans* Life Cycle. *Genetics* **216**, 837–878 (2020).

16. Kaur, J. & Debnath, J. Autophagy at the crossroads of catabolism and anabolism. *Nat. Rev. Mol. Cell Biol.* **16**, 461–472 (2015).

17. Kang, C., You, Y. & Avery, L. Dual roles of autophagy in the survival of Caenorhabditis elegans during starvation. *Genes Dev.* **21**, 2161–2171 (2007).

18. Kraft, C., Deplazes, A., Sohrmann, M. & Peter, M. Mature ribosomes are selectively degraded upon starvation by an autophagy pathway requiring the Ubp3p/Bre5p ubiquitin protease. *Nat. Cell Biol.* **10**, 602–610 (2008).

19. An, H. & Harper, J. W. Systematic analysis of ribophagy in human cells reveals bystander flux during selective autophagy. *Nat. Cell Biol.* **20**, 135–143 (2018).

20. An, H., Ordureau, A., Körner, M., Paulo, J. A. & Harper, J. W. Systematic quantitative analysis of ribosome inventory during nutrient stress. *Nature* **583**, 303–309 (2020).

21. Wyant, G. A. *et al.* NUFIP1 is a ribosome receptor for starvation-induced ribophagy. *Science* **360**,
751–758 (2018).

22. Gretzmeier, C. *et al.* Degradation of protein translation machinery by amino acid starvation-induced
macroautophagy. *Autophagy* **13**, 1064–1075 (2017).

23. López, A. R. *et al.* Autophagy-mediated control of ribosome homeostasis in oncogene-induced
senescence. *Cell Rep.* **42**, 113381 (2023).

24. Beese, C. J., Brynjólfsdóttir, S. H. & Frankel, L. B. Selective Autophagy of the Protein Homeostasis
Machinery: Ribophagy, Proteaphagy and ER-Phagy. *Front. Cell Dev. Biol.* **0**, (2020).

25. Schulenburg, H. & Félix, M.-A. The Natural Biotic Environment of *Caenorhabditis elegans*. *Genetics*
206, 55–86 (2017).

26. Arike, L. *et al.* Comparison and applications of label-free absolute proteome quantification methods
on Escherichia coli. *J. Proteomics* **75**, 5437–5448 (2012).

27. Schwanhäusser, B. *et al.* Global quantification of mammalian gene expression control. *Nature* **473**,
337–342 (2011).

28. Cox, J. *et al.* Accurate Proteome-wide Label-free Quantification by Delayed Normalization and
Maximal Peptide Ratio Extraction, Termed MaxLFQ. *Mol. Cell. Proteomics MCP* **13**, 2513–2526
(2014).

29. Hibshman, J. D. *et al.* Nonselective autophagy reduces mitochondrial content during starvation in
Caenorhabditis elegans. *Am. J. Physiol. - Cell Physiol.* **315**, C781–C792 (2018).

30. Perez, M. F. & Lehner, B. Vitellogenins - Yolk Gene Function and Regulation in Caenorhabditis elegans.
Front. Physiol. **10**, 1067 (2019).

31. Schwinn, M. K. *et al.* CRISPR-Mediated Tagging of Endogenous Proteins with a Luminescent Peptide.
ACS Chem. Biol. **13**, 467–474 (2018).

32. Sands, B. *et al.* A toolkit for DNA assembly, genome engineering and multicolor imaging for C.
elegans. *Transl. Med. Aging* **2**, 1–10 (2018).

33. Liu, J. *et al.* Essential Roles for Caenorhabditis elegans Lamin Gene in Nuclear Organization, Cell
Cycle Progression, and Spatial Organization of Nuclear Pore Complexes. *Mol. Biol. Cell* **11**, 3937–
3947 (2000).

34. Stojanovski, K., Großhans, H. & Towbin, B. D. Coupling of growth rate and developmental tempo
reduces body size heterogeneity in C. elegans. *Nat. Commun.* **13**, 3132 (2022).

35. Moore, B. T., Jordan, J. M. & Baugh, L. R. WormSizer: High-throughput Analysis of Nematode Size and
Shape. *PLOS ONE* **8**, e57142 (2013).

36. Takacs, Z. *et al.* ATG-18 and EPG-6 are Both Required for Autophagy but Differentially Contribute to
Lifespan Control in Caenorhabditis elegans. *Cells* **8**, 236 (2019).

37. Stroustrup, N. *et al.* The Caenorhabditis elegans Lifespan Machine. *Nat. Methods* **10**, 665–670
(2013).

38. Rennick, J. J., Nowell, C. J., Pouton, C. W. & Johnston, A. P. R. Resolving subcellular pH with a
quantitative fluorescent lifetime biosensor. *Nat. Commun.* **13**, 6023 (2022).

39. Kneen, M., Farinas, J., Li, Y. & Verkman, A. S. Green Fluorescent Protein as a Noninvasive Intracellular
pH Indicator. *Biophys. J.* **74**, 1591–1599 (1998).

40. Pankiv, S. *et al.* p62/SQSTM1 Binds Directly to Atg8/LC3 to Facilitate Degradation of Ubiquitinated
Protein Aggregates by Autophagy*. *J. Biol. Chem.* **282**, 24131–24145 (2007).

41. Kostich, M., Fire, A. & Fambrough, D. M. Identification and molecular-genetic characterization of a
LAMP/CD68-like protein from Caenorhabditis elegans. *J. Cell Sci.* **113**, 2595–2606 (2000).

42. Uppaluri, S. & Brangwynne, C. P. A size threshold governs Caenorhabditis elegans developmental
progression. *Proc. Biol. Sci.* **282**, 20151283 (2015).

43. Olmedo, M. *et al.* Prolonged quiescence delays somatic stem cell-like divisions in Caenorhabditis
elegans and is controlled by insulin signaling. *Aging Cell* **19**, e13085 (2020).

44. Gollwitzer, P., Grützmacher, N., Wilhelm, S., Kümmel, D. & Demetriades, C. A Rag GTPase dimer code
defines the regulation of mTORC1 by amino acids. *Nat. Cell Biol.* **24**, 1394–1406 (2022).

45. Al-Refaie, N. *et al. An mTOR/RNA Pol I Axis Shapes Chromatin Architecture in Response to Fasting*.
http://biorxiv.org/lookup/doi/10.1101/2023.07.22.550032 (2023) doi:10.1101/2023.07.22.550032.

46. Saxton, R. A. & Sabatini, D. M. mTOR Signaling in Growth, Metabolism, and Disease. *Cell* **168**, 960–
976 (2017).

47. Eden, E. *et al.* Proteome Half-Life Dynamics in Living Human Cells. *Science* **331**, 764–768 (2011).

48. de la Calle Arregui, C. *et al.* Limited survival and impaired hepatic fasting metabolism in mice with
constitutive Rag GTPase signaling. *Nat. Commun.* **12**, 3660 (2021).

49. Gurvich, Y., Leshkowitz, D. & Barkai, N. Dual role of starvation signaling in promoting growth and
recovery. *PLoS Biol.* **15**, e2002039 (2017).

50. Scharf, I., Filin, I. & Ovadia, O. A trade-off between growth and starvation endurance in a pit-building
antlion. *Oecologia* **160**, 453–460 (2009).

51. Stamps, J. A. Growth-mortality tradeoffs and 'personality traits' in animals. *Ecol. Lett.* **10**, 355–363
(2007).

52. Roux, A. E., Langhans, K., Huynh, W. & Kenyon, C. Reversible Age-Related Phenotypes Induced during
Larval Quiescence in C. elegans. *Cell Metab.* **23**, 1113–1126 (2016).

53. Birgisdottir, Å. B., Lamark, T. & Johansen, T. The LIR motif – crucial for selective autophagy. *J. Cell Sci.*
126, 3237–3247 (2013).

54. Korem Kohanim, Y. *et al.* A Bacterial Growth Law out of Steady State. *Cell Rep.* **23**, 2891–2900 (2018).

55. Erickson, D. W. *et al.* A global resource allocation strategy governs growth transition kinetics of
Escherichia coli. *Nature* **551**, 119 (2017).

56. Steffen, K. K. & Dillin, A. A Ribosomal Perspective on Proteostasis and Aging. *Cell Metab.* **23**, 1004–
1012 (2016).

57. Large-Scale Screening for Targeted Knockouts in the Caenorhabditis elegans Genome. *G3*
GenesGenomesGenetics **2**, 1415–1425 (2012).

58. Meeuse, M. W. *et al.* Developmental function and state transitions of a gene expression oscillator in
Caenorhabditis elegans. *Mol. Syst. Biol.* **16**, e9498 (2020).

59. Hills-Muckey, K. *et al.* An engineered, orthogonal auxin analog/AtTIR1(F79G) pairing improves both
specificity and efficacy of the auxin degradation system in Caenorhabditis elegans. *Genetics* **220**,
iyab174 (2021).

60. Cox, J. & Mann, M. MaxQuant enables high peptide identification rates, individualized p.p.b.-range
mass accuracies and proteome-wide protein quantification. *Nat. Biotechnol.* **26**, 1367–1372 (2008).

61. Tyanova, S. *et al.* The Perseus computational platform for comprehensive analysis of (prote)omics
data. *Nat. Methods* **13**, 731–740 (2016).

62. Turek, M., Besseling, J. & Bringmann, H. Agarose Microchambers for Long-term Calcium Imaging of
Caenorhabditis elegans. *JoVE J. Vis. Exp.* e52742 (2015) doi:10.3791/52742.

63. Ronneberger, O., Fischer, P. & Brox, T. U-Net: Convolutional Networks for Biomedical Image
Segmentation. Preprint at https://doi.org/10.48550/arXiv.1505.04597 (2015).

64. Perez-Riverol, Y. *et al.* The PRIDE database resources in 2022: a hub for mass spectrometry-based
proteomics evidences. *Nucleic Acids Res.* **50**, D543–D552 (2021).

Acknowledgements

We thank Sebastian Leidel, Anat Bren, Peter Lenart, and Helge Grosshans for helpful feedback on the manuscript. We acknowledge support by the Microscopy Imaging Center at the University of Bern. Some strains were provided by the CGC, which is funded by NIH Office of Research Infrastructure Programs (P40 OD010440). Figures 1a-c and 3a were created with BioRender.com. Alleles *xe232, syb3768*, *syb1945*, *syb1880* and *syb1903* were created by SunnyBiotech.

Funding

This work received support from an SNSF Eccellenza Grant (PCEFP3_181204) to BDT and the Bern Research Foundation (34/2022).

Author contributions

BDT and JT conceived the study, wrote the manuscript. JD, NS, BDT and JT edited the manuscript. JD and DSS conducted mass spectrometry measurements. JT and JP conducted all other experiments. JT and BDT performed computational and statistical analyses. BDT performed fitness modeling. SP implemented the U-net based image segmentation. NS designed alleles *syb1945*, *syb1880* and *syb1903*. JT, NS and JP created and validated strains used in the study.

Competing interests

The authors declare no competing interest.

Data and Code availability

The mass spectrometry proteomics data have been deposited to the ProteomeXchange Consortium via the PRIDE[64] partner repository with the dataset identifier PXD050951. Volume and fluorescence data from microscopy experiments is available in supplemental tables. Raw images are available upon request from the authors. Code for image analysis was as used in[34] (10.5281/zenodo.10032867). Code for the population growth model is available as supplementary information.

Supplementary Materials

Figs. S1 to S7

Tables S1 to S6