Columnar cholinergic neurotransmission onto T5 cells of *Drosophila*

Eleni Samara[1,2*], Tabea Schilling[1], Inês M. A. Ribeiro[1,3], Juergen Haag[1], Maria-Bianca Leonte[1,2], Alexander Borst[1,4*]

[1] Max Planck Institute for Biological Intelligence, Department of Circuits-Computation-Models, 82152 Planegg, Germany

[2] Graduate School of Systemic Neurosciences, LMU Munich, 82152 Planegg, Germany

[3] Institute of Medical Psychology, Medical Faculty, LMU Munich, 80336 Munich, Germany

[4] Lead contact

*Correspondence: eleni.samara@bi.mpg.de (E.S.), alexander.borst@bi.mpg.de (A.B.)

Summary

Several nicotinic and muscarinic acetylcholine receptors (AChRs) are expressed in the brain of *Drosophila melanogaster*. However, the contribution of different AChRs to visual information processing remains poorly understood. T5 cells are the primary motion-sensing neurons in the OFF pathway and receive input from four different columnar cholinergic neurons, Tm1, Tm2, Tm4 and Tm9. We reasoned that different AChRs in T5 postsynaptic sites might contribute to direction selectivity, a central feature of motion detection. We show that the nicotinic nAChRα1, nAChRα4, nAChRα5 and nAChRα7 subunits localize on T5 dendrites. By targeting synaptic markers specifically to each cholinergic input neuron, we find a prevalence of the nAChRα5 in Tm1-, Tm2- and Tm4-to-T5 synapses and of nAChRα7 in Tm9-to-T5 synapses. Knock-down of nAChRα4, nAChRα5, nAChRα7, or mAChR-B individually in T5 cells alters the optomotor response and reduces T5 directional selectivity. Our findings indicate a differential contribution of postsynaptic receptors to input visual processing and, thus, to the computation of motion direction in T5 cells.

Introduction

Neurotransmission is a key process of neuronal communication that entails the release of neurotransmitter from the presynaptic terminal and its binding by

receptors on the postsynaptic membrane. In *Drosophila melanogaster*, the

neurotransmitters acetylcholine, GABA, glutamate, dopamine, serotonin,

octopamine[1] and histamine[2] are used for various neuronal tasks. Amongst them, the

prime excitatory neurotransmitter is acetylcholine[3–5], which can bind to a large

number of structurally diverse acetylcholine receptors (AChRs)[6–9]. How this

structural variety supports specific neural computations is not understood.

AChRs comprise two categories in bilaterians and cnidarians[10]: the fast ionotropic

nicotinic receptors (nAChRs) and the slow metabotropic muscarinic receptors

(mAChRs). nAChRs are activated by acetylcholine (ACh) and the agonist nicotine

and belong to the cys-loop ligand-gated ion channel superfamily, forming functional

pentamers of various subunit stoichiometries, homomeric or heteromeric. In

Drosophila the pentameric nAChR formation is achieved from a pool of seven α and

three β subunits[11–13]. mAChRs are G protein-coupled receptors (GPCRs), activated

by ACh and the agonist muscarine and comprise three types, A, B and C in

Drosophila[14–16]. These mAChR types activate different signaling cascades, with the

mAChR-A and mAChR-C being coupled to the $G_{q/11}$ signaling cascade and the

mAChR-B to the $G_{i/0}$[15,17,18]. Although the variety of AChRs has been thoroughly

addressed at the biochemical level[18,19], our understanding at the circuit level in terms

of neural computations and their behavioral output is only starting to advance[14,15,20–

23].

Among sensory modalities, vision contributes the most synaptic input to the

Drosophila central brain[24]. Visual processing starts at the two compound eyes,

whose retinotopic organization leads to a repetitive circuitry across its approximately

850 facets[25,26]. The major neuroanatomical area dedicated to visual information

processing is the optic lobe, which is divided in four neuropils, the lamina, the

medulla, the lobula and the lobula plate. From all the different visual cues such as

color and polarization vision[27], motion vision has been the one thoroughly

investigated, mainly due to its importance in courtship and navigation of the

individual[28–30]. The photoreceptor-derived information is split into two pathways: the

ON-pathway for luminance increments and the OFF-pathway for luminance

decrements[31,32]. Along the motion vision circuitry, the first neurons that compute

directional information are T4 cells in the ON-pathway and T5 cells in the OFF-
pathway[33,34].

Both T4 and T5 cells share the same bifurcation morphology, with the dendrite giving
rise to the axonal terminal fiber and to the soma fiber[35]. The T4 dendrites are located
in medulla layer ten (M10) and the T5 dendrites in lobula layer one (LO1). T4 and T5
cells exist in four subtypes per column: a, b, c and d. Each subtype has its dendrite
oriented in one of the four cardinal directions and responds selectively to front-to-
back, back-to-front, upward or downward motion, respectively[34–37]. Furthermore,
each subtype has its axonal terminals in one of the four layers in lobula plate[35]. For
their directional responses, T4 and T5 cells rely on the visual information they
receive from input neurons. The major upstream columnar neurons of T4 cells are
the excitatory cholinergic Mi1 and Tm3 neurons, the glutamatergic Mi9 and the
inhibitory GABAergic CT1, C3 and Mi4 neurons[38–46]. For T5 cells, the main columnar
input neurons are the excitatory cholinergic Tm1, Tm2, Tm4 and Tm9 cells and the
inhibitory GABAergic CT1 neuron[40,42–44,47–52]. T4 and T5 cells collect their inputs in a
distinct spatial order[40] reflecting their specific directional tuning[53,54].

T4 and T5 cells achieve their response selectivity for a particular direction by a
combination of two mechanisms: preferred direction enhancement and null direction
suppression[54,55]. In both processes, the luminance signal from one point in visual
space is delayed and interacts with the instantaneous signal from a neighboring
image point. For preferred direction enhancement, the delayed signal is from the
preferred side of the cell, i.e. where a stimulus moving in the preferred direction
enters the receptive field of the cell, and amplifies the central signal. For null
direction suppression, the delayed signal is from the opposite, the so-called null side,
and is used to suppress the central signal. In T4 cells, preferred direction
enhancement is biophysically implemented by a release from shunting inhibition,
mediated by the glutamatergic, inhibitory OFF-center cell Mi9, which amplifies the
central excitatory input from Mi1 and Tm3 neurons[45,56]. For null direction
suppression, the inhibitory neurons CT1, Mi4 and C3 act in concert to inhibit the Mi1-
Tm3 signal. In T5 cells, the CT1 cell is responsible for null direction suppression[51]. It
is, however, unclear how the different cholinergic inputs accomplish a preferred
direction enhancement. While the functional consequences of the different T5

cholinergic input neurons have been addressed in the past[48,52,57], the functional

implications of a possible input-specific AChR composition on T5 dendrites have not

been considered.

In this study, we address the structural and functional variety of different AChRs in

T5 cells. We first show the existence of the nAChRα1, nAChRα4, nAChRα5 and

nAChRα7 subunits on T5 dendrites. We target synaptic markers to each T5

columnar cholinergic input and find a differential distribution of AChRs in T5

postsynaptic sites. Through a combinatorial approach of neuronal input spatial wiring

and synaptic receptor mapping, we demonstrate the complex columnar cholinergic

spatial wiring on T5 dendrites. Finally, we show that the nAChRα4, nAChRα5 and

nAChRα7 subunits, as well as the mAChR-B type, differentially affect the optomotor

response and the directional responses of T5 cells.

Results

Expression of AChRs in T5 cells

Information about the expression of AChRs in T5 cells is available at the RNA

level[43,58,59], but is lacking at the protein level. We first evaluated the endogenous

expression of the nAChRα1, nAChRα2, nAChRα4, nAChRα5, nAChRα6 and

nAChRα7 subunits on T5 dendrites (Figure 1A,B). We used nAChRα subunits

endogenously tagged with an enhanced version of the green fluorescent protein

(EGFP) generated by Pribbenow et al.[22], thus minimizing the mistargeting error that

can occur with receptor overexpression and allowing for the assessment of protein

expression levels. We verified the localization of nAChRs in T5 neurons by co-

localizing the receptor puncta with the membrane-bound tdTomato expressed in T5

dendrites. We found the nAChRα1, nAChRα4, nAChRα5, nAChRα6 and nAChRα7

subunits on T5 dendrites, while the nAChRα2 subunit was not detected (Figure 1C-

I). To ensure that the lack of nAChRα2 expression was T5-specific, we confirmed the

nAChRα2 expression in other LO layers in the optic lobe (Figure S1A). By

quantifying the nAChRα subunit density in regions of interest (ROIs) on T5 dendrites

in LO1, we observed a prevalence of the nAChRα5 subunit and low co-localization

levels of the nAChRα2 and the nAChRα6 subunits with T5 dendrites, which is in

agreement with previous nAChRα RNA expression levels[43,58] (Figure 1J, Figure

S1B-D). A recent study[60] also confirmed the T5 dendritic localization of the nAChRα3

and nAChRβ1 subunits, which together with our nAChRα1, nAChRα4, nAChRα5,

nAChRα6 and nAChRα7 subunit data, emphasizes the large variety of nAChRs

present on T5 dendrites. Next, we sought to identify the nAChRα expression patterns

in the T5 axonal terminals by searching for the co-localization of receptor puncta with

the membrane-bound tdTomato expressed in T5 terminals. Both T4 and T5 neuronal

populations were targeted. Due to the lack of neuropile distinction between T4 and

T5 axonal terminals, our results cannot be attributed explicitly to T5 cells. The

nAChRα1, nAChRα5, nAChRα6 and nAChRα7 subunits were detected in T4-T5

axonal terminals, while the nAChRα2 and nAChRα4 were not (Figure S2A-G).

Additionally, we observed the reduction of the nAChRα1, nAChRα5 and nAChRα7

subunits in LOP after a T4-T5 specific subunit knock-down, indicative of their T4-T5

axonal synaptic localization (Figure S2J, S3A,C,D).

To study the mAChR expression on T5 dendrites, we used a line expressing the

mAChR-B type under control of UAS. We observed the overexpressed mAChR-B

protein localizing exclusively on T5 dendrites (Figure 1K, S2H). The encountered

dendritic T5 mAChR-B type targeting might be the outcome of ectopic expression

leading to dendritic protein targeting. We reasoned that if mAChR-B was mistargeted

to T5 dendrites, this mistargeting would be observed in other cells and that over-

expression of mAChR-B with a pan-neuronal driver would result in the presence of

ectopic mAChR-B in all the dendrites of neurons in the fly brain (Figure S2I). Instead,

the mAChR-B type followed an enrichment in layer 10, where T4 dendrites reside,

and in LO1, where T5 dendrites reside, while being absent from other medulla layers

and the lobula plate. In summary, the dendritic and axonal terminal protein

expression patterns differ between ionotropic and metabotropic AChRs and even

within the same AChR (Figure S2K).

To investigate the expression of those AChRs where we lacked endogenously

tagged lines, and to verify the expression profiles reported in Figure 1, we used

enhancer-trap Gal4 lines[61] (Figure S4A-L). By comparing RNA sequencing

datasets[43,58] with the enhancer trap and the endogenously tagged experiments here

and in previous studies[60], we concluded that the nicotinic nAChRα1, nAChRα3,

nAChRα4, nAChRα5, nAChRα7 and nAChRβ1 subunits as well as the muscarinic

mAChR-A and mAChR-B types are expressed in T5 cells, while the nAChRα2,
nAChRα6, nAChRβ2 and nAChRβ3 subunits exhibit low expression levels or are not
expressed at all (Figure S4M). Transcriptomics suggest low mAChR-C expression
levels in the fly brain[14,20]. Together our data show that the dendritic and axonal
protein localization patterns of ionotropic and metabotropic AChRs in T5 cells are
diverse and complex.
176
Morphological identification of cholinergic T5 input neurons and synapse
visualization
The nAChRα1, nAChRα4, nAChRα5 and nAChRα7 subunits might contribute to the
directional responses of T5 subtypes by localizing to all cholinergic synapses linking
Tm1, Tm2, Tm4 and Tm9 to T5 dendrites, or to only a subset of these. To discern
between these possibilities, we assessed the protein localization of nAChRα
subunits in specific synaptic pairs. We first needed genetic access to the four T5
cholinergic neuronal inputs: Tm1, Tm2, Tm4 and Tm9. The morphology of these
input neurons has been extensively studied via Golgi staining[35] and electron
microscopy reconstructions[39,40,47,50,62−64]. We used this previously attained EM
morphological knowledge to cross-validate our light microscopy screening of
potential Tm-specific enhancer lines (Figure 2A,D,G,J). Both Gal4 and LexA driver
lines[65,66], previously used in functional studies[48,51], were tested and their efficacy was
assessed in terms of neuronal specificity and whole population targeting. Neuronal
specificity was evaluated based on the arborization pattern of each neuronal type
and the lobula layer where the axonal terminals are localized. To achieve sparse and
stochastic neuronal labelling, we used the Multi-color FlpOut approach[67] and
observed axonal terminals in lobula layer 1 for Tm1 (Figure 2A), up to layer 2 for
Tm2 (Figure 2D), up to layer 4 for Tm4 (Figure 2G), and in layer 1 for Tm9 (Figure
2J), as expected from previous studies[35,40]. To exclude the developmental bias as
well as the non-neuronal targeting induced by the heat-shock-activated flipase[68] in
the Multi-color FlpOut approach (Figure 2G), we supplemented our single cell
labelling with whole population labelling of the Tm-Gal4 lines (Figure 2B,E,H,K). This
approach revealed the Gal4 driver lines expressing in a cell-type specific way in the
whole population of Tm1, Tm2, Tm4 and Tm9 cells, respectively. For the respective
LexA[66,69] driver lines, the whole Tm1 population was successfully targeted, but after
comparing the results with the analogous Gal4 driver line, dendritic ramifications in

layer 4 that do not correspond to Tm1, were observed (Figure 2C). The Tm2, Tm4

and Tm9 LexA lines displayed sparse neuronal labelling, indicating that the driver

lines could not target the whole neuronal population (Figure 2F,I,L). Based on this

morphological screening, we therefore chose the Gal4 driver lines for the Tm-T5

synapse visualization.

To visualize synapses between Tm1, Tm2, Tm4, Tm9 and T5 cells, we screened for

the available reporter tools. First, we sought to explore methods of trans-synaptic

labelling such as GFP Reconstitution Across Synaptic Partners (GRASP)[70], where

the GFP_{1-10} fragment is targeted to the cell membrane of the presynaptic

compartment, while the GFP_{11} fragment is targeted to the cell membrane of the

postsynaptic compartment. The targeted GRASP (t-GRASP)[71] allowed to capture all

synapses between Tm and T5 cells (Figure S5A). The presence of reconstituted

GFP signal in the inner optic chiasm, likely due to the ectopic reconstitution in the

Tm4-T5 synapses, led us to exclude this genetic method. The next reporter tool was

that of the activity GRASP (syb-GRASP)[72]. The reconstituted GFP signal was

indicative of the given synaptic activity at the point of the dissection, as the GFP_{1-10}

fragment is tethered to the vesicular protein synaptobrevin, thus resulting in the syb-

GRASP reconstitution being sparser than the t-GRASP reconstitution (Figure S5B).

This approach did not allow for a quantitative analysis; therefore syb-GRASP was

not chosen for our experiments. Finally, the truncated version of the presynaptic

marker Brp, one of the major protein participants of the protein-dense T-bar[73,74],

named Brp[short 75,76] was also tested (Figure S5C). This Brp[short]::GFP line labelled only

the presynaptic sites of all the Tm synapses, regardless of their activity, without

labelling their postsynaptic sites. In conclusion, our reporter tool screening for

synaptic labelling pointed to Brp[short] as a suitable marker to evaluate the synaptic

localization of the nAChRα receptor subunits.

To validate the synaptic specificity of Brp[short], we assessed two different, well

described synapses: the cholinergic Tm3-T4 and the glutamatergic Mi9-T4 synapse

(Figure S6A,B). Brp[short::mCherry 77] successfully showed the expected lack of trans-

synaptic alignment with the GABAergic receptor Rdl in Tm3-T4 synapses (Figure

S6A), while it revealed trans-synaptic alignment with the glutamatergic receptor

GluClα[45] in Mi9-T4 synapses (Figure S6B). To confirm the synaptic localization of

Brp$^{short::mStraw}$, a chimera that has the same truncated form of Brp as in Brp$^{short::mCherry}$

but is instead fused with mStrawberry by Owald et al.,[78], we used an antibody

against the endogenous Brp having affinity to amino acids absent from the truncated

Brp^{short} form (Figure S6C). This approach revealed the pre-synaptic alignment of the

endogenous Brp (Anti-Brp) with the Tm1, Tm2, Tm4 and Tm9 boutons

(Brp$^{short::mStraw}$). These results, together with previous studies[75], speak in favor of

Brp$^{short::mStraw}$ as a valid pre-synaptic marker in our circuit of interest.

The synaptic localization of the nAChRα5 subunit differs among the four
cholinergic inputs

With Brp$^{short::mStraw}$ as a presynaptic marker, we evaluated the localization of the

most highly expressed subunits, nAChRα1, nAChRα4, nAChRα5 and nAChRα7,

along the Tm-T5 synapses on T5 dendrites. We focused on the Tm1, Tm2, Tm4 and

Tm9 synapses in LO1, the major cholinergic inputs to T5. The regions of interest

were restricted to LO1 so as to include the Tm-T5 synaptic connections (Figure

S7A). We found that the nAChRα1, nAChRα4, nAChRα5 and nAChRα7 subunits

localize to Tm1, Tm2, Tm4 and Tm9 synapses in LO1 (Figure 3A-D, Figure S7B).

The synapses of Tm cells in LO1 might account for other output neurons apart from

T5 cells. To test this, we used the FlyWire connectome data explorer (Codex) for the

full adult fly brain (FAFB) dataset[62–64] in order to find the output neurons of the Tm1,

Tm2, Tm4 and Tm9 synapses in LO1. This analysis revealed T5 cells as the

principal output neuronal type across all four Tm cell types in LO1 (Figure S8A).

Therefore, the quantification of the nAChRα subunit synaptic localization in LO1

principally corresponded to Tm-T5 synapses. Importantly, we found that not all Tm

input neurons had the same synaptic receptor profiles. Tm1-, Tm2- and Tm4-to-T5

synapses primarily used the nAChRα5 subunit, whereas Tm9-to-T5 synapses mainly

used the nAChRα7 subunit (Figure 3E-H). As the GABAergic neuron CT1 was a

considerable Tm9 output in terms of synapse number in LO1 (Figure S8A), we

wondered if the Tm9-nAChRα7 subunit trans-synaptic alignment (Figure 3H) could

actually be attributed to Tm9-T5 synapses as well. We generated new flies to employ

a C-RASP approach. C-RASP follows the activity GRASP principles, but instead of a

GFP$_{1-10}$ fragment uses a CFP$_{1-10}$ fragment. This method showed the nAChRα7

subunit localization in Tm9-T5 synapses (Figure S8B).

Whether Tm1, Tm2, Tm4 and Tm9 cells differ from one another by differentially using muscarinic receptors as well, is currently unknown. Overexpression of the mAChR-B type in T5 cells led to the mAChR-B localization in Tm9-T5 synapses (Figure S9A), which was more evident when compared to the nAChRα5 subunit (Figure S9B). The apparent lack of a generalized cellular mistargeting of the mAChR-B overexpression line (Figure S2I), does not exclude the existence of synaptic mistargeting events. Consequently, the Tm9-mAChR-B results are indicative rather than definitive. Overall, our nAChRα subunit mapping across Tm synapses in LO1 revealed that Tm1, Tm2 and Tm4 cells differ from Tm9 at the postsynaptic level[48,57].

Spatial organization of Tm-nAChRα synapses in T5 dendrites

The compartmentalization of inputs and subsequently of receptors on T5 dendrites is essential for the T5 direction-selectivity[53,54]. Thus far, receptor localization in Tm-to-T5 synapses has been addressed by combining single-cell receptor localization and neuronal input spatial wiring information[60,79]. To advance this, we analyzed both T5a and T5c dendrites as representatives of the two directional systems from the FAFB[62] dataset via the FlyWire (flywire.ai)[63,64] interface. Manual proofreading of predicted synapses[80,81] revealed the highest number of Tm9 synapses, compared to Tm1, Tm2 and Tm4, in both T5 subtypes as previously reported[40,51] (Figure 4A,B). To account for these Tm-to-T5 differences in synapse counts, we measured the synapse occupancy index (Figure 3E-H). Next, we sought to identify the spatial wiring of the four Tm neurons on T5 dendrites. Tm1, Tm2, Tm4 and Tm9 synapses localize to the central dendritic compartment, with Tm9 synapses extending to the distal compartment as well, as previously shown[40,51] (Figure 4C,D). Together with our subunit synaptic localization results, we conclude that the nAChRα1, nAChRα4 and nAChRα7 subunits localize across the dendrite to Tm1, Tm2, Tm4 and Tm9-T5 synapses, while the nAChRα5 subunit localizes primarily in Tm1, Tm2 and Tm4-T5 synapses in the central compartment of T5 dendrites (Figure 4E).

AChRs differentially affect the optomotor response

When stimulated by rotatory visual motion, flies perform corrective steering maneuvers in their flight or walking trajectories. This behavior is called the optomotor response[30,82,83] and T4 and T5 cells are indispensable for its manifestation in the ON

and the OFF pathway accordingly[84–87]. Therefore, molecular alterations that are

crucial for the proper T5 function should be reflected in the fly's optomotor response

behavior. We used reversed genetics to test the role of AChRs in the OFF-motion

pathway. Using RNAi knock-down, we tested whether nAChRα1, nAChRα4,

nAChRα5, nAChRα7 and mAChR-B were required for the optomotor response on

the 'fly-on-ball' set up. Given that each receptor has a unique developmental onset of

expression (Figure S1D), we used a T4-T5 driver line that was active from the third

larval stage onwards[84,88] and we raised flies in 29°C[89,90] in order to achieve a strong

knock-down of protein levels. The driver line targeted all T5 subtypes and allowed for

the RNAi induction prior to the peak of AChR expression (Figure S3G,H). Finally, we

confirmed the RNAi-induced reduction in AChR subunit expression with

immunohistochemistry (Figure S3A-F).

Flies walking on an air-suspended ball were exposed to a set of OFF edges moving

in sixteen directions, which elicit a typical profile of angular walking velocities, with

the highest amplitudes to horizontal motion and lowest amplitudes to vertical

motion[45] (Figure 5A). Knock-down of the nAChRα1 subunit did not alter the angular

velocity response profile to the OFF edges (Figure 5B,G). Knock-down of the

nAChRα4 subunit or the mAChR-B type resulted in an increase of the angular

velocity in response to OFF edges moving in both horizontal directions (Figure

5C,F,G). In contrast, knock-down of the nAChRα5 and the nAChRα7 subunits led to

a decrease in the angular velocity amplitude in response to horizontal moving edges

(Figure 5D,E,G). The nAChRα3 and nAChRβ1 subunits were recently found to

localize on T5 dendrites[60], while the mAChR-A type exhibited high RNA expression

levels[43,58] and expression in T4-T5 cells based on our enhancer-trap screen (Figure

S1C,D, S4K). Knock-down of nAChRα3, nAChRβ1 and mAChR-A did not affect the

angular velocity profile in response to OFF edges (Figure S10A-E). However, the

RNAi efficiency against the nAChRα3, nAChRβ1 and the mAChR-A could not be

verified. Therefore, these AChRs should not be excluded from playing an important

role in the optomotor response. In summary, we found that AChRs play different

roles in the optomotor response.

Directional tuning of T5 cells changes after AChR knock-down

The nAChRα5 and nAChRα7 subunits localize both on T5 dendrites and in T5
axonal terminals (Figure S2K). The effects on the behavioral responses to OFF-edge
horizontal motion that were observed upon nAChRα5 and nAChRα7 knock-down
(Figure 5D,E,G) could either be due to functional roles in T5 dendrites or in T5
axonal terminals. To discern between these two possibilities, we measured calcium
responses in T5 axonal terminals. We performed two-photon calcium imaging in T5c
neurons, which are sensitive to upward motion as the preferred motion direction[34].
nAChRα4, nAChRα5, nAChRα7 or mAChR-B knock-down flies were stimulated by
dark edges moving in sixteen different directions, while calcium responses were
monitored in T5c axonal terminals in layer 3 of the lobula plate (LOP3) (Figure 6A,
Figure S11A-C). As in the behavioral experiments, RNA interference was induced
prior to peak receptor expression with the use of a T4-T5 promoter that is active from
the third larval stage[88], and raising flies in 29°C[89,90]. Knock-down of the nAChRα4
subunit led to a slight increase in calcium responses in response to the non-
preferred direction (270°) of T5c neurons (Figure 6B). Similarly, knock-down of the
nAChRα5 subunit resulted in higher calcium responses to the edge moving in the
non-preferred direction (Figure 6C), suggesting nAChRα5, and to a lesser extent,
nAChRα4 contribute to directional tuning in T5 neurons. Knock-down of the
nAChRα7 subunit resulted in an increase in calcium responses to both the null and
preferred motion direction (Figure 6D). Knock-down of the mAChR-B caused an
overall increase of calcium T5 responses across all presented stimuli directions
(Figure 6E). Individual fly responses shown in Figure S11D, show that the nAChRα4
and nAChRα5 subunit knock-down flies show high inter-fly variation, whereas the
nAChRα7, mAChR-B and mCherry RNAi flies exhibit more stable response patterns.
In general, the nAChRα4, nAChRα5 subunits and mAChR-B type significantly
influenced the T5 directional tuning (Figure 6F). Together these experiments show
that an interplay of nicotinic and muscarinic receptors shapes the T5 directional
responses.
367
Cholinergic neurotransmission in T5 cells takes place in two distinct locations: on the
dendrite in LO1, where T5-T5 dendro-dendritic synapses[40] reside, and on the axonal
terminal niche in LOP3, where T4-T5 and T5-T5 axo-axonic synapses reside[91].
Calcium responses in nAChRα5 subunit knock-down flies exhibited different profiles
in LO1 and LOP3 (Figure S12A-C). Calcium responses in T5 dendrites were of lower

amplitude, while responses from T5 axonal terminals increased in response to the
non-preferred direction (270°) in nAChRα5 knock-down compared to control
mCherry RNAi flies (Figure S12B,C). Such differences in calcium responses imply
distinct roles of AChRs in the two cholinergic neurotransmission niches of T5
neurons.

Discussion

In this study we investigated the cholinergic neurotransmission onto T5 neurons of
Drosophila and found that T5 cells express a variety of AChRs on their dendrites,
which are surprisingly differentially distributed across the cholinergic Tm1-, Tm2-,
Tm4-, Tm9-to-T5 synapses. This acetylcholine receptor variety translated into a
functional variety, as portrayed by both the fly optomotor response and the T5
directional tuning.

In mammals, nicotinic AChRs comprise nine α subunits (α1-7,9,10), four β subunits
(β1-4), one γ, one δ and one ε subunits[92], whilst muscarinic AChRs consist of five M
(M1-5) types[93]. This AChR variety is also found, but to a lesser extent, in *Drosophila*
melanogaster, with its ten nAChR subunits and three mAChR types. Does this
structural variety allow for differences in neuronal function? The nAChRα5 subunit
was found to be required in M4/6 mushroom body output neurons (MBONs) for
coding immediate appetitive memories[22]. The mAChR-A type was observed on the
dendrites of Kenyon cells, mediating aversive olfactory learning in adult flies[14].
Moreover, a study of the simple visual neural circuit in *Drosophila* larvae showed that
the glutamatergic interneuron glu-lOLP mediates OFF detection via mAChR-B
signaling[23], while the interplay of nAChRs and mAChRs in regulating acetylcholine
modulation of larval locomotion has been pharmacologically assessed with bath
application of AChR agonists and antagonists[15]. Our study provides another example
of diverse AChR functions in neuronal circuits, where an interplay between nicotinic
and muscarinic receptors is observed in the visual system of the adult fly.

AChR expression on T5 dendrites

For a long time, the expression of AChRs in the visual system of the fruit fly has
been under investigation[60,79,94–96]. Advancements in genomics and molecular
engineering now allow access to the AChRs of interest by endogenously tagging

them with fluorescent proteins[22,60,79]. This allows for qualitatively and quantitatively

assessing the AChR neuronal expression, and avoiding the ambiguity that was

introduced via combinatorial studies of electrophysiology and antagonist application

in terms of qualitative AChR identification[97]. We identified the nAChRα1, nAChRα4,

nAChRα5 and nAChRα7 subunit protein localization on T5 dendrites (Figure 1). The

nAChRβ1 subunit has recently been found to co-localize with the nAChRα1 subunit

on T5 dendrites[60]. Whether this co-localization depicts a pentameric co-organization

as previously speculated in lobula[94], remains to be proven, but it certainly indicates

the stoichiometrical nAChR complexity[98,99] that exists in the optic lobe. For the

mAChR-mediated cholinergic neurotransmission, our results together with previous

RNA-seq studies[43,58,59] indicate the expression of the mAChR-B type in T5 cells.

nAChRα subunits in Tm-to-T5 synapses

Sparse neuronal labelling allowed us to verify the explicit genetic access to Tm1,

Tm2, Tm4 and Tm9 neurons, which then could be combined with synaptic labelling

by Brp. While mapping the most highly expressed subunits nAChRα1, nAChRα4,

nAChRα5 and nAChRα7 across Tm1, Tm2, Tm4 and Tm9 synapses in LO1, we

found the nAChRα5 subunit preferentially localizing to Tm1, Tm2, Tm4 rather than

Tm9 synapses, implying that the functional differences observed among Tm1, Tm2,

Tm4 and Tm9 cells[48,57] are extended to the postsynaptic receptor level. Tm9 is

functionally different from Tm1, Tm2 and Tm4 by acting as a low-pass filter[48,49,57]

and Tm9-to-T5 synapses use a different nAChR set for their cholinergic

neurotransmission. This suggests that the preferred direction enhancement in T5

cells might not only be shaped by the different time-courses of the input signals they

receive, but it additionally relies on postsynaptic molecular mechanisms. Moreover,

the mAChR-B localization in Tm9-T5 synapses, together with the nAChRα1,

nAChRα4, nAChRα7 and to a lesser extend nAChRα5 localization, points towards

an interplay between ionotropic and metabotropic receptors. The role of such

receptor interactions is still unclear, but recent examples indicate that it might

potentiate the nicotinic AChR responses[20].

Attempts to test the role of neuronal input compartmentalization onto T5 dendrites

has led to combinatorial approaches, where the spatial wiring of inputs together with

the receptor dendritic localization were addressed. Together with previous

reports[40,51], our EM analysis shows a spatial overlap of Tm synapses on T5 dendrites, specifically on the central dendritic compartment. It is therefore risky to draw conclusions about receptor localization in Tm-to-T5 synapses via single-cell receptor localization experiments, especially due to the shared cholinergic nature of the input neurons and their spatial overlap. In this study, we introduce a new methodology, where EM-derived neuronal input spatial wiring and synaptic receptor mapping are combined to extract pre-, post-synaptic and spatial wiring information. Ultimately, single synapse-to-single dendrite approaches will improve the resolution of connectivity studies, leading to a T5 AChR-connectome[100].

AChRs roles in the OFF-motion pathway

Do these individual nAChR subunits and mAChR types differentially control T5 motion detection? We found that both ionotropic and metabotropic AChR signaling contribute to T5 function at the behavioral and neuronal activity levels, implying that an unforeseen complexity of cholinergic neurotransmission likely underlies motion detection in the OFF pathway. Notably, the nAChRα5 and nAChRα7 localization in T5 axonal terminals in addition to dendrites, might result in behavioral alterations deriving from other than dendritic computations. As the behavioral output involves many T5 downstream partners[83,101,102], compensatory mechanisms might also be in place, potentially masking immediate effects in T5 cells.

Calcium imaging in T5c neurons revealed an increase in the null direction responses among all nAChRα4, nAChRα5, nAChRα7 and mAChR-B knock-down flies. In contrast, the nAChRα7 or mAChR-B knock-down led to amplified preferred direction responses. We expected that the nAChRα5 subunit knock-down, as being the most highly expressed subunit along Tm1-, Tm2-, Tm4- and Tm9-T5 synapses, would have severely attenuated the T5 preferred direction responses. However, this was not the case, suggesting an important role for the nAChRα1, nAChRα4 and nAChRα7 subunits in preserving Tm1-, Tm2-, Tm4- and Tm9-to-T5 function. Lastly, the mAChR-B knock-down-induced effects resemble that of a raised inhibition, which align with the receptor's reported inhibitory role[23]. Our results show that there is an interplay of nicotinic and muscarinic AChRs taking place on T5 dendrites. The distinct stoichiometries as well as the ionic properties of the different nAChR

474 subunits need to be identified in the future to help us better understand the molecular
475 mechanisms underlying direction selectivity in T5 cells.
476
Limitations of this study
Multiple aspects should be taken into account when interpreting RNA interference
experiments. Firstly, an AChR subunit compensation[103] might take place, concealing
the true receptor subunit function. To which extent such a mechanism is activated in
T5 neurons, and towards which subunits, is currently unknown. Secondly, the
imaging location (T5 dendrites in LO1 versus T5 axonal terminals in LOP) should be
considered. The T5-T5 dendro-dendritic synapses, the T5-T5 and T4-T5 axo-axonic
synapses[91] are cholinergic. Our RNA interference of nAChRα5 and nAChRα7
subunits will affect such synapses as well, which might confound our conclusions on
T5 dendrites. In an effort to attend to this, we observed different responses in T5
dendrites compared to T5 axonal terminals. The difference in dendrite and axon
responses could result from differences in the density of ionotropic AChRs and other
voltage gated calcium channels between the two T5 compartments or from an
uncharacterized influence of T5-T5 dendro-dendritic and T5-T5 and T4-T5 axo-
axonic synapses (Figure S12). Finally, even though we used standardized conditions
for our RNAi induction experiments, fly-to-fly variability was encountered in nAChRα4
and nAChRα5 knock-down flies (Figure S11). This led us to believe that the RNAi
efficiency may differ across individuals and compensation mechanisms at the
individual level might be activated.
496
497 **STAR+METHODS**
498
499 Detailed methods are provided in the online version of this paper and include the
500 following:
501
502 • **KEY RESOURCES TABLE**
503
• **RESOURCE AVAILABILITY**
o Lead contact
o Materials availability

o Data and code availability

• **EXPERIMENTAL MODEL AND SUBJECT DETAILS**

o Fly husbandry

• **METHOD DETAILS**

o Generation of UAS-CRASP line

o RNAi experiments

o Immunohistochemistry

o Confocal microscopy

o Behavior

o Two-photon microscopy

o Visual stimulation

o Trans-synaptic alignment

o EM analysis

• **QUANTIFICATION AND STATISTICAL ANALYSIS**

o nAChRα density

o Synapse occupancy

o RNAi efficiency

o Behavior

o Calcium imaging

o Statistical analysis

ACKNOWLEDGMENTS

The authors thank D. Owald, C.H. Lee, F. Pinto-Teixeira, the Bloomington
Drosophila Stock Center, the Vienna Drosophila Stock Center and the
Developmental Studies Hybridoma Bank for flies and antibodies. We are grateful to
the Princeton FlyWire team and members of the Murthy and Seung labs for
development and maintenance of FlyWire (supported by BRAIN Initiative grant
MH117815 to Murthy and Seung). We are thankful to B. Zuidinga for technical
assistance with behavioral experiments and calcium imaging analysis; M. Sauter for

technical assistance with fly dissections; S. Prech for technical assistance with
experimental setups; R.M. Vieira and J. Pujol-Martí for discussions. E. Samara is a
member of the Graduate school of Systemic Neurosciences (GSN) Munich. I.M.A.R.
was funded by LMU and the DFG grant nr RI 3644/2-1. This work was funded by the
Max Planck Society.
545

546 **AUTHOR CONTRIBUTIONS**
547

E.S. and A.B. conceived this study. E.S., T.S. and I.M.A.R. designed the
experiments. E.S. performed all experiments with the help of T.S. and M.B.L. T.S.
contributed to the receptor synaptic localization experiments and analysis. M.B.L
contributed to the optomotor response experiments. J.H. contributed to the calcium
imaging analysis. E.S. wrote the manuscript with the help of I.M.A.R. and A.B. All
authors have read and commented on this manuscript.
554

555 **DECLARATION OF INTERESTS**
556

557 The authors declare no competing interests.
558

559 **FIGURE TITLES AND LEGENDS**
560

Figure 1. AChR expression on T5 dendrites
(A) Schematic representation of T5 dendrites residing in lobula layer 1 (LO1) of the
Drosophila optic lobe (OL).
(B) Schematic representation of nicotinic ionotropic (magenta) and muscarinic
metabotropic AChRs (cyan) in *Drosophila*. Out of the seven nAChRα subunits, six
were tagged with EGFP[22]. The mAChR-B was tagged with the HA peptide tag
(C.H.Lee)
(C) OL (grey) with T4 and T5 dendrites (R42F06-Gal4>UAS-myr::tdTomato, blue) in
M10 and LO1 accordingly. Scale bar 40µm.
(D-I) nAChRα1, nAChRα2, nAChRα4, nAChRα5, nAChRα6 and nAChRα7
(nAChRα::EGFP, magenta) subunit expression on T5 dendrites (R42F06-
Gal4>UAS-myr::tdTomato, blue). Scale bar 10µm, inset 5µm.

(J) Density of nAChRα subunit expression on T5 dendrites across 30 ROIs in LO1
(optic lobes n=6). nAChRα3, nAChRβ1, nAChRβ2 and nAChRβ3 subunits were not
addressed. Subunits with the lowest expression levels are in grey. Data is mean ±
SEM.
(K) Overexpression of mAChR-B (UAS-mAChR-B::HA, cyan) on T5 dendrites
(R42F06-Gal4>UAS-myr::tdTomato, blue). Scale bar 10µm, inset 5µm.
579
Figure 2. Morphological identification of the four columnar T5 cholinergic input
neurons
(A) Electron microscopy (EM)(flywire.ai) reconstruction of a Tm1 neuron reveals two
loci of arborization in medulla. Schematic representation of the Tm1 axonal terminal
localization in LO1. Tm1-Gal4 expression (cyan) of the MultiColor FlpOut cassette
(V5, HA, FLAG) and axonal localization in LO. Scale bar 40µm and 10µm.
(B) Whole population labelling of Tm1 neurons (R74G01-Gal4>UAS-CD8::GFP).
Scale bar 40µm.
(C) Whole population labelling of Tm1-LexA (purple) neurons (VT041034-
LexAGAD>LexAop-myr::mCherry). Arrows represent additional dendritic
ramifications that lack in (B). Scale bar 40µm.
(D-F) As in (A-C), but for Tm2 neurons (VT012282-Gal4, VT058650-LexAGAD).
(G-I) As in (A-C), but for Tm4 neurons (R35H01-Gal4, R53C02-LexAGAD).
(J-L) As in (A-C), but for Tm9 neurons (VT065303-Gal4, R24C08-LexAp65).
594
Figure 3. Differential synaptic localization of nAChRα subunits in Tm synapses
(A-D) Trans-synaptic alignment of Tm1, Tm2, Tm4 and Tm9 boutons (Tm-Gal4>
UAS-Brp$^{short::mStraw}$, grey) with the nAChRα1, nAChRα4, nAChRα5 and nAChRα7
subunits (nAChRα::EGFP, magenta) in LO1. Arrows indicate the Brp$^{short::mStraw}$-
receptor punctum trans-synaptic alignment. Scale bar 1µm.
(E-H) Synapse occupancy of Tm1, Tm2, Tm4 and Tm9 boutons with nAChRα1 (pale
magenta), nAChRα4 (light magenta), nAChRα5 (magenta) and nAChRα7 (dark
magenta) subunits in LO1 (optic lobes n=4, ROIs N=20, for Tm4-nAChRα5 n=3,
N=15). Data is mean ± SEM.
604
Figure 4. Wiring of Tm1, Tm2, Tm4 and Tm9 on T5 dendrites

(A) Counts of Tm1, Tm2, Tm4 and Tm9 synapses in T5a dendrites (T5a cells n=5,

flywire.ai) (left). Counts of predicted (n=663) to proofread (n=479) Tm synapses in

T5a dendrites (right). Data is mean ± SEM.

(B) As in (A) for T5c dendrites (T5c cells n=5, flywire.ai) (left). Counts of predicted

(n=621) to proofread (n=488) Tm synapses in T5c dendrites (right). Data is mean ±

SEM.

(C) Dendritic localization of Tm synapses across the T5a anterior-posterior axis (T5a

cells n=5). Root is considered the first dendritic branch point.

(D) Dendritic localization of Tm synapses across the T5c dorso-ventral axis (T5c

cells n=5).

(E) Schematic representation of Tm-T5a dendritic synapses and the observed

postsynaptic receptor heterogeneity in terms of the most abundant nAChR subunit

as in Figure 3E-H.

Figure 5. Fly optomotor response post-AChR knock-down

(A) Schematic workflow representation for the behavioral assessment of AChR

knock-down flies.

(B) Trajectory (angular velocity) of nAChRα1 knock-down flies (R42F06-p65.AD;

VT043070-Gal4.DBD>UAS-nAChRα1-RNAi, pale magenta, flies n=20) while

presented to a sequence of upward to rightward to downward to leftward OFF edge

motion compared to the parental control flies (UAS-nAChRα1-RNAi, grey n=21). The

dark lines represent the mean ± SEM.

(C-F) Same as in (B) for the nAChRα4 (light magenta n=19, grey n=20), nAChRα5

(magenta n=18, grey n=21), nAChRα7 (dark magenta, n=21, grey n=20), mAChR-B

(cyan n=20, grey n=20) knock-down flies.

(G) Angular velocity of control and AChR knock-down flies. The normality of

distribution was assessed with the use of Shapiro-Wilk test. *$p<0.05$, **$p<0.01$,

***$p<0.001$. Data is mean ± SEM.

Figure 6. Directional tuning of T5 cells post-AChR knock-down

(A) Schematic workflow representation for the two-photon calcium imaging in AChR

knock-down T5 cells.

(B) Polar plot of the maximum responses (ΔF/F) across sixteen OFF edge directions

(º) of T5c cells for mCherry controls (black, flies n=7, ROIs N=85, shown as grey in

C, D and E) and nAChRα4 RNAi (light magenta, n=7, N=61) conditions (R39H12-Gal4>AChR-RNAi, VT50384-LexA>LexAop-IVS-GCaMP6m). The dark lines represent the mean (68% CI).

(C) Same as in (B) for nAChRα5 RNAi flies (magenta, n=7, N=82).

(D) Same as in (B) for nAChRα7 RNAi flies (dark magenta, n=7, N=60).

(E) Same as in (B) for mAChR-B RNAi flies (cyan, n=7, N=54).

(F) Directional tuning index (Ldir). Flies n=7. The normality of distribution was assessed with the use of Shapiro-Wilk test. *$p<0.05$, ***$p<0.001$. Data is mean ± SEM.

STAR⬜Methods

RESOURCE AVAILABILITY

Lead contact
Further information and requests for data should be directed to and will be fulfilled by the Lead Contact, Alexander Borst (alexander.borst@bi.mpg.de).

Materials availability
The study generated a new fly line and is available from lead contact upon request.

Data and code availability
- This paper analyzes existing, publicly available data from FlyWire (flywire.ai).
- Data and code for analysis are publicly available at the at the Edmond Open Research Data Repository of the Max Planck Society.
- Any additional information required to reanalyze the data reported in this paper is available from the lead contact upon request.

EXPERIMENTAL MODEL AND SUBJECT DETAILS

Fly husbandry
All flies were raised on corneal agar medium under standard conditions (25°C, 60% humidity, 12h light/dark cycle). For the interference experiments (RNAi), flies (control

and RNAi) at early pupa stages were transferred from 25 to 29°C. For the Multi-color
FlpOut experiments, flies at early pupa stages were heat-shocked for 30 minutes at
37°C.

METHOD DETAILS

Generation of UAS-CRASP line

The pQUAST-syb-spCFP1-10 vector[72] and the pJFRC7-20XUAS-IVS-mCD8::GFP
vector[104] were digested with XhoI and XbaI restriction enzymes and the syb-
spCFP1-10 DNA fragment was extracted and T4-ligated in the pJFRC7 vector. The
AGGCACACCGAAACGACTAA and CAGTTCCATAGGTTGGAATCTAAAA primers
were used for sequencing the successful insertion of the DNA fragment. Embryo
injections were performed by BestGene Inc (Chino Hills, CA, USA) in the VK00027
attp landing site.

RNAi experiments

To ensure high interference efficiency, given that both long dsRNA (nAChRα1,
nAChRα4 and nAChRβ1) and short shRNA (nAChRα3, nAChRα5 and nAChRα7,
mAChR-A, mAChR-B) were used, flies were raised (at early pupa stages) at
29°C[89,90] and the RNAi was induced prior to the receptor expression peak. Dicer was
not used for the long dsRNA experiments as previous studies[22,105,106] showed
positive AChR-RNAi induction without dicer co-expression. Our
immunohistochemical screening spoke in favor of the nAChRα1, nAChRα4,
nAChRα5, nAChRα7 and mAChR-B RNAi effectiveness (Figure S3).

Immunohistochemistry

Fly brains (aged 2-5 days, for G-and C-RASP experiments aged 10-12 days) were
dissected in cold phosphate-buffered saline (PBS) and fixed in 4%
paraformaldehyde (in PBS with 0.1% Triton X-100) for 24 minutes at room
temperature, followed by three 10-minute washes in PBT (PBS and 0.3% Triton X-
100). Brains were then incubated with primary antibodies in PBT for 48 hours at 4°C.
After being incubated for 2 hours at room temperature, brains were washed four
times for 15 minutes each in PBT and then incubated with secondary antibodies
diluted in PBT for 48 hours at 4°C. After being incubated for 2 hours at room

temperature and four 15 minutes PBT washes, brains were mounted for immediate
sample viewing. For the subunit mapping (Figure 3), brains were blocked in 5%
normal goat serum (NGS) in PBT for 1 hour at room temperature and then incubated
with the primary and secondary antibodies for 24 hours instead of 48 hours.
711

Confocal microscopy

Images were acquired in a Leica Stellaris 5 laser scanning confocal microscope with
a 20x glycerol 0.75 NA HC Plan-apochromat and 63x glycerol immersive 1.3 NA HC
Plan-apochromat objective at 2048 x 2048 x 0.4µm image resolution. The
deconvolution setting (Lightning) of this microscope was used for the acquisitions in
Figure 3. Image analysis was performed with Fiji (ImageJ)[107]. For the acquisitions in
Figure S4, a Leica SP8 laser scanning confocal microscope with a 63x oil immersive
1.3 NA HCX Plan-apochromat objective was used at 2048 x 2048 x 0.4µm image
resolution.
721

Behavior

To avoid behavioral effects introduced by sexual dimorphism[26], only female flies, 1-3
days old, were used in our experiments. Flies were attached to a pin post-cold
anaesthetization with the use of light-curing glue and dental curing light (440 nm,
New Woodpecker). Each locomotor recorder, out of the four that were
simultaneously used, consisted of an air-suspended sphere (6mm in diameter and
40 mg weight) made of polyurethane foam and coated with polyurethane spray[85,108].
A constant airflow by a rotary vane pump (G6/01-K-EB9L Gardner Denver Thomas)
allowed for the free rotation of the sphere on the ball-shaped sphere holder. The
sphere's rotation was tracked by two optical tracking sensors, focusing two 1-mm^2
equatorial spots at ±30° from the center of the infrared LED-illuminated (JET-800-10,
Roithner Electronics) sphere. Rotational data were tracked at 4 kHz and digitized at
200 Hz[109]. To achieve a successful fly positioning on the sphere, a camera (GRAS-
20S4M-C, Point Grey Research) was used. A custom-made Peltier system (QC-127-
1.4-6.0MS, Quick-Cool) controlled the airflow temperature at 34 ± 0.1°C, based on
the readouts deriving from a thermometer placed just below the sphere, so as to
ensure prolonged walking. A U-shaped visual arena comprising of three 120-Hz LCD
screens (Samsung 2233 RZ) (reaching a maximum luminance of 131 cd m^{-2}),
allowed for the presentation of visual stimuli spanning approximately 270° in azimuth

and 120° in elevation of the fly's visual field at a resolution of <0.1°. Panda3D was
used in Python v.2.7 for the visual stimulus generation[85].

Two-photon microscopy

For functional imaging experiments, we used a custom-build two photon laser
scanning microscope equipped with a 40x water 0.80 NA IR-Achroplan objective
(Zeiss)[34]. Flies (1-3 days) were cold-anaesthetized and the legs and thorax were
glued on a Plexiglas holder with the use of light-curing glue. The head was bend
down, fitted in an aluminum opening and the holder was clamped in a recording
chamber. Saline solution (PBS; 137 mM NaCl, 3 mM KCl, 8 mM Na_2HPO_4, 1.5 mM
KH_2PO_4, pH 7.3) was introduced and the posterior side of the fly's optic lobe was
exposed by a small incision of the head. Muscles, adipose tissue and trachea were
manually removed. Images were acquired at a 64 x 64-pixel resolution and a frame
rate of 15 Hz in Matlab R2013b (MathWorks) using ScanImage 3.8 software (Vidrio
Technologies, LLC).

Visual stimulation

For the behavioral experiments, OFF edges (50% contrast) moved at 16 evenly
spaced directions with a velocity of 60° s^{-1}, were randomized and crossed the whole
arena span within 5s. Each experiment lasted for ~55 min, including 35 trials of OFF
edges, of which the first 15 trials were excluded from analysis, as they were
appointed to temperature stabilization and fly accommodation. The same inclusion
criteria as in a previous study[45] were used.

For the calcium imaging experiments, we used a custom-built projector-based
arena[57], where two micro-projectors (TI DLP Lightcrafter 3000) projected visual
stimuli, with a refresh rate of 180 Hz and maximum luminance of 276 ± 48 cd/m^2,
onto the back of a cylindrical screen. The arena covered 180° in azimuth and 105° in
elevation of the fly's visual field. OFF edges (92% contrast) moved at 16 evenly
spaced directions with a velocity of 30° s^{-1}, were randomized and repeated four
times.

Trans-synaptic alignment

To assess the trans-synaptic alignment between the receptors and the synapses of
interest, we calculated the fluorescence intensity values (later normalized to the
maximum pixel intensity value) in a 1 µm line positioned across the synapse
punctum in a single z plane using Fiji-ImageJ[107]. Data analysis was performed with
Python v.3.9.18 with the use of seaborn 0.12.2, pandas 1.5.3 and numpy 1.23.3.

EM analysis

For the Tm-T5 wiring, we used the FAFB volume[62] via the FlyWire (flywire.ai)
proofreading environment[63,64] and chose five reconstructed T5a and five T5c
neurons from the right optic lobe (Table S3). T5 and Tm1, Tm2, Tm4, Tm9 identities
were verified by users' annotations and by comparison with previous electron
microscopy[40] and our light microscopy reconstructions. Synapse number Buhmann
predictions[80,81] between the T5 neuron of interest and its respective input neurons
were acquired from FlyWire's connectivity viewer. Cleft score was set to 50, so as to
eliminate synaptic redundancy resulting from the combination of synaptic
connections between the same neurons when their presynaptic locations are within
$100nm^2$. For each synapse, we sought four morphological markers: a. synaptic
vesicles, b. the protein dense T-bar structure, c. synaptic cleft, and d. postsynaptic
densities (or postsynaptic domains). Every synaptic locus was assessed in a volume
of five up to seven brain slices, 40nm each. Only synapses that displayed all four
morphological markers were included in our dataset. We treated proofread synapses
as points in T5 dendrite space by collecting the x,y,z coordinates of each T-bar (pre-
synapse). Synapse numbers in this study correspond to manually proofread
synapses.

For synapse distributions across the anterior-posterior and dorso-ventral dendritic
arborization axis (Figure 4C,D), we measured the Euclidean distance of every
synapse from the root (first branching point) of the dendrite. Distances in µm were
calculated by multiplying the y and z coordinates of each T-bar with 4 and 40
accordingly (voxel=4x4x40nm). To correct for anterior-posterior and dorsal-ventral
dendrite length differences, we normalized each synapse-to-root Euclidean distance
to the longest T5a and T5c dendrite.

For the identification of Tm1, Tm2, Tm4 and Tm9 synaptic partners in LO1 (Figure
S8), we used Codex[62,63,64]. We searched for all the Tm output neurons in five Tm1
(FlyWire ID: 720575940620976493,720575940608883465,720575940623583428,
720575940632600647,720575940621146733), Tm2 (FlyWire ID: 72057594063214
2904,720575940640453437,720575940640230259,720575940630551670,7205759
40622364961), Tm4 (FlyWire ID: 720575940637976666, 720575940620745163,
720575940621582401, 720575940626612228, 720575940620782171) and Tm9
(FlyWire ID: 720575940613374873, 720575940620814356, 720575940609254851,
720575940626242348, 720575940623771061) cells, and selected only those
outputs whose Tm synapses were restricted in LO1 (excluding all Tm medullary
output neurons). For the output neurons that only formed connections with Tm
neurons in LO1, we used the predicted synapse number, but for those output
neurons that made Tm connections in other LO layers as well, we manually
proofread the LO1 synapses. The reconstructed neurons for Figure 2 correspond to
FlyWire IDs 720575940621338038, 720575940614527742, 720575940616041926
and 720575940616813142 for Tm1, Tm2, Tm4 and Tm9 respectively.
823
824 **QUANTIFICATION AND STATISTICAL ANALYSIS**
825
826 **nAChRα density**
827 Receptor puncta that localized in T5 dendrites were quantified along 8μm x 8μm x
828 4μm voxels in Fiji-ImageJ[107]. The total number of puncta divided to the analyzed LO1
829 volume resulted in the nAChRα density.
830
831 **Synapse occupancy**
832 The total number of Brp-receptor co-localizing puncta in voxels of 8μm x 8μm x 4μm
833 in Fiji-ImageJ[107], was divided to the number of Brp puncta, resulting in the synapse
834 occupancy index.
835
836 **RNAi efficiency**
837 The mean grey value of ten 8μm x 8μm ROIs per optic lobe in LO1 and LOP (layer 1
838 and 2) was measured via Fiji-ImageJ[107] and was normalized by subtracting the mean
839 grey value of one 8μm x 8μm ROI in the inner optic chiasm.
840

Behavior

The optomotor response was quantified as the absolute average rotational velocity during 5 s of edge motion in each direction. Data analysis was performed with Python v.3.9.18 using seaborn 0.12.2, pandas 1.5.3 and numpy 1.23.3.

Calcium imaging

Calcium imaging data were analyzed as described in[57] with a custom written software in Python v.2.7. ROIs were drawn manually across LO layer 1 and LOP layer 3. Only ROIs that repeatedly responded to the given visual stimulus and exhibited consistent-across the four repetitions-responses, were included in the dataset. For each ROI, the time courses of relative fluorescence changes (ΔF/F) were calculated and responses to stimulus were baseline-subtracted and averaged over repetitions. Maximum responses were aligned to 90°. The directional tuning index (Figure 6F, Ldir) was calculated as the magnitude of the resultant vector divided by the sum of the individual vectors' magnitudes:

$$Ldir = \left| \frac{\sum_{\varphi} \overrightarrow{v(\varphi)}}{\sum_{\varphi} \overrightarrow{|v(\varphi)|}} \right|$$

Statistical analysis

Detailed analysis is reported in figure legends and table S2 and was performed in GraphPad Prism v.9.3.0.

SUPPLEMENTAL INFORMATION TITLES AND LEGENDS

Figure S1. AChR expression in T5 cells-Related to Figure 1

(A) nAChRα2 expression in LO. Arrows indicate the LO layers of expression. Scale bar 10μm, inset 5μm.

(B) Example ROIs in LO1 for quantification as in Figure 1J. Scale bar 10μm, inset 5μm.

(C) Bulk RNAseq of AChR expression (transcripts per million, TPM) in T5 cells and probability of expression (p)[43].

(D) scRNAseq of AChR expression in T4 (grey) and T5 (blue) cells across five developmental stages (hours correspond to APF-after puparium formation)[58].

Figure S2. AChR expression in T4-T5 axonal terminals-Related to Figure 1

(A) Schematic representation of T5 axonal terminals residing in lobula plate (LOP1-4) of the *Drosophila* optic lobe (OL).

(B-G) Expression of nAChRα1, nAChRα2, nAChRα4, nAChRα5, nAChRα6 and nAChRα7 (nAChRα::EGFP, magenta) in T4-T5 axonal terminals (R42F06-Gal4>UAS-myr::tdTomato, blue). Scale bar 10µm.

(H) Overexpression of mAChR-B (cyan) in T4-T5 axonal terminals (R42F06-Gal4>UAS-myr::tdTomato, UAS-mAChR-B::HA). Scale bar 10µm.

(I) Pan-neuronal expression of mAChR-B in OL (GMR57C10-Gal4>UAS-mAChR-B::HA). Scale bar 40µm.

(J) Normalized mean grey value of nAChR subunits in LOP (layers 1 and 2) before and after RNAi induction (optic lobes n=2, ROIs N=20, for nAChRα1 n=1). The normality of distribution was assessed with the use of Shapiro-Wilk test. $*p<0.05$, $****p<0.0001$. Data is mean ± SEM.

(K) Summarized expression patterns of nAChRα1, nAChRα2, nAChRα4, nAChRα5, nAChRα6 and nAChRα7 subunits and the mAChR-B type on T5 dendrites and in T4-T5 axonal terminals.

Figure S3. RNA interference against AChRs in T5 cells-Related to Figure 1 and 5

(A) Expression of the nAChRα1 subunit (nAChRα1::EGFP) in LO1 before (left panel) and after (right panel) the RNAi induction (R42F06-Gal4>AChR-RNAi). Scale bar 10µm.

(B-E) As in (A) for the nAChRα4, nAChRα5, nAChRα7 subunits and the mAChR-B type.

(F) Normalized mean grey value of AChRs in LO1 before and after RNAi induction (optic lobes n=2, ROIs N=20). The normality of distribution was assessed with the use of Shapiro-Wilk test. $**p<0.01$, $***p<0.001$, $****p<0.0001$. Data is mean ± SEM.

(G) Developmental onset (P18) of the R42F06 promoter used for the split T4-T5 line in all behavioral experiments (R42F06-Gal4> UAS-myr::tdTomato). Scale bar 40µm, inset 10µm.

(H) Split T4-T5 line (R42F06-p65.AD; VT043070-Gal4.DBD>UAS-CD4::tdGFP) used

in all behavioral experiments. Arrows indicate the four LOP layers. Scale bar 40µm,

inset 10µm.

Figure S4. Enhancer-trap verification of AChR expression in T4-T5 cells-

Related to Figure 1

(A-L) Enhancer-trap lines for the nAChRα1, nAChRα2, nAChRα3, nAChRα4,

nAChRα5, nAChRα6, nAChRα7, nAChRβ1, nAChRβ2, nAChRβ3 subunits and

mAChR-A and mAChR-B receptors (AChR-Mi or T2A-Gal4>UAS-CD8::GFP,

magenta and cyan). Signal overlap with T4-T5 somata (R42F06-LexA>LexAop-

CD8::RFP, blue) denoted the T4-T5 AChR expression. The nAChRα4 line did not

show expression across all brain areas. Images are z-projections. Scale bar 40µm,

inset 10µm.

(M) AChR expression in T5 cells across scRNA-seq[58], Bulk-seq[43], enhancer trap and

endogenously tagged receptor[60] experiments.

Figure S5. Screening of genetic approaches for synapse visualization-Related

to Figure 3

(A) t-GRASP expression in Tm1, Tm2, Tm4 and Tm9-T5 synapses (Tm1, Tm2, Tm4,

Tm9-Gal4>UAS-pre-t-GRASP,   VT25965-LexAp65>LexAop-post-t-GRASP).   Scale

bar 10µm. Ectopic reconstitution is observed in Tm4-T5 synapses.

(B) Activity GRASP (syb-GRASP) in Tm1, Tm2, Tm4 and Tm9-T5 synapses (Tm1,

Tm2, Tm4, Tm9-Gal4>UAS-nSyb-spGFP1-10, VT25965-LexAp65>LexAop-CD4-

spGFP11). Scale bar 10µm.

(C) Brp[short]::GFP expression in Tm1, Tm2, Tm4 and Tm9 boutons (Tm1, Tm2, Tm4,

Tm9-Gal4>UAS-Brp[short]::GFP). Scale bar 10µm.

Figure S6. Validation of Brp[short] as a precise presynaptic marker-Related to

Figure 3

(A) Trans-synaptic alignment between Brp[short::mCherry](blue) and the GABAergic Rdl

receptor (grey) in the cholinergic Tm3-T4 synapse (GMR13E12-LexA>LexAop-

Brp[short]::mCherry, R42F06-Gal4>UAS-Rdl::GFP). Scale bar 10µm.

(B) Trans-synaptic alignment between Brp$^{short::mCherry}$(blue) and the glutamatergic

GluClα receptor (grey) in the glutamatergic Mi9-T4 synapse (GMR48A07-

LexA>LexAop-Brp^{short}::mCherry, R42F06-Gal4>UAS-GluClα::GFP). Scale bar 10μm.

(C) Localization of Brp$^{short::mStraw}$ in Tm1, Tm2, Tm4 and Tm9 boutons (blue)

and endogenous Brp (grey) identified by the anti-Brp staining. Arrows correspond to

the signal overlap between the Tm-Brp^{short} and the Anti-Brp. Scale bar 5 and 1μm.

Figure S7. Localization of nAChRα subunits in Tm synapses-Related to Figure

3

(A) LO1 loci that were used for Figure 3A-D. Scale bar 10μm.

(B) Line profile analysis for the assessment of trans-synaptic alignment of

Brp$^{short::mStraw}$ puncta (grey) and nAChRα puncta (magenta) across Tm1, Tm2, Tm4

and Tm9 synapses (analyzed puncta n=10). The dark lines represent the mean

values.

Figure S8. Neuronal outputs of Tm1, Tm2, Tm4 and Tm9 cells in LO1-Related

to Figure 3

(A) Schematic representation of Tm1, Tm2, Tm4 and Tm9 axonal terminals in

relation to LO1. Primary Tm1, Tm2, Tm4 and Tm9 output neuronal types in LO1 (Tm

cells n=5, Codex). Data is mean ± SEM.

(B) C-RASP for the Tm9-T5 synapse visualization (Tm9-Gal4>UAS-IVS-syb-

spCFP1-10, VT25965-LexAp65>LexAop-CD4-spGFP11, grey) of the nAChRα7

subunit (nAChRα7::EGFP, magenta). Arrows correspond to Tm9-nAChRα7 trans-

synaptic alignment. Scale bar 10μm.

Figure S9. mAChR-B receptor localization in T5 dendrites-Related to Figure 3

(A) mAChR-B localization (R42F06-Gal4>UAS-mAChR-B::HA, cyan) in Tm9-T5

synapses (Tm9-LexA>LexAop-Brp^{short}::mCherry, grey). Scale bar 10μm.

(B) nAChRα5 subunit (nAChRα5::EGFP, magenta) localization pattern compared to

mAChR-B (R42F06-Gal4>UAS-mAChR-B::HA, cyan) in LO1. Scale bar 10μm.

Figure S10. Fly optomotor response after nAChRα3, nAChRβ1 and mAChR-A

knock-down-Related to Figure 5

(A) Schematic representation of the behavioral setup for the optomotor response

assessment in nAChRα3 (pale magenta), nAChRβ1 (violet) subunits and mAChR-A

(teal) type knock-down flies.

(B) Trajectory (angular velocity) of nAChRα3 knock-down flies (42F06-p65.AD;

VT043070-Gal4.DBD>UAS-nAChRα3-RNAi, pale magenta n=18) while presented to

a sequence of upward to rightward to downward to leftward OFF edge motion

compared to the parental control flies (UAS-nAChRα3-RNAi, grey n=20). The dark

lines represent the mean ± SEM.

(C-D) Same as previously for the nAChRβ1 (violet n=18, grey n=19) and mAChR-A

(teal n=20, grey n=20) knock-down flies.

(E) Angular velocity of control and AChR knock-down flies. The normality of

distribution was assessed with the use of Shapiro-Wilk test. Data is mean ± SEM.

Figure S11. Two-photon calcium imaging in T5 cells-Related to Figure 6

(A) Schematic representation of LO1 (where T5 dendrites reside) and of LOP3

(where T5c axonal terminals reside).

(B) Two-photon view of LOP3 (VT50384-LexA>LexAop-IVS-GCaMP6m).

(C) T4-T5c specific targeting (VT50384-Gal4>UAS-CD8::GFP). Arrows indicate the

four LOP layers. Scale bars 40 and 10µm.

(D) Polar plots of the maximum responses (ΔF/F) -along seven flies- across sixteen

OFF edge directions (^{o}) of T5c cells under nAChRα4 RNAi (light magenta),

nAChRα5 RNAi (magenta), nAChRα7 RNAi (dark magenta), mAChR-B RNAi (cyan)

and control mCherry RNAi (black) conditions.

Figure S12. Two-photon calcium imaging in T5 dendrites and T5 axonal

terminals-Related to Figure 6

(A) Schematic representation of LO1 (where T5 dendrites reside) and of LOP3

(where T5c axonal terminals reside).

(B) Two-photon view of the LO1 and LOP3 layers (VT50384-LexA>LexAop-IVS-

GCaMP6m).

(C) Polar plots of the maximum responses (ΔF/F) across sixteen OFF edge

directions (^{o}) of T5c cells under mCherry controls (black, flies n=5, ROIs N=10) and

nAChRα5 RNAi (magenta, n=5, N=13) conditions in LO1 (left) and in LOP3 (right).

The dark lines represent the mean (68% CI).

Table S1. List of fly experimental genotypes, related to STAR Methods.

Table S2. Statistical analysis, related to STAR Methods.

Table S3. FlyWire neuronal IDs, related to STAR Methods.

REFERENCES

1. Shin, M., Copeland, J.M., and Venton, B.J. (2018). *Drosophila* as a model system for neurotransmitter measurements. ACS Chem. Neurosci. *9*, 1872–1883. https://doi.org/ 10.1021/acschemneuro.7b00456.

2. Stuart, A.E. (1999). From fruit flies to barnacles, histamine is the neurotransmitter of arthropod photoreceptors. Neuron *22*, 431–433. https://doi.org/10.1016/S0896-6273 (00)80699-6.

3. Kolodziejczyk, A., Sun, X., Meinertzhagen, I.A., and Nässel, D.R. (2008). Glutamate, GABA and acetylcholine signalling components in the lamina of the *Drosophila* visual system. PLoS ONE *3*, e2110. https://doi.org/10.1371/journal.pone.0002110.

4. Yasuyama, K., and Salvaterra, P.M. (1999). Localization of choline acetyltransferase-expressing neurons in *Drosophila* nervous system. Microsc. Res. Tech. *45*, 65–79. https://doi.org/10.1002/(SICI)1097-0029(19990415)45:2<65::AID-JEMT2>3.0.CO;2-0.

5. Eckstein, N., Bates, A.S., Champion, A., Du, M., Yin, Y., Schlegel, P., Lu, A.K.-Y., Rymer, T., Finley-May, S., Paterson, T., et al. (2024). Neurotransmitter classification from electron microscopy images at synaptic sites in *Drosophila melanogaster*. Cell *187*, 2574-2594.e23. https://doi.org/10.1016/j.cell.2024.03.016.

6. Breer, H., and Sattelle, D.B. (1987). Molecular properties and functions of insect acetylcholine receptors. J. Insect Physiol. *33*, 771–790. https://doi.org/10.1016/0022-1910(87)90025-4.

7. Sattelle, D.B., Jones, A.K., Sattelle, B.M., Matsuda, K., Reenan, R., and Biggin, P.C. (2005). Edit, cut and paste in the nicotinic acetylcholine receptor gene family of *Drosophila melanogaster*. BioEssays *27*, 366–376. https://doi.org/10.1002/bies.20207.

8. Rosenthal, J.S., and Yuan, Q. (2021). Constructing and tuning excitatory cholinergic synapses: The multifaceted functions of nicotinic acetylcholine receptors in *Drosophila* neural development and physiology. Front. Cell. Neurosci. *15*, 720560. https://doi.org/10. 3389/fncel.2021.720560.

9. Thany, S.H., Lenaers, G., Raymond-Delpech, V., Sattelle, D.B., and Lapied, B. (2007). Exploring the pharmacological properties of insect nicotinic acetylcholine receptors. Trends Pharmacol. Sci. *28*, 14–22. https://doi.org/10.1016/j.tips.2006.11.006.

10. Faltine-Gonzalez, D.Z., and Layden, M.J. (2018). The origin and evolution of acetylcholine
signalling through AChRs in metazoans. Preprint, https://doi.org/10.1101/424804.

11. Littleton, J.T., and Ganetzky, B. (2000). Ion channels and synaptic organization: analysis
of the *Drosophila* genome. Neuron *26*, 35–43. https://doi.org/10.1016/s0896-
6273(00)811 35-6.

12. Thompson, A.J., Lester, H.A., and Lummis, S.C.R. (2010). The structural basis of function
in Cys-loop receptors. Q. Rev. Biophys. *43*, 449–499. https://doi.org/10.1017/S00335835
10000168.

13. Dupuis, J., Louis, T., Gauthier, M., and Raymond, V. (2012). Insights from honeybee (*Apis*
mellifera) and fly (*Drosophila melanogaster*) nicotinic acetylcholine receptors: From
genes to behavioral functions. Neurosci. Biobehav. Rev. *36*, 1553–1564. https://doi.org/
10.1016/j.neubiorev.2012.04.003.

14. Bielopolski, N., Amin, H., Apostolopoulou, A.A., Rozenfeld, E., Lerner, H., Huetteroth, W.,
Lin, A.C., and Parnas, M. (2019). Inhibitory muscarinic acetylcholine receptors enhance
aversive olfactory learning in adult *Drosophila*. eLife *8*, e48264. https://doi.org/10.7554/
eLife.48264.

15. Malloy, C.A., Somasundaram, E., Omar, A., Bhutto, U., Medley, M., Dzubuk, N., and
Cooper, R.L. (2019). Pharmacological identification of cholinergic receptor subtypes:
modulation of locomotion and neural circuit excitability in *Drosophila* larvae.
Neuroscience *411*, 47–64. https://doi.org/10.1016/j.neuroscience.2019.05.016.

16. Xia, R. -Y., Li, M. -Q., Wu, Y. -S., Qi, Y. -X., Ye, G. -Y., and Huang, J. (2016). A new family of
insect muscarinic acetylcholine receptors. Insect Mol. Biol. *25*, 362–369. https://doi.org/
10.1111/imb.12229.

17. Collin, C., Hauser, F., De Valdivia, E.G., Li, S., Reisenberger, J., Carlsen, E.M.M., Khan, Z.,
Hansen, N.Ø., Puhm, F., Søndergaard, L., et al. (2013). Two types of muscarinic
acetylcholine receptors in *Drosophila* and other arthropods. Cell. Mol. Life Sci. *70*, 3231–
3242. https://doi.org/10.1007/s00018-013-1334-0.

18. Ren, G.R., Folke, J., Hauser, F., Li, S., and Grimmelikhuijzen, C.J.P. (2015). The A- and B-
type muscarinic acetylcholine receptors from *Drosophila melanogaster* couple to
different second messenger pathways. Biochem. Biophys. Res. Commun. *462*, 358–364.
https://doi.org/10.1016/j.bbrc.2015.04.141.

19. Perry, T., Chen, W., Ghazali, R., Yang, Y.T., Christesen, D., Martelli, F., Lumb, C., Bao
Luong, H.N., Mitchell, J., Holien, J.K., et al. (2021). Role of nicotinic acetylcholine
receptor subunits in the mode of action of neonicotinoid, sulfoximine and spinosyn
insecticides in *Drosophila melanogaster*. Insect Biochem. Mol. Biol. *131*, 103547.
https://doi.org/10. 1016/j.ibmb.2021.103547.

20. Rozenfeld, E., Tauber, M., Ben-Chaim, Y., and Parnas, M. (2021). GPCR voltage
dependence controls neuronal plasticity and behavior. Nat. Commun. *12*, 7252.
https://doi.org/10.1038/s41467-021-27593-x.

21. Manoim, J.E., Davidson, A.M., Weiss, S., Hige, T., and Parnas, M. (2022). Lateral axonal
modulation is required for stimulus-specific olfactory conditioning in *Drosophila*. Curr.
Biol. CB *32*, 4438-4450.e5. https://doi.org/10.1016/j.cub.2022.09.007.

22. Pribbenow, C., Chen, Y., Heim, M.-M., Laber, D., Reubold, S., Reynolds, E., Balles, I.,
Fernández-d V Alquicira, T., Suárez-Grimalt, R., Scheunemann, L., et al. (2022).
Postsynaptic plasticity of cholinergic synapses underlies the induction and expression of
appetitive and familiarity memories in *Drosophila*. eLife *11*, e80445. https://doi.org/
10.7554/eLife.80445.

23. Qin, B., Humberg, T.-H., Kim, A., Kim, H.S., Short, J., Diao, F., White, B.H., Sprecher, S.G.,
and Yuan, Q. (2019). Muscarinic acetylcholine receptor signaling generates OFF
selectivity in a simple visual circuit. Nat. Commun. *10*, 4093.
https://doi.org/10.1038/s41467-019-12104-w.

24. Schlegel, P., Yin, Y., Bates, A.S., Dorkenwald, S., Eichler, K., Brooks, P., Han, D.S., Gkantia,
M., Dos Santos, M., Munnelly, E.J., et al. (2023). Whole-brain annotation and multi-
connectome cell typing quantifies circuit stereotypy in *Drosophila*. Preprint,
https://doi.org/10.1101/2023.06.27.546055.

25. Borst, A., and Euler, T. (2011). Seeing things in motion: Models, circuits, and
mechanisms. Neuron *71*, 974–994. https://doi.org/10.1016/j.neuron.2011.08.031.

26. Surkova, S., Görne, J., Nuzhdin, S., and Samsonova, M. (2021). Interplay between sex
determination cascade and major signaling pathways during *Drosophila* eye
development: Perspectives for future research. Dev. Biol. *476*, 41–52.
https://doi.org/10. 1016/j.ydbio.2021.03.005.

27. Behnia, R., and Desplan, C. (2015). Visual circuits in flies: beginning to see the whole
picture. Curr. Opin. Neurobiol. *34*, 125–132.
https://doi.org/10.1016/j.conb.2015.03.010.

28. Ribeiro, I.M.A., Drews, M., Bahl, A., Machacek, C., Borst, A., and Dickson, B.J. (2018).
Visual projection neurons mediating directed courtship in *Drosophila*. Cell *174*, 607-
621.e18. https://doi.org/10.1016/j.cell.2018.06.020.

29. Hindmarsh Sten, T., Li, R., Otopalik, A., and Ruta, V. (2021). Sexual arousal gates visual
processing during *Drosophila* courtship. Nature *595*, 549–553.
https://doi.org/10.1038/s4 1586-021-03714-w.

30. Mauss, A.S., and Borst, A. (2020). Optic flow-based course control in insects. Curr. Opin.
Neurobiol. *60*, 21–27. https://doi.org/10.1016/j.conb.2019.10.007.

31. Joesch, M., Schnell, B., Raghu, S.V., Reiff, D.F., and Borst, A. (2010). ON and OFF
pathways in *Drosophila* motion vision. Nature *468*, 300–304.
https://doi.org/10.1038/nature09545.

32. Strother, J.A., Nern, A., and Reiser, M.B. (2014). Direct observation of ON and OFF
pathways in the *Drosophila* visual system. Curr. Biol. *24*, 976–983. https://doi.org/10.10
16/j.cub.2014.03.017.

33. Fisher, Y.E., Silies, M., and Clandinin, T.R. (2015). Orientation selectivity sharpens motion
detection in *Drosophila*. Neuron *88*, 390–402.
https://doi.org/10.1016/j.neuron.2015.09. 033.

34. Maisak, M.S., Haag, J., Ammer, G., Serbe, E., Meier, M., Leonhardt, A., Schilling, T., Bahl,
A., Rubin, G.M., Nern, A., et al. (2013). A directional tuning map of *Drosophila*
elementary motion detectors. Nature *500*, 212–216.
https://doi.org/10.1038/nature12320.

35. Fischbach, K.-F., and Dittrich, A.P.M. (1989). The optic lobe of *Drosophila melanogaster*.
I. A Golgi analysis of wild-type structure. Cell Tissue Res. *258*. https://doi.org/10.1007/
BF00218858.

36. Pinto-Teixeira, F., Koo, C., Rossi, A.M., Neriec, N., Bertet, C., Li, X., Del-Valle-Rodriguez,
A., and Desplan, C. (2018). Development of concurrent retinotopic maps in the fly
motion detection circuit. Cell *173*, 485-498.e11.
https://doi.org/10.1016/j.cell.2018.02.053.

37. Apitz, H., and Salecker, I. (2018). Spatio-temporal relays control layer identity of
direction-selective neuron subtypes in *Drosophila*. Nat. Commun. *9*, 2295.
https://doi.org/10.1038/ s41467-018-04592-z.

38. Takemura, S.-Y., Nern, A., Chklovskii, D.B., Scheffer, L.K., Rubin, G.M., and
Meinertzhagen, I.A. (2017). The comprehensive connectome of a neural substrate for
"ON" motion detection in *Drosophila*. eLife *6*, e24394.
https://doi.org/10.7554/eLife.24394.

39. Shinomiya, K., Takemura, S., Rivlin, P.K., Plaza, S.M., Scheffer, L.K., and Meinertzhagen,
I.A. (2015). A common evolutionary origin for the ON- and OFF-edge motion detection
pathways of the *Drosophila* visual system. Front. Neural Circuits *9*. https://doi.org/10.
3389/fncir.2015.00033.

40. Shinomiya, K., Huang, G., Lu, Z., Parag, T., Xu, C.S., Aniceto, R., Ansari, N., Cheatham, N.,
Lauchie, S., Neace, E., et al. (2019). Comparisons between the ON- and OFF-edge motion
pathways in the *Drosophila* brain. eLife *8*, e40025. https://doi.org/10.7554/eLife.40025.

41. Takemura, S., Bharioke, A., Lu, Z., Nern, A., Vitaladevuni, S., Rivlin, P.K., Katz, W.T.,
Olbris, D.J., Plaza, S.M., Winston, P., et al. (2013). A visual motion detection circuit
suggested by *Drosophila* connectomics. Nature *500*, 175–181.
https://doi.org/10.1038/nature12450.

42. Meier, M., and Borst, A. (2019). Extreme compartmentalization in a *Drosophila* amacrine
cell. Curr. Biol. *29*, 1545-1550.e2. https://doi.org/10.1016/j.cub.2019.03.070.

43. Davis, F.P., Nern, A., Picard, S., Reiser, M.B., Rubin, G.M., Eddy, S.R., and Henry, G.L.
(2020). A genetic, genomic, and computational resource for exploring neural circuit
function. eLife *9*, e50901. https://doi.org/10.7554/eLife.50901.

44. Behnia, R., Clark, D.A., Carter, A.G., Clandinin, T.R., and Desplan, C. (2014). Processing
properties of ON and OFF pathways for *Drosophila* motion detection. Nature *512*, 427–
430. https://doi.org/10.1038/nature13427.

45. Groschner, L.N., Malis, J.G., Zuidinga, B., and Borst, A. (2022). A biophysical account of
multiplication   by   a   single   neuron.   Nature   *603*,   119–123.
https://doi.org/10.1038/s41586-022-04428-3.

46. Ammer, G., Leonhardt, A., Bahl, A., Dickson, B.J., and Borst, A. (2015). Functional
specialization of neural input elements to the *Drosophila* ON motion detector. Curr. Biol.
25, 2247–2253. https://doi.org/10.1016/j.cub.2015.07.014.

47. Takemura, S., Karuppudurai, T., Ting, C.-Y., Lu, Z., Lee, C.-H., and Meinertzhagen, I.A.
(2011). Cholinergic circuits integrate neighboring visual signals in a *Drosophila* motion
detection pathway. Curr. Biol. CB *21*, 2077–2084. https://doi.org/10.1016/j.cub.2011.
10.053.

48. Serbe, E., Meier, M., Leonhardt, A., and Borst, A. (2016). Comprehensive
characterization of the major presynaptic elements to the *Drosophila* OFF motion
detector. Neuron *89*, 829–841. https://doi.org/10.1016/j.neuron.2016.01.006.

49. Meier, M., Serbe, E., Maisak, M.S., Haag, J., Dickson, B.J., and Borst, A. (2014). Neural
circuit components of the *Drosophila* OFF motion vision pathway. Curr. Biol. *24*, 385–
392. https://doi.org/10.1016/j.cub.2014.01.006.

50. Shinomiya, K., Karuppudurai, T., Lin, T.-Y., Lu, Z., Lee, C.-H., and Meinertzhagen, I.A.
(2014). Candidate neural substrates for OFF-edge motion detection in *Drosophila*. Curr.
Biol. *24*, 1062–1070. https://doi.org/10.1016/j.cub.2014.03.051.

51. Braun, A., Borst, A., and Meier, M. (2023). Disynaptic inhibition shapes tuning of OFF-
motion   detectors   in   *Drosophila*.   Curr.   Biol.   CB   *33*,   2260-2269.e4.
https://doi.org/10.1016/ j.cub.2023.05.007.

52. Ramos-Traslosheros, G., and Silies, M. (2021). The physiological basis for contrast
opponency in motion computation in *Drosophila*. Nat. Commun. *12*, 4987.
https://doi.org/10.1038/s41467-021-24986-w.

53. Borst, A. (2018). A biophysical mechanism for preferred direction enhancement in fly
motion vision. PLOS Comput. Biol. *14*, e1006240. https://doi.org/10.1371/journal.
pcbi.1006240.

54. Haag, J., Arenz, A., Serbe, E., Gabbiani, F., and Borst, A. (2016). Complementary
mechanisms create direction selectivity in the fly. eLife *5*, e17421. https://doi.org/10.
7554/eLife.17421.

55. Haag, J., Mishra, A., and Borst, A. (2017). A common directional tuning mechanism of
1190   *Drosophila* motion-sensing neurons in the ON and in the OFF pathway. eLife *6*, e29044.
1191   https://doi.org/10.7554/eLife.29044.

1192   56. Gruntman, E., Romani, S., and Reiser, M.B. (2018). Simple integration of fast excitation
1193   and offset, delayed inhibition computes directional selectivity in *Drosophila*. Nat.
1194   Neurosci. *21*, 250–257. https://doi.org/10.1038/s41593-017-0046-4.

1195   57. Arenz, A., Drews, M.S., Richter, F.G., Ammer, G., and Borst, A. (2017). The temporal
1196   tuning of the *Drosophila* motion detectors is determined by the dynamics of their input
1197   elements. Curr. Biol. *27*, 929–944. https://doi.org/10.1016/j.cub.2017.01.051.

1198   58. Hoermann, N., Schilling, T., Haji Ali, A., Serbe, E., Mayer, C., Borst, A., and Pujol-Martí, J.
1199   (2020). A combinatorial code of transcription factors specifies subtypes of visual motion-
1200   sensing neurons in *Drosophila*. Development, dev.186296. https://doi.org/10.1242/
1201   dev.186296.

1202   59. Özel, M.N., Simon, F., Jafari, S., Holguera, I., Chen, Y.-C., Benhra, N., El-Danaf, R.N.,
1203   Kapuralin, K., Malin, J.A., Konstantinides, N., et al. (2021). Neuronal diversity and
1204   convergence in a visual system developmental atlas. Nature *589*, 88–95.
1205   https://doi.org/10.1038/s41586-020-2879-3.

1206   60. Sanfilippo, P., Kim, A.J., Bhukel, A., Yoo, J., Mirshahidi, P.S., Pandey, V., Bevir, H., Yuen,
1207   A., Mirshahidi, P.S., Guo, P., et al. (2024). Mapping of multiple neurotransmitter
1208   receptor subtypes and distinct protein complexes to the connectome. Neuron *112*, 942-
1209   958.e13. https://doi.org/10.1016/j.neuron.2023.12.014.

1210   61. Kondo, S., Takahashi, T., Yamagata, N., Imanishi, Y., Katow, H., Hiramatsu, S., Lynn, K.,
1211   Abe, A., Kumaraswamy, A., and Tanimoto, H. (2020). Neurochemical organization of the
1212   *Drosophila* brain visualized by endogenously tagged neurotransmitter receptors. Cell
1213   Rep. *30*, 284-297.e5. https://doi.org/10.1016/j.celrep.2019.12.018.

1214   62. Zheng, Z., Lauritzen, J.S., Perlman, E., Robinson, C.G., Nichols, M., Milkie, D., Torrens, O.,
1215   Price, J., Fisher, C.B., Sharifi, N., et al. (2018). A complete electron microscopy volume of
1216   the brain of adult *Drosophila melanogaster*. Cell *174*, 730-743.e22. https://doi.org/10.1
1217   016/j.cell.2018.06.019.

1218   63. Dorkenwald, S., McKellar, C.E., Macrina, T., Kemnitz, N., Lee, K., Lu, R., Wu, J., Popovych,
1219   S., Mitchell, E., Nehoran, B., et al. (2022). FlyWire: online community for whole-brain
1220   connectomics. Nat. Methods *19*, 119–128. https://doi.org/10.1038/s41592-021-01330-
1221   0.

1222   64. Dorkenwald, S., Matsliah, A., Sterling, A.R., Schlegel, P., Yu, S., McKellar, C.E., Lin, A.,
1223   Costa, M., Eichler, K., Yin, Y., et al. (2023). Neuronal wiring diagram of an adult brain.
1224   Preprint, https://doi.org/10.1101/2023.06.27.546656.

1225   65. Jenett, A., Rubin, G.M., Ngo, T.-T.B., Shepherd, D., Murphy, C., Dionne, H., Pfeiffer, B.D.,
1226   Cavallaro, A., Hall, D., Jeter, J., et al. (2012). A GAL4-driver line resource for *Drosophila*
1227   neurobiology. Cell Rep. *2*, 991–1001. https://doi.org/10.1016/j.celrep.2012.09.011.

66. Tirian, L., and Dickson, B.J. (2017). The VT GAL4, LexA, and split-GAL4 driver line
collections for targeted expression in the *Drosophila* nervous system. Preprint,
https://doi.org/10.1101/198648 https://doi.org/10.1101/198648.

67. Nern, A., Pfeiffer, B.D., and Rubin, G.M. (2015). Optimized tools for multicolor stochastic
labeling reveal diverse stereotyped cell arrangements in the fly visual system. Proc. Natl.
Acad. Sci. *112*, E2967–E2976. https://doi.org/10.1073/pnas.1506763112.

68. Meissner, G.W., Nern, A., Dorman, Z., DePasquale, G.M., Forster, K., Gibney, T.,
Hausenfluck, J.H., He, Y., Iyer, N.A., Jeter, J., et al. (2023). A searchable image resource
of *Drosophila* GAL4 driver expression patterns with single neuron resolution. eLife *12*,
e80660. https://doi.org/10.7554/eLife.80660.

69. Diegelmann, S., Bate, M., and Landgraf, M. (2008). Gateway cloning vectors for the
LexA-based binary expression system in *Drosophila*. Fly (Austin) *2*, 236–239.
https://doi.org/ 10.4161/fly.6817.

70. Feinberg, E.H., VanHoven, M.K., Bendesky, A., Wang, G., Fetter, R.D., Shen, K., and
Bargmann, C.I. (2008). GFP reconstitution across synaptic partners (GRASP) defines cell
contacts and synapses in living nervous systems. Neuron *57*, 353–363. https://doi.org/
10.1016/j.neuron.2007.11.030.

71. Shearin, H.K., Quinn, C.D., Mackin, R.D., Macdonald, I.S., and Stowers, R.S. (2018). t-
GRASP, a targeted GRASP for assessing neuronal connectivity. J. Neurosci. Methods *306*,
94–102. https://doi.org/10.1016/j.jneumeth.2018.05.014.

72. Macpherson, L.J., Zaharieva, E.E., Kearney, P.J., Alpert, M.H., Lin, T.-Y., Turan, Z., Lee, C.-
H., and Gallio, M. (2015). Dynamic labelling of neural connections in multiple colours by
trans-synaptic fluorescence complementation. Nat. Commun. *6*, 10024. https://doi.org/
10.1038/ncomms10024.

73. Wagh, D.A., Rasse, T.M., Asan, E., Hofbauer, A., Schwenkert, I., Dürrbeck, H., Buchner, S.,
Dabauvalle, M.-C., Schmidt, M., Qin, G., et al. (2006). Bruchpilot, a protein with
homology to ELKS/CAST, is required for structural integrity and function of synaptic
active zones in *Drosophila*. Neuron *49*, 833–844.
https://doi.org/10.1016/j.neuron.2006.02.008.

74. Mosca, T.J., and Luo, L. (2014). Synaptic organization of the *Drosophila* antennal lobe
and its regulation by the Teneurins. eLife *3*, e03726.
https://doi.org/10.7554/eLife.03726.

75. Schmid, A., Hallermann, S., Kittel, R.J., Khorramshahi, O., Frölich, A.M.J., Quentin, C.,
Rasse, T.M., Mertel, S., Heckmann, M., and Sigrist, S.J. (2008). Activity-dependent site-
specific changes of glutamate receptor composition *in vivo*. Nat. Neurosci. *11*, 659–666.
https://doi.org/10.1038/nn.2122.

76. Christiansen, F., Zube, C., Andlauer, T.F.M., Wichmann, C., Fouquet, W., Owald, D.,
Mertel, S., Leiss, F., Tavosanis, G., Farca Luna, A.J., et al. (2011). Presynapses in Kenyon

cell dendrites in the mushroom body calyx of *Drosophila*. J. Neurosci. *31*, 9696–9707.
https://doi.org/10.1523/JNEUROSCI.6542-10.2011.

77. Berger-Müller, S., Sugie, A., Takahashi, F., Tavosanis, G., Hakeda-Suzuki, S., and Suzuki, T.
(2013). Assessing the role of cell-surface molecules in central synaptogenesis in the
Drosophila visual system. PLoS ONE *8*, e83732. https://doi.org/10.1371/journal.pone.
0083732.

78. Owald, D., Fouquet, W., Schmidt, M., Wichmann, C., Mertel, S., Depner, H., Christiansen,
F., Zube, C., Quentin, C., Körner, J., et al. (2010). A Syd-1 homologue regulates pre- and
postsynaptic maturation in *Drosophila*. J. Cell Biol. *188*, 565–579. https://doi.org/
10.1083/jcb.200908055.

79. Fendl, S., Vieira, R.M., and Borst, A. (2020). Conditional protein tagging methods reveal
highly specific subcellular distribution of ion channels in motion-sensing neurons. eLife
9, e62953. https://doi.org/10.7554/eLife.62953.

80. Buhmann, J., Sheridan, A., Malin-Mayor, C., Schlegel, P., Gerhard, S., Kazimiers, T.,
Krause, R., Nguyen, T.M., Heinrich, L., Lee, W.-C.A., et al. (2021). Automatic detection of
synaptic partners in a whole-brain *Drosophila* electron microscopy data set. Nat.
Methods *18*, 771–774. https://doi.org/10.1038/s41592-021-01183-7.

81. Heinrich, L., Funke, J., Pape, C., Nunez-Iglesias, J., and Saalfeld, S. (2018). Synaptic cleft
segmentation in non-isotropic volume electron microscopy of the complete *Drosophila*
brain. Preprint at arXiv, https://doi.org/10.48550/ARXIV.1805.02718 https://doi.org/
10.48550/ARXIV.1805.02718.

82. Götz, K.G. (1968). Flight control in *Drosophila* by visual perception of motion. Kybernetik
4, 199–208. https://doi.org/10.1007/BF00272517.

83. Ryu, L., Kim, S.Y., and Kim, A.J. (2022). From photons to behaviors: Neural
implementations of visual behaviors in *Drosophila*. Front. Neurosci. *16*, 883640.
https://doi.org/10.3389/fnins.2022.883640.

84. Leonte, M.-B., Leonhardt, A., Borst, A., and Mauss, A.S. (2021). Aerial course
stabilization is impaired in motion-blind flies. J. Exp. Biol. *224*, jeb242219.
https://doi.org/10.1242/jeb.242219.

85. Bahl, A., Ammer, G., Schilling, T., and Borst, A. (2013). Object tracking in motion-blind
flies. Nat. Neurosci. *16*, 730–738. https://doi.org/10.1038/nn.3386.

86. Schnell, B., Raghu, S.V., Nern, A., and Borst, A. (2012). Columnar cells necessary for
motion responses of wide-field visual interneurons in *Drosophila*. J. Comp. Physiol. A
198, 389–395. https://doi.org/10.1007/s00359-012-0716-3.

87. Schilling, T., and Borst, A. (2015). Local motion detectors are required for the
computation of expansion flow-fields. Biol. Open *4*, 1105–1108.
https://doi.org/10.1242/bio.012690.

88. Schilling, T., Ali, A.H., Leonhardt, A., Borst, A., and Pujol-Martí, J. (2019). Transcriptional
control of morphological properties of direction-selective T4/T5 neurons in *Drosophila*.
Development, dev.169763. https://doi.org/10.1242/dev.169763.

89. Ni, J.-Q., Liu, L.-P., Binari, R., Hardy, R., Shim, H.-S., Cavallaro, A., Booker, M., Pfeiffer,
B.D., Markstein, M., Wang, H., et al. (2009). A *Drosophila* resource of transgenic RNAi
lines   for   neurogenetics.   Genetics   *182*,   1089–1100.
https://doi.org/10.1534/genetics.109.103630.

90. Ni, J.-Q., Zhou, R., Czech, B., Liu, L.-P., Holderbaum, L., Yang-Zhou, D., Shim, H.-S., Tao,
R., Handler, D., Karpowicz, P., et al. (2011). A genome-scale shRNA resource for
transgenic   RNAi   in   *Drosophila*.   Nat.   Methods   *8*,   405–407.
https://doi.org/10.1038/nmeth.1592.

91. Shinomiya, K., Nern, A., Meinertzhagen, I.A., Plaza, S.M., and Reiser, M.B. (2022).
Neuronal circuits integrating visual motion information in *Drosophila melanogaster*.
Curr. Biol. CB *32*, 3529-3544.e2. https://doi.org/10.1016/j.cub.2022.06.061.

92. Albuquerque, E.X., Pereira, E.F.R., Alkondon, M., and Rogers, S.W. (2009). Mammalian
nicotinic acetylcholine receptors: from structure to function. Physiol. Rev. *89*, 73–120.
https://doi.org/10.1152/physrev.00015.2008.

93. Abrams, P., Andersson, K.-E., Buccafusco, J.J., Chapple, C., de Groat, W.C., Fryer, A.D.,
Kay, G., Laties, A., Nathanson, N.M., Pasricha, P.J., et al. (2006). Muscarinic receptors:
their distribution and function in body systems, and the implications for treating
overactive   bladder.   Br.   J.   Pharmacol.   *148*,   565–578.
https://doi.org/10.1038/sj.bjp.0706780.

94. Schuster, R., Phannavong, B., Schröder, C., and Gundelfinger, E.D. (1993).
Immunohistochemical localization of a ligand-binding and a structural subunit of
nicotinic acetylcholine receptors in the central nervous system of *Drosophila*
melanogaster. J. Comp. Neurol. *335*, 149–162. https://doi.org/10.1002/cne.903350202.

95. Gundelfinger, E.D. (1992). How complex is the nicotinic receptor system of insects?
Trends Neurosci. *15*, 206–211. https://doi.org/10.1016/0166-2236(92)90035-7.

96. Jonas, P.E., Phannavong, B., Schuster, R., Schröder, C., and Gundelfinger, E.D. (1994).
Expression of the ligand-binding nicotinic acetylcholine receptor subunit Dα2 in the
Drosophila   central   nervous   system.   J.   Neurobiol.   *25*,   1494–1508.
https://doi.org/10.1002/ neu.480251203.

97. Thany, S.H. (2010). Electrophysiological studies and pharmacological properties of insect
native nicotinic acetylcholine receptors. In insect nicotinic acetylcholine receptors
advances in experimental medicine and biology., S. H. Thany, ed. (Springer New York),
pp. 53–63. https://doi.org/10.1007/978-1-4419-6445-8_5.

98. Chamaon, K., Smalla, K., Thomas, U., and Gundelfinger, E.D. (2002). Nicotinic
acetylcholine receptors of *Drosophila*: three subunits encoded by genomically linked

genes can co-assemble into the same receptor complex. J. Neurochem. *80*, 149–157.
https://doi.org/10.1046/j.0022-3042.2001.00685.x.

99. Chamaon, K., Schulz, R., Smalla, K.-H., Seidel, B., and Gundelfinger, E.D. (2000). Neuronal
nicotinic acetylcholine receptors of *Drosophila melanogaster*: the α-subunit Dα3 and the
β-type subunit ARD co-assemble within the same receptor complex. FEBS Lett. *482*,
189–192. https://doi.org/10.1016/S0014-5793(00)02057-3.

100.Deng, B., Li, Q., Liu, X., Cao, Y., Li, B., Qian, Y., Xu, R., Mao, R., Zhou, E., Zhang, W., et al.
(2019). Chemoconnectomics: Mapping chemical transmission in *Drosophila*. Neuron
101, 876-893.e4. https://doi.org/10.1016/j.neuron.2019.01.045.

101.Bahl, A., Serbe, E., Meier, M., Ammer, G., and Borst, A. (2015). Neural mechanisms for
Drosophila contrast vision. Neuron *88*, 1240–1252. https://doi.org/10.1016/j.neuron.
2015.11.004.

102.Namiki, S., Dickinson, M.H., Wong, A.M., Korff, W., and Card, G.M. (2018). The
functional organization of descending sensory-motor pathways in *Drosophila*. eLife *7*,
e34272. https://doi.org/10.7554/eLife.34272.

103.Komori, Y., Takayama, K., Okamoto, N., Kamiya, M., Koizumi, W., Ihara, M., Misawa, D.,
Kamiya, K., Yoshinari, Y., Seike, K., et al. (2023). Functional impact of subunit
composition and compensation on *Drosophila melanogaster* nicotinic receptors-targets
of neonicotinoids. PLoS Genet. *19*, e1010522. https://doi.org/10.1371/journal.pgen.101
0522.

104.Pfeiffer, B.D., Ngo, T.-T.B., Hibbard, K.L., Murphy, C., Jenett, A., Truman, J.W., and
Rubin, G.M. (2010). Refinement of tools for targeted gene expression in *Drosophila*.
Genetics *186*, 735–755. https://doi.org/10.1534/genetics.110.119917.

105.Raccuglia, D., Huang, S., Ender, A., Heim, M.-M., Laber, D., Suárez-Grimalt, R., Liotta, A.,
Sigrist, S.J., Geiger, J.R.P., and Owald, D. (2019). Network-specific synchronization of
electrical slow-wave oscillations regulates sleep drive in *Drosophila*. Curr. Biol. CB *29*,
3611-3621.e3. https://doi.org/10.1016/j.cub.2019.08.070.

106.Jin, X., Tian, Y., Zhang, Z.C., Gu, P., Liu, C., and Han, J. (2021). A subset of DN1p neurons
integrates thermosensory inputs to promote wakefulness via CNMa signaling. Curr. Biol.
CB *31*, 2075-2087.e6. https://doi.org/10.1016/j.cub.2021.02.048.

107.Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T.,
Preibisch, S., Rueden, C., Saalfeld, S., Schmid, B., et al. (2012). Fiji: an open-source
platform for biological-image analysis. Nat. Methods *9*, 676–682. https://doi.org/
10.1038/nmeth.2019.

108.Buchner, E. (1976). Elementary movement detectors in an insect visual system. Biol.
Cybern. *24*, 85–101. https://doi.org/10.1007/BF00360648.

109.Seelig, J.D., Chiappe, M.E., Lott, G.K., Dutta, A., Osborne, J.E., Reiser, M.B., and
Jayaraman, V. (2010). Two-photon calcium imaging from head-fixed *Drosophila* during

optomotor walking behavior. Nat. Methods *7*, 535–540. https://doi.org/10.1038/nmeth.
1468.

Figure 1. AChR expression on T5 dendrites

(A) Schematic representation of T5 dendrites residing in lobula layer 1 (LO1) of the *Drosophila* optic lobe (OL).

(B) Schematic representation of nicotinic ionotropic (magenta) and muscarinic metabotropic AChRs (cyan) in *Drosophila*. Out of the seven nAChRα subunits, six were tagged with EGFP[22]. The mAChR-B was tagged with the HA peptide tag (C.H.Lee)

(C) OL (grey) with T4 and T5 dendrites (R42F06-Gal4>UAS-myr::tdTomato, blue) in M10 and LO1 accordingly. Scale bar 40μm.

(D-I) nAChRα1, nAChRα2, nAChRα4, nAChRα5, nAChRα6 and nAChRα7 (nAChRα::EGFP, magenta) subunit expression on T5 dendrites (R42F06-Gal4>UAS-myr::tdTomato, blue). Scale bar 10μm, inset 5μm.

(J) Density of nAChRα subunit expression on T5 dendrites across 30 ROIs in LO1 (optic lobes n=6). nAChRα3, nAChRβ1, nAChRβ2 and nAChRβ3 subunits were not addressed. Subunits with the lowest expression levels are in grey. Data is mean ± SEM.

(K) Overexpression of mAChR-B (UAS-mAChR-B::HA, cyan) on T5 dendrites (R42F06-Gal4>UAS-myr::tdTomato, blue). Scale bar 10μm, inset 5μm.

Figure 2. Morphological identification of the four columnar T5 cholinergic input neurons

(A) Electron microscopy (EM)(flywire.ai) reconstruction of a Tm1 neuron reveals two loci of arborization in medulla. Schematic representation of the Tm1 axonal terminal localization in LO1. Tm1-Gal4 expression (cyan) of the MultiColor FlpOut cassette (V5, HA, FLAG) and axonal localization in LO. Scale bar 40µm and 10µm.

(B) Whole population labelling of Tm1 neurons (R74G01-Gal4>UAS-CD8::GFP). Scale bar 40µm.

(C) Whole population labelling of Tm1-LexA (purple) neurons (VT041034-LexAGAD>LexAop-myr::mCherry). Arrows represent additional dendritic ramifications that lack in (B). Scale bar 40µm.

(D-F) As in (A-C), but for Tm2 neurons (VT012282-Gal4, VT058650-LexAGAD).

(G-I) As in (A-C), but for Tm4 neurons (R35H01-Gal4, R53C02-LexAGAD).

(J-L) As in (A-C), but for Tm9 neurons (VT065303-Gal4, R24C08-LexAp65).

Figure 3. Differential synaptic localization of nAChRα subunits in Tm synapses
(A-D) Trans-synaptic alignment of Tm1, Tm2, Tm4 and Tm9 boutons (Tm-Gal4> UAS-Brp[short::mStraw], grey) with the nAChRα1, nAChRα4, nAChRα5 and nAChRα7 subunits (nAChRα::EGFP, magenta) in LO1. Arrows indicate the Brp[short::mStraw]-receptor punctum trans-synaptic alignment. Scale bar 1µm. (E-H) Synapse occupancy of Tm1, Tm2, Tm4 and Tm9 boutons with nAChRα1 (pale magenta), nAChRα4 (light magenta), nAChRα5 (magenta) and nAChRα7 (dark magenta) subunits in LO1 (optic lobes n=4, ROIs N=20, for Tm4-nAChRα5 n=3, N=15). Data is mean ± SEM.

Figure 4. Wiring of Tm1, Tm2, Tm4 and Tm9 on T5 dendrites

(A) Counts of Tm1, Tm2, Tm4 and Tm9 synapses in T5a dendrites (T5a cells n=5, flywire.ai) (left). Counts of predicted (n=663) to proofread (n=479) Tm synapses in T5a dendrites (right). Data is mean ± SEM.

(B) As in (A) for T5c dendrites (T5c cells n=5, flywire.ai) (left). Counts of predicted (n=621) to proofread (n=488) Tm synapses in T5c dendrites (right). Data is mean ± SEM.

(C) Dendritic localization of Tm synapses across the T5a anterior-posterior axis (T5a cells n=5). Root is considered the first dendritic branch point.

(D) Dendritic localization of Tm synapses across the T5c dorso-ventral axis (T5c cells n=5).

(E) Schematic representation of Tm-T5a dendritic synapses and the observed postsynaptic receptor heterogeneity in terms of the most abundant nAChR subunit as in Figure 3E-H.

Figure 5. Fly optomotor response post-AChR knock-down

(A) Schematic workflow representation for the behavioral assessment of AChR knock-down flies.

(B) Trajectory (angular velocity) of nAChRα1 knock-down flies (R42F06-p65.AD; VT043070-Gal4.DBD>UAS-nAChRα1-RNAi, pale magenta, flies n=20) while presented to a sequence of upward to rightward to downward to leftward OFF edge motion compared to the parental control flies (UAS-nAChRα1-RNAi, grey n=21). The dark lines represent the mean ± SEM.

(C-F) Same as in (B) for the nAChRα4 (light magenta n=19, grey n=20), nAChRα5 (magenta n=18, grey n=21), nAChRα7 (dark magenta, n=21, grey n=20), mAChR-B (cyan n=20, grey n=20) knock-down flies.

(G) Angular velocity of control and AChR knock-down flies. The normality of distribution was assessed with the use of Shapiro-Wilk test. *p<0.05, **p<0.01, ***p<0.001. Data is mean ± SEM.

Figure 6. Directional tuning of T5 cells post-AChR knock-down

(A) Schematic workflow representation for the two-photon calcium imaging in AChR knock-down T5 cells.

(B) Polar plot of the maximum responses (ΔF/F) across sixteen OFF edge directions (º) of T5c cells for mCherry controls (black, flies n=7, ROIs N=85, shown as grey in C, D and E) and nAChRα4 RNAi (light magenta, n=7, N=61) conditions (R39H12-Gal4>AChR-RNAi, VT50384-LexA>LexAop-IVS-GCaMP6m). The dark lines represent the mean (68% CI).

(C) Same as in (B) for nAChRα5 RNAi flies (magenta, n=7, N=82).

(D) Same as in (B) for nAChRα7 RNAi flies (dark magenta, n=7, N=60).

(E) Same as in (B) for mAChR-B RNAi flies (cyan, n=7, N=54).

(F) Directional tuning index (Ldir). Flies n=7. The normality of distribution was assessed with the use of Shapiro-Wilk test. *p<0.05, ***p<0.001. Data is mean ± SEM.

Figure S1. AChR expression in T5 cells-Related to Figure 1

(A) nAChRα2 expression in LO. Arrows indicate the LO layers of expression. Scale bar 10μm, inset 5μm.

(B) Example ROIs in LO1 for quantification as in Figure 1J. Scale bar 10μm, inset 5μm.

(C) Bulk RNAseq of AChR expression (transcripts per million, TPM) in T5 cells and probability of expression (p)[43].

(D) scRNAseq of AChR expression in T4 (grey) and T5 (blue) cells across five developmental stages (hours correspond to APF-after puparium formation)[58].

Figure S2. AChR expression in T4-T5 axonal terminals-Related to Figure 1

(A) Schematic representation of T5 axonal terminals residing in lobula plate (LOP1-4) of the *Drosophila* optic lobe (OL).

(B-G) Expression of nAChRα1, nAChRα2, nAChRα4, nAChRα5, nAChRα6 and nAChRα7 (nAChRα::EGFP, magenta) in T4-T5 axonal terminals (R42F06-Gal4>UAS-myr::tdTomato, blue). Scale bar 10μm.

(H) Overexpression of mAChR-B (cyan) in T4-T5 axonal terminals (R42F06-Gal4>UAS-myr::tdTomato, UAS-mAChR-B::HA). Scale bar 10μm.

(I) Pan-neuronal expression of mAChR-B in OL (GMR57C10-Gal4>UAS-mAChR-B::HA). Scale bar 40μm.

(J) Normalized mean grey value of nAChR subunits in LOP (layers 1 and 2) before and after RNAi induction (optic lobes n=2, ROIs N=20, for nAChRα1 n=1). The normality of distribution was assessed with the use of Shapiro-Wilk test. *p<0.05, ****p<0.0001. Data is mean ± SEM.

(K) Summarized expression patterns of nAChRα1, nAChRα2, nAChRα4, nAChRα5, nAChRα6 and nAChRα7 subunits and the mAChR-B type on T5 dendrites and in T4-T5 axonal terminals.

Figure S3. RNA interference against AChRs in T5 cells-Related to Figure 1 and 5
(A) Expression of the nAChRα1 subunit (nAChRα1::EGFP) in LO1 before (left panel) and after (right panel) the RNAi induction (R42F06-Gal4>AChR-RNAi). Scale bar 10μm.
(B-E) As in (A) for the nAChRα4, nAChRα5, nAChRα7 subunits and the mAChR-B type.
(F) Normalized mean grey value of AChRs in LO1 before and after RNAi induction (optic lobes n=2, ROIs N=20). The normality of distribution was assessed with the use of Shapiro-Wilk test. **p<0.01, ***p<0.001, ****p<0.0001. Data is mean ± SEM.
(G) Developmental onset (P18) of the R42F06 promoter used for the split T4-T5 line in all behavioral experiments (R42F06-Gal4> UAS-myr::tdTomato). Scale bar 40μm, inset 10μm.
(H) Split T4-T5 line (R42F06-p65.AD; VT043070-Gal4.DBD>UAS-CD4::tdGFP) used in all behavioral experiments. Arrows indicate the four LOP layers. Scale bar 40μm, inset 10μm.

	scRNA-seq	Bulk-seq	Enhancer trap	Tagged
nAChRα1	+	+	+	+
nAChRα2	-	+	-	-
nAChRα3	+	+	+	+
nAChRα4	+	+		+
nAChRα5	+	+	+	+
nAChRα6	-	-	-	+
nAChRα7	-	+	+	+
nAChRβ1	+	+	-	+
nAChRβ2	-	+	-	
nAChRβ3	-	-	-	
mAChR-A	+	+	-	
mAChR-B	+	+	+	

Figure S4. Enhancer-trap verification of AChR expression in T4-T5 cells-Related to Figure 1

(A-L) Enhancer-trap lines for the nAChRα1, nAChRα2, nAChRα3, nAChRα4, nAChRα5, nAChRα6, nAChRα7, nAChRβ1, nAChRβ2, nAChRβ3 subunits and mAChR-A and mAChR-B receptors (AChR-Mi or T2A-Gal4>UAS-CD8::GFP, magenta and cyan). Signal overlap with T4-T5 somata (R42F06-LexA>LexAop-CD8::RFP, blue) denoted the T4-T5 AChR expression. The nAChRα4 line did not show expression across all brain areas. Images are z-projections. Scale bar 40μm, inset 10μm.

(M) AChR expression in T5 cells across scRNA-seq[58], Bulk-seq[43], enhancer trap and endogenously tagged receptor[60] experiments.

Figure S5. Screening of genetic approaches for synapse visualization-Related to Figure 3

(A) t-GRASP expression in Tm1, Tm2, Tm4 and Tm9-T5 synapses (Tm1, Tm2, Tm4, Tm9-Gal4>UAS-pre-t-GRASP, VT25965-LexAp65>LexAop-post-t-GRASP). Scale bar 10μm. Ectopic reconstitution is observed in Tm4-T5 synapses.

(B) Activity GRASP (syb-GRASP) in Tm1, Tm2, Tm4 and Tm9-T5 synapses (Tm1, Tm2, Tm4, Tm9-Gal4>UAS-nSyb-spGFP1-10, VT25965-LexAp65>LexAop-CD4-spGFP11). Scale bar 10μm.

(C) Brp^{short}::GFP expression in Tm1, Tm2, Tm4 and Tm9 boutons (Tm1, Tm2, Tm4, Tm9-Gal4>UAS-Brp^{short}::GFP). Scale bar 10μm.

Figure S6. Validation of Brp^{short} as a precise presynaptic marker-Related to Figure 3

(A) Trans-synaptic alignment between Brp$^{short::mCherry}$(blue) and the GABAergic Rdl receptor (grey) in the cholinergic Tm3-T4 synapse (GMR13E12-LexA>LexAop-Brp^{short}::mCherry, R42F06-Gal4>UAS-Rdl::GFP). Scale bar 10µm.

(B) Trans-synaptic alignment between Brp$^{short::mCherry}$(blue) and the glutamatergic GluClα receptor (grey) in the glutamatergic Mi9-T4 synapse (GMR48A07-LexA>LexAop-Brp^{short}::mCherry, R42F06-Gal4>UAS-GluClα::GFP). Scale bar 10µm.

(C) Localization of Brp$^{short::mStraw}$ in Tm1, Tm2, Tm4 and Tm9 boutons (blue) and endogenous Brp (grey) identified by the anti-Brp staining. Arrows correspond to the signal overlap between the Tm-Brp^{short} and the Anti-Brp. Scale bar 5 and 1µm.

Figure S7. Localization of nAChRα subunits in Tm synapses-Related to Figure 3
(A) LO1 loci that were used for Figure 3A-D. Scale bar 10μm.
(B) Line profile analysis for the assessment of trans-synaptic alignment of Brp$^{short::mStraw}$ puncta (grey) and nAChRα puncta (magenta) across Tm1, Tm2, Tm4 and Tm9 synapses (analyzed puncta n=10). The dark lines represent the mean values.

Figure S8. Neuronal outputs of Tm1, Tm2, Tm4 and Tm9 cells in LO1-Related to Figure 3

(A) Schematic representation of Tm1, Tm2, Tm4 and Tm9 axonal terminals in relation to LO1. Primary Tm1, Tm2, Tm4 and Tm9 output neuronal types in LO1 (Tm cells n=5, Codex). Data is mean ± SEM.

(B) C-RASP for the Tm9-T5 synapse visualization (Tm9-Gal4>UAS-IVS-syb-spCFP1-10, VT25965-LexAp65>LexAop-CD4-spGFP11, grey) of the nAChRα7 subunit (nAChRα7::EGFP, magenta). Arrows correspond to Tm9-nAChRα7 trans-synaptic alignment. Scale bar 10µm.

Figure S9. mAChR-B receptor localization in T5 dendrites-Related to Figure 3
(A) mAChR-B localization (R42F06-Gal4>UAS-mAChR-B::HA, cyan) in Tm9-T5 synapses (Tm9-LexA>LexAop-Brp^short::mCherry, grey). Scale bar 10μm.
(B) nAChRα5 subunit (nAChRα5::EGFP, magenta) localization pattern compared to mAChR-B (R42F06-Gal4>UAS-mAChR-B::HA, cyan) in LO1. Scale bar 10μm.

Figure S10. Fly optomotor response after nAChRα3, nAChRβ1 and mAChR-A knock-down-Related to Figure 5

(A) Schematic representation of the behavioral setup for the optomotor response assessment in nAChRα3 (pale magenta), nAChRβ1 (violet) subunits and mAChR-A (teal) type knock-down flies.

(B) Trajectory (angular velocity) of nAChRα3 knock-down flies (42F06-p65.AD; VT043070-Gal4.DBD>UAS-nAChRα3-RNAi, pale magenta n=18) while presented to a sequence of upward to rightward to downward to leftward OFF edge motion compared to the parental control flies (UAS-nAChRα3-RNAi, grey n=20). The dark lines represent the mean ± SEM.

(C-D) Same as previously for the nAChRβ1 (violet n=18, grey n=19) and mAChR-A (teal n=20, grey n=20) knock-down flies.

(E) Angular velocity of control and AChR knock-down flies. The normality of distribution was assessed with the use of Shapiro-Wilk test. Data is mean ± SEM.

Figure S11. Two-photon calcium imaging in T5 cells-Related to Figure 6

(A) Schematic representation of LO1 (where T5 dendrites reside) and of LOP3 (where T5c axonal terminals reside).

(B) Two-photon view of LOP3 (VT50384-LexA>LexAop-IVS-GCaMP6m).

(C) T4-T5c specific targeting (VT50384-Gal4>UAS-CD8::GFP). Arrows indicate the four LOP layers. Scale bars 40 and 10μm.

(D) Polar plots of the maximum responses (ΔF/F) -along seven flies- across sixteen OFF edge directions (º) of T5c cells under nAChRα4 RNAi (light magenta), nAChRα5 RNAi (magenta), nAChRα7 RNAi (dark magenta), mAChR-B RNAi (cyan) and control mCherry RNAi (black) conditions.

Figure S12. Two-photon calcium imaging in T5 dendrites and T5 axonal terminals-Related to Figure 6

(A) Schematic representation of LO1 (where T5 dendrites reside) and of LOP3 (where T5c axonal terminals reside).

(B) Two-photon view of the LO1 and LOP3 layers (VT50384-LexA>LexAop-IVS-GCaMP6m).

(C) Polar plots of the maximum responses (ΔF/F) across sixteen OFF edge directions (°) of T5c cells under mCherry controls (black, flies n=5, ROIs N=10) and nAChRα5 RNAi (magenta, n=5, N=13) conditions in LO1 (left) and in LOP3 (right). The dark lines represent the mean (68% CI).