Genetic suppression interactions are highly conserved across genetic backgrounds

Claire Paltenghi and Jolanda van Leeuwen[#]

Center for Integrative Genomics, University of Lausanne, Lausanne, Switzerland
[#] Correspondence: jolanda.vanleeuwen@unil.ch

ABSTRACT

Genetic suppression occurs when the phenotypic defects caused by a deleterious mutation are rescued by another mutation. Suppression interactions are of particular interest for genetic diseases, as they identify ways to reduce disease severity, thereby potentially highlighting avenues for therapeutic intervention. To what extent suppression interactions are influenced by the genetic background in which they operate remains largely unknown. However, a high degree of suppression conservation would be crucial for developing therapeutic strategies that target suppressors. To gain an understanding of the effect of the genetic context on suppression, we isolated spontaneous suppressor mutations of temperature sensitive alleles of *SEC17*, *TAO3*, and *GLN1* in three genetically diverse natural isolates of the budding yeast *Saccharomyces cerevisiae.* After identifying and validating the genomic variants responsible for suppression, we introduced the suppressors in all three genetic backgrounds, as well as in a laboratory strain, to assess their specificity. Ten out of eleven tested suppression interactions were conserved in the four yeast strains, although the extent to which a suppressor could rescue the temperature sensitive mutant varied across genetic backgrounds. These results suggest that suppression mechanisms are highly conserved across genetic contexts, a finding that is potentially reassuring for the development of therapeutics that mimic genetic suppressors.

INTRODUCTION

Predicting phenotype from genotype remains challenging. Although some mutations, such as Mendelian disease alleles, are detrimental in nearly all individuals, the phenotype of most mutations is influenced by their environmental or genetic context, complicating the prediction of a mutation's phenotype (Nadeau 2001, Chandler et al. 2013, Cooper et al. 2013, Busby et al. 2019, Turco et al. 2023). Genetic context-dependency arises when modifying mutations either increase the severity of a genetic trait or protect against the deleterious effects of a particular mutation (Genin et al. 2008, Harper et al. 2015). Protective modifiers, also called suppressors, can occur in the same gene as the detrimental mutation, or may affect another gene (Lehner 2011, Van Leeuwen et al. 2017). Because suppressors can rescue deleterious phenotypes, suppressors of disease alleles may reveal new therapeutic avenues for treating genetic diseases (Esrick et al. 2021, Frangoul et al. 2021, Ünlü et al. 2023). For example, loss-of-function variants in *BCL11A*, encoding a transcriptional repressor of fetal hemoglobin subunit γ,

lead to expression of this subunit in adults. The γ subunit can functionally replace the
hemoglobin β subunit, which is compromised in β-thalassemia patients, thereby protecting
carriers of *BCL11A* loss-of-function variants against severe β-thalassemia (Uda et al. 2008).
This finding led to the development of a gene editing therapy targeting *BCL11A* (Frangoul et al.
2021), which was recently approved for treating β-thalassemia. Despite the success of this
therapy aimed at a genetic suppressor, for suppressors to be widely adopted for clinical
targeting, they must be conserved across individuals with diverse genetic backgrounds.
However, to what extent suppression interactions are influenced by the genetic context in which
they operate remains unknown.
54   Previous studies, focused on the genetic context dependency of particular genetic
interactions of interest, have mainly described large differences in interactions between genetic
backgrounds (Chari and Dworkin 2013, Wang et al. 2013, Filteau et al. 2015, Mullis et al. 2018).
Similarly, a systematic study of the genetic interactions of three yeast genes involved in sterol
homeostasis in four genetically diverged yeast strains found that the vast majority of synthetic
sick or lethal interactions, in which the combination of two viable mutants leads to a severe
fitness defect or lethality, were unique to one genetic background (Busby et al. 2019). However,
the generality of these findings for genes involved in other cellular processes remains
uncertain. Furthermore, compared to other types of genetic or physical interactions, extragenic
suppression interactions are relatively rare and highly enriched for connecting genes that
function in the same protein complex or pathway (Van Leeuwen et al. 2016, Van Leeuwen et al.
2020). These properties of genetic suppression may lead to differences in genetic background
dependency compared to other types of interactions.
67   Here, we harnessed the powerful genetics of the budding yeast *Saccharomyces*
cerevisiae to study the genetic context-dependency of suppression interactions. We find that
the vast majority of identified interactions were conserved in the four tested genetic
backgrounds. Nonetheless, the strength of the suppression phenotype varied across contexts,
and was sometimes dependent on the sequence or expression level of the suppressor allele.
These results suggest that suppression interactions are highly conserved across genetic
backgrounds, but that the extent of suppression is influenced by additional genetic variants
present in the genome.
75
76
77   **RESULTS**
78
Systematic identification of genetic suppressors
To study the conservation of suppression interactions across yeast strains, we selected three
functionally diverse "query" genes (*SEC17*, *TAO3*, and *GLN1*). The three query genes are
essential for cell viability and are involved in the fusion of vesicles transiting between organelles
(*SEC17*) (Clary et al. 1990), regulation of the RAM signaling network for cell proliferation
(*TAO3*) (Nelson et al. 2003), and the synthesis of glutamine (*GLN1*) (Mitchell 1985). We used
six sequential backcrosses to transfer temperature sensitive (TS) alleles of the three query
genes from the S288C reference background into three genetically diverse budding yeast
strains of distinct geographical locations and sources: L-1374, UWOPS87-2421, and NCYC110
(**Fig. 1A**) (Liti et al. 2009). The three yeast strains have a nucleotide divergence of 0.40, 0.59,

and 0.69%, respectively, compared to the reference strain. After six backcrosses, ~98% of this
genetic divergence should be maintained. All TS alleles still showed a temperature sensitive
phenotype in the various strain backgrounds (**Fig. S1**). For *TAO3*, however, the restrictive
temperature of the *tao3-5010* allele varied from 30°C in UWOPS87-2421 to 38°C in S288C,
suggesting that the severity of the allele was affected by genomic variants present in these
strains.
We used the temperature sensitive phenotype of the constructed strains to isolate
spontaneous suppressor mutants that could rescue the growth defect at high temperature. For
each TS allele, we isolated three independent suppressor colonies per genetic background, for
a total of 27 suppressor strains (**Fig. 1B, S1**). To identify the suppressor mutations, we
sequenced the genomes of all 27 suppressor strains and the 9 corresponding parental strains.
We identified 23 SNPs and 23 segmental or full aneuploidies that were present in a suppressor
strain but not in the parental strain (**Fig. 1C**; **Data S1**, **S2**). Out of the 23 detected SNPs, 5
occurred in intergenic regions, 8 introduced premature stop codons or frameshifts that most
likely led to loss of gene function, and 10 encoded missense variants. Most strains that carried
nonsynonymous mutations did not carry aneuploidies, and vice versa. In total, 7 out of 27
suppressor strains were euploid and carried one or more nonsynonymous SNPs, 16 carried
(partial) chromosomal duplications and no nonsynonymous SNPs, and 3 carried both a
nonsynonymous SNP and an aneuploidy (**Data S3**). In the remaining suppressor strain, we
could not identify any SNPs or other genomic alterations.
109
Validating potential suppressor candidates
To determine which of the discovered genomic alterations were responsible for the suppression
phenotype, we tested the effect of deletion and/or overexpression of the mutated genes on the
temperature sensitivity of the query mutants (**Fig. 1D**, **S2-S4**; **Data S3**). In several cases,
multiple suppressor strains carrying the same TS allele showed identical chromosome
duplications (**Fig. 1C**, **Data S3**), suggesting that the suppression phenotype was caused by an
increased copy number of one or more genes encoded on the affected chromosome. In 17
suppressor strains, the aneuploid chromosome carried the query TS allele itself, suggesting that
increased dosage of the query allele caused the suppression. Indeed, transforming the parental
TS strain (without the suppressor) with a plasmid carrying an extra copy of the TS allele
improved fitness of all tested TS strains at elevated temperature (**Fig. 1D**; **Data S3**). However,
in addition to the query allele itself, we suspected that in some cases other genes on the
aneuploid chromosomes contributed to the suppression phenotype, as the fitness improvement
caused by *sec17-1* and *tao3-5010* overexpression was modest in some backgrounds (**Fig.**
S2A, **S3A**).
All *sec17-1* suppressor strains carried a duplication of chromosome II, which carries
sec17-1. A previous study found that overexpression of either *SEC18* or *SCT1*, both located on
chromosome II, could suppress *sec17-1* in S288C (Magtanong et al. 2011). We confirmed that
overexpression of the S288C alleles of *SEC18* and *SCT1* could also suppress the *sec17-1* TS
phenotype in the three natural genetic backgrounds (**Fig. 1D**; **Data S3**). Furthermore, the
NCYC110 suppressor strains also carried a duplication of chromosome XII. Although there are
no known dosage suppressors of *SEC17* located on this chromosome, it carries multiple genes
with roles in vesicular transport. We tested five of these genes and found that only

Figure 1. Systematic identification of genetic suppressors in diverse genetic backgrounds. (**A**) Phylogenetic tree of *Saccharomyces cerevisiae*, indicating the yeast isolates that were used in this study. Adapted from Batté et al., 2024. (**B**) Validating the suppression phenotype of isolated suppressor strains. Three TS alleles (*sec17-1*, *tao3-5010*, and *gln1-5007*) were introduced into three natural yeast isolates (L-1374, UWOPS87-2421, and NCYC110) and three independent, spontaneous suppressors of the TS phenotype were isolated in each background. An example of isolated suppressor strains of one TS allele

(*sec17-1*) in one genetic background (L-1374) is shown here, all other combinations are shown in **Fig. S1**.
Cultures of the isolated suppressor strains, as well as of the corresponding parental TS strain without a
suppressor, were grown until saturation, and a series of ten-fold dilutions was spotted on YPD plates. Plates
were incubated at the indicated temperatures for 2 days. The wild-type natural isolate (without TS allele)
was included as a control. (**C**) All suppressor colonies, as well as the parental TS strains without
suppressor, were sequenced whole genome. Shown is the average sequencing read depth for each of the
16 yeast chromosomes in each of the strains. Dark yellow indicates the presence of additional copies of
the affected chromosome. (**D**) Validation of candidate suppressor genes. For candidate suppressor genes
that were either located on one of the aneuploid chromosomes or that carried a nonsynonymous mutation,
we tested the effect of deletion and/or overexpression of the genes on the temperature sensitivity of the
query mutants. Blue squares indicate that the gene was found and validated as a suppressor of the TS
allele in a given strain. Details on detected SNPs and aneuploidies can be found in **Data S1** and **S2**. Spot
dilution assays of the suppressor validation experiments are shown in **Fig. 2**, **3**, and **S2**-**S4** and results are
summarized in **Data S3**.
154
overexpression of *SEC22* could suppress the *sec17-1* TS phenotype (**Fig. 1D**, **S2C**; **Data S3**).
Similarly, all NCYC110 *tao3-5010* suppressor strains carried an aneuploidy of chromosome IX,
which carries *tao3-5010*. We validated that overexpression of *SIM1*, also located on
chromosome IX and previously reported as a dosage suppressor of a *tao3* TS mutant in S288C
(Du and Novick 2002), could suppress the *tao3-5010* TS allele in the NCYC110 background
(**Fig. 1D**; **Data S3**).
To investigate a potential role for the identified nonsynonymous SNPs in the
suppression phenotype, we introduced a plasmid carrying the wild-type alleles of the potential
suppressor genes into the suppressor strains. If the suppressor mutation is recessive or semi-
dominant, overexpression of the wild-type allele of the suppressor gene is expected to reverse
the suppression and reduce the fitness of the suppressor strain. Using this strategy, we could
not confirm a role for *NMD2*, *ZDS2*, *CYR1*, or *VTS1* in the suppression of *GLN1*, or for *MED1* in
the suppression of *TAO3* (**Fig. 1D**; **Data S3**). However, expression of wild-type *SSD1* in L-1374
and NCYC110 *tao3-5010* suppressor strains carrying a missense variant in *SSD1* did revert the
suppression phenotype, validating *SSD1* as the suppressor gene (**Fig. 1D**; **Data S3**).
Furthermore, we deleted *SSD1* and *CWP2*, that carried potential loss-of-function variants in L-
1374 and/or NCYC110 *tao3-5010* suppressor strains, in the parental strains that carry the TS
allele but not the suppressor variant and confirmed that deletion of either of the genes could
suppress *tao3* in these genetic backgrounds (**Fig. 1D**; **Data S3**). Similarly, we validated that
deletion of *LUG1*, which carried loss-of-function variants in the NCYC110 *gln1-5007* suppressor
strains, could suppress the temperature sensitivity of the parental NCYC110 *gln1-5007* strain
(**Fig. 1D**; **Data S3**). Overall, we validated one or more suppressor genes in 22 out of 27
suppressor strains (**Fig. 1D**; **Data S3**).
178
Suppression interactions are highly conserved across genetic backgrounds
In several cases, a particular suppressor gene was identified in only one genetic background.
For example, *CWP2* was identified as a suppressor of *tao3-5010* only in the L-1374 background
and loss-of-function mutations in *LUG1* were identified only in NCYC110 *gln1-5007* strains.
However, these differences in observed suppressors across genetic backgrounds could be due
to random chance or experimental factors. To directly investigate whether the identified

suppressors were unique to a specific genetic context, we introduced deletion or
overexpression alleles of the suppressors into all three natural strains, as well as in S288C, all
carrying the query TS allele. For *GLN1*, we also tested for suppression by deletion of *PMR1*, a
suppressor gene we had previously identified in the S288C background (our unpublished
results) but not in any of the other backgrounds. Overexpression of *sec17-1*, *SEC18*, *SIM1*, or
gln1-5007 and deletion of *SSD1*, *CWP2*, or *LUG1* could suppress the corresponding query TS
alleles in all four genetic backgrounds (**Table 1**; **Fig. S2-S4**). We did not succeed in deleting
PMR1 in the NCYC110 *gln1-5007* strain, but suppression was observed in the three remaining
genetic backgrounds (**Table 1**; **Fig. S4E**). In contrast, overexpression of *tao3-5010* could
suppress *tao3-5010* temperature sensitivity in the three natural backgrounds but not in S288C,
possibly because of the high restrictive temperature of the allele in this genetic background
(**Table 1**; **Fig. S3A**).
197

Query allele	Suppressor allele	S288C	L-1374	UWOPS87-2421	NCYC110
sec17-1	sec17-1 o.e.	✓	✓	✓	✓
	SEC18 o.e.	✓	✓	✓	✓
	SCT1 o.e.	✓	✓	✓*	✓*
	SEC22 o.e.	✓	✓	✓*	✓
tao3-5010	tao3-5010 o.e.	-	✓	✓	✓
	ssd1Δ	✓	✓	✓	✓
	cwp2Δ	✓	✓	✓	✓
	SIM1 o.e.	✓	✓	✓	✓
gln1-5007	gln1-5007 o.e.	✓	✓	✓	✓
	lug1Δ	✓	✓	✓	✓
	pmr1Δ	✓	✓	✓	n.d.

198 **Table 1. Conservation of genetic suppression.** For each of the detected suppressor alleles, its
199 suppression phenotype was tested in all three natural yeast isolates, as well as in S288C. ✓ = suppression
200 was observed in the indicated background; - = suppression was not observed in the indicated background;
201 * = suppression was dependent on the expression level of the suppressor allele; n.d. = not determined; o.e.
202 = overexpression. Spot dilution assays of the suppression assays are shown in **Fig. 2**, **3**, and **S2-S4.**
203
For *SCT1* and *SEC22*, that could both suppress *sec17-1*, suppression was dependent
on the expression level of the suppressor gene. A *CEN*-plasmid (low-copy) containing *SCT1*
could suppress *sec17-1* in S288C and L-1374, but a *2μ*-plasmid (high-copy) was needed to see
suppression in the NCYC110 or UWOPS87-2421 backgrounds (**Fig. 2A**). Because we were
using S288C alleles in the overexpression experiments, we tested whether overexpression of
the *SCT1* allele from NCYC110 or UWOPS87-2421 could suppress *sec17-1* in these
backgrounds when expressed from a low-copy plasmid. However, also the NCYC110 and
UWOPS87-2421 *SCT1* alleles were not able to suppress *sec17-1* in these backgrounds when
expressed from a *CEN*-plasmid (**Fig. 2B**). Similarly, overexpression of *SEC22* from a *CEN*-
plasmid could rescue *sec17-1* strains in most genetic backgrounds, but not in UWOPS87-2421
(**Fig. 3A**). In this case, further increasing the level of overexpression of the *SEC22* S288C did
not rescue *sec17-1* in the UWOPS87-2421 background (**Fig. 3A**), but overexpression of the
SEC22 UWOPS87-2421 allele showed weak but reproducible suppression (**Fig. 3B**). Although

the sequence of the *SEC22* ORF is identical in UWOPS87-2421 and S288C, the UWOPS87-
2421 allele contains a C-to-T variant in the 5' UTR, 69 nucleotides upstream of the start codon.
We investigated the effect of this UWOPS87-2421-specific variant on *SEC22* mRNA levels
using RNA sequencing and found an ~25% increase in *SEC22* expression in the wild-type
UWOPS87-2421 strain compared to S288C (**Fig. 3C**; **Data S4**). Possibly, suppression of
sec17-1 by *SEC22* in the UWOPS87-2421 background is sensitive to small changes in *SEC22*
expression. Expressing the S288C *SEC22* allele from a *CEN-* or *2μ*-plasmid may lead to too
low or too high expression, respectively, while expression of the UWOPS87-2421 *SEC22* allele
from a *CEN* plasmid may achieve a level of expression that is just right for suppression to
occur.
227

Figure 2. Different levels of *SCT1* expression are needed for suppression of *SEC17* across genetic
backgrounds. (**A**) S288C, L-1374, UWOPS87-2421, and NCYC110 strains carrying the *sec17-1* TS allele
were transformed with a low-copy (*CEN*) or a high-copy (*2μ*) plasmid expressing *SCT1* or the
corresponding empty vector. Cultures of two to three independent transformants were grown until
saturation, and a series of ten-fold dilutions was spotted on SD-Ura (low-copy) or SD-Leu (high-copy)
plates. Plates were incubated at the indicated temperatures for 3 days. Pictures of one representative
transformant are shown for each genotype. Rare, larger colonies that appear at higher temperatures are
spontaneous suppressor mutants that sometimes occur during the experiments. (**B**) As in (A), but using
the *SCT1* alleles from the L-1374, UWOPS87-2421, and NCYC110 backgrounds, rather than the S288C
allele. UWOPS = UWOPS87-2421; NCYC = NCYC110.

Thus, 10 out of 11 suppression mechanisms tested in this study (8 out of 8 when
excluding suppression by overexpression of the query allele) were conserved in all tested
genetic backgrounds, with only the required expression level of the suppressor gene changing
with the genetic context. Despite the high conservation of suppressor genes, the relative

strength of the suppressors varied between genetic backgrounds. For example, overexpression
of *SIM1* could strongly suppress *tao3-5010* in the NCYC110 and L-1374 backgrounds, but only
weakly improved fitness in the S288C and UWOP87-2421 backgrounds (**Fig. S3B**). Such
differences in the intensity of suppression across genetic backgrounds were common and
observed for nearly all suppressor genes. A notable exception is *ssd1Δ*, which could strongly
suppress *tao3-5010* in all backgrounds (**Fig. S3E**).
250

251
Figure 3. Suppression of *SEC17* by *SEC22* is dependent on the allele sequence. (**A**) S288C, L-1374,
UWOPS87-2421, and NCYC110 strains carrying the *sec17-1* TS allele were transformed with a low-copy
(*CEN*) or a high-copy (*2μ*) plasmid expressing *SEC22* or the corresponding empty vector. Cultures of two
to three independent transformants were grown until saturation, and a series of ten-fold dilutions was
spotted on SD-Ura (low-copy) or SD-Leu (high-copy) plates. Plates were incubated at the indicated
temperatures for 3 days. Pictures of one representative transformant are shown for each genotype. (**B**) A
UWOPS87-2421 strain carrying the *sec17-1* TS allele was transformed with a low-copy vector carrying
either the S288C or the UWOPS87-2421 version of *SEC22*, or the corresponding empty vector. Spot
dilutions were performed as in (A). (**C**) Expression levels of the indicated genes in wild-type S288C or
UWOPS87-2421 strains were determined by RNA sequencing. Plotted are RPKM (reads per kilobase per
million mapped reads) values, normalized to the total number of reads in a sample and averaged over three
technical replicates. Error bars indicate the standard deviation. * $p<0.05$, two-sided Student's t-test.
UWOPS = UWOPS87-2421.

265
Multiple genes can contribute to the suppression phenotype
In a few instances, we had identified multiple genes on aneuploid chromosomes that could each
independently suppress the TS phenotype (**Data S3**). We hypothesized that in these cases the
suppressors could have an additive effect, and that the combined overexpression of multiple
suppressor genes may further improve the fitness of the TS mutant at higher temperature. To
test this, we combined overexpression of *SEC18*, *SEC22*, and *SCT1* in *sec17-1* strains. *SEC18*
and *SCT1* are both located on chromosome II, which was duplicated in all *sec17-1* suppressor
strains, and *SEC22* is located on chromosome XII, which was duplicated together with
chromosome II in the NCYC110 *sec17-1* suppressor strains. We constructed a collection of

CEN-plasmids, each expressing a natural allele of one of the three genes and a different
selectable marker and verified that each gene individually could suppress *sec17-1* in the same
backgrounds as described above (**Fig. S5A**). We then overexpressed all possible combinations
of the three suppressors in their respective genetic background in presence of the *sec17-1* TS
allele (**Fig. 4, S5B**). To be able to detect small differences in fitness, we quantified the size of
on average ~200 colonies per strain (**Fig. 4**).
281

Figure 4. Multiple genes can contribute to the suppression phenotype. Relative fitness of *sec17-1*
strains overexpressing *SEC18*, *SEC22*, and/or *SCT1* in the S288C, L-1374, UWOPS87-2421, and
NCYC110 genetic backgrounds. In each case, the *SEC18*, *SEC22*, and *SCT1* alleles matched the genetic
background in which they were transformed, such that S288C was transformed with S288C alleles and L-
1374 with L-1374 alleles, etc. Strains were spotted on SD-Leu-Ura-His and colony size was determined
after two days of growth at 33°C and normalized to the colony size of a wild-type strain with the same
genetic background. Boxplots show the normalized colony size (fitness) of, on average, ~200 colonies per
strain. Tukey's test was used to determine whether strains overexpressing two or three genes had a
significantly higher fitness than strains overexpressing one of the genes. + = strains were transformed with
the indicated plasmid; - = strains were transformed with the corresponding empty vector. * p<0.05; **
p<0.005; *** p<0.0005.

In S288C and UWOPS87-2421, the suppression was mainly driven by the strongest
suppressor of *sec17-1*, *SEC18*, and little further increase in suppression was observed when
SCT1 and/or *SEC22* were overexpressed simultaneously with *SEC18* (**Fig. 4**). In contrast, in L-
1374, overexpression of all three genes simultaneously resulted in stronger suppression than
overexpression of each of the genes alone (**Fig. 4**). Combining *SEC18* (chromosome II) and
SEC22 (chromosome XII) overexpression strongly improved fitness compared to
overexpression of *SEC18* or *SEC22* alone in the NCYC110 *sec17-1* strain, which is also the
genetic background in which combined duplication of chromosome II and XII was observed.
These results show that multiple genes can contribute to the suppression phenotype caused by
aneuploidies, and that the relative contribution of the individual genes to the overall suppression
varies between genetic backgrounds.

DISCUSSION

In this study, we investigated the conservation of genetic suppression interactions across
natural yeast isolates using three mutant alleles of functionally diverse query genes. Ten out of
eleven mechanisms of suppression that spontaneously occurred in the natural yeast strains
could be reproduced in all four tested genetic backgrounds, including the laboratory strain
S288C. Despite the high conservation of suppression interactions, the extent of suppression
was often variable between backgrounds, and sometimes depended on the expression level
(**Fig. 2**) or allele sequence (**Fig. 3**) of the suppressor gene. Similarly, a previous study
compared five suppressor genes of a *las17* TS allele in the yeast strains S288C and RM11-1a
and found that the strength of the suppression phenotype varied between the two strains and
was in some cases influenced by the particular suppressor mutation (Filteau et al. 2015). Some
of the differences in strength and required sequence of the suppressor genes could be due to
differences in the restrictive temperature of the TS allele between genetic backgrounds. A large
difference in restrictive temperature may also explain why the *tao3-5010* mutant could not be
rescued by overexpression of the TS allele in S288C, in which the mutant had a restrictive
temperature of 38°C, in contrast to the other genetic backgrounds where the restrictive
temperature was ~8-10 degrees lower (**Fig. S3A**). Possibly, the remaining functionality of the
tao3-5010 allele at 38°C is insufficient to support proliferation. Alternatively, these differences
may result from strain-specific variants in additional genes.

Most of the isolated suppressor strains in the natural backgrounds carried aneuploidies
(19 out of 27, 70%), and in 17 out of 19 cases we validated that one or more genes on the
aneuploid chromosome were responsible for the suppression phenotype (**Data S3**). This
frequency of aneuploidies is significantly higher than what we generally observe for suppressors
of TS alleles in S288C (~16% carry aneuploidies, our unpublished results). We suspect that this
difference in aneuploidy occurrence is due to natural yeast strains being relatively tolerant to
aneuploidies when compared to S288C (Hose et al. 2015). Aneuploidies are associated with a
growth defect in laboratory yeast strains, independently of which chromosome is duplicated
(Torres et al. 2007, Beach et al. 2017). In contrast, the sequencing of more than a thousand
yeast isolates showed that ~20% of natural *S. cerevisiae* strains carry at least one aneuploidy
(Peter et al. 2018). Because aneuploidies commonly occur during cell division (Gilchrist and
Stelkens 2019), the enhanced tolerance for aneuploidies may increase the frequency at which
suppression occurs, which could be an advantage in highly selective natural environments.
Furthermore, we showed that multiple genes on an aneuploid chromosome can contribute to
the suppression phenotype (**Fig. 4**), further increasing the benefit associated with an
aneuploidy.

Out of the eight extragenic suppressors that we identified in this study, four had not
been described previously. For example, we found that overexpression of SNARE protein
Sec22 could suppress mutants of the SNARE chaperone Sec17 (**Fig. 3**), likely by (partially)
restoring vesicle fusion (Liu and Barlowe 2002, Song et al. 2021). Furthermore, we discovered
loss-of-function mutations in *CWP2*, encoding a major cell wall mannoprotein, as suppressors
of the RAM signaling network member Tao3 (**Fig. S3E**). Cells with an inactive RAM network

display a separation defect of mother and daughter cell walls due to the inability to activate
Ace2, which is needed for the expression of cell separation genes (Nelson et al. 2003).
Possibly, the changes in cell wall composition induced by loss of *CWP2* (Van der Vaart et al.
1995, Li et al. 2020) can promote the separation of mother and daughter cells in the absence of
a functioning RAM network. We also found that loss-of-function mutations in *PMR1* could
suppress a TS mutant of the glutamine synthetase Gln1 in all tested genetic backgrounds (**Fig.
S4E**). Pmr1 shuttles calcium and manganese (Mn^{2+}) ions into the Golgi lumen, and loss of Pmr1
leads to increased intracellular levels of Mn^{2+} (Lapinskas et al. 1995, Durr et al. 1998).
Glutamine synthetases are activated by Mn^{2+} ions (Monder 1965, Tholey et al. 1987),
suggesting that loss of *PMR1* may suppress the *GLN1* mutant by boosting its activity. Finally,
we uncovered loss-of-function mutations in the poorly characterized *LUG1* gene as suppressors
of *GLN1* (**Fig. S4D**). Mutations in *GLN1* were previously described to suppress *lug1Δ* mutants,
indicating that this suppression interaction is reciprocal (Edskes et al. 2018). Overall, this high
frequency of newly identified suppressor genes indicates that despite several large-scale
suppressor mapping efforts (Magtanong et al. 2011, Patra et al. 2016, Van Leeuwen et al.
2016, Van Leeuwen et al. 2020), the yeast suppression interaction space remains largely
unexplored.
In conclusion, although different genetic backgrounds have the potential to reveal novel
suppression interactions and thus uncover previously unidentified functional connections
between genes (Filteau et al. 2015), our results suggest that genetic suppression interactions
are largely robust to changes in genetic context. While the extent of conservation of
suppression interactions across human genomes remains to be determined, our finding is
potentially reassuring for the development of new therapeutic strategies that target suppressor
genes (Esrick et al. 2021, Frangoul et al. 2021, Ünlü et al. 2023).
374
375
376 **METHODS**
377
Yeast strains, plasmids and growth
Yeast strains were grown using standard rich (YPD) or minimal (SD) media. For overexpression
assays using S288C alleles, plasmids from either the MoBY-ORF 1.0 (native promoter,
CEN/ARS, URA3, kanMX4) (Ho et al. 2009) or the MoBY-ORF 2.0 (native promoter, 2µ, *LEU2,
kanMX4*) (Magtanong et al. 2011) collection were used. All yeast strains and plasmids used in
this study are listed in **Data S5**.
384
Introducing TS alleles into multiple genetic backgrounds
The three natural yeast isolates, L-1374 (LY00010), UWOPS87-2421 (LY00011), and
NCYC110 (LY00015), were previously (partially) deleted for *HO, URA3, HIS3*, and *LEU2*,
resulting in the genotype MATa *hoΔ::hphMX6 ura3Δ::kanMX his3Δ1 leu2Δ0* (Cubillos et al.
2009, Parts et al. 2021). The three genetically diverse strains were each crossed with three
different S288C strains carrying a TS allele, TSQ48 (*sec17-1*), TSQ2433 (*gln1-5007*), and
TSQ2031 (*tao3-5010*), with genotype MATα *xxx-ts::natMX4 can1Δ::STE2pr-Sp_his5 lyp1Δ0
his3Δ1 leu2Δ0 ura3Δ0 met15Δ0*. The resulting diploids were driven through meiosis and haploid
MATα segregants carrying the TS allele were isolated and crossed again to their respective

natural parental strain. This process was repeated four more times for a total of six crosses per
strain background and TS allele, resulting in progeny with a genome that is 98% identical to the
natural parental strain. Two independent spores carrying the TS allele were isolated from the
final crosses and frozen at -80°C.
The S288C and L-1374 *sec17-1* strains carried the *lyp1Δ::STE3pr-LEU2* cassette,
complicating some of the suppressor validation experiments that used *LEU2*-plasmids. To
remove the *lyp1Δ::STE3pr-LEU2* cassette, we first cloned *LEU2*-targeting guide RNA
sequences into the pML104 vector, which carries Cas9 and a *URA3* selection marker (**Data S5**)
(Laughery et al. 2015). Next, we PCR-amplified *LYP1*, including promoter and terminator
sequences, from a wild-type strain and co-transformed the *lyp1Δ::STE3pr-LEU2* strains with the
LYP1 PCR-product and the pML104-LEU2-2 plasmid. Transformants were selected on SD-Ura
and subsequently streaked on SD-Leu and SD-Lys+LYP (thialysine) to confirm loss of *LEU2*
and proper integration of *LYP1*. The final genotypes and strain IDs of the resulting strains are
listed in **Data S5**.
408
Isolating spontaneous suppressor mutations
For each TS allele in each genetic background, ~25 million cells were spread onto three
YPD+NAT agar plates and incubated for three days at the restrictive temperature of the strain.
Most cells will not be able to grow at the restrictive temperature, except for those that have
acquired a spontaneous suppressor mutation, that will grow up to form a colony. When colonies
were observed, one colony per plate was isolated and its growth at the restrictive temperature
was compared to the parental TS strain to confirm the suppression phenotype. In total, three
independent suppressors per query allele and per genetic background were isolated.
417
Sequencing, read mapping, and SNP calling
All suppressor strains as well as the corresponding parental TS strains were sequenced on the
DNBseq platform using paired-end 100-bp reads, with an average read depth of ~100x. Reads
were aligned to the S288C reference genome version R64.2.1 using BWA v0.7.17 (Li and
Durbin 2009). Pileups were processed and variants were called using SAMtools/BCFtools v1.11
(Li et al. 2009). Variants that had a Phred quality score <200, that were present in one of the
parental strains, or that were found in more than three of the suppressor strains were removed
from consideration. The consequence of detected variants was determined using Ensembl's
VEP (McLaren et al. 2016). All whole-genome sequencing data are publicly available at NCBI's
Sequence Read Archive (http://www.ncbi.nlm.nih.gov/sra) under accession number
PRJNA1100912. Variants are listed in **Data S1**.
429
Aneuploidy and ploidy assessment
Qualimap v2.3 (Okonechnikov et al. 2016) was used to detect (partial) aneuploidies based on
variation in sequencing read depth across windows of 30,000 base pairs in the nuclear genome
(**Data S2**). We note that the smaller chromosomes I, III, and VI showed a higher variation in
read count between samples than other chromosomes, likely due to variation in the capture of
these small chromosomes during genomic DNA isolation. Because the relative increase in
coverage caused by an aneuploidy depends on the overall ploidy, we analyzed all suppressor
strains by flow cytometry to determine ploidy. Briefly, cells were grown until log-phase

(OD$_{600}$≈0.5) and fixed with 70% ethanol. Fixed cells were washed with water and subsequently
treated with RNase A (200 μg/ml, 2 h, 37°C) and proteinase K (2 mg/ml, 40 min, 50°C). Treated
cells were washed with 200 mM Tris–HCl, 200 mM NaCl, 78 mM MgCl2 (pH 7.5) and stained
with 2× SYBR Green (Life Technologies) in 50 mM Tris–HCl (pH 7.5). Aggregates of cells were
dispersed via sonication and cells were analyzed by flow cytometry using a SONY SH800
FACS machine. DNA content was compared to known haploid and diploid controls. Normalized
average read depth per genomic region was corrected based on the observed DNA content,
such that the average normalized read depth of a genomic region in a diploid strain was twice
that of a haploid strain. Detected aneuploidies are summarized in **Data S3**.
447
Predicting and validating suppressor genes
For suppressor strains that carried an aneuploidy, we predicted potential causal suppressor
genes based on the functional relationships between the query gene and the genes located on
the aneuploid chromosome. We used BioGRID 4.4 (Oughtred et al. 2021) to identify genes that
are known to interact with the query gene (either genetically or physically) and the
Saccharomyces Genome Database (Wong et al. 2023) to identify genes that function in similar
or related biological processes as the query. Identified candidate suppressors were validated by
transforming plasmids expressing the candidate gene into the parental TS strain, without the
suppressor, using standard transformation protocols (Gietz and Schiestl 2007). Overnight
cultures of three independent transformants were diluted to an OD$_{600}$ of 0.1, serially diluted 1:10
with sterile water, and spotted onto agar plates. Plates were incubated at a range of
temperatures between 26°C and 38°C. After 2-3 days of incubation, pictures were taken and
the relative fitness of the transformants was compared to empty vector controls.
To test whether detected nonsynonymous SNPs contributed to the suppression
phenotype, we introduced a plasmid carrying the wild-type allele of the potential suppressor
gene into the suppressor strain. If the suppressor mutation is recessive or semi-dominant,
overexpression of the wild-type allele of the suppressor gene is expected to reverse the
suppression and reduce the fitness of the suppressor strain. Transformations and spot dilutions
assays were performed as described above for the aneuploidy suppressors. Validated
suppressor genes are listed in **Data S3**.
468
Suppression by overexpression of natural alleles
To test for suppression by overexpression of the natural alleles of suppressor candidates, we
constructed plasmids carrying these alleles. We PCR-amplified the suppressor candidates
including ~1000 bp upstream of the start codon and ~500 bp downstream of the stop codon
from the various natural yeast strains, thereby including regions of homology to plasmid
pRS313, pRS315, or pRS316 (Sikorski and Hieter 1989) (**Data S5**). The PCR product was co-
transformed with the corresponding linearized vector into LY00004 (BY4742; **Data S5**). The
assembled plasmid was isolated from the yeast strain and correct insertion of the PCR product
was verified using whole plasmid sequencing. Plasmids were transformed into parental TS
strains and tested for suppression as described above ("*Predicting and validating suppressor*
genes").
480
Validation of SSD1, CWP2, PMR1, and LUG1

To test whether deletion of *SSD1* or *CWP2* could suppress *tao3* TS alleles in the S288C
genetic background, *ssd1Δ* (DMA1035; **Data S5**) and *cwp2Δ* (DMA2828; **Data S5**) strains were
crossed to *tao3-5005* (TSQ2026; **Data S5**) and *tao3-5014* (TSQ2035; **Data S5**). Diploids were
selected on YPD+NAT+G418, driven through meiosis, and haploid double mutant progeny were
isolated using tetrad dissection. In total, three independent double mutant spores were isolated
per cross.
To test whether deletion of *SSD1* or *CWP2* could suppress *tao3* TS alleles in the natural
genetic backgrounds, we PCR-amplified *CaURA3MX4* from plasmid pFA6:CaURA3MX4
(Goldstein et al. 1999) (**Data S5**), thereby introducing regions of homology to the genomic DNA
directly upstream and downstream of both suppressor genes. The PCR products were
transformed into the natural *tao3-5010* strains and deletion of *CWP2* and *SSD1* was verified by
PCR. A similar strategy was used to delete *PMR1* and *LUG1* in the *gln1-5007* strains, with the
exception that we cloned a guide RNA targeting *PMR1* or *LUG1* into the Cas9-expressing
vector pML107 (Laughery et al. 2015). We then co-transformed the cloned plasmids with the
CaURA3MX4 cassettes to increase the efficiency of gene deletion. All strains were tested for
suppression as described above ("*Predicting and validating suppressor genes*").

RNA sequencing
Overnight cultures of S288C and UWOPS87-2421 were diluted in 10 mL YPD to an OD_{600} of
0.1 and grown for 3-4 hours at 26°C until an OD_{600} of ~0.7-1.0. Cells were collected, washed
with water, snap frozen in liquid nitrogen, and stored at -80°C until RNA extraction. Total RNA
was extracted by first lysing the yeasts with glass beads in trizole, separating the protein-DNA-
RNA phases with chloroform, and precipitating the RNA with isopropanol and glycogen. The
resulting RNA was washed with 70% ethanol, dissolved in water, treated with DNAse, and
further cleaned using the Macherey-Nagel NucleoSpin RNA kit. RNA quality was assessed
using a Fragment Analyzer and mRNA was enriched via polyA-selection with the Illumina
Stranded mRNA Prep kit and sequenced on the Element Biosciences AVITI system using 150
base pair, single-end reads with ~20 million reads per sample. Adapters were trimmed from the
reads with Cutadapt v2.5 (Martin 2011) and reads with low complexity sequences were
removed with Reaper v15-065 (Davis et al. 2013). Reads corresponding to ribosomal RNAs
were removed with FastQ Screen v0.11.1 (Wingett and Andrews 2018). Remaining reads were
aligned with STAR v2.5.3a (Dobin et al. 2013) against reference genome R64.2.1. The number
of read counts per gene locus was summarized with HTSeq-count v0.9.1 (Anders et al. 2015)
and normalized to gene length and the total number of reads per sample. Normalized read
counts are listed in **Data S4**.

Quantifying strain fitness
To determine the fitness of strains overexpressing combinations of SEC18, SEC22, and/or
SCT1, saturated cultures of three independent transformants per genotype were diluted 1,000
to 100,000 fold and spotted onto SD-His-Leu-Ura plates. Plates were incubated at 26°C or 33°C
and imaged after two days. All images were edited in an identical way to achieve maximal
sharpness and to increase contrast by 10% and highlights by 5%. Images were then cropped,
and colony size was determined using CellProfiler version 4.2.8 (Carpenter et al. 2006). Colony

sizes were normalized to the median colony size of the wild-type strain of the same genetic background.

DATA AVAILABILITY

All whole-genome sequencing data are publicly available at NCBI's Sequence Read Archive (http://www.ncbi.nlm.nih.gov/sra) under accession number PRJNA1100912. All other data are available in the supplementary data files.

ACKNOWLEDGEMENTS

We thank Sabine van Schie for critical reading of the manuscript. This work was supported by an Eccellenza grant from the Swiss National Science Foundation (PCEGP3_181242 to J.v.L).

CONFLICT OF INTERESTS

The authors declare that they have no conflict of interest.

REFERENCES

Anders, S., P. T. Pyl and W. Huber (2015). "HTSeq--a Python framework to work with high-throughput sequencing data." Bioinformatics **31**(2): 166-169.

Beach, R. R., C. Ricci-Tam, C. M. Brennan, C. A. Moomau, P. H. Hsu, B. Hua, R. E. Silberman, M. Springer and A. Amon (2017). "Aneuploidy causes non-genetic individuality." Cell **169**(2): 229-242 e221.

Busby, B. P., E. Niktab, C. A. Roberts, J. P. Sheridan, N. V. Coorey, D. S. Senanayake, L. M. Connor, A. B. Munkacsi and P. H. Atkinson (2019). "Genetic interaction networks mediate individual statin drug response in Saccharomyces cerevisiae." NPJ Syst Biol Appl **5**: 35.

Carpenter, A. E., T. R. Jones, M. R. Lamprecht, C. Clarke, I. H. Kang, O. Friman, D. A. Guertin, J. H. Chang, R. A. Lindquist, J. Moffat, P. Golland and D. M. Sabatini (2006). "CellProfiler: image analysis software for identifying and quantifying cell phenotypes." Genome Biol **7**(10): R100.

Chandler, C. H., S. Chari and I. Dworkin (2013). "Does your gene need a background check? How genetic background impacts the analysis of mutations, genes, and evolution." Trends Genet **29**(6): 358-366.

Chari, S. and I. Dworkin (2013). "The conditional nature of genetic interactions: the consequences of wild-type backgrounds on mutational interactions in a genome-wide modifier screen." PLoS Genet **9**(8): e1003661.

Clary, D. O., I. C. Griff and J. E. Rothman (1990). "SNAPs, a family of NSF attachment proteins involved in intracellular membrane fusion in animals and yeast." Cell **61**(4): 709-721.

Cooper, D. N., M. Krawczak, C. Polychronakos, C. Tyler-Smith and H. Kehrer-Sawatzki (2013). "Where genotype is not predictive of phenotype: towards an understanding of the molecular basis of reduced penetrance in human inherited disease." Hum Genet **132**(10): 1077-1130.

Cubillos, F. A., E. J. Louis and G. Liti (2009). "Generation of a large set of genetically tractable haploid and diploid Saccharomyces strains." FEMS Yeast Res **9**(8): 1217-1225.

Davis, M. P., S. van Dongen, C. Abreu-Goodger, N. Bartonicek and A. J. Enright (2013). "Kraken: a set of
tools for quality control and analysis of high-throughput sequence data." Methods **63**(1): 41-49.

Dobin, A., C. A. Davis, F. Schlesinger, J. Drenkow, C. Zaleski, S. Jha, P. Batut, M. Chaisson and T. R.
Gingeras (2013). "STAR: ultrafast universal RNA-seq aligner." Bioinformatics **29**(1): 15-21.

Du, L. L. and P. Novick (2002). "Pag1p, a novel protein associated with protein kinase Cbk1p, is required
for cell morphogenesis and proliferation in Saccharomyces cerevisiae." Mol Biol Cell **13**(2): 503-514.

Durr, G., J. Strayle, R. Plemper, S. Elbs, S. K. Klee, P. Catty, D. H. Wolf and H. K. Rudolph (1998). "The
medial-Golgi ion pump Pmr1 supplies the yeast secretory pathway with Ca2+ and Mn2+ required for
glycosylation, sorting, and endoplasmic reticulum-associated protein degradation." Mol Biol Cell **9**(5):
1149-1162.

Edskes, H. K., M. Mukhamedova, B. K. Edskes and R. B. Wickner (2018). "Hermes transposon
mutagenesis shows [URE3] prion pathology prevented by a ubiquitin-targeting protein: evidence for
carbon/nitrogen assimilation cross talk and a second function for Ure2p in Saccharomyces cerevisiae."
Genetics **209**(3): 789-800.

Esrick, E. B., L. E. Lehmann, A. Biffi, M. Achebe, C. Brendel, M. F. Ciuculescu, H. Daley, B. MacKinnon,
E. Morris, A. Federico, D. Abriss, K. Boardman, R. Khelladi, K. Shaw, H. Negre, O. Negre, S. Nikiforow,
J. Ritz, S. Y. Pai, W. B. London, C. Dansereau, M. M. Heeney, M. Armant, J. P. Manis and D. A. Williams
(2021). "Post-Transcriptional genetic silencing of BCL11A to treat sickle cell disease." N Engl J Med
384(3): 205-215.

Filteau, M., V. Hamel, M. C. Pouliot, I. Gagnon-Arsenault, A. K. Dube and C. R. Landry (2015).
"Evolutionary rescue by compensatory mutations is constrained by genomic and environmental
backgrounds." Mol Syst Biol **11**(10): 832.

Frangoul, H., D. Altshuler, M. D. Cappellini, Y. S. Chen, J. Domm, B. K. Eustace, J. Foell, J. de la Fuente,
S. Grupp, R. Handgretinger, T. W. Ho, A. Kattamis, A. Kernytsky, J. Lekstrom-Himes, A. M. Li, F.
Locatelli, M. Y. Mapara, M. de Montalembert, D. Rondelli, A. Sharma, S. Sheth, S. Soni, M. H. Steinberg,
D. Wall, A. Yen and S. Corbacioglu (2021). "CRISPR-Cas9 gene editing for sickle cell disease and beta-
thalassemia." N Engl J Med **384**(3): 252-260.

Genin, E., J. Feingold and F. Clerget-Darpoux (2008). "Identifying modifier genes of monogenic disease:
strategies and difficulties." Hum Genet **124**(4): 357-368.

Gietz, R. D. and R. H. Schiestl (2007). "High-efficiency yeast transformation using the LiAc/SS carrier
DNA/PEG method." Nat Protoc **2**(1): 31-34.

Gilchrist, C. and R. Stelkens (2019). "Aneuploidy in yeast: Segregation error or adaptation mechanism?"
Yeast **36**(9): 525-539.

Goldstein, A. L., X. Pan and J. H. McCusker (1999). "Heterologous URA3MX cassettes for gene
replacement in Saccharomyces cerevisiae." Yeast **15**(6): 507-511.

Harper, A. R., S. Nayee and E. J. Topol (2015). "Protective alleles and modifier variants in human health
and disease." Nat Rev Genet **16**(12): 689-701.

Ho, C. H., L. Magtanong, S. L. Barker, D. Gresham, S. Nishimura, P. Natarajan, J. L. Koh, J. Porter, C. A.
Gray, R. J. Andersen, G. Giaever, C. Nislow, B. Andrews, D. Botstein, T. R. Graham, M. Yoshida and C.
Boone (2009). "A molecular barcoded yeast ORF library enables mode-of-action analysis of bioactive
compounds." Nat Biotechnol **27**(4): 369-377.

Hose, J., C. M. Yong, M. Sardi, Z. Wang, M. A. Newton and A. P. Gasch (2015). "Dosage compensation
can buffer copy-number variation in wild yeast." Elife **4**: e05462.

Lapinskas, P. J., K. W. Cunningham, X. F. Liu, G. R. Fink and V. C. Culotta (1995). "Mutations in PMR1
suppress oxidative damage in yeast cells lacking superoxide dismutase." Mol Cell Biol **15**(3): 1382-1388.

Laughery, M. F., T. Hunter, A. Brown, J. Hoopes, T. Ostbye, T. Shumaker and J. J. Wyrick (2015). "New
vectors for simple and streamlined CRISPR-Cas9 genome editing in Saccharomyces cerevisiae." Yeast
32(12): 711-720.

Lehner, B. (2011). "Molecular mechanisms of epistasis within and between genes." Trends Genet **27**(8):
323-331.

Li, H. and R. Durbin (2009). "Fast and accurate short read alignment with Burrows-Wheeler transform."
Bioinformatics **25**(14): 1754-1760.

Li, H., B. Handsaker, A. Wysoker, T. Fennell, J. Ruan, N. Homer, G. Marth, G. Abecasis, R. Durbin and
S. Genome Project Data Processing (2009). "The Sequence Alignment/Map format and SAMtools."
Bioinformatics **25**(16): 2078-2079.

Li, J., M. M. Zhang, C. Wan, R. Den Haan, F. W. Bai and X. Q. Zhao (2020). "Improved cellulase
production in recombinant Saccharomyces cerevisiae by disrupting the cell wall protein-encoding gene
CWP2." J Biosci Bioeng **129**(2): 165-171.

Liti, G., D. M. Carter, A. M. Moses, J. Warringer, L. Parts, S. A. James, R. P. Davey, I. N. Roberts, A.
Burt, V. Koufopanou, I. J. Tsai, C. M. Bergman, D. Bensasson, M. J. O'Kelly, A. van Oudenaarden, D. B.
Barton, E. Bailes, A. N. Nguyen, M. Jones, M. A. Quail, I. Goodhead, S. Sims, F. Smith, A. Blomberg, R.
Durbin and E. J. Louis (2009). "Population genomics of domestic and wild yeasts." Nature **458**(7236):
337-341.

Liu, Y. and C. Barlowe (2002). "Analysis of Sec22p in endoplasmic reticulum/Golgi transport reveals
cellular redundancy in SNARE protein function." Mol Biol Cell **13**(9): 3314-3324.

Magtanong, L., C. H. Ho, S. L. Barker, W. Jiao, A. Baryshnikova, S. Bahr, A. M. Smith, L. E. Heisler, J. S.
Choy, E. Kuzmin, K. Andrusiak, A. Kobylianski, Z. Li, M. Costanzo, M. A. Basrai, G. Giaever, C. Nislow,
B. Andrews and C. Boone (2011). "Dosage suppression genetic interaction networks enhance functional
wiring diagrams of the cell." Nat Biotechnol **29**(6): 505-511.

Martin, M. (2011). "Cutadapt removes adapter sequences from high-throughput sequencing reads."
EMBnet.journal **17**(1): 10-12.

McLaren, W., L. Gil, S. E. Hunt, H. S. Riat, G. R. Ritchie, A. Thormann, P. Flicek and F. Cunningham
(2016). "The Ensembl Variant Effect Predictor." Genome Biol **17**(1): 122.

Mitchell, A. P. (1985). "The GLN1 locus of Saccharomyces cerevisiae encodes glutamine synthetase."
Genetics **111**(2): 243-258.

Monder, C. (1965). "Metal ion interactions and glutamine synthetase activity." Biochemistry **4**(12): 2677-
2686.

Mullis, M. N., T. Matsui, R. Schell, R. Foree and I. M. Ehrenreich (2018). "The complex underpinnings of
genetic background effects." Nat Commun **9**(1): 3548.

Nadeau, J. H. (2001). "Modifier genes in mice and humans." Nat Rev Genet **2**(3): 165-174.

Nelson, B., C. Kurischko, J. Horecka, M. Mody, P. Nair, L. Pratt, A. Zougman, L. D. McBroom, T. R.
Hughes, C. Boone and F. C. Luca (2003). "RAM: a conserved signaling network that regulates Ace2p
transcriptional activity and polarized morphogenesis." Mol Biol Cell **14**(9): 3782-3803.

Okonechnikov, K., A. Conesa and F. Garcia-Alcalde (2016). "Qualimap 2: advanced multi-sample quality
control for high-throughput sequencing data." Bioinformatics **32**(2): 292-294.

Oughtred, R., J. Rust, C. Chang, B. J. Breitkreutz, C. Stark, A. Willems, L. Boucher, G. Leung, N. Kolas,
F. Zhang, S. Dolma, J. Coulombe-Huntington, A. Chatr-Aryamontri, K. Dolinski and M. Tyers (2021). "The
BioGRID database: A comprehensive biomedical resource of curated protein, genetic, and chemical
interactions." Protein Sci **30**(1): 187-200.

Parts, L., A. Batte, M. Lopes, M. W. Yuen, M. Laver, B. J. San Luis, J. X. Yue, C. Pons, E. Eray, P. Aloy,
G. Liti and J. van Leeuwen (2021). "Natural variants suppress mutations in hundreds of essential genes."
Mol Syst Biol **17**(5): e10138.

Patra, B., Y. Kon, G. Yadav, A. W. Sevold, J. P. Frumkin, R. R. Vallabhajosyula, A. Hintze, B. Ostman, J.
Schossau, A. Bhan, B. Marzolf, J. K. Tamashiro, A. Kaur, N. S. Baliga, E. J. Grayhack, C. Adami, D. J.

Galas, A. Raval, E. M. Phizicky and A. Ray (2016). "A genome wide dosage suppressor network reveals genomic robustness." Nucleic Acids Res **45**(1): 255-270.

Peter, J., M. De Chiara, A. Friedrich, J. X. Yue, D. Pflieger, A. Bergstrom, A. Sigwalt, B. Barre, K. Freel, A. Llored, C. Cruaud, K. Labadie, J. M. Aury, B. Istace, K. Lebrigand, P. Barbry, S. Engelen, A. Lemainque, P. Wincker, G. Liti and J. Schacherer (2018). "Genome evolution across 1,011 Saccharomyces cerevisiae isolates." Nature **556**(7701): 339-344.

Sikorski, R. S. and P. Hieter (1989). "A system of shuttle vectors and yeast host strains designed for efficient manipulation of DNA in Saccharomyces cerevisiae." Genetics **122**(1): 19-27.

Song, H., T. L. Torng, A. S. Orr, A. T. Brunger and W. T. Wickner (2021). "Sec17/Sec18 can support membrane fusion without help from completion of SNARE zippering." Elife **10**.

Tholey, G., S. Bloch, M. Ledig, P. Mandel and F. Wedler (1987). "Chick brain glutamine synthetase and Mn2+-Mg2+ interactions." Neurochem Res **12**(11): 1041-1047.

Torres, E. M., T. Sokolsky, C. M. Tucker, L. Y. Chan, M. Boselli, M. J. Dunham and A. Amon (2007). "Effects of aneuploidy on cellular physiology and cell division in haploid yeast." Science **317**(5840): 916-924.

Turco, G., C. Chang, R. Y. Wang, G. Kim, E. H. Stoops, B. Richardson, V. Sochat, J. Rust, R. Oughtred, N. Thayer, F. Kang, M. S. Livstone, S. Heinicke, M. Schroeder, K. J. Dolinski, D. Botstein and A. Baryshnikova (2023). "Global analysis of the yeast knockout phenome." Sci Adv **9**(21): eadg5702.

Uda, M., R. Galanello, S. Sanna, G. Lettre, V. G. Sankaran, W. Chen, G. Usala, F. Busonero, A. Maschio, G. Albai, M. G. Piras, N. Sestu, S. Lai, M. Dei, A. Mulas, L. Crisponi, S. Naitza, I. Asunis, M. Deiana, R. Nagaraja, L. Perseu, S. Satta, M. D. Cipollina, C. Sollaino, P. Moi, J. N. Hirschhorn, S. H. Orkin, G. R. Abecasis, D. Schlessinger and A. Cao (2008). "Genome-wide association study shows BCL11A associated with persistent fetal hemoglobin and amelioration of the phenotype of beta-thalassemia." Proc Natl Acad Sci U S A **105**(5): 1620-1625.

Ünlü, B., C. Pons, U. L. Ho, A. Batté, P. Aloy and J. van Leeuwen (2023). "Global analysis of suppressor mutations that rescue human genetic defects." Genome Med **15**(1): 78.

Van der Vaart, J. M., L. H. Caro, J. W. Chapman, F. M. Klis and C. T. Verrips (1995). "Identification of three mannoproteins in the cell wall of Saccharomyces cerevisiae." J Bacteriol **177**(11): 3104-3110.

Van Leeuwen, J., C. Pons, C. Boone and B. J. Andrews (2017). "Mechanisms of suppression: The wiring of genetic resilience." Bioessays **39**(7): 1700042.

Van Leeuwen, J., C. Pons, J. C. Mellor, T. N. Yamaguchi, H. Friesen, J. Koschwanez, M. Mattiazzi Usaj, M. Pechlaner, M. Takar, M. Usaj, B. VanderSluis, K. Andrusiak, P. Bansal, A. Baryshnikova, C. E. Boone, J. Cao, A. Cote, M. Gebbia, G. Horecka, I. Horecka, E. Kuzmin, N. Legro, W. Liang, N. van Lieshout, M. McNee, B. J. San Luis, F. Shaeri, E. Shuteriqi, S. Sun, L. Yang, J. Y. Youn, M. Yuen, M. Costanzo, A. C. Gingras, P. Aloy, C. Oostenbrink, A. Murray, T. R. Graham, C. L. Myers, B. J. Andrews, F. P. Roth and C. Boone (2016). "Exploring genetic suppression interactions on a global scale." Science **354**(6312): 599.

Van Leeuwen, J., C. Pons, G. Tan, J. Z. Wang, J. Hou, J. Weile, M. Gebbia, W. Liang, E. Shuteriqi, Z. Li, M. Lopes, M. Usaj, A. Dos Santos Lopes, N. van Lieshout, C. L. Myers, F. P. Roth, P. Aloy, B. J. Andrews and C. Boone (2020). "Systematic analysis of bypass suppression of essential genes." Mol Syst Biol **16**(9): e9828.

Wang, Y., C. D. Arenas, D. M. Stoebel and T. F. Cooper (2013). "Genetic background affects epistatic interactions between two beneficial mutations." Biol Lett **9**(1): 20120328.

Wingett, S. W. and S. Andrews (2018). "FastQ Screen: A tool for multi-genome mapping and quality control." F1000Res **7**: 1338.

Wong, E. D., S. R. Miyasato, S. Aleksander, K. Karra, R. S. Nash, M. S. Skrzypek, S. Weng, S. R. Engel and J. M. Cherry (2023). "Saccharomyces genome database update: server architecture, pan-genome nomenclature, and external resources." Genetics **224**(1): iyac191.