Prefrontal working memory signal primarily controls phase-coded information within extrastriate cortex

Mohsen Parto-Dezfouli[1*@], Isabel Vanegas[2*], Mohammad Zarei[3], William H. Nesse[2], Kelsey L. Clark[2], Behrad Noudoost[2,4@]

1 Ernst Strüngmann Institute (ESI) for Neuroscience in Cooperation with Max Planck Society, 60528 Frankfurt, Germany
2 Department of Ophthalmology and Visual Sciences, John Moran Eye Center, University of Utah, Salt Lake City, UT, United States
3 Biosciences Institute, Newcastle University, Newcastle upon Tyne, UK
4 Lead

* Co-first author
@ Corresponding authors: mohsen.parto-dezfouli@esi-frankfurt.de, behrad.noudoost@utah.edu

Abstract

In order to understand how prefrontal cortex provides the benefits of working memory (WM) for visual processing we examined the influence of WM on the representation of visual signals in V4 neurons in two macaque monkeys. We found that WM induces strong β oscillations in V4 and that the timing of action potentials relative to this oscillation reflects sensory information- i.e., a phase coding of visual information. Pharmacologically inactivating the Frontal Eye Field part of prefrontal cortex, we confirmed the necessity of prefrontal signals for the WM-driven boost in phase coding of visual information. Indeed, changes in the average firing rate of V4 neurons could be accounted for by WM-induced oscillatory changes. We present a network model to describe how WM signals can recruit sensory areas primarily by inducing oscillations within these areas and discuss the implications of these findings for a sensory recruitment theory of WM through coherence.

Keywords: neural phase code, working memory, top-down control, prefrontal cortex, neural oscillations

Introduction

Our capacity to dynamically interact with the world around us based on our own needs, priorities and goals is an example of our cognitive flexibility. The goals and plans preserved by working memory (WM) are capable of altering our perceptions of and actions toward the world around us. Thus, understanding how our plans alter the representation of sensory information can reveal the neural basis of cognitive flexibility. Prefrontal cortex (PFC) is believed to be one of the sources controlling sensory signals according to goals and plans[1-6]. In this study, we are specifically examining how WM information sent from PFC can influence the representation of sensory information in visual areas in macaque monkeys. We have already shown that the Frontal Eye Field (FEF) part of PFC sends a direct WM signal to extrastriate area V4[7], and visual areas manifest the WM content in their oscillatory behavior[8]. Within the FEF, WM alters the efficacy of V4 inputs as well, and these inputs mostly target neurons believed to be involved in the transformation of visual information into motor action[9]. The finding that extrastriate visual areas receive the content of WM, and that information sent from these areas to prefrontal cortex is undergoing a visuomotor transformation, provide a clear picture of the sequence of events giving rise to the benefits of WM. However, exactly how this top-down WM signal enhances the representation in visual areas is not known, considering that neurons in extrastriate visual areas show little or no change in their firing rate in response to WM content[10-14]. Thus, how WM impacts visual representations is crucial for understanding the neural code, since the behavioral consequences of WM[15-21] suggest that the representation altered by WM content is likely to be the representation that our behavior relies on.

Knowing that the FEF sends a spatially-specific signal to V4 carrying the content of WM[7], we studied the responses of V4 neurons while top-down WM is directed to their receptive field (RF) or elsewhere, under the conditions in which FEF activity is intact or disrupted using pharmacological manipulation. V4 neurons were provided with a bottom-up visual input with the goal of understanding how a top-down WM signal can alter the representation of bottom-up information. This arrangement revealed that WM primarily enhances the phase coding of visual information in V4. Pharmacological inactivation demonstrated that FEF activity is necessary for this phase-dependent representational enhancement in V4. We also show that average firing rate modulations within visual areas can be accounted for by WM-induced oscillations within these areas, supporting the primacy of WM-dependent oscillations in generating the signatures of WM within sensory areas. These results corroborate a WM model in which sensory areas can be recruited by higher areas without needing to change the average firing rate of sensory neurons.

Results

WM modulates oscillatory power and spike timing in V4, but not average firing rates

In order to assess the impact of WM on the visual representation, we recorded local field potentials (LFPs) and neuronal activity in extrastriate area V4 during a spatial WM task with visual signals presented to neurons independent of the WM demand (see Methods; Fig. 1A). The animal had to remember a visual cue presented either inside the V4 RF or in the opposite hemifield (memory IN and OUT conditions, respectively), and following a delay period, made a saccade to the remembered location to receive a reward. The background stimulus could have one of four orientations and three levels of contrasts (or no contrast, the classic memory-guided saccade (MGS) task), allowing us to examine the interaction between bottom-up sensory information and top-down WM signals. We recorded 145 V4 neurons across 88 recording sessions; most of our analysis focuses on their responses during the last 700ms of the delay period of the WM task.

As expected, V4 neurons were capable of signaling the properties of the background stimulus present in their RF. The left panel in figure 1B shows the response of a sample V4 neuron during the IN condition. During the fixation period and the cue period, the neuron exhibited sensitivity to different contrast levels of the background stimulus projected to its RF (main effect of contrast: $F_{Fixation}=41.744$, $p<10^{-16}$; $F_{Cue}=14.541$, $p<10^{-6}$; one-way ANOVAs). The neuron's contrast sensitivity was still manifested in its firing rate during the delay period ($F_{IN}=19.950$, $p<10^{-8}$; One-way). As shown on the right side of figure 1B, this neuron also showed contrast sensitivity during the delay period of the OUT condition ($F_{OUT}= 17.851$, $p<10^{-7}$; One-way ANOVA). Importantly, the contrast sensitivity during the delay period was not significantly different between the IN vs. OUT conditions ($F_{condition}= 5\times10^{-4}$, $p=0.981$; $F_{contrast}=37.540$, $p<10^{-16}$; $F_{interaction}=0.301$, $p=0.742$; Two-way ANOVA). Similarly, the neuron was selective for the orientation of the background stimulus. The neuron exhibited orientation sensitivity during the fixation period as well as the delay period in both IN and OUT conditions (main effect of orientation: $F_{Fixation}=16.401$, $p<10^{-8}$; $F_{IN}=10.410$, $p<10^{-5}$; $F_{OUT}=19.014$, $p<10^{-10}$; One-way ANOVAs; Fig. 1C). As with contrast, the content of WM (IN vs. OUT condition) did not significantly change the neuron's orientation sensitivity during the delay period ($F_{Condition}= 5\times10^{-4}$, $p=0.982$; $F_{Orientation}=28.082$, $p<10^{-16}$; $F_{Interaction}=0.810$, $p=0.492$; Two-way ANOVA). Thus, this sample neuron exhibited contrast and orientation sensitivity for the background stimulus; however, WM did not alter the stimulus information reflected in the neuron's firing rate.

The same pattern was observed in the population of 145 V4 neurons: neurons' firing rates reflected

the properties of the background stimulus, but this capacity was not affected by WM. The response to

different background stimuli was similar between the fixation period, delay IN, and delay OUT conditions,

both for contrast ($F_{Contrast}$=16.923, p<10^{-7}; $F_{Condition}$=0.311, p=0.733; $F_{Interaction}$=0.400, p=0.811; Two-way

ANOVA) and orientation ($F_{Orientation}$=23.294, p<10^{-14}; $F_{Condition}$=0.770, p=0.465; $F_{Interaction}$=0.202, p=0.976;

Two-way ANOVA) (Fig. 1D, 1E; see also Fig. S1A-B). Figure 1F shows the ability of V4 neurons to discriminate

between all 12 stimuli of various contrasts and orientations across time. Overall, the ability of V4 neurons

to discriminate between various background stimuli was not altered during the delay period of the IN vs.

OUT conditions (discriminability$_{IN}$=1.514±0.783, discriminability$_{OUT}$=1.482±0.680, p=0.462; Fig. 1G; see also

Fig. S1C-D). Therefore, whereas sensory information is reflected in the firing rate of V4 neurons, the impact

of spatial WM on the sensory representation is not traceable by this neural signature.

In contrast to the lack of WM-driven change in firing rates, we found that V4 LFP oscillations strongly

reflect the content of WM. Figure 1H shows the LFP power spectrum during the delay period for the same

channel of recording as of the example neuron shown in Figure 1B-C. The β (14-22 Hz; see Methods) band

power was 0.019±0.013 dB/Hz during the IN condition, which is 18.9% greater than the 0.016±0.012 dB/Hz

during the OUT condition (Wilcoxon ranksum, p=0.002). Therefore, although neurons' firing rates do not

change due to WM, the power of the LFP oscillation in the β range reflects the impact of WM in the same

recording channel. This phenomenon was observed across the 88 LFP recordings (Fig. 1I): β LFP power

during the delay was greater for the IN condition compared to the OUT condition (Power$_{IN}$=2.001±0.336,

Power$_{OUT}$=1.842±0.333, p<10^{-3}; Fig. 1J). Importantly, this WM-dependent enhancement of β power was

observed independent of the background stimuli (Fig. S2). In summary, a visual area known to receive the

spatial WM signal exhibits the signature of this top-down signal in its subthreshold LFP activity, but not in

neurons' firing rates.

The LFP reflects a combination of nearby currents, including synaptic inputs[22-25], and is not directly

transmitted along axons to other brain areas in the manner that spikes are. Therefore, we sought to identify

whether these WM-dependent oscillatory changes impact any other aspects of spiking activity within V4.

Although at the coarse scale of average firing rate there was no change due to WM, at a higher temporal

resolution it became evident that the timing of V4 action potentials depended on the phase of WM-induced

β oscillations. Figure 2A shows the distribution of spikes generated by a sample V4 neuron across various

phases of the β oscillation during the delay period, for memory IN and OUT conditions. The average delay

period FR was not different between the two memory conditions ($F_{Condition}$= 2.050, p=0.153; $F_{Contrast}$=11.511,

p<10^{-4}; $F_{Interaction}$=1.314, p=0.271; Two-way ANOVA), but the phase distribution of spikes during the delay

period of the IN condition was more concentrated (centered around 160-degree phase) compared to the

OUT condition. To quantify the phenomenon, we used the spike-phase locking (SPL) as a measure of how

consistently spikes of a neuron are generated at a certain phase. The SPL index varies between 0 (spikes

homogenously distributed across phases) to 1 (all spikes occurring at a certain phase). For the sample

neuron shown in Figure 2A, SPL changed from 0.063 for the OUT to 0.163 for the IN condition. Across the

population of 145 V4 neurons, we found a consistent impact of WM on SPL: SPL relative to the β oscillation

was significantly greater during the IN condition compared to the OUT condition (n=145 neurons;

SPL_{IN}=0.164±0.014, SPL_{OUT}=0.155±0.005, p<10^{-3}; Fig. 2B). These results indicate that although WM does not

change the average firing rate, it influences V4 spike timing to be more closely aligned with WM-dependent

oscillations. Conversely, evaluating the consistency of the LFP at the time of V4 spikes, the spike triggered

average (STA) LFP also shows this strong coupling of spikes to LFP phase in the β range (Fig. S3).

137 Spatial WM specifically enhances phase coding of visual information

To understand how WM-induced oscillations benefit the sensory representation within V4, we

examined whether V4 neurons' sensitivity was reflected in the timing of their spikes relative to these WM-

induced oscillations. As shown in figure 2C, analyzing the average normalized LFP at the time of spiking (the

STA) for an example neuron, we observed that in the presence of the high contrast stimulus in the

background during the memory IN condition, V4 spikes were associated with a steeper LFP change

compared to when a low contrast stimulus was in the background ($Slope_{High}$=0.052, $Slope_{Low}$=0.029). For

this neuron, this difference was less prominent during the OUT condition ($Slope_{High}$=0.042, $Slope_{Low}$=0.035).

We observed a similar phenomenon across the population. The LFP slope around the time of a spike was

significantly sharper when the preferred (high contrast) stimulus was presented in the background,

compared to the nonpreferred (low contrast) stimulus, during the IN condition ($Slope_{High}$=0.046±0.003,

$Slope_{Low}$=0.021±0.002, p<10^{-8}; Fig. 2D left). This difference was not observed during the OUT condition

($Slope_{High}$=0.030±0.002, $Slope_{Low}$=0.039±0.004, p=0.110, Fig. 2D right). This indicates that WM-induced

oscillations can facilitate the spiking activity in visual areas, consistent with what other groups have shown

regarding the role of oscillations as a boost for passing the spiking threshold [26]. Similar results were

observed for preferred versus nonpreferred orientations (Fig. S4).

The idea that a WM-induced oscillation can change the timing of spikes also suggests the possibility

that the timing of spikes relative to that oscillation could convey information, referred to as a neural phase

code. We used the mutual information (MI) to quantify visual information conveyed by either the phase or

rate of spikes to have a side-by-side comparison of a neural phase vs. rate code (see Methods). Figure 2E

left shows the average population MI measured based on phase coding across time for both the IN and OUT

conditions. The presence of the WM cue reduced both phase and rate coding of background information

during the visual period (Phase MI_{IN}=0.007±0.001 bits, Phase MI_{OUT}=0.014±0.002 bits; p=0.012; Rate

MI_{IN}=0.019±0.002 bits, Rate MI_{OUT}=0.031±0.003 bits; p=0.012). However, maintenance of WM information

during the delay period increased the phase coding capacity of the V4 neurons to represent information

about the stimulus in their RF, but did not alter their rate coding. Figure 2F shows the capacity of V4 neurons

to encode the background stimulus during the IN vs. OUT conditions under phase (left) and rate (right)

coding schemes. Phase MI during the delay period of the IN condition was 0.019±0.002 bits, significantly

greater than the 0.012±0.002 bits during the OUT condition ($p<10^{-3}$). However, delay period rate MI was

not significantly different between the IN and OUT conditions (MI_{IN}=0.024±0.003 bits; MI_{OUT}=0.022±0.003

bits, p=0.261). We also found that the WM-dependent enhancement of phase-dependent visual

representation was limited to the β range oscillations (Fig. 2G) It is imperative to note that despite the

seemingly small values of MI (e.g., 0.019 bits), an increase of 54% in phase MI between the IN and OUT

conditions means a huge boost in coding capacity due to WM. The MI value can be interpreted as a measure

of the rate of statistical learning from incoming data through which sensory decisions can be made. For

example, with 0.012 bits per 100 ms phase MI available in the OUT condition, for a population of 100

neurons firing independently it would take 294 ms to fully differentiate 12 stimuli. WM-induced

enhancement of phase MI means that the same discriminatory capacity can be achieved within 190 ms with

the same number of neurons, or within the same amount of time but with only 65 neurons. Altogether,

consistent with the finding that WM mainly modulated oscillations within the β range (Fig. 1I, J), we found

that WM mostly improves the phase coding in V4 within the same β range. Thus, WM specifically enhanced

β range phase coding in V4, without altering rate coding.

FEF activity is necessary for the phase-dependent representation within V4 during WM.

The FEF sends direct projections to V4 carrying the content of spatial WM[7]. To causally test whether

the observed WM-driven phase coding in V4 depended on signals received from the FEF, we recorded from

V4 neurons before and after pharmacologically inactivating a portion of the FEF using a small-volume

injection of the GABAa-agonist muscimol (Fig. 3A). Localized FEF inactivation is known to impair

performance on the MGS task in a spatially-specific manner[27,28]. As shown for an example inactivation

session in figure 3B: prior to inactivation, the animal performed well at all locations, and after FEF

inactivation performance was disrupted for conditions in which the cue appeared in the left hemifield,

contralateral to the inactivated FEF. Figure 3C shows average MGS performance over time at various

locations across 33 inactivation sessions: performance for the IN condition and neighboring locations

decreased over time following FEF inactivation. For the IN condition, the performance dropped from

90.32 ± 2.62 percent correct before to 68.49 ± 5.39 percent correct three hours after inactivation ($p<10^{-3}$).

Across the same 33 sessions, saccade error during the IN condition compared to the OUT condition was not

different prior to FEF inactivation ($Scatter_{IN}=1.012 \pm 0.022$ dva, $Scatter_{OUT}=0.989 \pm 0.023$ dva, p=0.520; Fig.

3D), but was significantly greater for the IN condition after FEF inactivation ($Scatter_{IN}=1.192 \pm 0.042$ dva,

$Scatter_{OUT}=1.033 \pm 0.032$ dva, p=0.001; Fig. 3D). Similarly, reaction time (RT) increased following FEF

inactivation for the IN condition compared to the OUT condition (before inactivation: $RT_{IN}=1.015 \pm 0.003$,

$RT_{OUT}=1.023 \pm 0.007$, p=0.741; after inactivation: $RT_{IN}=1.075 \pm 0.015$, $RT_{OUT}=1.014 \pm 0.006$, $p<10^{-4}$, Fig. 3E).

We recorded from 66 V4 neurons before and after FEF inactivation. Both the V4 LFP power spectrum

and SPL showed a reduction in the β range following FEF inactivation (Fig. S5). In these 66 neurons, prior to

inactivation, the impact of WM on phase coding was evident: during the delay period there was significantly

stronger phase coding of information in the β range for the memory IN condition (Fig. 2E-G; Phase

$MI_{IN}=0.023 \pm 0.004$ bits, Phase $MI_{OUT}=0.018 \pm 0.003$ bits, p=0.039). Consistent with figure 2E, WM did not alter

the strength of rate coding (Rate $MI_{IN}=0.022 \pm 0.005$ bits, Rate $MI_{OUT}=0.020 \pm 0.005$ bits, p=0.446).

Importantly, the phase coded information within the β range during the delay period of the task dropped

following FEF inactivation (Fig. 3F). Figure 3G shows the cross section of figure 3F at the β frequency,

depicting the dynamics of phase MI over the course of a trial before and after FEF inactivation. The MI

values for individual neurons during the delay period of the IN condition for each session are shown in figure

3H; following FEF inactivation, phase-coded MI in the β range dropped from 0.023 ± 0.004 bits to

0.014 ± 0.003 bits (n= 66 neurons, p=0.005; Fig. 3H). Rate coding, in contrast, was unaffected by FEF

inactivation (IN condition: Rate $MI_{Pre}=0.022 \pm 0.005$ bits, Rate $MI_{Post}0.020 \pm 0.004$ bits, p=0.880). Thus, WM's

enhancement of phase coding in V4 depended on activity within the FEF.

Primacy of a phase code

The finding that WM mainly modulates phase coded information within extrastriate areas fundamentally shifts our understanding of how the top-down influence of prefrontal cortex shapes the neural representation, suggesting that inducing oscillations is the main way WM recruits sensory areas. However, while this side-by-side comparison of rate and phase coding shows the strength of the latter, several studies have reported an impact of WM on the firing rate of visual neurons[7,14,29-31]. One can argue that a slight increase in firing rate at each stage of visual processing can gradually accumulate to eventually emerge in the form of a robust firing rate change[11], and that this will be sufficient to support WM. In order to determine the primary means by which WM alters neural representations, we constructed a neural field network model of visual areas during WM. In order to examine the impact of WM on oscillatory and firing rate changes in visual areas, we designed the model to consist of interconnected excitatory and inhibitory units (Fig. 4A, e-cells and i-cells) capable of generating oscillatory activity. To modulate this oscillatory activity, these neural field units received bottom-up and top-down type input. The units were tuned to different stimulus 'features' of the bottom-up input (analogous to orientation tuning of V4 neurons in the experimental data, with input strength analogous to contrast). The top-down input was not feature selective, providing a uniform input across the network, with stronger connections to e-cells than i-cells, consistent with what is known about the FEF-V4 circuitry and anatomy[7,32]. A higher strength of WM signal in the model corresponds to the memory IN condition, in comparison to the absence of WM input in the memory OUT equivalent. The model replicates several key features of the experimental data: units reflect sensory information in their phase and rate, WM-enhanced β power, and locking of units' activity to this oscillation under the influence of WM (Fig. S6). The model's phase coding of visual stimuli is evident in the relative timing of responses of differently tuned e-cells to an input stimulus (Fig. 4B). Using this model, we can directly compare the magnitude of information encoded by the phase and rate of model units, quantified via information gain (see Methods). We found that not only was information encoded by phase much greater than that encoded by rate, but also that phase and rate information were oppositely affected by changes in WM strength: phase information increased and rate information decreased as the WM signal increased (Fig. 4C). We also found this same divergent pattern between phase and rate codes when measuring coding performance using mutual information (see Fig. S7). Therefore, the quantification of information within a tangible network model revealed that in an oscillating network, a top-down induced oscillation can be detrimental to the rate-dependent representation of information.

The model revealed that stronger WM input increased both oscillation strength and peak frequency
(Fig. S6), and we hypothesized that this change in frequency could explain changes in firing rate. We tested
this idea both in the model and in the experimental data. In the model, we varied the strength of the WM
signal across 4 levels of stimulus input strength (resembling various levels of stimulus contrast). We then
divided oscillatory cycles occurring in the respective stimulus input levels into deciles based on their
oscillation frequency, and measured information encoded by phase and rate as a function of oscillation
frequency (Fig. 4D). Phase information increased with increasing oscillation frequency while rate
information decreased at higher oscillation frequencies, for all non-zero contrast levels (Model utility t-test
for contrasts 0-3 respectively; phase code: $p>0.05$, >0.05, $<0.01,0.001$; and rate code: $p>0.05$,
$<0.05,0.01,0.0001$). Importantly, we found that the network replicating the oscillatory and representational
characteristics of V4 during WM shows an increased firing rate as oscillation frequency increases. Figure 4E
shows how small variations in the strength of the WM signal resulted in correlated changes in oscillation
frequency and firing rate ($r = 0.901$; linear model utility t-test significant: $p<0.0001$). Thus, firing rate is
positively correlated with oscillation frequency, but information encoded by that rate is negatively
correlated (Fig. 4C vs. 4E). To confirm that such a relationship exists in the experimental data, for each
neuron we measured the peak LFP frequency and average firing rate across subsamples of trials, allowing
us to test the relationship between peak β frequency and evoked firing rate within a single condition. As
shown for two sample V4 neurons, such a relationship between peak frequency and average firing rate
existed between the IN and OUT conditions (for a single background stimulus; Fig. 4F & G). For the first
sample neuron, in which the peak frequency changed from 18.81 to 15.84 Hz between the IN and OUT
conditions, the firing rate changed from 1.084 to 0.869, and peak frequency and firing rate were correlated
across subsamples of trials (Pearson correlation, $r=0.716$, $p<10^{-32}$; Fig. 4F). Results from the second sample
neuron show that this relationship remains the same even in cases where WM reduces the peak frequency
(peak-frequency$_{IN}$ = 15.041 Hz, peak-frequency$_{OUT}$ = 16.827 Hz, $FR_{IN}=0.630$, $FR_{OUT}=0.816$): the correlation
between firing rate and peak frequency remains positive (Pearson correlation, $r=0.671$, $p<10^{-26}$; Fig. 4G). At
the population level, we looked at firing rate during the delay period across subsamples of trials for all 145
V4 neurons, sorted according to their peak frequency (Fig. 4H). As shown in Figure 4H left, this relationship
between peak frequency and firing rate was present across both IN and OUT conditions; more importantly,
this relationship remained the same between the two memory conditions ($F_{Condition}=9.649$, $p=0.003$;
$F_{Frequency}=399.566$, $p<10^{-31}$; $F_{Interaction}=2.935$, $p=0.091$, ANCOVA), suggesting that the frequency of WM-
induced oscillations can account for firing rate changes. Critically, a similar analysis for the visual period of

the task revealed that the presence or absence of a visual stimulus (IN vs. OUT condition) creates a much

larger change in firing rate, which cannot be accounted for by changes in β frequency ($F_{Condition}=5.113 \times 10^3$,

$p<10^{-72}$; $F_{Frequency}=103.220$, $p<10^{-15}$; $F_{Interaction}=71.427$, $p<10^{-11}$, ANCOVA) (Fig. 4H right). In other words,

changes in rate during the delay period may be a consequence of changes in phase locking frequency. We

also examined the relationship between peak frequency and firing rate as a function of stimulus efficacy.

Both firing rate and peak frequency vary with stimulus efficacy (Fig. S8A; two sample neurons), but there

was no difference for this relationship between the IN and OUT condition (Fig. S8B). Figure S9 shows that

this relationship between firing rate and peak frequency is specific to the β frequency range. While there

was no overall change in average firing rate due to WM across the V4 population (see Fig. 1, Fig. S1, and

related statistics), this analysis further suggests that any WM-related changes in the firing rates of individual

V4 neurons could be primarily driven by changes in oscillatory frequency.

Discussion

Prefrontal cortex modulates sensory and motor signals in order to guide our actions based on goals and

priorities maintained in WM[3,4,33,34]. Within prefrontal areas, FEF sends direct projections to extrastriate

visual areas carrying the content of WM[7]. We designed a paradigm in which neurons in extrastriate area V4

are provided with bottom-up input while the top-down signal carrying WM content can be directed to the

part of space represented by these neurons or elsewhere. This allowed us to examine which aspect of the

sensory representation within V4 is influenced by a top-down WM signal, and to causally test the role of

FEF activity in this WM-driven modulation. We found that a neural phase code representation of sensory

stimuli was strongly modulated by top-down WM signals coming from the FEF, while firing rates were

relatively unaffected, leading us to conclude that representations based on the average firing rate of

neurons are not the primary way that top-down signals enhance sensory processing. Using a combination

of computational modelling and experimental data analysis, we provided evidence that any changes in the

average firing rate of individual neurons might be a byproduct of small changes in the frequency of the WM-

induced oscillation.

The long[35] and growing[36-38] list of neural signatures of attention begs a unifying theory describing

the exact mechanisms involved in generating this plethora of neural signatures. Many of these signatures

are seen in both attention and WM, including enhanced visual responses[7,39], changes in inter-neuronal

correlations[40,41], decreased variability[7,42], and shifts in RFs[7,43,44]. In light of the present findings, we suggest

that by inducing an oscillation, top-down signals allow expression of sensory representations in the form of

a neural phase code: neurons emit action potentials in response to this induced oscillation with a relative

timing that reflects their sensitivity. Slight changes in the frequency of the oscillation might then account

for changes in the average firing rate of neurons in sensory areas, and one can imagine coherent oscillations

altering the dependent and independent variability of the neurons as well. Understanding whether the β

oscillations observed in our study function in the same way as the mostly gamma oscillations reported in

attention studies will require a more complete understanding of the characteristics of oscillators operating

in the presence and absence of visual information (see also [40]).

A framework in which top-down signals primarily alter the phase of the spikes faces an important

challenge: in communication between brain areas, LFP oscillations are not carried along with spikes down

axonal projections. A phase code without its oscillatory reference frame is likely unreadable. However,

studies in our lab and others have provided growing evidence that there are coherent oscillations between

brain areas during WM, which could provide the shared oscillatory frame of reference required to transfer

phase-coded information. For example, oscillatory coherence between FEF and inferotemporal cortex

exists and predicts performance on an object WM task[45]; similarly, synchrony between PFC and V4 is also

correlated with WM performance[46]. Oscillatory coherence between prefrontal and parietal areas also

reflects the content of WM[47-49]. For a more complete review of findings of inter-areal coherence during

WM and their relationship to performance, see [50]. The significance of enhancing the efficacy of signals by

generating a coherent signal [51] has previously been presented in the context of communication through

coherence (CTC)[52-55], as has the idea that phase in the receiving area can influence sensitivity to incoming

signals[56,57]. In addition to gating of efficacy by phase of the receiving area (as in CTC, where this gating can

make a downstream area more sensitive to input from one source than another[58]), the precise timing of

spikes relative to oscillations even within a coherently oscillating site (i.e. phase coding) could also be crucial

when this timing is going to be gated back into signal strength using a coherent oscillation in the receiving

area.

We found that WM signals allow expression of visual representations in the form of a neural phase

code, indicating that prefrontal cortex can recruit sensory areas using a WM-induced oscillation. This new

finding, along with the abundant evidence of coherent oscillations across brain areas during WM[50], lead to

a working hypothesis about how WM can recruit sensory areas. Consistent with sensory recruitment

theories of WM[33,59-61], these results suggest that sensory and memory signals can be preserved in sensory

areas without being expressed in their average firing rate. This latent information is expressed in the form

of a phase code in response to a WM-induced oscillation, and is potentially readable by other areas that

have oscillations coherent with the oscillatory frame of reference induced by WM. This proposed

recruitment through coherence framework of working memory[62] offers an explanation for how WM can

recruit highly feature-sensitive sensory areas in the absence of robust firing rate changes within them.

Materials and methods

Experimental model details

We recorded from two male rhesus monkeys (Macaca mulatta, 12 and 16Kg). All experiments and animal procedures in this study were in accordance with the National Institutes of Health Guide for the Care and Use of Laboratory Animals and the Society for Neuroscience Guidelines and Policies. Protocols for experimental and behavioral procedures were approved by the University of Utah Institutional Animal Care and Use Committee.

General and surgical procedures

All surgeries were performed under aseptic conditions, using standard techniques and gas anesthesia, with appropriate peri-surgical analgesia and monitoring. After the study's conclusion, both animals remained healthy and were subsequently utilized in other research endeavors. Stereotactic surgery coordinates for the PFC and V4 chambers (20mm diameter) were performed for monkey 1, right hemisphere, at (AP 25+(2), ML 15 (±0)) and (AP -5(-1), ML 20+(2)), and for monkey 2, left hemisphere, at (AP 30+(1-2), ML 15-(1-2)) and (AP -5-(1-2), ML 20+(1-2)).

Behavioral tasks

We programmed all behavioral tasks using the NIMH Monkeylogic toolbox (ML2) 55, on 64-bit Matlab software (The MathWorks, Inc., Natick, MA). We monitored eye position with an infrared optical eye-tracking (EyeLink 1000, SR Research, Ottawa, Canada). Visual tasks were presented on a VG248 ASUS LED monitor with a refresh rate of 144 Hz and resolution of 1920 x 1080 pixels.

V4 RF mapping

On a daily basis, we first identified V4 RFs using audible responses to oriented bars. Second, we presented a series of visual stimuli on a black background, to quantitatively estimate V4 RFs based on the neuron's firing rate response. Visual stimuli were white circles (1dva diameter), 100ms on, 100ms off, pseudorandomly presented in a 7x7 grid spaced 2.5 dva between stimuli. The monkeys fixated on a central white circle (1dva diameter) throughout the trial.

FEF RF mapping

We estimated FEF RFs using electrical stimulation within the anterior bank of the arcuate sulcus, in biphasic microcurrent pulses (50µA) using a S88 Grass stimulator. Stimulation was performed via tungsten microelectrodes (FHC, Bowdoin, ME). FEF sites were identified based on the landing point of the evoked eye movement following stimulation with currents ≤50µA.

Memory guided saccade tasks

To assess the influence of WM on the representation of sensory stimuli, we used a variant of MGS task with a background stimulus (Fig. 1A). The MGS-background task is similar to classic MGS task with a task-irrelevant full field stimulus in the background. The background stimulus was an oriented grating, which could appear in one of four orientations and 4 contrasts (0% contrast is just a classic MG task). The WM cue was placed either within the overlapping RF of FEF and V4 (IN condition) or 180 degrees away (OUT condition). During FEF inactivation experiments, a classic 8 location MGS task with no background was used to assess the behavioral consequences of drug injection over space and time (Fig. 3B-E).

Neurophysiological recording

We recorded the activity of 145 V4 neurons across 88 recording sessions (55 sessions Monkey E, 33 sessions Monkey O), including 66 V4 neurons during 33 FEF inactivation sessions (29 sessions Monkey E, 4 sessions Monkey O). We recorded neurophysiological activity using Neuralynx and Blackrock data acquisition systems. We digitized spike waveforms at 32 KHz, and performed offline spike sorting manually. We used single tungsten microelectrodes of 200µm diameter, with epoxylite insulation (FHC, Bowdoin, ME), and linear 16-channel arrays (Plexon, Dallas, TX). Electrodes were inserted using a hydraulic microdrive (Narishige, Japan).

V4 recordings: We simultaneously recorded from FEF (single electrode) and V4 (single or linear array electrodes). In this paper we only present the data from the V4 recordings.

FEF inactivation with V4 recording: FEF was pharmacologically inactivated through infusion of 0.5-1µL of the GABA-a agonist muscimol, using a custom microinjectrode system (described in [63,64]). Muscimol concentration was 5mg/ml (pH 6.5 to 7). V4 activity was recorded from a site with RFs overlapping the estimated FEF RF, before and after FEF inactivation. Performance on the memory guided saccade task was used to verify FEF inactivation.

Data analysis

Quantification and statistical analysis

For all analyses of V4 responses (main Figs. 1,2,4), we pooled V4 data from simultaneous FEF-V4 recording sessions with data from FEF inactivation sessions (using the V4 data prior to FEF inactivation). For the results of Figure 3, we assess the role of FEF on V4 coding using inactivation data in which we recorded from V4 before and after FEF inactivation. Evaluations of neural responses to background stimuli of varying contrast or orientation include the three non-zero contrast values. Wherever a statistical test is not specified it is Wilcoxon sign rank. P values are reported up to three decimal digits, and p values less than 0.001 are reported as $p<10^{-x}$. The β range here is 14-22 Hz.

LFP power spectrum, spike-phase locking, and STA

The power spectral density of the local field potentials (LFPs) was calculated using the multitaper method, employing three tapers (discrete prolate spheroidal (DPSS)-Slepian sequences) for each trial and channel. For population LFP power statistics, LFP power spectrums were normalized, [(X-min)/(max-min)]. In sessions with array recordings (51/88 sessions) power calculations were performed for each channel and then averaged across all channels in that session before calculating population statistics. To quantify the reliability of spike timing relative to the LFP of the same channel, we employed the Spike-phase locking (SPL) method [65], which measures the consistency or locking strength of spike phases to the LFPs. This is achieved by calculating the angular summation between phases of LFPs and spike times. The amplitude of the SPL indicates the strength of spike locking to LFP phase, while the angle reflects the phases of LFPs when spikes occurred. For the spike-triggered average (STA) of the LFP, we first normalized the LFP by taking the z-score of the LFP across timepoints within 100ms of a spike for each trial, then averaged those values across trials.

Rate and phase coding capacity

Our primary means of measuring coding capacity was the method developed by Panzeri and colleagues, which allowed us to quantify and compare information contained in rate and phase codes [66,67]. This calculation of coding capacity was done in four steps. 1) First, using the FIR filter, LFPs were filtered into ten different frequency bands (1-4; 4-8; 8-12; 12-17; 17-22; 22-27; 27-35; 35-55; 65-90; 90-120). 2) Next, based

on Hilbert transform, the phase of filtered LFPs were extracted. 3) Subsequently, the average of phases at

the time of spike occurrence were estimated for a window of 100ms duration with 100ms shift. 4) Finally,

mutual information (MI) was calculated between these average phases (phase code) or average spike rate

(rate code) and different stimuli. The MI was calculated across all stimulus contrasts and orientations. For

full mathematical details see [66]. The configuration we used was: direct method, biased naive estimates and

20 bootstraps [67].

Mathematical modelling methods

Details of this model were previously published in *Frontiers in Computational Neuroscience* [68].

Neural field model

Our neural field model is defined by be a periodic orientation tuning domain parameter $\theta \in [0, \pi)$.

This neural field model is intended to represent a hypercolumn-like population with a subset of cells within

the neural field preferentially responsive to a θ-oriented stimulus. This model has been studied in detail in

a previous article by our group [68]. The neural field model is described by $u(\theta, t), v(\theta, t)$, the e- and i-activity

for every θ-location on the ring, that solves the integro-differential equation:

$$\tau_e \frac{\partial u}{\partial t} = -u + f_\sigma(W * [w_{ee}u - w_{ei}v] + I_e),$$

$$\tau_i \frac{\partial v}{\partial t} = -v + f_\sigma(W * [w_{ie}u - w_{ii}v] + I_i),$$

446   The integral convolution "*" in the above, is over the θ-domain, with $W(\theta)$ being the von-Mises periodic

447   weight kernel

$$W(\theta) = \frac{1}{\pi I_0(\kappa)} exp(\kappa cos(2\theta)).$$

A mass-model at θ and θ' will be connected with weight $W(\theta - \theta')d\theta'$. The parameter κ is the inverse-

variance-like scale parameter that shapes the broadness/tightness of the distribution, and $I\_0(\kappa)$ is the

order-zero modified Bessel function of the first kind, which serves as the normalization constant. Note that

the half-circle orientation tuning domain $\theta \in [0, \pi)$ necessitates a "2" factor in the weight function to be

π-periodic. The same spatial scale κ is used for both e- and i-cell populations.

The $f_\sigma(I)$ function defines the output firing rate of each population as a function of its input I---an

F-I curve. We have used a sigmoidal-shaped F-I curve defined as the inverse mean first passage time, plus a

5ms refractory period, of a leaky integrate an fire LIF model neuron driven by uncorrelated Gaussian white

noise $\sigma\xi(t)$ with standard deviation σ (see for example [68,69]:

458
$$C\frac{dV}{dt} = g_l(V_l - V) + I + \sigma\xi(t),$$

with capacitance $C = 1$ micro-Farads, spike threshold voltage $V_t = -50$mV, and reset and leak voltages

$V_r = V_l = -65$mV (all parameters are listed in Table S1). With these parameters, $I = 1$nA of current

induces the membrane voltage to approach spike threshold in the absence of noise. Increasing noise

parameter σ has the effect of reducing the overall gain of the F-I curve (see [68], for more details on this

model).

Three types of external inputs were given to the neural field: working memory inputs, stimulus

inputs, and random inputs. WM inputs are uniform current inputs, added to I_e and I_i, over the entire neural

field (equal for all θ-values). These uniform inputs raised very slightly the mean firing rate and oscillation

frequency and represent an WM-like or attentional-like enhancement of hypercolumn activity. Stimulus

inputs are orientation-tuned given by Von Mises-like distribution functions

469
$$S(\theta) = e^{\kappa_s \cos(2(\theta - \theta_0)) - \kappa_s}$$

in which the peak strength (set to unity) of the stimulus located at orientation θ_0. We fix $\theta_0 = \pi/2$--a 90-

degree (vertical) orientated stimulus, without loss of generality. Finally, to capture the temporal variations

in network oscillations observed in real cortical tissues, on simulations in which we assessed sensory coding

(see below), we included slow-timescale Ornstein-Ulenbeck noise $y(t)$ to both e- and i-cell input currents

I_e and I_i globally to the entire network (uniformly across all θ-values). The dynamics of y are given by the

stochastic differential equation

476
$$\tau_z \frac{dy}{dt} = -y + \sigma_z \sqrt{\tau_z}\,\xi(t),$$

where $\xi(t)$ is uncorrelated zero-mean unit-variance gaussian white noise. This equation results in a normal

stationary distribution of y-values, with zero-mean, and standard deviation σ_z, and a temporal

autocorrelation decay timescale $\tau_z = 50$ms, so that the network oscillations, which were typically in the

20 Hz range (50 ms oscillation cycles), showed robust cycle to cycle variability but little long-timescale multi-

cycle correlation.

In the absence of any external input, we set neural field model to be very near the a supercritical

Hopf instability (see [68]) in which additional current above a current threshold I^*, elicited oscillations with

amplitude emerging continuously from zero, and oscillation frequency in the β-band around 18-20 Hz. From

this I^* parameter starting point we ran simulations from over four levels of WM input (uniform current)

and four levels of orientation-selective stimulus input (contrast levels), starting from zero. We call these

WM 0,1,2,3 levels, and contrast levels 0,1,2,3. Altogether, the input to cells can be represented by

488
$$I_e(\theta, t) = I_{e0} + a\,\Delta_{stim}S(\theta) + b\,\Delta_{WM} + y(t),$$

489
$$I_i(\theta, t) = I_{i0} + a\,\Delta_{stim}S(\theta) + b\,\Delta_{WM} + y(t),$$

490 where Δ_{stim} and Δ_{WM} are the current increments for the respective input levels of stimulus $a = 0,1,2,3$

491 and WM $b = 0,1,2,3$. In addition to the input current changes that occur for our model, it is common to

492 accept that increased stimulus input comes with increased input current fluctuations. We modeled this by

493 adjusting the σ -parameter the F-I curve as a function of contrast input:

494
$$\sigma = \sigma_0 + a\Delta_\sigma.$$

Coding performance

The phase- and rate-based responses of the neural field model can be used to discriminate the input

stimuli. We have chosen to discriminate the neural field model responses at the neural field locations $\theta =$

$\pi/2$ and $\pi/4$. We computed the coding performance for two competing codes: a rate code (i.e., a spike

count code) and a phase code. To define the phase variable in the phase code, we examined the proxy LFP

signal formed by averaging e-cell rate responses over the entire field domain. We derived a phase angle

$\varphi(t) \in [-\pi, \pi]$ of oscillation via the Hilbert transform of this LFP signal. After segmenting the simulation

run time into oscillation cycles $\varphi(t) \in [-\pi, \pi]$, for $t \in [0, T]$, where $T = 1/f$ is oscillation period. The mean

rate response is simply $\lambda_\theta = \frac{1}{T}\int_0^T u(\theta, t)dt$, from which we assume Poisson-distributed n number of spikes

are emitted:

505
$$p_\theta(n) = \frac{(\lambda_\theta T)^n e^{-\lambda_\theta T}}{n!},$$

506 which constituted the rate code distribution.

507 The phase code distribution is obtained from the rate response $u(\theta, t)$, by using a change-of-

508 variables between time and phase $t = g(\varphi)$, where $g(\varphi)$ is the inverse of the Hilbert phase angle:

509
$$q_\theta(\varphi) = \frac{1}{\lambda_\theta T}\, u\big(\theta, g(\varphi)\big)g'(\varphi).$$

510 Using the spike count and phase distributions as the basis of the rate and phase codes, respectively,

511 we computed two different measures of coding performance. First, to measure the amount of information

512 gained from the code at $\theta = \pi/2$, given one assumes the data are distributed according to $\theta = \pi/4$, we

513 computed the information gain rate (IG)--the Kullback Libler divergence (in natural units of information,

514 nats):

515
$$IG_\varphi = D_{KL}\left(q_{\frac{\pi}{4}} \,||\, q_{\frac{\pi}{2}}\right),$$

516
$$IG_n = D_{KL}\left(p_{\frac{\pi}{4}} \,||\, p_{\frac{\pi}{2}}\right),$$

517 where $D_{KL}(p||q) = \int p\, ln\, ln\left(\frac{p}{q}\right)dx$. Second, we computed the mutual information (MI) between

518 stimulus feature $\theta = \pi/2$ or $\pi/4$, and the phase data φ. We assumed the two stimuli were equally likely

519   on a given ``trial" in which case the probability of each stimulus was 1/2. The phase mutual information

520   MI_φ is then defined as the Kullback Libler divergence D_{KL} (using log-base-two, in this case) between the

521   pooled distributions $\frac{1}{2}q_{\pi/2} + \frac{1}{2}q_{\pi/4}$ to the product distribution $q_{\pi/2}q_{\pi/4}$; and similarly, for the rate codes:

$$MI_\varphi = D_{KL}\left(\frac{1}{2}q_{\frac{\pi}{2}} + \frac{1}{2}q_{\frac{\pi}{4}} \,||\, q_{\frac{\pi}{2}}q_{\frac{\pi}{4}}\right),$$
$$MI_n = D_{KL}\left(\frac{1}{2}p_{\frac{\pi}{2}} + \frac{1}{2}p_{\frac{\pi}{4}} \,||\, p_{\frac{\pi}{2}}p_{\frac{\pi}{4}}\right).$$

Data availability

The code for mathematical modelling are publicly available at https://osf.io/dhcr2/. Further information for data and resources should be directed to Lead Contact, Behrad Noudoost (behrad.noudoost@utah.edu).

Acknowledgements

This work was supported by NIH grants EY026924 & NS113073 to B.N, and an Unrestricted Grant from Research to Prevent Blindness, New York, NY, to the Department of Ophthalmology & Visual Sciences, University of Utah.

Author contributions

Conceptualization, M.P., I.V., and B.N.; methodology, I.V., K.C., and B.N.; software, M.P., M.Z.; formal analysis, M.P. and M.Z., B.N.; modeling: W.N.; writing – original draft, M.P. and M.Z., K.C., B.N.; writing – review & editing, all authors; visualization, M.P., W.N., K.C.; supervision, B.N.; funding acquisition, B.N.

Declaration of interests

The authors declare no competing interests.

Supplemental information

Document S1. Figures S1-S9 and Table S1.

Figure legends

Figure 1. WM alters β oscillatory power but not firing rates in V4.

A) Memory-guided saccade with background (MGS-background) task. The monkey fixated and a peripheral visual cue appeared (Cue). The monkey maintained fixation while remembering the cue location (~1s; Delay), and after the fixation point disappeared, executed a saccadic eye movement to the remembered location (Response) to receive a reward. Throughout the task, there was a task-irrelevant, full-field oriented bar background; the background contrast ranged from 0-64%, in one of 4 orientations. The memory location was either inside the extrastriate RF (IN condition, shown) or 180 degrees away (OUT condition). Neurophysiological recordings of spiking and LFP activity were made from extrastriate visual area V4, with linear array or single electrodes.

B-C) Mean firing rate of a sample neuron over time for three different contrasts (B) or orientations (C) of the background stimulus, for the IN (left) and OUT (right) conditions. Shaded areas in all panels show standard error of mean (SEM).

D-E) Mean firing rate of the population of 145 neurons over time for three different contrasts (low, medium, and high contrast, D) and four different orientations (preferred, nonpreferred, and middle 1 & 2 orientations, E) for the IN condition (left) and OUT (right) conditions.

F) Time course of mean F-statistic values across 145 neurons, based on a one-way ANOVA for discrimination between 12 stimulus conditions for the IN (red) and OUT (black) conditions.

G) Scatter plot of F-statistic averaged in the last 700ms of the delay period for each session, for the IN vs. OUT conditions. Histogram in the upper right shows the distribution of change in F-statistic (OUT-IN) across sessions.

H) Mean power spectrum of the LFP recorded from the same channel as the sample V4 neuron in (B), during the delay period for the IN (red) vs. OUT (black) conditions. Inset panel shows 8-25 Hz. Asterisk indicates a significant difference ($p < 0.05$) in the range shown.

I) Mean power spectrum of population of the V4 LFPs (88 sessions) during the delay period for the IN (red) vs. OUT (black) conditions. Inset shows the power spectrum between 14-22 Hz.

J) Scatter plot of power spectrum averaged in the β range for each session, for the IN vs. OUT conditions. Histogram in the upper right shows the distribution of change in power (OUT-IN) across sessions (***, p <0.001).

Figure 2. WM alters the sensory representation in extrastriate cortex.

A) The distribution of spikes generated by a sample V4 neuron across various phases of β oscillations during the delay period. Arrows show the average of phase distributions for the IN (red) and OUT (black) conditions.

B) Scatter plot of SPL in the β range for each neuron, for the IN vs. OUT conditions. Histogram in the upper right shows the distribution of change in SPL (OUT-IN) across neurons.

C) Spike-triggered average of the normalized LFP of a sample neuron during the delay period, for the high contrast (red) and low contrast (blue) background stimuli, for the IN (left) and OUT (right) conditions. Shaded bars indicate the slopes in the falling phase.

D) Histogram of the distribution of STA slopes (abs $(V_{peak} - V_{trough})/(time_{peak} - time_{trough})$) across neurons, for high contrast (red) and low contrast (blue) stimuli, for the IN (left) and OUT (right) conditions.

E) Population phase (left) and rate (right) coding over time, for the IN (red) and OUT (black) conditions, based on mutual information (MI) between 12 stimulus conditions. MI was measured in 100ms windows with steps of 100ms. Shaded areas show standard error of mean (SEM).

F) Scatter plot of MI using a phase code in the β range (left) and rate code (right) for each neuron, for the IN vs. OUT conditions. Red crosses indicate population mean. Histograms in the upper right show the distribution of differences in MI (IN-OUT) across neurons.

G) Phase coding, measured by MI (colorbar), as a function of frequency and time, for memory IN (left) and OUT (right). Black rectangle shows the time and frequency range selected for phase code analysis. (*, $p < 0.05$; **, $p < 0.01$; ***, $p < 0.001$; ns, $p > 0.05$)

Figure 3. FEF inactivation alters WM behavioral performance and phase coding in visual areas.

A) V4 recordings were made before and after infusion of muscimol into FEF. Muscimol injections into FEF were made with a custom microinjectrode, at sites with stimulation-evoked saccade endpoints overlapping with simultaneous V4 recording site RFs.

B) Eye traces for 8 MGS target locations, before (left) and after (right) FEF inactivation, for an example session where 0.5 microliter of muscimol was injected into the FEF; performance deficits were localized to the infusion hemifield.

C) Average behavioral performance across sessions, at different locations over time following FEF inactivation (red pre-inactivation; green, blue, and black, 1, 2, and 3 hours after inactivation, respectively). Data from each session is aligned so that 0 degrees corresponds to the FEF RF.

D-E) Normalized saccade error (D) and reaction times (E) for the memory IN (red) and OUT (black) conditions, over time relative to the FEF inactivation. Black bar indicates times with a significant difference between IN and OUT. Shaded areas show SEM across sessions.

F) Heatmap shows phase coding (MI, colorbar) over time and frequency for 66 V4 neurons, for the IN condition, before (left) and after (right) FEF inactivation. Black rectangle indicates time and frequency range considered in (G-H): 14-22Hz, 200-800ms after start of delay period.

G) Strength of β phase coding over time, for memory IN, before (red) and after inactivation (blue). The phase-code MI is averaged in the β range. Shading shows SEM across neurons. Gray area indicates time window plotted in (H).

H) Scatter plot of β phase MI during the delay period (shaded area in G) of the IN condition for each V4 neuron, before vs. after FEF inactivation. Red square shows population mean. The histogram in the upper right shows the distribution of difference in MI (Pre-Post) across neurons. (*, $p < 0.05$; **, $p < 0.01$; ***, $p < 0.001$; ns, $p > 0.05$)

Figure 4. Experimental and computational dependence of firing rate, phase-coded information, and rate-coded information on changes in peak oscillation frequency.

A) Schematic of dynamical neural field network architecture. Excitatory and inhibitory units are interconnected and organized to respond to different input stimuli (theta). Units also receive a global WM input (not shown); see Methods for description of connectivity weights.

B) Example activity of excitatory units in the model over time, in response to an input at $\pi/2$. Excitatory units are plotted along the y-axis according to their input tuning, which ranges from 0 to π. Activity reflects both a beta-frequency oscillation across the entire population, and an earlier and stronger response of units whose preference matches the input feature (i.e., phase and rate coding).

C) Information about the input stimulus feature coded by phase and rate (left and right y-axes; see Methods) in the neural field model, as a function of WM input strength.

D) Phase information (shades of orange) and rate information (shades of blue) as a function of contrast levels and oscillation frequency, for the neural field model. Data for each contrast is divided into deciles

based on oscillation frequency, variability in which comes from noise in the WM input strength. Note that

rate code values are several orders of magnitude smaller than phase code values (left vs. right y-axis). Error

bars show standard deviation.

E) Correlation between oscillation frequency and firing rate in the neural field model under conditions of

noisy WM input strength.

F) Relationship between frequency of peak LFP power and firing rate during the delay period for the

memory IN (red) and OUT (black) conditions, for two example neurons with increased (left) or decreased

(right) firing rate during the IN condition. Each dot shows the average frequency of max power and

normalized firing rate for a subsample consisting of 50% of trials (n=100 subsamples per neuron).

G) Average normalized response as a function of peak frequency, pooled across subsamples of trials from

each of 145 V4 neurons (100 subsamples/neuron), during the IN (red) and OUT (blue) conditions, during

the delay period (left) and the cue period (right). Plot shows mean±SE for all subsamples with the peak

frequency indicated on the x-axis.

References

1.   Squire, R.F., Noudoost, B., Schafer, R.J., and Moore, T. (2013). Prefrontal Contributions to Visual
647   Selective Attention. Annual Review of Neuroscience *36*, 451–466. 10.1146/annurev-neuro-062111-
648   150439.

649   2.   Miller, E.K., and Cohen, J.D. (2001). An integrative theory of prefrontal cortex function. Annual
650   review of neuroscience *24*, 167–202. 10.1146/annurev.neuro.24.1.167.

651   3.   Miller, B.T., and D'Esposito, M. (2005). Searching for "the top" in top-down control. Neuron *48*, 535–
652   538.

653   4.   Paneri, S., and Gregoriou, G.G. (2017). Top-Down Control of Visual Attention by the Prefrontal
654   Cortex. Functional Specialization and Long-Range Interactions. Frontiers in Neuroscience *11*, 545.
655   10.3389/fnins.2017.00545.

656   5.   Gazzaley, A., and Nobre, A.C. (2012). Top-down modulation: bridging selective attention and
657   working memory. Trends in Cognitive Sciences *16*, 129–135. 10.1016/j.tics.2011.11.014.

658   6.   Knight, R.T., Grabowecky, M.F., and Scabini, D. (1995). Role of human prefrontal cortex in attention
659   control. Advances in neurology *66*, 21–34; discussion 34–26.

660   7.   Merrikhi, Y., Clark, K.L., Albarran, E., Parsa, M., Zirnsak, M., Moore, T., and Noudoost, B. (2017).
661   Spatial working memory alters the efficacy of input to visual cortex. Nature communications *8*,
662   15041. 10.1038/ncomms15041.

663   8.   Bahmani, Z., Daliri, M.R., Merrikhi, Y., Clark, K., and Noudoost, B. (2018). Working Memory Enhances
664   Cortical Representations via Spatially Specific Coordination of Spike Times. Neuron *97*, 967–
665   979.e966. 10.1016/j.neuron.2018.01.012.

666   9.   Noudoost, B., Clark, K.L., and Moore, T. (2021). Working memory gates visual input to primate
667   prefrontal neurons. eLife *10*, e64814. 10.7554/eLife.64814.

668   10.   Chelazzi, L., Duncan, J., Miller, E.K., and Desimone, R. (1998). Responses of neurons in inferior
669   temporal cortex during memory-guided visual search. Journal of neurophysiology *80*, 2918–2940.

670   11.   Mendoza-Halliday, D., Torres, S., and Martinez-Trujillo, J.C. (2014). Sharp emergence of feature-
671   selective sustained activity along the dorsal visual pathway. Nature neuroscience *17*, 1255–1262.
672   10.1038/nn.3785.

673   12.   Leavitt, M.L., Mendoza-Halliday, D., and Martinez-Trujillo, J.C. (2017). Sustained Activity Encoding
674   Working Memories: Not Fully Distributed. Trends in Neurosciences *40*, 328–346.
675   10.1016/j.tins.2017.04.004.

676   13.   Chelazzi, L., Miller, E.K., Duncan, J., and Desimone, R. (2001). Responses of neurons in macaque area
677   V4 during memory-guided visual search. Cerebral cortex (New York, N.Y. : 1991) *11*, 761–772.

678 14. Zaksas, D., and Pasternak, T. (2006). Directional signals in the prefrontal cortex and in area MT during
679   a working memory for visual motion task. The Journal of neuroscience *26*, 11726–11742.
680   10.1523/JNEUROSCI.3420-06.2006.

681 15. Soto, D., Wriglesworth, A., Bahrami-Balani, A., and Humphreys, G.W. (2010). Working memory
682   enhances visual perception: Evidence from signal detection analysis. Journal of Experimental
683   Psychology: Learning, Memory, and Cognition *36*, 441–456. 10.1037/a0018686.

684 16. Pan, Y., and Zhang, X. (2020). Visual working memory enhances target discrimination accuracy with
685   single-item displays. Attention, Perception & Psychophysics *82*, 3005-3012. 10.3758/s13414-020-
686   02041-y.

687 17. Hallenbeck, G.E., Sprague, T.C., Rahmati, M., Sreenivasan, K.K., and Curtis, C.E. (2021). Working
688   memory representations in visual cortex mediate distraction effects. Nature Communications *12*,
689   4714. 10.1038/s41467-021-24973-1.

690 18. Hollingworth, A., and Luck, S.J. (2009). The role of visual working memory (VWM) in the control of
691   gaze during visual search. Attention, perception & psychophysics *71*, 936–949.
692   10.3758/APP.71.4.936.

693 19. Hollingworth, A., Matsukura, M., and Luck, S.J. (2013). Visual working memory modulates rapid eye
694   movements to simple onset targets. Psychological science *24*, 790–796.
695   10.1177/0956797612459767.

696 20. Pan, Y., Lin, B., Zhao, Y., and Soto, D. (2014). Working memory biasing of visual perception without
697   awareness. Attention, Perception & Psychophysics *76*, 2051-2062. 10.3758/s13414-013-0566-2.

698 21. Salahub, C.M., and Emrich, S.M. (2016). Tuning perception: Visual working memory biases the
699   quality of visual awareness. Psychonomic Bulletin & Review *23*, 1854-1859. 10.3758/s13423-016-
700   1064-z.

701 22. Logothetis, N.K., Pauls, J., Augath, M., Trinath, T., and Oeltermann, A. (2001). Neurophysiological
702   investigation of the basis of the fMRI signal. Nature *412*, 150–157. 10.1038/35084005.

703 23. Buzsáki, G., Anastassiou, C.A., and Koch, C. (2012). The origin of extracellular fields and currents--
704   EEG, ECoG, LFP and spikes. Nature Reviews. Neuroscience *13*, 407-420. 10.1038/nrn3241.

705 24. Einevoll, G.T., Kayser, C., Logothetis, N.K., and Panzeri, S. (2013). Modelling and analysis of local field
706   potentials for studying the function of cortical circuits. Nature Reviews Neuroscience *14*, 770–785.
707   10.1038/nrn3599.

708 25. Herreras, O. (2016). Local Field Potentials: Myths and Misunderstandings. Frontiers in Neural
709   Circuits *10*, 101. 10.3389/fncir.2016.00101.

710 26. McCormick, D.A., Connors, B.W., Lighthall, J.W., and Prince, D.A. (1985). Comparative
711   electrophysiology of pyramidal and sparsely spiny stellate neurons of the neocortex. Journal of
712   neurophysiology *54*, 782–806. 10.1152/jn.1985.54.4.782.

27.   Sommer, M.A., and Tehovnik, E.J. (1997). Reversible inactivation of macaque frontal eye field.
Experimental brain research. *116*, 229–249.

28.   Dias, E.C., and Segraves, M.A. (1999). Muscimol-induced inactivation of monkey frontal eye field:
effects on visually and memory-guided saccades. Journal of neurophysiology *81*, 2191–2214.

29.   Fuster, J.M. (1990). Inferotemporal units in selective visual attention and short-term memory.
Journal of Neurophysiology *64*, 681-697. 10.1152/jn.1990.64.3.681.

30.   Supèr, H., Spekreijse, H., and Lamme, V.A.F. (2001). A neural correlate of working memory in the
monkey primary visual cortex. Science *293*, 120–124. 10.1126/science.1060496.

31.   Hayden, B.Y., and Gallant, J.L. (2013). Working memory and decision processes in visual area v4.
Front Neurosci *7*, 18. 10.3389/fnins.2013.00018.

32.   Anderson, J.C., Kennedy, H., and Martin, K.A.C. (2011). Pathways of attention: synaptic relationships
of frontal eye field to V4, lateral intraparietal cortex, and area 46 in macaque monkey. The Journal
of neuroscience : the official journal of the Society for Neuroscience *31*, 10872–10881.
10.1523/JNEUROSCI.0622-11.2011.

33.   Lara, A.H., and Wallis, J.D. (2015). The Role of Prefrontal Cortex in Working Memory: A Mini Review.
Frontiers in Systems Neuroscience *9*, 173. 10.3389/fnsys.2015.00173.

34.   Zanto, T.P., Rubens, M.T., Thangavel, A., and Gazzaley, A. (2011). Causal role of the prefrontal cortex
in top-down modulation of visual processing and working memory. Nature Neuroscience *14*, 656–
661. 10.1038/nn.2773.

35.   Noudoost, B., Chang, M.H., Steinmetz, N.A., and Moore, T. (2010). Top-down control of visual
attention. Current opinion in neurobiology *20*, 183–190.

36.   Snyder, A.C., Yu, B.M., and Smith, M.A. (2021). A Stable Population Code for Attention in Prefrontal
Cortex Leads a Dynamic Attention Code in Visual Cortex. Journal of Neuroscience *41*, 9163-9176.
10.1523/JNEUROSCI.0608-21.2021.

37.   Kempen, J.v., Gieselmann, M.A., Boyd, M., Steinmetz, N.A., Moore, T., Engel, T.A., and Thiele, A.
(2021). Top-down coordination of local cortical state during selective attention. Neuron *109*, 894-
904.e898. 10.1016/j.neuron.2020.12.013.

38.   Fiebelkorn, I.C., and Kastner, S. (2021). Spike Timing in the Attention Network Predicts Behavioral
Outcome Prior to Target Selection. Neuron *109*, 177-188.e174. 10.1016/j.neuron.2020.09.039.

39.   Moran, J., and Desimone, R. (1985). Selective attention gates visual processing in the extrastriate
cortex. Science *229*, 782–784.

40.   Merrikhi, Y., Clark, K.L., and Noudoost, B. (2018). Concurrent influence of top-down and bottom-up
inputs on correlated activity of Macaque extrastriate neurons. Nature Communications *9*, 5393.
https://doi.org/10.1038/s41467-018-07816-4.

41. Cohen, M.R., and Maunsell, J.H. (2009). Attention improves performance primarily by reducing
interneuronal correlations. Nature neuroscience *12*, 1594–1600. 10.1038/nn.2439.

42. Mitchell, J.F., Sundberg, K.A., and Reynolds, J.H. (2007). Differential attention-dependent response
modulation across cell classes in macaque visual area V4. Neuron *55*, 131–141.
10.1016/j.neuron.2007.06.018.

43. Womelsdorf, T., Anton-Erxleben, K., and Treue, S. (2008). Receptive field shift and shrinkage in
macaque middle temporal area through attentional gain modulation. The Journal of neuroscience :
the official journal of the Society for Neuroscience *28*, 8934–8944. 10.1523/JNEUROSCI.4030-
07.2008.

44. Womelsdorf, T., Anton-Erxleben, K., Pieper, F., and Treue, S. (2006). Dynamic shifts of visual
receptive fields in cortical area MT by spatial attention. Nature neuroscience *9*, 1156–1160.
10.1038/nn1748.

45. Rezayat, E., Dehaqani, M.-R.A., Clark, K., Bahmani, Z., Moore, T., and Noudoost, B. (2021).
Frontotemporal coordination predicts working memory performance and its local neural signatures.
Nature Communications *12*, 1103. 10.1038/s41467-021-21151-1.

46. Liebe, S., Hoerzer, G.M., Logothetis, N.K., and Rainer, G. (2012). Theta coupling between V4 and
prefrontal cortex predicts visual short-term memory performance. Nature Neuroscience *15*, 456–
462. 10.1038/nn.3038.

47. Antzoulatos, E.G., and Miller, E.K. (2016). Synchronous beta rhythms of frontoparietal networks
support only behaviorally relevant representations. eLife *5*, e17822. 10.7554/eLife.17822.

48. Salazar, R.F., Dotson, N.M., Bressler, S.L., and Gray, C.M. (2012). Content-specific fronto-parietal
synchronization during visual working memory. Science *437*, 1097–1101.

49. Dotson, N.M., Salazar, R.F., and Gray, C.M. (2014). Frontoparietal correlation dynamics reveal
interplay between integration and segregation during visual working memory. The Journal of
Neuroscience: The Official Journal of the Society for Neuroscience *34*, 13600-13613.
10.1523/JNEUROSCI.1961-14.2014.

50. Rezayat, E., Clark, K., Dehaqani, M.-R.A., and Noudoost, B. (2022). Dependence of Working Memory
on Coordinated Activity Across Brain Areas. Frontiers in Systems Neuroscience *15*, 164.
10.3389/fnsys.2021.787316.

51. Jia, X., Tanabe, S., and Kohn, A. (2013). γ and the coordination of spiking activity in early visual cortex.
Neuron *77*, 762-774. 10.1016/j.neuron.2012.12.036.

52. Fries, P. (2015). Rhythms for Cognition: Communication through Coherence. Neuron *88*, 220–235.
10.1016/j.neuron.2015.09.034.

53. Bastos, A.M., Vezoli, J., and Fries, P. (2015). Communication through coherence with inter-areal
delays. Current Opinion in Neurobiology *31*, 173–180. 10.1016/j.conb.2014.11.001.

54. Fries, P. (2005). A mechanism for cognitive dynamics: neuronal communication through neuronal
coherence. Trends in cognitive sciences *9*, 474–480. 10.1016/j.tics.2005.08.011.

55. Parto-Dezfouli, M., Vezoli, J., Bosman, C.A., and Fries, P. (2023). Enhanced behavioral performance
through interareal gamma and beta synchronization. Cell Rep *42*, 113249.
10.1016/j.celrep.2023.113249.

56. Cardin, J.A., Carlén, M., Meletis, K., Knoblich, U., Zhang, F., Deisseroth, K., Tsai, L.-H., and Moore, C.I.
(2009). Driving fast-spiking cells induces gamma rhythm and controls sensory responses. Nature
459, 663–667. 10.1038/nature08002.

57. Womelsdorf, T., Schoffelen, J.-M., Oostenveld, R., Singer, W., Desimone, R., Engel, A.K., and Fries, P.
(2007). Modulation of neuronal interactions through neuronal synchronization. Science *316*, 1609–
1612. 10.1126/science.1139597.

58. Bosman, C.A., Schoffelen, J.-M., Brunet, N., Oostenveld, R., Bastos, A.M., Womelsdorf, T., Rubehn,
B., Stieglitz, T., De Weerd, P., and Fries, P. (2012). Attentional stimulus selection through selective
synchronization between monkey visual areas. Neuron *75*, 875–888.
10.1016/j.neuron.2012.06.037.

59. Harrison, S.A., and Tong, F. (2009). Decoding reveals the contents of visual working memory in early
visual areas. Nature *458*, 632–635. 10.1038/nature07832.

60. Pasternak, T., and Greenlee, M.W. (2005). Working memory in primate sensory systems. Nature
reviews. Neuroscience *6*, 97–107. 10.1038/nrn1603.

61. D'Esposito, M., and Postle, B.R. (2015). The Cognitive Neuroscience of Working Memory. Annual
Review of Psychology *66*, 115–142. 10.1146/annurev-psych-010814-015031.

62. Comeaux, P., Clark, K., and Noudoost, B. (2023). A recruitment through coherence theory of working
memory. Progress in Neurobiology *228*, 102491. 10.1016/j.pneurobio.2023.102491.

63. Vanegas, M.I., Hubbard, K.R., Esfandyarpour, R., and Noudoost, B. (2019). Microinjectrode System
for Combined Drug Infusion and Electrophysiology. Journal of Visualized Experiments: JoVE.
10.3791/60365.

64. Noudoost, B., and Moore, T. (2011). A reliable microinjectrode system for use in behaving monkeys.
Journal of neuroscience methods *194*, 218–223. 10.1016/j.jneumeth.2010.10.009.

65. Zarei, M., Parto Dezfouli, M., Jahed, M., and Daliri, M.R. (2020). Adaptation Modulates Spike-Phase
Coupling Tuning Curve in the Rat Primary Auditory Cortex. Front Syst Neurosci *14*, 55.
10.3389/fnsys.2020.00055.

66. Kayser, C., Montemurro, M.A., Logothetis, N.K., and Panzeri, S. (2009). Spike-phase coding boosts
and stabilizes information carried by spatial and temporal spike patterns. Neuron *61*, 597–608.
10.1016/j.neuron.2009.01.008.

67.   Magri, C., Whittingstall, K., Singh, V., Logothetis, N.K., and Panzeri, S. (2009). A toolbox for the fast
817   information analysis of multiple-site LFP, EEG and spike train recordings. BMC Neurosci *10*, 81.
818   10.1186/1471-2202-10-81.

819   68.   Nesse, W.H., Clark, K.L., and Noudoost, B. (2023). Information representation in an oscillating neural
820   field model modulated by working memory signals. Front Comput Neurosci *17*, 1253234.
821   10.3389/fncom.2023.1253234.

822   69.   Sanzeni, A., Histed, M.H., and Brunel, N. (2020). Response nonlinearities in networks of spiking
823   neurons. PLoS Comput Biol *16*, e1008165. 10.1371/journal.pcbi.1008165.
824

A) FEF, Injectrode, V4 RF, FEF-evoked saccade vector, V4

B) Pre, Post (3hr), 5 dva, 5 dva

C) 90°, 180°, 0°, -90°, 0%, 100% Performance, Pre, Post(1hr), Post(2hr), Post(3hr)

D) Saccade error (norm. unit), IN, OUT, 33 sessions, Time from injection (min)

E) Reaction time (norm. unit), IN, OUT, Time from injection (min)

F) IN, Pre, IN, Post, Frequency (Hz), Cue, WM, Cue, WM, Phase MI (bits)

G) Phase MI (bits), Pre, Post, Cue, WM, 66 neurons

H) Phase MI [Post], Phase MI [Pre], p=0.00521