Isotope analysis of birds' eye lens provides early-life information

Emi Arai Hasegawa *[1]

Email: ehasegawa3@chikyu.ac.jp

Jun Matsubayashi*[2]

Email: matsuj@fpu.ac.jp

*These two authors contributed equally to this work.

Ichiro Tayasu[1]

Email: ichirotayasu@chikyu.ac.jp

Tatsuhiko Goto[3,4]

Email: tats.goto@obihiro.ac.jp

Haruka Inoue[5]

Axel G. Rossberg[6]

Email: a.rossberg@qmul.ac.uk

Chikage Yoshimizu[1]

Email: chikage@chikyu.ac.jp

Masaru Hasegawa[7]

Email : perorobomusadiobe@gmail.com

Takumi Akasaka[5]

Corresponding author

Email : mail2taq@gmail.com

(Institutional addresses)

[1] Research Institute for Humanity and Nature, 457-4 Motoyama, Kamigamo, Kita-ku, Kyoto,

603-8047, Japan

[2] Faculty of Marine Science and Technology, Fukui Prefectural University, 1-1 Gakuen-cho,

Obama, Fukui 917-0003, Japan

[3] Department of Life and Food Sciences, Obihiro University of Agriculture and Veterinary

Medicine, 11-2 Inadamachi-nishi, Obihiro, Hokkaido 080-8555, Japan

[4] Research Center for Global Agromedicine, Obihiro University of Agriculture and Veterinary

Medicine, 11-2 Inadamachi-nishi, Obihiro, Hokkaido 080-8555, Japan

[5] Laboratory of Conservation Ecology, Obihiro University of Agriculture and Veterinary

Medicine, 11-2 Inadamachi-nishi, Obihiro, Hokkaido 080-8555, Japan

[6] School of Biological and Chemical Sciences, Queen Mary University of London, Mile End

Road, London E1 4NS, UK

[7] Department of Environmental Science, Ishikawa Prefectural University, 1-308 Suematsu,

Nonoichi, Ishikawa 921-8836, Japan

Abstract

Early-life environment has a long-lasting effect on later life, though its estimation is often

prevented in the wild because of a lack of available methods. Recently, isotope analysis of eye

lenses has attracted considerable interest as a means to reconstruct the environmental conditions

experienced by animals during the developmental period. This analysis has mostly been

confined to fish for practical reasons and remains to be resolved for application to other animals.

In this study, we broadened its applicability by developing a novel approach and verifying its

usability for the reconstruction of early-life environments. We performed a feeding experiment

using Japanese quail (*Coturnix japonica*), in which we administered two diets: one composed

mainly of C3 plants (low δ^{13}C and high δ^{15}N) and the other of C4 plants (high δ^{13}C and low

δ^{15}N). Quails in the control group were continuously fed a C3-based diet from hatching until

200 days old, whereas those in the treatment groups (T10, T15, T20, and T40) were switched

from the C3 to the C4-based diet at 10, 15, 20, and 40 days after hatching, respectively. We

found that the δ^{13}C in the eye lenses of the treatment groups decreased from the center layer to

the middle layer of the lens and then increased toward the outer layer, thus reflecting the diet

change. In contrast, those of the control group exhibited a decreasing trend and equilibrated at

the middle layer of the eye lens, with no increase thereafter. This novel approach revealed the

postnatal feeding histories of the diet-shift experiment. The high δ^{13}C values observed in the

center of the eye lenses would reflect the prenatal feeding environment, i.e., the C4-based diet

consumed by their mothers, which is further reinforced by higher δ^{15}N values at this position

due to the consumption of egg yolk-derived nutrition. These results indicate that the avian eye

lens can be used as an "isotopic chronicle," which is a useful tool for reconstructing

chronological isotopic information about their early-life history.

Keywords

Eye lens, Feeding experiment, Chronology, Stable isotope, Japanese quail

1 Introduction

Early-life environment has a long-lasting effect on the later life of animals and is known as the "silver spoon hypothesis" (Monaghan 2008). Food availability, habitat quality, or physiological stress during early life can affect future development, survival, and reproduction (Carlisle et al. 2015; Hayward et al. 2013; Hamel et al. 2009; Nussey et al. 2007). Understanding the early-life environments of organisms is important in pure science to clarify the exact causality of phenotypic development as well as in applied science for the conservation of declining species (Sergio et al. 2022). However, early-life information of grown animals is often challenging to obtain because continuous long-term research is required to track the same individual from natal to adult period, and thus a retrospective approach for the reconstruction of early-life environments is required.

Isotope analysis is a widely used technique for investigating the trophic status and migratory origins of organisms across numerous taxa (Hobson and Wassenaar 2018). For example, isotope analysis plays a pivotal role in elucidating the ecological dynamics of long-distance migrants, such as birds, because of the challenges associated with tracking their

migratory patterns and dietary preferences (Inger and Bearhop 2008; Tzadik et al. 2017). Inert

integumentary tissues, such as bird feathers, have often been used as exemplary tissues for

retrospective isotope analysis (Hobson and Clark 1992; Wolf et al. 2013), as they provide

information on their development. In particular, isotope analysis of feathers has played an

important role in avian ecology by explaining migratory patterns with a relatively small cost

when compared to other methodologies, such as geo-loggers and GPS tracking (Hobson et al.

2012; Guiterrez-Exposito et al. 2015).

However, feathers and other inert integumentary tissues are not flawless because

information may be lost after their renewal, i.e., after molting (Carvalho et al. 2022). Therefore,

these analyses cannot provide permanent information about early-life environments, such as

their birthplaces. Early-life information, which is particularly important, needs to be collected

by other, more permanent inert tissues.

One tissue that can potentially preserve the environmental conditions experienced

during early life is the eye lens, as eye lenses are composed of metabolically inert proteins that

are sequentially deposited (Vecchio and Peebles 2020). Previous studies considering fish eyes

indicated that eye lenses can be a valuable tool for reconstructing past environments. However,

unlike fish and mammals, avian eye lenses are challenging to analyze because of their softness

and water solubility (Simpson et al. 1994); therefore, there is a need to develop a method for

avian eye lens analysis.

Avian lens maturation follows a pattern that is common to vertebrates, with new fibers

continuously emerging from the equatorial epithelial cells (Sivak 2004). The avian eye lens

develops during embryonic stages, and it is postulated that only the outermost lens harbors post-

embryonic information (Peebles and Hollander 2020). In addition, during the embryonic stage,

synthesized proteins are transported to the aged and metabolically inactive cells at the center of

the lens, referred to as the organelle-free zone (OFZ) (Shestopalov and Bassnett 2000), which

can be challenging to clarify via isotope analysis of the early-life environment. On the other

hand, unlike fish, the avian eye lens does not grow larger for more than 50 days, and also does

not undergo compaction at the center of the lens (Augusteyn 2008). Thus, once we can develop

a protocol to analyze avian eye lenses, a relatively stable lens size would make a useful tool to

reconstruct the developmental environment. By establishing a diet-shift experiment in early life,

we can assess whether the eye lens reveals the diet history and, if any, develop a formula to

reconstruct the history of the early-life environment from the isotopic information while

considering the substantial mixing of the isotopes in the radial direction (i.e., diffusion by the

above-mentioned biological transport process and sample preparation process) (Passey and

Cerling 2002).

The purpose of this study is to determine whether the environment experienced during

the early-life stage can be reconstructed from an avian eye lens. To achieve this, we performed

a feeding experiment on captive Japanese quails (*Coturnix japonica*) for 200 days, using diets

derived from C3 and C4 plants. After hatching, chicks received a C3-based diet (hereafter C3

food). The C3 food of the control group was maintained throughout the experiment, while the

treatment groups switched to a C4-based diet (hereafter C4 food) at specific points (Fig. 1).

Following the feeding period, we analyzed a time series of the isotopic ratios in both the eye

lens and primary feathers. Based on the data, we aim to (1) test whether birds' lenses record the

chronological isotopic information and (2) determine the duration of such information retention.

Furthermore, we investigate the correspondence between the position in the eye lens and the

age at which the information is deposited by developing a formula.

2 Methods/Experimental

2.1 Diet-shift experiments

For the C3 food (Kusanagi Farm Limited Company, Obihiro, Japan), food residues from potato

peels, confectionery waste, cotton and pumpkin seeds, sake lees, and other sources were utilized.

This mixture was fermented with lactic acid bacteria and combined with wheat, soybeans, yams,

scallops, rice bran, starch flour, fishmeal, and beet cake (Mori et al. 2020). In contrast, the C4

food employed a commercially available mixed feed (Runkeeper, Marubeni Nisshin Feed Co.,

Ltd., Japan) (Goto et al. 2023), containing imported corn and additional components (Mori et

al. 2020). Feeding trials for both diets were performed at a dedicated facility in the Obihiro

University of Agriculture and Veterinary Medicine between June 1 and October 20, 2019.

Japanese quail eggs were collected from parent birds that were raised on the above-

mentioned C4 food (Runkeeper) (Goto et al. 2023) and were incubated at 36.5°C until hatching

($n = 31$). Hatched chicks were housed individually and fed C3 food for 10 days. They were then

divided into control ($n = 6$) and treatment ($n = 25$) groups. The control group was fed C3 food

throughout the trial. The treatment group switched to the C4 food at specific time points: 10

days (T10; $n = 6$); 15 days (T15; $n = 10$); 20 days (T20; $n = 7$); and 40 days (T40; $n = 2$) after

the start of rearing. The final rearing period for the treatment groups included time points at 40,

80, 120, 160, 168, and 200 days (Table 1). The control group's final rearing periods were 17,

40, 80, 120, 160, 168, and 200 days.

2.2 Eye lens dehydration and dissection

To perform the continuous dissection of the fragile avian eye lens, which is susceptible to

liquefaction at room temperature, we established the following dehydration and drying protocol,

that was modified from the method described by Matsubayashi et al. (in preparation):

1.   After a minor incision in the sclera with scissors, the frozen eyeballs were immersed in 10

mL of absolute ethanol within a screw-capped plastic tube on ice. This initial cut enabled

ethanol access to the eye lens while maintaining its structural integrity.

2. Following immersion for 10 min, the eyeball was extracted, the scleral incision was

enlarged, and the eyeball was returned to fresh absolute ethanol. This process was repeated

with incisions made along the equatorial plane of the sclera. This step enhanced surface

dehydration compared to the initial immersion because of the improved ethanol penetration.

3. Thirty minutes after the initial immersion, the eyeball was removed from the ethanol, and

the eye lens was carefully extracted using a spatula, tweezers, and scissors. It was then

placed in fresh absolute ethanol on ice for 30 min, followed by 60 min at room temperature.

This ensured the through dehydration of the entire eye lens through continued ethanol

exposure.

4. For complete internal eye lens desiccation, the eye lenses were dried overnight at 40°C in a

dedicated drying oven (DO-600FPA, As ONE, Osaka, Japan). The 40°C drying temperature

was selected to adjust to the general body temperature of quails (Underwood et al. 1999).

The dried eye lens was then dissected according to the protocol described by Matsubayashi

et al. (in preparation):

1. The lens was positioned on a Petri dish, photographed under a dissecting microscope (Leica

M125, Leica Microsystems GmbH, Wetzlar, Germany) equipped with a Moticam S3 camera

(Shimadzu Rika Corporation, Tokyo, Japan), and its diameter was measured from the

circumference.

2. The lens was moistened with a minimal amount of deionized water. Using fine-tipped

tweezers, the soft material was carefully collected from across the entire surface and

transferred to a new designated container.

3. After scraping off approximately 0.5–200.0 µg of eye lens sample for isotope analysis while

preserving the overall lens shape, any remaining water was wiped away, and the used

instruments were washed or replaced as required.

4. The above steps 1–3 were repeated until the central core of the eye lens was reached.

Because of the inherent aspherical nature and unevenness of the ethanol-dehydrated quail ocular

lenses, the short axis was used as the basis for size determination. This defines the lens size as

the shortest diameter at the equator (mm). Ethanol treatment for 2 h should not have a significant

effect on the stable isotope analysis of epithelial tissue or amino acids, including the eye lens

(Borrow et al. 2008; Chua et al. 2020). The analysis excluded the outermost lens section

containing the attached fibrae zonulares (Fitzgerald 2018). This discrepancy between lens size

and the outermost analyzed position may be due to this exclusion. Finally, individual lens

sections were freeze-dried using a freeze dryer (FDU-2100, EYELA, Tokyo, Japan). We

excluded samples of the 160-day-old control quail because of sample degradation.

2.3 Feather processing

As the first primary feathers had already completely grown by 40-days old, we only analyzed

the first primary (P1) feathers (Paritte and Kelly 2009; Wolf et al. 2013) from 40, 120, and 160-

day-old birds (control, T10, T15, and T20). The samples were washed with water, soaked in a

2:1 chloroform: methanol solution for 24h, and then rinsed twice with methanol to remove any

surface contaminants and oils. After being air-dried in a fume food for more than 24h (Paritte

and Kelly 2009), the samples were moved to a drying oven (DO-600FPA, As ONE, Osaka,

Japan) at 60°C for more than 24h. The feather vanes were cut using surgical scissors from the

tip to the base, approximately 2–3 mm each but approximately 10 mm for the narrow feather

tip, which was sufficient for isotope analysis.

2.4 Stable isotope analysis

We performed continuous dissection of the fragile avian eye lenses that are susceptible to

liquefaction and placed them along with 0.05–0.16 mg of feathers and lens sections, and 0.65

mg of C3 food, and 0.86 mg of C4 food into tin capsules for the carbon and nitrogen stable

isotope analyses. Isotope ratios were measured using a Delta V Advantage mass spectrometer

(Thermo Fisher Scientific, Waltham, MA, USA) connected to a Flash EA 2000 elemental

analyzer via a Conflo IV interface (Thermo Fisher Scientific). The stable carbon ($^{13}C/^{12}C$) and

nitrogen ($^{15}N/^{14}N$) isotopic compositions of the sample (δ_{sample}) were expressed in standard delta

(δ) notation according to the following formula (Couplen 2011):

$$\delta_{sample} = (R_{sample}/R_{std}) - 1$$

where R_{sample} is the corresponding isotope ratio $^{13}C/^{12}C$ or $^{15}N/^{14}N$ for the sample. R_{std} is the

defined isotope ratio by international standard scales, where $^{13}C/^{12}C$ is the Vienna Pee Dee

Belemnite (VPDB) for $^{13}C/^{12}C$ and atmospheric nitrogen for $^{15}N/^{14}N$. The isotope ratios of

carbon and nitrogen were calibrated against those of the DL-alanine ($\delta^{13}C = -25.36‰$, $\delta^{15}N =$

$-2.89‰$; CERKU-01), L-alanine ($\delta^{13}C = -19.04‰$, $\delta^{15}N = 22.71‰$; CERKU-02), and L-

threonine ($\delta^{13}C = -9.45‰$, $\delta^{15}N = -2.88‰$; CERKU-05) laboratory standards, which were

calibrated against international standards (Tayasu et al. 2011). The analytical standard

deviations (SDs) of the DL-alanine standard ($n = 53$) were 0.09‰ ($\delta^{13}C$) and 0.04‰ ($\delta^{15}N$);

those of the L-alanine standard ($n = 50$) were 0.10‰ ($\delta^{13}C$, $n = 51$) and 0.06‰ ($\delta^{15}N$, $n = 56$);

and those of the threonine standard ($n = 54$) were 0.12‰ ($\delta^{13}C$) and 0.07‰ ($\delta^{15}N$).

2.5 Data analysis

To investigate the correspondence between the position in the eye lens and the age at which the

information is deposited, we developed the following model, which has as its starting point a

piecewise constant function (PCF), which represents the signature in the eye lens of the

switching of diets from the maternal diet (C4 food) to the initial diet (C3 food) and then back

to C4 food. This function was convoluted with a normalized Gaussian profile with a standard

deviation σ to model the mixing of isotope ratios in the radial direction, which could be caused

by contamination of layers with different time frames during the sample preparation or through

biological transport process. The convolution evaluates to:

$$\delta = \delta_1 + \frac{1}{2}(\delta_1 - \delta_2)\left[\mathrm{erf}\left(\frac{\frac{r_h}{1+ca} - r}{\sqrt{2}\sigma}\right) - \mathrm{erf}\left(\frac{\frac{r_h + r_T}{1+ca} - r}{\sqrt{2}\sigma}\right)\right]$$

where $\mathrm{erf}(\cdot)$ denotes the conventional error function; δ_1 is the asymptotic value for C4 food,

i.e., the value where an isotopic equilibrium state is achieved after a change in diet (Ayliffe et

al. 2004); δ_2 is the asymptotic value for C3 food; a is the age of quails at the time of

dissection; r_h is the radius corresponding to the time of hatching; c is the rate of lens

compaction; r_T is the radius corresponding to the start of the C4 feeding period ($r_T = \infty$ for

the control); and r is the actual distance from the lens center.

We fitted this model to the complete $\delta^{13}C$ dataset (Supplementary data by minimizing

the mean squared difference between $\delta^{13}C$ observed at radius r and the model prediction of δ

by adjusting the model parameters σ, δ_1, δ_2, a, r_h, c, and the four independent parameters

of r_T corresponding to the four diet switching times at $T = 10, 15, 20$ and 40 days. From this,

we estimated the transition lens radius ($r_h + r_T$) for each of the dissection date (Supplementary

Figure S3). We fitted two model variants, with and without ($c = 0$) lens compaction. In the eye

lens of vertebrates, there is a mechanism called compaction, in which the central part of the lens

shrinks and does not become larger (Augusteyn 2008), which we account for through the c

parameter. To date, no studies have examined whether compaction occurs over time in the avian

eye lens. Therefore, we constructed models both with and without the compaction parameter

(c) and compared them using Akaike information criterion (AIC).

To confirm whether quail lens sizes increased with the dissection date, the

relationship between lens size and age (17– 200 day old) was analyzed using a linear model

(LM). The isotopic offset between the tissue and the diet (e.g., $\delta^{13}C_{lens - diet}$) was calculated by

subtracting the isotope ratios of the diet from those of the equilibrated tissue.

For the $\delta^{15}N$ isotope ratio, the asymptotic value was also estimated by following the

PCF model:

265
$$\delta = \frac{1}{2}\left[(\delta_1 + \delta_3) + (\delta_1 - \delta_2)\,\mathrm{erf}\left(\frac{\frac{r_h}{1+ca} - r}{\sqrt{2}\sigma}\right) + (\delta_2 - \delta_3)\,\mathrm{erf}\left(\frac{\frac{r_h + r_T}{1+ca} - r}{\sqrt{2}\sigma}\right)\right]$$

266   where δ_1 is the asymptotic value of maternal nutrition, i.e., a high value should be assumed

267   by bioenrichment irrespective of food contents; δ_2 is the asymptotic value for C3 food; δ_3 is

268   the asymptotic value for C4 food. The other numerical statements are the same as the above-

269   mentioned PCF model for $\delta^{13}C$. All data analyses were performed using the R statistical

270   computing environment version 4.1.2 (R Core Team, 2021).

3 Results

3.1 Stable isotope ratios of diet and eye lens

The $\delta^{13}C$ values of the C3 and C4 food were $-26.4‰$ and $-17.1‰$, respectively. The PCF model, including compaction, provided a better fit (AIC = 804.5) over the model without controlling for lens compaction (AIC = 837.3, Supplementary Fig. S1). Therefore, the pattern of isotope shift was inferred from the model with compaction (Supplementary Table S1, showing all fitted parameters). In particular, the control quails initially had a high $\delta^{13}C$ (average ± SD = $-19.7‰$ ± 1.4‰, with a range of -18.4 to $-22.1‰$), which decreased with the lens radius (i.e., age), and stabilized at $-23.9‰ ± 0.2‰$ (Supplementary Table S1). The treatment groups exhibited a peak $\delta^{13}C$ value in the innermost lens sections, which then dropped and rose again towards the outer sections (Fig. 2). Treatment groups achieved maximum $\delta^{13}C$ values in the outermost lens sections, which approached the equilibrated $\delta^{13}C$ value of the C4 food (Supplementary Fig. S1, $-17.6‰ ± 0.3‰$). Individual variation in $\delta^{13}C$ existed even among quails with the same rearing conditions (Fig. 2).

The $\delta^{15}N$ values of the C3 and C4 foods were $4.6‰$ and $2.1‰$, respectively. The PCF model that included the compaction parameter c provided a better fit (AIC = 388.5) compared to the model without c (AIC = 416.9, Supplementary Fig. S2). Therefore, the isotope shift

pattern was inferred from the model with compaction (Supplementary Table S2, showing all

fitted parameters). The $\delta^{15}N$ values in the center of the eye lens were approximately 5.7‰ ±

0.1‰, which decreased with age and reached equilibrium at 4.5‰ ± 0.1‰ (Supplementary

Table S2, Fig. 3(a)). The offset of $\delta^{15}N$ between the C3 food (4.6 ‰) and eye lens ($\delta^{15}N_{lens\text{-}diet}$)

was estimated to be −0.1 ‰. The treatment group also showed a $\delta^{15}N$ shifts (i.e., decrease) from

the center to outer eye lens sections, achieving equilibrium at 3.9‰ ± 0.1‰ (Supplementary

Fig. S2, Supplementary Table S2), with individual variations, especially at the innermost

sections (Fig. 3). The offset of the $\delta^{15}N$ between the C4 food (2.1 ‰) and eye lens ($\delta^{15}N_{lens\text{-}diet,}$

3.9 ‰) was estimated to be 1.8 ‰.

3.2 Lens size and compaction

The mean lens size was 1.00 ± 0.07 mm at 17 days ($n = 2$), 1.23 ± 0.13 mm at 40 days ($n = 8$),

1.21 ± 0.15 mm at 80 days ($n = 8$), 1.22 ± 0.15 mm at 120 days ($n = 6$), 1.34 ± 0.11 mm at 160

days ($n = 7$), 1.24 ± 0.07 mm at 168 days ($n = 3$), and 1.10 ± 0.13 mm at 200 days ($n = 7$). There

was no significant increase in the eye lens radius with age (from 17 days to 200 days old, $n =$

41, LM: Coef. ± SE = 0.000082 ± 0.0004, t = 0.21, $p = 0.83$). From the PCF models, the lens

compaction rate was estimated to be 0.0012 mm per day for the $\delta^{13}C$ dataset (Supplementary

Fig. S3(a), Supplementary Table S1) and 0.0013 mm per day for the $\delta^{15}N$ dataset

(Supplementary Fig. S3(b), Supplementary Table S2).

3.3 Stable isotope ratios of primary feathers

We determined the $\delta^{13}C$ and $\delta^{15}N$ values of the P1 feathers from quails at 40 days, 120 days,

and 160 days (control, T10, T15, and T20 for each). The feather $\delta^{13}C$ values in the control group

remained almost constant throughout the feather section (Fig. 4, mean $\pm$ SD $= -23.6‰ \pm 0.2$ ‰,

$n = 3$). In the treatment group, all 40-day-old quails exhibited lower $\delta^{13}C$ values in the tip

sections, which increased with growth and reached equilibrium around the $\delta^{13}C$ value of the C4

food (Fig. 4). The other birds showed almost constant $\delta^{13}C$ values (Fig. 4, $-16.7‰ \pm 0.4$ ‰).

The feather $\delta^{15}N$ values of the 120 and 160-day-old quails in the control and treatment

groups exhibited individual differences but remained consistent within the individuals (mean $\pm$

SD; control group: 3.9‰ $\pm$ 0.5‰, treatment groups: 3.4‰ $\pm$ 0.5‰, Supplementary Fig. S3). In

the two 40-day-old quails, the initial $\delta^{15}N$ value was high and decreased as they grew, eventually

reaching equilibrium around the similar value of the 120 and 160-day-old quails of the same

group.

3.4 Trophic discrimination factors (TDF) of the quail eye lens and primary feather

We estimated the trophic discrimination factors (TDFs) (i.e., the offset) between each food and

tissue. The offset of $\delta^{13}C$ values between the eye lens ($-23.9‰$ for the constant value of the

control group) and the C3 food ($\delta^{13}C_{\text{lens-diet}}$) was estimated to be $2.5‰$ (Supplementary Table

S1). The offset of $\delta^{13}C$ values between the eye lens ($-17.6‰$ for the constant value of the

treatment group) and the C4 food ($\delta^{13}C_{\text{lens-diet}}$) was estimated to be $-0.5‰$ (Supplementary

Table S1). The offset of $\delta^{15}N$ values between the eye lens and the C3 food ($\delta^{15}N_{\text{lens-diet}}$) was

estimated to be $-0.1‰$ ($4.5‰$, Supplementary Table S1). The offset of $\delta^{15}N$ values between

the eye lens and the C4 food ($\delta^{15}N_{\text{lens-diet}}$) was estimated to be $1.8‰$ ($3.9‰$ for constant value

of treatment group, Supplementary Table S1).

The offset of the $\delta^{13}C$ values between the feather ($-23.6‰$ for the constant value of

the control group, Fig. 4 (a)) and the C3 food ($\delta^{13}C_{\text{feather-diet}}$) was estimated to be $2.8‰$

(Supplementary Table S1). The offset of the $\delta^{13}C$ values between the feather ($16.7‰$ for the

constant value of the treatment group, Fig. 4 (b)−(d)) and the C4 food ($\delta^{13}C_{\text{feather-diet}}$) was

estimated to be $0.4‰$. The offset of the $\delta^{15}N$ values between the feather and the C3 food

($\delta^{15}N_{\text{feather-diet}}$) was estimated to be $-0.7‰$ ($3.9‰$ for the control group, Supplementary Fig. S4

(a)). The offset of $\delta^{15}N$ values between the feather and the C4 food ($\delta^{13}C_{\text{feather-diet}}$) was estimated

to be $1.3‰$ ($3.4‰$ for the constant value of the treatment group, Supplementary Fig. S4 (b)

−(d), Supplementary Table S1).

4 Discussion

In this study, we found that the early-life environment could be reconstructed using the eye lens from quails. The high $\delta^{13}C$ values observed in the center of the eye lens for all individuals may reflect the isotope ratios of yolk-derived nutrition, specifically the C4 food fed to their mother (Fig. 2). Subsequently, in the control group of quail that was continuously fed the C3 food, the $\delta^{13}C$ values in the eye lens transitioned to lower values and equilibrated. In the quails of the treatment groups, the transition to lower C3 food-derived $\delta^{13}C$ was followed by a shift to higher C4 food-derived values, and this U-shaped shift was preserved in the quails at all switching times (10, 15, 20, and 40 days, Fig. 2). Thus, we confirmed the diet shifts up to 40 days, and hence the quail eye lens should retain isotopic information for at least 40 days after hatching. Furthermore, the decreasing $\delta^{13}C$ shift from the center to the outer lens was consistent within the control group from 40- to 200-day-old quails (Fig. 2), indicating that the early-life environment is conserved for at least 200 days.

To estimate the timing at which the most recent isotopic information is obtained from the avian eye lens, it is necessary to understand their growth pattern. No significant increase in lens radius was found in individuals after 40 days of age, which is supported by the previous studies in which the avian eye lens did not enlarge after 50 days of age (e.g., house sparrows, *Passer domesticus*, Augusteyn, 2014). Even if lens growth does not occur after its size reaches

the asymptote, lens compaction allows for the further preservation of isotopic records

(Augusteyn 2008). In our study, the PCF model of both $\delta^{13}C$ and $\delta^{15}N$ datasets indicated the

presence of lens compaction (Supplementary Fig. S3), but a previous study reported that the

absence of lens compaction in chickens (*Gallus domesticus*) has also been observed (Augusteyn

2014). In this study, the longest time to diet shift was 40 days; thus, to determine how long the

isotopic record can be stored, it is necessary to conduct experiments with a longer time to diet

shift.

368   We also revealed differences in the isotope ratios between feathers, which have been

the most commonly used tissue for isotopic studies in birds (Hobson and Wassenaar 2018), and

the eye lens, which we propose as a new target for retrospective isotope analysis. Unlike eye

lenses, feathers could not recover embryonic $\delta^{13}C$ records (Fig. 4), thereby indicating that lenses

can recreate the early-life environment, which cannot be obtained from feathers. In addition, in

feathers, the transitions from C3 to C4 foods were reflected only in the isotope ratios of quails

in the 40-day-old treatment group but not in the 120- and 160-day-old groups, indicating that

their early-life environment was lost due to molting. This can be a major obstacle in inferring

the isotope ratios (Hobson and Wassenaar 2018). On the other hand, feathers provide isotopic

information during feather development, which can be directly linked to information on feather

growth. Thus, we showed that the sequential isotope analysis of avian eye lenses provides a

new method to complement information on their prenatal phase that is challenging to obtain

from feathers.

Although the feeding trial of the current study was designed primarily for $\delta^{13}C$, the

results of $\delta^{15}N$ generally supported those of $\delta^{13}C$ (Fig. 3). The higher $\delta^{15}N$ values near the center

of the eye lens can be explained by the trophic enrichment associated with the consumption of

egg yolk-derived nutrients (Hobson1995). When purposely analyzing $\delta^{15}N$ of the eye lens to

understand the trophic ecology via the relationship between animals and their food, this effect

of trophic enrichment should be considered. The continuous decline reflecting the diet shift

from C3 food to C4 food (Fig. 3) indicated that the avian eye lens also preserves a chronological

record of $\delta^{15}N$ from the prenatal to early postnatal periods. For the $\delta^{15}N$ of feathers, there was

a large variation in the isotope ratios at equilibrium, stored period, and the direction of isotopic

ratio among individuals (Supplementary Fig. S4). These irregularities may be linked to feather

wear, such as mechanical abrasion and UV light causing a loss of melanin (Johnson 2019), and

the source of amino acids, such as endogenous reserves and fed proteins (Cherel et al. 2005),

but they do not affect the conclusions obtained for $\delta^{13}C$.

The TDFs between the food and tissue in the target organisms are important

information when using isotope analysis in ecological studies, which can help predict the

isotopic composition of the diets (Hobson and Clark 1992). Because this factor is tissue-

dependent, it is worthwhile to estimate the TDFs of eye lenses and feathers to reconstruct the

history of diet consumption in a comparable form (Martinez del Rio et al., 2009). The TDFs in

quail eye lens and feathers determined in this study (C3 food: $\delta^{13}C_{lens-diet}$ = 2.5‰, $\delta^{13}C_{feather-diet}$

= 2.8 ‰, Supplementary Table S1) were similar to those of feathers from a total of 25 bird

species ($\delta^{13}C_{feather-diet}$ = 2.2‰, Swan et al. 2019), but higher than those of whole blood (0.4‰,

Swan et al. 2019). The TDFs based on $\delta^{15}N$ in the current study (C3 food: $\delta^{15}N_{lens-diet}$ = −0.1‰,

$\delta^{13}C_{feather-diet}$ = −0.7‰, Supplementary Table S1) were estimated to have a lower value than

those of previous studies ($\delta^{15}N_{feather-diet}$ = 4.1‰, Swan et al. 2019, Supplementary Table S1).

This lower TDF estimation may partly be related to the difference in the reflection speed of the

isotope ratio of the components that are used as a carbon and nitrogen resources via exogenous

(mixed feed) and endogenous (stored nutrients) sources depending on their developmental stage

(Ayliffe et al. 2004). Furthermore, the mixed feed used in the current study included multiple

food sources; therefore, variation in the chemical characteristics (carbon chain length, number

of double bonds, degree of unsaturation) between the food components could affect digestibility

and energy use in quails (Valentim et al. 2023). These are unavoidable and an intrinsic problem

in the TDF estimation using mixed feed. Future research should consider the developmental

stages of birds and the turnover rate of their dietary sources (West et al. 2004; Ayliffe et al.

2004).

5 Conclusions

In the current study, we developed a methodology for the isotope analysis of avian eye lenses

and validated its retrospective use for the reconstruction of early-life environments. Through

diet-shift experiment, we were able to reconstruct the early-life history deposited in the avian

eye lens and develop a new formula. Our method of dehydration by ethanol and dissection using

fine-tipped tweezers is essential to prevent the disintegration of the avian lens. This method can

be applied not only to pure science studies but also to applied studies for the reconstruction of

the early-life environment for declining species and the implementation of conservation efforts.

Although there were large individual differences in the isotope ratios within each treatment

group, the control group converged to a single asymptotic value, indicating its reproducibility.

Generally, hatchlings (both in precocial and altricial birds) remain in their birthplace, are not

expected to disperse within a few weeks, and thus have sufficient time to store their early-life

environment within their eye lens, thus making a birds' eye lens as a valuable tool to reconstruct

their early-life environment. Further research is required to improve the accuracy and

reproducibility of reconstructing the early-life environments left in the bird eye lens.

Abbreviations

Air: Atmospheric nitrogen; OFZ: Organelle-Free Zone; PCF: Piecewise Constant Function;

TDFs: Trophic Discrimination Factors; VPDB: Vienna Pee Dee Belemnite

Declarations

Availability of data and material

The datasets supporting the conclusions of this article are available in the odf.io repository after

acceptance.

Competing interests

The authors declare no competing interests.

Funding

This work was supported by grants from JSPS KAKENHI (grant numbers 21H03579 for JM,

21H04784 and 20KK0165 for IT and 23KJ2160 for EA).

Authors' contributions

JM and EA contributed equally to this work and are co-first authors. JM and TA proposed the

topic and conceived and designed the study. TG, and HI performed the feeding trial. JM, EA,

and IT developed the method for eye lens dissection and dehydration. EA, CY, JM, and IT

performed isotope analysis. AGR developed the formula for data analysis. JM and EA wrote

the initial draft of the manuscript. JM, EA, MH, AGR and IT collaborated on the data analysis

and its interpretation and the construction of the manuscript. All authors have read and approved

the final manuscript.

Authors' information

EAH was employed by a grant from JSPS KAKENHI 21H04784 for IT to conduct this research.

Acknowledgements

Thanks to the laboratory members of the Research Institute for Humanity and Nature for their

assistance in our experiments. EA gratefully acknowledges the JSPS KAKENHI 20KK0165 for

the opportunity to attend the IsoEcol 2024 held at the University of New Brunswick, Canada.

Ethics statement

This study was approved by the Experimental Animal Committee of the Obihiro University of

Agriculture and Veterinary Medicine (authorization number 19-32).

References

Augusteyn RC (2008) Growth of the lens: in vitro observations. Clin Exp Optom 91:226-239

Augusteyn RC (2014) Growth of the eye lens: I. Weight accumulation in multiple species. Mol

Vis 20:410-426

Ayliffe LK, Cerling TE, Robinson T, West AG, Sponheimer M, Passey BH, Hammer J, Roeder

B, Dearing MD, Ehleringer JR (2004) Turnover of carbon isotopes in tail hair and breath

CO_2 of horses fed an isotopically varied diet. Oecologia 139:11-22

Barrow LM, Bjorndal KA, Reich KJ (2008) Effects of preservation method on stable carbon

and nitrogen isotope values. Physiol Biochem Zool 81:688-693

Carlisle AB, Goldman KJ, Litvin SY, Madigan DJ, Bigman JS, Swithenbank AM, Kline TC,

Block BA (2015) Stable isotope analysis of vertebrae reveals ontogenetic changes in habitat

in an endothermic pelagic shark. P Roy Soc B Biol Sci 282:20141446. doi:

10.1098/rspb.2014.1446 .

Carvalho PC, Ronconi RA, Bugoni L, Davoren GK (2022) Moult chronology and strategies of

sympatric Great (*Ardenna gravis*) and Sooty (*A. grisea*) Shearwaters based on stable isotope

analysis. Ibis 164:998-1012

Cherel Y, Hobson KA, Bailleul F, Groscolas R (2005) Nutrition, physiology, and stable

isotopes: new information from fasting and molting penguins. Ecology 86:2881-2888

Chua KW, Liew JH, Shin KH, Yeo DC (2020) Effects of ethanol preservation and formalin

fixation on amino acid stable isotope analysis (δ^{13}C and δ^{15}N) and its ecological applications.

Limnol Oceanogra-Meth 18:77-88

Fitzgerald BW (2018) Using Hawkeye from the Avengers to communicate on the eye. Adv

Physiol Educ 42:90-98

Goto T, Konno S, Konno M (2023) Establishment of Wild-Derived Strains of Japanese Quail

(*Coturnix japonica*) in Field and Laboratory Experiments. Biology 12:1080. doi:

10.3390/biology12081080

Gutierrez-Exposito C, Ramirez F, Afan I, Forero MG, Hobson KA (2015) Toward a deuterium

feather isoscape for sub-Saharan Africa: Progress, challenges and the path ahead. PloS one

10:e0135938. doi:10.1371/journal.pone.0135938

Hamel S, Gaillard JM, Festa-Bianchet M, Côté SD (2009) Individual quality, early-life

conditions, and reproductive success in contrasted populations of large herbivores. Ecology

90:1981-1995

Hayward AD, Rickard IJ, Lummaa V (2013) Influence of early-life nutrition on mortality and

reproductive success during a subsequent famine in a preindustrial population. Proc Natl

Acad Sci 110:13886-13891

Hobson KA (1995) Reconstructing avian diets using stable-carbon and nitrogen isotope analysis

of egg components: patterns of isotopic fractionation and turnover. Condor 97:752-762

Hobson KA, Clark RG (1992) Assessing avian diets using stable isotopes I: turnover of ^{13}C in

tissues. Condor 94:181-188

Hobson KA, Møller AP, Van Wilgenburg SL (2012) A multi-isotope (δ^{13}C, δ^{15}N, δ^2H) approach

to connecting European breeding and African wintering populations of barn swallow

(*Hirundo rustica*). Anim Migr 1:8-22

Hobson KA, Wassenaar LI (eds) (2018) Tracking animal migration with stable isotopes.

Academic Press, Cambridge

Inger R, Bearhop S (2008) Applications of stable isotope analyses to avian

ecology. Ibis 150:447-461

Johnson L (2019) Assessing the effect of feather wear on carbon and nitrogen stable isotope

ratios, and the use of stable isotopes to determine predator diets in the Namibian Islands

marine protected area. Master Thesis, University of Cape Town,

http://hdl.handle.net/11427/31248

Martínez del Rio C, Wolf N, Carleton SA, Gannes LZ (2009) Isotopic ecology ten years after a

call for more laboratory experiments. Biol Rev 84:91-111

Monaghan P (2008) Early growth conditions, phenotypic development and environmental

change. Philosophical Transactions of the Royal Society B: Biological Sciences, 363:1635-

1645

Mori H, Takaya M, Nishimura K, Goto T (2020) Breed and feed affect amino acid contents of

egg yolk and eggshell color in chickens. Poult Sci 99:172-178

Nussey DH, Kruuk LE, Morris A, Clutton-Brock TH (2007) Environmental conditions in early

life influence ageing rates in a wild population of red deer. Curr Biol 17:R1000-R1001

Paritte JM, Kelly JF (2009) Effect of cleaning regime on stable-isotope ratios of feathers in

Japanese Quail (*Coturnix japonica*). Auk 126:165-174

Passey BH, Cerling TE (2002) Tooth enamel mineralization in ungulates: implications for

recovering a primary isotopic time-series. Geochim Cosmochim Ac 66:3225-3234

Peebles EB, Hollander DJ (2020) Combining isoscapes with tissue-specific isotope records to

recreate the geographic histories of fish. Scenarios and Responses to Future Deep Oil Spills:

Fighting the Next War, Springer Nature, Switerland, Cham, p203-218

R Core Team (2021) R: A language and environment for statistical computing. R Foundation

for Statistical Computing, Vienna, Austria. URL: https://www.R-project.org/.

Sergio F, Tavecchia G, Blas J, Tanferna A, Hiraldo F, Korpimaki E, Beissinger SR (2022)

Hardship at birth alters the impact of climate change on a long-lived predator. Nat Commun

13:5517. doi: 10.1038/s41467-022-33011-7

Shestopalov VI, Bassnett S (2000) Expression of autofluorescent proteins reveals a novel

protein permeable pathway between cells in the lens core. J Cell Sci 113:1913-1921

Simpson A, Bateman O, Driessen H, Lindley P, Moss D, Mylvaganam S, Narebor E, Slingsby

C (1994) The structure of avian eye lens δ-crystallin reveals a new fold for a superfamily of

oligomeric enzymes. Nat Struct Biol 1:724-734

Sivak JG (2004) Through the lens clearly: phylogeny and development: the Proctor

lecture. Invest Ophthalmol Vis Sci 45:740-747

Swan GJ, Bearhop S, Redpath SM, Silk MJ, Goodwin CE, Inger R, McDonald RA (2020)

Evaluating Bayesian stable isotope mixing models of wild animal diet and the effects of

trophic discrimination factors and informative priors. Methods Ecol Evol 11:139-149

Tayasu I, Hirasawa R, Ogawa NO, Ohkouchi O, Yamada K (2011) New organic reference

materials for carbon-and nitrogen-stable isotope ratio measurements provided by Center for

Ecological Research, Kyoto University, and Institute of Biogeosciences, Japan Agency for

Marine-Earth Science and Technology. Limnology 12:261-266

Tzadik OE, Curtis JS, Granneman JE, Kurth BN, Pusack TJ, Wallace AA, Hollander DJ,

Peebles EB, Stallings CD (2017) Chemical archives in fishes beyond otoliths: A review on

the use of other body parts as chronological recorders of microchemical constituents for

expanding interpretations of environmental, ecological, and life-history changes. Limnol

Oceanogr Methods 15:238-263

Underwood H, Steele CT, Zivkovic B (1999) Effects of fasting on the circadian body

temperature rhythm of Japanese quail. Physiol Behav 66:137-143

Valentim JK, Garcia RG, Burbarelli MFDC, Caldara FR, Komiyama CM, Serpa FC, Zenella J,

Heiss VARDC, Polycarpo GDV, Albino LFT (2023) Energy values and metabolizability of

lipid sources of plant and animal origin in the diet of Japanese quail. Rev Bras Zootec

52:e20220105. doi: 10.37496/rbz5220220105

Vecchio JL, Peebles EB (2020) Spawning origins and ontogenetic movements for demersal

fishes: An approach using eye-lens stable isotopes. Estuar Coast Shelf Sci 246:107047

West AG, Ayliffe LK, Cerling TE, Robinson TF, Karren B, Dearing MD, Ehleringer JR (2004)

Short-term diet changes revealed using stable carbon isotopes in horse tail-hair. Funct Ecol

18:616-624

Wolf N, Newsome SD, Fogel ML, Del Rio CM (2013) The relationship between drinking water

and the hydrogen and oxygen stable isotope values of tissues in Japanese Quail (*Cortunix*

japonica). Auk 130:323-330

Figure legends

Figure 1. Diet-shift experiment design. Japanese quail eggs were collected from parent birds that were raised on C4 food. The hatched chicks were fed C3 food during the first 10 days, and they were then divided into control group ($n = 6$) and treatment (total $n = 25$) groups. The control group continued on the C3 food until the end of the experiment (i.e., dissection date at days 17, 40, 80, 120, 160, and 200). The treatment group switched to the C4 food at specific time points: 10 days (T10, $n = 6$), 15 days (T15, $n = 10$), 20 days (T20, $n = 7$), and 40 days (T40, $n = 2$) after the start of rearing. The final rearing period (i.e., dissection date) for the treatment groups included time points at 40, 80, 120, 160, 168, and 200 days.

Figure 2. ^{13}C values of quail eye lenses. (a) Control group ($n = 5$); (b) T10 group ($n = 6$); (c) T15 group ($n = 10$); (d)T20 group ($n = 7$); and (e) T40 group ($n = 2$; 168 days). The lamina from each eye lens (right or left) was included.

Figure 3. δ^{15}N values of quail eye lenses: (a) Control group ($n = 5$); (b) T10 group ($n = 6$); (c) T15 group ($n = 10$); (d) T20 group ($n = 7$); and (e) T40 group ($n = 2$; 168 days). The lens sections from each of eye lens (right or left) were included.

Figure 4. δ^{13}C values of quail feather sections. (a) Control group ($n = 3$); (b) T10 group ($n = 3$); (c) T15 group ($n = 3$); and (d) T20 group ($n = 3$).

593   Fig.1

595   Fig.2

597   Fig.3

599   Fig.4