The influence of sleep disruption on learning and memory in fish

Sowersby W[1]., Kobayahsi T[1]., Awata, S[1]., Sogawa S[1]., Kohda M[1].

1.Graduate School of Science, Osaka Metropolitan University, Osaka 558-8585, Japan

ABSTRACT

Sleep is ubiquitous across animal taxa. Strong evolutionary pressures have conserved sleep
over the evolutionary history of animals, yet our understanding of the functions of sleep still
largely derives from mammals and select laboratory models. Sleep is considered to play an
important role in mental processes, including learning and memory consolidation, but how
widespread this relationship occurs across taxa remains unclear. Here, we test the impact of
sleep disruption on the ability of the cleaner fish (*Labroides dimidiatus*) to both learn and
then remember a novel cognitive task. We found a significant negative relationship between
sleep disruption and the ability to learn a food reward choice system. Specifically, we found
that fish in a disturbed sleep treatment took significantly longer and made more incorrect
decisions when finding the food reward, compared to individuals in a non-disturbed/normal
sleep treatment. In contrast, the differences between the two treatment groups were non-
significant when fish where tasked with remembering the food reward several days later.
Our results demonstrate a negative impact of sleep disruption on performance in a cognitive
challenging task and that the effects were strongest when fish were first exposed to the
challenge. Importantly, we show that the association between sleep and mental processes,
such as learning, may be widespread across vertebrate taxa and potentially have an early
origin in the evolutionary history of animals.

Introduction

Sleep is a ubiquitous trait that has been conserved across animal taxa, with positive links to
immune function, neurogenesis, wellbeing, and cognitive performance (Tononi & Cirelli
2012; Czeisler 2015; Elvsåshagen et al. 2015). There are also costs associated with sleeping,
such as potentially being more vulnerable to predators and reducing the time available to
secure food and mates (Lima et al. 2005; Joiner 2016; Tworkowski & Lesku 2019).
Nevertheless, there is currently no evidence to suggest that animals can completely
renounce sleep without detrimental consequences, including death (Van Dongen et al.
2003; Cirelli & Tononi, 2008). Indeed, a commonly observed response to sleep disruption, is
longer and deeper sleep, indicating that sleep is maintained homeostatically (Tobler 2011;
Tobler and Borbély 1985; Nath et al. 2017; Yokogawa et al. 2007). Despite seemingly strong
selective pressure to conserve sleep across animal evolutionary history, how sleep has
coevolved with other traits and processes is still not often considered in evolutionary
research (Cirelli and Tononi 2008; Mignot 2008; Roth, Rattenborg, and Pravosudov 2010;
Rattenborg et al. 2017).

One exciting but still controversial proposed function of sleep is the important role it plays
in brain processes. For example, compelling evidence points to a strong positive link
between sleep, learning and memory consolidation (Huber et al. 2004; Walker and Stickgold

2004b; Peigneux et al. 2001; Smith 2001; Walker and Stickgold 2006; Anafi, Kayser, and
Raizen 2019; Siegel 2001; but see Vertes 2004; Rattenborg et al. 2017), neural maintenance
(Kavanau 1996; Tononi and Cirelli 2006) and cognitive ability (Scullin and Bliwise 2015;
Wagner et al. 2004). In animals, learning and memory are critical cognitive functions for a
range of tasks, including foraging or food-caching (Shettleworth 1995; Sherry and Hoshooley
2010), navigation and territory maintenance (Shettleworth 1998) and for recalling positive
and negative social interactions (Mateo and Johnston 2000). The link between sleep and
memory consolidation has been demonstrated in humans (Born, Rasch, and Gais 2006;
Diekelmann and Born 2010; Huber et al. 2004) and other mammals (Capellini et al. 2009;
Smith and Kelly 1988; Ribeiro et al. 2004; Walker and Stickgold 2004b) and to a lesser extent
in birds (Jackson et al. 2008; Derégnaucourt et al. 2005). To date, research investigating this
relationship has largely been restricted to these few taxa (Keene and Duboue 2018), despite
some evidence indicating the importance of sleep for memory formation in some
invertebrates (Beyaert, Greggers, and Menzel 2012; Krishnan, Noakes, and Lyons 2016) and
fishes (Pinheiro-da-Silva, Tran, and Luchiari 2018; Rawashdeh et al. 2007).

The function and in places the structure of the vertebrate brain, especially the forebrain,
appear to have been largely conserved throughout the evolution of vertebrate taxa (Vargas
et al. 2006; Xie & Dorsky 2017). For example, the mammalian and avian hippocampus, along
with the homologous pallial areas of the reptile and fish brain, share a central role in
supporting and encoding spatial information (Rodriguez et al. 2002; Portavella et al. 2002).
Vargas et al. (2006) demonstrated that lesions to the lateral pallium of goldfish (*Carassius*
auratus) can impair the encoding of geometric spatial information, comparable to the effect
of lesions on the hippocampus in mammals and birds. Interestingly, comparable patterns of
sleep have also been observed across these taxonomic groups, including in fishes, which
display both characteristic sleep behaviours (in cartilaginous fishes: Kelly et al. 2020, 2022)
and deep slow-wave and rapid eye like movements (REM; in bony fishes: Leung et al. 2019;
M. Kohda et al. unpublished data). Evidence suggesting an association between learning and
active memory consolidation with sleep in non-mammalian species would imply that sleep
has played an important role in these mental processes across vertebrate evolutionary
history (Vorster and Born 2015). Any link between sleep, learning and memory in fishes or
other non-mammalian taxa is however not well established (Pinheiro-da-Silva, Tran, and
Luchiari 2018; Rawashdeh et al. 2007) and this large and diverse group of vertebrates has
largely been neglected from studies on sleep (Campbell and Tobler 1984).

Here, we investigate the impact of sleep disruption on learning ability and subsequent
memory consolidation in the bluestreak cleaner wrasse (cleaner fish; *Labroides dimidiatus*).
This fish species has a wide distribution on coral reefs across the Indo-Pacific, where in a
mutualistic relationship, it eats parasites and dead tissue off "client" species. Hence, this
highly social fish, regularly engages in cognitively challenging tasks, such as recalling both
favourable and antagonistic interactions with both conspecifics and numerous client
individuals from a range of different species. Cleaner fish are highly amenable to captive
experimental conditions and have demonstrated several complex cognitive abilities, some
which may be impacted by sleep quality (e.g. Bshary & Brown 2014; Kohda et al. 2019,
2023). We predict that when subjected to interrupted sleep schedules, cleaner fish will
demonstrate poorer performance in novel cognitively challenging tasks, in comparison to
fish that have experienced uninterrupted, normal sleep regimes. We also predict that
cleaner fish subjected to disruptions in normal sleep regimes will take longer to rouse each

morning and react slower to the presence of food, compared to fish that have experienced
uninterrupted sleep.
90
91
92 **Methods**
93

Animal husbandry

Cleaner fish (n=12) were purchased from reputable tropical fish specialists and were
individually housed (aquaria 40L) under laboratory conditions (14:10 hr light regime; 27
degrees C) at the Department of Biology, Osaka Metropolitan University, Japan. All aquaria
were furnished with coral sand substrate and a small opaque piece of PVC pipe to provide
shelter. Water changes and water parameter checks (including salinity) were conducted
regularly. Fish were fed daily with commercially bought shrimp (Pandalidae), mashed into a
paste and suspended in the tank on a grey plastic plate. A pilot trial of the experimental set-
up took place in early 2020, with the main experiment then conducted between September
and October 2023. All experiments adhered to the animal ethics guidelines of Osaka
Metropolitan University.

Sleep Treatment

Fish were moved into their treatment tanks for a 48-hour acclimation period prior to the
start of the experiment. Treatment tanks were identical to housing aquaria, except
treatment tanks were divided into two by a clear plastic barrier with a door and had a lamp
overhead (Figure 1). Fish were placed into the back half of the tank and were provided with
food in the front half of the tank, once per day. Specifically, shrimp mashed into a paste was
put on a grey plastic square which was hung into the front half of the tank. When the door
in the partition was raised the fish could enter the front half of the tank and became
accustomed to accessing that half of the tank via the door to feed. After three hours the
grey plastic square was removed, and fish were gently ushered back into the second half of
the tank if they had not already returned voluntarily.

After the acclimation period, half of the fish (n = 6) were assigned to the interrupted sleep
treatment and the remaining half to the non-interrupted sleep treatment (n = 6). Fish in the
non-interrupted sleep treatment were kept in complete darkness for a period of 10 hours
(22:00-8:00) each night, a period which corresponds to the typical night length in the
tropical regions where these fish naturally occur. In contrast, fish in the interrupted sleep
treatment were exposed to a light that came on above the tank every two hours, for a
period of one hour (controlled by a timer), meaning that fish in this treatment were exposed
to less periods of darkness per night, broken into two-hour blocks. Using light to supress
sleep has successfully been used in zebrafish research (Yokogawa et al. 2007). Moreover,
the opaque pipe shelter was removed in the interrupted sleep treatment and replaced with
a transparent plastic pipe of identical size, which would provide the fish with a secure place
to shelter, but not offer refuge from the light above.

To prevent fish in the non-interrupted sleep treatment from being disturbed by light during
the night, fish in the sleep interrupted treatment were isolated at night via a black-out
sheet. Camcorders (SONY Handycam HDR-CX480) positioned in front of each tank (n = 4)
were quietly switched on 10 minutes prior to the morning light coming on (at 8:00). The
room was blacked out to prevent natural light entering, and researchers entered using a dim
red light. Fish were recorded for ~20 minutes after the morning light automatically switched
on at 8.00am, to capture the time it took each fish to wake up (i.e. when they left the
shelter and started swimming), and their initial activity behaviour (i.e. time spent moving
and distance travelled). This sleep treatment occurred every night for four nights during the
learning phase of the experiment. During the subsequent rest and memory phase of this
experiment all fish were exposed to darkness for the normal 10-hour period and were
provided with an opaque tube for shelter. A subset of the fish were recorded during the
night, confirming that their sleep was disturbed by the light treatment.

Food reward choice: Learning phase

We presented fish with a food reward choice at ~9.00am each morning, for three days,
beginning from the first morning after the night of the first sleep treatment. Each fish was
presented with a row of four plastic squares, each a different colour (red, green, black and
yellow) suspended in a row in the front half of the tank (Figure 1). Shrimp paste had been
placed on the back of only the red square, so fish were tasked with finding the food on the
back of this square. Each morning for the three days, fish were presented with the same
four coloured squares, but with the order of the squares in the row different each time.
Once the squares were in place the door separating the front and back halves of the tank
was lifted and we recorded the following, 1). time to exit the door, 2). number of incorrect
choices, 3). time to find food on the back of the red square. We considered an incorrect
choice to be any deliberate touching of a square with the mouth or head that was not on
the backside of the red square. After three mornings, fish in the sleep interrupted treatment
were exposed to a final night of disrupted sleep before all fish entered a six-day rest period.

Food reward choice: Memory phase

After the rest period, fish were again exposed to the same food reward choice experiment
for another three days. Like the rest period, no fish were exposed to the non-interrupted
sleep treatment. This experiment tested the ability of the cleaner wrasse to recall the
correct coloured square to locate the food reward. As previously, we recorded, 1) time to
exit the door, 2) number of incorrect choices, 3) time to find food on the back of the red
square. After the third day each fish was weighed and photographed and placed back in its
home tank.

Figure 1. Set-up for the food choice reward experiment.

Statistical Analysis

We performed separated repeated measures ANOVA to test the effect of sleep disruption
on wake time, time to exit door, number of incorrect choices and time to correct choice. In
each model we included the variables sleep treatment (disrupted or non-disrupted) and day
(day 1, 2 or 3) with experimental tank added as a random effect. All statistical analyses were
conducted in R v4.2.1 (R Core Team, 2022).

Results

Impact of sleep disruption on ability to learn cognitive task

In contrast to our first prediction, we did not observe any significant difference in time to
first movement between fish in the disturbed and non-disturbed sleep treatments (rousing
time; df=1, F=0.018, p=0.896) across the four mornings after sleep disturbance. On average,
fish in the disturbed sleep treatment first moved from their shelter after 197 seconds and
non-disturbed treatment fish 220 seconds once the light automatically turned on at 8.00am.

Non-disturbed sleep fish did exit the door quicker than disturbed fish, however this
difference was not significant (df=1, F=1.552, p=0.241).

In line with our second prediction, we did find a significant negative effect of sleep
disruption on the ability to learn the correct location of the food reward (df=1, F=5.5972,
p=0.0346. Figure 2, supp table 1). Specifically, fish in the sleep disturbed treatment
performed on average significantly more incorrect choices before finding the food reward
(5.65 incorrect choices) compared to fish in the non-disturbed sleep treatment (2.66

incorrect choices). As the days progressed, the mean number of incorrect choices by both
treatment groups declined (Fig 2).

Similarly, we found a significant negative effect of sleep disruption on the time to find the
correct food reward square (df=1, F=5.117, p=0.0472. Figure 3, supp table 2). Sleep
disturbed fish took on average 794 seconds to find the food reward, while non-disturbed
fish took 51.5 seconds. Again, fish in both groups improved over time and could find the
food reward in a similar time by the third day.

Figure 2. The mean number of incorrect decisions made by cleaner fish in two different sleep
treatments (red: sleep disrupted; blue: sleep not disturbed) to find a food reward during the learning
phase of the experiment. Mean incorrect choices per day; day 1: sleep disturbed fish 11, non-
disturbed 3; day 2: sleep disturbed fish 3.3, non-disturbed 2.5; day 3: sleep disturbed fish 3.5, non-
disturbed 2.5.
210
211
212

214 **Figure 3.** The mean time taken to find the food reward by cleaner fish in two different sleep
215 treatments (red: sleep disrupted; blue: sleep not disturbed) during the learning phase of the
216 experiment. Mean time per day; day 1: sleep disturbed fish 2278 sec, non-disturbed 120 sec; day 2:
217 sleep disturbed fish 74.5 sec, non-disturbed 17.7 sec; day 3: sleep disturbed fish 29.7 sec, non-
218 disturbed 16.8 sec.
219
220

Impact of past sleep disruption on the ability to remember a cognitive task

We again found no significant difference in mean time to exit the door between the two
treatments (df=1, F= 0.581, P=0.465).

In contrast to part of our second prediction, after the rest phase, we did not find a
significant difference in the number of incorrect choices made before finding the food
reward (df=1, F=1.937, p=0.197. Figure 4, supp table 3). Individuals that had been sleep
disrupted during the previous learning phase performed on average 2.4 incorrect choices,
while non-disturbed sleep individuals performed on average 1.5 incorrect choices.

Likewise, we found no significant difference in the time to find the food reward (df=1,
F=2.549, p=0.145. Figure 5, supp table 4). Individuals that had been sleep disrupted during
the previous learning phase took on average 5.6 sec, whereas non-disturbed sleep
individuals took on average 2.6 sec to find the food reward.

In both groups, individuals performed better at the food reward task in the memory phase
compared to the learning phase. Generally, performances continued to get better as the
memory phase progressed, however the pattern was not as clear as the continued
improvement in completing the task in the learning phase.

Figure 4. The mean number of incorrect decisions made by cleaner fish in two different sleep treatments (red: sleep disrupted; blue: sleep not disturbed) to find a food reward during the memory phase of the experiment. Mean incorrect choices per day; day 1: sleep disturbed fish 2.3, non-disturbed 1.4; day 2: sleep disturbed fish 2.8, non-disturbed 1.8; day 3: sleep disturbed fish 2, non-disturbed 1.3.

Figure 5. The mean time taken to find the food reward by cleaner fish in two different sleep treatments (red: sleep disrupted; blue: sleep not disturbed) during the memory phase of the experiment. Mean time per day; day 1: sleep disturbed fish 13.8 sec, non-disturbed 7.2 sec; day 2: sleep disturbed fish 21.2 sec, non-disturbed 8.8 sec; day 3: sleep disturbed fish 8.8 sec, non-disturbed 7.8 sec.

Discussion

We clearly demonstrate that disruptions to a normal sleep pattern can negatively affect how quickly and accurately an animal performs a novel cognitive task. We used as our model species the cleaner fish, to provide further evidence of an important function of sleep in non-mammalian and non-avian taxa. Despite retaining small mean differences overtime, we did not observe any significant sleep treatment differences in ability to perform the cognitive task during the memory phase of the experiment. Indeed, the ability to perform the task continued to improve in both treatment groups throughout both the learning phase and to a lesser extent the memory phase of the experiment. We demonstrate that any negative effects of sleep disruption could be most apparent when first encountering a new cognitive challenge, potentially because fish are tired and consequentially unmotivated. Nevertheless, our results support the claim that sleep plays an important role in mental processes, such as learning, across vertebrate taxa.

Sleep is predicted to maintain optimal brain functioning to support waking cognition (Alhola & Polo-Kantola 2007). The ability to learn is a critical function that allows strategic adaptation to changing environmental demands. Cleaner fish inhabit structurally complex physical environments and dynamic social environments and to be successful they require the ability to learn and to adapt their behavioural responses to changes accordingly. For example, it is estimated that wild cleaner fish remove up to 1200 ectoparasites each day (Grutter,1996) from anywhere between 800 to 3000 client fish (Grutter, 1995; Triki et al.

2018; Wismer et al., 2014). In this context, cleaner fish must learn and remember
information about their clients, including past positive or negative interactions, and
identifying which clients are residents or visitors to their cleaning territories. We know that
cleaner fish are also capable of social learning, with juvenile fish learning strategies to
interact with clients by observing cleaner adults interacting with clients (Truskanov et al.
2020). We also know that cleaner fish are not the only fish species capable of learning, for
instance archer fish (*Toxotes* sp.) learn complex skills from observing other group members
(Schuster et al. 2006), while guppies are capable of learning and anticipating the path of a
mechanical fish (Bierbach et al. 2022) and successfully completing reverse learning choice
tasks (Buechel et al. 2018). It is therefore likely that other fish will also be negatively
impacted by sleep disruption when learning novel tasks.
288
In cleaner fish, the negative effects of sleep disruption were most apparent at the beginning
of the learning phase of the experiment. Our results may suggest differences in the timing of
the performance of operant behaviours, i.e. those that that produce consequences and may
be repeated if they provide a reward (Toates 2012), with sleep-interrupted fish taking longer
to find the food reward on the first day after sleep disruption. An explanation for our finding
could be that fish in the sleep-disrupted treatment are less active or motivated. Indeed, a
similar finding has been observed in Australian magpies, where sleep loss impaired
motivation and cognitive performance (Johnsson et al. 2022). In contrast, sleep disruption
appeared to have less of an effect on reinforcement learning on subsequent days, with the
time taken to find the food reward similar between both sleep treatments on days two and
three of the experiment. Reinforcement learning influences behaviour via either a positive
or negative feedback response and can be somewhat affected by sleep loss (Whitney
et al. 2015; Gerhardsson et al. 2020). We recommend that future experiments investigating
the association between sleep and learning should include a reverse learning element,
which has shown in humans to amplify the impairment caused by sleep loss (Whitney
et al. 2015).
305
We did not find a significant negative effect of sleep disruption on the ability of cleaner fish
to recall the food reward cue after several days post learning phase. In humans, memory
functions comprise three major subprocesses, i.e., encoding, consolidation, and retrieval
(Straube 2012). Research has indicated that sleep after learning can have an important
influence on memory consolidation. Rather we found that the performance of fish in both
sleep treatments increased with time and exposure to the task in both the learning and to a
lesser extent the memory stages of the experiment. Our findings do not rule out an
important role for sleep in memory consolidation in fish, specifically non-declarative
memory, but potentially represents a limitation in our experimental design. For example,
the cognitive task may have been too simple or was repeated too often during the
experimental period. Another possible explanation is that our periods of sleep disruption
were too short to affect the consolidation and retrieval of memories associated with
reinforcement learning. In our experiment fish were still allowed several hours sleep each
night, albeit broken into discreet periods of time by the automated light. It is possible that a
more complex cognitive task, including a reverse learning component, or longer periods of
sleep deprivation each night may have increased cognitive and recall impairment.

The involvement of sleep in memory processing has been tested for at least a century (Rasch & Born 2013; Jenkins & Dallenbach 1924). Yet, sleep research is still overwhelmingly biased towards humans, other mammals, and laboratory model species (Toth & Bhargava 2013; Blumberg et al. 2020). Experimental evidence, has demonstrated behavioural, anatomical, genetic, and pharmacological conservation of sleep between fish (largely in zebrafish) and mammals, suggesting that research in fish can inform our understanding of mammalian sleep (Lee et al. 2022). Despite difficulties finding neuronal sleep signatures in fishes, due to the absence of a conventional neocortex, bony fishes do possess a homologue, the dorsal pallium, and neural signatures of sleep have been observed (Leung et al. 2019). Our evidence that sleep has an impact on mental processing in fish, in combination with the conservation of sleep and the presence of homologous structures in the vertebrate brain, suggests that sleep has played an important role in animal cognitive processes since early in the evolutionary history of vertebrate taxa.

Our study provides further evidence that sleep is functionally important in non-mammalian animals. Unlike previous studies on non-mammal species, we use an ecologically relevant stimulus to test learning ability and memory consolidation/retention using the cleaner fish. We found that the negative impacts of sleep disruption were most apparent initially, likely impacting cognition, activity levels and motivation, which in turn slowed the learning process. Our results do not uncover which aspects of cognition were most affected by sleep loss. However, we speculate that the association between sleep and mental processes is widespread in vertebrates and may have coevolved or facilitated the ability of animals to perform complex cognitive behaviours.

References

Acerbi, A. & Nunn, C.L. 2011. Predation and the Phasing of Sleep: An Evolutionary Individual-Based Model. *Animal Behaviour* 81 (4): 801–11. https://doi.org/10.1016/j.anbehav.2011.01.015.

Anafi, R.C., Kayser, M.S. & Raizen, D.M. 2019. Exploring Phylogeny to Find the Function of Sleep. *Nature Reviews Neuroscience* 20 (2): 109–16. https://doi.org/10.1038/s41583-018-0098-9.

Arble, D.M., Bass, J., Behn, C.D., Butler, M.P., Challet, E., Czeisler, C., Depner, C.M. et al. 2015. Impact of Sleep and Circadian Disruption on Energy Balance and Diabetes: A Summary of Workshop Discussions. *Sleep* 38 (12): 1849–60. https://doi.org/10.5665/sleep.5226.

Backhaus, J., Junghanns, K., Born, J., Hohaus, K., Faasch, F., & Hohagen, F. (2006). Impaired declarative memory consolidation during sleep in patients with primary insomnia: influence of sleep architecture and nocturnal cortisol release. *Biological psychiatry*, *60*(12), 1324-1330.

Beyaert, L., Greggers U., & Menzel, R. 2012. Honeybees Consolidate Navigation Memory during Sleep. *Journal of Experimental Biology* 215 (22): 3981–88. https://doi.org/10.1242/jeb.075499.

Bierbach D, Gómez-Nava L, Francisco FA, Lukas J, Musiolek L, Hafner VV, Landgraf T,
Romanczuk P. & Krause J. 2022. Live fish learn to anticipate the movement of a fish-
like robot. *Bioinspir Biomim.* 17(6).
Born, J., Rasch, B. & Gais, S. 2006. Sleep to Remember. The Neuroscientist: A *Review Journal*
Bringing Neurobiology, Neurology and Psychiatry 12 (5): 410–24.
https://doi.org/10.1177/1073858406292647.
Buechel S.D., Boussard A., Kotrschal A., van der Bijl W. & Kolm N. 2018. Brain size affects
performance in a reversal-learning test. *Proc Biol Sci.* 31;285(1871):20172031
Campbell, S.S., & Tobler, I. 1984. Animal Sleep: A Review of Sleep Duration across
Phylogeny. *Neuroscience & Biobehavioral Reviews* 8 (3): 269–300.
https://doi.org/10.1016/0149-7634(84)90054-X.
Capellini, I., McNamara, P., Preston, B.T., Nunn, C.L. & Barton, R.A. 2009. Does Sleep Play a
Role in Memory Consolidation? A Comparative Test. *PLoS ONE* 4 (2).
https://doi.org/10.1371/journal.pone.0004609.
Cappuccio, F.P., D'Elia, L., Strazzullo, P. & Miller M.A. 2010. Sleep Duration and All-Cause
Mortality: A Systematic Review and Meta-Analysis of Prospective Studies. *Sleep* 33
(5): 585–92. https://doi.org/10.1093/sleep/33.5.585.
Cirelli, C., & Tononi, G. 2008. Is Sleep Essential? *PLOS Biology* 6 (8): e216.
https://doi.org/10.1371/journal.pbio.0060216.
Derégnaucourt, S., Mitra, P.P., Fehér, O., Pytte, C. & Tchernichovski, O. 2005. How Sleep
Affects the Developmental Learning of Bird Song. *Nature* 433 (7027): 710–16.
https://doi.org/10.1038/nature03275.
Diekelmann, S. & Born, J. 2010. The Memory Function of Sleep. *Nature Reviews.*
Neuroscience 11 (2): 114–26. https://doi.org/10.1038/nrn2762.
Dinges, D.F., Pack, F., Williams, K., Gillen, K.A., Powell, J.W., Ott, G.E., Aptowicz, C. & Pack,
A.I. 1997. Cumulative Sleepiness, Mood Disturbance, and Psychomotor Vigilance
Performance Decrements During a Week of Sleep Restricted to 4–5 Hours per Night.
Sleep 20 (4): 267–77. https://doi.org/10.1093/sleep/20.4.267.
Frank, M.G., Waldrop, R.H., Dumoulin, M., Aton, S. & Boal, J.G. 2012. A Preliminary Analysis
of Sleep-Like States in the Cuttlefish Sepia Officinalis. *PLOS ONE* 7 (6): e38125.
https://doi.org/10.1371/journal.pone.0038125.
Gerhardsson, A., Porada, D.K., Lundström, J.N., Axelsson, J., Schwarz, J. Does insufficient
sleep affect how you learn from reward or punishment? Reinforcement learning
after 2 nights of sleep restriction. J Sleep Res,
2020,. https://doi.org/10.1111/jsr.13236
Hairston, Ilana S., Milton T. M. Little, Michael D. Scanlon, Monique T. Barakat, Theo D.
Palmer, Robert M. Sapolsky, and H. Craig Heller. 2005. "Sleep Restriction Suppresses
Neurogenesis Induced by Hippocampus-Dependent Learning." Journal of
Neurophysiology 94 (6): 4224–33. https://doi.org/10.1152/jn.00218.2005.
Hartse, K.M. 1994. "Sleep in Insects and Nonmammalian Vertebrates." In Principles and
Practice of Sleep Medicine, edited by M. H. Kryger, T Roth, and W.C. Dement, 1:95–
104. Philadelphia: Elsevier Saunders.
Huber, Reto, M. Felice Ghilardi, Marcello Massimini, and Giulio Tononi. 2004. "Local Sleep
and Learning." Nature 430 (6995): 78–81. https://doi.org/10.1038/nature02663.
Jackson, Claire, Brian J. McCabe, Alister U. Nicol, Amanda S. Grout, Malcolm W. Brown, and
Gabriel Horn. 2008. "Dynamics of a Memory Trace: Effects of Sleep on

Consolidation." Current Biology 18 (6): 393–400.
https://doi.org/10.1016/j.cub.2008.01.062.

Jenkins J, Dallenbach K. Obliviscence during sleep and waking. *Am J Psychol* 1924; **35**: 605–612.

Johnsson RD, Connelly F, Gaviraghi Mussoi J, Vyssotski AL, Cain KE, Roth TC 2nd, Lesku JA. Sleep loss impairs cognitive performance and alters song output in Australian magpies. Sci Rep. 2022 Apr 22;12(1):6645. doi: 10.1038/s41598-022-10162-7. PMID: 35459249; PMCID: PMC9033856.

Joiner, William J. 2016. "Unraveling the Evolutionary Determinants of Sleep." Current Biology: CB 26 (20): R1073–87. https://doi.org/10.1016/j.cub.2016.08.068.

Kayser, Matthew S., Zhifeng Yue, and Amita Sehgal. 2014. "A Critical Period of Sleep for Development of Courtship Circuitry and Behavior in Drosophila." Science 344 (6181): 269–74. https://doi.org/10.1126/science.1250553.

Keene, Alex C., and Erik R. Duboue. 2018. "The Origins and Evolution of Sleep." Journal of Experimental Biology 221 (11). https://doi.org/10.1242/jeb.159533.

Krishnan, Harini C., Eric J. Noakes, and Lisa C. Lyons. 2016. "Chronic Sleep Deprivation Differentially Affects Short and Long-Term Operant Memory in Aplysia." Neurobiology of Learning and Memory 134 (October): 349–59. https://doi.org/10.1016/j.nlm.2016.08.013.

Lee, D. A., Oikonomou, G., & Prober, D. A. (2022). Large-scale analysis of sleep in Zebrafish. *Bio-protocol*, *12*(3), e4313-e4313.

Lee Kavanau, J. 1996. "Memory, Sleep, and Dynamic Stabilization of Neural Circuitry: Evolutionary Perspectives." Neuroscience & Biobehavioral Reviews 20 (2): 289–311. https://doi.org/10.1016/0149-7634(95)00019-4.

Lesku, John A., and Niels C. Rattenborg. 2014. "Avian Sleep." Current Biology 24 (1): R12–14. https://doi.org/10.1016/j.cub.2013.10.005.

Leung, Louis C., Gordon X. Wang, Romain Madelaine, Gemini Skariah, Koichi Kawakami, Karl Deisseroth, Alexander E. Urban, and Philippe Mourrain. 2019. "Neural Signatures of Sleep in Zebrafish." Nature 571 (7764): 198–204. https://doi.org/10.1038/s41586-019-1336-7.

Lima, Steven L. 1998. "Stress and Decision Making under the Risk of Predation: Recent Developments from Behavioral, Reproductive, and Ecological Perspectives." In Advances in the Study of Behavior, edited by Anders Pape Møller, Manfred Milinski, and Peter J. B. Slater, 27:215–90. Stress and Behavior. Academic Press. https://doi.org/10.1016/S0065-3454(08)60366-6.

Mateo, Jill M., and Robert E. Johnston. 2000. "Retention of Social Recognition after Hibernation in Belding's Ground Squirrels." Animal Behaviour 59 (3): 491–99. https://doi.org/10.1006/anbe.1999.1363.

Mazzotti, Diego Robles, Camila Guindalini, Walter André dos Santos Moraes, Monica Levy Andersen, Maysa Seabra Cendoroglo, Luiz Roberto Ramos, and Sergio Tufik. 2014. "Human Longevity Is Associated with Regular Sleep Patterns, Maintenance of Slow Wave Sleep, and Favorable Lipid Profile." Frontiers in Aging Neuroscience 6. https://doi.org/10.3389/fnagi.2014.00134.

Mignot, Emmanuel. 2008. "Why We Sleep: The Temporal Organization of Recovery." PLOS Biology 6 (4): e106. https://doi.org/10.1371/journal.pbio.0060106.

Nath, Ravi D., Claire N. Bedbrook, Michael J. Abrams, Ty Basinger, Justin S. Bois, David A. Prober, Paul W. Sternberg, Viviana Gradinaru, and Lea Goentoro. 2017. "The Jellyfish

Cassiopea Exhibits a Sleep-like State." Current Biology 27 (19): 2984-2990.e3.
https://doi.org/10.1016/j.cub.2017.08.014.

Peigneux, Philippe, Steven Laureys, Xavier Delbeuck, and Pierre Maquet. 2001. "Sleeping
Brain, Learning Brain. The Role of Sleep for Memory Systems." NeuroReport 12 (18):
A111.

Pinheiro-da-Silva, Jaquelinne, Steven Tran, and Ana Carolina Luchiari. 2018. "Sleep
Deprivation Impairs Cognitive Performance in Zebrafish: A Matter of Fact?"
Behavioural Processes 157 (December): 656–63.
https://doi.org/10.1016/j.beproc.2018.04.004.

Portavella, M, J. P Vargas, B Torres, and C Salas. 2002. "The Effects of Telencephalic Pallial
Lesions on Spatial, Temporal, and Emotional Learning in Goldfish." Brain Research
Bulletin 57 (3): 397–99. https://doi.org/10.1016/S0361-9230(01)00699-2.

Preston, Brian T., Isabella Capellini, Patrick McNamara, Robert A. Barton, and Charles L.
Nunn. 2009. "Parasite Resistance and the Adaptive Significance of Sleep." BMC
Evolutionary Biology 9 (1): 7. https://doi.org/10.1186/1471-2148-9-7.

Rasch B, Born J. About sleep's role in memory. Physiol Rev. 2013 Apr;93(2):681-766. doi:
10.1152/physrev.00032.2012. PMID: 23589831; PMCID: PMC3768102.

Rattenborg, Niels C., Horacio O. de la Iglesia, Bart Kempenaers, John A. Lesku, Peter Meerlo,
and Madeleine F. Scriba. 2017. "Sleep Research Goes Wild: New Methods and
Approaches to Investigate the Ecology, Evolution and Functions of Sleep."
Philosophical Transactions of the Royal Society B: Biological Sciences 372 (1734).
https://doi.org/10.1098/rstb.2016.0251.

Rawashdeh, Oliver, Nancy Hernandez de Borsetti, Gregg Roman, and Gregory M. Cahill.
2007. "Melatonin Suppresses Nighttime Memory Formation in Zebrafish." Science
318 (5853): 1144–46. https://doi.org/10.1126/science.1148564.

Rechtschaffen, Allan, and Bernard M. Bergmann. 2002. "Sleep Deprivation in the Rat: An
Update of the 1989 Paper." Sleep 25 (1): 18–24.
https://doi.org/10.1093/sleep/25.1.18.

Ribeiro, Sidarta, Damien Gervasoni, Ernesto S. Soares, Yi Zhou, Shih-Chieh Lin, Janaina
Pantoja, Michael Lavine, and Miguel A. L. Nicolelis. 2004. "Long-Lasting Novelty-
Induced Neuronal Reverberation during Slow-Wave Sleep in Multiple Forebrain
Areas." PLoS Biology 2 (1): E24. https://doi.org/10.1371/journal.pbio.0020024.

Romer, Alfred Sherwood. 1959. The Vertebrate Story. University of Chicago Press.

Roth, Timothy C., Niels C. Rattenborg, and Vladimir V. Pravosudov. 2010. "The Ecological
Relevance of Sleep: The Trade-off between Sleep, Memory and Energy
Conservation." Philosophical Transactions of the Royal Society B: Biological Sciences
365 (1542): 945–59. https://doi.org/10.1098/rstb.2009.0209.

Sauer, S., M. Kinkelin, E. Herrmann, and W. Kaiser. 2003. "The Dynamics of Sleep-like
Behaviour in Honey Bees." Journal of Comparative Physiology A 189 (8): 599–607.
https://doi.org/10.1007/s00359-003-0436-9.

Schuster S, Wöhl S, Griebsch M, Klostermeier I. Animal cognition: how archer fish learn to
down rapidly moving targets. Curr Biol. 2006 Feb 21;16(4):378-83. doi:
10.1016/j.cub.2005.12.037. PMID: 16488871.

Scullin, Michael K., and Donald L. Bliwise. 2015. "Is Cognitive Aging Associated with Levels of
REM Sleep or Slow Wave Sleep?" Sleep 38 (3): 335–36.
https://doi.org/10.5665/sleep.4482.

Shaw, P. J., C. Cirelli, R. J. Greenspan, and G. Tononi. 2000. "Correlates of Sleep and Waking in Drosophila Melanogaster." Science (New York, N.Y.) 287 (5459): 1834–37. https://doi.org/10.1126/science.287.5459.1834.

Shaw, Paul J., Giulio Tononi, Ralph J. Greenspan, and Donald F. Robinson. 2002. "Stress Response Genes Protect against Lethal Effects of Sleep Deprivation in Drosophila." Nature 417 (6886): 287–91. https://doi.org/10.1038/417287a.

Sherry, David F., and Jennifer S. Hoshooley. 2010. "Seasonal Hippocampal Plasticity in Food-Storing Birds." Philosophical Transactions of the Royal Society B: Biological Sciences 365 (1542): 933–43. https://doi.org/10.1098/rstb.2009.0220.

Shettleworth, Sara J. 1995. "Comparative Studies of Memory in Food Storing Birds: From the Field to the Skinner Box." In Behavioural Brain Research in Naturalistic and Semi-Naturalistic Settings, edited by E Alleva, A Fasolo, Hans-Peter Lipp, L Nadel, and L Ricceri, 159–92. The Netherlands: Kluwer Academic.

———. 1998. Cognition, Evolution, and Behavior. 2nd ed. Oxford University Press.

Siegel, Jerome M. 2001. "The REM Sleep-Memory Consolidation Hypothesis." Science 294 (5544): 1058–63. https://doi.org/10.1126/science.1063049.

Smith, C. 2001. "Sleep States and Memory Processes in Humans: Procedural versus Declarative Memory Systems." Sleep Medicine Reviews 5 (6): 491–506. https://doi.org/10.1053/smrv.2001.0164.

Smith, C., and G. Kelly. 1988. "Paradoxical Sleep Deprivation Applied Two Days after End of Training Retards Learning." Physiology & Behavior 43 (2): 213–16. https://doi.org/10.1016/0031-9384(88)90240-5.

Stickgold, Robert. 2006. "A Memory Boost While You Sleep." Nature 444 (7119): 559–60. https://doi.org/10.1038/nature05309.

Stickgold, Robert, and Matthew P. Walker. 2005. "Memory Consolidation and Reconsolidation: What Is the Role of Sleep?" Trends in Neurosciences 28 (8): 408–15. https://doi.org/10.1016/j.tins.2005.06.004.

Straube, B. An overview of the neuro-cognitive processes involved in the encoding, consolidation, and retrieval of true and false memories. *Behav Brain Funct* **8**, 35 (2012). https://doi.org/10.1186/1744-9081-8-35

Toates, F. (2012). Operant Behavior. In: Seel, N.M. (eds) Encyclopedia of the Sciences of Learning. Springer, Boston, MA. https://doi.org/10.1007/978-1-4419-1428-6_992

Tobler, I. 2011. "Phylogeny of Sleep Regulation." In Principles and Practice of Sleep Medicine, edited by M. H. Kryger, T Roth, and W.C. Dement, 5th ed., 112–25. Philadelphia: Elsevier Saunders.

Tobler, Irene. 1995. "Is Sleep Fundamentally Different between Mammalian Species?" Behavioural Brain Research, The Function of Sleep, 69 (1): 35–41. https://doi.org/10.1016/0166-4328(95)00025-O.

Tobler, Irene, and Alexander A. Borbély. 1985. "Effect of Rest Deprivation on Motor Activity of Fish." Journal of Comparative Physiology A 157 (6): 817–22. https://doi.org/10.1007/BF01350078.

Tononi, Giulio, and Chiara Cirelli. 2006. "Sleep Function and Synaptic Homeostasis." Sleep Medicine Reviews 10 (1): 49–62. https://doi.org/10.1016/j.smrv.2005.05.002.

Vargas, Juan Pedro, Verner Peter Bingman, Manuel Portavella, and Juan Carlos López. 2006. "Telencephalon and Geometric Space in Goldfish." European Journal of Neuroscience 24 (10): 2870–78. https://doi.org/10.1111/j.1460-9568.2006.05174.x.

Vertes, Robert P. 2004. "Memory Consolidation in Sleep: Dream or Reality." Neuron 44 (1):
135–48. https://doi.org/10.1016/j.neuron.2004.08.034.
Vorster, Albrecht P., and Jan Born. 2015. "Sleep and Memory in Mammals, Birds and
Invertebrates." Neuroscience and Biobehavioral Reviews 50 (March): 103–19.
https://doi.org/10.1016/j.neubiorev.2014.09.020.
Wagner, Ullrich, Steffen Gais, Hilde Haider, Rolf Verleger, and Jan Born. 2004. "Sleep
Inspires Insight." Nature 427 (6972): 352–55. https://doi.org/10.1038/nature02223.
Walker MP. The role of slow wave sleep in memory processing. J Clin Sleep Med. 2009 Apr
15;5(2 Suppl):S20-6. PMID: 19998871; PMCID: PMC2824214.
Walker, MP., and R. Stickgold. 2004a. "Sleep-Dependent Learning and Memory
Consolidation." Neuron 44 (1): 121–33.
https://doi.org/10.1016/j.neuron.2004.08.031.
———. 2004b. "Sleep-Dependent Learning and Memory Consolidation." Neuron 44 (1):
121–33. https://doi.org/10.1016/j.neuron.2004.08.031.
———. 2006. "Sleep, Memory, and Plasticity." Annual Review of Psychology 57: 139–66.
https://doi.org/10.1146/annurev.psych.56.091103.070307.
Whitney, P., Hinson, J. M., Jackson, M. L., & Van Dongen, H. P. A. (2015). Feedback blunting:
Total sleep deprivation impairs decision making that requires updating based on
feedback. Sleep, **38**, 745–754.
Sharon Wismer, Ana I Pinto, Zegni Triki, Alexandra S Grutter, Dominique G Roche, Redouan
Bshary 2019. Cue-based decision rules of cleaner fish in a biological market task.
Animal behaviour 158: 249-260
Worley, Susan L. 2018. "The Extraordinary Importance of Sleep." Pharmacy and
Therapeutics 43 (12): 758–63.
Yokogawa, Tohei, Wilfredo Marin, Juliette Faraco, Guillaume Pézeron, Lior Appelbaum, Jian
Zhang, Frédéric Rosa, Philippe Mourrain, and Emmanuel Mignot. 2007.
"Characterization of Sleep in Zebrafish and Insomnia in Hypocretin Receptor
Mutants." PLoS Biology 5 (10). https://doi.org/10.1371/journal.pbio.0050277.
Zimmerman, John E., Nirinjini Naidoo, David M. Raizen, and Allan I. Pack. 2008.
"Conservation of Sleep: Insights from Non-Mammalian Model Systems." Trends in
Neurosciences 31 (7): 371–76. https://doi.org/10.1016/j.tins.2008.05.001.
Yuanyuan Xie and Richard I. Dorsky (2017) Development of the hypothalamus:
conservation, modification and innovation. 2017. Development.144(9): 1588–1599.
doi: 10.1242/dev.139055

Supplementary material

Table 1. Number of incorrect choices before food reward (Learning Phase)

	Df	Sum Sq	Mean Sq	F value	Pr(>F)	
sleep_treatment	1	77.66	77.66	5.972	0.0346	*
day	1	0.68	0.68	0.052	0.8240	
Residuals	10	130.04	13.00			

596   Table 2. Time to correct choice (Learning Phase)

```
                   Df   Sum Sq Mean Sq F value Pr(>F)
sleep_treatment    1  4958787 4958787   5.117 0.0472 *
day                1      353     353   0.000 0.9852
Residuals         10  9690710  969071
```

603   Table 3. Number of incorrect choices before food reward (Memory Phase)

```
                   Df Sum Sq Mean Sq F value Pr(>F)
sleep_treatment    1  6.458   6.458   1.937  0.197
day                1  0.502   0.502   0.151  0.707
Residuals          9 30.011   3.335
```

610   Table 4. Time to correct choice (Memory Phase)

```
                   Df Sum Sq Mean Sq F value Pr(>F)
sleep_treatment    1  389.0   389.0   2.549  0.145
day                1    2.5     2.5   0.017  0.900
Residuals          9 1373.3   152.6
```

615