Systematic evaluation of *de novo* mutation calling tools using whole genome sequencing data

Anushi Shah[1,2], Steven Monger[1,2], Michael Troup[1], Eddie KK Ip[1,2] and Eleni Giannoulatou[1,2,*]

[1] Victor Chang Cardiac Research Institute, NSW 2010 Sydney, Australia

[2] School of Clinical Medicine, St Vincent's Healthcare Clinical Campus, Faculty of Medicine and Health, UNSW Sydney, Australia

* To whom correspondence should be addressed. Tel: +61 2 9295 8669; Email: e.giannoulatou@victorchang.edu.au

ABSTRACT

De novo mutations (DNMs) are genetic alterations that occur for the first time in an offspring. DNMs have been found to be a significant cause of severe developmental disorders. With the widespread use of next-generation sequencing (NGS) technologies, accurate detection of DNMs is crucial. Several bioinformatics tools have been developed to call DNMs from NGS data, but no study to date has systematically compared these tools. We used both real whole genome sequencing (WGS) data from a trio from the 1000 Genomes Project (1000G) and simulated trio data to evaluate four DNM calling tools: DeNovoGear, TrioDeNovo, PhaseByTransmission, and VarScan 2. For DNMs called in the real dataset, we observed 8.7% concordance between all tools, while up to 36.2% of DNMs were unique to each caller. For simulated trio WGS datasets spiked with 100 DNMs, the concordance rate was also low at 4.1%. DeNovoGear achieved the highest F1-score on both the real and simulated datasets. Our study provides valuable recommendations for the selection and application of DNM callers on WGS trio data.

INTRODUCTION

De novo mutations (DNMs) are new mutations that arise in an individual, as opposed to being inherited from the parents. The human germline *de novo* single nucleotide variant (SNV) mutation rate is estimated at 1.0 to 1.8 x 10^{-8} which equates to 44 to 82 DNMs per genome per generation[1,2]. DNMs can occur during the formation of the gametes or post-zygotically (somatic mutations). These mutations have been associated with congenital malformations[3-5], early-onset diseases such as intellectual disability[6] and other developmental diseases[3,7], and late-onset diseases such as Parkinson's disease[8], schizophrenia[9], and bipolar disorder[10].

Next generation sequencing (NGS) technologies have made it possible to study whole genomes or exomes at a single base pair resolution. In disease studies, trio-based sequencing involves whole exome sequencing (WES) or whole genome sequencing (WGS) of an affected offspring and their parents. This approach allows for the identification of genetic variants unique to the affected offspring and is commonly used to study sporadic genetic disorders caused by DNMs[11-17].

In standard approaches, DNMs are detected by first inferring the genotypes of each individual in a trio separately using standard variant calling algorithms such as Genome Analysis Toolkit (GATK)[18] or Samtools[19]. The putative DNMs are then identified by comparing the genotypes of the offspring to those of the parents. However, this approach is inefficient and yields many false positives. The error rates in NGS data are much higher than the expected DNM rate, and any miscalled genotypes in either the parents or the offspring will result in false positives. In recent years, many computational methods have been developed specifically for calling DNMs from NGS data. These methods use the raw read-alignment data or genotype likelihoods for each possible genotype to determine a DNM call. Four widely-used DNM calling tools are DeNovoGear[20], VarScan 2[21], TrioDeNovo[22] and PhaseByTransmission[23].

Independent benchmarking and evaluation of genomics software, such as variant callers, is critical for establishing best practices. Several studies have compared variant calling pipelines for detecting SNVs and insertions/deletions (indels)[24-26], including somatic mutations[27-29], across different sequencing technologies. One prior study evaluated DNM callers in humans[30] using WES data from

the Ashkenazi Jewish trio set (NA12878). However, this comparison did not include the latest state-of-the-art DNM callers such as TrioDeNovo, DeNovoGear and PhaseByTransmission. Additionally, the evaluation of this study did not incorporate any validated DNMs. Instead, GC (guanine-cytosine) content, substitution type and single nucleotide variant (SNV) density were used as proxy evaluation metrics.

In this study, we present a systematic evaluation of current state-of-the-art and commonly used DNM calling tools for WGS trio data. Our assessment incorporates a dataset comprising validated DNMs as well as a simulated trio dataset. *De novo* single nucleotide variants (*dn*SNVs) and *de novo* insertions/deletions (*dn*INDELs) are compared separately for characteristics affecting DNM detection, such as read depth and allelic fraction. Our objective is to offer recommendations of best practices for the identification of DNMs from WGS trio data.

MATERIAL AND METHODS

Real dataset

We used the Illumina HiSeq WGS data of one CEU (Central European from Utah) trio from the 1000 Genomes Project (1000G) for evaluation[31]. The trio samples we used were NA12891 (father), NA12892 (mother), and NA12878 (female offspring). Their aligned CRAM files (based on the GRCh38 reference genome) were downloaded from

ftp://ftp.1000genomes.ebi.ac.uk/vol1/ftp/data_collections/illumina_platinum_pedigree/data/CEU/NA12878/ and converted to BAM files for further bioinformatics processing. The depth of coverage of BAM files for father, mother, and offspring was 60x, 60x, and 66x, respectively. The genome co-ordinates of variants were lifted over from hg19 to hg38 using UCSC genome browser's hgLiftOver feature[32]. The dataset contains experimentally validated DNMs, alongside other variant types. Specifically, the dataset comprises 48 germline DNMs, 888 cell line somatic DNMs, 125 inherited polymorphisms and 1,256 false positive DNMs, all originating from autosomal chromosomes. For the DNM variant caller comparison, the combined germline and cell line somatic DNMs (936) will be used as positive DNMs, while the inherited polymorphisms and false positive DNMs (1381) will be used as negative DNMs. A

Variant Call Format (VCF) containing all samples in the trio was generated by executing the best practices standard variant calling pipeline of GATK (version 4.0)[37] using default parameters.

Simulated dataset

To generate our simulated trio datasets, we first simulated variants for both the mother and father based on allele frequencies observed in the European population data available in the 1000G project. We used the open-source software SIMdrom[33] to do this, which generates random variants based on a provided VCF file. The two resulting VCF files were merged to create one parental VCF file encompassing variants from both parents. The offspring VCF file was then generated by applying the Mendelian inheritance law to the merged parental VCF file, to infer the genotypes of the offspring.

Next, 100 DNMs were randomly sampled from the public database denovo-db[34] and spiked into the offspring VCF file. These DNMs comprised 92 *dn*SNVs and 8 *dn*INDELs. All simulated and spiked-in variants originated from autosomal chromosomes, consistent with the real data. Lastly, Illumina-based NGS reads were simulated using the NEAT[35] open-source software package with the GRCh38 reference genome, at 30x coverage depth. The final output was three independent BAM files representing the trio (mother, father, and offspring). To ensure data integrity and quality, the generated BAM files underwent post-processing procedures in adherence to GATK's best practices guidelines. A VCF file utilising the simulated trio's generated BAMs was created using GATK (version 4.0), with default parameters. The VCF file was checked for Mendelian inconsistencies using the +mendelian plug-in provided by bcftools[19].

De novo mutation calling

DeNovoGear DeNovoGear is a software application that detects DNMs from familial and somatic tissue using NGS trio data[20]. Its model consists of individual genotype likelihoods, transmission probabilities and prior probabilities to model the likelihood of observing a polymorphism or a DNM at any given site (Table 1). Users can specify the prior probability of observing a DNM which can be used to increase or decrease calling sensitivity. It employs beta-binomial distributions that are fitted to alternate read frequencies for both SNPs and indels to facilitate mutation detection. The "dng dnm" subcommand of DeNovoGear was executed to detect DNMs in trios. DeNovoGear required as inputs

a Pedigree (PED) format file and a trio based Binary Variant Call Format (BCF) file. The output is a single text file containing a row for each putative DNM, with various fields including event type (Point Mutation/INDEL), physical location and the posterior probability of the most likely *de novo* genotype configuration.

VarScan 2 VarScan 2 is a variant calling tool that detects somatic and germline mutations from NGS trio data[21]. The DNM calling "trio" subcommand of VarScan 2 detects high confidence DNMs by leveraging the familial relationship present in trio datasets, thereby identifying apparent Mendelian Inheritance Errors (MIEs) (Table 1). It first calls all variants across all three samples using the standard way of mpileup2snp, using default values for parameters "--min-var-freq" and "--p-value". Subsequently, it identifies any variants with apparent MIEs, i.e., variants that are present in the offspring but absent in parental samples. In such circumstances, it re-calls the samples with relaxed parameters and reclassifies with some evidence (--adj-var-freq and --adj-p-value). Often, this corrects MIEs, and corrected genotypes are reported. If not, the site is reported as "mendelError" (in the "FILTER" field) and/or "DENOVO" (in the "INFO" field). The VarScan 2 "trio" subcommand utilises a trio BCF file as input and produces two output VCF files: one for SNVs and another for indels. The output VCF files contain "INFO" fields for all called SNVs and indels. If a DNM is called, the "STATUS VCF" field will contain a value of "3". For high confidence DNM calls, the variant is also annotated with the text label "DENOVO".

TrioDeNovo TrioDeNovo is a program which uses a Bayesian framework to detect DNMs from NGS trio data without specifying a prior mutation rate[22] (Table 1). Firstly, it evaluates the evidence of DNMs using trio data and then adjusts the effect of mutation rates on DNM calling through prior odds. For input, the user provides a PED file, and a VCF file with PL (Phred-scaled genotype likelihood) or GL (genotype likelihood) fields. The output is a VCF file containing the detected DNMs.

PhaseByTransmission PhaseByTransmission is a software module integrated with GATK (version 3.8) to detect DNMs from NGS trio data[23]. By utilising genotype likelihoods and a mutation prior, it calculates the most likely genotype combination and phases trios and parent/offspring pairs (Table 1). The software requires a PED file and a VCF file as input, and it outputs a VCF file and a Mendelian

violations file. PhaseByTransmission computes genotype combinations of the trio (i.e. mother-father-offspring) and supports bi-allelic cases, with 27 possible genotype-trio combinations. For our comparison, we have used the trio genotype combination of "AA-AA-AB" (i.e. mother-father-offspring), where the A is the reference allele. PhaseByTransmission is included in the GATK software, version 3.8 release.

Performance Evaluation

To evaluate the performance of the four DNM callers in detecting DNMs, we generated confusion matrices and calculated F1 scores, along with other evaluation metrics. For each caller, we obtained results using two methods: DNMs called using the program's default setting, and DNMs based on a "benchmarking-set maximal F1 threshold". We determined the "benchmarking-set maximal F1 threshold" for each caller by calculating F1 scores across the range of possible thresholds (the range of each caller's confidence scores) and identifying the threshold which resulted in the highest F1 score (Fig. S1). Results using the "benchmarking-set maximal F1 threshold" represent the best performance for each caller, for the dataset. This provides us with a level playing field in comparing results between the four callers.

RESULTS

We evaluated the DNM calling methods DeNovoGear, VarScan 2, TrioDeNovo, and PhaseByTransmission using both real and simulated datasets of WGS trios.

Real dataset

PhaseByTransmission detected the largest number of DNMs (60,079), a large portion being *dn*INDELs (44,463), while VarScan 2 detected the lowest number of DNMs (10,517) from the 1000G CEU trio dataset (Table 2). In evaluating the results, we found that some DNMs detected by the callers had missing genotypes for one or both parents. To remove these falsely classified DNMs, we applied genotype-based filters using the genotype pattern of "0/0" (homozygous for the reference allele) for the mother and father, and "0/1" (heterozygous for the alternative allele) for the offspring. After filtering, PhaseByTransmission had the lowest number of DNM calls for *dn*SNVs (2,302),

followed by VarScan 2 (7,841) and DeNovoGear (10,272) (Table 3). DeNovoGear (1,231) and TrioDeNovo (8,726) detected the lowest and highest number of *dn*INDELs, respectively.

We investigated the concordance of DNMs called by the four DNM callers for *dn*SNVs (Fig. 1a) and *dn*INDELs (Fig. 1b). The concordance was extremely low, with only 2,072 *dn*SNVs (8.7%) and 113 *dn*INDELs (0.9%) called by all callers. TrioDeNovo gave the largest number of unique calls (not identified by any other callers), with 8,661 *dn*SNVs and 6,862 *dn*INDELs. In contrast, PhaseByTransmission gave no unique *dn*SNVs calls and only 7 unique *dn*INDELs.

Using the DNM callers' default score thresholds, TrioDeNovo called the highest number of true positive calls (TPs) and had the highest sensitivity (0.99252) (Table 4). However, it also had the highest number of false positive calls (FPs=11,064) and had the lowest F1-score (0.14371). In comparison, PhaseByTransmission called a slightly lower number of TPs (918), but it called the lowest number of FPs (1,384), resulting in the highest F1-score (0.56702).

We calculated the "benchmarking-set maximal F1 threshold" for each DNM caller (see Methods – Performance evaluation). Using "benchmarking-set maximal F1 thresholds", DeNovoGear achieved the highest F1 score (0.58887), and notably had the fewest FPs (1,168) (Table 5). PhaseByTransmission's performance was a very close second, with an F1 score of 0.57838. VarScan 2 followed with an F1 score of 0.53965, producing the highest number of FNs (82). Finally, TrioDeNovo achieved an F1 score of 0.41015, producing the highest number of FPs (2541).

Properties of True Positive and False Positive DNMs identified by DNM callers

The accuracy of each algorithm in detecting DNMs depends on both the sequencing coverage and the true variant allele frequency (VAF) of the mutations. We investigated the relationship between read depth at the DNM positions and the estimated VAF of DNMs identified by all callers. To maintain consistent values between the DNM callers, VAF was calculated using allelic counts obtained using the GATK tool "CollectAllelicCounts". For true positive *dn*SNVs, all the DNM callers had similar VAFs with a combined mean value of 0.46 and standard deviation of 0.08 and were identified in positions with read depths between 27 and 80 (Fig. 2a-d).

DeNovoGear and TrioDeNovo identified false positive *dn*SNVs across a wide range of VAFs and read depths, especially at lower VAF levels when the read depth was high (>100 reads) (Fig. 2e-h, y-axis cropped at 200). By comparison, PhaseByTransmission and VarScan 2 did not detect false positive *dn*SNVs at high read depths. All the callers detected *dn*SNVs across the full range of VAFs, from 0-1. We note that the DNM callers can use different methods of filtering lower quality reads for use in the VAF calculation (compared to "CollectAllelicCounts"), which explains some discrepancies, for example the presence of DNM calls with VAF values of 0. There were no true positive *dn*INDELs called, so only false positive *dn*INDELs were analysed. For false positive *dn*INDELs, we observed that all callers identified *dn*INDELs with lower VAF, but only TrioDeNovo exhibited more *dn*INDEL calls at the higher VAF range (Fig. S2).

We also investigated the VAF distribution within the trio samples for both true positive and false positive DNMs, as a factor influencing whether DNMs are called. For true positive *dn*SNVs, we observed that the parent's VAFs in the DNM positions were extremely low or zero (father's mean VAF = 0.0002 and standard deviation = 0.002, mother's mean VAF = 0.0001 and standard deviation = 0.001, across all callers) (Fig. 3a-d). This aligns with a DNM parental and offspring profile. In the case of false positive *dn*SNVs, we observed that for all callers, there was a large concentration of lower VAFs in the parents, which may account for the false call of an offspring's *dn*SNV, especially for callers such as DeNovoGear and TrioDeNovo (Fig. 3e-h). A VAF value of 0.5 (expected for a germline heterozygote variant) was observed in the offspring for many DNMs detected by all the callers, with PhaseByTransmission DNMs showing the highest number. This high VAF indicates that some of these false positive *dn*SNVs are either systematic errors seen across the trio, or are inherited variants observed with lower VAFs in the parents. For false positive *dn*INDELs, we observed a similarly large concentration of very low VAFs for the mother, father, and offspring across all callers (Fig. S3). No peak around 0.5 in VAF (expected for a germline heterozygote variant) was observed for false positive *dn*INDELs.

We further investigated the relationship between VAF and read depth using the consensus of different numbers of callers for *dn*SNVs (Fig. S4). Out of 936 validated true positive *dn*SNVs, 905 were

detected by all four DNM callers. These 905 dnSNVs had VAFs ranging between 0.24 and 0.70 and a mean read depth of 45, which was very similar to the false positive *dn*SNVs called by all four DNM callers (VAFs ranging between 0.25 to 0.75, and a mean read depth of 46). As the number of DNM callers decreased, the VAF range increased.

We also investigated patterns in the output confidence scores of TP and FP DNMs for each caller (Fig. 4). The scores for DeNovoGear and VarScan 2 showed a clear separation between TP and FP DNMs, indicating better calibration. DeNovoGear uses posterior probability for its DNM score (PPDM). The majority (916/921) of TP DNMs called by DeNovoGear had PPDM > 0.99, while 83.5% (7809/9351) of FP DNMs had PPDM < 0.99. For VarScan 2, which uses a normalised P-value, all 916 TP DNMs had p-value < 9.4048 x 10^{-4}, while 84.5% of FP DNMs had p-value > 0.05. TrioDeNovo, which uses a normalised Denovo Quality score, showed overlap between TP and FP DNMs, though TPs were concentrated in the higher quality score range. For PhaseByTransmission, which uses a normalised Phred score, there was no clear distinction between the scores of TP and FP DNMs.

Simulated Datasets

We simulated trio WGS data with 100 spiked-in DNMs using a custom framework, as described in the methods section. Of these 100 spiked-in DNMs, 92 were *dn*SNVs and 8 were *dn*INDELs. We applied the four DNM callers to the simulated data and evaluated their performance.

DeNovoGear called the largest number of DNMs (1,431), followed by TrioDeNovo, then VarScan 2, and finally PhaseByTransmission, which called only 315 DNMs (Table 6). DeNovoGear also identified the most *dn*SNVs (1,391), while PhaseByTransmission identified the fewest (117). TrioDeNovo detected the largest number of *dn*INDELs (824), while DeNovoGear detected the smallest number (40). DeNovoGear, VarScan 2, and TrioDeNovo successfully identified all 92 TP *dn*SNVs, while PhaseByTransmission identified only 35 of them. DeNovoGear also reported the greatest number of false positives, followed by TrioDeNovo, VarScan 2, and PhaseByTransmission.

We calculated concordance and evaluation metrics for *dn*SNVs following the same procedure as with the real 1000G CEU trio dataset. We found that the concordance between the 4 callers was exceptionally low at 4.1% (Fig 5). This low concordance was consistent with the results observed from the real dataset. DeNovoGear had the highest number of unique DNM calls.

For performance evaluation, we calculated the "benchmarking-set maximal F1 threshold" for the callers (see Methods – Performance evaluation). DeNovoGear achieved by far the highest F1 score of 0.56089 (Table 7). The next best performer was TrioDeNovo which achieved an F1-score of 0.42703. Following this, PhasebyTransmission's F1 score was considerably lower at 0.33493, owing to its high number of FNs (52). Finally, VarScan 2 achieved the lowest F1 score of 0.23349, mainly due to its high number of FPs (483).

DISCUSSION

In this study, we present a systematic evaluation of four widely used DNM calling tools: DeNovoGear, VarScan 2, TrioDeNovo, and PhaseByTransmission. We developed a software framework for generating simulated NGS trio datasets. Using a simulated dataset, as well as an experimentally validated 1000G CEU trio dataset, we investigated the performance of each caller.

We found that the raw number of calls varied dramatically between the different callers, and the concordance among the four callers was strikingly low, with 8.7% for *dn*SNVs and 0.9% for *dn*INDELs in the 1000G CEU dataset. For both the real and simulated datasets, all DNM callers showed high sensitivity, which means they were able to detect the majority of true DNMs.

When detecting DNMs from NGS data, the large scale of WGS data eclipses the small number of true DNMs present. In this setting, a tendency to call FPs is particularly costly as it can lead to overwhelmingly large numbers of (mostly false positive) DNM calls. To provide a more realistic classification setting, we designed evaluation datasets with significant imbalances in favour of negative variants. In this imbalanced setting, F1-score was chosen as an appropriate evaluation metric.

We provide performance results using each caller's default thresholds, as we expect that many users would adopt the default settings, so performance at these thresholds is of interest. We also provide results using "benchmarking-set maximal F1 thresholds", which allows a fairer comparison of each caller's ability to identify DNMs. Overall, we identified DeNovoGear as the best performing DNM caller, as it achieved the highest F1 score on both the real and simulated datasets, using "benchmarking-set maximal F1 thresholds".

We found that F1 scores were relatively low for all callers, with the highest F1 scores achieved on the real and simulated datasets being 0.58887 and 0.56089, respectively. The calling of large numbers of false positive DNMs is a major issue. Taken together, these results are indicative of significant performance issues for all callers. We propose that an ensemble classifier using the scores of multiple DNM callers would likely provide significant performance improvements, especially given the low concordance among individual callers.

The performance of the DNM callers is dependent on both the sequencing coverage and the VAFs of the DNMs. We observed clear distinctions in the patterns of VAF and read depth between true positive and false positive *dn*SNVs called by each caller. As noted earlier, both TrioDeNovo and DeNovoGear call large numbers of false positives. A considerable proportion of these calls are made within a very low VAF range and at a very high or low read depth, indicating technical artifacts such as sequencing errors. This is corroborated by investigating the trio VAF density patterns, which clearly show parental VAFs at or near 0 values for true positive *dn*SNVs, as they are absent in parents. In contrast, parental VAFs of false positive *dn*SNVs showed a large proportion at detectable VAF values.

The VAF and read depth patterns of false positive *dn*SNVs obtained by consensus of all four callers in the real 1000G CEU dataset were similar to those of the true positive *dn*SNVs. It is possible that some of the false DNMs that were called by all the callers could be real somatic DNMs that were not validated by the original DNM study[31]. This is a limitation of our study that relies on this experimentally validated set of DNMs.

The use of confidence scores from the callers could be an appropriate quality control step to filter out false positive DNMs. For DeNovoGear (PPDM threshold) and VarScan 2 (p-value), there were clear distinctions in confidence scores between true positive and false positive DNMs. In the case of DeNovoGear, a PPDM threshold of 0.99 yielded the majority of true positives. For VarScan 2, all true positive calls have p-value $< 9.4048 \times 10^{-4}$. However, for TrioDeNovo and PhaseByTransmission, the distinction in confidence scores between true and false positive DNMs is unclear.

Many research groups have already incorporated standard variant processing pipelines, such as GATK, as part of their routine WGS data analyses. Hence, they may have VCF output files readily available for further processing. In such situations, it may be beneficial to use VCF-based input DNM callers such as TrioDeNovo and PhaseByTransmission. On the other hand, DeNovoGear and VarScan 2 have only one pre-processing step; that of generating mpileup files (BCFs) from BAMs using a samtools mpileup command, which is an additional step in the pipeline.

DeNovoGear and PhaseByTransmission provide phased genotype information, which can be used to infer haplotypes from genotype data. However, as TrioDeNovo and PhaseByTransmission do not provide this information, in our analysis we only considered unphased genotype information in comparing the DNM callers.

Recently, collaborative efforts such as the Genome in a Bottle (GIAB) consortium have aimed to establish a reference standard for human genome sequence data[38]. Data for this purpose has been collected using 12 different sequencing technologies and library preparation methods. While gold standard calls have been generated for the Ashkenazim Trio data, these consist of gold standard truth variants within specific, high-confidence regions of the GIAB genome. There has not been a focus on generating a set of gold standard DNMs across the entire genome. There is a strong need for such public datasets containing validated DNMs, which could be extremely beneficial for the evaluation of existing callers and pipelines, as well as for the development of new algorithms.

Finally, the presence of post-zygotic DNMs in developmental disorders has recently received attention[39]. Determining the proportion of affected cells and tissue types can help to determine the

timing of formation of DNMs, which plays a crucial role in the clinical phenotype. In the future, researchers will need to develop computational methods that can distinguish between germline and post-zygotic DNMs. This will require high-depth sequencing and computational methods that can confidently analyse the distribution of variant allele frequencies to detect differences between germline and post-zygotic DNMs that occur early enough in development to reach elevated levels.

DATA AVAILABILITY

The 1000 Genome CEU trio we used was downloaded from ftp://ftp.1000genomes.ebi.ac.uk/vol1/ftp/data_collections/illumina_platinum_pedigree/data/CEU/NA12878/

The validated DNMs were taken from Conrad *et al.*[31]. Our trio simulated framework is available at https://github.com/VCCRI/TrioSim.

FUNDING

This work was supported by the NSW Health [Early Mid-Career Cardiovascular Grant to EG], National Health and Medical Research Council [Investigator Grant 2018360 to EG] and a Research Training Program (RTP) scholarship by the University of New South Wales [to AS].

CONFLICT OF INTEREST

The authors declare no conflict of interest.

REFERENCES

1.   Roach JC, Glusman G, Smit AF, et al. Analysis of genetic inheritance in a family quartet by whole-genome sequencing. Science 2010;328(5978):636-9, doi:10.1126/science.1186802
2.   Acuna-Hidalgo R, Veltman JA, Hoischen A. New insights into the generation and role of de novo mutations in health and disease. Genome Biol 2016;17(1):241, doi:10.1186/s13059-016-1110-1

3.	Homsy J, Zaidi S, Shen Y, et al. De novo mutations in congenital heart disease with neurodevelopmental and other congenital anomalies. Science 2015;350(6265):1262-6, doi:10.1126/science.aac9396

4.	Hoischen A, van Bon BW, Gilissen C, et al. De novo mutations of SETBP1 cause Schinzel-Giedion syndrome. Nat Genet 2010;42(6):483-5, doi:10.1038/ng.581

5.	Hoischen A, van Bon BW, Rodríguez-Santiago B, et al. De novo nonsense mutations in ASXL1 cause Bohring-Opitz syndrome. Nat Genet 2011;43(8):729-31, doi:10.1038/ng.868

6.	Hamdan FF, Srour M, Capo-Chichi JM, et al. De novo mutations in moderate or severe intellectual disability. PLoS Genet 2014;10(10):e1004772, doi:10.1371/journal.pgen.1004772

7.	Iossifov I, O'Roak BJ, Sanders SJ, et al. The contribution of de novo coding mutations to autism spectrum disorder. Nature 2014;515(7526):216-21, doi:10.1038/nature13908

8.	Kun-Rodrigues C, Ganos C, Guerreiro R, et al. A systematic screening to identify de novo mutations causing sporadic early-onset Parkinson's disease. Hum Mol Genet 2015;24(23):6711-20, doi:10.1093/hmg/ddv376

9.	Fromer M, Pocklington AJ, Kavanagh DH, et al. De novo mutations in schizophrenia implicate synaptic networks. Nature 2014;506(7487):179-84, doi:10.1038/nature12929

10.	Nishioka M, Kazuno AA, Nakamura T, et al. Systematic analysis of exonic germline and postzygotic de novo mutations in bipolar disorder. Nat Commun 2021;12(1):3750, doi:10.1038/s41467-021-23453-w

11.	Girard SL, Gauthier J, Noreau A, et al. Increased exonic de novo mutation rate in individuals with schizophrenia. Nat Genet 2011;43(9):860-3, doi:10.1038/ng.886

12.	Jin ZB, Wu J, Huang XF, et al. Trio-based exome sequencing arrests de novo mutations in early-onset high myopia. Proc Natl Acad Sci U S A 2017;114(16):4219-4224, doi:10.1073/pnas.1615970114

13.	Michaelson JJ, Shi Y, Gujral M, et al. Whole-genome sequencing in autism identifies hot spots for de novo germline mutation. Cell 2012;151(7):1431-42, doi:10.1016/j.cell.2012.11.019

14.	O'Roak BJ, Deriziotis P, Lee C, et al. Exome sequencing in sporadic autism spectrum disorders identifies severe de novo mutations. Nat Genet 2011;43(6):585-9, doi:10.1038/ng.835

15.	Jin ZB, Li Z, Liu Z, et al. Identification of de novo germline mutations and causal genes for sporadic diseases using trio-based whole-exome/genome sequencing. Biol Rev Camb Philos Soc 2018;93(2):1014-1031, doi:10.1111/brv.12383

16.	Awotoye W, Mossey PA, Hetmanski JB, et al. Whole-genome sequencing reveals de-novo mutations associated with nonsyndromic cleft lip/palate. Sci Rep 2022;12(1):11743, doi:10.1038/s41598-022-15885-1

17.	Hunt D, Leventer RJ, Simons C, et al. Whole exome sequencing in family trios reveals de novo mutations in PURA as a cause of severe neurodevelopmental delay and learning disability. J Med Genet 2014;51(12):806-13, doi:10.1136/jmedgenet-2014-102798

18.	McKenna A, Hanna M, Banks E, et al. The Genome Analysis Toolkit: a MapReduce framework for analyzing next-generation DNA sequencing data. Genome Res 2010;20(9):1297-303, doi:10.1101/gr.107524.110

19.	Li H, Handsaker B, Wysoker A, et al. The Sequence Alignment/Map format and SAMtools. Bioinformatics 2009;25(16):2078-9, doi:10.1093/bioinformatics/btp352

20.	Ramu A, Noordam MJ, Schwartz RS, et al. DeNovoGear: de novo indel and point mutation discovery and phasing. Nat Methods 2013;10(10):985-7, doi:10.1038/nmeth.2611

21.	Koboldt DC, Larson DE, Wilson RK. Using VarScan 2 for Germline Variant Calling and Somatic Mutation Detection. Curr Protoc Bioinformatics 2013;44(15.4.1-17, doi:10.1002/0471250953.bi1504s44

22.	Wei Q, Zhan X, Zhong X, et al. A Bayesian framework for de novo mutation calling in parents-offspring trios. Bioinformatics 2015;31(9):1375-81, doi:10.1093/bioinformatics/btu839

23.   Francioli LC, Cretu-Stancu M, Garimella KV, et al. A framework for the detection of de novo mutations in family-based sequencing data. Eur J Hum Genet 2017;25(2):227-233, doi:10.1038/ejhg.2016.147

24.   Kumaran M, Subramanian U, Devarajan B. Performance assessment of variant calling pipelines using human whole exome sequencing and simulated data. BMC Bioinformatics 2019;20(1):342, doi:10.1186/s12859-019-2928-9

25.   Bian X, Zhu B, Wang M, et al. Comparing the performance of selected variant callers using synthetic data and genome segmentation. BMC Bioinformatics 2018;19(1):429, doi:10.1186/s12859-018-2440-7

26.   Supernat A, Vidarsson OV, Steen VM, et al. Comparison of three variant callers for human whole genome sequencing. Sci Rep 2018;8(1):17851, doi:10.1038/s41598-018-36177-7

27.   Chen Z, Yuan Y, Chen X, et al. Systematic comparison of somatic variant calling performance among different sequencing depth and mutation frequency. Sci Rep 2020;10(1):3501, doi:10.1038/s41598-020-60559-5

28.   Chen J, Li X, Zhong H, et al. Systematic comparison of germline variant calling pipelines cross multiple next-generation sequencers. Sci Rep 2019;9(1):9345, doi:10.1038/s41598-019-45835-3

29.   Wang Q, Kotoula V, Hsu PC, et al. Comparison of somatic variant detection algorithms using Ion Torrent targeted deep sequencing data. BMC Med Genomics 2019;12(Suppl 9):181, doi:10.1186/s12920-019-0636-y

30.   Liang Y, He L, Zhao Y, et al. Comparative Analysis for the Performance of Variant Calling Pipelines on Detecting the. Front Pharmacol 2019;10(358, doi:10.3389/fphar.2019.00358

31.   Conrad DF, Keebler JE, DePristo MA, et al. Variation in genome-wide mutation rates within and between human families. Nat Genet 2011;43(7):712-4, doi:10.1038/ng.862

32.   Kent WJ, Sugnet CW, Furey TS, et al. The human genome browser at UCSC. Genome Res 2002;12(6):996-1006, doi:10.1101/gr.229102

33.   Schubach M. SIMdrom. 2016. Available from: https://github.com/visze/simdrom.

34.   Turner TN, Yi Q, Krumm N, et al. denovo-db: a compendium of human de novo variants. Nucleic Acids Res 2017;45(D1):D804-D811, doi:10.1093/nar/gkw865

36.   Stephens ZD, Hudson ME, Mainzer LS, et al. Simulating Next-Generation Sequencing Datasets from Empirical Mutation and Sequencing Models. PLoS One 2016;11(11):e0167047, doi:10.1371/journal.pone.0167047

36.   Van der Auwera GA, Carneiro MO, Hartl C, et al. From FastQ data to high confidence variant calls: the Genome Analysis Toolkit best practices pipeline. Curr Protoc Bioinformatics 2013;43(11.10.1-11.10.33, doi:10.1002/0471250953.bi1110s43

37.   Zook JM, Catoe D, McDaniel J, et al. Extensive sequencing of seven human genomes to characterize benchmark reference materials. Sci Data 2016;3(160025, doi:10.1038/sdata.2016.25

38.   Acuna-Hidalgo R, Bo T, Kwint MP, et al. Post-zygotic Point Mutations Are an Underrecognized Source of De Novo Genomic Variation. Am J Hum Genet 2015;97(1):67-74, doi:10.1016/j.ajhg.2015.05.008

TABLE AND FIGURES

Table 1. Features of DNM calling tools

	DeNovoGear	VarScan 2	TrioDeNovo	PhaseByTransmission
Method/Algorithm	Beta-binomial based likelihood model and evaluation of joint trio likelihood.	Statistical significance test for variant calling and heuristics.	Bayesian model selection framework using individual genotype likelihoods.	Bayesian framework using individual genotype likelihoods.
Input data	BCF (mpileup) and PED files	BCF (mpileup)	VCF file	VCF file
Output data	Text file	VCF files	VCF file	VCF file
Output Confidence score for DNM	Posterior probability	Phred-scaled P-value from Fisher's exact test	Bayes Factor	Phred-scaled posterior probability
Phasing information	Yes	No	No	Yes

Table 2. Total counts of DNMs, *dn*SNVs and *dn*INDELs called by each DNM caller using the 1000G CEU trio, using default caller settings.

DNM Tool	Total DNMs	Total *dn*SNVs	Total *dn*INDELs
DeNovoGear	16,367	15,081	1,286
VarScan 2	10,517	7,841	2,676
TrioDenovo	26,863	14,882	11,981
PhaseByTransmission	60,079	15,616	44,463

Table 3. Total counts of DNMs, *dn*SNVs and *dn*INDELs called by each DNM caller using the 1000G CEU trio, following genotype filtering. Genotype filters were applied against the callers' results from the default setting run. (Father, Mother, Offspring: 0/0, 0/0, 0/1).

DNM Tool	Total DNMs	Total *dn*SNVs	Total *dn*INDELs
DeNovoGear	11,503	10,272	1,231
VarScan 2	10,517	7841	2,676
TrioDeNovo	20,719	11,993	8,726
PhaseByTransmission	4,008	2,302	1,706

Table 4. Performance results using default DNM caller thresholds for 1000G CEU trio dataset.
The four callers were evaluated on the combined dataset of 1000G CEU trio *dn*SNVs (combining all 25,280 putative *dn*SNVs called by all callers). The following metrics were calculated: True Positives (TP), True Negatives (TN), False Negatives (FN), False Positives (FP), Sensitivity and F1-score.

DNM caller	TP	TN	FN	FP	Sensitivity	F1-score
DeNovoGear	921	14,993	15	9351	0.98397	0.16435
VarScan 2	916	17,419	20	6925	0.97863	0.20873
TrioDeNovo	929	13,280	7	11,064	0.99252	0.14371
PhaseByTransmission	918	22,960	18	1,384	0.98077	0.56702

Table 5. Performance results using "benchmarking-set maximal F1 thresholds" for 1000G CEU trio dataset. The four callers were evaluated on the combined dataset of 1000G CEU trio *dn*SNVs (combining all 25,280 putative *dn*SNVs called by all callers). The following metrics were calculated: True Positives (TP), True Negatives (TN), False Negatives (FN), False Positives (FP), Sensitivity and F1-score.

DNM caller	Benchmarking-set maximal F1 threshold	TP	TN	FN	FP	Sensitivity	F1-score
DeNovoGear	0.9999981	878	23,176	58	1168	0.93803	0.58887
VarScan 2	3.28e-06	854	22,969	82	1375	0.91239	0.53965
TrioDeNovo	12.35	897	21,803	39	2541	0.95833	0.41045
PhaseByTransmission	16	880	23,117	56	1227	0.94017	0.57838

Table 6. Total DNMs, *dn*SNVs, *dn*INDELs, True Positives (TP) and False Positives (FP) for each DNM caller using NEAT simulated data using default DNM caller settings.

DNM Tool	Total DNMs	Total dnSNVs	Total dnINDELs	TP dnSNVs	FP dnSNVs
DeNovoGear	1431	1391	40	92	1299
VarScan 2	759	676	83	92	584
TrioDeNovo	1350	526	824	92	434
PhaseByTransmission	315	117	198	35	82

Table 7. Performance results using "benchmarking-set maximal F1 thresholds" on the NEAT simulated dataset. The following metrics were calculated: True Positives (TP), True Negatives (TN), False Negatives (FN), False Positives (FP), Sensitivity and F1-score.

DNM caller	Benchmarking-set maximal F1 threshold	TP	TN	FN	FP	Sensitivity	F1-score
DeNovoGear	0.99102	76	1553	16	103	0.82609	0.56089
VarScan 2	0.00012	76	1173	16	483	0.82609	0.23349
TrioDeNovo	9.32	79	1457	13	199	0.8587	0.42703
PhaseByTransmission	1	35	1574	57	82	0.38043	0.33493

Figure 1. Concordance between DNM callers on the 1000G CEU trio dataset. The Venn diagrams show counts of DNMs called by combinations of the 4 DNM callers for (a) *dn*SNVs and (b) *dn*INDELs. The numbers in parentheses indicate the percentages of total DNMs.

Figure 2. Variant Allele Frequency (or Fraction) vs. Read Depth for *dn*SNVs for all DNM callers using CEU trio from 1000G. True positives (a-d) and false positives (e-h) are shown separately. The colours indicate different DNM callers.

Figure 3. Densities of Variant Allele Frequency (or Fraction) of *dn*SNVs in the 1000G CEU trio.

VAF density is shown for (a-d) true positives and (e-f) false positives for all callers. In (a-d), the VAF

distributions for the parents are visible only as a spike close to VAF=0.

Figure 4. Distribution of the DNM scores for each DNM caller using the 1000G CEU trio. True positive and False positive *dn*SNVs are shown for each DNM caller. (a) Posterior Probability of *de novo* mutation (PPDM) by DeNovoGear (b) Denovo Quality log 10 (Bayes Factor) by TrioDeNovo (c) P-value by VarScan 2 (d) Phred score by PhaseByTransmission. The raw output scores have been normalized.

Figure 5. Concordance between DNM callers on the NEAT-simulated data. The Venn diagram

illustrates total counts for *dn*SNVs called by each DNM caller. The numbers in the parentheses

indicate the percentages of total *dn*SNVs.