Wild jackdaws learn to tolerate juveniles to exploit new information

Josh J. Arbon[1,2*], Noa Truskanov[1,3], Emily Stott[1], Guillam E. McIvor[1,4], Alex Thornton[1]

[1]Centre for Ecology and Evolution, University of Exeter, Cornwall, UK

[2]School of Biological Sciences, University of Bristol, Bristol, UK

[3]School of Zoology, Tel Aviv University, Tel Aviv, Israel

[4]Department of Behavioural and Cognitive Biology, University of Vienna, Vienna, Austria

*Correspondence to J.J.A.: School of Biological Sciences, Life Sciences Building, Tyndall Avenue,
University of Bristol, BS81TQ. josh.arbon@bristol.ac.uk.

Data Accessibility: All data and code required to reproduce the analyses presented in the text are
available to download from https://figshare.com/s/34ce199d3d06a0be20b1

Keywords: Social, tolerance, learning, generalisation, flexibility, social information, social learning,
social structure, aggression.

Article Type: Research Article.

Word count: Abstract – 130, Main Text – 3295.

Number of Figures: 2; Number of Tables: 0; Number of Text Boxes: 0.

Abstract

Social tolerance can enhance access to resources and is thought to be crucial in facilitating the

evolution of cooperation, social cognition and culture, but it is unknown whether animals can

optimise their social tolerance through learning. We presented wild jackdaws (*Corvus*

monedula) with a novel social information problem using automated feeders: to access food,

adults had to inhibit their tendency to displace juveniles and instead show tolerance by

occupying an adjacent perch. Adults learned to tolerate juveniles, generalising across juveniles

as a cohort, demonstrating learning of a new information-use strategy. The ability to learn to

tolerate those that provide valuable information, and generalise across cohorts of informed

individuals, may facilitate adaptive responses in the face of environmental change and help to

explain the success of jackdaws in human-dominated environments.

Introduction

Social tolerance, broadly defined as close proximity to conspecifics with limited aggression [1], is thought to be crucial to the evolution of prosociality [2] and social learning [3] in non-human animals, and underpins distinctive forms of cognition, cooperation and culture in humans [4]. Tolerating others can provide opportunities to learn important information such as how to access food [3], and has even been shown to help buffer the costs of ecological disasters [5], but social tolerance can also entail costs such as foraging competition [6]. General heuristics can aid in responding adaptively to environmental change[5], such as preferentially tolerating kin when resource availability is low [7]. However, if potential partners vary in the benefits they provide, then learning to tolerate specific individuals, or classes of individual, could be vital in facilitating access to information [8] or resources [9]. Despite these important implications, studies have yet to explicitly test whether animals in natural social networks can learn to tolerate others based on their value as sources of information.

Biases in social information use and social learning are well described across taxa [10]. For instance, many animals preferentially learn from older, more experienced individuals [11,12] that are more likely to provide reliable information. However, sources of useful information may change: in humans, for instance, younger generations commonly learn from their elders, but the digital revolution has flipped this pattern with older people seeking vital skills from younger "digital natives" [13]. Similar patterns could occur in animals. Juveniles are often easily displaced by adults and generally not used as preferential information sources [14,15], but juveniles are often more exploratory and innovative [16–19]. Adults may therefore benefit from learning to attend to and tolerate juveniles, especially in changeable or heterogenous environments where reliable models can change quickly [20–22]. Whilst some recent work indicates that animals adjust their social associations based on information about specific partners [23–25], whether these associations can be generalised to classes of

individuals that share characteristics (e.g. juveniles) remains unknown. This combination of

learning and generalisation that guide who to tolerate would represent the learning of a social

information-use strategy [26]. We examined this possibility using an automated field experiment

on wild jackdaws.

Jackdaws are social corvids that forage in fission-fusion social groups [27] and vary in

their social tolerance of different partners, generally showing particularly high tolerance to close

associates and kin [28]. They are proficient social learners about novel foods [29] and

anthropogenic dangers [30], and have been shown to generalise learnt information in foraging

contexts [15]. These abilities may contribute to their success in human-dominated habitats

[22]. Previous work [23] shows jackdaws can learn to associate with individuals based on

payoffs, but whether they can alter their tolerance or generalise this learning across individuals

remains unknown. To answer these outstanding questions, we presented adults with a novel

social information-use problem. Under normal conditions foraging adults do not preferentially

associate with juveniles other than their own offspring (Supplementary Material: Age-class

Assortment), nor attend to juveniles as sources of information [15]. We simulated a scenario

where juveniles became 'knowledgeable' about a new foraging resource by allowing them free

access to automated experimental feeders containing high-quality food (mealworms). As it took

up to a couple of seconds for feeders to close, adults could often scrounge a small reward (max.

one beakful) by aggressively displacing a juvenile as the feeder door closed. However, if adults

refrained from displacing juveniles and instead tolerated their presence on the adjacent feeder,

they gained the far larger reward of feeding *ad libitum* for the duration that they occupied the

adjacent feeder (termed a cofeeding event: mean duration ± SE = 12.3 ± 1.5 s).

To test whether adults could learn to tolerate juveniles to exploit this novel scenario, we

modelled changes in both cofeeding with juveniles and displacement of juveniles relative to

adult experience. We included both the number of cofeeding events with a specific individual,

as well as the number with all juveniles, to test between learning to tolerate specific individuals

versus generalising learnt tolerance to juveniles more generally. We also placed pairs of feeders

with low-quality food (grain), freely accessible to all birds, at distinct locations at the field-site,

to test if changes observed at our experiment carried over into a related, but unrewarded,

context.

Methods

Study System

Experimental sessions ran for five hours from ~5:30 am between from 19 July to 13 August 2021,

at our long-term jackdaw study population in Cornwall, UK (0°11'23"N, 5°10'54"W). The Cornish

Jackdaw Project monitors a core nest-box population of ~80 boxes as part of the wider jackdaw

population, with more than 3000 individuals ringed over a 10-year period. Individuals are fitted

with a unique set of coloured leg rings, one of which contains a Radio Frequency Identification

(RFID) tag (IB Technologies, Leicester, UK), enabling remote detection via antennae at feeders.

Approximately 150 chicks from nest boxes are ringed each year before fledging, and non-

nestbox juveniles are ringed as part of routine trapping [31].

Experimental Apparatus

Feeders automatically recorded the identity of any visiting RFID-tagged bird. Access to food was

restricted by a motor-controlled door connected by a logger unit to an antenna in the perch

(Naturecounters, Maidstone, UK; Figure 1a; Supplementary Material: Experimental Setup).

Cofeeding events were determined from the temporal overlap between tag visits at adjacent,

interconnected feeders (Figure 1a). Temporal data recorded which individual arrived first, initiating an event, and which individual subsequently joined. Displacements were characterised as events where a bird was detected arriving at a feeder within 2 seconds of a different bird departing (validated through video; see Supplementary Material: Displacement Validation).

Feeders were split into two categories: *Experimental* feeders contained high-quality food (mealworms; known to be preferred by jackdaws [23]), whereas *passive* feeders contained low-quality food (grain). *Experimental* feeders were in connected pairs 0.5 m apart, with access to food determined by the tag identity and combination across both feeders (see Methods: *Treatments*). All tagged jackdaws could access *passive* feeders at all times. *Experimental* and *passive* feeders were visually distinguishable by visible food content and a green band present only on *experimental* feeders (Figure 1a).

We placed two pairs of *experimental* feeders at each of two sub-sites. These were flanked by a pair of *passive* feeders to create an experimental array. Together, the four feeders formed a shallow arc, each separated by 0.5 m (Supplementary Material: Experimental Setup). The *passive* feeders in experimental arrays were primarily designed to attract birds to the array, and to provide data on the individuals present in the area; data from these *passive* feeders within the experimental array are not presented in this study. For each experimental array, we also placed a corresponding pair of *passive* feeders in the same arrangement (0.5 m apart) elsewhere within the local area (~50m away). These *passive* feeders were used to detect associations and displacements outside of the direct context of the experimental setup. Unfortunately, during the onset of the experiment, a dead jackdaw appeared next to one of the experimental arrays, resulting in a near-complete abandonment of all feeders at one sub-site

(Supplementary Material: Feeder Visitation). As such, only data from feeders at one sub-site

was analysed (N = 2 experimental arrays, 2 passive feeder pairs).

Treatments

All RFID-tagged juveniles formed the 'knowledgeable' group, with unrestricted access to

experimental feeders. Treatment adults (70% of adults chosen randomly) could access

mealworm rewards from *experimental* feeders, but only if a juvenile was simultaneously present

on the adjacent *experimental* feeder. Control adults (remaining 30% of adults) were not able to

access food from the *experimental* feeders under any circumstances. All three groups had

unrestricted access to *passive* feeders, including those within experimental arrays and the

passive feeder pairs at nearby locations. The control adults were intended to provide baseline

data on adult–juvenile interactions but visited experimental arrays at rates too low to allow

meaningful analyses (Supplementary Material: Feeder Visitation). However, the use of a control

group is not necessary for our key aim of detecting changes in within-individual behaviour over

time (see Statistical Methods).

Ethical Note

Research was conducted under ethical permission from the Penryn Campus Ethics Committee

(eCORN000406), following Association for the Study of Animal Behaviour ethical guidelines

[32]. Birds were ringed under BTO permits C6449, C5752, C6923 and C6079.

Statistical Methods

We analysed data using Relational Event Models (REMs), a form of time-to-event model

that accounts for the structure of the social network, to compare changes in the rates of

cofeeding and displacement events based on individual experience (following Kings *et al.* 2023

[23]). REMs account for the order of events in a network, preventing data aggregation [33–35],

and test hypotheses by comparing observed events against a set of 'non-events', permuted

events that could hypothetically happen between individuals in the network; 10000

permutations were conducted ([36], see Supplementary Material: Relational Event Modelling

for details).

We used this approach to separately model whether adults changed their propensity to

(a) cofeed with juveniles (Supplementary Tables 1,3,5), and (b) displace juveniles

(Supplementary Tables 2,4,6), based on their previous experience in the experiment. When

investigating cofeeding likelihood, we considered the number of times an adult had previously

cofed with a *specific individual* (*IRR individual*), as well as the number of times an adult had

cofed with *any juvenile* (*IRR all juvs*) to investigate generalisation of learning. The likelihood of

displacement was modelled as for cofeeding, with the addition of a fixed factor for the number

of times an adult had displaced a juvenile (*IRR displacement*) to investigate whether learning

from displacements reinforced this behaviour. By interacting these factors with the age

(juvenile/adult) of the individual joined/displaced, we determined the likelihood of cofeeding

with/displacing a juvenile relative to that of an adult as a function of experience. Results from

experimental feeders were qualitatively identical when considering adult experience as either

(a) the number of successful cofeeds where the adult joined a juvenile at an experimental

feeder, or (b) the total number of successful cofeeds the adult participated in, both joining or

being joined by a juvenile (Supplementary Tables 1–2). We present the results from (a) in the

main text unless specified.

All results are reported as Incidence Rate Ratios, such that at IRR of 1.05 for a predictor

variable means that the event type in question in 5% more likely for every unit increase in that

predictor. We took the median model coefficient of the 10000 permuted models as a point

estimate for each parameter, with the 2.5 and 97.5 percentiles of each coefficient taken to

generate 95% confidence intervals.

Results

i) Experimental Results

As adult jackdaws gained experience, they increased their tolerance of juveniles. Over

the experiment, we detected 2431 cofeeding events, where both *experimental* feeders were

occupied simultaneously. Of those, 766 were successful cofeeds between 200 unique adult–

juvenile dyads, 174 of which involved adults joining juveniles. Overall, the proportion of adult

cofeeding events with juveniles (i.e., successes) roughly doubled across the course of the

experiment (Figure 1b, grey-blue line).

Figure 1. (a) Experimental feeders, showing an adult (right) in a successful cofeed with a juvenile. RFID tags (highlighted), determine access to mealworms by motor-controlled doors. (b) The daily proportion of visits where adults successfully co-fed with (grey-blue) or displaced (violet) a juvenile.

At the individual level, the likelihood an adult joined a juvenile at the feeders (as opposed to joining another adult) increased by 3.4% for each previous successful cofeed the adult had participated in with any juvenile (*IRR all juvs* = 1.034, *CI* = 1.021–1.049; Figure 2a). Displacements of juveniles also became slightly more common over the course of the experiment (Figure 1b, violet line) as adults learnt that when displacing juveniles they could scrounge a small reward before the feeder closed (*IRR displacement* = 1.004, *CI* = 1.002–1.006). However, the likelihood that an adult displaced a juvenile (relative to another adult) declined by 2.2% per successful cofeed with any juvenile (*IRR all juvs* = 0.978, *CI* = 0.970–0.987; Figure 2b).

Together, these results indicate that successful cofeeding with any juvenile leads to adults learning to change their behaviour towards juveniles in general. Indeed, the effects of

experience with specific juveniles (as opposed to any juvenile), are inconclusive on both the

relative likelihood of cofeeding with (*IRR individual* = 1.002, *CI* = 0.986–1.020), or displacing (*IRR*

individual = 0.992, *CI* = 0.983–1.001) a juvenile relative to another adult. Succeeding with a

specific juvenile, or failing with a specific adult, appears to therefore be less important than

overall successes with all juveniles.

Figure 2. The difference in likelihood of an adult (a) cofeeding with, and (b) displacing a juvenile

(orange) as opposed to another adult (blue) as adults acquired experience of successful

cofeeding with all juveniles. Nine adults reached 6 or more previous successes. Lines and

shaded area represent median and 95% REM estimates.

One adult was responsible for a large number of successful joining events (37/174). If

we exclude this adult from analyses, the effect of generalisation across juveniles becomes less

clear for patterns of cofeeding; the estimate for the effect of experience with *all juveniles* is still

positive, but the CI crosses 1 (i.e., no effect: *IRR all juvs* = 1.012, *CI* = 0.987–1.044), whilst the

specific individual effect becomes more clear (*IRR individual* = 1.039, *CI* = 1.007–1.079).

However, the effects for displacement, are qualitatively the same without the influential adult,

with an increase in magnitude for the effect size of experience with all juveniles (*IRR individual*

= 1.000, *CI* = 0.988–1.012, *IRR all juvs* = 0.952, *CI* = 0.938–0.968).

ii) Carry-over of experimental manipulation

We also found limited evidence that tolerance learnt at the experimental feeders carried

over to the non-experimental context. We did not see clear changes in cofeeding at passive

feeders (Supplementary Table 3), however, adults were less likely to displace juveniles if their

successes with all juveniles at the experimental feeders were higher (though only in models

where successes included those where a juvenile joined an adult Supplementary Table 4).

iii) Ruling out confounds

In principle, the apparent changes in adult behaviour with experience could arise as an artefact

if adults with a greater pre-existing propensity to either associate with juveniles, or to visit

experimental feeders, engaged with the experiment more and were more successful. We can

rule this out, however as (1) adults' preferences for cofeeding with juveniles at the start of the

experiment were unrelated to the final number of successes they had and (2) adults that visited

the experimental apparatus more frequently were not more successful overall (Supplementary

Information: Initial Propensity).

Discussion

Taken together, our results show that adult jackdaws can learn to tolerate juveniles when it is

advantageous to do so. As they gained experience in the experiment, adults increased

cofeeding with and reduced displacement of juveniles relative to adults, indicating they learned

to tolerate juveniles based on their information value. Although there were small rewards

available for displacing juveniles, these were largely outweighed by the benefits of tolerant

cofeeding. Moreover, our results suggest generalisation across individuals may facilitate rapid,

beneficial changes in tolerant behaviour, representing the learning of a new social information-

use strategy, namely 'tolerate juveniles'.

The potential benefits of such learned tolerance are abundant. Tolerating conspecifics

has been shown to facilitate cooperation in ravens *Corvus corax* [2], whilst tolerance is

implicated in the social learning of key foraging skills such as tool use in other corvid species

[37,38], even in species not known to use tools in the wild [39]. Whilst information can be

gathered from others without tolerance (e.g. broad-scale patch discovery by observing others

from a distance), social tolerance can increase the chances of co-occurring with valuable

individuals and observing their actions in close proximity. This may be especially valuable in

species without fixed social groups, such as jackdaws and other animals living in fission-fusion

societies, where individuals can switch to other social partners if they provide less aggression

or better payoffs [23,27]. Our study therefore highlights the ability of jackdaws to flexibly alter

information-use strategies through learning and generalisation. Whether the changes we

detected in the context of foraging behaviour might carry over into other contexts (e.g. predator

defence) is a major outstanding question. Addressing this will be challenging but important to

help us understand how flexibility may help animals like jackdaws thrive in changeable, human-

disturbed environments [21].

The findings that experience interacting with juveniles in general, over experience with

specific individuals, was an important driver of behavioural change suggests that adults were

able to generalise their learning to juveniles as a cohort. Whilst information-use strategies are

known to be influenced by developmental stress [40], our findings show that learning can allow

the rapid adjustment of strategies to novel conditions. This is consistent with suggestions that

rather than evolving as fixed, species-typical traits [10], social learning strategies use domain-

general associative learning processes to modify associations and inhibition of aggressive

behaviour in response to reinforcement, enabling flexibility in the face of changing payoffs [26].

This flexibility also illustrates the benefits of tracking and responding to others' behaviour in

dynamic social environments, a key assumption of the Social Intelligence Hypothesis [41].

However, generalising among similar individuals could facilitate access to valuable new

opportunities and information without needing to remember the outcomes of past interactions

with many different individuals, reducing cognitive burdens.

It is also possible that different adults implemented different learning strategies. The

nuanced changes in interpretation when removing the individual responsible for the largest

number of successful joining events – i.e., strong evidence for generalising across juveniles

when considering displacements but evidence for the importance of experience with specific

individuals when considering co-feeding – highlights the potential for variation in who is

attended to, and how. In practice it is likely that both remembering past interactions with

specific individuals and generalising across groups of individuals will be used together to guide

behaviour. Differences in how individuals integrate these processes could be underlaid by

differences in cognitive abilities or prior experience [42], but we do not have a sample size large

enough to investigate these differences robustly. Regardless of the specific strategy

implemented by individuals, the key finding that adults learn to modify their social tolerance is

clear: results cannot be explained by adults' pre-existing preference for associating with

juveniles, or their propensity to engage with the experiment in general.

Broadly, flexibly adjusting social tolerance, including through learning new information-

use strategies, could facilitate the flow of information through networks by permitting close

observation of a wide range of innovators. Indeed, the ability to capitalise on opportunities

generated by innovators through flexible information-use is thought to be a crucial element of

human culture [26]. Assessing the degree of flexibility in social information use and social

learning by non-human animals will be an important factor in understanding how they respond

to changes in the environment and increasing interactions with humans [43]. Our work also

provides a rare illustration of the bi-directional interplay between information use and social

structure in natural populations, contributing to our understanding of how sociality and

decision-making processes coevolve [44,45].

Acknowledgements

We would like to thank the Gluyas family and Odette Eddy for access to their land to conduct

the experiment. Thanks to Andy Young for many valuable discussions and Stephanie King for

comments.

Author Contributions: Conceptualization: J.J.A. and A.T. Methodology: J.J.A. and A.T. Validation:
J.J.A. Formal Analysis: J.J.A. Investigation: J.J.A, N.T., E.S. Resources: J.J.A., G.E.M., and A.T. Data
Curation: J.J.A. Writing – Original Draft: J.J.A and A.T. Writing – Review & Editing: J.J.A., N.T, E.S.,
G.E.M., and A.T. Visualization: J.J.A. Supervision: A.T. Project Administration: J.J.A., G.E.M., and A.T.
Funding Acquisition: J.J.A. and A.T.

Funding

J.J.A. was supported by the Biotechnology and Biological Sciences Research Council-funded

South West Biosciences (SWBio) Doctoral Training Partnership [680027356]. A.T. and G.E.M.

were supported by a Leverhulme Trust grant [RGP-2020-170] to A.T. and N.T. was supported by

an SNSF post-doctoral mobility fellowship [P400PB_194397/1].

Data accessibility

328 All data and code for this work can be found at https://figshare.com/s/34ce199d3d06a0be20b1.

References

1. Cronin KA, Sánchez A. 2012 Social dynamics and cooperation: The case of nonhuman
primates and its implications for human behavior. *Advances in Complex Systems* **15**,
1250066. (doi:10.1142/S021952591250066X)

2. Massen JJM, Ritter C, Bugnyar T. 2015 Tolerance and reward equity predict cooperation in
ravens (*Corvus corax*). *Scientific Reports* **5**, 1–11. (doi:10.1038/srep15021)

3. Coelho CG, Garcia-Nisa I, Ottoni EB, Kendal RL. 2024 Social tolerance and success-biased
social learning underlie the cultural transmission of an induced extractive foraging tradition in
a wild tool-using primate. *Proceedings of the National Academy of Sciences* **121**,
e2322884121. (doi:10.1073/pnas.2322884121)

4. Cieri RL, Churchill SE, Franciscus RG, Tan J, Hare B. 2014 Craniofacial feminization, social
tolerance, and the origins of behavioral modernity. *Current Anthropology* **55**, 419–443.
(doi:10.1086/677209)

5. Testard C *et al*. 2024 Ecological disturbance alters the adaptive benefits of social ties. *Science*
384, 1330–1335. (doi:10.1126/science.adk0606)

6. De la Fuente MF, Sueur C, Garber PA, Bicca-Marques JC, Souto A, Schiel N. 2022 Foraging
networks and social tolerance in a cooperatively breeding primate (*Callithrix jacchus*). *Journal*
of Animal Ecology **91**, 138–153. (doi:10.1111/1365-2656.13609)

7. Banks SC, Lindenmayer DB, Mcburney L, Blair D, Knight EJ, Blyton MDJ. 2011 Kin selection in
den sharing develops under limited availability of tree hollows for a forest marsupial.
Proceedings of the Royal Society B: Biological Sciences **278**, 2768–2776.
(doi:10.1098/rspb.2010.2657)

8. Van Schaik CP. 2003 Local traditions in orangutans and chimpanzees: social learning and
social tolerance. In *The Biology of Traditions* (eds DM Fragaszy, S Perry), pp. 297–328.
Cambridge: Cambridge University Press.

9. Miller R, Schiestl M, Whiten A, Schwab C, Bugnyar T. 2014 Tolerance and Social Facilitation in
the Foraging Behaviour of Free-Ranging Crows (*Corvus corone corone*; *C. c. cornix*). *Ethology*
120, 1248–1255. (doi:10.1111/eth.12298)

10. Kendal RL, Boogert NJ, Rendell L, Laland KN, Webster M, Jones PL. 2018 Social Learning
Strategies: Bridge-Building between Fields. *Trends Cognit. Sci.* **22**, 651–665.
(doi:10.1016/j.tics.2018.04.003)

11. Wood LA, Kendal RL, Flynn EG. 2012 Context-dependent model-based biases in cultural
transmission: Children's imitation is affected by model age over model knowledge state.
Evolution and Human Behavior **33**, 387–394. (doi:10.1016/j.evolhumbehav.2011.11.010)

12. Kendal RL, Hopper LM, Whiten A, Brosnan SF, Lambeth SP, Schapiro SJ, Hoppitt W. 2015
Chimpanzees copy dominant and knowledgeable individuals: Implications for cultural
diversity. *Evolution and Human Behavior* **36**, 65–72.
(doi:10.1016/j.evolhumbehav.2014.09.002)

13. Heyes C. 2016 Blackboxing: social learning strategies and cultural evolution. *Philosophical*
Transactions of the Royal Society of London B: Biological Sciences **371**.
(doi:10.1098/rstb.2015.0369)

14. Penndorf J, Aplin L. 2020 Environmental and life history factors, but not age, influence social
learning about food: a meta-analysis. *Animal Behaviour* **167**, 161–176.
(doi:10.1016/j.anbehav.2020.07.001)

15. Arbon JJ, Hahn LG, McIvor GE, Thornton A. 2023 Competition and generalization impede
cultural formation in wild jackdaws. *Proc. R. Soc. B.* **290**, 20230705.
(doi:10.1098/rspb.2023.0705)

16. Biro D, Inoue-Nakamura N, Tonooka R, Yamakoshi G, Sousa C, Matsuzawa T. 2003 Cultural
innovation and transmission of tool use in wild chimpanzees: Evidence from field experiments.
Animal Cognition **6**, 213–223. (doi:10.1007/s10071-003-0183-x)

17. Carter GG, Forss S, Page RA, Ratcliffe JM. 2018 Younger vampire bats (*Desmodus rotundus*)
are more likely than adults to explore novel objects. *PLoS ONE* **13**, 1–11.
(doi:10.1371/journal.pone.0196889)

18. Benson-Amram S, Heinen VK, Gessner A, Weldele ML, Holekamp KE. 2014 Limited social
learning of a novel technical problem by spotted hyenas. *Behavioural Processes* **109**, 111–120.
(doi:10.1016/j.beproc.2014.09.019)

19. Morand-Ferron J, Cole EF, Rawles JEC, Quinn JL. 2011 Who are the innovators? A field
experiment with 2 passerine species. *Behavioral Ecology* **22**, 1241–1248.
(doi:10.1093/beheco/arr120)

20. Sol D, Timmermans S, Lefebvre L. 2002 Behavioural flexibility and invasion success in birds.
Animal Behaviour **63**. (doi:10.1006/anbe.2001.1953)

21. Ducatez S, Sol D, Sayol F, Lefebvre L. 2020 Behavioural plasticity is associated with reduced
extinction risk in birds. *Nature Ecology & Evolution* , 1–6. (doi:10.1038/s41559-020-1168-8)

22. Lee VE, Thornton A. 2021 Animal Cognition in an Urbanised World. *Frontiers in Ecology and*
Evolution **9**, 1–20. (doi:10.3389/fevo.2021.633947)

23. Kings M, Arbon JJ, McIvor GE, Whitaker M, Radford AN, Lerner J, Thornton A. 2023 Wild
jackdaws can selectively adjust their social associations while preserving valuable long-term
relationships. *Nature Communications* **14**, 5103. (doi:10.1038/s41467-023-40808-7)

24. Kulahci IG, Ghazanfar AA, Rubenstein DI. 2018 Knowledgeable Lemurs Become More Central
in Social Networks. *Current Biology* **28**, R344–R346. (doi:10.1016/j.cub.2018.02.079)

25. O'Hearn WJ, Beckmann J, Von Fersen L, Dal Pesco F, Mundry R, Keupp S, Diakhate N,
Niederbremer C, Fischer J. 2025 Increased female competition for males with enhanced
foraging skills in Guinea baboons. *Proc. R. Soc. B.* **292**, 20242925.
(doi:10.1098/rspb.2024.2925)

26. Mesoudi A, Chang L, Dall SRX, Thornton A. 2016 The evolution of individual and cultural
variation in social learning. *Trends in Ecology and Evolution* **31**, 215–225.
(doi:10.1016/j.tree.2015.12.012)

27. Chen R, Spiegel O, Bartan Y, Nathan R. 2022 Resource limitation drives fission–fusion
dynamics of group composition and size in a social bird. *Animal Behaviour* **191**, 15–32.
(doi:10.1016/j.anbehav.2022.06.003)

28. Röell A. 1977 Social behaviour of the jackdaw, *Corvus mondeula*, in relation to its niche.
Behaviour **64**, 1–122. (doi:10.1163/156853978X00459)

29. Greggor AL, McIvor GE, Clayton NS, Thornton A. 2016 Contagious risk taking: Social
information and context influence wild jackdaws' responses to novelty and risk. *Scientific
Reports* **6**, 27764. (doi:10.1038/srep27764)

30. Lee VE, Régli N, McIvor GE, Thornton A. 2019 Social learning about dangerous people by wild
jackdaws. *Royal Society Open Science* **6**, 191031. (doi:10.1098/rsos.191031)

31. Greggor AL, McIvor GE, Clayton NS, Thornton A. 2018 Wild jackdaws are wary of objects that
violate expectations of animacy. *Royal Society Open Science* **5**, 181070.
(doi:10.1098/rsos.181070)

32. ASAB. 2020 Guidelines for the treatment of animals in behavioural research and teaching.
Animal Behaviour **159**, i–xi. (doi:10.1016/j.anbehav.2019.11.002)

33. Tranmer M, Marcum CS, Morton FB, Croft DP, de Kort SR. 2015 Using the relational event
model (REM) to investigate the temporal dynamics of animal social networks. *Animal
Behaviour* **101**, 99–105. (doi:10.1016/j.anbehav.2014.12.005)

34. Patison KP, Quintane E, Swain DL, Robins G, Pattison P. 2015 Time is of the essence: an
application of a relational event model for animal social networks. *Behavioral Ecology and
Sociobiology* **69**, 841–855. (doi:10.1007/s00265-015-1883-3)

35. Butts CT. 2008 A relational event framework for social action. *Sociological Methodology* **38**,
155–200. (doi:10.1111/j.1467-9531.2008.00203.x)

36. Lerner J, Lomi A. 2020 Reliability of relational event model estimates under sampling: How to
fit a relational event model to 360 million dyadic events. *Network Science* **8**, 97–135.
(doi:10.1017/nws.2019.57)

37. Holzhaider JC, Hunt GR, Gray RD. 2010 Social learning in New Caledonian crows. *Learning &*
Behavior **38**, 206–219. (doi:10.3758/LB.38.3.206)

38. Rutz C *et al.* 2016 Discovery of species-wide tool use in the Hawaiian crow. *Nature* **537**, 403–
407. (doi:10.1038/nature19103)

39. Bird CD, Emery NJ. 2009 Insightful problem solving and creative tool modification by captive
nontool-using rooks. *Proc. Natl. Acad. Sci. U.S.A.* **106**, 10370–10375.
(doi:10.1073/pnas.0901008106)

40. Farine DR, Spencer KA, Boogert NJ. 2015 Early-life stress triggers juvenile zebra finches to
switch social learning strategies. *Current Biology* **25**, 2184–2188.
(doi:10.1016/j.cub.2015.06.071)

41. Dunbar RIM. 1998 The social brain hypothesis. *Evolutionary Anthropology: Issues, News, and*
Reviews **6**, 178–190. (doi:10.1002/(SICI)1520-6505(1998)6:5<178::AID-EVAN5>3.0.CO;2-8)

42. Nöbel S, Wang X, Talvard L, Tariel J, Lille M, Cucherousset J, Roussigné M, Danchin É. 2022 The
importance of population heterogeneities in detecting social learning as the foundation of
animal cultural transmission. *Proceedings of the Royal Society B: Biological Sciences* **289**,
20220431. (doi:10.1098/rspb.2022.0431)

43. Arbon JJ, Boogert NJ, Jordan NR, Thornton A. 2025 The flexibility of social learning and its
conservation implications in mammals and beyond. *Philosophical Transactions of the Royal*
Society B: Biological Sciences In Press.

44. Hahn LG, Sergiou AS, Arbon JJ, Furtbauer I, King AJ, Thornton A. 2025 The co-evolution of
cognition and sociality. *Philosophical Transactions of the Royal Society B: Biological Sciences*
In Press.

45. Kulahci IG, Quinn JL. 2019 Dynamic Relationships between Information Transmission and
Social Connections. *Trends in Ecology and Evolution* **34**, 545–554.
(doi:10.1016/j.tree.2019.02.007)