Wild corvids learn social tolerance to exploit new information

Abstract

Social tolerance is crucial in facilitating the evolution of cooperation, social cognition and

culture, but it is unknown whether animals can optimise their social tolerance through

learning. We presented wild jackdaws (*Corvus monedula*) with a novel social information

problem using automated feeders: to access food, adults had to inhibit their tendency to

displace juveniles and instead show tolerance by occupying an adjacent perch. Adults learned

to tolerate juveniles, generalising across juveniles as a cohort, demonstrating learning of a

new information-use strategy.

Main

Social tolerance, broadly defined as close proximity to conspecifics with limited aggression[1], is thought to be crucial to the evolution of prosociality[2] and social learning[3] in animals, and underpins distinctive forms of cognition, cooperation and culture in humans[4]. However, tolerating others can entail costs such as foraging competition[5], so individuals may benefit from adjusting their tolerance in response to changing socio-ecological conditions[6]. Learning to tolerate high-value individuals, or classes of individual, may be beneficial in maximizing access to information[7] or resources[8]. For instance, juveniles are often easily displaced by adults and may be ignored as information sources[9], but as juveniles can be more exploratory and innovative[10] adults could benefit from learning to tolerate and attend to them. For example, in human societies, the digital revolution flips the tendency for youngsters to learn from their elders[11], as older people seek skills from younger "digital natives"[12]. Using learning and generalisation to adjust a social information-use strategy[13] and tolerate individuals with particular characteristics (e.g. age) may facilitate rapid, adaptive responses in changeable or heterogenous environments where reliable models can change quickly[14]. However, studies have yet to explicitly test whether animals in natural social networks can learn to tolerate others based on their information value.

We tested whether wild corvids (jackdaws: *Corvus monedula*) could learn to adjust their social tolerance to maximise rewards by presenting adults with a novel social information-use problem. Foraging adults do not preferentially associate with juveniles other than their own offspring (Supplementary Information: Figure S1) or attend to them as sources of information[9]. However, we simulated a scenario where juveniles became 'knowledgeable' about a new foraging resource by allowing them free access to automated experimental feeders containing high-quality food (mealworms). Feeders remained locked to adults unless there was a juvenile on an adjacent connected feeder, at which point access to adults was granted (hereafter a 'successful cofeeding event'; Figure 1a). As the feeder doors took approximately one second to close after a juvenile departed, adults could scrounge a small reward (max. one beakful) by displacing a juvenile. However, if adults refrained from displacing juveniles and instead tolerated their presence on the adjacent feeder, they gained the far larger reward of feeding for the duration of the cofeeding event (mean ± SE = 12.3 ± 1.5 s). To test whether effects carried over beyond the experimental feeders, we placed pairs of "passive" feeders with low-quality food (grain), freely accessible to all birds, at separate

locations (~50 m away). We assigned 30% of adults to a control group, but these adults
visited these feeders at low rates, prohibiting analysis (for details see Supplementary
Information: Figure S2). Following previous work[15], we analysed data using Relational Event
Models (REMs), a form of time-to-event model that accounts for the structure of the social
network. Models estimate an Incidence Rate Ratio (IRR), where an IRR of 1.3 means an event
is 30% more likely to occur per unit increase in the predictor (see Methods; model outputs in
Extended Data Tables 1–6).

Figure 1. (a) Experimental feeders, showing an adult (right) in a successful cofeed with a
juvenile. RFID tags (highlighted), determine access to mealworms by motor-controlled doors.
(b) Daily proportion of Daily proportion of visits where adults successfully co-fed with
(yellow) or displaced (blue) a juvenile.

As adult jackdaws gained experience, they increased their tolerant cofeeding with
juveniles. Over the experiment, we detected 2431 cofeeding events, where both perched were
occupied simultaneously. Of those, 766 were successful cofeeds between 200 unique adult–
juvenile dyads, 174 of which involved adults joining juveniles. Overall, the proportion of adult
cofeeding events with juveniles roughly doubled across the course of the experiment (Figure
1b, yellow line). At the individual level, the likelihood an adult joined a juvenile at the feeders
(as opposed to joining another adult) increased by 3.4% for each previous successful cofeed
the adult had participated in (*IRR* = 1.034, *CI* = 1.021–1.049; Figure 2a). Displacements of
juveniles also became slightly more common over the course of the experiment (Figure 1b,

blue line) as adults learnt that when displacing juveniles they could scrounge a small reward before the feeder closed (*IRR* = 1.004, *CI* = 1.002–1.006). However, the benefits of tolerant cofeeding with a juvenile (*ad libitum* access) far outweighed the benefits of scrounging. Consequently, the likelihood of displacements of juveniles, relative to that of displacements of other adults, declined by 2.2% per cofeeding success (*IRR* = 0.978, *CI* = 0.970–0.987; Figure 2b), indicating adults learned the rewards of tolerance.

Figure 2. The difference in likelihood of an adult (a) cofeeding with, and (b) displacing a juvenile (yellow) as opposed to another adult (blue) relative to the number of previous successful cofeeding events that adult had previously participated in. Nine adults reached 6 or more previous successes. Lines and shaded area represent median and 95% REM estimates.

Our results also suggest adults generalised across juveniles as a cohort, rather than learning to tolerate specific individuals. In the results above, adults' increases in cofeeding and reductions in displacement were predicted by their total previous successes with any juvenile. By contrast, the number of prior cofeeding events with a specific juvenile did not affect subsequent cofeeding with, or displacement of that juvenile (both estimates span 1: specific individual cofeed *IRR* = 1.002, *CI* = 0.986–1.020; specific individual displacement *IRR* = 0.992, *CI* = 0.983–1.001). Thus, rewarding interactions with juveniles in general were more important in adult decision making than experience with specific juvenile partners. If we exclude one adult that was responsible for a large number of successful joining events

(37/174) from analyses, the result that adults generalise to reduce displacements of juveniles is strengthened, though generalisation of cofeeding becomes less clear (Extended Data Tables 5-6). However, there was also some evidence that tolerance learnt at the experimental feeders may have carried over to a non-experimental foraging context at passive feeders, where no rules were required to access food. Although we did not see clear changes in cofeeding at passive feeders, there was some evidence that adults were less likely to displace juveniles if their successes at the experimental feeders was higher (Extended data table 4). This suggestion that some tolerance carried over into an unrewarded context provides further support to the idea that adults generalised their tolerance of juveniles.

Taken together, our findings show that adult jackdaws learn to adjust their social tolerance to access information and rewards from demographic group that is usually ignored[9]. The results are unlikely to be explained by pre-existing social preferences before the experiment, as performance (proportion of visits that were successful cofeeds) was unrelated to the total number of visits made to the experiment. Similarly, adults' propensity to cofeed with a juvenile when they had no prior experience was unrelated to their overall number of successful cofeeds at the end of the experiment (Supplementary Information: Figure S3). Instead, we suggest that adults modified their behaviour by learning the value of tolerating juveniles. Rather than modifying their responses towards the specific juveniles with whom they had successfully co-fed previously, adults appeared to apply a rule across all juveniles, potentially even in an unrewarded context. Thus, jackdaws learned a new social information-use strategy[13,16]: juveniles provide information about the availability of food.

Information-use strategies are known to be influenced by developmental stress[17], but we show that learning can allow the rapid adjustment of strategies to novel conditions. This is consistent with suggestions that rather than evolving as fixed, species-typical traits [16], social learning strategies use domain-general associative processes, enabling flexibility in the face of changing payoffs[13]. This flexibility also illustrates the benefits of tracking and responding to others' behaviour in dynamic social environments, a key assumption of the Social Intelligence Hypothesis[18]. However, generalising among similar individuals could facilitate access to valuable new information and opportunities without needing to remember the outcomes of past interactions with many different individuals, reducing cognitive burdens.

Flexibly adjusting social tolerance by learning new information-use strategies could facilitate the flow of information through networks by permitting close observation of a wide

range of innovators. Indeed, the ability to capitalise on opportunities generated by innovators through flexible information-use is thought to be a crucial element of human culture[13] and may help to explain how animals like jackdaws can thrive in human-disturbed environments[19]. Our work also provides a rare illustration of the bi-directional interplay between information use and social structure in natural populations; contributing to our understanding of how sociality and decision-making processes coevolve[20].

Methods

Study System

The experiment was conducted on a wild population of colour-ringed jackdaws at two field-sites (Site Y: 50°11'23"N, 5°10'54"W; Site Z: 50°11'55.5"N, 5°10'09.0"W) of the Cornish Jackdaw Project, a long-term field study in Cornwall, UK. The project actively monitors a core nest-box population of ~80 boxes as part of the wider community of birds in the area. Jackdaws are captured at nest-boxes, using walk-in traps and whoosh nets throughout the breeding season, with more than 3000 individuals ringed over a 10-year period. Every individual has a set of uniquely identifiable coloured leg rings, with one ring containing a Radio Frequency Identification (RFID) tag (IB Technologies, Leicester, UK) to enable remote detection via antennae at feeders. Individual age is known from plumage characteristics at ringing. Approximately 150 chicks from nest boxes are ringed each year before fledging, as well as non-nest-box juveniles ringed as part of routine trapping [21].

Experimental Apparatus

Automated feeders were attached to posts approximately 1 m above the ground at prominent locations across the two field sites. Feeders featured a perch with an embedded RFID-detecting antenna that recorded the identity of any visiting bird. Access to the food within the feeder was restricted by a servo-motor controlled perspex door, with the antennae and motor connected by a logger unit (Naturecounters, Maidstone, UK). Feeders were split into two categories, *experimental* and *passive*. *Experimental* feeders were filled with a high-quality food (mealworms)[15], whereas passive feeders were filled with a low-quality food (poultry corn). *Experimental* feeders were placed in pairs 0.5 m apart and directly connected via a networking cable, with access to food within determined by the tag identity and combination of tags across both feeders (see Methods: *Treatments*). Passive feeders opened in response to

the reading of any RFID tag, so all tagged jackdaws could access food at all times. *Experimental*
and *passive* feeders were identical units running different operational programs, and so were
visually distinguished both by food contents and by a green band present on *experimental*,
but not passive, feeders (Figure 1). All feeders were closed by default, with a tag read required
to trigger door opening.

We placed two pairs of *experimental* feeders at each site, which themselves were
flanked by a pair of *passive* feeders to create an experimental array. Together, the four feeders
formed a shallow arc, each separated by 0.5 m (setup: *Passive – Experimental – Experimental*
– Passive). The passive feeders in *experimental* arrays were primarily designed to attract birds
to the array, as well as to provide data on the individuals present in the local area. Data from
these *passive* feeders within the experimental array are not presented in this study. For each
experimental array, we placed a corresponding pair of *passive* feeders within the same local
area which was used to detect associations and displacements outside of the direct context of
the experimental setup. These *passive* pairs were placed in the same arrangement as the
experimental feeders, 0.5 m apart, roughly 50 m from the experimental array. Unfortunately,
during the onset of the experiment, a dead jackdaw appeared next to one of the feeder pairs
at site Z, resulting in a near-complete abandonment of all feeders at the site. As such, only
data from site Y was analysed.

Experimental Trials

Birds were given *ad libitum* access to high-quality food (mealworms) from the feeders for two
weeks leading up to the experiment to both facilitate habituation to the new apparatus, as
well as ensure that many adults knew that this food was theoretically available from these
feeders. The experimental treatment was implemented between the 19th July and 13th August
2021, with experimental sessions conducted in the morning for five hours from ~5:30 am.
Outside of these times the experimental feeders were covered in order to prevent access, with
passive feeders accessible at all times. All trials were recorded with Sony Handycam (Sony,
Tokyo, Japan) camcorders for validation purposes (see Supplementary Material:
Displacement Validation).

Treatments

All RFID-tagged juveniles formed the 'knowledgeable' group, giving them unrestricted access

to the *experimental* feeders. Treatment adults (70% of adult birds) were able to access the

mealworm reward from the *experimental* feeders, but only if a juvenile was present on the

connected adjacent *experimental* feeder. Control adults (30% of adult birds) were not able to

use the *experimental* feeders under any circumstances. All three groups could access *passive*

feeders in an unrestricted manner, including both the two *passive* feeders either side of the

experimental feeders and the *passive* feeder pairs at nearby locations. The control adults were

designated to provide baseline data on adult-juvenile interactions with which to compare the

learning of the treatment group and this split was chosen to preserve a meaningful

comparison between these groups whilst not unnecessarily reducing the sample size of

potential treatment adults. However, as seen in the statistical methods, the use of a control

group is not necessary for the detection of a change in within-individual behaviour over time.

Ultimately, adults in the control group visited experimental arrays at a rate too low to conduct

meaningful analyses.

Due to limits in the memory of the microcomputer that controlled the experiment,

only adults that had been seen in the habituation phase of the study were split amongst

treatments, with adults seen in this phase randomly assigned to either 'treatment' or 'control'

at the required 0.7/0.3 probability. Any adults that subsequently interacted with the

experiment joined the control treatment by definition as they were able to access the *passive*

feeders, but never the *experimental* feeders.

Ethical Note

Research was conducted under ethical permission from the Penryn Campus Ethics

Committee (eCORN000406), and in line with Association for the Study of Animal Behaviour

ethical guidelines [22]. Birds were ringed under BTO permit (permit nos. C6449, C5752, C6923

and C6079).

Statistical Methods

Relational Event Models (REMs), a type of time to event model for analysing dyadic network

data, were run to compare changes in the rates of cofeeding and displacement events based

on individual experience (as in [15]). REMs enable analysis to account for the order of events in

a network, preventing data aggregation [23–25], and test hypotheses by comparing events

against a set of 'non-events', unobserved events that could hypothetically happen between individuals in the network [26]. However, non-events are conventionally evenly sampled from the pool of individuals, which is not a realistic assumption for data of interactions between freely interacting wild animals, where some individuals are more represented in the data-stream than others. In line with conventional social network approaches [27], we used a permutation method to generate null datasets that maintained the structure of the data regarding individual activity at the experiment. This method enabled us to account for the varying individual levels of activity seen at the experiment, as well as the opportunities to succeed at the task based on the other individuals present, something we would be unable to incorporate into traditional modelling approaches. We created 10000 permuted datasets to generate predictors of individual experience (number of successful join events by an adult of any juvenile, number of cofeeding events between a specific dyad, number of displacements of any juvenile by an adult). We then ran cox proportional hazard models using the *survival* package [28] in R v4.1.2 [29] between each permuted dataset (non-events) and the observed data, including a strata term for day. We took the median model coefficient of the 10000 models as a point estimate for each parameter, with the 2.5 and 97.5 percentiles of each coefficient taken to generate 95% confidence intervals. All results are reported as Incidence Rate Ratios, which is calculated as the exponential of the cox coefficient.

For testing hypotheses about adult behaviour, all events where an adult was the source of an event (e.g. joined the cofeed or displaced the feeding individual) were randomised, conditioned on the target of the behaviour. This means the identities of the individuals that were joined or displaced were preserved, with the identity of the adult that either joined or displaced them randomised by shuffling. This preserved the structure of opportunities to join or displace, enabling us to ask questions about how individual experience influenced who individuals joined and displaced. Randomisations were constrained within day and feeder pair to preserve the visit structure across the experiment and prevent the generation of scenarios that weren't possible given individual visit history. As such, the estimated differences between permuted and observed datasets are likely to be conservative, producing smaller effects than if using a less constrained permutation procedure. Results from experimental feeders were qualitatively identical when considering adult success as either (a) the number of successful cofeeds the adult joined a juvenile at an experimental feeder, or (b) the total number of successful cofeeds the adult participated in, both joining or being joined by a juvenile (Extended Data Tables 1-2). We present the results from (a) in the main text.

Acknowledgements

We would like to thank the Gluyas family and Odette Eddy for access to their land to conduct the experiment. Thanks to Andy Young for many valuable discussions and Stephanie King for comments.

Funding

J.J.A. was supported by the Biotechnology and Biological Sciences Research Council-funded South West Biosciences (SWBio) Doctoral Training Partnership [680027356]. A.T. and G.E.M. were supported by a Leverhulme grant [RGP-2020-170] to A.T. and N.T. was supported by an SNSF post-doctoral mobility fellowship [P400PB_194397/1].

References

1. Cronin, K. A. & Sánchez, A. Social dynamics and cooperation: The case of nonhuman primates and its implications for human behavior. *Advances in Complex Systems* **15**, 1250066 (2012).

2. Massen, J. J. M., Ritter, C. & Bugnyar, T. Tolerance and reward equity predict cooperation in ravens (*Corvus corax*). *Scientific Reports* **5**, 1–11 (2015).

3. Coelho, C. G., Garcia-Nisa, I., Ottoni, E. B. & Kendal, R. L. Social tolerance and success-biased social learning underlie the cultural transmission of an induced extractive foraging tradition in a wild tool-using primate. *Proceedings of the National Academy of Sciences* **121**, e2322884121 (2024).

4. Cieri, R. L., Churchill, S. E., Franciscus, R. G., Tan, J. & Hare, B. Craniofacial feminization, social tolerance, and the origins of behavioral modernity. *Current Anthropology* **55**, 419–443 (2014).

5. De la Fuente, M. F. *et al.* Foraging networks and social tolerance in a cooperatively breeding primate (*Callithrix jacchus*). *Journal of Animal Ecology* **91**, 138–153 (2022).

6. Testard, C. *et al.* Ecological disturbance alters the adaptive benefits of social ties. *Science*
384, 1330–1335 (2024).

7. Van Schaik, C. P. Local traditions in orangutans and chimpanzees: social learning and
social tolerance. in *The Biology of Traditions* (eds. Fragaszy, D. M. & Perry, S.) 297–328
(Cambridge University Press, Cambridge, 2003).

8. Miller, R., Schiestl, M., Whiten, A., Schwab, C. & Bugnyar, T. Tolerance and Social
Facilitation in the Foraging Behaviour of Free-Ranging Crows (*Corvus corone corone*; *C. c.*
cornix). *Ethology* **120**, 1248–1255 (2014).

9. Arbon, J. J., Hahn, L. G., McIvor, G. E. & Thornton, A. Competition and generalization
impede cultural formation in wild jackdaws. *Proc. R. Soc. B.* **290**, 20230705 (2023).

10. Morand-Ferron, J., Cole, E. F., Rawles, J. E. C. & Quinn, J. L. Who are the innovators? A field
experiment with 2 passerine species. *Behavioral Ecology* **22**, 1241–1248 (2011).

11. Cavalli-Sforza, L. L. & Feldman, M. W. *Cultural Transmission and Evolution: A Quantitative*
Approach. (Princeton University Press, 1981).

12. Heyes, C. Blackboxing: social learning strategies and cultural evolution. *Philosophical*
Transactions of the Royal Society of London B: Biological Sciences **371**, (2016).

13. Mesoudi, A., Chang, L., Dall, S. R. X. & Thornton, A. The evolution of individual and
cultural variation in social learning. *Trends in Ecology and Evolution* **31**, 215–225
(2016).

14. Lee, V. E. & Thornton, A. Animal Cognition in an Urbanised World. *Frontiers in Ecology*
and Evolution **9**, 1–20 (2021).

15. Kings, M. *et al.* Wild jackdaws can selectively adjust their social associations while
preserving valuable long-term relationships. *Nature Communications* **14**, 5103 (2023).

16. Kendal, R. L. *et al.* Social Learning Strategies: Bridge-Building between Fields. *Trends*
Cognit. Sci. **22**, 651–665 (2018).

17. Farine, D. R., Spencer, K. A. & Boogert, N. J. Early-life stress triggers juvenile zebra finches

to switch social learning strategies. *Current Biology* **25**, 2184–2188 (2015).

18. Dunbar, R. I. M. The social brain hypothesis. *Evolutionary Anthropology: Issues, News, and*

Reviews **6**, 178–190 (1998).

19. Ducatez, S., Sol, D., Sayol, F. & Lefebvre, L. Behavioural plasticity is associated with

reduced extinction risk in birds. *Nature Ecology & Evolution* 1–6 (2020)

doi:10.1038/s41559-020-1168-8.

20. Kulahci, I. G. & Quinn, J. L. Dynamic Relationships between Information Transmission

and Social Connections. *Trends in Ecology and Evolution* **34**, 545–554 (2019).

21. Greggor, A. L., McIvor, G. E., Clayton, N. S. & Thornton, A. Wild jackdaws are wary of

objects that violate expectations of animacy. *Royal Society Open Science* **5**, 181070

(2018).

22. ASAB. Guidelines for the treatment of animals in behavioural research and teaching.

Animal Behaviour **159**, i–xi (2020).

23. Tranmer, M., Marcum, C. S., Morton, F. B., Croft, D. P. & de Kort, S. R. Using the relational

event model (REM) to investigate the temporal dynamics of animal social networks.

Animal Behaviour **101**, 99–105 (2015).

24. Patison, K. P, Quintane, E., Swain, D. L., Robins, G. & Pattison, P. Time is of the essence: an

application of a relational event model for animal social networks. *Behavioral Ecology*

and Sociobiology **69**, 841–855 (2015).

25. Butts, C. T. A relational event framework for social action. *Sociological Methodology* **38**,

155–200 (2008).

26. Lerner, J. & Lomi, A. Reliability of relational event model estimates under sampling: How

to fit a relational event model to 360 million dyadic events. *Network Science* **8**, 97–135

(2020).

27. Farine, D. R. & Whitehead, H. Constructing, conducting and interpreting animal social

network analysis. *Journal of Animal Ecology* **84**, 1144–1163 (2015).

28. Therneau, T. M. A package for survival analysis in R. R package version 3.3-0 (2022).

29. R Core Team. R: A language and environment for statistical computing. R Foundation for

Statistical Computing (2019).