Trait-space disparity in fish communities spanning 380 million years from the Late Devonian to present

John Llewelyn[1], John A. Long[2], Richard Cloutier[3], Alice M. Clement[2], Giovanni Strona[4], Frédérik Saltré[1], Michael S. Y. Lee[2,5], Brian Choo[2], Kate Trinajstic[6], Olivia Vanhaesebroucke[3], Corey J. A. Bradshaw[1]

[1]Global Ecology | *Partuyarta Ngadluku Wardli Kuu*, College of Science and Engineering Flinders University, GPO Box 2100, Adelaide, South Australia 5001, Australia
[2]College of Science and Engineering Flinders University, GPO Box 2100, Adelaide, South Australia 5001, Australia
[3]Université du Québec à Rimouski, Rimouski, Québec G5L 3A1, Canada
[4]European Commission, Joint Research Centre, Ispra, Italy
[5]South Australian Museum, North Terrace, Adelaide, South Australia 5000, Australia
[6]School of Molecular and Life Sciences, Curtin University, Bentley, Western Australia 6845, Australia

Abstract

The diversity and distribution of traits in an ecological community determine how it functions. Modern fish communities conserve trait space across similar habitats. However, little is known about trait-space variation through deep time. We examined two Devonian fish communities — a tropical reef (Gogo, Australia) and a tropical estuary (Miguasha, Canada) — and compares them with six modern communities from diverse habitats. Trait-space metrics reflecting the amount of diversity found in each community (functional richness) and the similarity among species in each community (nearest neighbour) indicated Devonian communities had similar scores to modern communities, but were consistently less functionally rich than their closest modern analogues. Metrics describing location in trait space (specialisation, centroid distances, hypervolume overlap) also showed distinctions between Devonian and modern communities. Indeed, location in trait space of Devonian and modern communities differed on several axes associated with a range of traits largely confined to modern communities. Thus, modern fish communities are more functionally diverse in the traits we considered, potentially due to greater evolutionary lability. These results highlight the importance of evolutionary constraints in shaping community trait space and raise questions regarding whether differences between Devonian and modern communities indicate differences in community structure and function.

Key words: functional diversity; palaeoecology; community assembly; evolutionary constraints; trait diversity; fish communities.

Introduction

The combination of a species' traits mediate its interactions with its environment [1,2], so the range and distribution of traits among interacting species is important in determining function and resilience of ecological communities [3–5]. Further, the diversity and distribution of traits in a community — also known as community 'trait space' — can provide insights into the rules of community assembly, including what roles evolution, species filtering, and niche differentiation play in controlling how specific communities form [6–8]. Indeed, recent research on the trait space of contemporary communities has addressed how communities develop and adjust to their ever-shifting environments [9,10]. However, it is still unclear whether patterns observed within and among modern communities also apply to ecological communities over deep time. Consistent spatial and temporal patterns in community trait space would suggest that community assembly rules are constant, whereas variation could indicate phylogenetic constraints [11], stochasticity in evolutionary pathways [12,13], and/or shifts in the environmental parameters that determine community composition [14].

A range of metrics are commonly used to describe and investigate community trait space [15]. Such trait-diversity metrics often indicate variation among communities in different habitats and climates [16,17]. For example, trait diversity in fish (non-tetrapod vertebrate) communities differs depending on substratum (e.g., reef *versus* seagrass *versus* sand) [18–20] and habitat complexity [21,22]. Despite this variation, there is also evidence for the conservation of trait space across similar habitats, even when there are substantial phylogenetic and taxonomic differences among the resident communities [23]. That evolutionarily distinct communities occupying similar habitats have similar trait spaces suggests convergence (through evolution and/or niche filtering) and highlights the importance of deterministic processes in shaping community trait space (i.e., environment determining trait space) [24,25].

Although trait space varies among habitats and is conserved within habitat types across locations in contemporary communities, it remains unclear how it varies through deep time. Few studies have examined changes in functional diversity and trait space over millions of years (but see [26] for impacts of extinction events on ecospace). By testing for patterns across time as well as across contemporaneous habitats, it is possible to test whether similar habitats select for similar community trait space irrespective of phylogeny and evolutionary history.

The Devonian Period (419.2 to 358.9 million years ago [Ma]) is popularly termed the "Age of Fishes". During that time, diverse fish communities established and radiated across the planet and gave rise to the first tetrapods (four-legged vertebrates). Four mass extinctions have since occurred (not including the current one) [27], which along with the passing of time and millions of 'background' extinction and speciation events, have caused massive phylogenetic turnover [28,29]. In addition, Devonian and modern fish communities differ in terms of the non-fish species and nutrient inputs that make up the wider ecological community [30,31]. For example, Devonian reefs were primarily built of stromatoporoid sponges, calcareous algae, and microbial communities rather than hexacorallian corals (but also included some rugose and tabulate corals) [32,33], and ancient estuaries generally had low invertebrate and algal diversity compared to modern estuaries [34,35]. Thus, although the environments Devonian fish communities experienced were physically similar to their modern counterparts, their vertebrate and non-vertebrate biota differed greatly. By comparing these ancient assemblages to modern communities, it is therefore possible to test if physical environment is the primary determinant of fish community trait diversity across space and time, or whether biological, phylogenetic, and evolutionary differences associated with the passing of hundreds of millions of years have a dominant effect on trait space. Such knowledge will improve our understanding of how communities have responded to change in the past, and how they could change in response to anthropogenic and non-anthropogenic disturbances in the future.

We compared the trait space of two of the best-described Devonian fish communities — a tropical reef (Gogo Formation, northern Australia) and a tropical estuary (Miguasha, eastern Canada) — to that of six modern fish communities from different habitats and climate zones, to identify how trait diversity varies through space and time and thereby shedding light on the processes of community assembly. We compared these communities in terms of two trait-space metrics that reflect diversity and distribution of traits within each community (functional richness and nearest neighbour) and three that quantify each community's position in trait space relative to other communities (specialisation, distance between hypervolume centroids, overlap of hypervolumes) [36]. We hypothesise that (*i*) if community trait space is primarily determined by habitat, there will be larger differences between communities from different habitats (i.e., reef *versus* estuary *versus* freshwater) than between Devonian and modern communities from the same habitat (e.g., Devonian reef community similar to modern reef community) or between tropical and temperate/subtropical communities from the same habitat (e.g., tropical reef similar to temperate reef). (*ii*) If

community trait space is primarily determined by climate, there will be larger differences

between tropical *versus* temperate/subtropical communities than between communities from

different periods of time or different habitat types within the same climate zone. (*iii*) If

community trait space is primarily determined by evolutionary history and the phylogenetic

makeup of its constituent species, there will be larger differences between Devonian and

modern communities compared to between communities from different habitat types or

different climate zones within the same period. (*iv*) If time, habitat, and climate all

substantially affect community trait space, there will be consistent groupings/patterns across

all these variables (e.g., community metric scores will group together based on time, habitat,

and climate); and (*v*) If community trait space is independent of time, habitat, and climate,

trait space metrics will not be grouped by period (Devonian *versus* modern), habitat or

climate zone. By identifying which of these five patterns are observed in trait space metrics,

we provide insight into the factors shaping community trait space (i.e., community assembly)

in the past, present and future.

Methods

Sites and species lists

We compared two Late Devonian (Frasnian) and six modern fish communities. The Late

Devonian communities of Miguasha (Escuminac Formation) in north-eastern Canada and the

Gogo Formation in north-western Australia are two of only six recognised fish Lagerstätten

from this period [37]. We selected these communities based on (1) their fossil records

capturing most of the *in situ* fish diversity, and (2) the availability of trait data for a range of

traits and species that make up these communities (see electronic supplementary material

Table S1). Miguasha (modern: 48° 48′ N, 66° 16′ W) was a tropical estuary in Laurussia 385

to 374 million years ago [38–40]. Thousands of fish specimens have been preserved and

recovered from Miguasha, representing at least 19 species [39]. The Gogo Formation

(modern: 18° 26′ S, 125° 55′ E) represents the basinal and channel facies adjacent to a

tropical, stromatoporoid sponge-dominated reef that was on the edge of the Gondwanan

landmass 384 to 382 million years ago; the fish preserved in the Gogo Formation were

deposited under anoxic marine conditions and came from the adjacent reef [33]. Fifty-three

fish species have been identified from Gogo (including 47 described and 6 undescribed

species; Electronic supplementary material Tables S1 and S2) [33,41]. Discovery curves

suggest all (fossilised) fish species from Miguasha have been described [34] and that most (>

95%) fish species from Gogo have been identified (see electronic supplementary material Fig. S1; Table S2).

We selected six modern fish communities for comparison to capture fish assemblages from a diversity of environments (different habitat types and climate zones). Our selection was further guided by the availability of species lists for the communities and the presence of detailed information on trophic interactions (allowing for later development of trophic-network models of those communities). We included reef, estuarine, and freshwater communities, with tropical and temperate or subtropical representatives of each (i.e., we included each habitat-type/climate-zone combination). The modern communities were: Caribbean reefs (tropical; 18° 3′ N, 65° 28′ W) [42]; Chilean reefs (temperate/subtropical; 31° 30′ S, 71° 35′ W) [43]; Santa Cruz Channel, north-eastern Brazil (tropical estuary; 7° 46′ S, 34° 53′ W) [44,45]; Ythan Estuary, Scotland (temperate estuary; 57° 19′ N, 1° 59′ W) [46–48]; Braço Morto Acima and Abaixo, central Brazil (tropical freshwater oxbow lakes; 19° 41′ S, 56° 59′ W) [49–58]; and the Nepean River, New South Wales, Australia (temperate/subtropical freshwater river section; 33° 47′ S, 150° 38′ E; ala.org.au; see electronic supplementary material Table S3) [59].

Traits

We collected information on 13 traits from the 488 species in our study (416 extant species and 72 Devonian species; see electronic supplementary material Table S4). We chose traits based on their ecological relevance and their availability for both living and extinct species. The traits were: sagittal body shape, transverse body shape, maximum total body length (tip of snout to tip of caudal fin), standard body length, head length, eye diameter, pre-orbital length, body depth, position of mouth, position of eyes, presence of mandible, presence of spiracle, and caudal fin shape (Fig. 1; see electronic supplementary material Table S4). We expressed all morphometric variables (except maximum total body length) as a proportion of maximum total body length as a size standardisation and to reduce correlations among these traits. Due to inconsistencies in how standard body length is measured across different body shapes and taxonomic groups, we later removed that measure from our analyses. We also removed 'presence of mandible' because these data were highly uneven (4 without mandibles *versus* 483 with), which causes problems when calculating distances in multidimensional space. Thus, our final database included 11 traits for 488 species.

Figure 1. Fish traits used in the trait space analysis. An example from each category is shown for the categorical

traits, whereas trait extremes are shown for continuous traits. Sagittal body shapes include (a) short and deep,

(b) elongated, (c) eel-like, (d) fusiform, and (e & f) other. Transverse body shapes include (g) circular, (h) oval,

(i) compressed, (j) flattened, and (k) compressed and lies on side. Eye positions included (l & m) lateral, (n)

raised/top of head, and (o) migrated to one side and raised. Large spiracles could be present (p) or absent.

Caudal fin shapes included (q) heterocercal, (r) homocercal, (s) hypocercal, (t) isocercal, (u) protocercal, (v)

leptocercal, (w) reduced or absent, and (x) whip-like. Mouth positions included (y) terminal, (z) superior, and

(aa) subterminal/inferior.

We collected trait data for Devonian species from the literature, specimens, and

photographs of specimens. However, it was not possible to collect all traits for the species

from the Gogo Formation because some of them are only known from incomplete or

disarticulated body parts (due to the preservational environment) [60]. We addressed these

gaps by estimating traits using: (1) expert opinion in cases where traits could be confidently

inferred (inferences made by J. Long, A. Clement, B. Choo, and K. Trinajstic) or (2) multiple

imputation using species' traits, coarse taxonomy, and the `missForest` R package [61,62].

Evaluating multiple imputation performance indicated out-of-bag error estimates of 0.30

(normalised root-mean-square error) and 0.11(proportion of falsely classified) for continuous

and categorical variables, respectively. For modern species, we extracted trait data from

Fishbase (using the `rFishbase` package) [63], photographs, and scientific literature [62].

We visually inspected the distribution of continuous traits, identifying two as right-skewed

(total length and pre-orbital length); we $\log_e$-transformed these traits to normalise them.

Analyses

We built a Gower dissimilarity matrix [64] to quantify trait differences among species using

the `gawdis` package in R [65]. Gower dissimilarity can handle mixed data types (our trait

data included continuous and categorical variables) and can normalise the contribution of

variables [6,64]. Including all traits resulted in an unbalanced distribution, so we also

calculated Gower distances with the maximum number of traits that did not result in uneven

distributions (there were three such combinations of nine traits). Observed patterns in

functional diversity metrics were consistent across these trait sets (see electronic

supplementary material Fig. S2 and S3). We therefore only present results from the 'all traits'

Gower distance trait set in the main text.

We applied principal coordinate analysis (PCoA) to the Gower distances to ordinate

species within a lower-dimensional space and calculate functional diversity metrics using the

`mFD` R package [66]. The first eight principal coordinate axes explained 73% of the total

variation [62]; we identified that eight was the optimal number of axes based on scaled mean

absolute deviation and scaled root of mean square error (minimising the deviation between

trait-based distances and functional space distances) [62]. We therefore used these eight axes

to calculate each community's functional richness (proportion of functional space filled by

the community), nearest neighbour (average distance in trait space to the

nearest functional neighbour in the same community), and specialisation (average distance to

the global centroid). These metrics calculated from PCoA coordinates are unitless 'scores'

that can be used to compare communities.

Species diversity varied substantially among the included communities, ranging from 17

species in Ythan Estuary to 208 species in Caribbean reefs (Electronic supplementary

material Tables S1 and S3). Functional diversity metrics can be correlated with species

diversity [15,67], so we recalculated the three functional diversity metrics (functional

richness, nearest neighbour, and specialisation) controlling for species diversity to test which,

if any, patterns in these metrics persisted after removing species diversity-linked associations.

To control for species diversity, we sub-sampled 10 species from each community and

calculated the functional diversity metrics based on these sub-samples, replicating the process

100 times and taking the mean score for each metric for each community across the

replicates. We plotted both the 'raw' and species-diversity controlled metrics, allowing us to

evaluate visually whether period, habitat type, or climate zone grouped communities in any

of these metrics.

We also used the principal coordinates to build trait hypervolumes for each community

using Gaussian kernel density estimation and the `hypervolume` package in R [68]. We set

the requested quantile to 0.95 and estimated the bandwidth for each axis in each community

separately. We then used the mean estimated bandwidth for each axis across the eight

communities as fixed bandwidths to recalculate hypervolumes (i.e., ensuring the same

bandwidths for each community). The resulting hypervolumes were used to quantify

distances between community centroids (geometric centre) and overlap in multidimensional

trait space (Jaccard index). We made these comparisons between all pairwise community

combinations to test if communities were grouped/clustered according to their period, habitat,

or climate zone (see hypotheses listed in the introduction). We did not apply the sub-

sampling technique as for the functional diversity metrics to these hypervolumes because (a)

the average centroid from replicated sub-samples of a community is approximately the same

as the centroid when including all species in the community (i.e., no change in distances

between centroids), (b) when comparing overlap in trait space between communities (Jaccard

index) to assess their similarity, it is more intuitive to use all species instead of sub-sampling

species, and (c) the Jaccard index is a proportionate measure. Because distances between

centroids suggested the fish community trait spaces were segregated, we identified the

specific traits responsible by determining along which hypervolume axes the centroids

differed, and which traits were correlated with these axes. We then investigated how these

traits were distributed among the fish communities.

Results

Functional richness of the Devonian communities fell within the range observed in modern

communities, irrespective of whether we controlled for species richness (Fig. 2a,b; not

controlling for species richness: Gogo = 2.4×10^{-3}, Miguasha = 4.7×10^{-4}, modern

assemblages = 2.6×10^{-5} to 0.29; controlling for species richness: Gogo = 3.6×10^{-7},

Miguasha = 4.1×10^{-6}, modern assemblages = 2.2×10^{-7} to 4.6×10^{-6}). However, the

Devonian communities had lower functional richness than their closest modern counterparts

(Gogo *versus* Caribbean: 2.4×10^{-3} *vs.* 0.29 [not controlling for species richness] and $5.6 \times$

10^{-7} *vs.* 3.8×10^{-6} [controlling for species richness]; Miguasha *versus* Santa Cruz Estuary 4.7

$\times 10^{-4}$ *vs.* 4.5×10^{-2} [not controlling for species richness] and 4.1×10^{-6} *vs.* 4.6×10^{-6}

[controlling for species richness]). There was habitat-linked variation in functional richness,

but the pattern differed depending on whether we controlled for species diversity (Fig. 2a,b).

Reef communities had the highest score followed by estuarine communities when all species

were used to calculate functional richness (Fig. 2a), whereas estuarine communities had the

highest scores followed by reefs after controlling for species diversity (Fig. 2b). This habitat-

functional richness pattern, including the switching between reefs and estuaries as the most

functionally rich communities when controlling for species diversity, emerged in both

modern and Devonian communities (Fig. 2a,b). Among the modern assemblages, tropical

communities consistently had higher functional richness than their temperate/subtropical

counterparts (mean richness tropical *versus* temperate/subtropical: 0.11 vs. 0.0006 and 3.0 $\times$

10^{-6} vs. 2.3×10^{-6} for richness calculated without controlling for species diversity and

controlling for species diversity, respectively; Fig. 2a,b).

Figure 2. Functional diversity metrics for two Devonian and six modern fish communities. Three functional diversity metrics are shown, including: functional richness (a. and b.; volume of functional trait space filled by species in a community); nearest neighbour (c. and d.; mean distance in trait space to nearest neighbour in the same community); and functional specialisation (e. and f.; mean distance of species in a community to the mean position of species from the global pool). Panels in the left column show results when all species in each community are included to calculate these metrics (i.e., not controlling for species diversity), whereas panels in the right column show results controlling for species diversity using resampling. Pink background indicates the five tropical communities (ancient and modern), yellow background indicates the three temperate/subtropical communities (modern only). For display purposes, we scaled functional richness values in the controlled-diversity plot by multiplying by 100,000 (functional metrics calculated from principal coordinates are unitless and indicate relative differences).

Nearest-neighbour scores, reflecting the distance in functional space of each species to the nearest species in its community, were similar for Devonian and modern communities (Fig. 2c,d). Miguasha had a slightly higher score than the other communities when not controlling for species diversity (Fig. 2c; Miguasha = 0.421, Gogo = 0.207; modern assemblages = 0.181 to 0.411). After accounting for species diversity, nearest-neighbour scores for Gogo and Miguasha fell within the range observed in the modern communities (Fig. 2d; Gogo = 0.414, Miguasha = 0.544; modern assemblages = 0.413 to 0.554).

There were differences among habitats in nearest-neighbour scores within periods and climate zones. Freshwater communities consistently had the lowest nearest-neighbour scores (Fig. 2c,d). Estuarine communities consistently had the highest scores followed by reef communities, except in the tropical, species diversity-controlled comparison where these communities had similar scores (Fig. 2 c,d). Further, there was a latitudinal pattern among modern communities when not controlling for species diversity — species in temperate and subtropical communities were farther from their nearest neighbour than were species in tropical communities (Fig. 2c) — but this latitudinal pattern disappeared after controlling for species diversity (Fig. 2d).

The Devonian communities had the highest functional specialisation, irrespective of whether or not we controlled for species diversity (Fig. 2e,f; not controlling for species diversity: Miguasha = 0.555, Gogo = 0.496, modern communities = 0.393 to 0.445; controlling for species diversity: Miguasha = 0.556, Gogo = 0.500, modern communities = 0.391 to 0.444). High specialisation indicates species in these communities are farther from the global (combined communities) mean position/centroid. Similarly, the distances between hypervolume centroids of Devonian *versus* modern communities were greater than the distances between centroids of communities from the same period (Fig. 3a,b; distance between Devonian and modern: 0.16 ± 0.01, distance between communities from the same period: 0.07 ± 0.02; all values are means $\pm$ SD). Centroids of communities from the same

habitat type or the same climate zone were not closer together than centroids from different

habitats or climate zones (Fig. 3c,d).

Figure 3. Distances between centroids of functional trait space hypervolumes for fish communities. Two
Devonian (Gogo and Miguasha) and six modern fish communities from different habitats and climates are
compared. On the left, plot a. is a heatmap showing distances between the centroids of each community. On the
right are three violin plots showing distances between communities from either the same (within) or different
(between): b. periods, c. habitat types, and d. climate zones. Hypervolumes were built using species coordinates
on the first 8 axes from a principal coordinate analysis applied to a Gower dissimilarity matrix of the species'
traits. Plot points in the violin plots represent individual comparisons between communities, jittered on the *x*
axis for clarity.

The hypervolume Jaccard index measuring trait-space overlap also indicated less similarity

between Devonian and modern communities compared to between communities from the

same period (Fig. 4a,b; overlap between Devonian and modern: 0.05 ± 0.02, overlap between

communities from the same period: 0.18 ± 0.07; all values are means $\pm$ SD). The overlap

index between Devonian communities was within the range observed among modern

communities (Devonian overlap = 0.16, modern community overlap = 0.09-0.37; Fig. 4a).

There was no evidence of greater overlap between hypervolumes of communities from the

same *versus* different habitat types or climate zones (Fig. 4c,d).

333
Figure 4. Jaccard Index comparing overlap in functional trait space hypervolumes for fish communities. Two
Devonian (Gogo and Miguasha) and six modern fish communities from different habitats and climates are
compared. On the left, plot a. is a heatmap showing Jaccard indices for all community comparisons. On the right
is a series of violin plots showing Jaccard indices between communities from either the same (within) or
different (between): b. periods, c. habitat types, and d. climate zones. Hypervolumes were built using species
coordinates on the first 8 axes from a principal coordinate analysis applied to a Gower dissimilarity matrix of
the species' traits. Plot points in the violin plots represent individual comparisons between communities, jittered
on the *x* axis for clarity.

342
343
344 The centroids for both Devonian communities were outside the range observed in modern

345 communities on axis 1 (Devonian ≤ -0.089, modern ≥ -0.045), axis 2 (Devonian ≥ 0.061,

346 modern ≤ 0.009), and axis 7 (Devonian ≥ 0.026, modern ≤ -0.003; see electronic

347 supplementary material Table S5).

348
349
Although directly linking the principal coordinate axes to traits is not possible because the

axes are based on Gower distances between species, correlations between species coordinates

along these axes and traits suggest they mostly reflect eye diameter, body depth, and caudal

fin shape on axis 1; position of mouth and eye position on axis 2; and transverse body shape

and presence of spiracles on axis 7 (Table 1). Accordingly, Devonian communities were

distinct from modern communities in these traits. Devonian fishes tended to have smaller eye

diameters and their bodies were not as deep relative to their length (axis 1; Fig. 5c,e). None

of the fish in these Devonian communities had superior mouth position whereas all the

modern communities had some fish with this mouth position (axis 2; Fig. 6e). None of the

Devonian fish had eyes migrated to one side of the body (as in flatfish; axis 2; Fig. 6f),

spiracles and heterocercal tails were more prevalent in the Devonian than modern

communities (axis 1 and 7, respectively; Fig. 6b,d), and the proportion of fish with a

compressed transverse body shape is lower in the Devonian fishes (axis1; Fig. 6c).

Gogo's centroid was also extreme on axes 5 and 6 (with values of 0.056 and 0.019,

respectively, compared to values ≤ 0.002 and ≤ 0.017 for the other communities on the same

axes; see electronic supplementary material Table S5), while Miguasha's centroid was

extreme on axis 4 (Miguasha = -0.049, other communities $\geq$ -0.026; see electronic

supplementary material Table S5). Sagittal (not transverse) body shape was the most strongly

correlated trait with these axes, showing a slightly stronger correlation with axis 5 (Table 1)

than with axes 4 and 6 (sagittal body shape-axis correlation: $\eta^2 = 0.311$, p < 0.000 and $\eta^2 =$

0.252, p < 0.000 for axes 4 and 6, respectively; see electronic supplementary material Table

S6). These correlations suggest that the extreme positions of Gogo and Miguasha could be

attributed to Gogo's lack of eel-shaped and elongate (e.g., anguilliform) fish and to

Miguasha's higher occurrence of fish with unconventional ('other') body shapes (Fig. 6a),

which included the tadpole-shaped *Escuminaspis laticeps* and *Levesquaspis patteni*.

Table 1. Correlation between traits *versus* species coordinates on principal coordinate analysis axes, showing
each trait and the axes with which it had its highest correlation (for all trait/axis correlations, see electronic
supplementary material Table S6). We applied Kruskal-Wallis tests to categorical traits and linear models
(regressions) to continuous traits. r^2 and η^2 describe the proportion of variance explained by the Kruskal-Wallis
and linear models, respectively. Type I error rates (p-values) were < 0.0001 in all cases. See Fig. 1 for trait
details.

Trait	axis	test	statistic	value
body depth	PC1	linear model	r^2	0.537
sagittal body shape	PC5	Kruskal-Wallis	η^2	0.397
transverse body shape	PC7	Kruskal-Wallis	η^2	0.378
caudal fin shape	PC1	Kruskal-Wallis	η^2	0.339
eye diameter	PC1	linear model	r^2	0.623
eye position	PC2	Kruskal-Wallis	η^2	0.371
head length	PC3	linear model	r^2	0.335
$\log_e$(pre-orbital length)	PC3	linear model	r^2	0.493
$\log_e$(total body length)	PC3	linear model	r^2	0.431
mouth position	PC2	Kruskal-Wallis	η^2	0.415
spiracle	PC7	Kruskal-Wallis	η^2	0.221

386
387 **Figure 5.** Violin plots of five continuous fish traits from two Devonian and six modern communities. Total
388 body length is in centimetres and transformed using the natural logarithm (log_e) to normalise. All other traits are
389 expressed as a proportion of total body length (log_e-transformed for pre-orbital length). Pink background
390 indicates the five tropical communities (ancient and modern), yellow background indicates the three
391 temperate/subtropical communities (modern only).

Figure 6. Stacked bar charts of six categorical fish traits from two Devonian and six modern communities: a. sagittal body shape, b. presence/absence of spiracles, c. transverse body shape, d. caudal fin shape, e. mouth position, and f. eye position. The *y* axis indicates the percentage of species in each community with each category of the trait.

Discussion

Our analysis of modern and ancient fish communities revealed patterns in community trait space associated with period, habitat type, and climate zone. The importance of these variables differed among trait-space metrics, indicating that different factors are important for different trait-space attributes. The two metrics that reflect diversity and distribution of traits within a community (functional richness and nearest neighbour) suggest: (1) broad similarity between Devonian and modern assemblages, but with Devonian communities having consistently lower functional richness than their closest modern counterparts, (2) an association with habitat type, and (3) an association with climate zone (Fig. 2; consistent with hypothesis *iv*). In contrast, period was the only variable that clearly differentiated fish assemblages in the three metrics that reflect a community's position in trait space relative to other communities (functional specialisation, distance between hypervolume centroids, and Jaccard index); we observed no associations between these metrics and habitat type or climate zone (Fig. 3e,f, 4, 5; consistent with hypothesis *iii*). The trait-space positions of Devonian communities were distinct from modern communities on several axes (see electronic supplementary material Table S5) associated with seven of the eleven traits we included in our analyses (Table 1); such trait differences could indicate disparity between Devonian and modern communities in terms of their structure and function. Further, that communities from the same period have similar trait spaces, while those from different periods are dissimilar (consistent with hypothesis *iii*), suggests evolutionary constraints and stochasticity in evolutionary pathways are potentially important in determining community trait space and community assembly.

Period

If the fish community trait space has changed through time, we would expect to see communities from different periods segregated in terms of their trait-space metrics (i.e., modern communities similar to each other and different to Devonian communities). This pattern is clear in four of the five metrics we calculated: Devonian communities consistently differed from modern communities in terms of functional richness, specialisation, centroid location, and hypervolume overlap (Fig. 2, 3, 4).

Although functional richness of Devonian communities was within the range spanned by

all modern communities, the scores of Devonian communities in this metric were consistently

lower than those observed in modern communities from the same habitat type and climate

zone (i.e., Gogo Reef *versus* Caribbean Reef, Miguasha *versus* Santa Cruz Estuary; Fig.

2a,b). Despite the Devonian period being popularly named the 'Age of Fishes', Devonian fish

communities tended to have lower species diversity compared to modern fish communities;

our results suggest they also had lower trait diversity compared to their modern counterparts

(Fig. 2a,b). Discovery curves confirm that most species in the Devonian communities we

examined have been described (see electronic supplementary material Table S2, Fig. S1 and

[34]), and, therefore, the relatively low functional and species richness in Devonian

communities compared to their modern analogues are probably real features instead of

preservational artefacts. Indeed, functional richness was lower for Devonian communities

even after controlling for species diversity (Fig. 2b), providing further support for lower

functional richness in those communities. However, it is not clear whether the entire

'missing' trait space in these Devonian fish assemblages was vacant. It is possible that non-

fish taxa filled some of that space. For example, conodonts (early, non-fish, but fish-like,

jawless vertebrates) might have occupied the trait space of modern lampreys, hagfish, and

small eels [69,70], while invertebrates such as Eurypterids (sea scorpions) might have filled

trait and trophic spaced occupied by predatory fishes in modern communities [71]. Similarly,

non-fish taxa, such as sea snakes, turtles, and marine mammals, make unique contributions to

the trait space in modern aquatic communities [72].

Functional specialisation, centroid distances, and the Jaccard index (trait-space overlap)

are metrics that reflect the location of a community's trait space relative to other

communities. The fact that Devonian and modern communities are separated by these metrics

indicates variation in trait values (continuous traits), trait status (categorical traits), and/or

trait combinations. By examining which hypervolume axes separated modern and Devonian

communities, and which traits were correlated with these axes, we were able to identify the

traits involved in differentiating the trait spaces of these communities. Seven of the eleven

traits we included in our analysis appear to contribute to this differentiation, including: eye

diameter (Fig. 5c); body depth (Fig. 5e); presence of large spiracles (Fig. 6b); transverse body

shape (Fig. 6c); caudal fin shape (Fig. 6d); mouth position (Fig. 6e); and eye position (Fig.

6f). Of these traits, eye diameter, body depth, mouth position and eye position had lower

diversity in Devonian compared to modern communities (Fig. 5,6). Thus, differences

between modern and Devonian communities as indicated by specialisation, centroid

distances, and the Jaccard index reflect, at least in part, trait space that is absent in the

Devonian fish, consistent with the observed differences in functional richness between

communities from these periods (Fig. 2a,b).

Community trait space is linked to community structure and function [73,74]. Our trait

space results therefore suggest substantial ecological differences between Devonian and

modern fish communities. For example, the dearth of Devonian fish with a superior mouth

position — an adaptation to feeding from above [75,76] — indicates differences in trophic

interactions. While superior mouth position has been documented in Devonian species

[77,78], the rarity of this trait in the Devonian contrasts with its widespread occurrence in

modern communities (Fig. 6e). Greater diversity of mouth position does not only indicate a

greater variety of feeding strategies used today, but also suggests that species in modern

communities experience different predation pressures to those present in the Devonian (e.g.,

in terms of predator archetypes) [79], which could lead to the evolution of distinct

antipredator strategies. Similarly, the lack of fish with deep, laterally compressed bodies in

the Devonian communities (indicated by the body depth and transverse body shape traits;

Fig. 5e and 6c) likely indicates ecological differences associated with microhabitat use and

how a species interacts with conspecifics and other species [80,81]. Deep-bodied, laterally

compressed jawless fish have been documented in older communities (Late Silurian and

Early Devonian) [82], but they were less widespread compared to deep-bodied fish in modern

communities, and they appear to have gone extinct before the Late Devonian (i.e., the epoch

we examined). Thus, the relatively low variation of traits in Late Devonian communities

suggests they were less ecologically/functionally diverse overall than modern fish

communities.

The greater trait variation we observed in modern *versus* Devonian fish communities

could result from (*i*) increased habitat/environmental complexity through time [83], (*ii*) the

longer time modern fish assemblages have had to evolve/diversify [84], and/or (*iii*) the

morphological flexibility of teleost fish, a group that did not evolve until the Triassic Period

(> 100 million years after the Devonian; although greater variation in actinopterygian traits

started to emerge immediately following the Hangenberg extinction event at the end of the

Devonian) [85]. In modern fish communities, there is a positive correlation between habitat

complexity and functional richness [22,86]. Thus, given that Devonian habitats appear to

have been less complex than modern habitats (e.g., even the most complex Devonian reefs

tended to be less structurally complex than modern coral reefs [87] and diversity of

benthic/habitat-forming organisms was lower [88]), this factor might contribute to the lower

trait diversity in Devonian fish. Conversely, teleosts have several traits that foster

morphological flexibility. For example, the prehensile mouth parts of teleosts give great

flexibility in mouth-linked traits, leading to the evolution of different mouth positions and

head shapes [89]. Other attributes that promote morphological flexibility in teleosts and other

osteichthyans — fish that are more prevalent in modern communities compared to Devonian

ones — include the presence of a swim bladder for buoyancy control and bony

endoskeletons, facilitating diversification in movement patterns and body shape [90,91].

While our results suggest higher trait diversity in modern fish communities, we acknowledge

that the choice of traits and the groupings used for categorical variables could influence

conclusions regarding which taxa or communities are most diverse [6]. For example, if we

had focused on bony armour plates and lobe-paired fin structure, Devonian communities

would be classified as more trait-diverse than modern communities [92]. Regardless, our

main conclusion — that trait space of modern and Devonian communities differ — should

hold true for any broad sample of traits across the entire phenotype.

Habitat

Functional nearest-neighbour scores did not show consistent differences between periods, but

did display variation with habitat type: freshwater communities consistently had the lowest

scores (their constituent species were the most similar to each other), while estuarine

communities had the highest nearest-neighbour scores in most cases (Fig. 2c,d). The low

functional richness and low nearest-neighbour scores in the freshwater communities we

examined (i.e., indicating low trait diversity and high similarity among species; Fig. 2a,b,c,d)

is likely the result of strong niche filtering, corroborating previous conclusions that habitat

filters determine species (and therefore trait) composition in freshwater communities [93,94].

Reef communities in both periods have high functional richness and low-to-intermediate

nearest-neighbour scores when not controlling for species diversity (Fig. 2a, c), suggesting

that many, densely concentrated niches are facilitated by complex reef habitats (corroborating

research on modern communities) [95]. Estuarine communities have the highest nearest-

neighbour scores in most cases (Fig. 2c,d), indicating that their species are distinct from each

other in trait space. The trait variation among fish in estuarine communities might reflect that

these assemblages are an admixture of species from different origins — they include marine,

freshwater, and diadromous fish, as well as species that complete their life cycle in estuaries

— and therefore include species that have experienced different niche/environmental filters

[96,97]. The association between nearest-neighbour scores and habitat type, and the lack of a

consistent difference between communities from different periods, suggests that
environmental factors are the primary determinants of how similar species within a
community can be to each other. Furthermore, the high nearest-neighbour scores in Devonian
and modern estuarine communities (Fig. 2c,d) suggest low trait and functional redundancy
[98,99]. Communities with low redundancy might be particularly sensitive to the loss of
species [100], depending on the network of species' interdependencies supporting the
community [101,102]. Indeed, previous studies have reported low function redundancy in
reef-fish communities [5,18]; our results suggest even lower redundancy in estuarine
communities.

We also found that functional richness varied with habitat type, but the functional
richness-habitat pattern depended on whether we controlled for species diversity (Fig. 2a,b).
Across climate zones, reef communities had the highest functional richness followed by
estuarine communities when all species were included, whereas estuarine communities had
the highest richness followed by reef communities when controlling for species diversity
(Fig. 2a,b). We observed this switching between which habitat type had the highest
functional richness in modern and Devonian communities (Fig. 2a,b). High habitat
complexity could explain the elevated species and functional richness in reef communities by
allowing greater niche differentiation among resident fish [95]. Indeed, that species richness
was higher for reef *versus* estuarine communities (e.g., 208 *versus* 86 species for tropical
communities, and 27 *versus* 17 species for temperate/subtropical communities) also
potentially explains why controlling for species diversity switches which habitat type had the
highest functional richness. The difference might result from saturation of trait space in
species-rich reef communities (i.e., the expansion of a community's trait space associated
with the addition of a species diminishes as more species are added to the community)
[6,103–106], a hypothesis supported by the low nearest-neighbour scores for the reef
compared to estuarine communities when ignoring species diversity (Fig. 2c).

Climate
Functional richness of modern tropical communities was higher than that of modern
temperate and subtropical communities, consistent with the previously reported pattern of
increasing functional richness with decreasing latitude in fish, other vertebrates, and
invertebrates [106–111]. Our results also revealed a latitudinal/climate pattern among modern
communities in terms of their nearest-neighbour scores, with tropical communities having
lower scores than their temperate/subtropical counterparts when ignoring species diversity

(Fig. 2c). Together, functional richness and nearest-neighbour scores indicate that fish

communities in the tropics have greater variation in traits and their constituent species are

packed more closely together in trait space than observed in temperate and subtropical

communities (where species richness also tends to be lower) [107,112]. We were unable to

assess whether latitudinal patterns occurred among the Devonian fish communities because

both Devonian communities in our study were tropical at deposition. Thus, future research

could test whether latitude (climate) has been important in determining community trait space

through time, or whether this pattern developed recently.

Conclusion

The differences we detected between modern and Devonian fish communities indicate that

the trait space has changed through time. Although the Devonian period is known for its

diverse and abundant fish fauna, our results suggest that modern fish communities are in fact

more trait-diverse than their Devonian counterparts, and that their trait spaces are centred in

different locations in trait space. These differences could indicate several phenomena,

including the longer time modern communities have had to develop and diversify [84],

greater variation in habitats today [88,113], phylogenetic constraints of different fish lineages

[114], and/or stochasticity in evolutionary pathways [115,116]. Irrespective of the root

cause(s), the observed differences in trait space suggests Devonian communities were

structured and functioned differently to modern communities. In other words, fish in these

Devonian communities likely interacted with each other and their environment in different

ways to that observed in modern communities. Further research on how community structure

and function has changed through time, combining information from ancient and modern

communities [117], could provide important insights into community-assembly rules and the

ecological and evolutionary responses of communities to environmental disturbances in the

past and the future.

Acknowledgements

We thank S. Libralato, R. Angelini, A. Pérez-Matus , S. Villéger, C. Brauer for help

accessing species lists and trait data for modern fish communities. We thank T. Llewelyn for

providing helpful feedback. We also acknowledge the traditional owners of the lands on

which this research was done, including the Gooniyandi, Kaurna, and Mi'kmaq peoples. This

work was supported by the Australian Research Council [DP220100825].

References

599   1. McGill BJ, Enquist BJ, Weiher E, Westoby M. 2006 Rebuilding community ecology from functional traits. *Trends in Ecology & Evolution* **21**, 178–185. (doi:10.1016/j.tree.2006.02.002)

2. Kiørboe T, Visser A, Andersen KH. 2018 A trait-based approach to ocean ecology. *ICES Journal of Marine Science* **75**, 1849–1863. (doi:10.1093/icesjms/fsy090)

3. Cadotte MW. 2017 Functional traits explain ecosystem function through opposing mechanisms. *Ecology Letters* **20**, 989–996. (doi:10.1111/ele.12796)

4. McLean M, Auber A, Graham NAJ, Houk P, Villéger S, Violle C, Thuiller W, Wilson SK, Mouillot D. 2019 Trait structure and redundancy determine sensitivity to disturbance in marine fish communities. *Global Change Biology* **25**, 3424–3437. (doi:10.1111/gcb.14662)

5. Micheli F *et al.* 2014 High vulnerability of ecosystem function and services to diversity loss in Caribbean coral reefs. *Biological Conservation* **171**, 186–194. (doi:10.1016/j.biocon.2013.12.029)

6. Mouillot D *et al.* 2021 The dimensionality and structure of species trait spaces. *Ecology Letters* **24**, 1988–2009. (doi:10.1111/ele.13778)

7. Schöb C, Butterfield BJ, Pugnaire FI. 2012 Foundation species influence trait-based community assembly. *New Phytologist* **196**, 824–834. (doi:10.1111/j.1469-8137.2012.04306.x)

8. Weiher E, Clarke GDP, Keddy PA. 1998 Community assembly rules, morphological dispersion, and the coexistence of plant species. *Oikos* **81**, 309–322. (doi:10.2307/3547051)

9. Frimpong EA, Angermeier PL. 2010 Trait-based approaches in the analysis of stream fish communities. *American Fisheries Society Symposium* **73**, 109–136.

10. Vogel A *et al.* 2019 Chapter Three - Lost in trait space: Species-poor communities are inflexible in properties that drive ecosystem functioning. In *Advances in Ecological Research* (eds N Eisenhauer, DA Bohan, AJ Dumbrell), pp. 91–131. Academic Press. (doi:10.1016/bs.aecr.2019.06.002)

11. Kohli BA, Rowe RJ. 2019 Beyond guilds: The promise of continuous traits for mammalian functional diversity. *Journal of Mammalogy* **100**, 285–298. (doi:10.1093/jmammal/gyz054)

12. Chase JM, Myers JA. 2011 Disentangling the importance of ecological niches from stochastic processes across scales. *Philosophical Transactions of the Royal Society B: Biological Sciences* **366**, 2351–2363. (doi:10.1098/rstb.2011.0063)

13. Zhou J, Ning D. 2017 Stochastic community assembly: Does it matter in microbial ecology? *Microbiology and Molecular Biology Reviews* **81**, 10.1128/mmbr.00002-17. (doi:10.1128/mmbr.00002-17)

14. Powell JR, Karunaratne S, Campbell CD, Yao H, Robinson L, Singh BK. 2015 Deterministic processes vary during community assembly for ecologically dissimilar taxa. *Nat Commun* **6**, 8444. (doi:10.1038/ncomms9444)

15. Mammola S, Carmona CP, Guillerme T, Cardoso P. 2021 Concepts and applications in functional diversity. *Functional Ecology* **35**, 1869–1885. (doi:10.1111/1365-2435.13882)

16. Pease AA, González-Díaz AA, Rodiles-Hernández R, Winemiller KO. 2012 Functional diversity and trait–environment relationships of stream fish assemblages in a large tropical catchment. *Freshwater Biology* **57**, 1060–1075. (doi:10.1111/j.1365-2427.2012.02768.x)

17. Ingram T, Shurin JB. 2009 Trait-based assembly and phylogenetic structure in northeast

Pacific rockfish assemblages. *Ecology* **90**, 2444–2453. (doi:10.1890/08-1841.1)

18. Rincón-Díaz MP, Pittman SJ, Arismendi I, Heppell SS. 2018 Functional diversity metrics detect spatio-temporal changes in the fish communities of a Caribbean marine protected area. *Ecosphere* **9**, e02433. (doi:10.1002/ecs2.2433)

19. Pecuchet L, Törnroos A, Lindegren M. 2016 Patterns and drivers of fish community assembly in a large marine ecosystem. *Marine Ecology Progress Series* **546**, 239–248. (doi:10.3354/meps11613)

20. Henseler C, Nordström MC, Törnroos A, Snickars M, Pecuchet L, Lindegren M, Bonsdorff E. 2019 Coastal habitats and their importance for the diversity of benthic communities: A species- and trait-based approach. *Estuarine, Coastal and Shelf Science* **226**, 106272. (doi:10.1016/j.ecss.2019.106272)

21. Sgarlatta MP, Ramírez-Valdez A, Ladah LB, Calderón-Aguilera LE. 2023 Fish functional diversity is modulated by small-scale habitat complexity in a temperate ecosystem. *Hydrobiologia* **850**, 747–759. (doi:10.1007/s10750-022-05061-x)

22. Quirino BA, Lansac-Tôha FM, Thomaz SM, Heino J, Fugi R. 2021 Macrophyte stand complexity explains the functional α and β diversity of fish in a tropical river-floodplain. *Aquat Sci* **83**, 12. (doi:10.1007/s00027-020-00768-2)

23. McLean M, Stuart-Smith RD, Villéger S, Auber A, Edgar GJ, MacNeil MA, Loiseau N, Leprieur F, Mouillot D. 2021 Trait similarity in reef fish faunas across the world's oceans. *Proceedings of the National Academy of Sciences* **118**, e2012318118. (doi:10.1073/pnas.2012318118)

24. Vellend M. 2010 Conceptual synthesis in community ecology. *The Quarterly Review of Biology* **85**, 183–206. (doi:10.1086/652373)

25. Triantis KA *et al.* 2022 Deterministic assembly and anthropogenic extinctions drive convergence of island bird communities. *Global Ecology and Biogeography* **31**, 1741–1755. (doi:10.1111/geb.13556)

26. Reeves JC, Moon BC, Benton MJ, Stubbs TL. 2021 Evolution of ecospace occupancy by Mesozoic marine tetrapods. *Palaeontology* **64**, 31–49. (doi:10.1111/pala.12508)

27. Ceballos G, Ehrlich PR, Barnosky AD, García A, Pringle RM, Palmer TM. 2015 Accelerated modern human–induced species losses: Entering the sixth mass extinction. *Science Advances* **1**, e1400253. (doi:10.1126/sciadv.1400253)

28. Sallan LC, Coates MI. 2010 End-Devonian extinction and a bottleneck in the early evolution of modern jawed vertebrates. *Proceedings of the National Academy of Sciences* **107**, 10131–10135. (doi:10.1073/pnas.0914000107)

29. McGhee GR, Clapham ME, Sheehan PM, Bottjer DJ, Droser ML. 2013 A new ecological-severity ranking of major Phanerozoic biodiversity crises. *Palaeogeography, Palaeoclimatology, Palaeoecology* **370**, 260–270. (doi:10.1016/j.palaeo.2012.12.019)

30. Beerling DJ, Chaloner WG, Woodward FI, Algeo TJ, Scheckler SE. 1998 Terrestrial-marine teleconnections in the Devonian: Links between the evolution of land plants, weathering processes, and marine anoxic events. *Philosophical Transactions of the Royal Society of London. Series B: Biological Sciences* **353**, 113–130. (doi:10.1098/rstb.1998.0195)

31. Brett CE, Walker SE. 2002 Predators and predation in Paleozoic marine environments. *Paleontol. Soc. pap.* **8**, 93–118. (doi:10.1017/S1089332600001078)

32. Copper P. 2011 100 million years of reef prosperity and collapse: Ordovician to Devonian interval. *The Paleontological Society Papers* **17**, 15–32. (doi:10.1017/S1089332600002424)

33. Trinajstic K, Briggs DEG, Long JA. 2021 The Gogo Formation Lagerstätte: A view of Australia's first great barrier reef. *Journal of the Geological Society* **179**, jgs2021-105. (doi:10.1144/jgs2021-105)

34. Cloutier R. 2013 Great Canadian Lagerstätten 4. The Devonian Miguasha biota (Québec): UNESCO world heritage site and a time capsule in the early history of vertebrates. *geocan* **40**, 149–163.

35. Gess RW, Whitfield AK. 2020 Estuarine fish and tetrapod evolution: Insights from a Late Devonian (Famennian) Gondwanan estuarine lake and a southern African Holocene equivalent. *Biological Reviews* **95**, 865–888. (doi:10.1111/brv.12590)

36. Mouchet MA, Villéger S, Mason NWH, Mouillot D. 2010 Functional diversity measures: An overview of their redundancy and their ability to discriminate community assembly rules. *Functional Ecology* **24**, 867–876. (doi:10.1111/j.1365-2435.2010.01695.x)

37. Anderson PSL, Friedman M, Brazeau MD, Rayfield EJ. 2011 Initial radiation of jaws demonstrated stability despite faunal and environmental change. *Nature* **476**, 206–209. (doi:10.1038/nature10207)

38. Cloutier R, Loboziak S, Candilier A-M, Blieck A. 1996 Biostratigraphy of the Upper Devonian Escuminac Formation, eastern Québec, Canada: A comparative study based on miospores and fishes. *Review of Palaeobotany and Palynology* **93**, 191–215. (doi:10.1016/0034-6667(95)00126-3)

39. Cloutier R, Proust J-N, Tessier B. 2011 The Miguasha Fossil-Fish-Lagerstätte: a consequence of the Devonian land–sea interactions. *Palaeobio Palaeoenv* **91**, 293–323. (doi:10.1007/s12549-011-0058-0)

40. Chevrinais M, Jacquet C, Cloutier R. 2017 Early establishment of vertebrate trophic interactions: Food web structure in Middle to Late Devonian fish assemblages with exceptional fossilization. *Bulletin of Geosciences* , 491–510. (doi:10.3140/bull.geosci.1651)

41. Mory AJ, Hocking RM. 2011 *Permian, carboniferous and upper devonian geology of the Northern Canning Basin, Western Australia: A field guide*. Perth: Government of Western Australia.

42. Bascompte J, Melián CJ, Sala E. 2005 Interaction strength combinations and the overfishing of a marine food web. *Proc. Natl. Acad. Sci. U.S.A.* **102**, 5443–5447. (doi:10.1073/pnas.0501562102)

43. Pérez-Matus A *et al.* 2017 Temperate rocky subtidal reef community reveals human impacts across the entire food web. *Marine Ecology Progress Series* **567**, 1–16. (doi:10.3354/meps12057)

44. Lira AS, Lucena-Frédou F, Lacerda CHF, Eduardo LN, Ferreira V, Frédou T, Ménard F, Angelini R, Le Loc'h F. 2022 Effect of fishing effort on the trophic functioning of tropical estuaries in Brazil. *Estuarine, Coastal and Shelf Science* **277**, 108040. (doi:10.1016/j.ecss.2022.108040)

45. Ferreira V, Le Loc'h F, Ménard F, Frédou T, Frédou FL. 2019 Composition of the fish fauna in a tropical estuary: The ecological guild approach. *Sci. Mar.* **83**, 133. (doi:10.3989/scimar.04855.25A)

46. Cohen JE, Schittler DN, Raffaelli DG, Reuman DC. 2009 Food webs are more than the sum of their tritrophic parts. *Proc. Natl. Acad. Sci. U.S.A.* **106**, 22335–22340. (doi:10.1073/pnas.0910582106)

47. Hall SJ, Raffaelli D. 1991 Food-web patterns: Lessons from a species-rich web. *The Journal of Animal Ecology* **60**, 823. (doi:10.2307/5416)

48. Huxham M, Beaney S, Raffaelli D. 1996 Do Parasites Reduce the Chances of Triangulation in a Real Food Web? *Oikos* **76**, 284. (doi:10.2307/3546201)

49. Angelini R, De Morais RJ, Catella AC, Resende EK, Libralato S. 2013 Aquatic food webs of the oxbow lakes in the Pantanal: A new site for fisheries guaranteed by alternated control? *Ecological Modelling* **253**, 82–96. (doi:10.1016/j.ecolmodel.2013.01.001)

50. Resende EKD. 2000 Trophic structure of fish assemblages in the Lower Miranda River, Pantanal, Mato Grosso do Sul State, Brazil. *Rev. Bras. Biol.* **60**, 389–403. (doi:10.1590/S0034-71082000000300004)

51. Pereira RAC, de Resende EK. 1998 Peixes detritívoros da planície inundável do Rio Miranda, Pantanal, Mato Grosso do Sul, Brasil. *EMBRAPA-CPAP. Research Bulletin* **12**, 1–50.

52. de Resende EK, Pereira RAC. 1998 Peixes herbívoros da planície inundável do rio Miranda, Pantanal, Mato Grosso do Sul, Brasil. *EMBRAPA-CPAP. Research Bulletin* **10**, 1–24.

53. de Resende EK, Pereira RAC. 2000 Peixes onívoros da planície inundável do rio Miranda, Pantanal, Mato Grosso do Sul, Brasil. *EMBRAPA-CPAP. Research Bulletin* **16**, 1–60.

54. de Resende EK, Pereira RAC. 2000 Peixes insetívoros e zooplanctófagos da planície inundável do rio Miranda, Mato Grosso do Sul, Brasil. *EMBRAPA-CPAP. Research Bulletin* **17**, 1–42.

55. Ferreira FS, Duarte GS do V, Severo-Neto F, Froehlich O, Súarez YR. 2017 Survey of fish species from plateau streams of the Miranda River Basin in the Upper Paraguay River Region, Brazil. *Biota Neotrop.* **17**, e20170344. (doi:10.1590/1676-0611-BN-2017-0344)

56. Severo-Neto F, Tencatt LFC, Costa-Pereira R, Tavares LER. 2015 Fishes from Baía da Medalha, southern Pantanal, Brazil: A 20 years review. *Biota Neotrop.* **15**. (doi:10.1590/1676-06032015011614)

57. Costa-Pereira R, Severo-Neto F, Yule TS, Pereira APT. 2011 Fruit-eating fishes of Banara arguta (Salicaceae) in the Miranda River floodplain, Pantanal wetland. *Biota Neotrop.* **11**, 373–376. (doi:10.1590/S1676-06032011000400033)

58. Ferreira FS, Duarte GS do V, Severo-Neto F, Froehlich O, Súarez YR. 2017 Survey of fish species from plateau streams of the Miranda River Basin in the Upper Paraguay River Region, Brazil. *Biota Neotrop.* **17**, e20170344. (doi:10.1590/1676-0611-BN-2017-0344)

59. Growns I, Gehrke PC, Astles KL, Pollard DA. 2003 A comparison of fish assemblages associated with different riparian vegetation types in the Hawkesbury–Nepean River system. *Fisheries Management Eco* **10**, 209–220. (doi:10.1046/j.1365-2400.2003.00337.x)

60. Long JA, Trinajstic K. 2010 The Late Devonian Gogo Formation lägerstatte of Western Australia: Exceptional early vertebrate preservation and diversity. *Annu. Rev. Earth Planet. Sci.* **38**, 255–279. (doi:10.1146/annurev-earth-040809-152416)

61. Stekhoven DJ. 2022 missForest: Nonparametric missing value imputation using Random Forest. **R package version 1.5**.

62. Llewelyn J. 2024. Trait-space disparity. *GitHub* and *Zenodo*. https://github.com/JohnLlewelyn/Trait-space-disparity (doi: 10.5281/zenodo.13440287)

63. Boettiger C, Lang DT, Wainwright PC. 2012 rfishbase: Exploring, manipulating and visualizing FishBase data from R. *Journal of Fish Biology* **81**, 2030–2039. (doi:10.1111/j.1095-8649.2012.03464.x)

64. Gower JC, Legendre P. 1986 Metric and Euclidean properties of dissimilarity coefficients. *Journal of Classification* **3**, 5–48. (doi:10.1007/BF01896809)

65. de Bello F, Botta-Dukát Z, Lepš J, Fibich P. 2021 Towards a more balanced combination of multiple traits when computing functional differences between species. *Methods in Ecology and Evolution* **12**, 443–448. (doi:10.1111/2041-210X.13537)

66. Magneville C *et al.* 2022 mFD: An R package to compute and illustrate the multiple facets of functional diversity. *Ecography* **2022**. (doi:10.1111/ecog.05904)

67. Ricklefs RE, Miles DB. 1994 Ecological and evolutionary inferences from morphology: An ecological perspective. In *Ecological morphology: Integrative organismal biology*, pp. 13–41. The University of Chicago Press.

68. Blonder B, Morrow CB, Brown S, Butruille G, Chen D, Laini A, Harris DJ. 2023 hypervolume: High dimensional geometry, set operations, projection, and inference   using kernel density estimation, support vector machines, and convex hulls. *R package version 3.1.3* , <https://CRAN.R-project.org/package=hypervolume>.

69. Aldridge RJ, Purnell MA. 1996 The conodont controversies. *Trends in Ecology & Evolution* **11**, 463–468. (doi:10.1016/0169-5347(96)10048-3)

70. Aldridge RJ, Donoghue PCJ. 1998 Conodonts: A sister group to hagfishes? In *The Biology of Hagfishes* (eds JM Jørgensen, JP Lomholt, RE Weber, H Malte), pp. 15–31. Dordrecht: Springer Netherlands. (doi:10.1007/978-94-011-5834-3_2)

71. McCoy VE, Lamsdell JC, Poschmann M, Anderson RP, Briggs DEG. 2015 All the better to see you with: Eyes and claws reveal the evolution of divergent ecological roles in giant pterygotid eurypterids. *Biology Letters* **11**, 20150564. (doi:10.1098/rsbl.2015.0564)

72. Pimiento C, Leprieur F, Silvestro D, Lefcheck JS, Albouy C, Rasher DB, Davis M, Svenning J-C, Griffin JN. 2020 Functional diversity of marine megafauna in the Anthropocene. *Sci. Adv.* **6**, eaay7650. (doi:10.1126/sciadv.aay7650)

73. Le Bagousse-Pinguet Y *et al.* 2021 Functional rarity and evenness are key facets of biodiversity to boost multifunctionality. *Proceedings of the National Academy of Sciences* **118**, e2019355118. (doi:10.1073/pnas.2019355118)

74. Schleuning M *et al.* 2020 Trait-based assessments of climate-change impacts on interacting species. *Trends in Ecology & Evolution* , S0169534719303556. (doi:10.1016/j.tree.2019.12.010)

75. Brind'Amour A, Boisclair D, Dray S, Legendre P. 2011 Relationships between species feeding traits and environmental conditions in fish communities: A three-matrix approach. *Ecological Applications* **21**, 363–377. (doi:10.1890/09-2178.1)

76. Moyle PB, Cech JJ. 2004 *Fishes: An introduction to ichthyology*. 5th edition. New Jersey, USA: Pearson Prentice Hall.

77. Jobbins M, Rücklin M, Sánchez Villagra MR, Lelièvre H, Grogan E, Szrek P, Klug C. 2024 Extreme lower jaw elongation in a placoderm reflects high disparity and modularity in early vertebrate evolution. *Royal Society Open Science* **11**, 231747. (doi:10.1098/rsos.231747)

78. Janvier P. 1996 *Early vertebrates*. Oxford, U.K.: Clarendon Press.

79. Ehlman SM, Trimmer PC, Sih A. 2019 Prey responses to exotic predators: Effects of old risks and new cues. *The American Naturalist* **193**, 575–587. (doi:10.1086/702252)

80. Schakmann M, Korsmeyer KE. 2023 Fish swimming mode and body morphology affect the energetics of swimming in a wave-surge water flow. *Journal of Experimental Biology* **226**, jeb244739. (doi:10.1242/jeb.244739)

81. Kelley JL, Fitzpatrick JL, Merilaita S. 2013 Spots and stripes: Ecology and colour pattern evolution in butterflyfishes. *Proceedings of the Royal Society B: Biological Sciences* **280**, 20122730. (doi:10.1098/rspb.2012.2730)

82. Wilson MVH, Caldwell MW. 1993 New Silurian and Devonian fork-tailed 'thelodonts' are jawless vertebrates with stomachs and deep bodies. *Nature* **361**, 442–444. (doi:10.1038/361442a0)

83. Girard C, Renaud S. 2012 Disparity changes in 370 Ma Devonian fossils: The signature of ecological dynamics? *PLOS ONE* **7**, e36230. (doi:10.1371/journal.pone.0036230)

84. Friedman M, Sallan LC. 2012 Five hundred million years of extinction and recovery: A phanerozoic survey of large-scale diversity patterns in fishes. *Palaeontology* **55**, 707–742. (doi:10.1111/j.1475-4983.2012.01165.x)

85. Henderson S, Dunne EM, Fasey SA, Giles S. 2023 The early diversification of ray-finned fishes (Actinopterygii): Hypotheses, challenges and future prospects. *Biological Reviews* **98**, 284–315. (doi:10.1111/brv.12907)

86. Richardson LE, Graham NAJ, Pratchett MS, Hoey AS. 2017 Structural complexity mediates functional structure of reef fish assemblages among coral habitats. *Environ Biol Fish* **100**, 193–207. (doi:10.1007/s10641-016-0571-0)

87. Majchrzyk A, Jakubowicz M, Berkowski B, Król JJ, Zatoń M, Zapalski MK. 2024 Modern-type reef in ancient time - Palaeoecology of a Middle Devonian coral community from Madène el Mrakib (Anti-Atlas, Morocco). *Palaeogeography, Palaeoclimatology, Palaeoecology* **633**, 111876. (doi:10.1016/j.palaeo.2023.111876)

88. Benton MJ, Emerson BC. 2007 How did life become so diverse? The dynamics of diversification according to the fossil record and molecular phylogenetics. *Palaeontology* **50**, 23–40. (doi:10.1111/j.1475-4983.2006.00612.x)

89. Hill JJ, Puttick MN, Stubbs TL, Rayfield EJ, Donoghue PCJ. 2018 Evolution of jaw disparity in fishes. *Palaeontology* **61**, 847–854. (doi:10.1111/pala.12371)

90. He S *et al.* 2023 Genetic innovations underpin morphological diversity and radiation in teleost fish. (doi:10.21203/rs.3.rs-3148441/v1)

91. Witten PE, Hall BK. 2015 Teleost skeletal plasticity: Modulation, adaptation, and remodelling. *Copeia* **103**, 727–739. (doi:10.1643/CG-14-140)

92. Long JA, Choo B, Clement A. 2018 The evolution of fishes through geological time. In *Evolution and development of fishes* (eds Z Johanson, C Underwood, M Richter), pp. 3–29. Cambridge University Press. (doi:10.1017/9781316832172.002)

93. Peres-Neto PR. 2004 Patterns in the co-occurrence of fish species in streams: The role of site suitability, morphology and phylogeny versus species interactions. *Oecologia* **140**, 352–360. (doi:10.1007/s00442-004-1578-3)

94. Grossman GD, Ratajczak Jr. RE, Crawford M, Freeman MC. 1998 Assemblage organization in stream fishes: Effects of environmental variation and interspecific interactions. *Ecological Monographs* **68**, 395–420. (doi:10.1890/0012-9615(1998)068[0395:AOISFE]2.0.CO;2)

95. Gratwicke B, Speight MR. 2005 The relationship between fish species richness, abundance and habitat complexity in a range of shallow tropical marine habitats. *Journal of Fish Biology* **66**, 650–667. (doi:10.1111/j.0022-1112.2005.00629.x)

96. Passos CVB, Fabré NN, Malhado ACM, Batista VS, Ladle RJ. 2016 Estuarization increases functional diversity of demersal fish assemblages in tropical coastal ecosystems. *Journal of Fish Biology* **89**, 847–862. (doi:10.1111/jfb.13029)

97. Potter IC, Tweedley JR, Elliott M, Whitfield AK. 2015 The ways in which fish use estuaries: A refinement and expansion of the guild approach. *Fish and Fisheries* **16**, 230–239. (doi:10.1111/faf.12050)

98. Lamothe KA, Alofs KM, Jackson DA, Somers KM. 2018 Functional diversity and redundancy of freshwater fish communities across biogeographic and environmental gradients. *Diversity and Distributions* **24**, 1612–1626. (doi:10.1111/ddi.12812)

99. Leitão RP, Zuanon J, Villéger S, Williams SE, Baraloto C, Fortunel C, Mendonça FP, Mouillot D. 2016 Rare species contribute disproportionately to the functional structure of species assemblages. *Proceedings of the Royal Society B: Biological Sciences* **283**, 20160084. (doi:10.1098/rspb.2016.0084)

100. Biggs CR *et al.* 2020 Does functional redundancy affect ecological stability and resilience? A review and meta-analysis. *Ecosphere* **11**, e03184. (doi:10.1002/ecs2.3184)

101. Landi P, Minoarivelo HO, Brännström Å, Hui C, Dieckmann U. 2018 Complexity and stability of ecological networks: A review of the theory. *Population Ecology* **60**, 319–345. (doi:10.1007/s10144-018-0628-3)

102.   Strona G. 2022 *Hidden Pathways to Extinction*. Switzerland: Springer.
(doi:10.1007/978-3-030-86764-5)
103.   Cardoso P, Pekár S, Jocqué R, Coddington JA. 2011 Global Patterns of Guild
Composition and Functional Diversity of Spiders. *PLOS ONE* **6**, e21710.
(doi:10.1371/journal.pone.0021710)
104.   Mouillot D *et al.* 2014 Functional over-redundancy and high functional vulnerability
in global fish faunas on tropical reefs | PNAS. *PNAS* **111**, 13757–13762.
(doi:10.1073/pnas.1317625111)
105.   Pigot AL, Trisos CH, Tobias JA. 2016 Functional traits reveal the expansion and
packing of ecological niche space underlying an elevational diversity gradient in
passerine birds. *Proceedings of the Royal Society B: Biological Sciences* **283**, 20152013.
(doi:10.1098/rspb.2015.2013)
106.   Myers EMV, Anderson MJ, Liggins L, Harvey ES, Roberts CD, Eme D. 2021 High
functional diversity in deep-sea fish communities and increasing intraspecific trait
variation with increasing latitude. *Ecology and Evolution* **11**, 10600–10612.
(doi:10.1002/ece3.7871)
107.   Stuart-Smith RD *et al.* 2013 Integrating abundance and functional traits reveals new
global hotspots of fish diversity. *Nature* **501**, 539–542. (doi:10.1038/nature12529)
108.   Berke SK, Jablonski D, Krug AZ, Valentine JW. 2014 Origination and immigration
drive latitudinal gradients in marine functional diversity. *PLOS ONE* **9**, e101494.
(doi:10.1371/journal.pone.0101494)
109.   Schumm M, Edie SM, Collins KS, Gómez-Bahamón V, Supriya K, White AE, Price
TD, Jablonski D. 2019 Common latitudinal gradients in functional richness and
functional evenness across marine and terrestrial systems. *Proceedings of the Royal
Society B: Biological Sciences* **286**, 20190745. (doi:10.1098/rspb.2019.0745)
110.   Jarzyna MA, Quintero I, Jetz W. 2021 Global functional and phylogenetic structure of
avian assemblages across elevation and latitude. *Ecology Letters* **24**, 196–207.
(doi:10.1111/ele.13631)
111.   Lamanna C *et al.* 2014 Functional trait space and the latitudinal diversity gradient.
Proc. Natl. Acad. Sci. U. S. A. **111**, 13745–13750. (doi:10.1073/pnas.1317722111)
112.   Hillebrand H. 2004 On the generality of the latitudinal diversity gradient. *The
American Naturalist* **163**, 192–211. (doi:10.1086/381004)
113.   Benton MJ. 2001 Biodiversity on land and in the sea. *Geological Journal* **36**, 211–
230. (doi:10.1002/gj.877)
114.   McKitrick MC. 1993 Phylogenetic constraint in evolutionary theory: Has it any
explanatory power? *Annu. Rev. Ecol. Evol. Syst.* , 307–330.
115.   Champagnat N, Ferrière R, Méléard S. 2006 Unifying evolutionary dynamics: From
individual stochastic processes to macroscopic models. *Theoretical Population Biology*
69, 297–321. (doi:10.1016/j.tpb.2005.10.004)
116.   Mani GS, Clarke BC. 1997 Mutational order: A major stochastic process in evolution.
Proceedings of the Royal Society of London. B. Biological Sciences **240**, 29–37.
(doi:10.1098/rspb.1990.0025)
117.   Fritz SA, Schnitzler J, Eronen JT, Hof C, Böhning-Gaese K, Graham CH. 2013
Diversity in time and space: Wanted dead and alive. *Trends in Ecology & Evolution* **28**,
509–516. (doi:10.1016/j.tree.2013.05.004)