# An affordable solution for investigating zebra finch intracranial electroencephalography (iEEG) signals.

Mohammad-Mahdi Abolghasemi

Institute for Cognitive and Brain Sciences, Shahid Beheshti University, Tehran, Iran.

E-Mail: mma.abolghasemi@gmail.com

Shahriar Rezghi Shirsavar

School of Cognitive Sciences, Institute for Research in Fundamental Sciences (IPM), Tehran 1956836484, Iran.

E-Mail: shahriar.rezghi@ut.ac.ir

Amirreza Bahramani

School of Cognitive Sciences, Institute for Research in Fundamental Sciences (IPM), Tehran 1956836484, Iran.

Department of Electrical Engineering, Sharif University of Technology, Tehran 1458889694, Iran.

E-Mail: bahramani@ipm.ir

Milad Yekani[*]

School of Cognitive Sciences, Institute for Research in Fundamental Sciences (IPM), Tehran 1956836484, Iran.

E-Mail: miladyekani@ipm.ir

Author contributions:

Conceptualization M.Y.; Methodology M.Y., M.M.A., and A.B.; Software S.R.S. and M.M.A.; Validation M.Y.; Formal Analysis M.M.A, .A.B. and S.R.S.; Resources M.Y., M.M.A., and S.R.S.; Data Curation M.M.A., S.R.S., and A.B.; Writing – Original Draft M.Y., M.M.A and A.B.; Writing – Review & Editing M.Y., M.M.A. and A.B.; Visualization M.M.A. and M.Y.; Supervision M.Y.; Project Administration M.Y.

Abstract:

The zebra finch is a well-studied animal model for investigating the neural mechanisms of vocal learning, and electrophysiology is the primary technique for understanding their song system. Most of the studies on zebra finches have focused on intracerebral recordings. However, these methods are only affordable for limited laboratories. Recently, different open-source hardware for acquiring electroencephalography (EEG) signals has been developed. It's unclear whether these solutions suit zebra finch studies as they have not been evaluated. Electrocorticography signals can provide a preliminary guide for more in-depth inquiries and also aid in understanding the global behavior of the bird's brain, as opposed to the more common localized approach. We present a detailed protocol for acquiring intracranial electroencephalography (iEEG) data from zebra finches using the OpenBCI Cyton board, an open-source device. We implemented stainless steel electrodes on the brain's surface and recorded the brain signals from two recording sites above two auditory-responsive nuclei. To validate our method, we ran two different experiments. In the first experiment, we recorded neural activity under various concentrations of isoflurane and extracted the suppression duration to measure anesthesia depth. In the second experiment, we head-fixed the birds and, under light anesthesia, presented them with various auditory stimuli to evaluate event-related potentials (ERP) and generate spectrograms. The results showed a significant increase in suppression duration with deeper anesthesia, and the ERP and spectrogram responses to auditory stimuli differed accordingly. These findings indicate that using our methodology, one can successfully collect iEEG signals from zebra finches. These findings pave the way for future studies to use iEEG to investigate bird cognition in a more affordable way.

Keywords: Zebra finch, animal iEEG, open source, Songbird

## Introduction:

The investigation of the avian neural system provides a promising avenue for illuminating the underlying neural mechanisms of vocal learning and even speech acquisition in humans (1). As an exceptional species that can acquire singing via vocal imitation from parents, humans exhibit noteworthy similarities with songbirds regarding complex sound production and learning. Thus, songbirds have emerged as effective models for neuroscientific inquiries concerning vocal learning (2). A range of research methods is used in the extensive investigations of the bird's vocal system, including electrophysiology (3), histology (4), genetics (5,6), and behavioral studies (7). The bulk of current knowledge concerning the avian vocal system has been derived from experimental studies conducted on the zebra finch (*Taeniopygia guttata*), a widely accepted avian laboratory model species (8). Invasive neural recordings in zebra finches have revealed valuable insights into the song system and mechanisms of vocal learning. The ultimate goal in this line of research is to understand the neural mechanism of creating complicated motor sequences and vocal production and translate these findings to the human brain.

Human brain studies are restricted by a limited capacity for intervention due to ethical and technical considerations. These limitations force human research to use noninvasive methods like electroencephalography (EEG). Nevertheless, testing these methods on animal models provides valuable insights, establishing a robust foundation for verifying their applicability. These signals are clinically relevant to consciousness alterations, seizure activity, anesthesia, and sleep states. As an alternative approach, since animals enable invasive practices, studies utilizing the intracranial version of EEG (iEEG) have provided more valuable insight into the nature of acquired signals than EEG due to a higher signal-to-noise ratio (9). Animal iEEG studies also have an extensive history that provides complementary data to human research. Specifically, animal iEEG models exploring pharmacological agents' impact on the brain represent a crucial translational inquiry area (10).

IEEG recording has been conducted to study sleep and migratory behaviors in birds (11). Research on birds' sleep patterns, which are essential from an evolutionary perspective, has also been conducted using

iEEG. Furthermore, brain activity related to memory consolidation during the critical period of song

learning in juvenile finches can be detected by this method (12). The research literature on working with

birds is full of extracellular recording studies, in which we can examine the behavior of a neuron during

the production or hearing of a song with a very high accuracy. Despite the importance of animal iEEG,

avian iEEG has been studied very limited compared to other animals.

IEEG studies in birds have great translational potential. In this concept, reproducing bird songs from

iEEG signals can be a suitable model for the brain-computer interface of speech. The advantage of this

method is its ease of implementation, which allows it to be performed before other neuronal recording

experiments and during behavioral studies. Consequently, the higher cost and energy of conducting neural

recording experiments, which make them less feasible, will be evaded.

With the increasing demand for neural data, scientists seek more accessible ways to democratize scientific

research (13). To this end, many entities produce tools, programs, and open-source data sets. Given the

importance of EEG studies, open-source hardware and software solutions are being introduced in this

field. OpenBCI offers a versatile open-source hardware platform for various biosensing applications,

including EEG, EMG, and ECG signals (14). At its core, the OpenBCI Cyton board integrates several key

components: a PIC32MX250F128B microcontroller, a ChipKIT UDB32-MX2-DIP bootloader, an

ADS1299 analog-to-digital converter with eight channels (expandable to 16) and a maximum sampling

rate of 16 kHz, and a 3-axis LIS3DH accelerometer. Users can configure EEG channels in either

monopolar or bipolar modes, while the board also accommodates five external digital inputs and three

analog inputs. Moreover, the device supports data streaming over Wi-Fi via Transmission Control

Protocol (TCP) (15).

We conducted two experiments to test the efficacy of our method. In the first experiment, we recorded

signals under different concentrations of isoflurane. The suppression duration was used as a proxy for

anesthesia depth.  In the second experiment, following the recovery and adaptation period, we recorded

iEEG from two electrodes above two auditory-responsive nuclei while the birds were under light

anesthesia. We presented various auditory stimuli, including the bird's own song (BOS) and its reversed

version. Analysis of the recorded data revealed robust neural responses to all stimuli. Notably, the neural

response patterns to BOS were distinct from those elicited by the other stimuli, supporting the validity of

our method. Our study highlights the usefulness of OpenBCI in recording the iEEG activity of a bird's

brain, particularly the zebra finch, for investigating auditory responses.

## 110   Methods:

111     A detailed explanation of the procedure has been provided on protocol.io.

## 112   Animal subjects:

113     We used five male zebra finches (> 60 days old) for experiment one and two male zebra finches for

114     experiment two. All animals were obtained from the IPM birds garden aviary and kept in a local animal

115     house after surgery. All procedures were approved by the Research Ethics Committee of the School of

116     Cognitive Sciences (SCS) of the Institute for Research in Fundamental Sciences (IPM, protocol number

117     1402/40/1/2841).

## 118   Electrodes and head-post implantation:

Animals were kept under restricted food and water conditions before surgery. Anesthesia was induced

using 2% isoflurane, and the animal was secured into the stereotaxic device after induction. We used our

designed 3D-printed anesthesia mask compatible with the stereotaxic device to sustain gas delivery to the

head-fixed bird. The head feathers were removed, and the incision site was prepared for surgery. After

exposing the skull and identifying the bifurcation landmark, we determined the coordinates for the

craniotomy sites relative to this bifurcation. Our goal was to position electrodes on two auditory-

responsive nuclei: HVC (proper name) and the caudomedial mesopallium (CMM). The HVC site was

located 0.2 mm anterior and 2.3 mm lateral, while the CMM site was positioned 2 mm anterior and 0.4

mm lateral from the bifurcation. Craniotomy sites were drilled, and after placing the electrodes, the skull

was prepared for cement adhesion. After the cement was applied to the area, the headpost was installed.

The cement was allowed to be cured completely before proceeding. Detailed instructions can be found in

the protocol section.

## Experimental Procedures

### Experiment 1: Suppression Duration Recording under Anesthesia

We collected iEEG data from two recording electrodes implanted beneath the skull of the zebra finches

during the different depths of anesthesia with isoflurane. After the surgery and electrode implantation, the

OpenBCI recording was set up, and the signals were recorded with a 250 Hz sampling rate for about 2

minutes in each of 2.5%, 2%, 1.5%, 1%, 0.6%, and 0.4% of isoflurane (Figure 1A). Before analysis, the

data were preprocessed to ensure data quality and reliability. Standard preprocessing steps were

employed, including a high-pass filter at 1 Hz, a low-pass filter at 80 Hz, and a notch filter at 50 Hz to

remove noise, and the signal was re-referenced during recording by placing electrodes on the cerebellum.

After that, the data was extracted based on the different levels of isoflurane for further analysis in

suppression detections.

### Experiment 2: Auditory Response Recording under Light Anesthesia

To record neural activity from HVC and CMM regions, five auditory stimuli were each presented 60

times with silent intervals randomly ranging from 2 to 4 seconds between them. The stimuli included

BOS, reversed BOS, conspecific song, a 5 kHz tone, and white noise (Figure 1B). A highly synchronized

trigger signal was essential during recording to accurately annotate each trial. Audio playback and trigger

output to the OpenBCI device were managed using a Raspberry Pi 3. An optocoupler circuit was

employed to isolate the OpenBCI device from the Raspberry Pi, preventing noise induction from the

trigger signal. The task was programmed in C++ and optimized to achieve a latency of 1 ms, ensuring the

required accuracy.

Analysis

Experiment 1: Suppression duration

Bursts are characterized by short episodes of partial reactivity or temporary spikes in neural firing,

followed by a significant decrease or suppression of neural activity. We measured suppression duration in

zebra finches under varying levels of isoflurane anesthesia. Suppression duration is a crucial metric for

evaluating brain activity and detecting instances of burst suppression, which may indicate fluctuations in

brain function or anesthesia-induced alterations in neural activity. Our objective was to explore the neural

dynamics and anesthesia effects on zebra finches while administering different levels of isoflurane. The

previous method for defining burst suppression relied on setting a fixed threshold for EEG voltage

(typically between 0.5 and 20 µV) and classifying suppressions as segments where the voltage remained

below this threshold for at least 0.5 seconds. This approach was problematic due to variations in

amplitude and spectral characteristics across different pathological conditions and anesthetic effects. In

2013, Westover et al. introduced a new method involving an empirical approach in which two

experienced electroencephalographers manually segmented EEG data based on visual pattern recognition

and clinical judgment. Additionally, the study developed an automatic segmentation method using

recursive variance estimation, which classifies bursts and suppressions based on the adaptive mean and

variance of the EEG signal, optimized to match expert consensus. As a result, a real-time algorithm was

created to automatically differentiate suppressions from non-suppressions (bursts) in EEGs. Using this

algorithm, we rely on the total suppression duration for each task condition and compare these durations

statistically. At each level of isoflurane (low, medium, and high), we compute the suppression duration for

each 2-minute segment of the session and perform statistical analyses to compare the conditions (16).

To perform further statistical analyses, we sorted the dataset based on the isoflurane concentration into

three distinct categories: low, medium, and high levels of anesthesia. Specifically, isoflurane below one

percent was classified as low-level anesthesia, those above two indicated a high level, and values falling

between 1 and 2 were classified as medium levels of anesthesia. To compare suppression durations across

the three experimental conditions, a Friedman test was conducted as a nonparametric alternative for

repeated measures. When significant differences were found, post hoc pairwise comparisons were

performed using Wilcoxon signed-rank tests with Bonferroni correction to adjust for multiple

comparisons.

Experiment 2: Spectrogram and Event-Related Potential (ERP)

The data collected from OpenBCI was saved as .txt files. After importing the data, we applied a bandpass

filter between 1 and 40 Hz to focus on the frequency range of interest and reduce noise. We then

segmented the continuous data into individual trials based on trigger times corresponding to stimulus

presentations. For the event-related potential (ERP) analysis, we performed baseline correction by

subtracting the mean activity during the 500 ms preceding the stimulus presentation from each trial. This

normalization step ensures that the post-stimulus activity is relative to the pre-stimulus baseline. We then

averaged the baseline-corrected trials for each stimulus type to obtain the ERP waveforms. For the time-

frequency analysis, we computed spectrograms using the Stockwell Transform (17,18), which provides a

detailed time-frequency representation of the signal. To normalize each trial's spectrogram, we calculated

the mean power over time and used it to adjust the trial's response, effectively controlling for inter-trial

variability in power. After normalization, we averaged the spectrograms by stimulus type to produce the

final time-frequency plots. All analyses were conducted using Python with the MNE 1.8 toolbox. To

ensure the reliability of our results, we also replicated the analyses using custom scripts in MATLAB

R2023b.

Results

Anesthetized recording

Our analysis revealed distinct changes in suppression duration across different stages of isoflurane

anesthesia. As isoflurane was administered, we observed a gradual increase in the suppression duration,

indicating an increase in burst suppression events (Figure 2 C) and reduced brain activity, as evidenced by

computed fast Fourier transforms in each of low, medium, and high anesthesia for both anterior and

posterior electrodes (Figure 2 A). Additionally, we statistically compared the suppression durations across

nine blocks of 120-second recordings at low, medium, and high anesthesia levels. The analysis of

suppression durations across the three experimental conditions (low, medium, and high isoflurane) was

conducted using a Friedman test. Significant differences were identified among the conditions for both

electrode sites. The p-value for the Friedman test in the anterior electrode was 0.00841, while for the

posterior electrode, it was 0.0015. Post hoc pairwise comparisons were performed using Wilcoxon signed-

rank tests with Bonferroni correction to account for multiple comparisons. For the anterior electrode, the

adjusted p-values for the pairwise comparisons were as follows: high vs. medium: 0.75, low vs. medium:

0.16, and low vs. high: 0.02  (Figure 2 B-top). For the posterior electrode, the adjusted p-values for the

pairwise comparisons were 0.93, 0.023, and 0.023, respectively  (Figure 2B-bottom ).

Auditory Response Recording

All five auditory stimuli elicited neural responses, evident in the ERP recordings from both electrodes

(Figure 3A, B). Notably, a sharp response occurred approximately 200 ms after the stimulus

presentation. This delay is expected, as the neural activity is primarily generated by the underlying

nuclei—namely, HVC and the CMM—which are both responsive to higher-level auditory stimuli.

Interestingly, the ERP response for the BOS differs significantly from the other stimuli; it exhibits a later

rise with a smaller slope. This may be due to the distinct responses of these nuclei to the BOS. These

differences in response were also distinctly observed in the HVC and CMM. A larger amplitude in the

CMM compared to the HVC may result from the dominant role of the CMM in processing auditory

stimuli, as is evident across all conditions (Figure 3A, B).

Figure 4A, B shows the spectrogram of the neural responses. There is a rapid, short response in the Theta

and Alpha frequency bands for all stimuli. These early responses can be attributed to neural activity

generated by the primary auditory areas. Moreover, the neural response for BOS is again distinct,

displaying more pronounced and sustained activity in the Delta band. An interesting phenomenon is that

the increase in Delta band power is followed by an immediate decrease, which can indicate the underlying

neural dynamics.

Differences are also evident in the HVC and CMM. Specifically, the CMM shows a greater response to

conspecific and Reverse BOS stimuli than BOS alone. However, the HVC response to the BOS is more

prominent than the reverse BOS. Interestingly, the response to the BOS in HVC is also greater than this

response in the CMM. Overall, both areas respond more strongly to song-like stimuli than to white noise

or the 5 kHz tone, as seen in the spectrogram. For better illustration, the spectrograms of the presented

audio stimuli are shown in Figures 4C and 3C.

Discussion:

Songbirds are the primary animal models for studying vocal perception, production, and learning (19). In the past decades, zebra finch has been utilized as a significant model in these studies, and spike recording was the primary method in these experiments. However, the techniques used to track the activity of a single neuron are expensive (20) and complicated and provide a narrow field of view to capture the brain's overall activity (21). Recently, there has been a growing interest in studying the cognitive processes in birds as they show fascinating abilities in doing high-level mental functions (22). These functions are processed in a distributed manner in the brain, and to study them, we need to have a larger field of view. Our experiment tries to address the problems mentioned above. In summary, we attempt to solve two issues. First, to reduce the cost of the avian studies by using open-source solutions, and second, to capture the brain's dynamic state during mental activity by recording iEEG. This paper presents a detailed protocol for recording iEEG from an anesthetized zebra finch. Previous experiments on birds' EEG under isoflurane anesthesia showed similar signal patterns to those of mammals (23). This pattern is entirely detectable by visually inspecting a bird's EEG signal under surgical anesthesia, characterized by brief periods of burst activity and silent suppressed intervals between them (24). We used suppression duration as an indicator of the anesthesia's depth in our experiment to test the efficiency of our method. The results show that this index is significantly higher in the high concentrations of gas than the low concentrations, indicating that our method successfully captures the brain's dynamic. Although we chose this experiment to show the validity of our method as Anesthesia produces a characteristic feature of brain dynamics that has been previously observed, there are some scientifically valuable facts in showing this phenomenon. The emergence of burst suppression in this species under anesthesia shows an evolutionarily

conserved arousal mechanism. These findings can open future studies to understand a shared

mechanism behind producing a conscious state in the brain. Comparative Studies related to

conscious states in other animals can create insight into the serious questions about the

evolutionary aspect of consciousness (25), and studies on birds' cognition regarding their high

cognitive ability are in this category. Our ultimate aim in developing this method is to use it in

studies related to the cognitive process in birds, particularly those connected to vocal perception

and production. We conducted a second experiment with male zebra finches under light

anesthesia, and they were presented with five different stimuli. Our results demonstrated an ERP

response to all five stimuli, which were composed of two parts: an early activity in higher

frequency bands, which probably originated from field L. This was followed by an increase in

the power of the Delta band, which was later and specifically different for BOS. We think that

this second phase is mostly generated by higher areas, especially HVC and CMM, in which the

electrodes were implanted above them. Moreover, these responses in the Delta band were

followed by a decrease in power, which can be the subject of further analysis of circuit-level

mechanisms. Altogether, these observations can validate our method and showcase its use for

investigating the neural correlates of vocal perception in an affordable way.

Although we observed notable differences between the responses to BOS and reverse BOS in the

HVC and CMM electrodes, we should consider that localization can be improved by using

higher-impedance electrodes and limiting the contact area. These modifications could also allow

us to implant electrodes directly within the nuclei, enabling us to capture LFP signals with

greater spatial resolution.

Our method offers a way to record chronic data from animals, as they can carry implanted

materials for their lives. This recording approach is helpful when we want to capture the changes

in brain function during aging and maturation (e.g., changes in vocal perception during the

transition from the juvenile stage to adulthood) (26). Given that the main noninvasive current

method in BCI uses EEG signals to control effectors in humans, we think the zebra finch iEEG

can be an appropriate model for the vocal prosthesis. Previous studies that have utilized the

biomechanical model of the birds' vocal organs and fed them by neural signals gained

tremendous success (27). Future studies can improve this method fed those models and create a

strong basic model for vocal prosthesis controlled by EEG signals.  Finally, the wireless

communication protocol employed by OpenBCI presents an exciting opportunity for developing

small-scale devices capable of chronic recording in freely behaving animals, especially in flying

birds. Thanks to its open-source nature, research groups can modify these devices to improve

their reliability for animal studies. While OpenBCI has proven useful in human studies, its

application in animal studies has been limited until now. This study paves the way for future

research endeavors to leverage this technology to collect a broader range of signals, especially

local field potential.  Birds are increasingly used in neuroscience and cognitive studies; however,

the EEG signal has not been used extensively in this line of research. Our findings can trigger

future studies to use iEEG in bird cognitive studies.

Acknowledgements

We express our deepest gratitude to Dr. Mahdi Khademian for his pivotal role in founding the Birds Lab
and for his continued support. We also extend our thanks to the IPM School of Cognitive Sciences for
their support in the establishment of the Birds Lab.

## References:

1.  Zhang Y, Zhou L, Zuo J, Wang S, Meng W. Analogies of human speech and bird song: From vocal learning behavior to its neural basis. Front Psychol. 2023 Feb 22;14:1100969.

2.  Doupe AJ, Kuhl PK. BIRDSONG AND HUMAN SPEECH: Common Themes and Mechanisms. Annu Rev Neurosci. 1999 Mar;22(1):567–631.

3.  Lewicki MS. Intracellular Characterization of Song-Specific Neurons in the Zebra Finch Auditory Forebrain. J Neurosci. 1996 Sep 15;16(18):5854–63.

4.  Farries MA, Perkel DJ. A Telencephalic Nucleus Essential for Song Learning Contains Neurons with Physiological Characteristics of Both Striatum and Globus Pallidus. J Neurosci. 2002 May 1;22(9):3776–87.

5.  Gordon RL, Ravignani A, Hyland Bruno J, Robinson CM, Scartozzi A, Embalabala R, et al. Linking the genomic signatures of human beat synchronization and learned song in birds. Phil Trans R Soc B. 2021 Oct 11;376(1835):20200329.

6.  Xiao L, Merullo DP, Koch TMI, Cao M, Co M, Kulkarni A, et al. Expression of FoxP2 in the basal ganglia regulates vocal motor sequences in the adult songbird. Nat Commun. 2021 May 11;12(1):2617.

7.  Brazas ML, Shimizu T. Significance of visual cues in choice behavior in the female zebra finch (Taeniopygia guttata castanotis). Anim Cogn. 2002 Jun;5(2):91–5.

8.  Mello CV. The Zebra Finch, *Taeniopygia guttata*□: An Avian Model for Investigating the Neurobiological Basis of Vocal Learning. Cold Spring Harb Protoc. 2014 Dec;2014(12):pdb.emo084574.

9.  Parvizi J, Kastner S. Promises and limitations of human intracranial electroencephalography. Nat Neurosci. 2018 Apr;21(4):474–83.

10. Drinkenburg WHIM, Ruigt GSF, Ahnaou A. Pharmaco-EEG Studies in Animals: An Overview of Contemporary Translational Applications. Neuropsychobiology. 2015;72(3–4):151–64.

11. Rattenborg NC, Voirin B, Cruz SM, Tisdale R, Dell'Omo G, Lipp HP, et al. Evidence that birds sleep in mid-flight. Nat Commun. 2016 Aug 3;7(1):12468.

12. Shank SS, Margoliash D. Sleep and sensorimotor integration during early vocal learning in a songbird. Nature. 2009 Mar;458(7234):73–7.

13. White SR, Amarante LM, Kravitz AV, Laubach M. The Future Is Open: Open-Source Tools for Behavioral Neuroscience Research. eNeuro. 2019 Jul;6(4):ENEURO.0223-19.2019.

14. Laport F, Vazquez-Araujo FJ, Iglesia D, Castro PM, Dapena A. A Comparative Study of Low Cost Open Source EEG Devices. Proceedings. 2019;21(1):40.

15. Cardona-Álvarez YN, Álvarez-Meza AM, Cárdenas-Peña DA, Castaño-Duque GA, Castellanos-Dominguez G. A Novel OpenBCI Framework for EEG-Based Neurophysiological Experiments. Sensors. 2023 Apr 6;23(7):3763.

16. Brandon Westover M, Shafi MM, Ching S, Chemali JJ, Purdon PL, Cash SS, et al. Real-time segmentation of burst suppression patterns in critical care EEG monitoring. Journal of Neuroscience Methods. 2013 Sep;219(1):131–41.

17. Moukadem A, Bouguila Z, Abdeslam DO, Dieterlen A. Stockwell transform optimization applied on the detection of split in heart sounds. In: 2014 22nd European Signal Processing Conference (EUSIPCO) [Internet]. 2014 [cited 2024 Oct 29]. p. 2015–9. Available from: https://ieeexplore.ieee.org/document/6952743

18. Rodino L, Schulze BW, Wong M, editors. Pseudo-Differential Operators: Partial Differential Equations and Time-Frequency Analysis [Internet]. Providence, Rhode Island: American Mathematical Society; 2007 [cited 2024 Oct 29]. Available from: http://www.ams.org/fic/052

19. Sakata JT, Woolley SC, Fay RR, Popper AN, editors. The Neuroethology of Birdsong [Internet]. Cham: Springer International Publishing; 2020 [cited 2023 Jan 23]. (Springer Handbook of Auditory Research; vol. 71). Available from: http://link.springer.com/10.1007/978-3-030-34683-6

20. Cheng N, Murari K. OSERR: an open-source standalone electrophysiology recording system for rodents. Sci Rep. 2020 Oct 12;10(1):16996.

21. Sejnowski TJ, Churchland PS, Movshon JA. Putting big data to good use in neuroscience. Nat Neurosci. 2014 Nov;17(11):1440–1.

22. Emery NJ. Cognitive ornithology: the evolution of avian intelligence. Phil Trans R Soc B. 2006 Jan 29;361(1465):23–43.

23. McIlhone AE, Beausoleil NJ, Kells NJ, Johnson CB, Mellor DJ. Effects of halothane on the electroencephalogram of the chicken. Veterinary Medicine & Sci. 2018 May;4(2):98–105.

24. Swank RL, Watson CW. EFFECTS OF BARBITURATES AND ETHER ON SPONTANEOUS ELECTRICAL ACTIVITY OF DOG BRAIN. Journal of Neurophysiology. 1949 Mar 1;12(2):137–60.

25. Irwin LN, Chittka L, Jablonka E, Mallatt J. Editorial: Comparative animal consciousness. Front Syst Neurosci. 2022 Oct 19;16:998421.

26. Funabiki Y, Konishi M. Long Memory in Song Learning by Zebra Finches. J Neurosci. 2003 Jul 30;23(17):6928–35.

27. Arneodo EM, Chen S, Brown DE, Gilja V, Gentner TQ. Neurally driven synthesis of learned, complex vocalizations. Current Biology. 2021 Aug;31(15):3419-3425.e5.

370 Figures:

## Experiment 1

## Experiment 2

**Figure 1: Graphical abstract of two experiments.** Implanted parts on the bird's skull have been identified in different colors. The cream area on the skull shows the cemented region on top of the head. As demonstrated, all other compartments are fixed by cement on the skull. The curly wires represent recording and ground electrodes, and the perpendicularly attached bar (gray) indicates the head post. The electrodes are connected to the Cyton recording board (for a detailed explanation of connections, see the protocol.io). (A) In the first experiment, varying concentrations of isoflurane were used to achieve different depths of anesthesia. (B) In the second experiment, the finch was head-fixed under light anesthesia, and intracranial EEG (iEEG) data were recorded from implanted

 electrodes                     during                     an                     auditory                     task.

**Figure 2: FFT analysis of isoflurane levels, suppression duration comparison across conditions, and sample oscillations at different isoflurane levels.** (A) FFT analysis under low, medium, and high levels of isoflurane anesthesia, showing a decrease in power across various frequency ranges as isoflurane levels increase, with observations at both posterior (Right) and frontal (Left) sites. (B) Statistical pairwise comparisons of three different isoflurane levels, with Bonferroni-adjusted p-values for anterior (top) and posterior (bottom) sites. (C) A 20-second

387    frame of raw signal oscillations under low (top), medium (middle), and high (bottom) isoflurane anesthesia,

388    illustrating a higher amplitude in the frequency range for low-level isoflurane.

389
390    **Figure 3**: **Event-Related Potentials (ERPs) for all stimuli gathered, separated, and spectrogram of presented**

391    **stimuli.** (A) Displays the ERPs of all presented auditory stimuli for HVC (Right) and CMM (Left). (B) Shows the

392    separated ERPs of stimuli in HVC (Right) and CMM (Left). (C) Presents the spectrogram of the audio stimuli

393    during the task.

**Figure 4: Spectrograms for each stimulus condition in HVC, CMM, and audio stimuli.** (A) Spectrogram of HVC responses to the five stimuli: BOS, Reverse BOS, Conspecific, White Noise, and 5 kHz Tone. A distinct, selective response to BOS is observed in the HVC site. (B) Spectrogram of auditory stimuli responses in the CMM area, showing a greater response to Reverse BOS and Conspecific compared to the HVC area. (C) Spectrogram of the presented audio stimuli during the task.

401