Response diversity is a major driver of temporal stability in complex food webs

Alain Danet (1), Sonia Kéfi (2,3), Thomas F. Johnson (1), Andrew P. Beckerman (1)

- 1 School of Biosciences, Ecology and Evolutionary Biology, University of Sheffield, Western Bank, Sheffield, S10 2TN, UK
- 2 ISEM, CNRS, Univ. Montpellier, IRD, EPHE, Montpellier, France
- 3 Santa Fe Institute, 1399 Hyde Park Road, Santa Fe, NM 87501, USA

Abstract

Global change constitutes a suite of major threats to biodiversity and ecosystem functioning. These threats can materialise via changes in the temporal stability of ecological communities and the services they provide. However, the majority of research on stability has focused on single trophic level communities and has not yet integrated classic theory about species richness and food web structure with more recent theory centred on response diversity and stochasticity. Using a stochastic bioenergenetic food-web model, we reveal that response diversity is a major driver of community stability. Moreover, positive stability-richness relationships emerge only in the presence of response diversity. In contrast to previous work, food-web structural properties are only secondary drivers of overall community stability, but interact with response diversity to determine the sign of the stability-richness relationship. Our study highlights the complex pathways by which food-web structure and response diversity drive community stability, and raises concerns about how the loss of response diversity may lead to a breakdown of stability and the capacity for these communities to deliver functions and services to human societies.

Introduction

Global change, characterised by multiple, simultaneous threats to biodiversity[1,2], is having impacts on the temporal stability of complex ecological communities[3–7]. A breakdown of stability impacts the capacity for these communities to deliver functions and services to human societies[8]. Predicting and managing the stability and services provided by biodiverse communities requires reconciling multiple concepts and theories about temporal stability in species diverse and structurally complex communities. These include classic ideas about the role of trophic structure and interaction strengths[9–15], newer concepts of response diversity, average population stability, asynchrony and portfolio effects and integrative theories about the relationship between stability, structure and diversity.

An abundant literature has established that species diversity can buffer the effect of environmental perturbations on ecological communities, thereby increasing the temporal stability (hereafter "stability") of communities[16–21]. Because different species have diverse niches and environmental preferences, they may react differently to the same environmental change[22,23]. This diversity of species responses to environmental change, coined as "response diversity", has been identified as a crucial mechanism that buffers the response of ecological communities to environmental perturbations[16,20,24]. For example, in a given community, some species will respond positively and others negatively to changes in temperature, depending on their thermal niche[25]. Furthermore, it has also been shown recently that the stability of communities can be decomposed into the average population stability and the asynchrony of population fluctuations[21,24]. Average population stability quantifies the average variability in species abundances, while asynchrony refers to how much the temporal fluctuations of the different species of the community are buffering each other. Recent studies have shown that asynchrony can be further resolved into "portfolio effects", the statistical averaging of independent species fluctuations in response to environmental perturbations, and "compensatory dynamics", where one species' decrease can be compensated for by the increase in another species[17,21,24].

These mechanisms have been well-established in data from single trophic level (e.g. competitive) communities, such as grasslands, butterflies, birds, and bats communities[3,19]. In these studies, response diversity increases asynchrony, and is a more important driver of community stability than population stability[3]. In these

same studies, the portfolio effect was found to play a key role in the emergence of positive diversity-stability relationships in experimental grasslands[21].

Despite these data, there is limited evidence and understanding for how these collections of theories integrate to define stability in food webs with substantial diversity and vertical complexity (i.e. trophic levels)[26–29]. Classic theory and concepts about stability-complexity[9,10], stability-structure[11–13,28] and stability-richness[13,14,30] offer some expectation. They suggest that population stability is expected to decrease with increasing interaction strength[9,11,12] and food web connectance[13] but to increase with average trophic level[27,31]. As a result, the distribution and strength of species interactions, as well as the average trophic level, can have strong effects on community stability as well as on stability-richness relationships[13,14]. These relationships indicate that the structure of food-webs strongly influences population stability and stability-richness relationships.

The more recent theory and concepts centred on response diversity and asynchrony add further predictions[21,24]. With vertical structure and trophic interactions, consumers can synchronise the dynamics of their prey[32,33], can offset the effects of response diversity on asynchrony and can generate substantial compensatory dynamics via prey switching and trophic cascades[11,34,35].

Despite this collection of historical and more modern theory, we don't know what determines stability in vertically structured, species rich communities. Against this gap in understanding, we specifically investigate the effects of response diversity and food-web structure on community stability and on stability-richness relationships in a variable environment. The key question is whether the new concepts of environmental stochasticity, response diversity and asynchrony are the main mechanisms driving community stability versus more classic concepts tied to food-web structure and diversity.

We explore this key question by extending the bioenergetic food web model[36,37], to incorporate environmental stochasticity and response diversity[38–40]. We introduced environmental stochasticity by making species death rates stochastic[38]. Response diversity is introduced as the correlation of stochasticity among species, so that species that experience environmental changes in the same way show positively correlated variations in their death rates.

We generated over 45000 in-silico communities with diverse richness, trophic levels and connectance and

experimentally simulated variable degrees of interaction strength, response diversity and environmental

stochasticity. For each community, we evaluated the effects on stability of species richness, response diver-

sity, community structure and stochasticity. Stability was itself partitioned in our analyses into population

stability and asynchrony (i.e. portfolio and compensatory effects). Finally, we also evaluated the occurrence

of negative and positive relationships between community stability and species richness.

Results

Applying a structural equation model on the outcomes of food web simulations (see Methods) revealed

that the temporal stability of community biomass is positively influenced by both population stability and

asynchrony, and that population stability has twice the effect of asynchrony (resp. $r_\delta = 0.9$ and 0.57, r_δ being

the standardised coefficient, Fig. 1a).

We further found that asynchrony is driven more by portfolio effects than by compensatory dynamics (resp.

$r_\delta = 0.92$ and 0.36). In more details, response diversity has strong positive effects on asynchrony, mainly

through portfolio effects ($r_\delta = 0.85$), at the expense of compensatory dynamics ($r_\delta = -0.12$). All other studied

properties were found to have relatively much weaker effects on asynchrony. Food web structural properties

– namely average trophic level, connectance and average interaction strength – have consistent negative

effects on asynchrony, through both portfolio and compensatory effects (Fig. 1a). Indeed, as average trophic

level and connectance increase compensatory effects are reduced (resp. $r_\delta = -0.17$ and -0.16, Fig. 1a), while

average interaction strength has a small negative effect on portfolio effects ($r_\delta = -0.05$). Species richness has

a positive effect on both portfolio and compensatory effects (resp. $r_\delta = 0.18$ and 0.14). Finally, as expected,

environmental stochasticity has little effect on asynchrony.

Focusing on population stability, environmental stochasticity was found to have a strong negative effect on

it ($r_\delta = -0.91$). All other studied properties were found to have little effect on population stability, except for

the average trophic level which has a small positive effect ($r_\delta = 0.17$, Fig. 1a, but see Fig. S1) and response

diversity a small negative effect ($r_\delta = -0.07$).

Figure 1: Structural Equation Model linking food-web structure and stressors to the temporal stability of community biomass. a, Continuous blue arrows display positive and dashed red arrows display negative standardised effects, width of the arrows being proportional to the absolute values of the standardised effects. Only the standardised coefficients whose absolute values are superior to 0.05 are displayed. The full table of coefficients including control variables is displayed in Table S2. b, Total effects of food-web structure on community stability, asynchrony, and population stability derived from the structural equation model. N = 46880 simulations.

Deriving total effects from the sum of direct and indirect effects in the structural equation model (See Method), we showed that both environmental stochasticity and species response diversity have by far the strongest total effects on community stability (resp. r_δ= -0.8 and 0.36, Fig. 1b). Species richness and average trophic level have similar small but positive total effects on community stability (r_δ= 0.08), but through opposite pathways. While species richness increases community stability through a strong positive effect on asynchrony (r_δ= 0.21), it also weakly decreases population stability (r_δ= -0.04). In contrast, average trophic level increases community stability through strong positive effect on population stability (r_δ= 0.17), but is dampened by a negative total effect on asynchrony (r_δ= -0.1). Interestingly, average interaction strength also has a positive effect on population stability and a negative one on asynchrony (resp. r_δ= 0.02 and -0.06). This results in a total negative effect on community stability (r_δ= -0.02). Connectance, instead, has a negative effect on community stability (r_δ= -0.06) through negative effects on both asynchrony and population stability (resp. r_δ= -0.07 and -0.02).

We further assessed how environmental stochasticity and response diversity interacted with food-web structural properties to determine the sign of the stability-richness relationship, using a linear model including main effects and all interactions among food-web structural properties, response diversity and environmental stochasticity (see Methods). We found that there are strong interactions between response diversity and food-web structure, which jointly determine the sign of the stability-richness relationship.

In the absence of response diversity, we found only negative stability-richness relationships, regardless of the food-web structure (Fig. 2a). Response diversity enables the rise of positive stability-richness relationships, enhanced by average interaction strength and dampened by average trophic level and connectance. Higher interaction strength leads to positive stability-richness relationships for lower values of response diversity, and interaction strength enhances asynchrony-richness relationships (Fig. S2), so that the stronger positive stability-richness relationships are found at both high interaction strength and high response diversity levels. In contrast, higher average trophic level and connectance both lead to negative stability-richness relationships, because they drive more negative population stability-richness relationships and weaker positive asynchrony-richness relationships (Fig. S2). Overall, our results highlight a strong contrast between the relatively small effects of food-web structural properties on overall community stability (Fig. 1b) but their strong effects on

Figure 2: Slope of the community stability - species richness relationship based on food-web structure and response diversity. a, Diagrams displaying the slope coefficients of the effect of species richness on community stability. The average values of the control variables were used to generate the predictions: connectance = 0.14, average interaction strength = 0.025, average trophic level = 1.49, predator interference = 0.5, Predator-Prey Mass Ratio = 33, and environmental stochasticity = 0.3. b, c: Examples of predictions from the model for two combinations of average trophic level and response diversity (values displayed in panel a). Lines display the mean predictions of the model. The coefficients of the linear model are displayed in Fig. S3. Relationships between population stability, asynchrony and species richness are displayed in Fig. S2.

the slope of the stability-richness relationship (Fig. 2a).

Discussion

With the projected changes in climate and land use, and the expected resulting changes in community composition, it is important to understand the mechanisms which allow ecological communities to remain stable despite perturbations. The two components of temporal community stability are the average stability of the populations composing the community on the one hand, and the asynchrony of the population fluctuations on the other hand[21,24]. In simplified assemblages, such as single trophic level communities, community stability has been shown to be driven by asynchrony in species fluctuations, rather than by population stability[3]. However, whether this holds for more complex ecological communities, such as food webs, remains largely unknown. Here, we tackled this question using simulations of complex food-web dynamics with a stochastic bioenergetic model. Our model results show that, in complex food-webs, community stability is more driven by population stability than by asynchrony, in contrast with what was observed in single trophic level communities but in agreement with a previous food-web data study[27]. Our results thereby contribute to reconcile models and data analyses. The study further reveals that the main drivers of community stability are environmental stochasticity and response diversity, respectively acting on population stability and asynchrony. Surprisingly, food-web structural properties have in turn much weaker effects, a result consistent across methodological choices (Fig. S1). Structural properties were however found to play an important role in the sign of the stability-richness relationship, which is strongly mediated by the interaction between food-web structure and response diversity. These results highlight the importance of response diversity to improve our understanding of the stability of ecological communities in variable environments.

We found environmental stochasticity and response diversity to be the main overall drivers of community stability. While environmental stochasticity decreases population stability, response diversity generates asynchrony, thereby buffering environmental stochasticity. This result is in accordance with previous theoretical and empirical findings in competitive communities[13,19,20,24,30,41]. In turn, response diversity was found to operate on asynchrony mainly through portfolio effects in agreement with empirical findings in competi-

tive assemblages[20,21]. This suggests that response diversity generates asynchrony mainly by the statistical averaging of random independent species fluctuations[17].

A consequence of the large effect of response diversity is that asynchrony in species fluctuations is driven more by the statistical averaging of independent species fluctuations (portfolio effects) than by compensatory dynamics emerging from species interactions. This finding is also coherent with previous studies in competitive communities[20,21], but it was less expected to happen in food-webs. Trophic interactions are indeed expected to result in more compensatory dynamics resulting from oscillations between predators and prey[42] or prey switching from predators[11,34]. However, pioneering theoretical studies have shown that environmental stochasticity can dampen prey switching[38] and that trophic cascades between trophic levels in species rich food-webs are weak compared to food-chains and species poor food-web modules[35]. So, the question of the importance of compensatory dynamics in complex food webs remains an open and interesting avenue for future research.

In contrast to empirical evidence in competitive communities[3], population stability (not asynchrony) was found to be the dominant driver of community stability in our food web simulations. These results are in agreement with previous findings in empirical food-webs[27] and food-web models[28], which suggests fundamental differences between competitive communities and food webs in the relative contributions of population stability and asynchrony to community stability. Our findings suggest that the presence of trophic links contributes to synchronise the species of the community because we found that the food-web metrics studied – namely average trophic level, connectance, and average interaction strength – all decrease asynchrony. These results are in line with empirical evidence that predators can synchronise their prey[32,33], and that species pairs involved in trophic interactions are more synchronous than those that are not[27]. A complementary explanation for the relatively lower importance of asynchrony in food webs (compared to single trophic level communities) could be related to the biomass structure of food-webs. Food-webs typically include a larger range of body masses than single trophic level assemblages[43]. Such highly uneven biomass distribution can translate in uneven species temporal variability, thereby creating uneven species contributions to asynchrony[24] and limiting portfolio effects and compensatory dynamics in species-rich communities[21].

Investigating in more detail the drivers of community stability, our results suggest that species richness and the network structural properties investigated only have a weak effect on overall community stability. We found that higher average trophic level leads to higher community stability by increasing average population stability, confirming previous theoretical[28,31,36] and empirical[27] results. However, this effect disappeared when we simulated community dynamics with equal mortality rates for all species instead of allometric ones (Fig. S1). This suggests that the higher stability of food-webs with higher average trophic levels stems more from their lower mortality rates rather than from their trophic role directly. The overall effect of connectance on overall community stability was found to be quite low, in agreement with previous empirical and theoretical findings[27,28]. Similarly, we found very low effects of the average interaction strength on community stability, which is in apparent disagreement with the predominant role of interaction strength and connectance on stability[9].

Our results also allow us to revisit the old question of the relationship between the complexity of ecological communities and their stability[9,10]. We found that response diversity enables the rise of positive stability-richness relationships, enhanced by average interaction strength and dampened by average trophic level and connectance. In the absence of response diversity, only negative relationships between community stability and species richness are observed. This reinforces the idea that increases in species richness increase community stability only if it adds species that respond differently to environmental perturbations, i.e. if there is response diversity[13,19]. Indeed, a community with numerous species which respond differently to environmental change is more likely to buffer a wide array of perturbations[16,30]. Despite its relatively weaker effect on community stability (than response diversity and environmental stochasticity), food-web structure interacts strongly with response diversity to determine the sign of stability-richness relationships. Surprisingly, we find that higher average interaction strength enhances positive stability-richness relationships by enhancing asynchrony-richness relationships. Conversely, higher average trophic level and higher connectance both led to negative stability-richness relationships by dampening asynchrony-richness relationships. Our results resonate with findings that some community structure can lead to negative stability-richness relationships, as previously reported in freshwater food-webs[27]. Response diversity might therefore be one of the mechanisms that explains why positive relationships between stability and species richness are so prevalent in empirical

settings[22,24], while food-web theoretical models that do not often include environmental stochasticity and response diversity typically find negative complexity-stability relationships[39].

In conclusion, our study highlights the complex but consistent effects of response diversity and food-web structure on the temporal stability of species rich communities. At the core of the insurance hypothesis[16,44], our results suggest that environmental stochasticity and response diversity are the main mechanisms driving community stability, while food-web structural properties are only mediating them. This aligns with the idea that, in species-rich communities, community structure might be secondary in understanding the generic effects of biodiversity on ecosystem functioning[30,45]. Our results further add to mounting evidence that population stability is a stronger driver of community stability than asynchrony in food-webs, in contrast with what was observed in competitive communities. The fact that environmental stochasticity was found to be the main driver of overall community stability raises concerns about the consequences of the current rise in the frequency and severity of environmental extreme events for ecological communities[1,46]; indeed our results suggest that these changes are unlikely to be compensated by biotic mechanisms alone. Documenting how rapid biodiversity changes[47,48] driven by human pressures[49] – and in particular the reported homogenisation[50,51] in community composition – affect response diversity will be key to accurately predict how it will affect, in turn, the stability of communities.

Methods

Bioenergenetic model

We simulated the dynamics of complex food-webs using the allometric bioenergetic model[36,37,52]. This model is widely used to simulate the dynamics of complex ecological communities because of the simplicity of its parametrisation, using the metabolic theory of ecology[53]. The model describes species biomass dynamics over time with two master equations, one for the primary producers (eq. (2)) and one for the consumers (eq.(1)):

$$\frac{dB_i}{dt} = \sum_{j \in \{\text{res}\}_i} x_i y_i B_k F_{ij} - \sum_{j \in \{\text{cons}\}_i} \frac{x_j y_j B_j F_{ji}}{e_{ij}} - x_i B_i - d_i B_i \tag{1}$$

$$\frac{dB_i}{dt} = r_i B_i G_i - \sum_{j \in \{\text{cons}\}_i} \frac{x_j y_j B_j F_{ji}}{e_{ij}} - d_i B_i \tag{2}$$

The consumers gain biomass by consuming resources (i.e. their preys) at a rate that depends on their mass specific metabolic rate (x_i), maximum consumption rate (y_i) and on the functional response (F_{ij}) describing how the rate of consumption of a consumer i on a resource j vary with the biomass of this resource. Consumers then lose biomass by being consumed, e_{ij} being the assimilation efficiency of the consumer j on the resource i. Consumers also lose biomass over time through metabolic losses ($-x_i B_i$) and natural mortality ($-d_i B_i$). The primary producers gain biomass over time with a growth rate (r_i, eq. (2)) and a functional response (G_i) describing how the growth of the producers vary with their biomass. They lose biomass by being consumed by consumers and by natural mortality.

$$G_i = (1 - \frac{\sum_{j \in \{\text{prod}\}} \alpha_{ij} B_j}{K_i}) \tag{3}$$

$$F_{ij} = \frac{\omega_{ij} B_j^h}{B_0^h + c_i B_i + \sum_{k \in \{\text{res}\}_i} \omega_{ik} B_k^h} \tag{4}$$

The growth of the primary producers (eq. (3)) is logistic where the growth rate is maximal and null when $\sum_{j \in \{\text{prod}\}} \alpha_{ij} B_j$ are respectively close to 0 and 1. α_{ij} is the per capita effect of the producer j on the producer i, K_i being the carrying capacity for the producer j. The functional response of a consumer feeding on a resource (eq. (4)) depends on the relative preference of the consumer on the resource ($\omega_i j$) is limited by a half-saturation rate (B_0), intraspecific interference coefficient (c_i), and on the availability of its other resources ($\sum_{k \in \{\text{res}\}_i} \omega_{ik} B_k^h$). Finally, the h exponent controls the shape of the functional response from a saturating function ($h = 1$, Holling type II) to a sigmoid ($h = 2$, Holling type III).

Environmental stochasticity and response diversity

We added a stochastic natural mortality rate to the species dynamics, representing a fluctuating environment. Based on the literature and previous models[38,54], we assumed that natural mortality rates scales inversely with the species body mass ($d_i = d_0 M_i^{-1/4}$), as all other physiological parameters. We used a basal natural mortality rate of $d_0 = 0.4$, as a previous study[38].

We introduced environmental stochasticity by considering stochastic variation in mortality rates. Doing so allows to consider stochasticity without modifying species interactions themselves[38]. The stochastic part of the mortality rates ($\epsilon_{i,t}$) follow a normal distribution 0 centered ($\epsilon_{i,t} \sim N(0, \sigma_e^2)$) with a variance ($\sigma_e^2$). The mortality rates of each species at the time t equals to $d_{i,t} = d_i e^{\epsilon_{i,t}}$, which ensured that the mortality rates never become negative[38]. We simulated ϵ using an Ornstein-Uhlenbeck process, a modified Brownian motion where the stochastic values tend to come back to the central value (i.e. $\epsilon = 0$) such as ϵ_t varies according to this stochastic differential equation: $d\epsilon_t = (0 - \epsilon_t)d_t + \sigma_e dW_t$, where dW_t is a Brownian motion. The strength of environmental stochasticity was then controlled by σ_e.

Response diversity was controlled by the correlation among species stochastic mortality rates ($\rho_{ij,i\neq j}$) such as response diversity is maximal when the stochastic component of species mortality rates ($\epsilon_{i,t}$) are uncorrelated ($\rho = 0$), and response diversity is null when the stochastic component of species mortality rates are perfectly correlated[38,40,55] ($\rho = 1$). So the species mortality rates were a product of the stochastic mortality rates generated by the Ornstein-Uhlenbeck process and a variance-covariance matrix such as:

$$
\epsilon_{i,t} = \begin{bmatrix} \rho_{11}\sigma_e^2 & \rho_{12}\sigma_e^2 & \cdots & \rho_{1n}\sigma_e^2 \\ \rho_{21}\sigma_e^2 & \rho_{22}\sigma_e^2 & \cdots & \rho_{2n}\sigma_e^2 \\ \vdots & \vdots & \ddots & \vdots \\ \rho_{n1}\sigma_e^2 & \rho_{n2}\sigma_e^2 & \cdots & \rho_{nn}\sigma_e^2 \end{bmatrix} \begin{bmatrix} a_{1,t} \\ a_{2,t} \\ \vdots \\ a_{n,t} \end{bmatrix} \tag{5}
$$

Where $a_{i,t}$ the stochastic value generated by the Ornstein-Uhlenbeck process at time t, $\rho_{ii} = 1.0$ while all $\rho_{ij,i\neq j}$ were equal and control the level of response diversity.

Simulation

We generated food-web structure using the niche model[56] with an initial species richness from 10 to 60, and connectance from 0.02 to 0.38 (Table S1). We generated food-webs without cannibalistic links and discarded food-webs that contained disconnected species.

We varied the strength of environmental stochasticity by varying σ_e from 0.1 to 0.6, the range of mortality rates observed in protists[38]. We decreased response diversity by increasing ρ from 0 to 1 (i.e. response diversity being $1 - \rho$) (Fig. S4).

To vary interaction strength and mortality rates (i.e. allometrically scaled mortality rate d) in the community, we varied the Predator-Prey Mass Ratio (PPMR) from 1 to 100 (Fig. S5b). We set $h = 2$, i.e. a functional response of type III (eq. (4)), and varied predator interference (i.e. from $c = 0$ to $c = 1$, eq. (4)), as predator interference was shown to have a tremendous role on population stability[36]. We set a global carrying capacity of $K' = 10$ to ensure that consumers were not limited by the biomass input from primary producers. We standardised the carrying capacity by the number of primary producers and by the interspecific competition among producers to ensure that the effects of species richness on temporal stability are not driven by the increase in the number of primary producers, thereby trivially increase biomass input for consumers. We then standardised K' such as $K_i = \frac{1+\alpha_{ij}(S-1)}{S}K'^{30}$. However, we found that our results were not affected when removing the standardisation of carrying capacity (Fig. S1).

We numerically solved the stochastic differential equation systems during at least 2000 timesteps and collected the last 500 timesteps, similarly to previous studies[36]. We numerically solve the system with $\Delta t = 0.1$ or $\Delta t = 0.05$ when instability was detected during the simulations due to stiff changes in biomass. We set extinction threshold to 10^{-6} and if an extinction happened in the last 500 timesteps, we continued to run the simulation for another 1000 timesteps until there was no extinction events during the last 500 timesteps. When we found disconnected species at the end of the simulation (i.e. a primary producer without consumer or a consumer without prey), we set their biomass to 0 and re-run the simulation for another 1000 timesteps. Except in the case of disconnected species, we re-ran simulations with species starting biomass equal to their biomass at the last timestep of the previous simulation. Supplementary analysis showed that alternative

choices of simulation processes, such as longer simulations, rebuilding food-webs (i.e. resetting consumer

preferences, recomputing species bodymass according to the new food-web) after removing disconnected

species did not affect our results (Fig. S1). The food-web generation and simulations were done in Julia by

developing an extension of the package `EcologicalNetworkDynamics.jl`[57].

In sum, we generated 46880 simulations generated a wide range of food-web structure (Median (5%,

95%), species richness: 13.0 (5.0,26.0), connectance: 0.12 (0.058,0.3), average interaction strength: 0.02

(0.0031,0.071), maximum trophic level: 2.5 (2.0,3.5)) (Table S3).

Simulation analysis

We measured food-web properties that have been linked with stability in the literature. We measured total

biomass, species richness, connectance, average trophic level weighted by biomass, and average interaction

strength. Food-web properties were measured at each of the 500 timesteps and then averaged, except

connectance and trophic level that were static. Connectance was computed as $C = L/S(S-1)$, L being

the number of trophic links and S being the number of species in the community. Species trophic level

was computed recursively from the bottom of the food-web, such as the trophic level of a consumer was

equal to the average trophic level of its resources added to 1, the trophic level of primary producers being

set to 1[58]. Average trophic level was then equal to: $1/\hat{B}_i \sum_i \mathrm{TL}_i$, TL_i being the trophic level and $\hat{B}_i$ the

average biomass of the species i. Interaction strength was computed was quantified as the biomass fluxes

going from a resource to a consumer, such as: $I = x_i y_i B_k F_{ij}$ (eq. (1)), and averaged over time. Our way of

measuring interaction strength is very similar to previous metrics used in previous studies, at the difference

here that we average fluxes over time and not the maximal interaction strength[11]. Finally, the average

interaction strength of a community was computed as the average interaction strength at the exclusion of

null interaction strengths (i.e. excluding absence of trophic interactions).

We assessed the effects of food-web structure, environmental stochasticity and response diversity on temporal

stability of biomass. We measured temporal stability of biomass as the inverse of the coefficient of variation.

We further partitioned stability into population stability (S_{pop}) and asynchrony (ϕ) such as:

$$S_{com} = S_{pop} \times \phi \tag{6}$$

with $S_{com} = \mu_{tot}/\sigma_{tot}$, $S_{pop} = \mu_{tot}/\sum_i \sigma_i$, $\phi = \sum_i \sigma_i/\sigma_{tot}$; μ_{tot} and σ_{tot} being respectively average total biomass and standard deviation of total biomass.

To further get a mechanistic insight about the effects of stressors and food-web structure on stability, we partitioned asynchrony into portfolio effects and compensatory dynamics stemming from species interactions[21] (CPE_{int}). The measurement of portfolio effects has been a topic of debate[21,44]. Doak's definition[17] defines portfolio effects as the product of statistical averaging effects (SAE) and compensatory effects arising from response diversity[17,21] (CPE_{env}). It then quantities the portfolio effect ($PFE = SAE \times CPE_{env}$) emerging from independent species fluctuations, i.e. the statistical averaging effect (SAE), weighted by the effects of response diversity of species to environmental fluctuations, such as portfolio effects are dampened if species have correlated responses to environmental fluctuations.

$$S_{com} = S_{pop} \times SAE \times CPE \tag{7}$$

Statistical averaging effect is the share of asynchrony assuming that species are independent, meaning that the variance total of the community is equal to the sum of species variances only (i.e. species covariances are null). We can define community stability ($S_{com,IP}$) in that scenario and then derive SAE.

$$S_{com,IP} = \frac{\mu_{tot}}{\sqrt{\sum_i \sigma_i^2}} = SAE \times S_{pop}$$
$$SAE = \frac{S_{com,IP}}{S_{pop}} = \frac{\sum_i \sigma_i}{\sqrt{\sum_i \sigma_i^2}} \tag{8}$$

Compensatory dynamics in the food-webs is then the remaining part of asynchrony:

$$CPE = \frac{S_{com}}{SAE \times S_{pop}} = \frac{\sqrt{\sum_i \sigma_i^2}}{\sigma_{tot}} \tag{9}$$

Compensatory dynamics can arise from predator-prey interactions (CPE_{int}) or from differential response of species to environmental stochasticity (CPE_{env}). From the deterministic and stochastic parts of mortality rates, we computed the compensatory dynamics linked to the environment such as:

$$CPE_{env} = \mathbf{B} \times \text{diag}(d_i) \times e^{\mathbf{E}} \tag{10}$$

With $\mathbf{B}$ and $\mathbf{E}$ being the matrices time x species for species biomass and stochastic mortality rates respectively. Then compensatory dynamics raising from species interactions were computed as: $CPE_{int} = CPE/CPE_{env}$. Finally, portfolio effects was computed as: $PFE = SAE \times CPE_{env}$ and compensatory dynamics raising from species interactions as CPE_{int}. The final stability decomposition reads as follows:

$$S_{com} = S_{pop} \times PFE \times CPE_{int} \tag{11}$$

Statistical analysis

Using a structural equation model, we disentangled the complex effects of food-web structure, environmental stochasticity and response diversity on the different facets of temporal stability of biomass (i.e. population stability, asynchrony, portfolio effects and compensatory dynamic effects). We tested the effects of food-web structure (average interaction strength, average trophic level, connectance) on population stability and asynchrony partitions (portfolio and compensatory effects). Grounded in theoretical ecology, we expected that higher average trophic level, lower interaction strength, lower connectance result in higher population stability and lower asynchrony[11,12,34,35]. We further added simulation parameters as control variables because higher Predator-Prey Mass Ratio and predator interference were also expected to increase population stability[36]. Finally, we tested the effects of environmental stochasticity and response diversity on population stability and asynchrony partitions, although there were predicted to mainly affect to population stability (i.e. by increasing amplitude of population fluctuations) and asynchrony (i.e. by generating independent population fluctuations) respectively. We completed the structural equation model with the mathematical relationship linking compensatory dynamics due to species interactions (CPE_{int}) and statistical averag-

ing effects ($SAE \times CPE_{env}$) to asynchrony, and from asynchrony and population stability to community stability.

Prior to test the structural equation model, we logged all the stability components to transform their relation from multiplicative to additive (eq. (7)). We ensured that all the linear models composing the structural equation model presented low multicollinearity (Variance Inflation Factor < 3, Table S4), although interaction strength and species richness were highly correlated (Fig. S5). The structural equation model using the R package `PiecewiseSEM`[59]. We computed the sum of the direct and indirect effects using `semEff` R package[60]. In the main text, we reported standardised coefficients which were obtained by scaling the coefficients by the standard deviation of the response and predictor variable. Finally, we reported only the direct standardised coefficients with an absolute value equal or above 0.05, because almost all effects were statistically significant given the high numbers of simulations.

In a second analysis, we tested how food-web structure, response diversity, and environmental stochasticity modulate stability-species richness relationships. We modelled temporal stability of community biomass according to food-web structure: species richness, weighted average trophic level, connectance and average interaction strength, as well as environmental stochasticity and response diversity. We further included the two-way interactions between species richness and food-web structure, between species richness and response diversity, between species richness and environmental stochasticity. Finally, we added the three-way interactions between species richness, food-web structure and response diversity, between species richness, food-web structure and environmental stochasticity. We added predator interference, predator-prey mass ratio as control variables. We further added the ID of the food-web as a random intercept. We then used the linear equation of the model and associated slope coefficients to predict the shape of the relationship between community stability and species richness according to a gradient of response diversity and of food-web structure. To do so, we summed all the coefficients involving species richness in the model, i.e. the one, two, and three-way terms. The values of the other variables were set to the values indicated in the Fig. 2. The coefficients of the models are displayed in Fig. S3. The VIF were inferior to 3, indicating low multicollinearity (Table S5), we checked the distribution of the residuals (Fig. S6) using `DHARMa` R package. The model was implemented using the R package `glmmTMB`[61].

Code availability

The code used to run the simulations, analyse the data and write the manuscript is available on Github (https://github.com/alaindanet/xStressorsStabBEFW) and archived on Zenodo (https://zenodo.org/recor ds/13454946).

References

1. Calvin, K. *et al.* *IPCC, 2023: Climate Change 2023: Synthesis Report. Contribution of Working Groups I, II and III to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change [Core Writing Team, H. Lee and J. Romero (eds.)]. IPCC, Geneva, Switzerland.* https://www.ipcc.ch/report/ar6/syr/ (2023) doi:10.59327/IPCC/AR6-9789291691647.

2. Simmons, B. I. *et al.* Refocusing multiple stressor research around the targets and scales of ecological impacts. *Nature Ecology & Evolution* **5**, 1478–1489 (2021).

3. Olivier, T., Thébault, E., Elias, M., Fontaine, B. & Fontaine, C. Urbanization and agricultural intensification destabilize animal communities differently than diversity loss. *Nature Communications* **11**, 2686 (2020).

4. De Boeck, H. J. *et al.* Patterns and drivers of biodiversity–stability relationships under climate extremes. *Journal of Ecology* **106**, 890–902 (2018).

5. Hansen, B. B. *et al.* Climate Events Synchronize the Dynamics of a Resident Vertebrate Community in the High Arctic. *Science* **339**, 313–315 (2013).

6. Blüthgen, N. *et al.* Land use imperils plant and animal community stability through changes in asynchrony rather than diversity. *Nature Communications* **7**, 10697 (2016).

7. Harris, R. M. B. *et al.* Biological responses to the press and pulse of climate trends and extreme events. *Nature Climate Change* **8**, 579 (2018).

8. Loreau, M., Hector, A. & Isbell, F. *The Ecological and Societal Consequences of Biodiversity Loss.* (Wiley, 2022). doi:10.1002/9781119902911.

9. May, R. M. Will a Large Complex System be Stable? *Nature* **238**, 413–414 (1972).

10. Angelis, D. L. D. Stability and Connectance in Food Web Models. *Ecology* **56**, 238–243 (1975).

11. McCann, K., Hastings, A. & Huxel, G. R. Weak trophic interactions and the balance of nature. *Nature* **395**, 794–798 (1998).

12. McCann, K. & Hastings, A. Re–evaluating the omnivory–stability relationship in food webs. *Proceedings of the Royal Society of London. Series B: Biological Sciences* **264**, 1249–1254 (1997).

13. Thébault, E. & Loreau, M. Trophic Interactions and the Relationship between Species Diversity and Ecosystem Stability. *The American Naturalist* **166**, E95–E114 (2005).

14. Thébault, E. & Loreau, M. The relationship between biodiversity and ecosystem functioning in food webs. *Ecological Research* **21**, 17–25 (2006).

15. Thébault, E. & Fontaine, C. Stability of Ecological Communities and the Architecture of Mutualistic and Trophic Networks. *Science* **329**, 853–856 (2010).

16. Yachi, S. & Loreau, M. Biodiversity and ecosystem productivity in a fluctuating environment: The insurance hypothesis. *Proceedings of the National Academy of Sciences* **96**, 1463–1468 (1999).

17. Doak, D. F. *et al.* The Statistical Inevitability of Stability-Diversity Relationships in Community Ecology. *The American Naturalist* **151**, 264–276 (1998).

18. Tilman, D. THE ECOLOGICAL CONSEQUENCES OF CHANGES IN BIODIVERSITY: A SEARCH FOR GENERAL PRINCIPLES [101]. *Ecology* **80**, 1455–1474 (1999).

19. De Mazancourt, C. *et al.* Predicting ecosystem stability from community composition and biodiversity. *Ecology Letters* **16**, 617–625 (2013).

20. Thibaut, L. M., Connolly, S. R. & Sweatman, H. P. A. Diversity and stability of herbivorous fishes on coral reefs. *Ecology* **93**, 891–901 (2012).

21. Zhao, L. *et al.* Biodiversity stabilizes plant communities through statistical-averaging effects rather than compensatory dynamics. *Nature Communications* **13**, 7804 (2022).

22. Elmqvist, T. *et al.* Response diversity, ecosystem change, and resilience. *Frontiers in Ecology and the Environment* **1**, 488–494 (2003).

23.   Ross, S. R. P.-J., Petchey, O. L., Sasaki, T. & Armitage, D. W. How to measure response diversity. *Methods in Ecology and Evolution* **14**, 1150–1167 (2023).

24.   Thibaut, L. M. & Connolly, S. R. Understanding diversity–stability relationships: Towards a unified model of portfolio effects. *Ecology Letters* **16**, 140–150 (2013).

25.   Fründ, J., Zieger, S. L. & Tscharntke, T. Response diversity of wild bees to overwintering temperatures. *Oecologia* **173**, 1639–1648 (2013).

26.   Yang, Q., Fowler, M. S., Jackson, A. L. & Donohue, I. The predictability of ecological stability in a noisy world. *Nature Ecology & Evolution* **3**, 251–259 (2019).

27.   Danet, A., Mouchet, M., Bonnaffé, W., Thébault, E. & Fontaine, C. Species richness and food-web structure jointly drive community biomass and its temporal stability in fish communities. *Ecology Letters* **24**, 2364–2377 (2021).

28.   Eschenbrenner, J. & Thébault, É. Diversity, food web structure and the temporal stability of total plant and animal biomasses. *Oikos* **2023**, e08769 (2023).

29.   Srednick, G. & Swearer, S. E. Understanding diversity–synchrony–stability relationships in multi-trophic communities. *Nature Ecology & Evolution* 1–11 (2024) doi:10.1038/s41559-024-02419-3.

30.   Ives, A. r., Klug, J. l. & Gross, K. Stability and species richness in complex communities. *Ecology Letters* **3**, 399–411 (2000).

31.   Shanafelt, D. W. & Loreau, M. Stability trophic cascades in food chains. *Royal Society Open Science* **5**, 180995 (2018).

32.   Raimondo, S., Turcáni, M., Patoèka, J. & Liebhold, A. M. Interspecific synchrony among foliage-feeding forest Lepidoptera species and the potential role of generalist predators as synchronizing agents. *Oikos* **107**, 462–470 (2004).

33.   Korpimäki, E., Norrdahl, K., Huitu, O. & Klemola, T. Predator–induced synchrony in population oscillations of coexisting small mammal species. *Proceedings of the Royal Society B: Biological Sciences* **272**, 193–202 (2005).

34.   McCann, K. S. The diversity–stability debate. *Nature* **405**, 228–233 (2000).

35. Fahimipour, A. K., Anderson, K. E. & Williams, R. J. Compensation masks trophic cascades in complex food webs. *Theoretical Ecology* **10**, 245–253 (2017).

36. Brose, U., Williams, R. J. & Martinez, N. D. Allometric scaling enhances stability in complex food webs. *Ecology Letters* **9**, 1228–1236 (2006).

37. Yodzis, P. & Innes, S. Body Size and Consumer-Resource Dynamics. *The American Naturalist* **139**, 1151–1175 (1992).

38. Vasseur, D. A. & Fox, J. W. Environmental fluctuations can stabilize food web dynamics by increasing synchrony. *Ecology Letters* **10**, 1066–1074 (2007).

39. Ives, A. R. & Carpenter, S. R. Stability and Diversity of Ecosystems. *Science* **317**, 58–62 (2007).

40. Ripa, J. & Ives, A. R. Food web dynamics in correlated and autocorrelated environments. *Theoretical Population Biology* **64**, 369–384 (2003).

41. Loreau, M. & Mazancourt, C. de. Species Synchrony and Its Drivers: Neutral and Nonneutral Community Dynamics in Fluctuating Environments. *The American Naturalist* **172**, E48–E66 (2008).

42. Vandermeer, J. Coupled Oscillations in Food Webs: Balancing Competition and Mutualism in Simple Ecological Models. *The American Naturalist* **163**, 857–867 (2004).

43. Brose, U. *et al.* Predator traits determine food-web architecture across ecosystems. *Nature Ecology & Evolution* **3**, 919–927 (2019).

44. Loreau, M. *et al.* Biodiversity as insurance: From concept to measurement and application. *Biological Reviews* **96**, 2333–2354 (2021).

45. Barbier, M., Arnoldi, J.-F., Bunin, G. & Loreau, M. Generic assembly patterns in complex ecological communities. *Proceedings of the National Academy of Sciences of the United States of America* **115**, 2156–2161 (2018).

46. Ghosh, S., Matthews, B. & Petchey, O. L. Temperature and biodiversity influence community stability differently in birds and fishes. *Nature Ecology & Evolution* 1–12 (2024) doi:10.1038/s41559-024-02493-7.

47. Dornelas, M. *et al.* Assemblage Time Series Reveal Biodiversity Change but Not Systematic Loss. *Science* **344**, 296–299 (2014).

48. Blowes, S. A. *et al.* The geography of biodiversity change in marine and terrestrial assemblages. *Science* **366**, 339–345 (2019).

49. Danet, A., Giam, X., Olden, J. D. & Comte, L. Past and recent anthropogenic pressures drive rapid changes in riverine fish communities. *Nature Ecology & Evolution* 1–12 (2024) doi:10.1038/s41559-023-02271-x.

50. Olden, J. D., Comte, L. & Giam, X. The Homogocene: A research prospectus for the study of biotic homogenisation. *NeoBiota* **37**, 23–36 (2018).

51. Wang, S. *et al.* Biotic homogenization destabilizes ecosystem functioning by decreasing spatial asynchrony. *Ecology* **102**, e03332 (2021).

52. Delmas, E., Brose, U., Gravel, D., Stouffer, D. B. & Poisot, T. Simulations of biomass dynamics in community food webs. *Methods in Ecology and Evolution* **8**, 881–886 (2017).

53. Brown, J. H., Gillooly, J. F., Allen, A. P., Savage, V. M. & West, G. B. Toward a metabolic theory of ecology. *Ecology* **85**, 1771–1789 (2004).

54. Hatton, I. A., Dobson, A. P., Storch, D., Galbraith, E. D. & Loreau, M. Linking scaling laws across eukaryotes. *Proceedings of the National Academy of Sciences* **116**, 21616–21622 (2019).

55. Gouhier, T. C., Guichard, F. & Gonzalez, A. Synchrony and Stability of Food Webs in Metacommunities. *The American Naturalist* **175**, E16–E34 (2010).

56. Williams, R. J. & Martinez, N. D. Simple rules yield complex food webs. *Nature* **404**, 180–183 (2000).

57. Lajaaiti, I. *et al.* EcologicalNetworksDynamics.jl A Julia package to simulate the temporal dynamics of complex ecological networks. (2024) doi:10.1101/2024.03.20.585899.

58. Christensen, V. & Pauly, D. ECOPATH II — a software for balancing steady-state ecosystem models and calculating network characteristics. *Ecological Modelling* **61**, 169–185 (1992).

59. Lefcheck, J. S. piecewiseSEM: Piecewise structural equation modelling in r for ecology, evolution, and systematics. *Methods in Ecology and Evolution* **7**, 573–579 (2016).

438   60.   Murphy, M. semEff: Automatic Calculation of Effects for Piecewise Structural Equation Models. (2021).

439   61.   Brooks, M. E. *et al.* glmmTMB Balances Speed and Flexibility Among Packages for Zero-inflated Generalized Linear Mixed Modeling. *The R Journal* **9**, 378–400 (2017).