Extensive transmission and variation in a functional receptor for praziquantel resistance in endemic *Schistosoma mansoni*

Duncan J. Berger[1]*, Sang-Kyu Park[2], Thomas Crellen[1,3], Tushabe John Vianney[4], Narcis B. Kabatereine[5], James A. Cotton[1,3], Richard Sanya[6] , Alison Elliot[6,7], Edridah M. Tukahebwa[5], Moses Adriko[5], Claire J. Standley[8], Anouk Gouvras[9], Safari Kinung'hi[10], Helmut Haas[11], Muriel Rabone[12], Aidan Emery[12], Poppy H. L. Lamberton[3], Bonnie L. Webster[12], Fiona Allan[12], Sarah Buddenborg[1], Matthew Berriman[1,13]*, Jonathan S. Marchant[2]*, Stephen R. Doyle[1]*, Joanne P. Webster[14]*

1. Wellcome Sanger Institute, Wellcome Genome Campus, Hinxton, Cambridgeshire, CB10 1SA, UK
2. Department of Cell Biology, Neurobiology and Anatomy, Medical College of Wisconsin, Milwaukee, Wisconsin 53226, USA
3. School of Biodiversity, One Health, and Veterinary Medicine, University of Glasgow, Glasgow, UK
4. Division of Biomedical Engineering, University of Glasgow, G12 8LT Glasgow, UK.
5. Vector Borne & Neglected Tropical Disease Control Division, Ministry of Health, Kampala, Uganda
6. Immunomodulation and Vaccines Programme, Medical Research Council/Uganda Virus Research Institute and London School of Hygiene and Tropical Medicine Uganda Research Unit, Entebbe, Uganda.
7. Department of Infection Biology, London School of Hygiene and Tropical Medicine, London, UK
8. Center for Global Health Science and Security, Georgetown University, 3900 Reservoir Rd NW, Washington DC 20007, USA
9. Global Schistosomiasis Alliance, Podium Space - Ealing Cross, 85 Uxbridge Road, London, W5 5BW, UK
10. National Institute for Medical Research (NIMR) Mwanza Centre, P.O Box 1462, Mwanza, United Republic of Tanzania
11. helminGuard, Süelfeld, Germany
12. Department of Life Sciences, The Natural History Museum, London, SW7 5BD, UK
    Wolfson Wellcome Biomedical Laboratories, Department of Life Sciences, The Natural History Museum, London, SW7 5BD, UK, London Centre for Neglected Tropical Disease Research (LCNTDR), London, UK
13. Current address: School of Institute of Infection & Immunity, College of Medical, Veterinary & Life Sciences, University of Glasgow, 120 University Place, Glasgow, G12 8TA, UK
14. Department of Pathology and Pathogen Biology, Royal Veterinary College, University of London, Herts, UK

* Corresponding authors

Underline - co-supervised this work

Abstract

Mass-drug administration (MDA) of human populations using praziquantel monotherapy has become the primary strategy for controlling and potentially eliminating the major neglected tropical disease schistosomiasis. To understand how long-term MDA impacts schistosome populations, we analysed whole-genome sequence data of 570 *Schistosoma mansoni* samples (and the closely related outgroup species, *S. rodhaini)* from eight countries incorporating both publicly-available sequence data and new parasite material. This revealed broad-scale genetic structure across countries but with extensive transmission over hundreds of kilometres. We characterised variation across the transient receptor potential melastatin ion channel, $TRPM_{PZQ}$, a target of praziquantel, which has recently been found to influence praziquantel susceptibility. Functional profiling of $TRPM_{PZQ}$ variants found in endemic populations identified four mutations that reduced channel sensitivity to praziquantel, indicating standing variation for resistance. Analysis of parasite infrapopulations sampled from individuals pre- and post-treatment identified instances of treatment failure, further indicative of potential praziquantel resistance. As schistosomiasis is targeted for elimination as a public health problem by 2030 in all currently endemic countries, and even interruption of transmission in selected African regions, we provide an in-depth genomic characterisation of endemic populations and an approach to identify emerging praziquantel resistance alleles.

Introduction

Schistosomiasis is a neglected tropical disease which currently infects 250 million people across 78 endemic nations (Control of Neglected Tropical Diseases (NTD) Guidelines Review Committee, 2022; Lo *et al.*, 2022). Infections are prevalent among children, including pre-school-aged children, as well as adults in low and middle-income countries, with 90% of infections occurring in sub-Saharan Africa. The etiological agents of schistosomiasis are freshwater snail-borne parasitic trematodes of the genus *Schistosoma* (principally *Schistosoma mansoni, S. japonicum,* and *S. haematobium*). These parasitic worms dwell in the host's blood vessels, often for years, where they lay eggs, many of which become trapped in host tissues, resulting in a spectrum of pathologies including anaemia, stunted growth, genital lesions, fever and irreversible organ damage (Colley *et al.*, 2014). Recognising the severe and widespread impact of this disease, the World Health Organization (WHO) endorsed praziquantel monotherapy, distributed as part of mass drug administration (MDA) programmes, as the primary strategy for schistosomiasis control (WHO Expert Committee on the Control of Schistosomiasis, 2002; Webster *et al.*, 2014). In 2020 alone, 76.9 million people (representing 44.9% of those requiring treatment), predominantly school-aged children, were treated with praziquantel for schistosomiasis (World Health Organization, 2021). Such treatment campaigns have resulted in an overall decrease in the prevalence of schistosomiasis among school-aged children by approximately 60% (Deol *et al.*, 2019; Kokaliaris *et al.*, 2022). Based on such successes, the WHO launched its 2021-2030 NTD roadmap and revised Guidelines for the Control and Elimination of Schistosomiasis (Control of Neglected Tropical Diseases (NTD) Guidelines Review Committee, 2022). These set ambitious goals to eliminate schistosomiasis as a public health problem in all endemic countries (defined as reducing the proportion of heavy-intensity infections to <1%) and complete interruption of transmission in selected regions by 2030, primarily through the escalating use of praziquantel MDA (World Health Organization, 2020).

Although MDA programmes have, in general, resulted in large-scale reductions in schistosomiasis prevalence and morbidity in endemic regions (Kabatereine *et al.*, 2007; Bronzan *et al.*, 2018; Deol *et al.*, 2019), some of these prevalence reductions have proven to be reversible over short timescales and, despite years of repeated MDA, persistent hotspots of infection remain (Kittur *et al.*, 2019; Mutuku *et al.*, 2019; Mawa *et al.*, 2021; Lim *et al.*, 2023). Ongoing surveillance projects have also revealed substantial heterogeneity in infection prevalence, intensity and morbidity within and between endemic regions (Meurs *et al.*, 2013; Aemero *et al.*, 2015; Mari *et al.*, 2017). This includes potential variability in parasite

response to praziquantel, which, given the current lack of available vaccines or alternative
antischistosomal drugs, represents a major threat to the control and elimination of
schistosomiasis (Stelma *et al.*, 1995; Melman *et al.*, 2009; Crellen, Walker, *et al.*, 2016). In
combination with other public health measures, MDA programmes are expected to
dramatically alter patterns of schistosome transmission and exert selective pressure for
praziquantel resistance. However, despite over twenty years of MDA in some areas, there is
no consistent evidence that sustained praziquantel administration has impacted parasite
population genetic structure or diversity (Blanton *et al.*, 2011; Huyse *et al.*, 2013; Lelo *et al.*,
2014; Faust *et al.*, 2019). Likewise, while there is clear evidence that resistance to
praziquantel can be rapidly selected for in laboratory settings (Fallon and Doenhoff, 1994;
Couto *et al.*, 2011; Mwangi *et al.*, 2014; Lamberton, Faust and Webster, 2017), there is
limited evidence of established praziquantel resistance in endemic *S. mansoni* populations
(Fukushige *et al.*, 2021). This contrasts with that of emerging or established heritable
resistance, particularly in the veterinary helminths, to every other class of anthelmintic used
(Schwab *et al.*, 2005; Kaplan and Vidyashankar, 2012; Rose *et al.*, 2015; Doyle *et al.*, 2022).
Part of the explanation lies in the fact that, until recently, we neither knew the precise mode
of action of praziquantel against schistosomes nor had any molecular markers to monitor
potential praziquantel resistance amongst natural *Schistosoma* spp. populations. Recent
efforts have, for the first time, identified the molecular target of praziquantel, a transient
receptor potential (TRP) melastatin ion channel (*Sm.*TRPM$_{PZQ}$) (Park *et al.*, 2019, 2021; Le
Clec'h *et al.*, 2021; Marchant, 2024). This discovery permits evaluation of praziquantel
efficacy and resistance risk, although such resistance-conferring alleles have only been
identified in a single isolate from endemic populations (Le Clec'h *et al.*, 2021). Characterising
the transmission and recent evolution of schistosome populations is, therefore, of broad
epidemiological importance as a means to understand how parasite populations are
structured and how they are changing in response to interventions aimed at controlling
schistosomiasis.
130
Here, we have used whole-genome sequencing to characterise genomic variation from
globally dispersed populations of the major human infective species *S. mansoni* (and the
closely related outgroup species, *S. rodhaini)* (Crellen, Allan, *et al.*, 2016; Berger *et al.*,
2021; Vianney *et al.*, 2022). Our analyses focus on populations within endemic regions of
East Africa, primarily those found around Lake Victoria, one of the largest foci of
schistosomiasis infections and a target of long-term MDA efforts (Crellen, Walker, *et al.*,
2016; Deol *et al.*, 2019). We aimed to quantify variation within the newly identified candidate
praziquantel resistance locus, *Sm.*TRPM$_{PZQ}$ (Le Clec'h *et al.*, 2021; Park *et al.*, 2021;
Chevalier *et al.*, 2024), to determine the prevalence of potential resistance-conferring

mutations. This also included extensive WGS of *S. mansoni* from within individual hosts

before and after praziquantel administration to quantify the immediate genetic impact of

treatment. These data raise significant implications regarding the standing variation of

potential praziquantel resistance in human schistosome population and represent an

important resource for assessing the efficacy of current interventions and guiding future

treatment strategies.

Results

The genetic diversity of geographically dispersed isolates supports high transmission

in endemic regions

We analysed whole-genome sequence data from 574 *Schistosoma* samples (n = 570 *S.*

mansoni and n = 4 *S. rodhaini* ; hereafter all referred to as accessions) isolated from eight

countries, including 207 new *Schistosoma* samples sequenced for this study (Figure 1a;

Table 1; Supplementary Data 1) (Crellen, Allan, *et al.*, 2016; Berger *et al.*, 2021; Vianney *et*

al., 2022). Most accessions were derived from Lake Victoria (88.3%), a major focus of *S.*

mansoni infection. These included published sequence data from miracidial samples isolated

from infected children between 2014 and 2017 from either the Koome & Damba Islands

(hereafter referred to as the Koome Islands; n = 174) (Vianney *et al.*, 2022) or Southern

Uganda (n = 164) (Berger *et al.*, 2021), with a median of two isolates sequenced per child

(range 1-11). We sequenced an additional 164 miracidia from just three children from

Southern Uganda (n = 89, 51 & 23 miracidia per child), including both pre- (n = 82) and post-

praziquantel treatment (n = 82) sampling (Crellen, Walker, *et al.*, 2016). The only other Lake

Victoria accessions were derived from cercariae shed from snails captured in Northern

Tanzania (n = 31) or Southern Uganda (n = 2). Of the remaining samples, two were

cercariae from the shoreline of Lake Albert, four were cercariae from a passaged strain

originally isolated in Lake Albert, 17 were miracidia collected from children living in Eastern

Uganda, and ten were published adult-stage isolates of passaged laboratory strains

originally sampled from Kenya (n = 1), Senegal (n = 3), Cameroon (n = 1), Guadeloupe (n =

4) or Puerto Rico (n = 1) (Crellen, Allan, *et al.*, 2016). As an outgroup, we included four *S.*

rodhaini accessions, two cercarial isolates from Tanzania and two adult-stage isolates of

passaged lab strains from isolates originally sampled in Burundi. We mapped sequence

reads from all 574 accessions to the *S. mansoni* reference genome, and after variant calling

and quality control, we identified 35,146,249 single-nucleotide polymorphisms (SNPs) and

6,632,156 indels across all accessions (Supplementary Data 2).

To characterise population structure, we produced a maximum likelihood phylogeny and
performed principal component analysis using unlinked autosomal variants. These analyses
distinguished accessions sampled in East Africa from those in West Africa and the
Caribbean (Figure 1b,c). The only exception was a single laboratory-passaged isolate from
coastal Kenya that clustered with West African accessions, consistent with previous
analyses of this sample (Crellen, Allan, *et al.*, 2016; Platt *et al.*, 2021). Within Eastern Africa,
populations from Lake Victoria, Lake Albert and Eastern Uganda formed distinct but partially
overlapping clusters. Pairwise measures of the fixation index (F_{ST}) showed limited population
differentiation between Lake Victoria, Lake Albert and Eastern Uganda (range F_{ST} = 2.49×10^{-}
^{2} - 4.59×10^{-2}) and negligible differentiation between Lake Victoria populations, despite the
geographic separation of up to 300 km (range F_{ST} = 2.41×10^{-3} - 4.07×10^{-3}, Supplementary
Figure 1a). East African populations had comparable levels of nucleotide diversity (range
median π = 2.35×10^{-3} - 3.35×10^{-3}; Supplementary Figure 1b) and effective population size
(N_e) estimates (range N_e = 58,004-64,063; Supplementary Figure 2; Supplementary Data 3)
suggesting shared recent population histories. Analyses of population ancestry for individual
accessions identified similar population compositions across all Lake Victoria populations
with variable ancestry contributions from Lake Albert and West African populations (Figure
1f, Supplementary Data 4). Accessions from Eastern Uganda displayed variable, mixed
ancestry from Lake Victoria, Lake Albert and West Africa populations, suggesting migration
from multiple regions.
196
Ancestry analyses revealed low levels of *S. rodhaini* admixture (range: 0.00-2.32%),
potentially indicative of interspecific hybridisation, across most unrelated *S. mansoni*
accessions (Figure 1d; Supplementary Data 4). In East Africa, *S. mansoni* and *S. rodhaini*
are broadly sympatric and share both intermediate and definitive hosts, with *S. mansoni*
infecting both specific primate and rodent species and *S. rodhaini* infecting rodents, allowing
opportunities for inter-species pairings and hybridisation. However, due to the small number
of *S. rodhaini* isolates included in this study (*n* = 4) and the lack of a non-admixed, outgroup
population, the incidence of recent admixture and/or historic introgression was not
investigated further.

Figure 1: Population structure of *Schistosoma* accessions. a) Global distribution of the 574 *Schistosoma*

accessions used in this study. Sampling locations are coloured by geographical divisions: Puerto Rico (PR;

brown), Guadeloupe (GP; orange), Senegal (SN; dark purple), Cameroon (pink), Coastal Kenya (KE; teal), Lake

Albert (dark blue), Eastern Uganda (red), Southern Uganda (light blue), Koome group islands (yellow), Northern

Tanzania (dark green). Black points represent the sampling coordinates of *S. rodhaini* accessions. **b)** The

maximum-likelihood phylogenetic tree was inferred using 188,923 autosomal single-nucleotide polymorphisms

(SNPs) and all 574 accessions. Branches are coloured based on the subdivisions described in a), and the tree is

rooted on *S. rodhaini*. Highlighted clade 'C1' represents all West African (Senegalese and Cameroonian),

Caribbean (Puerto Rican and Guadeloupean) and Coastal Kenyan accessions. **c)** Principal component analysis

(PCA) of genetic differentiation between 505 unrelated *S. mansoni* accessions using 214,445 autosomal SNPs.

Points are coloured and shaped based on the groups described in a). **d)** ADMIXTURE plots illustrating the

inferred ancestry of 505 unrelated *S. mansoni* and four *S. rodhaini* accessions. Here, we assume five populations

(K) are present, inferred using 10-fold cross-validation and a standard error estimation with 250 bootstraps. Y-

axis values show the admixture proportions for each accession, and colours for each population were assigned

based on the majority ancestry of each geographical division.

Table 1: Sample information and history of praziquantel treatment.

Species	Country	Region	Development Stage			Sampling stage			Reference
			Adults	Cercariae	Miracidia	Pre-treatment	Post-treatment	Not applicable	
S. mansoni	Uganda	Lake Albert	1	6	0	0	0	7	Herein; Crellen et al. 2016
	Uganda	Eastern Uganda	0	0	17	17	0	0	Berger et al. 2021
	Uganda	Southern Uganda	1	2	328	201	127	3	Herein; Berger et al. 2021
	Uganda	Koome Islands	0	0	174	123	51	0	Vianney et al. 2022
	Tanzania	Northern Tanzania	0	31	0	0	0	31	Herein
	Kenya	Coastal Kenya	1	0	0	0	0	1	Crellen et al. 2016
	Cameroon		1	0	0	0	0	1	Crellen et al. 2016
	Senegal		3	0	0	0	0	3	Herein; Crellen et al. 2016
	Puerto Rico		1	0	0	0	0	1	Crellen et al. 2016
	Guadeloupe		4	0	0	0	0	4	Crellen et al. 2016
S. rodhaini	Tanzania	Northern Tanzania	0	2	0	0	0	2	Herein
	Burundi		2	0	0	0	0	2	Herein; Crellen et al. 2016

This study included 574 accessions, sequenced from parasite samples derived from snail hosts (cercariae), humans (miracidia) or laboratory-passaged strains (adults). Publicly-available data (*n* = 339) and new sequencing data (*n* = 208; defined as "herein") were generated for this study. Miracidia accessions were sampled pre-treatment with praziquantel (40 mg/kg) or four weeks post-treatment.

Functionally relevant genomic variation influences praziquantel sensitivity

A transient receptor potential (TRP) melastatin ion channel (*Sm.*TRPM$_{PZQ}$; *Smp_246790*) has recently been proposed as the target of praziquantel (Park *et al.*, 2019, 2021; Le Clec'h *et al.*, 2021; Marchant, 2024). This candidate gene underlies variation in praziquantel susceptibility, at least within schistosome laboratory populations (Park *et al.*, 2019, 2021; Le Clec'h *et al.*, 2021; Summers *et al.*, 2022). Mutagenesis of key residues within the praziquantel binding site of TRPM$_{PZQ}$ (located in the transmembrane voltage-sensor-like domain, VSLD) resulted in the loss of sensitivity to praziquantel (Park *et al.*, 2021).

Consistent with previous genomic surveys (Berger *et al.*, 2021; Vianney *et al.*, 2022), analysis of haplotype diversity using the integrated haplotype score (iHS) did not suggest that *Sm.*TRPM$_{PZQ}$ was under positive selection in Lake Victoria populations (Supplementary Figure 3, Supplementary Data 5,6). However, analysis of variation within *Sm.*TRPM$_{PZQ}$ identified a high degree of variability with a predicted 496 amino acid changes at 433/2268 residues (Figure 2), with mutations at 174 of these residues found only in single accessions (Supplementary Data 7). We generated point mutations to modify 12 *Sm.*TRPM$_{PZQ}$ residues, and used an *in vitro* Ca^{2+} reporter assay (Park *et al.*, 2019) to assess their impact on *Sm.*TRPM$_{PZQ}$ function. Eight of these mutants, all conservative amino acid substitutions, exhibited similar praziquantel sensitivity (range EC$_{50}$ = 0.49-0.96 μM) to the wild-type *Sm.*TRPM$_{PZQ}$ channel (EC$_{50}$ = 0.74 μM [standard error (SE) = 0.17]; Figure 3a; Supplementary Data 8; Supplementary Figure 4). Mutants p.T1624K and p.R1843Q exhibited a decreased sensitivity to praziquantel (EC$_{50}$ = 1.41 μM [SE = 0.23] and EC$_{50}$ = 1.11 μM [SE = 0.08], respectively) and two mutants (p.Y1554C, p.Q1670K) caused a complete loss in praziquantel sensitivity (Figure 3b; Supplementary Data 8; Supplementary Figure 5).

266
Figure 2: Genetic variation in the candidate mediator of praziquantel susceptibility, the transient potential
receptor channel *Sm*.TRPM_{PZQ}. Mutational frequency along the protein structure of *Sm*.TRPM_{PZQ}
(*Smp_246790*). Frequencies of mutations are reported across 550 *S. mansoni* accessions (y-axis), representing
samples from Eastern Uganda ($n = 17$), Southern Uganda ($n = 328$), the Koome islands ($n = 174$) and Northern
Tanzania ($n = 31$). X-axis values represent the location of the mutations on the protein, and bars (and terminal
points) are coloured by the predicted impact of each mutation. Structure of *Sm*.TRPM_{PZQ}: N-terminal TRPM
homology region (MHR) domain, ankyrin-like repeat domain (AD), pre-S1 helix (shaded), TM-spanning helices
(1-6), pore helices (PH), TRP domain (TRP), rib helices (R), pole helices (P), coiled-coil (CC) region and the
COOH terminal NUDT9H domain (NUDT9H).

Seven of the twelve mutations mapped within an existing homology model for the
transmembrane-spanning region of *Sm*.TRPM_{PZQ}, enabling structural insight into these
functional effects (Figure 3c) (Park *et al*., 2021). Four mutations had no impact on
praziquantel action (p.V1405I, p.L1476I, p.S1341N & p.F1399Y; Figure 3a): p.V1405I and
p.L1476I are found within extracellular loops of the channel, p.S1341N lies the COOH-
terminal cytoplasmic domain and p.F1399Y, while within the voltage-sensor like domain
(VSLD) of the channel that harbours the praziquantel binding site, is remote (~10Å away)
from the praziquantel binding poise (Figure 3c). Mutation p.T1624K (~2-fold lower sensitivity;
Figure 3b) projects into the extracellular milieu before the start of S6 and may cause sub-
optimal orientation of the S6 helix. The other variant associated with lower sensitivity
(p.R1843Q; Figure 3b) lies outside the existing TRPM_{PZQ} homology model but is predicted to

localise within the COOH-terminal coiled-coil region of the channel, which is involved in

subunit interaction (Mei *et al.*, 2006). Tetramer misfolding and lower channel expression

would explain the decreased sensitivity of this variant.

The first of the two mutations that caused a complete loss of praziquantel sensitivity (Figure

3b), p.Y1554C, is found in the S5 helix (Figure 3c). This residue projects toward S6, where it

is predicted to interact with a tyrosine residue (Y1636) in the S6 helix. This cysteine mutant

would be expected to disrupt this interaction, which is likely important for the appropriate

orientation of the helices that form the pore-forming domain. The second, p.Q1670K,

introduces a positive charge within the intracellular TRP helix close to the bilayer interface.

This is a critical region for channel gating, and the mutation likely impacts the orientation of

the TRP helix relative to the bilayer and interactivity with the S4/S5 loop of $Sm.$TRPM$_{PZQ}$,

which regulates TRPM channel activation (Yin *et al.*, 2019). The deleterious effects of

p.Y1554C and p.Q1670K are, therefore, consistent with the known importance of these

regions for channel function (Park *et al.*, 2021).

Of the four mutations that decreased or eliminated praziquantel sensitivity, three (p.Y1554C,

p.Q1670K, p.T1624K) were only found in a heterozygous state in single accessions

(Supplementary Data 7,9,10). p.R1843Q was far more common, with homozygous variants

(c.5528G>A) identified in 3.68% ($n = 21$ accessions across the entire dataset; $n = 15$

miracidia from 12 individuals) of accessions and heterozygous in 22.6% of accessions ($n =$

129 accessions across the entire dataset; $n = 120$ miracidia from 68 individuals). This variant

was also more prevalent in post-treatment populations (5.05% of post-treatment accessions;

$n = 9$ miracidia from 8 individuals) compared to pre-treatment populations (1.75% of pre-

treatment accessions; $n = 6$ miracidia from 4 individuals; Supplementary Data 10), although

accessions with this variant (and all other evaluated variants) did not form a distinct

subpopulation (Supplementary Figures 6-9). In addition to the mutations analysed via

targeted mutagenesis, we identified a second high-prevalence homozygous frameshift

mutation, p.Q1915fs, in 94.0% of accessions. Park and colleagues (2021) reported that

truncation mutations (Δ1914) of the C-terminal NUDT9H domain do not impact the TRP

channels' responsiveness to praziquantel.

321
322

Figure 3: Functional profiling of variants of the transient potential receptor channel *Sm*.TRPM$_{PZQ}$.

Concentration-response relationships for consensus Sm.TRPM$_{PZQ}$ sequence compared with twelve Sm.TRPM$_{PZQ}$

variants, which exhibit an average EC$_{50}$ to ±PZQ of **a)** <1 µM (green) or **b)** >1 µM (orange). Constructs for which

no praziquantel-evoked activity was observed (Y1554C, Q1670K) are shown in (b) in red. Results represent

mean±sem from at least three independent transfections. **c)** *Inset*, homology model of the transmembrane

spanning region (residues 1100 to 1800) of a *Sm*.TRPM$_{PZQ}$ monomer from [2]. The enlarged view shows the

location of the functionally profiled variants. Praziquantel and residues within 5Å of the praziquantel binding poise

(white) are shown at the base of the voltage-sensor-like domain (VSLD) of the channel, with the S5 and S6 pore-

forming helices to the left.

Le Clec'h and colleagues recently identified three putative marker variants for praziquantel

resistance in selected laboratory lines (Le Clec'h *et al.*, 2021). The first of these was a

homozygous SNP predicted to result in a synonymous mutation ($Sm.$TRPM$_{PZQ}$-2723187C;

p.I1020l), which we identified in 6.14% ($n = 35$, $n = 30$ miracidia from 21 individuals) of

accessions (Supplementary Data 7,9,10). This mutation was nearly twice as common in

post-treatment populations (8.43% of accessions; $n = 15$ miracidia from 10 individuals)

compared to pre-treatment populations (4.40% of accessions; $n = 15$ miracidia from 13

individuals). A further 24.9% of accessions had a single copy of this marker. The second and

third markers were 150 kb deletions adjacent to *Smp_246790* (~1.22 Mb) and *Smp_345310*

(~3.18-3.33 Mb), respectively. Structural variant genotyping did not identify the former

variant in our accessions, but we did find a series of long homozygous deletions (69.9-215.0

kb) located between 3.02-3.36 Mb in 43.3% of genotyped accessions ($n = 170$ of 393

genotyped accessions; Supplementary Data 11, Supplementary Figure 10). However, the

150 kb deletion was not enriched in post-treatment populations, and neither the deletion nor

p.I1020l clustered phylogenetically (Supplementary Figure 9).

Host infrapopulations reveal the extent of genetic relationships and evidence of

treatment failure

Sampling and analysis of the parasite population from within a single host (defined as the

'infrapopulation') across specific time points has the potential to identify instances of

treatment failure, which may be indicative of praziquantel resistance. However, these

infrapopulations are thought to be highly heterogeneous (Agola *et al.*, 2009; Faust *et al.*,

2019; Neves, Webster and Walker, 2019), and previous genomic surveys of *S. mansoni*

have only included small numbers of parasites from the same individuals (max $n = 11$)

(Chevalier *et al.*, 2016, 2019; Berger *et al.*, 2021; Vianney *et al.*, 2022) reducing the

likelihood of identifying related parasites. To provide greater resolution of infrapopulation

structure and evaluate the impact of praziquantel treatment, we analysed pairwise kinship

between 164 sequenced miracidia sampled pre- or post-praziquantel treatment from a total

of three donors: Bb1 ($n = 47$ pre-treatment, $n = 42$ post-treatment), Bu3 ($n = 33$ pre-

treatment, $n = 19$ post-treatment) and Bu1 ($n = 2$ pre-treatment, $n = 21$ post-treatment) and

reanalysed the 174 accessions from the Vianney et al. (Vianney *et al.*, 2022) survey (98

donors, $n = 123$ pre-treatment, $n = 51$ post-treatment).

Figure 4: Relatedness between *Schistosoma mansoni* accessions from endemic regions. **a)** Ternary plots

representing pairwise relationships between accessions using the three relatedness coefficients K_0, K_1, and K_2,

representing the probabilities that at a given locus, the two accessions shared zero, one or two-allele identity-by-

descent, respectively. Each point represents a pairwise relationship between two accessions, showing whether

each accession was from the same (triangles) or different (circles) donor. Points are coloured by the inferred

relationship: first-degree (pink), second-degree (green), third-degree (orange) or unrelated (blue). Only

accessions from three Southern Ugandan donors and all Koome Island donors are shown. **b)** Inferred

relationships within and between donors. The outer lines represent each donor; within each line, individual

accessions are shown coloured by whether they were sampled before (dark grey) or after (light grey) treatment

with praziquantel. Coloured lines represent the same pairwise relationships as a). For donors Bb1 and Bu3, all

relationships are shown below the circular plot and shown in boxes where they are grouped into pre-treatment

samples, post-treatment samples, and pre- and post-treatment samples (spanning treatment). Within the boxes,

dotted lines represent relationships inferred by NgsRelate but were found to have a lower degree or no-relation

by Sequoia. **c)** The effect of praziquantel treatment on nucleotide diversity (π) was calculated for each donor

infrapopulation (Bb1, Bu1 and Bu3) and all Koome group island samples. Nucleotide diversity was calculated in 5

kb non-overlapping windows across each autosome for each pre-treatment (purple) and post-treatment (yellow)

population (including related samples). For all boxplots, the central line indicates the median, and the top and

bottom edges of the box indicate the 25th and 75th percentiles, respectively. The maximum whisker lengths are

specified as 1.5 times the interquartile range. **d)** Identification of a circulating lineage of *S. mansoni* on Damba

Island (the northernmost island of the Koome group islands). We identified a cluster of nine accessions, all with

third-degree identical by descent relationships, which clustered phylogenetically with accessions from Lake

Albert.

We identified 49 first-degree relationships, 13 second-degree relationships and 34 third-

degree relationships (Figure 4a; Supplementary Data 12). Forty-seven of the 49 first-degree

relationships were between accessions sampled from the same donor; the remaining two

were between donors and have previously been identified as potentially mislabelled

accessions (Vianney *et al.*, 2022). Estimates of the coefficient of inbreeding (*f*) using the

condensed identity coefficients also revealed low levels of inbred relatedness (median *f* =

0.000227; range *f* = 1×10^{-6} - 0.35) in 18.8% of parents (Supplementary Data 13) (Jacquard,

1966). Within infrapopulations of donors Bb1, Bu1 and Bu3, we identified three first-degree

relationships between pre-treatment accessions and 23 between post-treatment accessions.

Most of these were found in infrapopulations from Bb1, represented by three independent

clusters of four accessions (Figure 4b,c). We also identified one first-degree and four

second-degree relationships spanning treatment arms, suggesting bi- and uni-parental

survival, respectively. Both of the first-degree relatives were also homozygous for the

*Sm.*TRPM$_{PZQ}$-2723187C marker; however, none of the second-degree relatives had a

homozygous or heterozygous copy. Additionally, we did not identify any homo- or

heterozygous variants of p.R1843Q in any of the first- or second-degree relatives spanning

treatment. We also found no evidence of reduced nucleotide diversity in post-treatment

populations (Figure 4c) and low genetic differentiation between pre- and post-treatment

populations (range mean F_{ST} = 1.24×10^{-3} - 1.75×10^{-3}, Supplementary Figure 11).

Examination of third-degree relationships within the Vianney and colleagues (Vianney *et al.*,

2022) dataset identified a cluster of nine related accessions, all localised to Damba Island

and found in all four of the sampled villages (Figure 4b,d). Phylogenetic analysis showed all

nine formed a distinct clade with six accessions from Lake Albert (Supplementary Figure 12),

suggesting that these are derived from a recently imported lineage from Lake Albert, recently

dispersed across Damba Island.

Discussion

As efforts to interrupt transmission and eliminate schistosomiasis as a public health problem intensify over the next decade, the escalating use of MDA is expected to result in increased selective pressures, altered transmission dynamics and widespread reductions in schistosome populations (World Health Organization, 2020; Control of Neglected Tropical Diseases (NTD) Guidelines Review Committee, 2022). The consequences of such large-scale changes to schistosome populations are unclear but provide a strong incentive for genomic surveillance to detect and monitor such changes. In this study, we have, to our knowledge, assembled the most comprehensive collection of *Schistosoma* whole-genome sequencing data to date, including samples from across eight countries and two major foci of infection.

Praziquantel resistance represents a credible threat to schistosomiasis control if it were to establish in endemic populations. Despite over 50 years of use, $TRPM_{PZQ}$ has only recently been identified as a direct target of praziquantel and a genetic determinant of reduced praziquantel susceptibility (Le Clec'h *et al.*, 2021; Park *et al.*, 2021). We identified extensive, low-frequency variation within this gene and used this variation database to inform the targeted mutagenesis of 12 residues, focusing on mutations at potential key residues. We identified four mutations in our population genetic analyses that reduced or ablated channel responsiveness to praziquantel functional genetic *in vitro* assays, and while only one (p.R1843Q) was present in more than a single accession, all represent standing variation for praziquantel resistance in endemic populations. Both p.R1843Q and p.I1020l (the latter a marker for praziquantel resistance in passaged laboratory lines (Le Clec'h *et al.*, 2021)) variants were present in multiple accessions, indicating potential variability in praziquantel efficacy in these populations. Both mutations were also more prevalent in post-treatment parasite populations than their pre-treatment counterparts. However, our analyses and others (Huyse *et al.*, 2013; Faust *et al.*, 2019; Berger *et al.*, 2021) have shown that post-treatment populations do not typically represent subpopulations of pre-treatment parasites, which may result from reinfection or under-sampling. It is possible that the increased prevalence of these mutations is unrelated to praziquantel administration. It is clear, nevertheless, that despite the asymmetrical per-host sampling of this study, these variants remain widespread in parasite populations sampled from a wide range of hosts, showing that they are prevalent across Lake Victoria. Further work should also be undertaken to determine to what extent the mutations we identified impact the *in vivo* efficacy of praziquantel and parasite fitness. Profiling these specific SNPs in the context of other

changes within $TRPM_{PZQ}$ in the same samples, reflecting broader allelic variation, is also

necessary.

Our analyses have demonstrated how genomic surveillance of parasites can help identify

variants contributing to reduced praziquantel efficacy in endemic populations and provide

structural and functional insights into $TRPM_{PZQ}$ that will permit a better understanding of

praziquantel efficacy in schistosomes. While we only characterised a small proportion of the

total variation we observed, our results survey genetic variation within endemic regions

where praziquantel has seen extensive use. These data provide candidate variants for

functional investigations and serve as a database of variation for retrospective surveys once

further resistance-conferring mutations are identified. However, it is notable that a large

proportion of our dataset was derived from donors living in regions of long-term praziquantel

administration.

Targeted mutagenesis identified four mutations in endemic populations that reduced channel

sensitivity to praziquantel, indicating standing variation for resistance. Whilst we did identify

both standing variation for future resistance evolution and instances of treatment failures

here, the apparent lack of obvious high-frequency resistance-conferring mutations is

encouraging as these populations have been under long-term MDA pressure. The variable

coverage of MDA may have allowed substantial refugia populations to exist in untreated

humans, snails and animal reservoirs. Refugia would reduce the selection pressure for

resistance, an approach that has been deliberately employed in veterinary systems (Kenyon

et al., 2009; Leathwick, Ganesh and Waghorn, 2015; Hodgkinson *et al.*, 2019). Alternatively,

reduced praziquantel susceptibility might be linked to regulatory changes in expression or

RNA splicing (Le Clec'h *et al.*, 2021), neither of which would be identified by our analyses.

Considering how recently *Sm.*$TRPM_{PZQ}$ was recognised as a mediator of praziquantel

sensitivity, future functional investigations will likely reveal additional relevant variants.

Despite involving a small number of donors, our analyses of host infrapopulation dynamics

represent the most thorough genomic characterisation of a host infrapopulation of any

parasitic helminth. Our analyses found that host infrapopulations are exceptionally diverse,

consistent with both autopsy studies and genetic surveys, which have found that individual

worm burdens range from one up to hundreds or thousands of worm pairs per individual in

low and high-endemicity regions, respectively (Gryseels and De Vlas, 1996; Neves *et al.*,

2021). Despite this, we did find evidence of sibling relationships within both treatment arms,

with indications of a higher degree of post-treatment relatedness. We also found evidence of

post-treatment parental survival, suggesting either ineffective treatment or higher

praziquantel tolerance within these populations. However, the limited number of these

accessions limited any formal analyses of the possible genetic basis of reduced praziquantel

efficacy. Consistent with previous genetic studies, these only represented a small proportion

of the overall post-treatment populations (Faust *et al.*, 2019), suggesting that most post-

treatment samples represent either infection with additional parasites and/or parasites

missed in pre-treatment sampling.

To conclude, we have characterised endemic *S. mansoni* populations at multiple spatial

scales ranging from comparisons across major foci of infection to individual parasite

infrapopulations. Our variation analysis within a candidate praziquantel resistance loci

enabled us to identify multiple novel resistance-conferring mutations. These remain at low

frequencies in studied populations but demonstrate standing variation from praziquantel

resistance in endemic populations undergoing MDA. Our analyses also provide a resource

for retrospective or confirmatory support of functional analyses of this recently identified

locus. Finally, our extensive sequencing of host infrapopulations provided an in-depth

genomic characterisation of parasite diversity and how this is impacted by praziquantel

treatment. Overall, our study summarises *S. mansoni* genomic diversity and represents a

resource to assess the efficacy of current interventions and guide future treatment

strategies.

Methods

The data analysed here incorporated whole-genome accessions from published datasets from eight countries and 205 new *Schistosoma* sample data (Figure 1a; Table 1; Supplementary Data 1). The sections below describe the origin, collection, and ethical approval of samples, focusing only on the parasite samples sequenced for this study and not previously published accessions.

Collection and ethical approval for the sampling of Ugandan miracidia

The collection of all Ugandan miracidial samples was undertaken as part of the monitoring, evaluation and disease control activities conducted by the Vector Control Division of the Ministry of Health (Uganda), the Schistosomiasis Control Initiative and Imperial College London. All methods and data collection were approved by the Uganda National Council for Science and Technology (Memorandum of Understanding: sections 1.4, 1.5, 1.6) and the Imperial College Research Ethics Committee (EC NO: 03.36. R&D No: 03/SB/033E). The Head of the Vector Control Division informed local district officials, and the headteachers of each school were informed about the study and requested to provide informed consent to allow sampling to be performed within the school. Parents of the children were informed of the study through school meetings, where they were provided with detailed information regarding the purpose of the study and technical staff were present to answer questions. Parents were requested to provide informed consent for their children to participate in the study, and in addition, any children aged ten or older were asked to give informed consent after receiving complete information about the study. Participation was voluntary, access to treatment was not dependent on participation in the study, and children could withdraw or be withdrawn from the study at any time.

Stool samples were collected from each child, and duplicate Kato-Katz thick smears were conducted (Katz, Chaves and Pellegrino, 1972). A Pitchford–Visser funnel was used to wash and filter parasite eggs from the remaining stool sample, and the filtrate was stored overnight (Pitchford and Visser, 1975). Miracidia were hatched the following day and were transferred into two sequential dishes of nuclease-free water to dilute bacterial contaminants before being individually fixed onto Whatman FTA-indicating classic cards (Gower *et al.*, 2007; Emery *et al.*, 2012). Between 1-3 days following testing, children with evidence of parasitic infection were treated with praziquantel (40 mg/kg) for schistosomiasis and albendazole (400 mg) for soil-transmitted helminths. Children were retested 25-27 days following treatment, and in cases of incomplete *S. mansoni* clearance, miracidia were sampled again, and treatment was re-administered.

Collection and ethical approval for the sampling of Ugandan *S. mansoni* cercariae

The FTA-preserved Ugandan *S. mansoni* cercariae were collected as part of snail collection surveys conducted between 2007 and 2010 as part of the EU-CONTRAST programme, a consortium of European and African researchers (CONTRAST EU/INCO.Dev contract No. 032203) in association with the Schistosomiasis Control Initiative (Adriko *et al.*, 2013) and provided via SCAN. The Ethical Review Board of the Uganda National Council of Science & Technology approved the surveys.

Collection and ethical approval for the sampling of Tanzanian *S. mansoni* cercariae

Tanzanian *S. mansoni* cercariae were originally collected as part of snail collection surveys undertaken in the Mwanza region and Geita regions of Northern Tanzania between January 2012 and December 2015 (2017) as part of the Schistosomiasis Consortium for Operational Research and Evaluation (SCORE) snail project (Ezeamama *et al.*, 2016) and archived within the Schistosomiasis Collection at the Natural History Museum (SCAN) (Emery et al., 2007). Survey sites were picked based on their proximity (within 5-15 metres) to schools and were identified based on local information about water activities (bathing, fishing, water collection) (Ezeamama *et al.*, 2016). *Biomphalaria* spp. snails were collected by scooping using handheld metal sieve scoops or dredging using a metal dredge dropped from a boat and dragged 10 m back to shore. Scooped/dredged snails were hand-collected with forceps and placed into collection jars. All snails were placed in 24-well ELISA plates under direct light for at least four hours to induce shedding, and cercariae from each shedding snail were individually collected in 3 µL of water using a Gilson pipette and fixed on Whatman FTA cards (Gower *et al.*, 2007; Emery *et al.*, 2012). COX-1 identification of *S. mansoni* was performed as described in Gouvras *et al. (2017)*.

Origin of the Senegalese *S. mansoni* laboratory (adult worm) sample

Both *S. mansoni* Senegalese isolates (FS0001 and FS0002) originated from approximately 30 *Biomphalaria pfeifferi* with patent *S. mansoni* infections found in the Western principal irrigation canal in the Ndiengue District of Richard Toll (Fallon *et al.*, 1995). The strain was originally sampled in 1993 and then maintained at the School of Biological Science, University of Wales (Bangor, United Kingdom) and subsequently in the helminGuard laboratory (Süelfeld, Germany) (Fallon *et al.*, 1997).

Origins of the *S. rodhaini* (adult worms and cercariae) samples

The adult *S. rodhaini* adult worm sample (RZ0001) was provided by SCAN from a laboratory-passaged *S. rodhaini* strain originally isolated from infected *Biomphalaria* snails

collected in Burundi in 2000. The sample used here represented the 6th passage through

laboratory *Biomphalaria glabrata* and *Mus musculus*, under the Home Office project license

numbers 70/4687 (before 2003) and 70/5935 (2003-2008) (Emery *et al.*, 2012).

The two *S. rodhaini* cercariae analysed here originated from the Mwanza region of Tanzania.

They were collected as part of the SCORE xenomonitoring project and provided by SCAN.

Species identification was confirmed by Sanger sequence analysis of a partial region of the

cox1 gene, the ITS1+2 rDNA region, and a partial region of the 18S rDNA region using the

methods described in Pennance *et al.*, 2020.

DNA extraction and sequencing

The DNA from individual miracidia and cercariae were isolated from the Whatman FTA cards

using methods described by (Doyle *et al.*, 2019; Moore *et al.*, 2020). A 2mm Harris micro-

punch was used to punch out the FTA disc containing the DNA. Within a well of a 96 PCR

plate, individual samples on individual punches were lysed in 30 µL of the following buffer:

30 mM Tris–HCl pH 8.0 (Sigma Aldrich), 0.5% Tween 20 (Sigma Aldrich), 0.5%

NP40/IGEPAL CA-630 (Sigma Aldrich) and 1.25 µg/mL of Protease reagent (Qiagen; cat

19155). Punches were incubated at 50°C for 1 h, then heated to 75°C for 30 mins. DNA was

extracted from adult worms using the Qiagen MagAttract HMW kit (PN-67653) following the

manufacturer's instructions. The DNA from the two individual Senegalese adult worms was

extracted using the Qiagen MagAttract HMW kit (PN-67653) following the manufacturer's

instructions.

Library preparation was performed using a low-input enzymatic fragmentation-based library

preparation method (Lee-Six *et al.*, 2019). For each sample, 20 µL of lysate (or extracted

DNA) was mixed with 50 µL of TE buffer (Ambion 10 mM Tris-HCL, 1 mM EDTA) and 50 µL

of Ampure XP beads followed by a 5 min binding reaction at room temperature. Magnetic

bead separation was used to separate genomic DNA, which was then washed twice with

75% ethanol. Beads were resuspended in 26 µL of TE buffer. To perform DNA

fragmentation and A-tailing, each sample was immediately mixed with 7 µL of 5x Ultra II FS

buffer and 2 µL of Ultra II FS enzyme and incubated on a thermal cycler for 12 min at 37°C

followed by 30 min at 65°C. Adaptor ligated libraries were prepared by adding 30 µL of

ligation mix, 1 µL ligation enhancer (New England BioLabs), 0.9 µL nuclease-free water

(Ambion) and 0.1 µL duplexed adapters to each well followed by incubation for 20 min at

20°C. Libraries were purified and eluted by adding 65 µL of Ampure XP beads and 65 µL of

TE buffer. Libraries were amplified by adding 25 µL KAPA HiFi HotStart ReadyMix (KAPA

Biosystems) and 1 µL PE1.0 primer to 21.5 µL of the library. Each sample was thermal-

cycled as follows: 98 °C for 5 min, then 12 cycles of 98°C for 30 sec, 65°C for 30 sec, 72°C

for 1 min and finally 72°C for 5 min. Ampure beads were used to purify amplified libraries

using a 0.7:1 volumetric ratio of beads to library. Each library was then eluted into 25 μL of

nuclease-free water. Library concentrations were adjusted to 2.4 nM and pooled, followed by

sequencing on the Illumina NovaSeq 6000 using 150 bp paired-end chemistry.

Variant discovery and annotation

In addition to the whole-genome sequencing of samples described above, sequence data for

407 *S. mansoni* accessions (Crellen, Allan, *et al.*, 2016; Berger *et al.*, 2021; Vianney *et al.*,

2022) and one *S. rodhaini* accession (Crellen, Allan, *et al.*, 2016) were included in this study.

Raw sequencing reads from all 682 accessions were trimmed using BBDuk

(https://jgi.doe.gov/data-and-tools/bbtools/bb-tools-user-guide/) to remove low-quality bases

and adapter sequences. Trimmed sequence reads were aligned to the *S. mansoni* (SM_V9,

WormBase ParaSite v16) (Buddenborg *et al.*, 2021) reference genome using BWA mem

(v.0.7.17) (Li, 2013). PCR duplicates were marked using PicardTools MarkDuplicates (as

part of GATK v.4.2.0.0) (McKenna *et al.*, 2010). Variant calling was performed per sample

using GATK HaplotypeCaller (v.4.2.0.0) in gVCF mode, retaining both variant and invariant

sites. Individual gVCFs were merged using GATK CombineGVCFs, and joint-call cohort

genotyping was performed using GATK GenotypeGVCFs. Variant sites with only single-

nucleotide polymorphisms (SNPs) were separated from indels and mixed sites (variant sites

that had both SNPs and indels called) using GATK SelectVariants. GATK VariantFiltration

was used to filter both groups independently. SNPs were retained if they met the following

criteria: QD $\geq$ 2.0, FS $\leq$ 60.0, MQ $\geq$ 40.0, MQRankSum $\geq$ −12.5,

ReadPosRankSum $\geq$ −8.0, SOR $\leq$ 3.0. Variant sites containing indels or mixed sites

were retained if they met the following criteria: QD $\geq$ 2.0, FS $\leq$ 200.0,

ReadPosRankSum $\geq$ −20.0, SOR $\leq$ 10.0.

VCFtools (v.0.1.15) was used to exclude accessions with a high rate of variant site

missingness (missing genotype called at >5% of sites) and subsequently used to remove

sites where >10% of accessions had a missing genotype (Danecek *et al.*, 2011). This

formed the primary VCF file used for almost all analyses. For analyses of nucleotide diversity

and fixation index (F_{ST}), we produced a second filtered VCF file. We used VCFtools to filter

both variant and invariant sites with >80% missing variants, enforced a minimum mean read

depth of 5, a maximum mean read depth of 500, removing variant sites found to be

significantly out of Hardy-Weinberg equilibrium ($p < 0.001$), and only retained SNPs.

Functional annotation of SNPs and indels in the primary VCF file was performed using

SnpEff (v.5.0e) (Cingolani *et al.*, 2012) with gene annotations (v.9) downloaded from

WormBase ParaSite (Howe *et al.*, 2017) V17. Repetitive elements in the *S. mansoni*

assembly were annotated using RepeatModeler and RepeatMasker (Smit, Hubley and

Green, 2015).

Depth of coverage

For each sample, the depth of read coverage was calculated in 2 kb windows along each

chromosome using bedtools coverage (v.2.30.0) (Quinlan and Hall, 2010).

Sample relatedness

We estimated pairwise relatedness between accessions from the same endemic regions.

We subsampled to only autosomal SNPs and ran NgsRelate (Hanghøj *et al.*, 2019) using

default parameters. As an additional line of evidence, we first used PLINK (v.2.0) (Chang *et*

al., 2015) to exclude SNPs found to be in strong linkage disequilibrium. The genome was

then scanned in sliding windows of 50 SNPs, increasing in steps of 10 SNPs and SNPs

within windows with squared correlation coefficients >0.2 were removed. Using this filtered

subset of SNPs, we performed pedigree reconstruction using Sequoia (Huisman, 2017).

Pairwise relationships were first classified based on the coefficient of kinship score (θ)

calculated by NgsRelate. NgsRelate was first run on all 570 *S. mansoni* samples to identify

relationships between datasets or populations. After confirming these relationships did not

occur, NgsRelate was rerun on individual populations from Southern Uganda, Northern

Tanzania and Koome Islands. Additionally, samples from Berger et al. (2021) were excluded

from these analyses due to both highly elevated inbreeding coefficients in this dataset.

Pairwise relationships with $\theta > 0.354$ were classified as monozygotic twins, $0.354 > \theta \geq$

0.177 were classified as first-degree relatives, $0.177 > \theta \geq 0.0884$ were classified as second-

degree relatives and $0.0884 > \theta \geq 0.0442$ were classified as third-degree relatives. In

addition, first-, second-, and third-degree relationships were only considered valid if the

maximum likelihood estimate of sharing 1 IBD allele (K_1) was greater than 0.05. Instances

where both NgsRelate and Sequoia identified the same first-degree relationships were

designated 'high-confidence' relationships. GGtern was used to plot all three maximum

likelihood estimates of sharing 0, 1 or 2 IBD alleles (K_0, K_1, and K_2), and Circos (Krzywinski

et al., 2009) and Gephi (Bastian, Heymann and Jacomy, 2009) were used to visualise

pairwise relationships between different donors.

Population genomic structure and diversity

We removed variants found at minor allele frequencies < 0.05 and excluded all variants

found on the Z chromosome, W chromosome and mitochondrial genome. We then removed

all variants found within repetitive regions identified using RepeatMasker and finally removed

variants found to be in strong linkage disequilibrium, as above. Principal component analysis

was performed with the remaining 188,923 autosomal SNPs using PLINK. Admixture

analyses were performed using ADMIXTURE (Alexander, Novembre and Lange, 2009) with

K values (number of hypothetical ancestral populations) ranging from 1 to 20, 10-fold cross-

validation and standard error estimation with 250 bootstraps. The lowest cross-validation

error (CV) value was found for K$\square$=$\square$5 (Supplementary Figure 13).

We used publicly-available scripts to convert all 188,923 autosomal SNPs into Phylip format

(https://github.com/edgardomortiz/vcf2phylip/vcf2phylip.py) and remove invariant sites

(https://github.com/btmartin721/raxml_ascbias/ascbias.py). Phylogenomic inference was

performed using IQ-TREE (Minh *et al.*, 2020), using the best-fit substitution model with

ascertainment bias correction selected by ModelFinder (GTR+F+ASC+R10) and 1000

ultrafast bootstraps. The resulting phylogeny was visualised using ggtree (Yu *et al.*, 2017).

pixy (v.1.2.3.beta1) (Korunes and Samuk, 2021) was used to calculate autosomal nucleotide

diversity (π) and the fixation index (F_{ST}) in 5 kb non-overlapping, sliding windows for each

population using the secondary (mixed variant and invariant sites) VCF. Negative F_{ST} values

were corrected to 0 before calculating genome-wide median values. Watterson's estimator

(Θ) was calculated using Scikit-allel (Miles *et al.*, 2020). Effective population size (N_e) was

using a per-generation mutation rate (μ) of 8.1x10^{-9} (Crellen, Allan, *et al.*, 2016) and the

following equation:

$$Ne = \Theta/4\mu$$

The coefficient of inbreeding (F) was calculated using VCFtools to assess per-sample

homozygosity. Comparisons of samples within and between study sites can be found in

Supplementary Figure 14a-c.

The impact of praziquantel treatment on population genomic diversity

pixy (v.1.2.3.beta1) was used to calculate autosomal nucleotide diversity (π) in 5 kb non-

overlapping, sliding windows for each pre- and post-treatment population using the

secondary (mixed variant and invariant sites) VCF and including related accessions

(Korunes and Samuk, 2021).

Structural variant genotyping and annotation

Structural variants for each sample were detected using LUMPY (v.0.2.13) and genotyped

using SVTyper (v.0.7.0) as implemented in the Smoove (v.0.2.7) pipeline

(https://github.com/brentp/smoove) (Layer *et al.*, 2014; Chiang *et al.*, 2015). Using this

pipeline, we first performed single-sample calling using the BWA-aligned reads to the *S.*
mansoni reference genome. This was followed by merging called variants and sample-wise
re-genotyping across all 570 *S. mansoni* accessions. Structural variants intersecting coding
regions were annotated using the reference GFF file. We then filtered variants based on the
Smoove author's recommendations, where heterozygous calls with MSHQ>=3, deletions
with DHFFC>=0.7 and duplications with DHFFC<=1.25 were excluded. Due to inconsistent
coverage and whole-genome amplification of the original libraries, all accessions from
Berger et al. (2021) were excluded from structural variant analyses after genotyping.
744
Inference of demographic history
We ran SMC++ (v.1.15.2) (Terhorst, Kamm and Song, 2017) on each autosome using a per-
generation mutation rate of $8.1x10^{-9}$ and a generation time of 85 days. For Southern
Ugandan, Koome Island, Northern Tanzanian, and Eastern Ugandan populations, we
randomly subset them down to $n = 12$ unrelated accessions, providing replicates where
populations had more than 24 accessions. For all other populations, only single accessions
were analysed.
752
Screening for recent admixture
Analysis of passaged laboratory strains also identified a single probable *S. mansoni-S.*
rodhaini hybrid (RZ0001), which displayed high heterozygosity, intermediate admixture
proportions and intermediate phylogenetic positioning, consistent with an early-generation
(F_1 - F_3) hybrid, this sample was excluded from further analyses (Supplementary Figure 15;
Supplementary Data 2) (Platt *et al.*, 2019; Berger *et al.*, 2022).
759
Analysis of candidate praziquantel resistance genes
We inspected SnpEff annotated variants within the candidate praziquantel resistance gene
(*Smp_24790*), focusing on isoform 5 (*Smp_246790.5* in the v9 annotation), the only full-
length isoform containing the predicted binding site. Frequencies of functionally impactful
mutations were calculated using only Lake Victoria accessions. The maximum-likelihood
phylogeny produced previously was annotated to highlight accessions containing specifically
functionally impactful mutations. We excluded a single indel, p.Met1fs, which was a
frameshift variant and a short repeat from the analysis; for completeness, information on this
variant is still reported in Supplementary Data 7,9,10.
769
Ca^{2+} reporter assays
Ca^{2+} imaging assays were performed using a Fluorescence Imaging Plate Reader
(FLIPR^{TETRA}, Molecular Devices). HEK293 cells (naïve or transfected with specific TRPM$_{PZQ}$

variants) were seeded (20,000 cells/well) in a black-walled clear-bottomed poly-d-lysine

coated 384-well plate (Corning) in DMEM growth media supplemented with 10% FBS. This

medium was removed after 24hrs, and the cells were loaded with a fluorescent Ca^{2+}

indicator (Fluo-4 NW dye, Invitrogen) by incubation (20 µL per well, 1 h at 37°C) in Hanks'

balanced salt solution (HBSS) assay buffer containing probenecid (2.5 mM) and HEPES

(20mM). Dilutions of praziquantel (Sigma) were prepared in this same buffer, but without

probenecid and dye, in flat shape 384-well plates (Greiner Bio-one, Germany). The Ca^{2+}

reporter assay was performed at room temperature by monitoring fluorescence (raw

fluorescence units) before (basal, 20s) and after the addition of praziquantel (an additional

250s). For quantitative analyses, peak fluorescence in each well was normalised to the

maximal response of the reference channel sequence and concentration-response curves

were plotted using the sigmoidal dose-response function in Origin.

Author contributions

DJB conceived and designed the study with input from JAC, MR, AE, FA, SB, MB, JSM and JPW. TC, NBK, EMT, PHLL and JPW planned and coordinated the collection of samples from Bugoto and Bukagabo Beach schools, which was carried out by TC, PHLL and MA. SCORE Tanzanian samples were collected by AG, BLW, FA, HN, JI, MR, RA, SK, and JPW in partnership with local technicians. CONTRAST samples were collected by CS, JRS, BW, and JPW in partnership with local technicians. JTV, RS and AE provided information on the Koome Island samples. MR, AE and FA curated the SCAN collection and coordinated sample delivery. HH coordinated the isolation and delivery of the two passaged Senegalese laboratory isolates, and NH performed DNA extraction and coordinated sequencing for those samples. For all other samples, DJB extracted and amplified DNA, and planned and coordinated the sequencing. JSM and SKP performed Ca^{2+} reporter assays and modelling analysis. DJB analysed all other data and wrote the paper with input from JSM, SRD and JPW. S-KP, TC, JAC, MR, AE, BLW, PHL, SB and MB provided comments on the text, and all authors viewed and approved the final manuscript.

Competing interests

The authors declare no competing interests.

Data availability

The sequencing data generated in this study have been deposited in the European Nucleotide Archive (ENA) repository under project accession codes PRJEB42451 and PRJEB29904. Individual sample accessions are listed in Supplementary Data 14. The genome assembly and annotation files are available through ENA (PRJEA36577; GCA_000237925).

Code availability

The code used for data analysis is available at https://github.com/duncanberger/SM_VARIATION.

Acknowledgements

Funding was via core funding of the Wellcome Sanger Institute (Wellcome grant 206194; PI: MB). Ugandan sampling was supported within the remit of Schistosomiasis Control Initiative (SCI) monitoring and evaluation activities funded through the Bill and Melinda Gates Foundation; PI: JPW, a European Commission 'Specific Research Project grant' CONTRAST (FP6 STREP contract no: 032203, http://www.eu-contrast.eu; PI: JPW), and a Wellcome Trust grant (no: 095778; PI: AE). Northern Tanzania sample collection was funded

by the Schistosomiasis Consortium for Operational Research (SCORE) from the University

of Georgia Research Foundation Inc., funded by the Bill & Melinda Gates Foundation (Sub-

awards RR37–053/4787466 & RR374-053/4785426; PI: JPW). FA, AE, MR and the

Schistosome Collection at the Natural History Museum (SCAN: AE) were supported by

Wellcome grant 1045958/Z/13/Z (PI: AE). JAC, MB and JPW were supported by a European

& Developing Countries Clinical Trials Partnership (EDCTP) grant FibroScHot (Ref

RIA2017NIM-1842, PIs JAC, MB & JPW). JSM and SKP were supported by the National

Institutes of Health (R01-AI145871, PI: JSM). SRD is supported by a UKRI Future Leaders

Fellowship (MR/T020733/1). For the purpose of Open Access, the authors have applied a

CC BY public copyright Licence to any Author Accepted Manuscript version arising from this

submission.

We would also like to thank all the technicians, nurses and drivers who supported

sample collections, including (but not exclusive to) Moses Arinaitwe, Aida Wamboko, Annet

Enzaru, Andrina Nankasi, Aaron Atuhaire and Fiddi Rugigana from the Vector Control

Division, Ugandan Ministry of Health, Uganda, and John Igogote, Honest Nagai, and

Revocatus Alphonce from the National Institute for Medical Research, Tanzania. We would

also like to thank the teachers from the schools we visited for their assistance and

cooperation, the officials in the Health and Education Departments in districts of Mayuge

(Uganda), and Alan Fenwick and Wendy Harrison of the Schistosomiasis Control Initiative

(renamed Unlimit Health) for their help organising and funding the Ugandan fieldwork.

Finally, we thank all the children recruited into this study and their parents and teachers for

their cooperation.

References

Adriko, M. *et al.* (2013) 'Compatibility of Ugandan *Schistosoma mansoni* isolates with
Biomphalaria snail species from Lake Albert and Lake Victoria', *Acta Tropica*, 128(2), pp.
303–308.

Aemero, M. *et al.* (2015) 'Genetic diversity, multiplicity of infection and population structure
of *Schistosoma mansoni* isolates from human hosts in Ethiopia', *BMC Genetics*, 16, p. 137.

Agola, L.E. *et al.* (2009) 'Genetic diversity and population structure of *Schistosoma mansoni*
within human infrapopulations in Mwea, central Kenya assessed by microsatellite markers',
Acta Tropica, 111(3), pp. 219–225.

Alexander, D.H., Novembre, J. and Lange, K. (2009) 'Fast model-based estimation of
ancestry in unrelated individuals', *Genome Research*, 19(9), pp. 1655–1664.

Bastian, M., Heymann, S. and Jacomy, M. (2009) 'Gephi: an open source software for

exploring and manipulating networks', *Proceedings of the International AAAI Conference on*
Web and Social Media, 3(1), pp. 361–362.

Berger, D.J. *et al.* (2021) 'Whole-genome sequencing of *Schistosoma mansoni* reveals
extensive diversity with limited selection despite mass drug administration', *Nature*
Communications, 12(1), p. 4776.

Berger, D.J. *et al.* (2022) 'Genomic evidence of contemporary hybridization between
Schistosoma species', *PLoS Pathogens*, 18(8), p. e1010706.

Blanton, R.E. *et al.* (2011) '*Schistosoma mansoni* population structure and persistence after
praziquantel treatment in two villages of Bahia, Brazil', *International Journal for Parasitology*,
41(10), pp. 1093–1099.

Bronzan, R.N. *et al.* (2018) 'Impact of community-based integrated mass drug administration
on schistosomiasis and soil-transmitted helminth prevalence in Togo', *PLoS Neglected*
Tropical Diseases, 12(8), p. e0006551.

Buddenborg, S.K. *et al.* (2021) 'Assembled chromosomes of the blood fluke *Schistosoma*
mansoni provide insight into the evolution of its ZW sex-determination system', *bioRxiv*
[Preprint]. Available at: https://www.biorxiv.org/content/10.1101/2021.08.13.456314.abstract.

Chang, C.C. *et al.* (2015) 'Second-generation PLINK: rising to the challenge of larger and
richer datasets', *GigaScience*, 4, p. 7.

Chevalier, F.D. *et al.* (2016) 'Independent origins of loss-of-function mutations conferring
oxamniquine resistance in a Brazilian schistosome population', *International Journal for*
Parasitology, 46(7), pp. 417–424.

Chevalier, F.D. *et al.* (2019) 'Oxamniquine resistance alleles are widespread in Old World
Schistosoma mansoni and predate drug deployment', *PLoS Pathogens*, 15(10), p.
e1007881.

Chevalier, F.D. *et al.* (2024) 'A single locus determines praziquantel response in
Schistosoma mansoni', *Antimicrobial Agents and Chemotherapy*, 68(3), p. e0143223.

Chiang, C. *et al.* (2015) 'SpeedSeq: ultra-fast personal genome analysis and interpretation',
Nature Methods, 12(10), pp. 966–968.

Cingolani, P. *et al.* (2012) 'A program for annotating and predicting the effects of single
nucleotide polymorphisms, SnpEff', *Fly*, 6(2), pp. 80–92.

Colley, D.G. *et al.* (2014) 'Human schistosomiasis', *The Lancet*, 383(9936), pp. 2253–2264.

Control of Neglected Tropical Diseases (NTD) Guidelines Review Committee (2022) *WHO*
Guideline on control and elimination of human schistosomiasis. World Health Organization.

Couto, F.F.B. *et al.* (2011) '*Schistosoma mansoni*: a method for inducing resistance to
praziquantel using infected *Biomphalaria glabrata* snails', *Memorias do Instituto Oswaldo*
Cruz, 106(2), pp. 153–157.

Crellen, T., Walker, M., *et al.* (2016) 'Reduced Efficacy of Praziquantel Against *Schistosoma*
mansoni Is Associated with Multiple Rounds of Mass Drug Administration', *Clinical infectious*
diseases: an official publication of the Infectious Diseases Society of America, 63(9), pp.
1151–1159.

Crellen, T., Allan, F., *et al.* (2016) 'Whole genome resequencing of the human parasite

Schistosoma mansoni reveals population history and effects of selection', *Scientific Reports*,
6, pp. 1–13.

Danecek, P. *et al.* (2011) 'The variant call format and VCFtools', *Bioinformatics* , 27(15), pp.
2156–2158.

Deol, A.K. *et al.* (2019) 'Schistosomiasis - assessing progress toward the 2020 and 2025
global goals', *The New England Journal of Medicine*, 381(26), pp. 2519–2528.

Doyle, S.R. *et al.* (2019) 'Evaluation of DNA Extraction Methods on Individual Helminth Egg
and Larval Stages for Whole-Genome Sequencing', *Frontiers in Genetics*, 10, p. 826.

Doyle, S.R. *et al.* (2022) 'Genomic landscape of drug response reveals mediators of
anthelmintic resistance', *Cell Reports*, 41(3), p. 111522.

Emery, A.M. *et al.* (2012) 'Schistosomiasis collection at NHM (SCAN)', *Parasites & Vectors*,
5(1), p. 1.

Ezeamama, A.E. *et al.* (2016) 'Gaining and sustaining schistosomiasis control: study
protocol and baseline data prior to different treatment strategies in five African countries',
BMC Infectious Diseases, 16(1). Available at: https://doi.org/10.1186/s12879-016-1575-2.

Fallon, P.G. *et al.* (1995) 'Short Report: Diminished Susceptibility to Praziquantel in a
Senegal Isolate of *Schistosoma mansoni*', *The American Journal of Tropical Medicine and*
Hygiene, 53(1), pp. 61–62.

Fallon, P.G. *et al.* (1997) '*Schistosoma mansoni*: maturation rate and drug susceptibility of
different geographic isolates', *Experimental Parasitology*, 86(1), pp. 29–36.

Fallon, P.G. and Doenhoff, M.J. (1994) 'Drug-resistant schistosomiasis: resistance to
praziquantel and oxamniquine induced in *Schistosoma mansoni* in mice is drug specific',
The American Journal of Tropical Medicine and Hygiene, 51(1), pp. 83–88.

Faust, C.L. *et al.* (2019) 'Two-year longitudinal survey reveals high genetic diversity of
Schistosoma mansoni with adult worms surviving praziquantel treatment at the start of mass
drug administration in Uganda', *Parasites & Vectors*, 12(1), p. 607.

Fukushige, M. *et al.* (2021) 'Efficacy of praziquantel has been maintained over four decades
(from 1977 to 2018): A systematic review and meta-analysis of factors influence its efficacy',
PLoS Neglected Tropical Diseases, 15(3), p. e0009189.

Gouvras, A.N. *et al.* (2017) 'Longitudinal survey on the distribution of *Biomphalaria sudanica*
and *B. choanomophala* in Mwanza region, on the shores of Lake Victoria, Tanzania:
implications for schistosomiasis transmission and control', *Parasites & Vectors*, 10(316).

Gower, C.M. *et al.* (2007) 'Development and application of an ethically and epidemiologically
advantageous assay for the multi-locus microsatellite analysis of *Schistosoma mansoni*',
Parasitology, 134(Pt 4), pp. 523–536.

Gryseels, B. and De Vlas, S.J. (1996) 'Worm burdens in schistosome infections',
Parasitology Today , 12(3), pp. 115–119.

Hanghøj, K. *et al.* (2019) 'Fast and accurate relatedness estimation from high-throughput
sequencing data in the presence of inbreeding', *GigaScience*, 8(5), p. giz034.

Hodgkinson, J.E. *et al.* (2019) 'Refugia and anthelmintic resistance: Concepts and
challenges', *International Journal for Parasitology: Drugs and Drug Resistance*, 10, pp. 51–

57.

Howe, K.L. *et al.* (2017) 'WormBase ParaSite - a comprehensive resource for helminth
genomics', *Molecular and Biochemical Parasitology*, 215, pp. 2–10.

Huisman, J. (2017) 'Pedigree reconstruction from SNP data: parentage assignment, sibship
clustering and beyond', *Molecular Ecology Resources*, 17(5), pp. 1009–1024.

Huyse, T. *et al.* (2013) 'Regular treatments of praziquantel do not impact on the genetic
make-up of *Schistosoma mansoni* in Northern Senegal', *Infection, Genetics and Evolution*,
18, pp. 100–105.

Jacquard, A. (1966) 'Logique du calcul des coefficients d'identité entre deux individus',
Population, 21(4), pp. 751–776.

Kabatereine, N.B. *et al.* (2007) 'Impact of a national helminth control programme on infection
and morbidity in Ugandan schoolchildren', *Bulletin of the World Health Organization*, 85(2),
pp. 91–99.

Kaplan, R.M. and Vidyashankar, A.N. (2012) 'An inconvenient truth: global worming and
anthelmintic resistance', *Veterinary Parasitology*, 186(1-2), pp. 70–78.

Katz, N., Chaves, A. and Pellegrino, J. (1972) 'A simple device for quantitative stool thick-
smear technique in *Schistosoma mansoni*', *Revista do Instituto de Medicina Tropical de Sao
Paulo*, 14(6), pp. 397–400.

Kenyon, F. *et al.* (2009) 'The role of targeted selective treatments in the development of
refugia-based approaches to the control of gastrointestinal nematodes of small ruminants',
Veterinary Parasitology, 164(1), pp. 3–11.

Kittur, N. *et al.* (2019) 'Persistent hotspots in schistosomiasis consortium for operational
research and evaluation studies for gaining and sustaining control of schistosomiasis after
four years of mass drug administration of praziquantel', *The American Journal of Tropical
Medicine and Hygiene*, 101(3), pp. 617–627.

Kokaliaris, C. *et al.* (2022) 'Effect of preventive chemotherapy with praziquantel on
schistosomiasis among school-aged children in sub-Saharan Africa: a spatiotemporal
modelling study', *The Lancet Infectious Diseases*, 22(1), pp. 136–149.

Korunes, K.L. and Samuk, K. (2021) 'pixy: Unbiased estimation of nucleotide diversity and
divergence in the presence of missing data', *Molecular Ecology Resources*, 21(4), pp. 1359–
1368.

Krzywinski, M. *et al.* (2009) 'Circos: an information aesthetic for comparative genomics',
Genome Research, 19(9), pp. 1639–1645.

Lamberton, P.H.L., Faust, C.L. and Webster, J.P. (2017) 'Praziquantel decreases fecundity
in *Schistosoma mansoni* adult worms that survive treatment: evidence from a laboratory life-
history trade-offs selection study', *Infectious Diseases of Poverty*, 6(1), p. 110.

Layer, R.M. *et al.* (2014) 'LUMPY: a probabilistic framework for structural variant discovery',
Genome Biology, 15(6), p. R84.

Leathwick, D.M., Ganesh, S. and Waghorn, T.S. (2015) 'Evidence for reversion towards
anthelmintic susceptibility in *Teladorsagia circumcincta* in response to resistance
management programmes', *International Journal for Parasitology: Drugs and Drug
Resistance*, 5(1), pp. 9–15.

Le Clec'h, W. *et al.* (2021) 'Genetic analysis of praziquantel response in schistosome
parasites implicates a transient receptor potential channel', *Science Translational Medicine*,
13(625), p. eabj9114.

Lee-Six, H. *et al.* (2019) 'The landscape of somatic mutation in normal colorectal epithelial
cells', *Nature*, 574, pp. 532–537.

Lelo, A.E. *et al.* (2014) 'No apparent reduction in schistosome burden or genetic diversity
following four years of school-based mass drug administration in Mwea, central Kenya, a
heavy transmission area', *PLoS Neglected Tropical Diseases*, 8(10), p. e3221.

Li, H. (2013) 'Aligning sequence reads, clone sequences and assembly contigs with BWA-
MEM', *arXiv [q-bio.GN]*. Available at: http://arxiv.org/abs/1303.3997.

Lim, R.M. *et al.* (2023) 'Defining schistosomiasis hotspots based on literature and
shareholder interviews', *Trends in Parasitology*, 39(12), pp. 1032–1049.

Lo, N.C. *et al.* (2022) 'Review of 2022 WHO guidelines on the control and elimination of
schistosomiasis', *The Lancet Infectious Diseases*, 22(11), pp. e327–e335.

Marchant, J.S. (2024) 'Progress interrogating TRPMPZQ as the target of praziquantel', *PLoS
Neglected Tropical Diseases*, 18(2), p. e0011929.

Mari, L. *et al.* (2017) 'Heterogeneity in schistosomiasis transmission dynamics', *Journal of
Theoretical Biology*, 432, pp. 87–99.

Mawa, P.A. *et al.* (2021) 'Schistosomiasis morbidity hotspots: roles of the human host, the
parasite and their interface in the development of severe morbidity', *Frontiers in
Immunology*, 12, p. 751.

McKenna, A. *et al.* (2010) 'The Genome Analysis Toolkit: a MapReduce framework for
analyzing next-generation DNA sequencing data', *Genome Research*, 20(9), pp. 1297–1303.

Mei, Z.-Z. *et al.* (2006) 'Intracellular coiled-coil domain engaged in subunit interaction and
assembly of melastatin-related transient receptor potential channel 2', *The Journal of
Biological Chemistry*, 281(50), pp. 38748–38756.

Melman, S.D. *et al.* (2009) 'Reduced Susceptibility to Praziquantel among Naturally
Occurring Kenyan Isolates of *Schistosoma mansoni*', *PLoS Neglected Tropical Diseases*,
3(8), p. e504.

Meurs, L. *et al.* (2013) 'Micro-geographical heterogeneity in *Schistosoma mansoni* and *S.
haematobium* infection and morbidity in a co-endemic community in northern Senegal', *PLoS
Neglected Tropical Diseases*, 7(12), p. e2608.

Miles, A. *et al.* (2020) *cggh/scikit-allel: v1.3.1*. Available at:
https://doi.org/10.5281/zenodo.3935797.

Minh, B.Q. *et al.* (2020) 'IQ-TREE 2: New Models and Efficient Methods for Phylogenetic
Inference in the Genomic Era', *Molecular Biology and Evolution*, 37(5), pp. 1530–1534.

Moore, L. *et al.* (2020) 'The mutational landscape of normal human endometrial epithelium',
Nature, 580(7805), pp. 640–646.

Mutuku, M.W. *et al.* (2019) 'A search for snail-related answers to explain differences in
response of *Schistosoma mansoni* to praziquantel treatment among responding and
persistent hotspot villages along the Kenyan shore of Lake Victoria', *The American Journal*

of Tropical Medicine and Hygiene, 101(1), pp. 65–77.

Mwangi, I.N. *et al.* (2014) 'Praziquantel sensitivity of Kenyan *Schistosoma mansoni* isolates
and the generation of a laboratory strain with reduced susceptibility to the drug', *International*
Journal for Parasitology: Drugs and Drug Resistance, 4(3), pp. 296–300.

Neves, M.I. *et al.* (2021) 'Revisiting density-dependent fecundity in schistosomes using
sibship reconstruction', *PLoS Neglected Tropical Diseases*, 15(5), p. e0009396.

Neves, M.I., Webster, J.P. and Walker, M. (2019) 'Estimating helminth burdens using sibship
reconstruction', *Parasites & Vectors*, 12(1), p. 441.

Park, S.-K. *et al.* (2019) 'The anthelmintic drug praziquantel activates a schistosome
transient receptor potential channel', *The Journal of Biological Chemistry*, 294(49).

Park, S.-K. *et al.* (2021) 'Mechanism of praziquantel action at a parasitic flatworm ion
channel', *Science Translational Medicine*, 13(625), p. eabj5832.

Pitchford, R.J. and Visser, P.S. (1975) 'A simple and rapid technique for quantitative
estimation of helminth eggs in human and animal excreta with special reference to
Schistosoma sp.', *Transactions of the Royal Society of Tropical Medicine and Hygiene*,
69(3), pp. 318–322.

Platt, R.N. *et al.* (2019) 'Ancient Hybridization and Adaptive Introgression of an Invadolysin
Gene in Schistosome Parasites', *Molecular Biology and Evolution*, 36(10), pp. 2127–2142.

Platt, R.N. *et al.* (2021) 'Genomic analysis of a parasite invasion: colonization of the
Americas by the blood fluke, *Schistosoma mansoni*', *Molecular Ecology*, 31(8), pp. 2242–
2263.

Quinlan, A.R. and Hall, I.M. (2010) 'BEDTools: a flexible suite of utilities for comparing
genomic features', *Bioinformatics* , 26(6), pp. 841–842.

Rose, H. *et al.* (2015) 'Widespread anthelmintic resistance in European farmed ruminants: a
systematic review', *The Veterinary Record*, 176(21), p. 546.

Schwab, A.E. *et al.* (2005) 'Detection of benzimidazole resistance-associated mutations in
the filarial nematode *Wuchereria bancrofti* and evidence for selection by albendazole and
ivermectin combination treatment', *The American Journal of Tropical Medicine and Hygiene*,
73(2), pp. 234–238.

Smit, A.F.A., Hubley, R. and Green, P. (2015) *RepeatMasker Open-4.0. 2013-2015*.
Available at: http://www.repeatmasker.org.

Stelma, F.F. *et al.* (1995) 'Efficacy and side effects of praziquantel in an epidemic focus of
Schistosoma mansoni', *The American Journal of Tropical Medicine and Hygiene*, 53(2), pp.
167–170.

Summers, S. *et al.* (2022) 'A review of the genetic determinants of praziquantel resistance in
Schistosoma mansoni: Is praziquantel and intestinal schistosomiasis a perfect match?',
Frontiers in Tropical Diseases, 3.

Terhorst, J., Kamm, J.A. and Song, Y.S. (2017) 'Robust and scalable inference of population
history from hundreds of unphased whole genomes', *Nature Genetics*, 49(2), pp. 303–309.

Vianney, T.J. *et al.* (2022) 'Genome-wide analysis of *Schistosoma mansoni* reveals limited
population structure and possible praziquantel drug selection pressure within Ugandan hot-

spot communities', *PLoS Neglected Tropical Diseases*, 16(8), p. e0010188.

Webster, J.P. *et al.* (2014) 'The contribution of mass drug administration to global health:
past, present and future', *Philosophical transactions of the Royal Society of London. Series*
B, Biological sciences, 369(1645), p. 20130434.

WHO Expert Committee on the Control of Schistosomiasis (2002) *Prevention and control of*
schistosomiasis and soil-transmitted helminthiasis : report of a WHO expert committee.
World Health Organ Tech Rep Ser. Available at: http://apps.who.int/iris/handle/10665/42588
(Accessed: 13 November 2018).

World Health Organization (2020) *A road map for neglected tropical diseases 2021–2030.*
World Health Organization. Available at: https://www.who.int/neglected_diseases/Ending-
the-neglect-to-attain-the-SDGs--NTD-Roadmap.pdf.

World Health Organization (2021) 'Schistosomiasis and soil-transmitted helminthiases:
progress report, 2020 - Schistosomiase et géohelminthiases: rapport de situation, 2020',
Weekly Epidemiological Record, 96(48), pp. 585–595.

Yin, Y. *et al.* (2019) 'Structural basis of cooling agent and lipid sensing by the cold-activated
TRPM8 channel', *Science*, 363(6430), p. eaav9334.

Yu, G. *et al.* (2017) 'ggtree : an r package for visualization and annotation of phylogenetic
trees with their covariates and other associated data', *Methods in Ecology and Evolution*,
8(1), pp. 28–36.

Adriko, M. et al. (2013) 'Compatibility of Ugandan Schistosoma mansoni isolates with
Biomphalaria snail species from Lake Albert and Lake Victoria', Acta Tropica, 128(2), pp.
303–308.

Aemero, M. et al. (2015) 'Genetic diversity, multiplicity of infection and population structure
of Schistosoma mansoni isolates from human hosts in Ethiopia', BMC Genetics, 16, p. 137.

Agola, L.E. et al. (2009) 'Genetic diversity and population structure of Schistosoma mansoni
within human infrapopulations in Mwea, central Kenya assessed by microsatellite markers',
Acta Tropica, 111(3), pp. 219–225.

Alexander, D.H., Novembre, J. and Lange, K. (2009) 'Fast model-based estimation of
ancestry in unrelated individuals', Genome Research, 19(9), pp. 1655–1664.

Bastian, M., Heymann, S. and Jacomy, M. (2009) 'Gephi: an open source software for
exploring and manipulating networks', Proceedings of the International AAAI Conference on
Web and Social Media, 3(1), pp. 361–362.

Berger, D.J. et al. (2021) 'Whole-genome sequencing of Schistosoma mansoni reveals
extensive diversity with limited selection despite mass drug administration', Nature
Communications, 12(1), p. 4776.

Berger, D.J. et al. (2022) 'Genomic evidence of contemporary hybridization between
Schistosoma species', PLoS Pathogens, 18(8), p. e1010706.

Blanton, R.E. et al. (2011) 'Schistosoma mansoni population structure and persistence after
praziquantel treatment in two villages of Bahia, Brazil', International Journal for Parasitology,
41(10), pp. 1093–1099.

Bronzan, R.N. et al. (2018) 'Impact of community-based integrated mass drug administration
on schistosomiasis and soil-transmitted helminth prevalence in Togo', PLoS Neglected

Tropical Diseases, 12(8), p. e0006551.

Buddenborg, S.K. et al. (2021) 'Assembled chromosomes of the blood fluke Schistosoma
mansoni provide insight into the evolution of its ZW sex-determination system', bioRxiv
[Preprint]. Available at: https://www.biorxiv.org/content/10.1101/2021.08.13.456314.abstract.

Chang, C.C. et al. (2015) 'Second-generation PLINK: rising to the challenge of larger and
richer datasets', GigaScience, 4, p. 7.

Chevalier, F.D. et al. (2016) 'Independent origins of loss-of-function mutations conferring
oxamniquine resistance in a Brazilian schistosome population', International Journal for
Parasitology, 46(7), pp. 417–424.

Chevalier, F.D. et al. (2019) 'Oxamniquine resistance alleles are widespread in Old World
Schistosoma mansoni and predate drug deployment', PLoS Pathogens, 15(10), p.
e1007881.

Chevalier, F.D. et al. (2024) 'A single locus determines praziquantel response in
Schistosoma mansoni', Antimicrobial Agents and Chemotherapy, 68(3), p. e0143223.

Chiang, C. et al. (2015) 'SpeedSeq: ultra-fast personal genome analysis and interpretation',
Nature Methods, 12(10), pp. 966–968.

Cingolani, P. et al. (2012) 'A program for annotating and predicting the effects of single
nucleotide polymorphisms, SnpEff', Fly, 6(2), pp. 80–92.

Colley, D.G. et al. (2014) 'Human schistosomiasis', The Lancet, 383(9936), pp. 2253–2264.

Control of Neglected Tropical Diseases (NTD) Guidelines Review Committee (2022) WHO
Guideline on control and elimination of human schistosomiasis. World Health Organization.

Couto, F.F.B. et al. (2011) 'Schistosoma mansoni: a method for inducing resistance to
praziquantel using infected Biomphalaria glabrata snails', Memorias do Instituto Oswaldo
Cruz, 106(2), pp. 153–157.

Crellen, T., Walker, M., et al. (2016) 'Reduced Efficacy of Praziquantel Against Schistosoma
mansoni Is Associated with Multiple Rounds of Mass Drug Administration', Clinical infectious
diseases: an official publication of the Infectious Diseases Society of America, 63(9), pp.
1151–1159.

Crellen, T., Allan, F., et al. (2016) 'Whole genome resequencing of the human parasite
Schistosoma mansoni reveals population history and effects of selection', Scientific Reports,
6, pp. 1–13.

Danecek, P. et al. (2011) 'The variant call format and VCFtools', Bioinformatics , 27(15), pp.
2156–2158.

Deol, A.K. et al. (2019) 'Schistosomiasis - assessing progress toward the 2020 and 2025
global goals', The New England Journal of Medicine, 381(26), pp. 2519–2528.

Doyle, S.R. et al. (2019) 'Evaluation of DNA Extraction Methods on Individual Helminth Egg
and Larval Stages for Whole-Genome Sequencing', Frontiers in Genetics, 10, p. 826.

Doyle, S.R. et al. (2022) 'Genomic landscape of drug response reveals mediators of
anthelmintic resistance', Cell Reports, 41(3), p. 111522.

Emery, A.M. et al. (2012) 'Schistosomiasis collection at NHM (SCAN)', Parasites & Vectors,

5(1), p. 1.

Ezeamama, A.E. et al. (2016) 'Gaining and sustaining schistosomiasis control: study
protocol and baseline data prior to different treatment strategies in five African countries',
BMC Infectious Diseases, 16(1). Available at: https://doi.org/10.1186/s12879-016-1575-2.

Fallon, P.G. et al. (1995) 'Short Report: Diminished Susceptibility to Praziquantel in a
Senegal Isolate of Schistosoma mansoni', The American Journal of Tropical Medicine and
Hygiene, 53(1), pp. 61–62.

Fallon, P.G. et al. (1997) 'Schistosoma mansoni: maturation rate and drug susceptibility of
different geographic isolates', Experimental Parasitology, 86(1), pp. 29–36.

Fallon, P.G. and Doenhoff, M.J. (1994) 'Drug-resistant schistosomiasis: resistance to
praziquantel and oxamniquine induced in Schistosoma mansoni in mice is drug specific',
The American Journal of Tropical Medicine and Hygiene, 51(1), pp. 83–88.

Faust, C.L. et al. (2019) 'Two-year longitudinal survey reveals high genetic diversity of
Schistosoma mansoni with adult worms surviving praziquantel treatment at the start of mass
drug administration in Uganda', Parasites & Vectors, 12(1), p. 607.

Fukushige, M. et al. (2021) 'Efficacy of praziquantel has been maintained over four decades
(from 1977 to 2018): A systematic review and meta-analysis of factors influence its efficacy',
PLoS Neglected Tropical Diseases, 15(3), p. e0009189.

Gouvras, A.N. et al. (2017) 'Longitudinal survey on the distribution of Biomphalaria sudanica
and B. choanomophala in Mwanza region, on the shores of Lake Victoria, Tanzania:
implications for schistosomiasis transmission and control', Parasites & Vectors, 10(316).

Gower, C.M. et al. (2007) 'Development and application of an ethically and epidemiologically
advantageous assay for the multi-locus microsatellite analysis of Schistosoma mansoni',
Parasitology, 134(Pt 4), pp. 523–536.

Gryseels, B. and De Vlas, S.J. (1996) 'Worm burdens in schistosome infections',
Parasitology Today , 12(3), pp. 115–119.

Hanghøj, K. et al. (2019) 'Fast and accurate relatedness estimation from high-throughput
sequencing data in the presence of inbreeding', GigaScience, 8(5), p. giz034.

Hodgkinson, J.E. et al. (2019) 'Refugia and anthelmintic resistance: Concepts and
challenges', International Journal for Parasitology: Drugs and Drug Resistance, 10, pp. 51–
57.

Howe, K.L. et al. (2017) 'WormBase ParaSite - a comprehensive resource for helminth
genomics', Molecular and Biochemical Parasitology, 215, pp. 2–10.

Huisman, J. (2017) 'Pedigree reconstruction from SNP data: parentage assignment, sibship
clustering and beyond', Molecular Ecology Resources, 17(5), pp. 1009–1024.

Huyse, T. et al. (2013) 'Regular treatments of praziquantel do not impact on the genetic
make-up of Schistosoma mansoni in Northern Senegal', Infection, Genetics and Evolution,
18, pp. 100–105.

Jacquard, A. (1966) 'Logique du calcul des coefficients d'identité entre deux individus',
Population, 21(4), pp. 751–776.

Kabatereine, N.B. et al. (2007) 'Impact of a national helminth control programme on infection

and morbidity in Ugandan schoolchildren', Bulletin of the World Health Organization, 85(2),
pp. 91–99.

Kaplan, R.M. and Vidyashankar, A.N. (2012) 'An inconvenient truth: global worming and
anthelmintic resistance', Veterinary Parasitology, 186(1-2), pp. 70–78.

Katz, N., Chaves, A. and Pellegrino, J. (1972) 'A simple device for quantitative stool thick-
smear technique in Schistosoma mansoni', Revista do Instituto de Medicina Tropical de Sao
Paulo, 14(6), pp. 397–400.

Kenyon, F. et al. (2009) 'The role of targeted selective treatments in the development of
refugia-based approaches to the control of gastrointestinal nematodes of small ruminants',
Veterinary Parasitology, 164(1), pp. 3–11.

Kittur, N. et al. (2019) 'Persistent hotspots in schistosomiasis consortium for operational
research and evaluation studies for gaining and sustaining control of schistosomiasis after
four years of mass drug administration of praziquantel', The American Journal of Tropical
Medicine and Hygiene, 101(3), pp. 617–627.

Kokaliaris, C. et al. (2022) 'Effect of preventive chemotherapy with praziquantel on
schistosomiasis among school-aged children in sub-Saharan Africa: a spatiotemporal
modelling study', The Lancet Infectious Diseases, 22(1), pp. 136–149.

Korunes, K.L. and Samuk, K. (2021) 'pixy: Unbiased estimation of nucleotide diversity and
divergence in the presence of missing data', Molecular Ecology Resources, 21(4), pp. 1359–
1368.

Krzywinski, M. et al. (2009) 'Circos: an information aesthetic for comparative genomics',
Genome Research, 19(9), pp. 1639–1645.

Lamberton, P.H.L., Faust, C.L. and Webster, J.P. (2017) 'Praziquantel decreases fecundity
in Schistosoma mansoni adult worms that survive treatment: evidence from a laboratory life-
history trade-offs selection study', Infectious Diseases of Poverty, 6(1), p. 110.

Layer, R.M. et al. (2014) 'LUMPY: a probabilistic framework for structural variant discovery',
Genome Biology, 15(6), p. R84.

Leathwick, D.M., Ganesh, S. and Waghorn, T.S. (2015) 'Evidence for reversion towards
anthelmintic susceptibility in Teladorsagia circumcincta in response to resistance
management programmes', International Journal for Parasitology: Drugs and Drug
Resistance, 5(1), pp. 9–15.

Le Clec'h, W. et al. (2021) 'Genetic analysis of praziquantel response in schistosome
parasites implicates a transient receptor potential channel', Science Translational Medicine,
13(625), p. eabj9114.

Lee-Six, H. et al. (2019) 'The landscape of somatic mutation in normal colorectal epithelial
cells', Nature, 574, pp. 532–537.

Lelo, A.E. et al. (2014) 'No apparent reduction in schistosome burden or genetic diversity
following four years of school-based mass drug administration in Mwea, central Kenya, a
heavy transmission area', PLoS Neglected Tropical Diseases, 8(10), p. e3221.

Li, H. (2013) 'Aligning sequence reads, clone sequences and assembly contigs with BWA-
MEM', arXiv [q-bio.GN]. Available at: http://arxiv.org/abs/1303.3997.

Lim, R.M. et al. (2023) 'Defining schistosomiasis hotspots based on literature and

shareholder interviews', Trends in Parasitology, 39(12), pp. 1032–1049.

Lo, N.C. et al. (2022) 'Review of 2022 WHO guidelines on the control and elimination of schistosomiasis', The Lancet Infectious Diseases, 22(11), pp. e327–e335.

Marchant, J.S. (2024) 'Progress interrogating TRPMPZQ as the target of praziquantel', PLoS Neglected Tropical Diseases, 18(2), p. e0011929.

Mari, L. et al. (2017) 'Heterogeneity in schistosomiasis transmission dynamics', Journal of Theoretical Biology, 432, pp. 87–99.

Mawa, P.A. et al. (2021) 'Schistosomiasis morbidity hotspots: roles of the human host, the parasite and their interface in the development of severe morbidity', Frontiers in Immunology, 12, p. 751.

McKenna, A. et al. (2010) 'The Genome Analysis Toolkit: a MapReduce framework for analyzing next-generation DNA sequencing data', Genome Research, 20(9), pp. 1297–1303.

Mei, Z.-Z. et al. (2006) 'Intracellular coiled-coil domain engaged in subunit interaction and assembly of melastatin-related transient receptor potential channel 2', The Journal of Biological Chemistry, 281(50), pp. 38748–38756.

Melman, S.D. et al. (2009) 'Reduced Susceptibility to Praziquantel among Naturally Occurring Kenyan Isolates of Schistosoma mansoni', PLoS Neglected Tropical Diseases, 3(8), p. e504.

Meurs, L. et al. (2013) 'Micro-geographical heterogeneity in Schistosoma mansoni and S. haematobium infection and morbidity in a co-endemic community in northern Senegal', PLoS Neglected Tropical Diseases, 7(12), p. e2608.

Miles, A. et al. (2020) cggh/scikit-allel: v1.3.1. Available at: https://doi.org/10.5281/zenodo.3935797.

Minh, B.Q. et al. (2020) 'IQ-TREE 2: New Models and Efficient Methods for Phylogenetic Inference in the Genomic Era', Molecular Biology and Evolution, 37(5), pp. 1530–1534.

Moore, L. et al. (2020) 'The mutational landscape of normal human endometrial epithelium', Nature, 580(7805), pp. 640–646.

Mutuku, M.W. et al. (2019) 'A search for snail-related answers to explain differences in response of Schistosoma mansoni to praziquantel treatment among responding and persistent hotspot villages along the Kenyan shore of Lake Victoria', The American Journal of Tropical Medicine and Hygiene, 101(1), pp. 65–77.

Mwangi, I.N. et al. (2014) 'Praziquantel sensitivity of Kenyan Schistosoma mansoni isolates and the generation of a laboratory strain with reduced susceptibility to the drug', International Journal for Parasitology: Drugs and Drug Resistance, 4(3), pp. 296–300.

Neves, M.I. et al. (2021) 'Revisiting density-dependent fecundity in schistosomes using sibship reconstruction', PLoS Neglected Tropical Diseases, 15(5), p. e0009396.

Neves, M.I., Webster, J.P. and Walker, M. (2019) 'Estimating helminth burdens using sibship reconstruction', Parasites & Vectors, 12(1), p. 441.

Park, S.-K. et al. (2019) 'The anthelmintic drug praziquantel activates a schistosome transient receptor potential channel', The Journal of Biological Chemistry, 294(49).

Park, S.-K. et al. (2021) 'Mechanism of praziquantel action at a parasitic flatworm ion
channel', Science Translational Medicine, 13(625), p. eabj5832.

Pitchford, R.J. and Visser, P.S. (1975) 'A simple and rapid technique for quantitative
estimation of helminth eggs in human and animal excreta with special reference to
Schistosoma sp.', Transactions of the Royal Society of Tropical Medicine and Hygiene,
69(3), pp. 318–322.

Platt, R.N. et al. (2019) 'Ancient Hybridization and Adaptive Introgression of an Invadolysin
Gene in Schistosome Parasites', Molecular Biology and Evolution, 36(10), pp. 2127–2142.

Platt, R.N. et al. (2021) 'Genomic analysis of a parasite invasion: colonization of the
Americas by the blood fluke, Schistosoma mansoni', Molecular Ecology, 31(8), pp. 2242–
2263.

Quinlan, A.R. and Hall, I.M. (2010) 'BEDTools: a flexible suite of utilities for comparing
genomic features', Bioinformatics , 26(6), pp. 841–842.

Rose, H. et al. (2015) 'Widespread anthelmintic resistance in European farmed ruminants: a
systematic review', The Veterinary Record, 176(21), p. 546.

Schwab, A.E. et al. (2005) 'Detection of benzimidazole resistance-associated mutations in
the filarial nematode Wuchereria bancrofti and evidence for selection by albendazole and
ivermectin combination treatment', The American Journal of Tropical Medicine and Hygiene,
73(2), pp. 234–238.

Smit, A.F.A., Hubley, R. and Green, P. (2015) RepeatMasker Open-4.0. 2013-2015.
Available at: http://www.repeatmasker.org.

Stelma, F.F. et al. (1995) 'Efficacy and side effects of praziquantel in an epidemic focus of
Schistosoma mansoni', The American Journal of Tropical Medicine and Hygiene, 53(2), pp.
167–170.

Summers, S. et al. (2022) 'A review of the genetic determinants of praziquantel resistance in
Schistosoma mansoni: Is praziquantel and intestinal schistosomiasis a perfect match?',
Frontiers in Tropical Diseases, 3.

Terhorst, J., Kamm, J.A. and Song, Y.S. (2017) 'Robust and scalable inference of population
history from hundreds of unphased whole genomes', Nature Genetics, 49(2), pp. 303–309.

Vianney, T.J. et al. (2022) 'Genome-wide analysis of Schistosoma mansoni reveals limited
population structure and possible praziquantel drug selection pressure within Ugandan hot-
spot communities', PLoS Neglected Tropical Diseases, 16(8), p. e0010188.

Webster, J.P. et al. (2014) 'The contribution of mass drug administration to global health:
past, present and future', Philosophical transactions of the Royal Society of London. Series
B, Biological sciences, 369(1645), p. 20130434.

WHO Expert Committee on the Control of Schistosomiasis (2002) Prevention and control of
schistosomiasis and soil-transmitted helminthiasis : report of a WHO expert committee.
World Health Organ Tech Rep Ser. Available at: http://apps.who.int/iris/handle/10665/42588
(Accessed: 13 November 2018).

World Health Organization (2020) A road map for neglected tropical diseases 2021–2030.
World Health Organization. Available at: https://www.who.int/neglected_diseases/Ending-
the-neglect-to-attain-the-SDGs--NTD-Roadmap.pdf.

World Health Organization (2021) 'Schistosomiasis and soil-transmitted helminthiases:
progress report, 2020 - Schistosomiase et géohelminthiases: rapport de situation, 2020',
Weekly Epidemiological Record, 96(48), pp. 585–595.

Yin, Y. et al. (2019) 'Structural basis of cooling agent and lipid sensing by the cold-activated
TRPM8 channel', Science, 363(6430), p. eaav9334.

Yu, G. et al. (2017) 'ggtree : an r package for visualization and annotation of phylogenetic
trees with their covariates and other associated data', Methods in Ecology and Evolution,
8(1), pp. 28–36.

1323