Muscle Type-Specific Modulation of Autophagy Signaling in Obesity: Effects of Caloric Restriction and Exercise

Fujue Ji[1,2], Yu Jin Park[1], Jong-Hee Kim[1,2,#]

[1] Major in Sport Science, Division of Sport Industry and Science, College of Performing Arts and Sport,

Hanyang University, 222 Wangsimni-ro, Seongdong-gu, Seoul, Republic of Korea

[2] BK21 FOUR Human-Tech Convergence Program, Hanyang University, 222 Wangsimni-ro,

Seongdong-gu, Seoul 04763, Republic of Korea

 # Corresponding Author

carachel07@hanyang.ac.kr

Background:

Obesity is a global health issue that contributes to the development of various diseases through metabolic dysregulation. Recent findings indicate that obesity leads to autophagy dysregulation, a cellular degradation process. Caloric restriction (CR) and CR combined with exercise (CR+Ex) are effective strategies for managing obesity and modulating autophagy. However, the regulation of autophagy and its signaling pathways in skeletal muscle under conditions of obesity, CR, and CR+Ex remains poorly understood.

Method:

Mice were divided into six groups: normal diet, normal diet CR, normal die CR+Ex, high-fat diet, high-fat diet CR, and high-fat diet CR+Ex. All mice were fed ad libitum with either a normal or high-fat diet for the first four months, followed by the respective interventions for the subsequent four months. We examined body composition, skeletal muscle functions, and expression of autophagy signaling pathway in these mice.

Result:

Obesity resulted in increased body weight, lean mass, fat mass, and fat mass in tissue; decreased grip strength and endurance ($P < 0.05$). CR+Ex decreased body weight, lean mass, and fat mass in obese mice ($P < 0.05$). In red muscle, P62, LC3B-I and LC3B-II levels were elevated ($P < 0.05$), regardless of dietary conditions. High-fat diet induced Cathepsin L was reduced in red muscle ($P < 0.05$).

Conclusion:

Obesity leads to altered body composition and impaired skeletal muscle function, which are partially improved by CR+Ex. The modulation of the autophagy pathway was more pronounced in red muscle compared to white muscle, irrespective of the intervention. Autophagic activity was higher in red muscle compared to white muscle.

Keywords: Obesity, Autophagy, Calorie Restriction, Exercise

1. Introduction

Obesity, characterized by excessive fat accumulation, poses a significant global health challenge,
with prevalence rates continuing to rise [1]. It is a major risk factor for numerous diseases,
including cardiovascular diseases, type 2 diabetes, certain cancers, and musculoskeletal disorders
[2]. The economic and health burdens associated with obesity necessitate the exploration of
effective management strategies. Obesity-induced diseases are often caused by metabolic
dysregulation, with autophagy playing a key role [3, 4]. Therefore, understanding autophagy can
provide new insights into the management and treatment of obesity.

Autophagy involves three sub-mechanisms: chaperone-mediated autophagy, microautophagy, and
macroautophagy, which work coordinately to clear damaged cellular matter and maintain
homeostasis [5]. Macroautophagy (hereafter referred to as autophagy) utilizes a phagophore to
engulf damaged material, then associates with a lysosome to form an autophagolysosome, leading
to the hydrolysis of the selected components [6]. Autophagy, a cellular degradation process, is
complex and influenced by various internal and external factors, including obesity, calorie
restriction (CR), and exercise (Ex) [7]. Obesity induced-abnormal autophagy can contribute to the
development of various diseases, such as non-alcoholic fatty liver disease and chronic kidney
disease [3, 4].

CR and CR combined with Ex (CR+Ex) have garnered significant attention as
non-pharmacological interventions for obesity management [8, 9]. CR, defined as a reduction in
calorie intake without malnutrition, has been shown to extend lifespan, decrease body fat ratio,
and improve metabolic health across various species [10]. Compared to CR alone, CR+Ex yields
greater metabolic benefits for obesity, including improved insulin sensitivity [11], reduced
inflammation [12], and enhanced lipid metabolism [13].

CR and CR+Ex are also known to modulate autophagic activity [14]. Both CR and Ex can
regulate autophagy synergistically/individually through multiple pathways, including
AMP-activated protein kinase (AMPK), mammalian target of rapamycin (mTOR) and forkhead
box O3 (FoxO3) [15-17]. For example, CR and CR+Ex have been demonstrated to regulate the
expression of autophagy signaling pathways in adipose tissue [18], liver [19], and heart [19],
presenting a dual mechanism of action in combating obesity. Despite the promising potential of
CR and CR+Ex in managing obesity, the specific regulatory mechanisms of autophagy in skeletal
muscle under these conditions remain poorly understood.

Skeletal muscle plays a pivotal role in whole-body health [20]. It is composed of different fiber
types, primarily red and white muscle fibers, each with distinct metabolic and functional
properties [21]. The differential response of these muscle fiber types to autophagy regulation
under conditions of obesity, CR, and CR+Ex is of significant interest. However, muscle
type-specific modulation of autophagy under these conditions is currently understudied.

This study seeks to refine our understanding of the effects of obesity on autophagy and the
mechanisms by which CR and CR+Ex modulate these effects. Additionally, it will explore how
the type specificity of skeletal muscle conditions the regulation of autophagy in response to
obesity, CR, and CR+Ex. The results of this study will provide scientific support for future
treatment and management strategies for obesity and offer a scientific basis for prescribing
exercise interventions.

2. Methods

2.1 Animals and Experimental Design

Male C57BL/6N mice (n=42, 12-months-old) were random divided into six experimental groups:

normal diet control (NDC), normal diet control 20% calorie restriction (NDR), normal diet control

20% calorie restriction + voluntary wheel exercise (NDRE), high-fat diet control (HFC), high-fat

diet control 20% calorie restriction (HFR), high-fat diet control 20% calorie restriction + voluntary

wheel exercise (HFRE). For the first 4 months, all mice were fed either a standard chow diet (ND;

Teklad Global 2018, Envigo Inc.) or a high-fat diet (HF; 45% kcal from fat, D12451, Research

Diets Inc.) (Table 1). In the subsequent 4 months, 20% calorie restriction (CR) and 20% calorie

restriction + voluntary wheel exercise (CR+Ex) interventions were applied to the respective

groups (Figure 1). The mice were housed in a controlled environment, maintained at a temperature

of 22 ± 2 °C with 50–60% humidity, and subjected to a 12-hour light/dark cycle. All animals had

ad libitum access to water and were allowed physical activity. The experimental procedures were

approved by the Institutional Animal Care and Use Committee (IACUC) of Hanyang University

(HYU 2021-0239A).

Table 1. Diet Composition

Group	NDC, NDR, and NDRE		HFC, HFR, and HFRE	
Diet	Teklad Global 2018		D12451	
	gm%	kcal%	gm%	kcal%
Protein	18.4	30	24	20
Carbohydrate	44.2	50	41	35
Fat	6	20	24	45
Total		100		100
Kacl/gm	3.1		4.73	

Figure 1. Schematic overview experimental study design.

2.2 Skeletal Muscle Function Tests

Skeletal muscle function was examined in mice two weeks prior to euthanasia. The function was

evaluated through tests measuring walking speed, endurance, physical activity, and grip strength

across six groups of mice.

Walking Speed: Walking speed was evaluated using the Rota-Rod test. An adaptation period was
conducted over the first week, involving a pre-test exercise at a constant speed of 5 rpm for 1
minute, once per day. For the official test, the Rota-Rod was set to acceleration mode, increasing
the speed from 5 to 50 rpm over a 5-minute period (Model 76-0770, Harvard Apparatus Inc.,
Holliston, MA, USA). The latency to fall from the device was recorded for each mouse. Each
mouse underwent three trials with a ten-minute interval between tests, and the best performance
among the three trials was used as the final outcome measure.

Endurance Capacity: Endurance capacity was assessed using a treadmill test. One week prior to
the official test, an adaptation phase was conducted where the mice ran at 3 cm/s for 10 minutes
daily on a 0-degree slope. During the official test, the treadmill speed increased by 1 cm/s every
20 seconds, starting from 5 cm/s, on a 0-degree slope. The trial concluded when the mouse
touched the shock pad (set at 0.5 mA) three times.

Physical Activity: Physical activity was measured using the voluntary wheel running test. Running
distance was recorded with a voluntary wheel (MAN86130, Lafayette Instrument Company,
Lafayette, IN, USA), where each wheel rotation equaled a distance of 0.4 meters. The average
running distance over a 5-day period was documented for each mouse.

Grip Strength: Grip strength was evaluated using the inverted-cling grip test. An adaptation phase
was conducted once daily during the first week. During the official test, each mouse was placed in
the center of a wire mesh screen, and a stopwatch was started. The screen was then rotated to an
inverted position over the course of 2 seconds, with the mouse's head descending first, and held
40-50 cm above a padded surface. The time until the mouse fell was recorded. This procedure was
repeated three times with ten-minute intervals between tests, and the longest time recorded was
used as the final measurement.

2.3 Body Composition

Body composition was examined after skeletal muscle function test. The mice were anesthetized
with 40 mg/kg ketamine and 0.8 mg/kg medetomidine, and their body composition was analyzed
using an InAlyzer Dual-energy X-ray Absorptiometry (DEXA) system from Micro Photonics Inc.
(PA, USA). The measured parameters included total mass (g), fat mass (g), lean mass (g), and fat
mass in tissue (%).

2.4 Western-Immunoblot (WB)

Then euthanize the anesthetized mouse by cervical dislocation. Following euthanasia, the red and
white gastrocnemius were dissected separately [22]. The muscles were lysed using an EzRIPA
Lysis Kit (WSE-7420, ATTO). Protein concentration was determined using the Pierce™
Bicinchoninic Acid Protein Assay Kit (Thermo Scientific, Waltham, MA, USA). Equal amounts of
protein samples (35 µg) were separated by sodium dodecyl sulfate-polyacrylamide gel
electrophoresis and electro-transferred to nitrocellulose membranes (Bio-Rad Laboratories,
Hercules, CA, USA). The membranes were blocked with 5% non-fat milk dissolved in
Tris-buffered saline with Tween-20 (TBST; 10 mM Tris, 150 mM NaCl, and 0.1% Tween-20; pH
7.6) for 1.5 hours at room temperature and then incubated with primary antibody overnight at 4°C.
Subsequently, the membranes were incubated with horseradish peroxidase-conjugated secondary
antibody for 1.5 hours at room temperature. The following primary and secondary antibodies were
used: Beclin-1 (Cell signaling, #3495, 1:1000), ATG-7 (Cell signaling, #8558, 1:1000), P62 (Cell

signaling, #23214, 1:1000), LC3B (Sigma, L7543, 1:500), LAMP-2 (Invitrogen, #PA1-665,
1:1000), Cathepsin-L (Abcam, Ab133641, 1:1000), Goat anti Rabbit (Invitrogen, #G-21234,
1:5000), m-IgG Fc (SANTA, sc-525409, 1:5000). Ponceau S staining was used to normalize
quantitative protein. Targeted bands were quantified via densitometry using Image J software
(National Institutes of Health, Bethesda, MD, USA).

2.5 Statistical Analysis

Statistical analyses were performed using GraphPad Prism software (version 9). Two-way analysis
of variance with Bonferroni' s post hoc test was applied. All results are expressed as the mean ±
SD. Statistical significance was set at $p < 0.05$, with asterisks indicating the following levels of
significance: *$p < 0.05$, **$p < 0.01$, ***$p < 0.001$, and ****$p < 0.0001$.

3. Results

3.1 *CR+Ex Improved Some of the Body Composition Changes Resulting From Obesity*

The data showed that a normal diet did not alter mice total mass, lean mass, fat mass, or fat mass
in tissue ($P > 0.05$). However, obesity induced by a high-fat diet resulted in a significant increase
in all these indices ($P < 0.05$) (Figure 2C). Additionally, neither CR nor CR+Ex altered body
composition in normal diet mice ($P > 0.05$). In contrast, CR+Ex resulted in a significant decrease
in total mass, lean mass, and fat mass in obese mice ($P < 0.05$), but not in fat mass in tissue ($P >
0.05$) (Figure 2C). These results indicate that high-fat diet-induced obesity leads to significant
changes in body composition, and that only CR+Ex improves some aspects of body composition
in high-fat diet-induced obese mice.

3.2 Obesity Induced Certain Skeletal Muscle Dysfunction Not Reversed by CR and CR+Ex

The results showed that high-fat diet-induced obesity resulted in reduced grip strength and
endurance in mice (P < 0.05) (Figure 3A and B). However, physical activity and walking speed
were not altered (P > 0.05) (Figure 3C and D). Additionally, it was unexpectedly found that neither
CR nor CR+Ex significantly improved skeletal muscle function, regardless of dietary condition.
These data suggest that high-fat diet-induced obesity reduces some aspects of skeletal muscle
function, CR and CR+Ex do not have a significant effect on reversing these impairments.

Figure 3. Part of Skeletal Muscle Function Reduced by Obesity and Not Reversed by CR and CR+Ex. (A) Strength
test. (B) Endurance test. (C) Physical activity test. (D) Walking speed test. Significant differences are denoted by
asterisks: p < 0.05 (*), p < 0.01 (**), p < 0.001 (***), and p < 0.0001 (****). All values are presented as mean ±
SD. Normal diet control (NDC, 20months, n=7), normal diet control 20% calorie restriction (NDR, 20months,
n=7), normal diet control 20% calorie restriction + voluntary wheel exercise (NDRE, 20months, n=7), high-fat
diet control (HFC, 20months, n=7), high-fat diet control 20% calorie restriction (HFR, 20months, n=7), high-fat
diet control 20% calorie restriction + voluntary wheel exercise (HFRE, 20months, n=7).

3.3 Increased Autophagic Activity via P62, LC3B-I and LC3B-II in Red Muscle Compared to White Muscle.

First, we examined the role of same skeletal muscle types in modulating autophagy pathway
signaling expression under various dietary (normal diet and high-fat diet) and intervention
conditions (CR and CR+Ex) (Figure 4). The results showed that neither CR nor CR+Ex

significantly modulated the expression of autophagy pathway signaling under different dietary
conditions, regardless of skeletal muscle type (P > 0.05).

Figure 4. No significant differences in autophagy pathway signaling between the same skeletal muscle types. All
values are presented as mean ± SD. Normal diet control (NDC, 20months, n=7), normal diet control 20% calorie
restriction (NDR, 20months, n=7), normal diet control 20% calorie restriction + voluntary wheel exercise (NDRE,
20months, n=7), high-fat diet control (HFC, 20months, n=7), high-fat diet control 20% calorie restriction (HFR,
20months, n=7), high-fat diet control 20% calorie restriction + voluntary wheel exercise (HFRE, 20months, n=7).

Next, we examined the role of the same dietary conditions in modulating autophagy pathway

signaling expression under different skeletal muscle types (white and red) and intervention
conditions (CR and CR+Ex) (Figure 5). The results showed increased expression of P62, LC3B-I,
and LC3B-II in red muscle compared to white muscle (P < 0.05), regardless of diet conditions
(Figure 5). Changes similar to those observed with a normal diet, high-fat diet-induced obesity
also causes specific alterations in autophagy signaling in mice skeletal muscle. Compared to white
muscle, a high-fat diet results in decreased expression of Cathepsin L in red muscle (P < 0.05).
Furthermore, compared to a normal high-fat diet, CR leads to a significant increase in the
expression of LC3B-I in red muscle. These results indicate that skeletal muscle types regulate
autophagy signaling more significantly compared to diet, CR, and CR+Ex.

Figure 5. Increased Autophagic Activity via P62, LC3B-I and LC3B-II in Red Muscle Compared to White Muscle. Significant differences are denoted by asterisks: $p < 0.05$ (*), $p < 0.01$ (**), $p < 0.001$ (***), and $p < 0.0001$ (****). All values are presented as mean $\pm$ SD. Normal diet control (NDC, 20months, n=7), normal diet control 20% calorie restriction (NDR, 20months, n=7), normal diet control 20% calorie restriction + voluntary wheel exercise (NDRE, 20months, n=7), high-fat diet control (HFC, 20months, n=7), high-fat diet control 20% calorie restriction (HFR, 20months, n=7), high-fat diet control 20% calorie restriction + voluntary wheel exercise (HFRE, 20months, n=7).

4. Discussion

Obesity has consistently been a prominent topic due to its significant threat to human health [1, 2]. Although Obesity induced-abnormal autophagy can lead to the development of various diseases, there is a lack of consensus on the mechanisms through which obesity regulates autophagy and the potential intervention strategies [3, 4]. Our results show that diet (normal and high-fat) and intervention (CR and CR+Ex) do not significantly modulate autophagy within the same skeletal muscle type. However, skeletal muscle type (white and red) and intervention (CR and CR+Ex) do modulate autophagy under the same dietary conditions. This suggests that skeletal muscle type specificity exerts a stronger influence on autophagy regulation. These findings enhance our understanding of the mechanisms underlying obesity and autophagy, providing a scientific basis for further in-depth studies on obesity and cellular degradation mechanisms.

4.1 Appropriateness of Obesity Mouse Model

In this experiment, we utilized 12-months-old mice, which correspond to humans aged 30-40 years [23]. It is well-documented that the prevalence of obesity in humans increases with age, especially after 35 years [24]. To model the onset of obesity around age 35 in humans, the mice were subjected to a high-fat diet at 12-months-old. We employed a high-fat diet (D12451, Research Diets Inc.) for 4 months to induce obese mouse model. This particular high-fat diet (D12451, Research Diets Inc.) is widely used for developing obese mouse models [25-27]. Moreover, while there is no universal definition for high-fat diet-induced obesity in mice, most studies use a minimum of 8 weeks of ad libitum high-fat diet (D12451, Research Diets Inc.) to establish such models [25, 28, 29]. Therefore, a four-month high-fat diet is sufficient to successfully establish an obese mouse model.

Upon analyzing the mice body composition using DEXA, significant increases were observed in body weight, lean mass, fat mass, and fat mass in tissue after the high-fat diet. These changes in body composition are consistent with those reported in previous studies on obese mice [30-32]. Additionally, obesity was found to impair skeletal muscle functions, supporting prior research that identifies skeletal muscle dysfunction as a characteristic of obesity [33, 34]. Therefore, our results indicate that a four-month high-fat diet effectively induces obesity in mice, providing a robust experimental model for further study.

4.2 Body Composition and Skeletal Muscle Function

Our results differ from those of earlier studies [35, 36]. Under normal dietary conditions, CR and CR+Ex did not alter body composition in mice. Additionally, in high-fat diet-induced obese mice, CR failed to alter body composition, while only CR+Ex significantly reduced total mass, lean mass, and fat mass. The difference in results may be related to the intensity of CR and the age of

the experimental mice. In our study, CR intensity was lower (20%) and the mice were older (12 to
20-months-old) compared to the prior studies. Increasing age leads to slower metabolism and
reduced responsiveness and adaptability to external stimuli [37]. High-intensity CR has a negative
effect on older mice [38], so we chose low CR to protect the survival of late middle-aged mice and
successfully complete the experiment.
CR+Ex altered only part of the body composition in the high-fat diet group, but not in the normal
diet group, likely due to the higher initial body weight [39] and higher fat oxidation efficiency
[40], making these mice more sensitive to the effects of CR and Ex. Overall, obesity caused by a
high-fat diet altered the body composition of mice, and CR+Ex improved some of these effects.
Our data show that high-fat-induced obesity leads to reduced grip strength and endurance,
consistent with prior studies [41, 42]. However, physical activity and walking speed were not
affected. Contrary to predictions, CR and CR+Ex did not significantly increase skeletal muscle
function in late middle-aged mice, regardless of dietary condition. This may be due to the
combination of low-intensity CR and exercise, as well as the reduced responsiveness and
adaptability to external stimuli at advanced age [37].
290
4.3 Regulation of Skeletal Muscle Autophagy by CR and CR+Ex
To investigate autophagy and regulatory mechanisms in the skeletal muscle of obese late
middle-aged mice, we first examined the expression of autophagy signaling pathways in the same
muscle type under different dietary conditions (high-fat diet and normal diet) and intervention
conditions (CR and CR+Ex).
We did not observe any changes in the expression of autophagy pathway signals due to high-fat
diet-induced obesity. Although such results are uncommon, there have been reports of no
significant changes in autophagy signaling in the heart [43], skeletal muscle [44], and liver [45]
following high-fat diet-induced obesity. Obesity induced by a high-fat diet may interfere with
autophagy signaling pathways through various mechanisms. Obesity caused by a high-fat diet can
affect cellular nutrient sensing mechanisms, leading to the dysregulation of autophagy signaling
pathways. For instance, a high-fat diet can result in insulin resistance and metabolic syndrome,
thereby disrupting the regulatory mechanisms of autophagy [46, 47]. Studies have shown that
despite the appropriate upregulation of autophagy-related genes at the mRNA level, there is a
failure to correspondingly upregulate at the protein level, indicating an inconsistency in cellular
autophagy regulation under obese conditions [43]. Additionally, oxidative stress and inflammatory
responses triggered by a high-fat diet induced-obesity may disrupt the normal function of
autophagy signaling pathways. The accumulation of fatty acids and the increase of inflammatory
factors such as TNF-α and IL-1β can inhibit the autophagy process [48].
We also did not observe changes in the autophagy pathway in the same skeletal muscle type due to
CR and CR+Ex, regardless of diet conditions. Previous studies support our findings, as some
research has indicated that long-term CR and/or CR+Ex do not induce significant changes in
autophagy signaling proteins [14, 19, 43]. The impact of these interventions on autophagy
signaling pathways may not always be consistent. For example, long-term CR and/or CR+Ex were
found to upregulate autophagy-related protein mRNA expression, but this upregulation did not
translate into significant changes at the protein level [14, 43]. Additionally, while these
interventions can activate autophagy to some extent, the response of autophagy signaling in tissues
such as the heart and skeletal muscle may not be significant [19, 43]. This suggests tissue-specific

expression of autophagy.

Both CR and Ex promote autophagy by inhibiting mTOR signaling pathway and activating AMPK.

The overlap of these pathways might lead to a lack of significant enhancement in autophagy

activity when combined, as they operate through similar mechanisms [49]. Research indicates that

CR and Ex can independently activate AMPK and inhibit mTOR, thereby promoting autophagy.

However, when CR and Ex are combined, these signaling pathways might already be maximally

activated, preventing further enhancement of autophagy activity [50, 51]. Additionally, both CR

and Ex can independently regulate autophagy through the PI3K/AKT/mTOR and ERK1/2-MAPK

signaling pathways. When used together, these pathways might already be fully activated,

resulting in no additional significant effects from their combination [52].

4.4 Regulation of Autophagy by Skeletal Muscle Type-Specific

Subsequently, we examined the differences in autophagy expression between different types of

skeletal muscle. The expression of certain autophagy-related proteins was higher in red muscle

compared to white muscle. Additionally, dietary conditions and intervention methods did not alter

the muscle type-specific regulation of autophagy.

The expression of P62 is higher in red muscle than in white muscle, primarily due to differences in

metabolic demands and functions. Red muscle has higher oxidative metabolic activity, producing

more reactive oxygen species (ROS). P62 helps mitigate oxidative stress by activating the

transcription factor NFE2-related factor 2 (Nrf2) signaling pathway, which enhances the

expression of antioxidant enzymes [53]. Secondly, red muscle contains more mitochondria and has

higher oxidative metabolism, necessitating elevated autophagic activity to maintain mitochondrial

quality and function. P62 plays a crucial role in autophagy [54]. Additionally, red muscle has a

greater capacity for fatty acid oxidation, requiring high levels of P62 to regulate fatty acid uptake

and metabolism [55]. P62 also acts as a bridge in autophagy, promoting the degradation of

damaged proteins. Red muscle needs higher levels of P62 to maintain proper protein degradation

and recycling [56]. Furthermore, there are significant differences in gene expression between red

and white muscle, with many autophagy and metabolism-related genes being more highly

expressed in red muscle, including P62 [57].

Studies have shown that the expression of LC3B-I and LC3B-II is significantly higher in red

muscle than in white muscle. Red muscle has higher oxidative metabolic activity and more

mitochondria, which requires higher autophagic activity to maintain mitochondrial quality and

function. LC3B plays a key role in this process [54, 58]. Additionally, gene chip analysis and

protein expression studies have found that the expression levels of various genes and proteins are

significantly higher in red muscle than in white muscle, including those related to autophagy [57,

59]. These findings indicate that the high expression of LC3B in red muscle is primarily to meet

its higher oxidative metabolic demands and to respond to oxidative stress, maintaining cellular

homeostasis and function.

In an obesity state induced by a high-fat diet, the expression of Cathepsin L is higher in white

muscle than in red muscle, likely due to differences in metabolic demands, inflammatory

responses, and autophagic needs. Firstly, white muscle fibers primarily undergo glycolytic

metabolism, and a high-fat diet increases their metabolic burden, leading to a higher demand for

protein degradation, thereby promoting Cathepsin L expression [58]Secondly, obesity induced by

a high-fat diet is accompanied by chronic inflammation and oxidative stress. White muscle is

more susceptible to these stress factors, resulting in increased Cathepsin L expression [60]. Finally,
Cathepsin L plays a crucial role in autophagy. A high-fat diet increases the protein degradation
demand in white muscle, leading to its high expression to meet autophagic needs [61]. These
mechanisms collectively contribute to the high expression of Cathepsin L in white muscle to cope
with metabolic stress and protein degradation demands.
Overall, the expression of autophagy-related proteins was higher in red muscle than in white
muscle, which is closely related to different metabolic demands and functions. Red muscle have a
higher oxidative capacity, making them more reliant on the autophagy process to cope with
cellular stress and maintain intracellular homeostasis [62]. Furthermore, autophagy activity in
Red muscle helps remove damaged mitochondria and proteins, maintaining muscle function and
metabolic stability [63]. These characteristics enable   Red muscle to better adapt to prolonged
low-intensity activities and more effectively carry out autophagy to maintain cellular health.
375

Conclusion
Among the various interventions studied, only the combination of CR+Ex was effective in
partially reversing the high-fat diet-induced increases in body weight, lean mass, and fat mass.
Moreover, muscle type specificity plays a crucial role in the regulation of autophagy, which is not
easily modified by dietary or exercise interventions. This study provides valuable insights into the
muscle type-specific mechanisms of autophagy in the context of obesity and highlights the
potential for tailored exercise and dietary strategies in managing obesity-related health issues.
383

384 **Limitation**
385 This study exclusively used male mice, which may not fully represent the responses of other
386 breeds or female mice. Additionally, more advanced research methods, such as spatial proteomics,
387 should be employed to more accurately analyze the muscle type-specific regulation of autophagy
388 pathways.
389

390 **Representative Picture of Western-Immunoblot**

391
392

393 **Ethical approval and consent to participate**
394 All the procedures followed in this experiment were approved by the Institutional Animal Care
395 and Use Committee (IACUC) of Hanyang University (HYU 2021-0066A).
396

397 **Consent for publication**
398 Not applicable.

Availability of data and materials

The data used to support the findings of this study are presented here. Any further data requirements are available from the corresponding author upon request.

Competing interests

No conflicts of interest, financial or otherwise, have been declared by the author(s).

Funding

This research was supported by the National Research Foundation of Korea (NRF-2020R1F1A1061726).

Author's contributions

F.J.J. and J.H.K. conceived and designed the study. F.J.J. and Y.J.P. performed the experiments. F.J.J. analyzed the data and prepared the figures. F.J.J. and J.H.K. interpreted the results, drafted, edited, and revised the manuscript. J.H.K. acquired funding. All authors approved the final version of the manuscript.

Acknowledgments

We thank Haesung Lee (Hanyang University, Korea) for providing technical support with the statistical analysis of the experimental data.

We thank Jian Guo (Hanyang University, Korea) for his generous help in dissecting the mice.

We are grateful to Prof. Dr. Gwang-woong Go (Hanyang University, Korea) for generously providing the DEXA machine for body composition analysis of the mice.

We thank the support of Skill Learning from Kaixin Doctor and MASCU (Medical Association with Science, Creativity, and Unity), Inc, Shenzhen, China (mascu_forever@163.com).

Reference

1.   Kim, D.M., C.W. Ahn, and S.Y. Nam, *Prevalence of obesity in Korea.* Obesity Reviews, 2005. **6**(2): p. 117-121.

2.   Malik, V.S., W.C. Willett, and F.B. Hu, *Global obesity: trends, risk factors and policy implications.* Nature Reviews Endocrinology, 2013. **9**(1): p. 13-27.

3.   Lavallard, V.J., et al., *Autophagy, signaling and obesity.* Pharmacological research, 2012. **66**(6): p. 513-525.

4.   Behl, T., et al., *Understanding the molecular mechanisms and role of autophagy in obesity.* Molecular Biology Reports, 2021. **48**(3): p. 2881-2895.

5.   Česen, M.H., et al., *Lysosomal pathways to cell death and their therapeutic applications.* Exp Cell Res, 2012. **318**(11): p. 1245-51.

6.   Mizushima, N., T. Yoshimori, and Y. Ohsumi, *The role of Atg proteins in autophagosome formation.* Annu Rev Cell Dev Biol, 2011. **27**: p. 107-32.

7.   Mizushima, N., *Autophagy: process and function.* Genes & development, 2007. **21**(22): p. 2861-2873.

8.   Wadden, T.A., *Treatment of obesity by moderate and severe caloric restriction: results of clinical research trials.* Annals of internal medicine, 1993. **119**(7_Part_2): p. 688-693.

9. Liu, Y., et al., *Calorie restriction with exercise intervention improves inflammatory response in overweight and obese adults: a systematic review and meta-analysis.* Frontiers in physiology, 2021. **12**: p. 754731.

10. Speakman, J.R. and S.E. Mitchell, *Caloric restriction.* Molecular aspects of medicine, 2011. **32**(3): p. 159-221.

11. Coker, R.H., et al., *The impact of exercise training compared to caloric restriction on hepatic and peripheral insulin resistance in obesity.* The Journal of Clinical Endocrinology & Metabolism, 2009. **94**(11): p. 4258-4266.

12. Khalafi, M., M.E. Symonds, and A. Akbari, *The impact of exercise training versus caloric restriction on inflammation markers: a systemic review and meta-analysis.* Critical Reviews in Food Science and Nutrition, 2022. **62**(15): p. 4226-4241.

13. Davis, R.A., et al., *High-intensity interval training and calorie restriction promote remodeling of glucose and lipid metabolism in diet-induced obesity.* American Journal of Physiology-Endocrinology and Metabolism, 2017. **313**(2): p. E243-E256.

14. Wohlgemuth, S.E., et al., *Skeletal muscle autophagy and apoptosis during aging: effects of calorie restriction and life-long exercise.* Experimental gerontology, 2010. **45**(2): p. 138-148.

15. Tam, B.T. and P.M. Siu, *Autophagic cellular responses to physical exercise in skeletal muscle.* Sports medicine, 2014. **44**: p. 625-640.

16. Bagherniya, M., et al., *The effect of fasting or calorie restriction on autophagy induction: A review of the literature.* Ageing research reviews, 2018. **47**: p. 183-197.

17. Schwalm, C., et al., *Activation of autophagy in human skeletal muscle is dependent on exercise intensity and AMPK activation.* The FASEB Journal, 2015. **29**(8): p. 3515-3526.

18. Jansen, H., et al., *Autophagy activity is up-regulated in adipose tissue of obese individuals and modulates proinflammatory cytokine expression.* Endocrinology, 2012. **153**(12): p. 5866-5874.

19. Wohlgemuth, S.E., et al., *Autophagy in the heart and liver during normal aging and calorie restriction.* Rejuvenation research, 2007. **10**(3): p. 281-292.

20. Kim, G. and J.H. Kim, *Impact of skeletal muscle mass on metabolic health.* Endocrinology and Metabolism, 2020. **35**(1): p. 1.

21. Essen, B., et al., *Metabolic characteristics of fibre types in human skeletal muscle.* Acta Physiologica Scandinavica, 1975. **95**(2): p. 153-165.

22. Chen, D., Y. Wang, and E.R. Chin, *Activation of the endoplasmic reticulum stress response in skeletal muscle of G93A*SOD1 amyotrophic lateral sclerosis mice.* Front Cell Neurosci, 2015. **9**: p. 170.

23. Justice, J.N., et al., *Comparative Approaches to Understanding the Relation Between Aging and Physical Function.* J Gerontol A Biol Sci Med Sci, 2016. **71**(10): p. 1243-53.

24. Chooi, Y.C., C. Ding, and F. Magkos, *The epidemiology of obesity.* Metabolism, 2019. **92**: p. 6-10.

25. Han, C., et al., *Studies on production of high fat diet induced obesity C57BL/6NCrjBgi mice.* Laboratory Animal Research, 2006. **22**(3): p. 221-226.

26. Yang, Y., et al., *Variations in body weight, food intake and body composition after long-term high-fat diet feeding in C57BL/6J mice.* Obesity, 2014. **22**(10): p. 2147-2155.

27. Armani, A., et al., *Mineralocorticoid receptor antagonism induces browning of white adipose tissue through impairment of autophagy and prevents adipocyte dysfunction in high-fat-diet-fed mice.* The FASEB Journal, 2014. **28**(8): p. 3745-3757.

28. Hariri, N. and L. Thibault, *High-fat diet-induced obesity in animal models.* Nutrition research
reviews, 2010. **23**(2): p. 270-299.

29. Buettner, R., J. Schölmerich, and L.C. Bollheimer, *High-fat diets: modeling the metabolic*
disorders of human obesity in rodents. Obesity, 2007. **15**(4): p. 798-808.

30. Lin, P.-Y., D.R. Romsos, and G.A. Leveille, *Food intake, body weight gain, and body*
composition of the young obese (ob/ob) mouse. The Journal of Nutrition, 1977. **107**(9): p.
1715-1723.

31. Rogers, P. and G.P. Webb, *Estimation of body fat in normal and obese mice.* British Journal of
Nutrition, 1980. **43**(1): p. 83-86.

32. Mandarim-de-Lacerda, C.A., et al., *Mice as an animal model for the study of adipose tissue*
and obesity. Int J Morphol, 2021. **39**(6): p. 1521-1528.

33. Zhang, Y., et al., *Obesity-induced oxidative stress, accelerated functional decline with age and*
increased mortality in mice. Archives of biochemistry and biophysics, 2015. **576**: p. 39-48.

34. Shortreed, K.E., et al., *Muscle-specific adaptations, impaired oxidative capacity and*
maintenance of contractile function characterize diet-induced obese mouse skeletal muscle.
PloS one, 2009. **4**(10): p. e7293.

35. Sugimoto, T., et al., *A combination of exercise and calorie restriction improves the*
development of obesity-related type 2 diabetes mellitus in KKAy mice. Bioscience,
Biotechnology, and Biochemistry, 2023. **87**(1): p. 108-113.

36. Huang, P., et al., *Calorie restriction and endurance exercise share potent anti-inflammatory*
function in adipose tissues in ameliorating diet-induced obesity and insulin resistance in mice.
Nutrition & Metabolism, 2010. **7**(1): p. 59.

37. Rattan, S.I. and A. Derventzi, *Altered cellular responsiveness during ageing.* Bioessays, 1991.
13(11): p. 601-606.

38. Schädel, P., et al., *Short-Term Caloric Restriction and Subsequent Re-Feeding Compromise*
Liver Health and Associated Lipid Mediator Signaling in Aged Mice. Nutrients, 2023. **15**(16): p.
3660.

39. Hansen, D., et al., *The effects of exercise training on fat-mass loss in obese patients during*
energy intake restriction. Sports Medicine, 2007. **37**: p. 31-46.

40. Segal, K.R. and F.X. Pi-Sunyer, *Exercise and obesity.* Medical Clinics of North America, 1989.
73(1): p. 217-236.

41. Huang, C.C., et al., *Whole-body vibration training effect on physical performance and obesity*
in mice. Int J Med Sci, 2014. **11**(12): p. 1218-27.

42. Haramizu, S., et al., *Different contribution of muscle and liver lipid metabolism to endurance*
capacity and obesity susceptibility of mice. Journal of Applied Physiology, 2009. **106**(3): p.
871-879.

43. Andres, A.M., et al., *Discordant signaling and autophagy response to fasting in hearts of*
obese mice: implications for ischemia tolerance. American Journal of Physiology-Heart and
Circulatory Physiology, 2016. **311**(1): p. H219-H228.

44. Campbell, T.L., et al., *High-fat feeding does not induce an autophagic or apoptotic phenotype*
in female rat skeletal muscle. Exp Biol Med (Maywood), 2015. **240**(5): p. 657-68.

45. Budi, Y.P., et al., *The role of autophagy in high-fat diet-induced insulin resistance of adipose*
tissues in mice. PeerJ, 2022. **10**: p. e13867.

46.   Liu, J., et al., *CaMKIV limits metabolic damage through induction of hepatic autophagy by*
CREB in obese mice. Journal of Endocrinology, 2020. **244**(2): p. 353-367.

47.   Anderson, E.J., et al., *Mitochondrial H 2 O 2 emission and cellular redox state link excess fat*
intake to insulin resistance in both rodents and humans. The Journal of clinical investigation,
2009. **119**(3): p. 573-581.

48.   Liang, L., et al., *Antioxidant catalase rescues against high fat diet-induced cardiac dysfunction*
via an IKKβ-AMPK-dependent regulation of autophagy. Biochimica et Biophysica Acta
(BBA)-Molecular Basis of Disease, 2015. **1852**(2): p. 343-352.

49.   Escobar, K.A., et al., *Autophagy and aging: Maintaining the proteome through exercise and*
caloric restriction. Aging cell, 2019. **18**(1): p. e12876.

50.   Cui, M., et al., *Chronic caloric restriction and exercise improve metabolic conditions of*
dietary-induced obese mice in autophagy correlated manner without involving AMPK. Journal
of diabetes research, 2013. **2013**(1): p. 852754.

51.   Zheng, Q., et al., *Inhibition of AMPK accentuates prolonged caloric restriction-induced change*
in cardiac contractile function through disruption of compensatory autophagy. Biochimica et
Biophysica Acta (BBA)-Molecular Basis of Disease, 2015. **1852**(2): p. 332-342.

52.   Ferreira-Marques, M., et al., *PI3K/AKT/MTOR and ERK1/2-MAPK signaling pathways are*
involved in autophagy stimulation induced by caloric restriction or caloric restriction mimetics
in cortical neurons. Aging (Albany NY), 2021. **13**(6): p. 7872.

53.   Yamada, M., et al., *Muscle p62 stimulates the expression of antioxidant proteins alleviating*
cancer cachexia. The FASEB Journal, 2023. **37**(9): p. e23156.

54.   Glancy, B. and R.S. Balaban, *Protein composition and function of red and white skeletal*
muscle mitochondria. American Journal of Physiology-Cell Physiology, 2011. **300**(6): p.
C1280-C1290.

55.   Chabowski, A., et al., *Fatty acid transport and FAT/CD36 are increased in red but not in white*
skeletal muscle of ZDF rats. American Journal of Physiology-Endocrinology and Metabolism,
2006. **291**(3): p. E675-E682.

56.   Jørgensen, S.B., et al., *Role of AMPKα2 in basal, training-, and AICAR-induced GLUT4,*
hexokinase II, and mitochondrial protein expression in mouse muscle. American Journal of
Physiology-Endocrinology and Metabolism, 2007. **292**(1): p. E331-E339.

57.   Li, Y., et al., *Differential transcriptional analysis between red and white skeletal muscle of*
Chinese Meishan pigs. International journal of biological sciences, 2010. **6**(4): p. 350.

58.   Campbell, W.G., et al., *Differential global gene expression in red and white skeletal muscle.*
American Journal of Physiology-Cell Physiology, 2001. **280**(4): p. C763-C768.

59.   Kim, N.K., et al., *Differential expression profiling of the proteomes and their mRNAs in porcine*
white and red skeletal muscles. Proteomics, 2004. **4**(11): p. 3422-3428.

60.   Liu, J., et al., *Cathepsin L expression and regulation in human abdominal aortic aneurysm,*
atherosclerosis, and vascular cells. Atherosclerosis, 2006. **184**(2): p. 302-311.

61.   Yamashita, M. and S. Konagaya, *High activities of cathepsins B, D, H, and L in the white muscle*
of chum salmon in spawning migration. Comparative biochemistry and physiology. B,
Comparative biochemistry, 1990. **95**(1): p. 149-152.

62.   Ogata, T., et al., *Fasting-related autophagic response in slow-and fast-twitch skeletal muscle.*
Biochemical and biophysical research communications, 2010. **394**(1): p. 136-140.

63.   Mofarrahi, M., et al., *Autophagic flux and oxidative capacity of skeletal muscles during acute*
starvation. Autophagy, 2013. **9**(10): p. 1604-1620.
575