Experimental capture of genomic islands defines a widespread class of genetic element capable of non-autonomous transfer

Yansong Zhao[1,*], Yue Ma[2], Andrew D. Farr[1], David W. Rogers[1] & Paul B. Rainey[1,3,*]

[1] Department of Microbial Population Biology, Max Planck Institute for Evolutionary Biology, Plön, Germany

[2] Guest Group Evolutionary Medicine, Max Planck Institute for Evolutionary Biology, Plön, Germany

[3] Laboratory of Biophysics and Evolution, CBI, ESPCI Paris, Université PSL, CNRS, Paris, France

* To whom correspondence may be addressed. Emails: Y.Z. (yansongzhao@evolbio.mpg.de), P.B.R. (rainey@evolbio.mpg.de).

Abstract

Bacteria acquire new genes by horizontal gene transfer (HGT). Acquisition is typically mediated by mobile genetic elements (MGEs), however, beyond plasmids, bacteriophages and certain integrative conjugative elements (ICEs), the nature and diversity of MGEs is poorly understood. The bacterium *Pseudomonas fluorescens* SBW25 was propagated by serial transfer in the presence of filtrate obtained from garden compost communities. Genome sequencing of derived colonies revealed acquisition of three different mobile elements, each integrated immediately downstream of *tmRNA*. All are flanked by direct repeats and harbour a tyrosine integrase (*intY*), followed by a cargo of accessory genes including putative phage defence systems. Although characteristic of genomic islands, MGE-classifiers showed no matches to mobile elements. Interrogation of DNA sequence databases showed that similar elements are widespread in the genus *Pseudomonas* and beyond, with *Vibrio* Pathogenicity Island-1 (VPI-1) from *V. cholerae* being a notable example. Bioinformatic analyses demonstrate frequent transfer among diverse hosts. With focus on a single 55 kb element (I55) we show that *intY* is necessary for excision and circularisation, that the element is incapable of autonomous horizontal transfer, but is mobilizable – in the absence of direct cell-cell contact – upon addition of community filtrate. Further analyses demonstrate that I55 enhances host fitness in the presence of community filtrate, which stems in part from ability to defend against phages.

Significance Statement

The impact of horizontal gene transfer on the evolution of bacteria outpaces that driven by spontaneous mutation, but knowledge of the range of mediators, the genes mobilised, and the mechanisms of movement have largely depended on inferences stemming from bioinformatics. Here we describe a real-time evolution experiment in which a single focal strain propagated with filtrate from a complex microbial community captures genetic elements carrying a diverse cargo of genes whose mobility was previously unknown. The elements are representative of a class of mobile DNA that depend on nothing

38   other than a tyrosine integrase that targets a highly conserved gene. The elements encode phage
39   resistance, require the integrase to excise from the genome, confer a significant fitness advantage, and
40   depend on a secondary element for transfer.

42   Key Words
43   community ecology – comparative genomics – experimental evolution – lateral transduction

Introduction

There is growing awareness of the role of mobile genetic elements (MGEs), such as phages, plasmids, and integrative conjugative elements (ICEs), in shaping the structure and function of microbial communities (Arnold et al., 2022). While encoding their own replicative capacity, MGEs oftentimes capture and move genes of ecological relevance among hosts (Arnold et al., 2022; Dmitrijeva et al., 2024; Quistad et al., 2020) by horizontal gene transfer (HGT). Recently, it has even been suggested that accessory genes define a set of "community genes" that could allow selection to operate on communities by promoting the efficacy of feedback cycles that underpin differential persistence (Doolittle & Inkpen, 2018; Lenton et al., 2021; Rainey & Quistad, 2020; Roughgarden, 2020).

While classical genetics has led to discovery of various mechanisms of HGT, including conjugation (Lederberg & Tatum, 1946), transduction (Zinder & Lederberg, 1952) and transformation (Griffith, 1928), the majority of recent discoveries stem from genome analysis (Dmitrijeva et al., 2024; Groussin et al., 2021; Hackl et al., 2023) . For example, analysis of closely related strains often reveals regions with distinct evolutionary histories – often referred to as "genomic islands" (GIs) – that are then marked as having likely arisen by HGT (Koutsovoulos et al., 2022; Yuan et al., 2023). While instructive, such bioinformatic methods rarely provide insight into the causes of horizontal movement. Notable exceptions include demonstration of the mobility of ICEs (Colombi et al., 2017; J. T. Sullivan & Ronson, 1998), of plasmids (Hall et al., 2016; Lilley & Bailey, 1997b), of gene transfer agents (Lang et al., 2012; Marrs, 1974), and of phage-inducible chromosomal islands (PICIs) (Barcia-Cruz et al., 2024; Fillol-Salom et al., 2019; Lindsay et al., 1998; Novick et al., 2010). The PICIs, a class of phage satellite (Ibarra-Chávez et al., 2021) , hitchhike with, or hijack, helper bacteriophages to facilitate their own intercellular transfer.

Experimental evolution, where populations or communities are propagated under controlled conditions (Lenski, 2017; Rainey et al., 2017; Rainey & Travisano, 1998), while not typically used to study HGT, offer new possibilities to understand the range of genes disseminated by MGEs, the mediators of transfer, and functional consequences (Frazão et al., 2019; Hall et al., 2016; Lilley & Bailey, 1997a; Quistad et al., 2020; Woods et al., 2020). For example, Quistad et al. (Quistad et al., 2020), focusing on complex microbial communities founded from garden compost, employed an experimental design in which the evolutionary life of MGEs was either enhanced, or curtailed. Comparative metagenomic analyses revealed a significant flux of horizontally disseminated DNA, including putative transfer agents (Quistad et al., 2020; van Dijk et al., 2023), but in the absence of culturable hosts carrying newly acquired DNA – underpinned by further analyses of fitness and genetics – understanding is limited to inferences stemming from bioinformatic analyses.

In this study, we use experimental evolution to detect, isolate and characterise novel MGEs from environmental sources. To achieve this, we simplified the experimental strategy of Quistad *et al.* (Quistad et al., 2020) to one in which a single bacterium, *Pseudomonas fluorescens* SBW25 (Fortmann-Grote et al., 2023; Silby et al., 2009), was propagated in the presence of filtrate derived from complex compost communities. This led to capture of three GIs that define a class of MGE unclassified by current identification tools. The elements integrate immediately downstream of *tmRNA*, are diverse in terms of size and gene content, with many encoding putative phage defence systems, but also metabolic genes, and many with no matches to sequences in databases. Bioinformatic analyses based on a minimal set of core genes demonstrate the elements to be representative of a widespread class of GI whose mobility

has not been previously demonstrated. Detailed analysis of a single element shows that the integrase is
necessary for spontaneous excision from the genome and concomitant circularisation, but intracellular
transfer occurs only in the presence of community filtrate. The element confers a fitness benefit in the
presence of community filtrate, but is weakly deleterious in pure culture. Further analysis shows that
the fitness benefit stems in part from enhanced resistance to bacteriophages present in the community
filtrate.

Results

Pseudomonas fluorescens SBW25 propagated in the presence of compost filtrate acquired mobile genetic elements

A simple mark-recapture approach (Fig. 1a(ii)) was used to see whether SBW25 introduced into garden compost communities might acquire MGEs. To this end, SBW25 was marked with gentamycin resistance (strain MPB28806), placed in static microcosms together with wash from compost and propagated under a regime involving transfer of a 1-in-100 aliquot every 48 hours. Samples were obtained at regular intervals, however marked SBW25 was not recovered after ~5 transfers (see Fig. S1). This most likely reflected inability of SBW25 to persist in the face of competition imposed by community members. We next propagated SBW25 (marked with either GFP or RFP, strains MPB29284 and MPB29300, respectively) with filtrate derived from the community, using two slightly different protocols referred to as "live-filtrate" and "fixed-filtrate" regimes (see Fig. 1a(i) and 1a(iii), respectively). Choice of the two regimes was motivated by concerns of potential negative effects of storing samples at -80 °C, but also the benefits stemming from having archived samples. In both cases, filtrate was obtained by passing community culture through a 0.2 µm filter, thus eliminating all cells with the exception of nanobacteria (Quistad et al., 2020). For the live-filtrate regime, material was obtained from a single synchronised parallel community (compost C) that was also subject to transfer every 48 hours. On each occasion 1 mL of filtrate was obtained with 50 µL being added to each of eight independent serially-propagated replicate populations of SBW25 (Fig. 1a(i)). Populations were transferred 14 times. For the fixed-filtrate regime, a single sample of filtrate was obtained from each of three independent, seven-day-old compost communities (composts B, C and D) and stored at -80 °C. An aliquot of 50 µL from the archived sample was added at each serial transfer (Fig. 1a(iii)). Ten replicate SBW25 populations received filtrate from each of the three independent communities. Populations were transferred seven times.

At the end of each experiment, samples of derived SBW25 were plated on agar. For the live-filtrate regime, two to three colonies were selected from each of the eight replicate lines and re-sequenced. For the fixed-filtrate regime, 24 colonies were chosen at random from each replicate, grown separately overnight and then pooled, with DNA extracted from the pooled sample being re-sequenced. Mapping DNA sequence reads back to the ancestral SBW25 genome sequence revealed the presence of DNA sequences that did not match the genome (Fig. 1b). Further analysis of these reads led to identification of three GIs that had been acquired during the course of the selection experiment. The elements, designated I23, I44 and I55 (approximately 23 kb, 44 kb and 55 kb in length, respectively), were identified on five independent occasions (see Fig. 1c). From the 19 colonies sequenced from the live-filtrate regime, one contained I55 (strain MPB33357), which was acquired from compost C. Of the 720 colonies sequenced from the fixed-filtrate regime, six contained I23 (strains MPB38392, MPB38393, MPB38394, MPB38395, MPB38396, MPB38895) and two contained I44 (strains MPB38906, MPB38907). I23 was acquired independently on three occasions from compost B; I44 was detected in just a single replicate receiving filtrate from compost B.

The three GIs are unrecognised by MGE-classifiers

Mapping the ends of contigs containing each element confirmed that all three GIs inserted at the same genomic locus, immediately downstream of *tmRNA* and immediately upstream of a putative mobilisable genomic island PFI-12 (see Fig. 1d). This was confirmed by PCR analysis. Comparison of the elements shows a basic level of similarity. Upon insertion, each of the newly acquired elements is flanked by a 14 bp direct repeat (CCC CGG CTC CAC CA), which defines the extent of each element and maps perfectly to

the second half of the *attP* sequence in *Pseudomonas aeruginosa* PAO1 (Fiedoruk et al., 2020). One of
the 14 bp sequences is located in the genome of SBW25 and spans the last 11 bp of *tmRNA* and 3 bp
immediately downstream. The second (a duplication) is introduced by the MGE upon insertion. Each
element harbours a gene encoding a tyrosine integrase (IntY) near the 5'-end and a gene encoding AlpA,
a putative transcriptional activator of *intY* (Barcia-Cruz et al., 2024; Wang et al., 2009), at the 3'-end (see
Fig. 1e). It is likely that IntY catalyses site-specific recombination of the MGE into the genome, with the
14 bp sequence being the common core O sequence of *attB* and *attP* sites (Grindley et al., 2006; Groth
& Calos, 2004). The integrases each show ~20% divergence at the nucleotide level; *alpA* in I44 and I55
are identical, but ~20% dissimilar to *alpA* from I23. Recombination thus impacts evolution of the
elements.

Beyond the duplicated integration site and presence of an IntY plus AlpA, the three elements share a
~18 kb region of sequence similarity at the 3'-end, although I44 and I55 contain gaps (see Fig. 1e for
alignment, and Table S4 for detailed annotations). The region encodes YqaJ (I23-*orf06*, I44-*orf23*, I55-
orf28), an HTH-containing protein (I23-*orf11*, I44-*orf28*, I55-*orf32*), inovirus Gp2 (I23-*orf19*, I44-*orf31*,
I55-*orf38*), AlpA (I23-*orf20*, I44-*orf32*, I55-*orf39*), along with many open reading frames predicted to
encode proteins with no matches to either DNA or protein sequence databases. In this region elements
I23 and I55 encode a putative DNA methyltransferase (I23-*orf18* and I55-*orf37*), which is predicted to be
part of a type II restriction-modification (RM) system (Tock & Dryden, 2005) with possible phage defence
activity. However, the endonuclease component of the RM system is not present, thus leaving in doubt
roles in phage defence.

Between *intY* and the region of similarity – and accounting for length differences – are various cargo
genes that differ among the three MGEs. In the hyper-variable region, I23 encodes a known phage
defence system, the Gabija system (*orf02-orf03*) (Cheng et al., 2023; Doron et al., 2018). By contrast, I55
encodes dynamins (*orf02-orf03*) and RecBCD (*orf23-orf25*), as well as many proteins that have no
matches to databases. I44 encodes two putative phage defence systems, the classic type II restriction-
modification system (*orf02-orf03*) (Tock & Dryden, 2005) and the DISARM system (*orf05-orf10*) (Ofir et
al., 2017) . In addition, I44 encodes a ~14 kb operon (*orf11-orf20*) composed of metabolic genes
including putative hydantoinases (Ishikawa et al., 1997), a peptidase M24, and an Asp/Glu/hydantoin
racemase.

Given that the three elements were acquired during the course of serial transfer propagation
experiments, they must be capable of either autonomous, or non-autonomously transfer. A search of
encoded genes shows no evidence of those typically associated with lateral transmission, such as
machinery for conjugation, or type IV pilli. Additionally, they bear no resemblance to prophages and lack
hallmarks of phage satellites (de Sousa et al., 2023; Ibarra-Chávez et al., 2021) including capsid-encoding
genes, and *cos* sites. Moreover, interrogation of databases using recognised bioinformatic pipelines,
including ISEScan (screening for prokaryotic insertion sequences) (Xie & Tang, 2017), geNomad
(screening for viruses and plasmids) (Camargo et al., 2023), SatelliteFinder (screening for phage
satellites) (de Sousa et al., 2023) and MacSyFinder v2 (including tools TFFscan, TXSScan, CasFinder and
CONJScan, screening for Type IV filament systems, bacterial secretion systems, CRISPR-Cas systems and
conjugative systems, respectively) (Néron et al., 2023), showed no matches to any recognised mobile
element.

The three GIs represent a widespread and diverse class of element in *Pseudomonas*

Given knowledge of shared features of the three elements, we turned to whole genome sequences and asked whether *Pseudomonas* genomes harbour elements with these basic features. Search criteria included the presence of *intY* downstream of *tmRNA*, as well as the presence of an *alpA*, or a 23 bp sequence (TTC AGA TTA CCC CGG CTC CAC CA, the last 14 bp being the direct repeat) that is common to the 3'-end of each element. Searches revealed 99 elements from 17,078 *Pseudomonas* genomes. Fig. 2a shows the bioinformatic workflow and Fig. 2b depicts the relationship among the elements, constructed based on previously established methods (Low et al., 2019; Nayfach et al., 2021). Briefly, 36 ORFs that are commonly present in the entire set of MGEs were selected as core genes. For each element, the core genes were concatenated in order, with gaps allowed where core genes are absent. The tree was then inferred based on the concatenated sequences. The cladogram should be interpreted in a topological sense; the edges represent evolutionary relationships only (and not distances).

Overall, the elements are highly diverse in terms of size and gene content (Fig. 2b). The largest elements are in excess of 70 kb with the smallest being less than 20 kb. No two elements are identical; however several are highly similar. For example, the element detected in *P. rhodesiae* B21-046 (NCBI RefSeq Assembly GCF_024748755.1) is 99.96% similar at the nucleotide level to I55 acquired by SBW25. As described more fully below, these two host genomes are highly divergent. Other examples of such pairs are denoted by pink boxes in Fig. 2b.

Many elements contain homologues of RadC, YqaJ, inovirus Gp2 and HTH-containing protein (Fig. 2b, 2c). These are also present in I23, I44 and I55 but were not included in the search criteria. Together, these genes, in addition to *intY* and *alpA*, likely define core element functions.

Fig. 2c lists the rank abundance of genes that occur at least 10 times across the elements, along with functional classes and corresponding COG categories (Tatusov, 2000). In terms of variable genes, those encoding functions related to DNA replication, recombination and repair and transcription are frequently represented. Furthermore, most elements contain recognisable defence systems against bacteriophages (Fig. 2b). Not surprisingly, many genes are found as singletons with 104 of the 241 functional gene classes identified via matches to protein databases being represented just once (Table S6). Additionally, most elements harbour at least one recognisable ORFs with no matches to genes in databases (Fig. 2b).

To explore the relationship among elements and host genomes, a comparison of pairwise distances between MGEs and hosts was performed. This involved construction of phylogenetic trees for both MGEs and hosts (the latter based on 16S rDNA sequences) with the results represented in Fig. 2d. The elements shown on the X- and Y- axes are ordered based on clustering shown in Fig. 2b. The region above the diagonal represents similarity among elements, with the region below showing similarity among hosts. For example, with focus on comparison between I55 (in *P. fluorescens* SBW25) and the element in assembly GCF_024748755.1 (in *P. rhodesiae* B21-046), which is marked by a pink box: the square below the diagonal is coloured dark blue indicating that the two elements are highly similar. When the same comparison is made above the diagonal, the corresponding box is light blue indicating that the host genomes carry divergent 16S rDNA sequences. Similar patterns are evident throughout; dark blue regions below the diagonal very often correspond to divergent hosts. These incongruent patterns are indicative of the transfer of similar elements across a diverse range of host genomes as

expected for elements capable of frequent horizontal movement (Colombi et al., 2024; Koutsovoulos et
al., 2022).

I55 transfers intercellularly only in the presence of community filtrate
Knowing that the three captured elements represent a widespread, frequently-transferred, but poorly-
studied class of MGEs, we sought understanding of mechanisms of transfer and fitness consequences
(the latter is addressed in the next section) with attention focussed on I55.

By design of the serial transfer experiment, I55 is mobile (it evidently transferred from the compost-
derived filtrate into the SBW25 genome). To ask whether I55 is capable of autonomous transfer from
one SBW25 cell to another, I55 was tagged with a kanamycin resistance (Kan[R]) marker (strain
MPB39465/MPB39467). The gene encoding resistance (including promoter and a synthesised
transcription terminator) was inserted in the intergenic region between two convergently transcribed
genes, *orf08* and *orf09*, on I55. This position was chosen in order to minimise chances that insertion
disrupted element function. SBW25 carrying a chromosomally-integrated tetracycline resistance (Tet[R])
gene (strain MPB39500) was used as the recipient. Donor and recipient strains were co-cultured in liquid
medium (without shaking as in the selection experiment, see Fig. 3a) and on solid agar plates (as per a
standard conjugation assay). After 48 hours of co-culture, mating mixtures were vortexed and aliquots
added to plates containing kanamycin and tetracycline. No colonies expressing both resistance markers
were detected. This suggests that I55 is incapable of autonomous intercellular movement.

In the original serial transfer experiment, filtrate from serially-propagated compost C had been added to
SBW25 cultures. We thus considered it possible that one or more components contained within the
filtrate might be necessary to mobilise the element. To this end, the experiment above was repeated in
liquid medium, but this time with the addition of 50 μL of filtrate from compost C (Fig. 3a). On plating
mixtures to selective agar an average ~10^4 colonies (per mL) resistant to kanamycin and tetracycline
were detected (Fig. 3b).

While the result strongly implicates a component of the community filtrate as mediator in the transfer
of I55 between bacterial cells, competing hypotheses exist. For example, the Tet[R] gene in the recipient
could in principle reverse-transfer to the donor (Pei et al., 2024). Alternatively, I55 might spontaneously
transfer to another cell at a rate below detection limit, with transfer events being amplified and
detected only when community filtrate imposes selection for I55. A test of these hypotheses requires
repetition of the experiment with an engineered form of I55 incapable of transfer. We reasoned that
deletion of the integrase *intY* might prevent excision of I55 from the genome (Groth & Calos, 2004) and
thus curtail transfer. PCR, using primers flanking the insertion junctions of I55, was performed on
overnight cultures of SBW25-I55 (strain MPB39024/MPB39026). The results, shown and described in Fig.
S2, demonstrated that a fraction of I55 is in circular form, consistent with excision of the element from
the genome. The integrase was then deleted (strain MPB39132/MPB39136) and the PCR analysis
repeated. Results show that SBW25-I55Δ*intY* is incapable of excising from the genome and generating a
circular form.

The transfer experiment was repeated using SBW25-I55Δ*intY*-Kan[R] (strain MPB39565/MPB39568), with
addition of culture filtrate from compost C. While on average ~10^2 colonies (per mL) resistant to both
antibiotics were detected, the number was significantly reduced ($p < 0.0001$) compared to the case
where SBW25-I55-Kan[R] was the donor (Fig. 3b). This not only confirms that the occurrence of double-

resistance cells was not due to artifacts, but also implies that the circularisation of I55 underpins
intercellular transfer and / or integration.

Next, we asked whether donor-recipient cell-cell contact is necessary for the transfer of I55. To test this,
the transfer process was broken into two steps (Fig. 3c). In the first step, SBW25-I55-Kan^R with filtrate
from compost C was cultured for 48 hrs and the filtrate collected. In the second step, recipient SBW25-
Tet^R was grown with addition of the filtrate obtained from the first step. The mixture was plated as
previously. On average, more than 10^6 colonies (per mL) resistant to both antibiotics were detected (Fig.
3d), indicating that transfer of I55 is independent of cell-cell contact. To test the significance of the
tyrosine integrase under this two-step transfer regime, SBW25-I55Δ*intY*-Kan^R was used in place of
SBW25-I55-Kan^R, and the assay repeated. The density of double-resistant colonies decreased to ~10^4
colonies (per mL) (*p* = 0.0152), with this 100-fold difference consistent with the fold change in the one-
step transfer regime (Fig. 3b, 3d).

To ask which step in the transfer process the compost C filtrate is necessary, we repeated the two-step
transfer regime, but with compost C filtrate added to the culture at the second step. No double-resistant
colonies were detected (Fig. 3d), indicating that transfer of I55 did not occur (or if it did then it remained
below the detection limit). This suggests that in order for I55 to be transferred, the compost filtrate
must be co-cultured with the donor. Therefore, I55 appears to require some component in the
community filtrate that mediates transfer to recipient cells (Fig. 3e).

The identity of the transfer-facilitating component of compost filtrate is unknown, but it is reasonable to
suspect the involvement of phages. Five phage isolates were purified from the compost C filtrate and
individually added to the intercellular transfer assay (Fig. 3a) in place of the community filtrate, but no
double-resistant cells were detected and hence no evidence of transfer of I55. Additionally, community
filtrate from composts B, C and D were amplified and sequenced. This was done on the off-chance that
we might identify I23, I44 or I55 as a co-integrate with some other element. The sequences of I23 and
I55 were not covered with sufficient quality by reads of any of the samples. The sequence of I44,
however, was covered by reads from compost B filtrate, which is the filtrate in the selection experiment
from which isolates carrying I44 were detected. Approximately 250 reads that partially map to the left
or right ends of I44 sequence were extracted and then mapped to the circular version of I44 sequence.
All such reads matched the circularisation junction. This suggests that most of I44 in compost B filtrate
was in circular form, although rolling-circle replication during amplification of DNA could cause an over-
estimation of the proportion of circular DNA. There was no evidence that I44 in the filtrate formed a co-
integrant with some other element.

I55 confers a fitness advantage in the presence of community filtrate, partially due to defence
against phages
While knowledge of the mechanism(s) of I55 transfer are unclear, experiments to determine fitness
consequence are readily achievable. The fitness of SBW25 with and without I55 were determined by
direct competition taking advantage of fluorescent reporters allowing the two types to be distinguished
by flow cytometry (Fig. 4a). However, because I55 can transfer in the presence of community filtrate (a
potentially confounding factor), fitness effects were determined using SBW25-I55Δ*intY*. Moreover, to
ensure the two competing strains have identical genetic backgrounds, ancestral SBW25 (strain
MPB39379/MPB39459) was re-constructed by deleting I55 from SBW25-I55Δ*intY*. Compared to the wild
type, I55 confers a significant fitness advantage (~6%) in the presence of compost C filtrate (*p* = 0.0104),

but is slightly costly (~2%) in the absence of the filtrate (p = 0.0033) (Fig. 4b). Addition of community
filtrate significantly increased the fitness benefit of I55 (p = 0.0006).

I55 encodes a putative phage defence system (*orf37*), and the community filtrate is rich in phages
(Quistad et al., 2020). We therefore hypothesised that benefit of I55 stems from defence against
phages. This hypothesis was tested by determining the ability of SBW25 and SBW25-I55Δ*intY* to resist
attack by phages. Compost C filtrate was added to both hosts, with the number of plaque forming units
on SBW25-I55Δ*intY* being significantly reduced compared to SBW25 (p = 0.0080) (Fig. 4d). Moreover, the
plaques on lawns of SBW25-I55Δ*intY* appeared visually opaque compared to those on SBW25 (see Fig.
4c), suggesting that infection and cell lysis is less severe when cells carry I55. Together, this result
indicates that I55 confers non-perfect, but significant protection against phages present in the
community filtrate.

As the community filtrate stock is limited, we purified two phages (P1 and P2) from the plaque assay of
community filtrate. Phage P1 was isolated by picking a random plaque from the SBW25 lawn followed
by iterative purification on SBW25 as the host. Phage P2 was isolated in the same way except the host
for the initial plaque assay was SBW25-I55Δ*intY*. The fluorescent competition assay and plaque assay
were repeated with the two phage isolates replacing the community filtrate, and the assay results were
qualitatively reproduced (see Fig. 4e, 4f). This indicates that phage defence partially explains the fitness
benefit of I55.

Discussion

HGT mediated by MGEs shapes the diversity and function of microbial communities, but recent studies indicate that there is much left to discover concerning both the effectors of transfer and the range of functional genes subject to horizontal movement. Motivating the work reported here, was a previous study (Quistad et al., 2020) in which frequent exposure of pooled supernatant from compost communities to fresh hosts caused a substantive fraction of community DNA to partition into the pool of horizontally disseminated material. Comparative metagenomics provided evidence of genes of ecological relevance and in some instances phages, plasmids and other known MGEs were identified, but in the absence of ability to capture unknown elements in readily culturable hosts, progress is severely curtailed.

Here we simplified our previous strategy, retaining the complexity of culture filtrate from compost communities, but exposing the diversity of elements contained within the filtrate to a single focal, culturable and genetically tractable strain of *P. fluorescens.* The focal bacterium acquired three new MGEs in a short period of time, each integrated immediately downstream of *tmRNA*. The I23, I44 and I55 family of elements (hereafter the I55 family) lack characteristics of recognised MGEs, with a range of purpose-built bioinformatic pipelines (Camargo et al., 2023; de Sousa et al., 2023; Néron et al., 2023; Xie & Tang, 2017) failing to detect similarities. While tempting to suggest that the elements are phage satellites (de Sousa et al., 2023), absent are genes encoding primases, genes for viral packaging and *cos* sites, which are regarded as defining features of this class.

Bioinformatic searches for similar elements defined by minimal set of criteria including the target site, 3'-end sequence and integrase revealed ~100 within the genus *Pseudomonas*, however, the class is more diverse. Relaxation of the strict requirement for the flanking sequence (*attP*) to match the *tmRNA* sequence from *Pseudomonas* shows that elements with essential features of I55 are widespread. In fact, the broadest manifestation of the class appears to be that of GIs, many of which insert at *tmRNA* (or *tRNA*) genes, are flanked by direct repeats, encode tyrosine recombinases, but lack genes known to be required for autonomous transfer. The so-named "phage defense elements" recently described by Hussain et al. (Hussain et al., 2021) are a case in point. Islander, a bioinformatic tool developed to search for integrases downstream of *tmRNA* or *tRNA* genes (Hudson et al., 2015) shows presence of I55-like elements (GIs) across eubacteria and archaea.

A particularly notable example is the 41 kb *Vibrio* Pathogenicity Island-1 (VPI-1) that encodes the toxin-co-regulated pilus (TCP), which is the receptor for the cholera toxin-encoding phage CTXΦ. Acquisition of VPI-1, followed by entry of CTXΦ, converts environmental *V. cholerae* strains into toxigenic strains (Faruque & Mekalanos, 2003, 2012). While it is evident from comparative genomics that VPI-1 horizontally transferred (Karaolis et al., 1999; O'Shea & Boyd, 2002), attempts to demonstrate mobility have proven controversial (Faruque et al., 2003) with the current consensus being that VPI-1 is incapable of autonomous movement. Armed with knowledge learned from acquisition of members of the I55 family by SBW25, combined with studies of I55 transfer, and the fact that VPI-1 shares all the hallmarks that define this new class, we suggest that VPI-1 is a non-autonomous MGE of the I55 family, whose movement is facilitated by a secondary factor likely common in the environment naturally occupied by environmental *V. cholerae*. Perhaps this discovery ignites attempts to define the helper element, thus paving the way for development of strategies that curtail opportunity for the agent to facilitate transfer of VPI-1.

Capture of I55 (and family members) by SBW25 combined with genetic tractability of the focal
bacterium facilitated studies of the mechanism of transfer. While much remains to be discovered,
unequivocal demonstration that I55 is incapable of autonomous transfer is an important negative
finding. Indeed, this result led us to consider the possibility that a secondary element present in
compost filtrate may act as a helper. That transfer did occur in the presence of compost filtrate, was
largely dependent on IntY (necessary for excision and circularisation of I55), and did not require cell-to-
cell contact, is informative. Obvious secondary factors are phages, but purified phages have thus far
failed to mediate transfer. Nonetheless, we suggest that I55 family elements – including the huge
diversity of similar GIs suspected of movement, but for which direct evidence of transfer is lacking – are
likely to depend on the process of lateral transduction (Chen et al., 2018; Fillol-Salom et al., 2021;
Humphrey et al., 2021) mediated by *pac*-type phages that package DNA using the headful mechanism.

A benefit to working on I55 in a genetically tractable host has been understanding of fitness
consequence upon acquisition. That I55 confers a significant advantage in the presence of compost
filtrate (but a minor cost in its absence) offers an explanation for why it was possible to detected these
elements in SBW25 given sequence analysis of so few derived colonies. Demonstration of fitness
advantage in the presence of purified phages supports the hypothesis that the advantage stems in part
from benefits derived from phage defence systems. Interestingly, bioinformatic predictors (Tesson et al.,
2022) failed to detect complete phage defence systems in I55, however orthologs of dynamins (Guo et
al., 2022) and RecBCD (Dillingham & Kowalczykowski, 2008) are present (see Table S4). More generally,
this supports the view that MGEs are a rich source of novel phage defence systems (Hussain et al., 2021;
Johnson et al., 2023; Rousset et al., 2022).

Experimental detection of I55 family elements, and with these elements being instrumental in discovery
of a plethora of similar MGEs across *Pseudomonas* and beyond, likely defines just a small quota of the
yet-to-be discovered mediators of HGT. Repetition of the experiment with filtrates from different
environments, or different hosts, is likely to result in further breakthroughs. This gives substance to the
notion that microbial communities, replete with frequently moving MGEs (and associated genetic
cargoes), might be shaped by a process akin to sex as understood in eukaryotic populations (Rainey &
Quistad, 2020). To test this hypothesis, it will be necessary to go beyond knowledge that MGEs move
ecologically relevant DNA, to a more complete description of the nature and diversity of mobilising
elements, including factors affecting capacity to capture and recombine cargo genes. Equally important
will be development of new approaches to directly measure the flux of horizontally disseminated DNA.
Finally, simplicity of the strategy outlined here stands as a useful and general approach for discovery of
new MGEs, and the functional genes to which these elements become linked.

Materials and Methods

Strains and growth conditions

All bacterial strains used in this study are listed in Table S1. All strains were preserved in -80 °C with 30% glycerol saline.

Isolation of foreign elements, plasmid construction, strain construction: Unless otherwise stated, *P. fluorescens* SBW25 strains were cultured in Luria broth (LB) at 28 °C, while *E. coli* strains were cultured in LB (with antibiotics where maintenance of plasmids was required) at 37 °C, all shaking at 220 rpm for 16-24 hrs.

Experimental evolution, intercellular transfer assay and fluorescent fitness assay: Unless otherwise stated, each pre-conditioning and incubation was performed in the absence of shaking at 28 °C, in 5 mL freshly made M9-glucose medium in a 15 mL Falcon tube. The M9-glucose medium was M9 salt solution supplemented with 2 mM $MgSO_4$, 0.1 mM $CaCl_2$ and 0.2% w/v D-glucose.

Plaque assay and bacteriophage isolation: Unless otherwise stated, bacterial host strains were cultured in S-LB medium (LB medium supplemented with 10 mM $MgCl_2$ and 5 mM $CaCl_2$) at 28 °C, shaking at 220 rpm for 16-24 hrs until stationary phase. Exponential phase bacterial culture was obtained by 1/25 diluting stationary phase culture in fresh S-LB medium and then shaking incubate at 28 °C, 220 rpm for 2 hrs. The semi-soft agar (SSA) used consisted of S-LB medium and 0.4% agar.

Bacterial culture was plated on respective growth medium supplemented with 1.5% agar. Unless otherwise stated, agar plates were incubated at 28 °C. Antibiotics were applied as per the following concentrations: kanamycin (Kan, 50 μg/mL), nitrofurantoin (100 μg/mL), tetracycline (Tet, 12.5 μg/mL), *Pseudomonas* CFC selective supplement (Millipore, as per the manual).

Compost community sampling, storage and retrieval

Three heaps of garden compost, named B, C and D, were sampled in Plön, Germany (54°9'44"N, 10°25'17"E), from May to August 2022. Upon collection, the samples were immediately transferred to laboratory and processed as following. For each sample, 30 g compost was mixed with 150 mL M9 salt solution, shaken vigorously for ~30 sec, and left static for ~5 min. The supernatant was distributed into 1 mL aliquots, to each of which 350 μL glycerol saline was added. The aliquots were frozen in -80 °C for storage.

To retrieve the compost wash, one aliquot from the frozen stock was kept on ice for ~10 min. After thawing, the mixture was pelleted by centrifuging at 15000 rcf for 1 min, resuspended in 1 mL M9 salt solution, and then pelleted by centrifuging again. The washed pellet was then resuspended in 500 μL M9 salts solution to inoculate microcosms.

Community filtrate (fixed) generation, storage and retrieval

Initially, 500 μL washed suspension from one compost wash aliquot was inoculated into 100 mL M9-glucose medium in a 500 mL microcosm. The microcosm was incubated without shaking at 28 °C for 7 days. At the end of incubation period, the whole culture was filtered with a 0.2 μm filter (Whatman). The filtrate was distributed into 600 μL aliquots. Aliquots were directly frozen at -80 °C for storage.

To retrieve the community filtrate, one aliquot from the frozen stock was kept at room temperature for
~10 min. After thawing, the aliquot was directly used to inoculate microcosms. Unless otherwise stated,
50 μL community filtrate was added to each microcosm.

As a negative control, *P. fluorescens* SBW25 was used in place of the washed suspension, and otherwise
followed exactly the same generation, storage and retrieval protocols to obtain the SBW25 filtrate to
inoculate microcosms.

Experimental evolution
Co-culture regime
Parallel lines of SBW25, cocultured with the compost community, were serially propagated. In detail,
SBW25-RFP-Gen^R (strain MPB28806) and 90 μL washed suspension from compost C were inoculated.
There were three parallel lines, with one control line where no washed suspension was added. For each
line, an 1/20 transfer was made every 24 hrs, for a total of five transfers. At each transfer, a dilution
series of the culture was spread plated on LB + CFC + Gen plates, which were then cultivated for 24 hrs.
The number of RFP colonies were counted and recorded.

Fixed filtrate regime
Parallel lines of SBW25 were serially propagated, with aliquots of filtrate from the frozen stock added at
each transfer. In detail, filtrate from communities B, C and D, and from SBW25, were separately used.
For each type of filtrate, there were five parallel lines of SBW25-GFP-Kan^R (strain MPB29284) and five
parallel lines of SBW25-RFP-Kan^R (strain MPB29300). For each line, an 1/100 transfer was made every 48
hrs, for a total of seven transfers. At each transfer, 50 μL filtrate was added to a fresh microcosm, and
aliquots from the old microcosm were stored at -80 °C.

Live filtrate regime
Here, parallel lines of SBW25 were serial propagated, with aliquots of filtrate from independently
propagated communities or SBW25 added at each transfer. In detail, 200 μL washed suspension from
compost C wash was added and transferred 1/100 every 48 hrs, yielding fresh filtrate by filtering 1mL of
the culture through a 0.2 μm filter (Whatman). As a negative control, *P. fluorescens* SBW25 was used in
place of the washed suspension, but otherwise the same protocol was followed to obtain SBW25
filtrate. At the same time, for each type of filtrate, there were four parallel lines of SBW25-GFP-Kan^R
(strain MPB29284) and four parallel lines of SBW25-RFP-Kan^R (strain MPB29300). For each line, a 1/100
transfer was made every 48 hrs, for a total of 14 transfers. At each transfer, 50 μL fresh filtrate was
added to the new microcosm, and a sample from the old microcosm were stored at -80 °C.

PCR and gel electrophoresis
465 All PCR primers used in this study are listed in Table S3. PCR was done with NEB Q5 polymerase, with
466 either extracted DNA or bacterial suspension as templates. Annealing temperatures and extension time
467 was determined as per NEB Tm Calculator (version 1.16.6). Unless otherwise stated, 35 cycles were
468 performed. Gel electrophoresis was performed for 5 μL PCR products with 1x DNA loading dye, in 1%
469 agarose gels.

Detection, isolation and annotation of foreign elements
For each line, frozen culture at the last transfer was serially diluted and plated out on LB plates. For the
live-filtrate regime, 2 to 3 colonies were picked and separately cultured overnight, from which aliquots
were frozen and DNA extracted with Qiagen DNeasy UltraClean Microbial kit. DNA samples were

sequenced by Illumina NextSeq for at least 300x coverage. For fixed-filtrate regime, 24 colonies were
randomly picked and cultured up overnight in 300 µL LB in 96-deep-well plates. Aliquots from each of
the 24 cultures were mixed in equal proportion and then DNA extractions were performed with Qiagen
DNeasy UltraClean Microbial kit. The leftovers in 96-deep-well plates were stored at -80 °C. DNA
samples were Illumina short-read sequenced (read length 150 bp) by Eurofins Scientific SE for at least
300x coverage.

Bioinformatic analysis of sequencing data was done in Geneious Prime 2022.2.1
(https://www.geneious.com). For each sample, raw reads were trimmed by BBDuk and mapped to *P.*
fluorescens SBW25 reference genome (Fortmann-Grote et al., 2023) under the default settings. The
unmapped reads were saved and then *de novo* assembled. The vast majority of contigs were short (< 1
kb) and apparently artefacts of sequencing technology. However, in some samples, long (> 10 kb) and
good quality contigs were obtained. To test if the contig inserted into the genome as a foreign element,
the first 50 bp and last 50 bp of the contig were blasted against the reference genome. If the two
regions mapped, manual comparison were used to confirm the insertion and its locus.

To confirm the insertion of the foreign element, two primer pairs were designed, flanking either the left
or the right junction of the insertion. For fixed-filtrate regime, the two primer pairs were additionally
used to exclude the cultured isolates (out of the 24 picked) that carried the foreign element. Such
isolates were then routinely whole-genome sequenced by Eurofins Scientific SE and Variations / SNPs
calling was subsequently performed. Table S5 shows the list of mutations detected in each isolate.

The three foreign elements were screened for open reading frames (ORFs) by Prokka (Seemann, 2014) in
KBase (Arkin et al., 2018). The ORFs were then annotated by Pfam in InterProScan (Jones et al., 2014).
The elements were also screened by DefenseFinder (Tesson et al., 2022) for potential phage defense
systems (only HMM-only hits were retained). Alignment of the 3 foreign elements was done by Easyfig
(M. J. Sullivan et al., 2011) based on blastn under the default settings.

Detection and analysis of similar elements in databases
The nucleic acid sequences of the *tmRNA*, *intY* (from the element I55) and the 23 bp sequence at the 3'-
end were compared to representative genomes in GTDB-tk (version R214) (Chaumeil et al., 2020).
Sequence comparisons revealed that 35 representative genomes contain all three sequences, and they
all belong to the family *Pseudomonadaceae*. This motivated a more focused and thorough search within
this family. Therefore, the nucleic acid sequences of the *tmRNA*, *intY* (from the element I55) and the 23
bp sequence at the 3'-end were compared to all available genomic sequences (17,078 in total) of strains
belonging to the family *Pseudomonadaceae*. Sequence comparisons revealed that 136 genomes in the
family *Pseudomonadaceae* contain all three sequences. Those 136 genomes were then subject to gene
prediction by Prodigal (v2.6.3) (Hyatt et al., 2010) and functional annotation by eggNOG-mapper (v2)
(Cantalapiedra et al., 2021). Based on annotations, 99 genomes fulfilled the following criteria: (1)
contained an annotated tmRNA; (2) contained an annotated tyrosine integrase (IntY); (3) contained at
least one of an annotated AlpA or the 23 bp sequence. These 99 genomes are derived from 67 species,
all in the genus *Pseudomonas*. In each genome, the MGEs were isolated by extracting the sequence from
the 3'-end of the *tmRNA* to the 3'-end of the 23 bp sequence or the 3'-end of *alpA*. All MGEs were
screened by DefenseFinder (Tesson et al., 2022) for potential phage defense systems.

To establish evolutionary relationships of the 99 detected MGEs and I23, I44 and I55, a cladogram of the
102 MGEs was built based on previously established methods (Low et al., 2019; Nayfach et al., 2021).
Briefly, 36 ORFs that are commonly present in the MGEs were selected as marker genes. For each MGE,
the marker genes were concatenated in order, with a gap introduced in cases where a marker gene was
absent. The tree of the MGEs was then inferred based on the concatenated sequences.

To assess host-MGE relationships, the 16S rDNA sequences of the hosts of the 102 MGEs were
extracted. Among them, 5 host genomes lack detectable 16S rDNA regions and hence were excluded
from the following analysis. For the 97 remaining genomes, phylogenetic trees were built separately
based on the 16S rDNA sequences by Clustal (v2.1) (Sievers et al., 2011), and based on the MGE
sequences by the same method as before (Low et al., 2019; Nayfach et al., 2021). The two trees were
converted to distance matrices in R (version 4.1.2), with rows clustered according to MGE distances.

Plasmid construction

All plasmids used in this study are listed in Table S2. Plasmid pUIsacB (plasmid MPB35522) backbone
was amplified by PCR. SBW25 genomic fragments flanking the region of interest, where a mutagenesis,
knock-out or knock-in was to be performed, were amplified by PCR. In particular, the Kan^R fragment
used to tag I55 was produced by cloning the Kan^R gene from plasmid pMRE-Tn7-152 (plasmid
MPB27708) followed by a synthesised transcription terminator (CCA ATT ATT GAA CAC CCT TCG GGG
TGT TTT TTT GTT TCT GGT CTC CC). Primers were designed with a 5' overhang to facilitate assembly.
Backbone and fragments were assembled (with NEBuilder HiFi DNA Assembly Master Mix) and
transformed into *E. coli* Top10 competent cells. Successful transformants were selected on LB + Tet
plates, and confirmed by PCR and Sanger sequencing across the construct.

Strain construction

Two-step allelic replacement (2SAR)

The method is adapted from Hmelo et al. (Hmelo et al., 2015) and extensively described by Farr et al.
(Farr et al., 2024). Briefly, recipient (*P. fluorescens* SBW25), donor (*E. coli* carrying the plasmid used for
allelic exchange) and helper (*E. coli* carrying plasmid pRK2013, plasmid MPB35623) strains were cultured
overnight until stationary phase. Then, 1 mL recipient culture was heat-shocked at 42 °C for 20 min, and
subsequently mixed with 500 µL donor and 500 µL helper culture. The mixture was pelleted and
resuspended in 100 µL LB, which was then spotted onto an LB plate for incubation overnight. The
cultivated cell layer was then plated out on an LB + nitrofurantoin + Tet plate and incubated overnight to
select for trans-conjugates. Single trans-conjugates were then picked and streaked onto TYS10 plates,
which were subsequently incubated overnight. Finally, PCR and Sanger sequencing were used to screen
for the desired mutants.

Fluorescent tagging

The method is adapted from Schlechter et al. (Schlechter et al., 2018). Briefly, recipient (*P. fluorescens*
SBW25) and donor (*E. coli* carrying the plasmid pMRE-Tn7-152 or pMRE-Tn7-155, MPB27708 or
MPB27709, respectively) strains were cultured overnight until stationary phase. Then, 1 mL recipient
culture was heat-shocked at 42 °C for 20 min, and subsequently mixed with 1 mL donor culture. The
mixture was pelleted and resuspended in 100 µL LB, which was then spotted onto an LB plate for
incubation overnight. The cultivated cell layer was then plated out on an LB + nitrofurantoin + Kan +
0.1% w/v arabinose plate and incubated overnight to select for trans-conjugates. Single trans-conjugates
were picked, and integration of the fluorescent tag was then confirmed by PCR.

Preparation of SBW25-I55 and SBW25 (reconstructed) for assays

DNA sequencing of the evolved isolate carrying I55 (strain MPB33357) revealed, in addition to presence of the mobile element, two SNPs elsewhere in the genome (see Table S5) that had arisen during the course of serial propagation. Additionally, this isolate also harboured the Tn*7*-152 cassette. While there was no reason to suspect any connection to I55, these SNPs were reverted to the ancestral SBW25 sequence, and Tn*7*-152 removed, by site-directed mutagenesis using the 2SAR strategy described above. Prior to these mutagenesis steps, two independent colonies were picked from the evolved isolate and treated as biological replicates in all subsequent genetic manipulations. The resulting genotype, SBW25-I55 (strain MPB39024/MPB39026) is in principle isogenic with SBW25 (strain MPB29212). However, to be more certain of this fact, the ancestral SBW25 genotype used in the fitness assays and plaque assays was re-generated from SBW25-I55 by removal of the element, giving reconstructed SBW25 (strain MPB39379/MPB39459).

Intercellular transfer assay

One-step regime: The donor, a SBW25 strain with a Kan^R-tagged I55 (with or without *intY*, strains MPB39465/MPB39467 or MPB39565/MPB39568, respectively), and the recipient, a SBW25 strain with a Tet^R marker (strain MPB39500), were cultivated separately for 48 hrs for pre-conditioning. Then 25 µL of each culture and 50 µL of community filtrate from frozen stocks (autoclaved or not) were inoculated into the same microcosm and then cultivated for a further 48 hrs. A dilution series of the culture was then spread plated on LB + Kan + Tet plates, which were then cultivated for 24 hrs. The number of colonies were counted and recorded.

Two-step regime: For the first step, the donor, a SBW25 strain with a Kan^R-tagged I55 (with or without *intY*, strains MPB39465/MPB39467 or MPB39565/MPB39568, respectively), were cultivated for 48 hrs for pre-conditioning. Then 50 µL of the donor culture, with or without 50 µL of community filtrate from frozen stocks, was added to microcosms and then cultivated for a further 48 hrs. A stock of donor-culture filtrate was obtained by filtering the culture through a 0.2 µm filter (Whatman). The filtrate was distributed into 100 µL aliquots. Aliquots were directly frozen in -80 °C for storage. To retrieve the donor-culture filtrate, one aliquot from the frozen stock was kept at room temperature for ~10 min.

For the second step, the recipient, a SBW25 strain with a Tet^R marker (strain MPB39500), was cultivated for 48 hrs for pre-conditioning. Then 50 µL of the recipient culture and 50 µL of the donor-culture filtrate from frozen stocks (obtained in the first step), with or without 50 µL of community filtrate from frozen stocks, were added into the same microcosm and then cultivated for a further 48 hrs. A dilution series of the culture was then spread plated on LB + Kan + Tet plates, which were then cultivated for 24 hrs. The number of colonies were counted and recorded.

Conjugation regime: The donor, a SBW25 strain with Kan^R-tagged I55 (strain MPB39465/MPB39467), recipient, a SBW25 strain with a Tet^R marker (strain MPB39500), and helper, *E. coli* carrying plasmid pRK2013 (plasmid MPB35623), were cultured overnight until stationary phase. Then, 1 mL recipient culture was heat-shocked at 42 °C for 20 min, and subsequently mixed with 500 µL donor and 500 µL helper culture. The mixture was pelleted and resuspended in 100 µL LB, which was then spotted onto an LB plate for incubation overnight. The cultivated cell layer was then plated out on an LB + Kan + Tet plate and incubated overnight to select for trans-conjugates.

Community filtrate sequencing and analysis

To sequence each community filtrate, a 600 μL aliquot of filtrate was extracted with Qiagen DNeasy PowerSoil Pro kit. The concentrations of eluted DNA were in general lower than ~1 ng/μL and insufficient for subsequent sequencing. To amplify the DNA, 10 μL of eluted DNA was amplified with Cytiva GenomiPhi V3 Ready-To-Go DNA Amplification kit. The amplified DNA was extracted again with Qiagen DNeasy PowerSoil Pro kit. The DNA samples Illumina short-read sequenced (read length 150 bp) by Eurofins Scientific SE for ~100M paired-end reads.

Bioinformatic analysis of sequencing data was done in Geneious Prime 2022.2.1 (https://www.geneious.com). For each sample, raw reads were trimmed by BBDuk and separately mapped to the sequences of I23, I44 and I55.

Fluorescent fitness assay

The protocol was adapted from Barnett et al. (Barnett et al., 2024). Two competitor strains, one with a GFP tag and the other with an RFP tag, were cultivated separately for 48 hrs for pre-conditioning. Then 25 μL of each culture, and 50 μL of community filtrate or 1 μL of phage lysate (see following sections) from frozen stocks were inoculated into the same microcosm, which was then cultivated for a further 48 hrs. At the beginning and end of the second incubation period, the ratios of GFP to RFP cells were determined by flow cytometry (MACSQuant VYB), denoted by r_0 and r_1, and total densities of cells in the microcosm were determined by spread plating on LB plates, denoted by n_0 and n_1. The relative fitness of GFP to RFP cells is the ratio of Malthusian parameters (Lenski et al., 1991), given by

$$1 + \frac{\log\left(\frac{r_1}{r_0}\right)}{\log\left(\frac{n_1(1 + r_0)}{n_0(1 + r_1)}\right)}.$$

Plaque assay

To determine the phage titre, a dilution series of community filtrate or phage lysate (see following sections) was made. Then, 50 μL diluted material and 400 μL bacterial host overnight culture were mixed into 4 mL SSA (kept at 55 °C). The mixture was immediately poured onto a LB plate. The plaque counts were determined after ~40 hrs incubation. Pictures of the plates were taken with Bio-Rad ChemiDoc Imager, under the colorimetric setting with exposure time 0.1 sec.

Bacteriophage isolation and storage

To purify phages from the community filtrate, a plaque assay (Clokie & Kropinski, 2009) using the filtrate was performed. A single isolated plaque was then picked, on which a plaque assay was performed. This was repeated at least three times. Finally, a single isolated plaque from the last assay was picked and amplified in exponential-phase bacterial host culture for ~3 hrs. The obtained phage lysate was chloroform extracted and stored in -80 °C with 50% glycerol saline. For the lysate extraction control, an exponential-phase bacterial host culture without added phages was extracted and stored in the same way.

Statistical analysis

All experiments were done for at least 4 independent biological replicates. Statistical analyses were performed with GraphPad Prism (v. 10.1.2). Welch's *t*-test and Mann-Whitney test were performed for

633   parametric and non-parametric pairwise comparisons, respectively. In each plot, *p* values are indicated
634   above the bars: *p* > 0.05 not significant (ns), *p* ≤ 0.05 (∗), *p* ≤ 0.01 (∗∗), p ≤ 0.001 (∗∗∗), p ≤ 0.0001 (∗∗∗∗).

Miscellaneous

Data availability

Supporting information, including sequencing raw reads, bioinformatic codes, gel images and plaque assay images, will be deposited in Edmond (https://edmond.mpg.de/).

Author contributions

Y.Z., A.D.F., D.W.R. and P.B.R. designed research; Y.Z. and Y.M. performed research; all authors analysed data; and Y.Z. and P.B.R. wrote the paper; all authors approved the final version.

Acknowledgements

We acknowledge the Max Planck Society for generous core funding. We thank members of Department of Microbial Population Biology (past and present) for support, technical assistance and discussion.

Competing interests

The authors declare no competing interest.

References

Arkin, A. P., Cottingham, R. W., Henry, C. S., Harris, N. L., Stevens, R. L., Maslov, S., Dehal, P., Ware, D.,

Perez, F., Canon, S., Sneddon, M. W., Henderson, M. L., Riehl, W. J., Murphy-Olson, D., Chan, S.

Y., Kamimura, R. T., Kumari, S., Drake, M. M., Brettin, T. S., … Yu, D. (2018). KBase: The United

States Department of Energy Systems Biology Knowledgebase. *Nature Biotechnology*, *36*(7),

566–569. https://doi.org/10.1038/nbt.4163

Arnold, B. J., Huang, I.-T., & Hanage, W. P. (2022). Horizontal gene transfer and adaptive evolution in

bacteria. *Nature Reviews Microbiology*, *20*(4), 206–218. https://doi.org/10.1038/s41579-021-

00650-4

Barcia-Cruz, R., Goudenège, D., Moura de Sousa, J. A., Piel, D., Marbouty, M., Rocha, E. P. C., & Le Roux,

F. (2024). Phage-inducible chromosomal minimalist islands (PICMIs), a novel family of small

marine satellites of virulent phages. *Nature Communications*, *15*(1), 664.

https://doi.org/10.1038/s41467-024-44965-1

Barnett, M., Zeller, L., & Rainey, P. B. (2024). Experimental evolution of evolvability. *BioRxiv*.

https://doi.org/10.1101/2024.05.01.592015

Camargo, A. P., Roux, S., Schulz, F., Babinski, M., Xu, Y., Hu, B., Chain, P. S. G., Nayfach, S., & Kyrpides, N.

C. (2023). Identification of mobile genetic elements with geNomad. *Nature Biotechnology*.

https://doi.org/10.1038/s41587-023-01953-y

Cantalapiedra, C. P., Hernández-Plaza, A., Letunic, I., Bork, P., & Huerta-Cepas, J. (2021). eggNOG-

mapper v2: Functional annotation, orthology assignments, and domain prediction at the

metagenomic scale. *Molecular Biology and Evolution*, *38*(12), 5825–5829.

https://doi.org/10.1093/molbev/msab293

Chaumeil, P.-A., Mussig, A. J., Hugenholtz, P., & Parks, D. H. (2020). GTDB-Tk: A toolkit to classify

genomes with the Genome Taxonomy Database. *Bioinformatics*, *36*(6), 1925–1927.

https://doi.org/10.1093/bioinformatics/btz848

Chen, J., Quiles-Puchalt, N., Chiang, Y. N., Bacigalupe, R., Fillol-Salom, A., Chee, M. S. J., Fitzgerald, J. R.,

& Penadés, J. R. (2018). Genome hypermobility by lateral transduction. *Science*, *362*(6411), 207–

212. https://doi.org/10.1126/science.aat5867

Cheng, R., Huang, F., Lu, X., Yan, Y., Yu, B., Wang, X., & Zhu, B. (2023). Prokaryotic Gabija complex senses

and executes nucleotide depletion and DNA cleavage for antiviral defense. *Cell Host & Microbe*,

31(8), 1331-1344.e5. https://doi.org/10.1016/j.chom.2023.06.014

Clokie, M. R. J., & Kropinski, A. M. (Eds.). (2009). *Bacteriophages* (Vol. 501). Humana Press.

https://doi.org/10.1007/978-1-60327-164-6

Colombi, E., Bertels, F., Doulcier, G., McConnell, E., Pichugina, T., Sohn, K. H., Straub, C., McCann, H. C.,

& Rainey, P. B. (2024). Rapid dissemination of host metabolism–manipulating genes via

integrative and conjugative elements. *Proceedings of the National Academy of Sciences*, *121*(11),

e2309263121. https://doi.org/10.1073/pnas.2309263121

Colombi, E., Straub, C., Künzel, S., Templeton, M. D., McCann, H. C., & Rainey, P. B. (2017). Evolution of

copper resistance in the kiwifruit pathogen *Pseudomonas syringae* pv. *Actinidiae* through

acquisition of integrative conjugative elements and plasmids. *Environmental Microbiology*,

19(2), 819–832. https://doi.org/10.1111/1462-2920.13662

de Sousa, J. A. M., Fillol-Salom, A., Penadés, J. R., & Rocha, E. P. C. (2023). Identification and

characterization of thousands of bacteriophage satellites across bacteria. *Nucleic Acids*

Research, *51*(6), 2759–2777. https://doi.org/10.1093/nar/gkad123

Dillingham, M. S., & Kowalczykowski, S. C. (2008). RecBCD enzyme and the repair of double-stranded

DNA breaks. *Microbiology and Molecular Biology Reviews*, *72*(4), 642–671.

https://doi.org/10.1128/MMBR.00020-08

Dmitrijeva, M., Tackmann, J., Matias Rodrigues, J. F., Huerta-Cepas, J., Coelho, L. P., & von Mering, C.

(2024). A global survey of prokaryotic genomes reveals the eco-evolutionary pressures driving

horizontal gene transfer. *Nature Ecology & Evolution*, *8*(5), 986–998.

https://doi.org/10.1038/s41559-024-02357-0

Doolittle, W. F., & Inkpen, S. A. (2018). Processes and patterns of interaction as units of selection: An

introduction to ITSNTS thinking. *Proceedings of the National Academy of Sciences*, *115*(16),

4006–4014. https://doi.org/10.1073/pnas.1722232115

Doron, S., Melamed, S., Ofir, G., Leavitt, A., Lopatina, A., Keren, M., Amitai, G., & Sorek, R. (2018).

Systematic discovery of antiphage defense systems in the microbial pangenome. *Science*,

359(6379), eaar4120. https://doi.org/10.1126/science.aar4120

Farr, A. D., Vasileiou, C., Lind, P. A., & Rainey, P. B. (2024). An extreme mutational hotspot in *nlpD*

depends on transcriptional induction of *rpoS*. *BioRxiv*.

https://doi.org/10.1101/2024.07.11.603030

Faruque, S. M., & Mekalanos, J. J. (2003). Pathogenicity islands and phages in *Vibrio cholerae* evolution.

Trends in Microbiology, *11*(11), Article 11. https://doi.org/10.1016/j.tim.2003.09.003

Faruque, S. M., & Mekalanos, J. J. (2012). Phage-bacterial interactions in the evolution of toxigenic *Vibrio*

cholerae. *Virulence*, *3*(7), Article 7. https://doi.org/10.4161/viru.22351

Faruque, S. M., Zhu, J., Asadulghani, Kamruzzaman, M., & Mekalanos, J. J. (2003). Examination of diverse

toxin-coregulated pilus-positive *Vibrio cholerae* strains fails to demonstrate evidence for *Vibrio*

pathogenicity island phage. *Infection and Immunity*, *71*(6), 2993–2999.

https://doi.org/10.1128/IAI.71.6.2993-2999.2003

Fiedoruk, K., Zakrzewska, M., Daniluk, T., Piktel, E., Chmielewska, S., & Bucki, R. (2020). Two lineages of

Pseudomonas aeruginosa filamentous phages: Structural uniformity over integration

preferences. *Genome Biology and Evolution*, *12*(10), 1765–1781.

https://doi.org/10.1093/gbe/evaa146

Fillol-Salom, A., Bacarizo, J., Alqasmi, M., Ciges-Tomas, J. R., Martínez-Rubio, R., Roszak, A. W., Cogdell,

R. J., Chen, J., Marina, A., & Penadés, J. R. (2019). Hijacking the hijackers: *Escherichia coli*

pathogenicity islands redirect helper phage packaging for their own benefit. *Molecular Cell*,

75(5), 1020-1030.e4. https://doi.org/10.1016/j.molcel.2019.06.017

Fillol-Salom, A., Bacigalupe, R., Humphrey, S., Chiang, Y. N., Chen, J., & Penadés, J. R. (2021). Lateral

transduction is inherent to the life cycle of the archetypical *Salmonella* phage P22. *Nature*

Communications, *12*(1), 6510. https://doi.org/10.1038/s41467-021-26520-4

Fortmann-Grote, C., Hugoson, E., Summers, J., Theodosiou, L., & Rainey, P. B. (2023). Genome update

for *Pseudomonas fluorescens* isolate SBW25. *Microbiology Resource Announcements*, *12*(2),

e00637-22. https://doi.org/10.1128/mra.00637-22

Frazão, N., Sousa, A., Lässig, M., & Gordo, I. (2019). Horizontal gene transfer overrides mutation in

Escherichia coli colonizing the mammalian gut. *Proceedings of the National Academy of Sciences*,

116(36), 17906–17915. https://doi.org/10.1073/pnas.1906958116

Griffith, F. (1928). The significance of pneumococcal types. *Journal of Hygiene*, *27*(2), 113–159.

https://doi.org/10.1017/S0022172400031879

Grindley, N. D. F., Whiteson, K. L., & Rice, P. A. (2006). Mechanisms of site-specific recombination.

Annual Review of Biochemistry, *75*(1), 567–605.

https://doi.org/10.1146/annurev.biochem.73.011303.073908

Groth, A. C., & Calos, M. P. (2004). Phage integrases: Biology and applications. *Journal of Molecular*

Biology, *335*(3), 667–678. https://doi.org/10.1016/j.jmb.2003.09.082

Groussin, M., Poyet, M., Sistiaga, A., Kearney, S. M., Moniz, K., Noel, M., Hooker, J., Gibbons, S. M.,

Segurel, L., Froment, A., Mohamed, R. S., Fezeu, A., Juimo, V. A., Lafosse, S., Tabe, F. E., Girard,

C., Iqaluk, D., Nguyen, L. T. T., Shapiro, B. J., … Alm, E. J. (2021). Elevated rates of horizontal gene

transfer in the industrialized human microbiome. *Cell*, *184*(8), 2053-2067.e18.

https://doi.org/10.1016/j.cell.2021.02.052

Guo, L., Sattler, L., Shafqat, S., Graumann, P. L., & Bramkamp, M. (2022). A bacterial dynamin-like

protein confers a novel phage resistance strategy on the population level in *Bacillus subtilis*.

MBio, *13*(1). https://journals.asm.org/doi/10.1128/mbio.03753-21

Hackl, T., Laurenceau, R., Ankenbrand, M. J., Bliem, C., Cariani, Z., Thomas, E., Dooley, K. D., Arellano, A.

A., Hogle, S. L., Berube, P., Leventhal, G. E., Luo, E., Eppley, J. M., Zayed, A. A., Beaulaurier, J.,

Stepanauskas, R., Sullivan, M. B., DeLong, E. F., Biller, S. J., & Chisholm, S. W. (2023). Novel

integrative elements and genomic plasticity in ocean ecosystems. *Cell*, *186*(1), 47-62.e16.

https://doi.org/10.1016/j.cell.2022.12.006

Hall, J. P. J., Wood, A. J., Harrison, E., & Brockhurst, M. A. (2016). Source–sink plasmid transfer dynamics

maintain gene mobility in soil bacterial communities. *Proceedings of the National Academy of*

Sciences, *113*(29), 8260–8265. https://doi.org/10.1073/pnas.1600974113

Hmelo, L. R., Borlee, B. R., Almblad, H., Love, M. E., Randall, T. E., Tseng, B. S., Lin, C., Irie, Y., Storek, K.

M., Yang, J. J., Siehnel, R. J., Howell, P. L., Singh, P. K., Tolker-Nielsen, T., Parsek, M. R.,

Schweizer, H. P., & Harrison, J. J. (2015). Precision-engineering the *Pseudomonas aeruginosa*

genome with two-step allelic exchange. *Nature Protocols*, *10*(11), 1820–1841.

https://doi.org/10.1038/nprot.2015.115

Hudson, C. M., Lau, B. Y., & Williams, K. P. (2015). Islander: A database of precisely mapped genomic

islands in tRNA and tmRNA genes. *Nucleic Acids Research*, *43*(D1), D48–D53.

https://doi.org/10.1093/nar/gku1072

Humphrey, S., Fillol-Salom, A., Quiles-Puchalt, N., Ibarra-Chávez, R., Haag, A. F., Chen, J., & Penadés, J. R.

(2021). Bacterial chromosomal mobility via lateral transduction exceeds that of classical mobile

genetic elements. *Nature Communications*, *12*(1), Article 1. https://doi.org/10.1038/s41467-

021-26004-5

Hussain, F. A., Dubert, J., Elsherbini, J., Murphy, M., VanInsberghe, D., Arevalo, P., Kauffman, K., Rodino-

Janeiro, B. K., Gavin, H., Gomez, A., Lopatina, A., Le Roux, F., & Polz, M. F. (2021). Rapid

evolutionary turnover of mobile genetic elements drives bacterial resistance to phages. *Science*,

374(6566), 488–492. https://doi.org/10.1126/science.abb1083

Hyatt, D., Chen, G.-L., LoCascio, P. F., Land, M. L., Larimer, F. W., & Hauser, L. J. (2010). Prodigal:

Prokaryotic gene recognition and translation initiation site identification. *BMC Bioinformatics*,

11(1), 119. https://doi.org/10.1186/1471-2105-11-119

Ibarra-Chávez, R., Hansen, M. F., Pinilla-Redondo, R., Seed, K. D., & Trivedi, U. (2021). Phage satellites

and their emerging applications in biotechnology. *FEMS Microbiology Reviews*, *45*(6), fuab031.

https://doi.org/10.1093/femsre/fuab031

Ishikawa, T., Watabe, K., Mukohara, Y., & Nakamura, H. (1997). Mechanism of stereospecific conversion

of DL-5-substituted hydantoins to the corresponding L-amino acids by *Pseudomonas* sp. Strain

NS671. *Bioscience, Biotechnology, and Biochemistry*, *61*(1), 185–187.

https://doi.org/10.1271/bbb.61.185

Johnson, M. C., Laderman, E., Huiting, E., Zhang, C., Davidson, A., & Bondy-Denomy, J. (2023). Core

defense hotspots within *Pseudomonas aeruginosa* are a consistent and rich source of anti-phage

defense systems. *Nucleic Acids Research*, *51*(10), 4995–5005.

https://doi.org/10.1093/nar/gkad317

Jones, P., Binns, D., Chang, H.-Y., Fraser, M., Li, W., McAnulla, C., McWilliam, H., Maslen, J., Mitchell, A.,

Nuka, G., Pesseat, S., Quinn, A. F., Sangrador-Vegas, A., Scheremetjew, M., Yong, S.-Y., Lopez, R.,

789  & Hunter, S. (2014). InterProScan 5: Genome-scale protein function classification.

790  *Bioinformatics*, *30*(9), 1236–1240. https://doi.org/10.1093/bioinformatics/btu031

791 Karaolis, D. K. R., Somara, S., Maneval, D. R., Johnson, J. A., & Kaper, J. B. (1999). A bacteriophage

792  encoding a pathogenicity island, a type-IV pilus and a phage receptor in cholera bacteria.

793  *Nature*, *399*(6734), Article 6734. https://doi.org/10.1038/20715

794 Koutsovoulos, G. D., Granjeon Noriot, S., Bailly-Bechet, M., Danchin, E. G. J., & Rancurel, C. (2022). AvP:

795  A software package for automatic phylogenetic detection of candidate horizontal gene

796  transfers. *PLOS Computational Biology*, *18*(11), e1010686.

797  https://doi.org/10.1371/journal.pcbi.1010686

798 Lang, A. S., Zhaxybayeva, O., & Beatty, J. T. (2012). Gene transfer agents: Phage-like elements of genetic

799  exchange. *Nature Reviews Microbiology*, *10*(7), 472–482. https://doi.org/10.1038/nrmicro2802

800 Lederberg, J., & Tatum, E. L. (1946). Gene recombination in *Escherichia coli*. *Nature*, *158*(4016), 558–

801  558. https://doi.org/10.1038/158558b0

802 Lenski, R. E. (2017). Experimental evolution and the dynamics of adaptation and genome evolution in

803  microbial populations. *The ISME Journal*, *11*(10), 2181–2194.

804  https://doi.org/10.1038/ismej.2017.69

805 Lenski, R. E., Rose, M. R., Simpson, S. C., & Tadler, S. C. (1991). Long-term experimental evolution in

806  *Escherichia coli*. I. Adaptation and divergence during 2,000 generations. *The American*

807  *Naturalist*, *138*(6), 1315–1341. https://doi.org/10.1086/285289

808 Lenton, T. M., Kohler, T. A., Marquet, P. A., Boyle, R. A., Crucifix, M., Wilkinson, D. M., & Scheffer, M.

809  (2021). Survival of the systems. *Trends in Ecology & Evolution*, *36*(4), 333–344.

810  https://doi.org/10.1016/j.tree.2020.12.003

811 Lilley, A. K., & Bailey, M. J. (1997a). Impact of plasmid pQBR103 acquisition and carriage on the

812  phytosphere fitness of *Pseudomonas fluorescens* SBW25: Burden and benefit. *Applied and*

Environmental Microbiology, *63*(4), 1584–1587. https://doi.org/10.1128/aem.63.4.1584-

1587.1997

Lilley, A. K., & Bailey, M. J. (1997b). The acquisition of indigenous plasmids by a genetically marked

pseudomonad population colonizing the sugar beet phytosphere is related to local

environmental conditions. *Applied and Environmental Microbiology*, *63*(4), 1577–1583.

https://doi.org/10.1128/aem.63.4.1577-1583.1997

Lindsay, J. A., Ruzin, A., Ross, H. F., Kurepina, N., & Novick, R. P. (1998). The gene for toxic shock toxin is

carried by a family of mobile pathogenicity islands in *Staphylococcus aureus*. *Molecular

Microbiology*, *29*(2), 527–543. https://doi.org/10.1046/j.1365-2958.1998.00947.x

Low, S. J., Džunková, M., Chaumeil, P.-A., Parks, D. H., & Hugenholtz, P. (2019). Evaluation of a

concatenated protein phylogeny for classification of tailed double-stranded DNA viruses

belonging to the order Caudovirales. *Nature Microbiology*, *4*(8), 1306–1315.

https://doi.org/10.1038/s41564-019-0448-z

Marrs, B. (1974). Genetic recombination in *Rhodopseudomonas capsulata*. *Proceedings of the National

Academy of Sciences*, *71*(3), 971–973. https://doi.org/10.1073/pnas.71.3.971

Nayfach, S., Páez-Espino, D., Call, L., Low, S. J., Sberro, H., Ivanova, N. N., Proal, A. D., Fischbach, M. A.,

Bhatt, A. S., Hugenholtz, P., & Kyrpides, N. C. (2021). Metagenomic compendium of 189,680

DNA viruses from the human gut microbiome. *Nature Microbiology*, *6*(7), 960–970.

https://doi.org/10.1038/s41564-021-00928-6

Néron, B., Denise, R., Coluzzi, C., Touchon, M., Rocha, E. P. C., & Abby, S. S. (2023). MacSyFinder v2:

Improved modelling and search engine to identify molecular systems in genomes. *Peer

Community Journal*, *3*, e28. https://doi.org/10.24072/pcjournal.250

Novick, R. P., Christie, G. E., & Penadés, J. R. (2010). The phage-related chromosomal islands of Gram-

positive bacteria. *Nature Reviews Microbiology*, *8*(8), 541–551.

https://doi.org/10.1038/nrmicro2393

Ofir, G., Melamed, S., Sberro, H., Mukamel, Z., Silverman, S., Yaakov, G., Doron, S., & Sorek, R. (2017).

DISARM is a widespread bacterial defence system with broad anti-phage activities. *Nature*

Microbiology, *3*(1), 90–98. https://doi.org/10.1038/s41564-017-0051-0

O'Shea, Y. A., & Boyd, E. F. (2002). Mobilization of the *Vibrio* pathogenicity island between *Vibrio*

cholerae isolates mediated by CP-T1 generalized transduction. *FEMS Microbiology Letters*,

214(2), 153–157. https://doi.org/10.1111/j.1574-6968.2002.tb11339.x

Pei, T.-T., Luo, H., Wang, Y., Li, H., Wang, X.-Y., Zhang, Y.-Q., An, Y., Wu, L.-L., Ma, J., Liang, X., Yan, A.,

Yang, L., Chen, C., & Dong, T. (2024). Filamentous prophage Pf4 promotes genetic exchange in

Pseudomonas aeruginosa. *The ISME Journal*, *18*(1), wrad025.

https://doi.org/10.1093/ismejo/wrad025

Quistad, S. D., Doulcier, G., & Rainey, P. B. (2020). Experimental manipulation of selfish genetic elements

links genes to microbial community function. *Philosophical Transactions of the Royal Society B:*

Biological Sciences, *375*, 20190681. https://doi.org/10.1098/rstb.2019.0681

Rainey, P. B., & Quistad, S. D. (2020). Toward a dynamical understanding of microbial communities.

Philosophical Transactions of the Royal Society B: Biological Sciences, *375*, 20190248.

https://doi.org/10.1098/rstb.2019.0248

Rainey, P. B., Remigi, P., Farr, A. D., & Lind, P. A. (2017). Darwin was right: Where now for experimental

evolution? *Current Opinion in Genetics & Development*, *47*, 102–109.

https://doi.org/10.1016/j.gde.2017.09.003

Rainey, P. B., & Travisano, M. (1998). Adaptive radiation in a heterogeneous environment. *Nature*,

394(6688), 69–72. https://doi.org/10.1038/27900

Roughgarden, J. (2020). Holobiont evolution: Mathematical model with vertical vs. horizontal

microbiome transmission. *Philosophy, Theory, and Practice in Biology*, *12*(20220112).

https://doi.org/10.3998/ptpbio.16039257.0012.002

Rousset, F., Depardieu, F., Miele, S., Dowding, J., Laval, A.-L., Lieberman, E., Garry, D., Rocha, E. P. C.,

Bernheim, A., & Bikard, D. (2022). Phages and their satellites encode hotspots of antiviral

systems. *Cell Host & Microbe*, *30*(5), 740-753.e5. https://doi.org/10.1016/j.chom.2022.02.018

Schlechter, R. O., Jun, H., Bernach, M., Oso, S., Boyd, E., Muñoz-Lintz, D. A., Dobson, R. C. J., Remus, D.

M., & Remus-Emsermann, M. N. P. (2018). Chromatic bacteria – A broad host-range plasmid and

chromosomal insertion toolbox for fluorescent protein expression in bacteria. *Frontiers in*

Microbiology, *9*, 3052. https://doi.org/10.3389/fmicb.2018.03052

Seemann, T. (2014). Prokka: Rapid prokaryotic genome annotation. *Bioinformatics*, *30*(14), 2068–2069.

https://doi.org/10.1093/bioinformatics/btu153

Sievers, F., Wilm, A., Dineen, D., Gibson, T. J., Karplus, K., Li, W., Lopez, R., McWilliam, H., Remmert, M.,

Söding, J., Thompson, J. D., & Higgins, D. G. (2011). Fast, scalable generation of high-quality

protein multiple sequence alignments using Clustal Omega. *Molecular Systems Biology*, *7*(539),

1–6. https://doi.org/10.1038/msb.2011.75

Silby, M. W., Cerdeño-Tárraga, A. M., Vernikos, G. S., Giddens, S. R., Jackson, R. W., Preston, G. M.,

Zhang, X.-X., Moon, C. D., Gehrig, S. M., Godfrey, S. A., Knight, C. G., Malone, J. G., Robinson, Z.,

Spiers, A. J., Harris, S., Challis, G. L., Yaxley, A. M., Harris, D., Seeger, K., … Thomson, N. R. (2009).

Genomic and genetic analyses of diversity and plant interactions of *Pseudomonas fluorescens*.

Genome Biology, *10*(5), R51. https://doi.org/10.1186/gb-2009-10-5-r51

Sullivan, J. T., & Ronson, C. W. (1998). Evolution of rhizobia by acquisition of a 500-kb symbiosis island

that integrates into a phe-tRNA gene. *Proceedings of the National Academy of Sciences*, *95*(9),

5145–5149. https://doi.org/10.1073/pnas.95.9.5145

Sullivan, M. J., Petty, N. K., & Beatson, S. A. (2011). Easyfig: A genome comparison visualizer.

Bioinformatics, *27*(7), 1009–1010. https://doi.org/10.1093/bioinformatics/btr039

Tatusov, R. L. (2000). The COG database: A tool for genome-scale analysis of protein functions and

evolution. *Nucleic Acids Research*, *28*(1), 33–36. https://doi.org/10.1093/nar/28.1.33

Tesson, F., Hervé, A., Mordret, E., Touchon, M., d'Humières, C., Cury, J., & Bernheim, A. (2022).

Systematic and quantitative view of the antiviral arsenal of prokaryotes. *Nature*

Communications, *13*(1), 2561. https://doi.org/10.1038/s41467-022-30269-9

Tock, M. R., & Dryden, D. T. (2005). The biology of restriction and anti-restriction. *Current Opinion in*

Microbiology, *8*(4), 466–472. https://doi.org/10.1016/j.mib.2005.06.003

van Dijk, B., Buffard, P., Farr, A. D., Giersdorf, F., Meijer, J., Dutilh, B. E., & Rainey, P. B. (2023).

Identifying and tracking mobile elements in evolving compost communities yields insights into

the nanobiome. *ISME Communications*, *3*(90). https://doi.org/10.1101/2023.02.02.526783

Wang, X., Kim, Y., & Wood, T. K. (2009). Control and benefits of CP4-57 prophage excision in *Escherichia*

coli biofilms. *The ISME Journal*, *3*(10), 1164–1179. https://doi.org/10.1038/ismej.2009.59

Woods, L. C., Gorrell, R. J., Taylor, F., Connallon, T., Kwok, T., & McDonald, M. J. (2020). Horizontal gene

transfer potentiates adaptation by reducing selective constraints on the spread of genetic

variation. *Proceedings of the National Academy of Sciences*, *117*(43), 26868–26875.

https://doi.org/10.1073/pnas.2005331117

Xie, Z., & Tang, H. (2017). ISEScan: Automated identification of insertion sequence elements in

prokaryotic genomes. *Bioinformatics*, *33*(21), 3340–3347.

https://doi.org/10.1093/bioinformatics/btx433

Yuan, L., Lu, H., Li, F., Nielsen, J., & Kerkhoven, E. J. (2023). HGTphyloDetect: Facilitating the

identification and phylogenetic analysis of horizontal gene transfer. *Briefings in Bioinformatics*,

24(2), bbad035. https://doi.org/10.1093/bib/bbad035

Zinder, N. D., & Lederberg, J. (1952). Genetic exchange in *Salmonella*. *Journal of Bacteriology*, *64*(5),

679–699. https://doi.org/10.1128/jb.64.5.679-699.1952

 Figure Captions

Fig. 1. Experimental evolution and detection of foreign elements

a. Three serial-propagation regimes to evolve SBW25 populations. (i) live filtrate regime; (ii) co-culture
regime; (iii) fixed filtrate regime. Components within each box are specific to each regime, and
components outside of boxes are common to all regimes.

b. The pipeline for screening and confirming SBW25 strains carrying foreign elements inserted into the
genome.

c. Basic information for the 3 foreign elements detected. The table lists the number of base pairs of each
foreign element, the name of the compost community from which the community filtrate was
generated, the serial-propagation regime, the number of parallel lines from which isolates with the
foreign element were detected, and the number of isolates containing the foreign element in each such
line.

d. Insertion site of the foreign elements in the SBW25 genome. Upon insertion, two 14 bp direct repeats
flank the foreign element. The direct repeats at the 5'-end and 3'-end of the element were treated as
from SBW25 genome and from the foreign element, respectively, despite the exact origin of each repeat
is unknown.

e. The aligned genetic maps of the 3 foreign elements, I23, I44 and I55. The alignment was created by
Easyfig (M. J. Sullivan et al., 2011). Parallelograms between MGEs indicate similar genomic regions based
on blastn. The ORFs were numbered serially from 5-end to 3'-end for each MGE, and coloured according
to annotations. Detailed annotations of each ORF are listed in Table S4.

Fig. 2. Bioinformatic analysis of similar elements

a. Bioinformatic pipeline for detection of similar elements to I55. Bullet points below 'sequence

comparison' and 'sequence selection' represent the criteria for comparison and searching, respectively.

b. Cladogram and genomic representations of I23, I44, I55 and similar MGEs (identified by NCBI RefSeq

assembly indices). Basis for construction of the tree is summarised in the text. Shape of the tree

represents inferred evolutionary relationship but not evolutionary distances.

c. Rank abundance of annotated genes of different functional classes in panel b. Genes occurring at least

10 times are illustrated. A full rank abundance table is in Table S6. Colours represent COG functional

categories.

d. Pairwise distances between the MGEs (shown in lower-left), and pairwise distances between their

corresponding hosts (shown in upper-right). MGEs and hosts are identified by NCBI RefSeq assembly

indices. Basis for construction and interpretation of the heatmap is summarised in the text. Pink boxes

highlight pairs of strains with almost identical MGE, but remarkably different 16S sequence. Annotations

of those MGEs are boxed in pink in panel b.

Fig. 3. Intercellular transfer of I55

a. Schematic of the one-step intercellular transfer assay.

b. The densities of CFUs on LB + Kan + Tet plates in the intercellular transfer assay. Data are mean ± s.d.
The labels A/C and D.L. represent autoclaved filtrate and the detection limit of the assay, respectively.
For each treatment, 8 to 12 biological replicates were used. Mann-Whitney test was applied to test the
effect of IntA ($p < 0.0001$) when the community filtrate was added.

c. Schematic of the two-step intercellular transfer assay.

d. The densities of CFUs on LB + Kan + Tet plates in the two-stage intercellular transfer assay. Data are
mean ± s.d. The label D.L. represents the detection limit of the assay. For each treatment, 8 biological
replicates were used. Mann-Whitney test was applied to test the effect of IntA ($p = 0.0152$) when the
community filtrate was added at the first step.

e. Proposed model for the transfer of I55 via an unknown component in the community filtrate.

Fig. 4. Fitness advantage and phage defence activity of I55

a. Schematic of fluorescent fitness assay. The diagram indicates the case where SBW25-I55Δ*intY* was
tagged with GFP and the wild type was tagged with RFP. The reverse tagging strategy, where SBW25-
I55Δ*intY* was tagged with RFP and the wild type was tagged with GFP, is also conducted to eliminate any
fitness difference due to the choice of fluorescent protein markers.

b. Fluorescent fitness assay results for SBW25-I55Δ*intY* relative to wild type, in presence or absence of
community filtrate. Data are mean ± 95% CI. In the Filtrate + and - treatments, stored filtrate aliquots
from compost C and from SBW25 were added, respectively. The colour of each data point corresponds
to the colour of the fluorescent marker added to SBW25-I55Δ*intY*. For each treatment, 8 biological
replicates (4 for either tagging strategy) were used. One sample *t*-tests were applied to test the relative
fitness values against 1 when the community filtrate was added (p = 0.0104) and not added (p = 0.0033).
Paired Welch's *t*-test was applied to test the effect of community filtrate (p = 0.0006).

c. Typical photos of the plaque assay of the community filtrate against SBW25-I55Δ*intY* or wild type as
hosts. Picture contrast was altered to facilitate visual realisation of the plaques. Photos of three
additional biological replicates can be found in supporting information.

d. Plaque forming unit (PFU) results of the plaque assay in panel c. Data are mean ± s.d. For each strain,
four biological replicates were used. Paired Welch's *t*-test was applied to test the effect of I55Δ*intY* (p =
0.0080).

e. Fluorescent fitness assay results for SBW25-I55Δ*intY* relative to wild type, in presence or absence of
the two phage isolates (P1 and P2). Data are mean ± 95% CI. In the P1, P2 and Ctrl treatments, stored
phage lysate P1, phage lysate P2 and lysate extraction control (see Materials and Methods) were added,
respectively. The colour of each data point corresponds to the colour of the fluorescent marker added to
SBW25-I55Δ*intY*. For each treatment, 4 biological replicates (2 for either tagging strategy) were used.

f. PFU results of plaque assay of P1 and P2 (in (i) and (ii) respectively) against SBW25-I55Δ*intY* or wild
type as hosts. Data are mean ± s.d. For each strain, 4 biological replicates were used. Paired Welch's *t*-
tests were applied to test the effect of I55Δ*intY* (p = 0.0011 in the case of P1, and p = 0.0082 in the case
of P2).

Supplementary Figure Captions and Tables

Fig. S1. SBW25 density in the co-culture regime (see Fig. 1a(ii) and Materials and Methods)

SBW25 co-cultured with compost C community decreased in abundance over transfers. By the 5^{th}

transfer the density was only $10^3 \sim 10^4$ cells (per mL) across the three replicate lines. By contrast, mono-

culture of SBW25 maintained a density of $10^8 \sim 10^9$ cells (per mL).

Fig. S2. Excision and circularisation of I55

a. The optical density of overnight LB shaking cultures of SBW25-I55, SBW25-I55Δ*intY* and wild type
(WT). Data are mean ± s.d. Brown-Forsythe and Welch ANOVA test was applied to test the difference
between the three strains (p = 0.3551).

b. PCR of overnight cultures of SBW25-I55, SBW25-I55Δ*intY* and WT, with combination of primers
①②③④. Photos of 3 more biological replicates can be found in supporting information.

c. A proposed explanation for the presence of bands for SBW25-I55 with primer pairs ①④ and ②③.
Primers ① and ② flank the left insertion junction, and primers ③ and ④ flanks the right. Primers ①
and ④ match the genome, and primers ② and ③ match I55. The presence of bands on SBW25-I55
with pairs ①+② and ③+④ confirms the integration of I55 in the genome. The presence of band on
SBW25-I55 with pair ①+④, at a length that is the same as WT, suggests that in the overnight culture
of SBW25-I55 there are some cells with I55 excised from the genome. The presence of band on SBW25-
I55 with pair ②+③ suggests that I55 somehow circularises so that primers ② and ③ flank the
circularisation junction. Sanger sequencing of the PCR products of ②+③ confirms that the circular I55
contains exactly one 14 bp sequence at the junction, and of ①+④ confirms that I55 excises in a scar-
free manner. We therefore proposed that there are two forms of I55 in the population of overnight
culture of SBW25-I55, the integrated form (which gives bands with pairs ①+② and ③+④) and the
circular form (which gives bands with pairs ①+④ and ②+③). To test whether the tyrosine integrase
(IntY) is necessary for excision, we deleted *intY* and did the same PCR. The strain SBW25-I55Δ*intY*
showed bands with pairs ①+② and ③+④, but no band with pairs ①+④ and ②+③. This
confirms that IntY is necessary for excision and circularisation of I55.

1015

1016 Table S1. List of bacterial strains

1017

Number	Name	Genotype	Construction notes
MPB29212	SBW25_Wild_Type	SBW25	Lab stock
MPB28806	SBW25 Tn7-PntpII-mScarlet-I-Gm	SBW25 *glmS*::Tn7-145(RFP, Gen^R)	Lab stock
MPB29284	Tn7_152_CFC_LB_SBW25	SBW25 *glmS*::Tn*7*-152(GFP, Kan^R)	Conjugation of MPB29212 with MPB27708
MPB29300	Tn7_155_CFC_LB_SBW25	SBW25 *glmS*::Tn*7*-155(RFP, Kan^R)	Conjugation of MPB29212 with MPB27709
MPB38392	SFF_b_GFP_4_iso24_23kb	SBW25 *tmRNA*::I23 *glmS*::Tn*7*-152(GFP, Kan^R) isolate1*	Isolated from fixed-filtrate experiment, compost b, GFP, line 4
MPB38393	SFF_b_GFP_5_iso02_23kb	SBW25 *tmRNA*::I23 *glmS*::Tn*7*-152(GFP, Kan^R) isolate2*	Isolated from fixed-filtrate experiment, compost b, GFP, line 5
MPB38394	SFF_b_GFP_5_iso04_23kb	SBW25 *tmRNA*::I23 *glmS*::Tn*7*-152(GFP, Kan^R) isolate3*	Isolated from fixed-filtrate experiment, compost b, GFP, line 5
MPB38395	SFF_b_GFP_5_iso10_23kb	SBW25 *tmRNA*::I23 *glmS*::Tn*7*-152(GFP, Kan^R) isolate4*	Isolated from fixed-filtrate experiment, compost b, GFP, line 5
MPB38396	SFF_b_GFP_5_iso13_23kb	SBW25 *tmRNA*::I23 *glmS*::Tn*7*-152(GFP, Kan^R) isolate5*	Isolated from fixed-filtrate experiment, compost b, GFP, line 5
MPB38895	SFF_b_RFP_3_iso13_23kb	SBW25 *tmRNA*::I23 *glmS*::Tn*7*-155(RFP, Kan^R) isolate6*	Isolated from fixed-filtrate experiment, compost b, RFP, line 3
MPB38906	SFF_b_RFP_4_iso06_44kb	SBW25 *tmRNA*::I44 *glmS*::Tn*7*-155(RFP, Kan^R) isolate1*	Isolated from fixed-filtrate experiment, compost b, RFP, line 4
MPB38907	SFF_b_RFP_4_iso23_44kb	SBW25 *tmRNA*::I44 *glmS*::Tn*7*-155(RFP, Kan^R) isolate2*	Isolated from fixed-filtrate experiment, compost b, RFP, line 4
MPB33357	SF_G_C_2_T14_iso04	SBW25 *tmRNA*::I55 *glmS*::Tn*7*-152(GFP, Kan^R) isolate1*	Isolated from live-filtrate experiment, compost c, GFP, line 2
MPB38103/MPB38109	Iso04_rev_PFLU4414	SBW25 *tmRNA*::I55 *glmS*::Tn*7*-152(GFP, Kan^R) *pflu3805*SNP	2SAR of two independent colonies of MPB33357 with MPB38033
MPB38572/MPB38574	Iso04_rev_PFLU4414_3805	SBW25 *tmRNA*::I55 *glmS*::Tn*7*-152(GFP, Kan^R)	2SAR of MPB38103/MPB38109 with MPB38391
MPB39024/MPB39026	Iso04_rev_PFLU4414_3805_del_Tn7	SBW25 *tmRNA*::I55	2SAR of MPB38572/MPB38574 with MPB38943
MPB39132/MPB39136	SBW25_I55_Δint	SBW25 *tmRNA*::(I55 Δ*intY*)	2SAR of MPB39024/MPB39026 with MPB38939
MPB39504/MPB39507	SBW25_I55_Δint_Tn7_GFP	SBW25 *tmRNA*::(I55 Δ*intY*) *glmS*::Tn*7*-152(GFP, Kan^R)	Conjugation of MPB39132/MPB39136 with MPB27708
MPB39514/MPB39516	SBW25_I55_Δint_Tn7_RFP	SBW25 *tmRNA*::(I55 Δ*intY*) *glmS*::Tn*7*-152(GFP, Kan^R)	Conjugation of MPB39132/MPB39136 with MPB27709
MPB39379/MPB39459	SBW25_reconstructed	SBW25	2SAR of MPB39132/MPB39136 with MPB39223
MPB39508/MPB39510	SBW25_reconstructed_Tn7_GFP	SBW25 *glmS*::Tn*7*-152(GFP, Kan^R)	Conjugation of MPB39379/MPB39459 with MPB27708
MPB39518/MPB39520	SBW25_reconstructed_Tn7_RFP	SBW25 *glmS*::Tn*7*-155(RFP, Kan^R)	Conjugation of MPB39379/MPB39459 with MPB27709
MPB39465/MPB39467	SBW25_I55_KanR_InsertS1	SBW25 *tmRNA*::(I55 (*orf08-orf09*)::Kan^R)	2SAR of MPB39024/MPB39026 with MPB39269
MPB39565/MPB39568	SBW25_I55_Δint_ KanR_InsertS1	SBW25 *tmRNA*::(I55 Δ*intY* (*orf08-orf09*)::Kan^R)	2SAR of MPB39132/MPB39136 with MPB39269
MPB39500	SBW25-TetR	SBW25 *glmS*::Tn7(AGACTGTTGCGCGCTGGCGGTGGCG, Tet^R)	Lab stock, see https://doi.org/10.1007/s00239-023-10103-6

1018

1019 * For variations/mutations, see Table S5.

1020

 Table S2. List of *E. coli* plasmids

Number	Name	Description	Source
MPB27708	pMRE-Tn7-152	Plasmid to tag *Pseudomonas* with GFP and Kan^R	https://www.addgene.org/118566/
MPB27709	pMRE-Tn7-155	Plasmid to tag *Pseudomonas* with RFP and Kan^R	https://www.addgene.org/118569/
MPB35522	pUIsacB	Empty plasmid for 2-step allelic replacement	Lab stock, see https://www.biorxiv.org/content/10.1101/2024.07.11.603030v1
MPB35623	pRK2013	Helper plasmid for 2-step allelic replacement	Lab stock, see https://www.biorxiv.org/content/10.1101/2024.07.11.603030v1
MPB38033	rev_PFLU4414	Plasmid to revert the SNP in *pflu4414*	this study
MPB38391	rev_PFLU3805	Plasmid to revert the SNP in *pflu3805*	this study
MPB38943	del_Tn7	Plasmid to delete the Tn*7* cassette	this study
MPB38939	del_55kbInt	Plasmid to delete the gene *intY* of I55	this study
MPB39223	del_tmIns	Plasmid to delete any inserts downstream of *tmRNA*	this study
MPB39269	KanR_InsertS1	Plasmid to insert a KanR cassette between *orf08* and *orf09* of I55	this study

Table S3. List of primers

Number	Name	Purpose	Sequence
YZ001	YZ_SBW_5797955F	To detect potential insertions downstream of tmRNA	GTTACCGCTGCGATGTTGAC
YZ002	YZ_SBW_5799313R	To detect potential insertions downstream of tmRNA	GCGTGCTGATGCTGCTAATC
YZ003	YZ_In_1016R	To confirm the presence of I55	GTGGCCAGATCAGGGAAGAC
YZ004	YZ_In_54359F	To confirm the presence of I55	ACACGACACACGAAGAGACC
YZ005	pUIsacB_fwd	To amplify the pUIsacB plasmid backbone	AGCACTACATCAACTGACTA
YZ006	pUIsacB_rev	To amplify the pUIsacB plasmid backbone	CTGAACCAAGATAGCTGTAC
YZ007	2SAR_4880278_Insert_fwd	To construct the plasmid for reverting the SNP in *pflu4414*	GTACAGCTATCTTGGTTCAGGAAATCAACCCATCGCCTG
YZ008	2SAR_4880278_Insert_rev	To construct the plasmid for reverting the SNP in *pflu4414*	TAGTCAGTTGATGTAGTGCTAACGCCTGTCCTATGTGG
YZ009	pUIsacB_seq_fwd	To Sanger sequence the assembled regions of the pUIsacB plasmid	ATTTGTCCTACTCAGGAGAGCGTTC
YZ010	4880278_Insert_4962R	To Sanger sequence the plasmid for reverting the SNP in *pflu4414*	GCTGAGTTCGAATCCCTGCT
YZ011	4880278_Insert_4804F	To Sanger sequence the plasmid for reverting the SNP in *pflu4414*	CTTGCCATGCAGTTGGTCG
YZ012	4880278_Insert_5486R	To Sanger sequence the plasmid for reverting the SNP in *pflu4414*	GGCCGGCGAAATTTTAGAGC
YZ013	4880278_Insert_5275F	To Sanger sequence the plasmid for reverting the SNP in *pflu4414*	CGAATCAACGTGACGCTCAC
YZ014	4880278_Insert_5956R	To Sanger sequence the plasmid for reverting the SNP in *pflu4414*	CGACTGGGGCCGTTTCAT
YZ015	4880278_Insert_5777F	To Sanger sequence the plasmid for reverting the SNP in *pflu4414*	ACACGCTTGATCACCTGACC
YZ016	pUIsacB_seq_rev	To Sanger sequence the assembled regions of the pUIsacB plasmid	AGCGTTCTGAACAAATCCAGATG
YZ084	SBW_23kb_insert_1	To confirm the presence of I23	CCGTTTCTCTTTGGGGTCGA
YZ085	SBW_23kb_insert_2	To confirm the presence of I23	ACTCAAGAGCCTCGCCAAAG
YZ086	2SAR_rev_3805_1	To construct the plasmid for reverting the SNP in *pflu3805*	GTACAGCTATCTTGGTTCAGCCAGCAACGGGTTGTTCTTG
YZ087	2SAR_rev_3805_2	To construct the plasmid for reverting the SNP in *pflu3805*	TAGTCAGTTGATGTAGTGCTCCACTGATCTTGACGCCAGA
YZ088	rev_PFLU3805_4997F	To Sanger sequence the plasmid for reverting the SNP in *pflu3805*	CCTGTTCGCTGAAAATGGCC
YZ089	rev_PFLU3805_5687R	To Sanger sequence the plasmid for reverting the SNP in *pflu3805*	ACACGATGACGACGGTTCTG
YZ091	PFLU4414_Final_Seq_2	To confirm the genotype of mutants after allelic replacement in *pflu4414*	GAACGCATCCACGGCTAATG
YZ092	PFLU3805_Final_Seq_1	To confirm the genotype of mutants after allelic replacement in *pflu3805*	GACACGTCGACCTTCTGGAAG
YZ093	PFLU3805_Final_Seq_2	To confirm the genotype of mutants after allelic replacement in *pflu3805*	CGAGTTGATTGGGGTGGCC
YZ098	PFLU4414_Final_Seq_1_1	To confirm the genotype of mutants after allelic replacement in *pflu4414*	CGCCGACCATGTTCATGATG
YZ102	SBW_44kb_insert_1	To confirm the presence of I44	GCATAGGCTGTGGTCACTGA
YZ103	SBW_44kb_insert_2	To confirm the presence of I44	AGACCAAGGCCGTCAAGAAG
YZ105	2SAR_del_55kbInt_1	To construct the plasmid for deleting the tyrosine integrase (*orf01*) of I55	GTACAGCTATCTTGGTTCAGGTTTAGGATTCGACGCCG
YZ106	2SAR_del_55kbInt_2	To construct the plasmid for deleting the tyrosine integrase (*orf01*) of I55	GAAGTCTCTCGGGTGTATTCCAGTGCTGTTTATG
YZ107	2SAR_del_55kbInt_3	To construct the plasmid for deleting the tyrosine integrase (*orf01*) of I55	TGGAATACACCCGAGAGACTTCAACCAGTGA

YZ108	2SAR_del_55kbInt_4	To construct the plasmid for deleting the tyrosine integrase (*orf01*) of I55	TAGTCAGTTGATGTAGTGCTTCTTCGTTCACCACATTCCG
YZ109	del_55kbInt_4941F	To Sanger sequence the plasmid for deleting the tyrosine integrase (*orf01*) of I55	TGGACGAAGCGGCTAAAACT
YZ110	del_55kbInt_5514R	To Sanger sequence the plasmid for deleting the tyrosine integrase (*orf01*) of I55	TCGCCATTCAGCTGTCCAAT
YZ111	del_55kbInt_Final_Seq_1	To confirm the genotype of mutants after allelic replacement in the tyrosine integrase (*orf01*) of I55	ACGTTAGCTCAGCCGAGGT
YZ112	2SAR_del_Tn7_1	To construct the plasmid for deleting the Tn*7* cassette	GTACAGCTATCTTGGTTCAGTGAGCTATCAGTTGCGCCTG
YZ113	2SAR_del_Tn7_2	To construct the plasmid for deleting the Tn*7* cassette	TAGTCAGTTGATGTAGTGCTAATCCGACCTGACCTTGCTG
YZ114	del_Tn7_4864F	To Sanger sequence the plasmid for deleting the Tn*7* cassette	TTGCGGGTTACGGATGACC
YZ115	del_Tn7_5273F	To Sanger sequence the plasmid for deleting the Tn*7* cassette	GACGTTGGTGGATCAGGGAG
YZ116	del_Tn7_5670F	To Sanger sequence the plasmid for deleting the Tn*7* cassette	CAATGGGCCGTGTTTGAGC
YZ117	del_Tn7_5084R	To Sanger sequence the plasmid for deleting the Tn*7* cassette	TTGGAAGGTGAGCTGCTGTC
YZ118	del_Tn7_5480R	To Sanger sequence the plasmid for deleting the Tn*7* cassette	CTGTCGCCGATCCTCTACAC
YZ119	del_Tn7_5845R	To Sanger sequence the plasmid for deleting the Tn*7* cassette	CAATGGACAGCACCGTGGAA
YZ120	del_Tn7_Final_Seq_1	To confirm the genotype of mutants after allelic replacement in the Tn*7* cassette	ACACCGCCCAACGTGATG
YZ121	del_Tn7_Final_Seq_2	To confirm the genotype of mutants after allelic replacement in the Tn*7* cassette	ACACCCTGTTCGTGACCATC
YZ124	2SAR_KanR_InsertS1_1	To construct the plasmid for inserting an Kan^R cassette between *orf08* and *orf09* of I55	GTACAGCTATCTTGGTTCAGGTCCGCAGAACTGTTGGAG
YZ125	2SAR_KanR_InsertS1_2	To construct the plasmid for inserting an Kan^R cassette between *orf08* and *orf09* of I55	GATTAGCTGTGTCGCTACTG
YZ126	2SAR_KanR_InsertS1_3	To construct the plasmid for inserting an Kan^R cassette between *orf08* and *orf09* of I55	CCGCTCACCGAAGCTGATTA
YZ127	2SAR_KanR_InsertS1_4	To construct the plasmid for inserting an Kan^R cassette between *orf08* and *orf09* of I55	TAGTCAGTTGATGTAGTGCTGGCGACTTCGGTATTCCAG
YZ128	2SAR_KanR_InsertS1_5	To construct the plasmid for inserting an Kan^R cassette between *orf08* and *orf09* of I55	CAGTAGCGACACAGCTAATCGGGAGACCAGAAACAAAAAAACACCCCGAAGGGTGTTCAATAATTGGTCCGCTTCCTTTAGCAGCC
YZ129	2SAR_KanR_InsertS1_6	To construct the plasmid for inserting an Kan^R cassette between *orf08* and *orf09* of I55	TAATCAGCTTCGGTGAGCGGTCCTAGGTGCTCGAGTGG
YZ130	KanR_InsertS1_5008F	To Sanger sequence the plasmid for inserting an Kan^R cassette between *orf08* and *orf09* of I55	TGGCACACGATACCGAAGAC
YZ131	KanR_InsertS1_5508F	To Sanger sequence the plasmid for inserting an Kan^R cassette between *orf08* and *orf09* of I55	AGCGTGCGAGCTCTAAAACA
YZ132	KanR_InsertS1_6011F	To Sanger sequence the plasmid for inserting an Kan^R cassette between *orf08* and *orf09* of I55	ACTCTTCCGAGCAAAGGACG
YZ133	KanR_InsertS1_6662F	To Sanger sequence the plasmid for inserting an Kan^R cassette between *orf08* and *orf09* of I55	GCAGGAAATGGGTCATCGGT
YZ134	KanR_InsertS1_5189R	To Sanger sequence the plasmid for inserting an Kan^R cassette between *orf08* and *orf09* of I55	TTGCAGCAGTGACAAGGACA
YZ135	KanR_InsertS1_5695R	To Sanger sequence the plasmid for inserting an Kan^R cassette between *orf08* and *orf09* of I55	TTGATCTTGGGAGAAGCGGC
YZ136	KanR_InsertS1_6177R	To Sanger sequence the plasmid for inserting an Kan^R cassette between *orf08* and *orf09* of I55	GGACAGCCGGTATAAAGGGA
YZ137	KanR_InsertS1_6853R	To Sanger sequence the plasmid for inserting an Kan^R cassette between *orf08* and *orf09* of I55	TCTCACACGTTCGTTCGACC
YZ144	2SAR_del_tmIns_1	To construct the plasmid for deleting MGEs inserted downstream of *tmRNA*	GTACAGCTATCTTGGTTCAGGAGATCGACACCGCCGTCTC
YZ145	2SAR_del_tmIns_2	To construct the plasmid for deleting MGEs inserted downstream of *tmRNA*	TAGTCAGTTGATGTAGTGCTGCGTGCTGATGCTGCTAATC
YZ146	del_tmIns_5225F	To Sanger sequence the plasmid for deleting MGEs inserted downstream of *tmRNA*	GGGTCGCTGAGGGTTAACTT
YZ147	del_tmIns_5699F	To Sanger sequence the plasmid for deleting MGEs inserted downstream of *tmRNA*	ATGACGCTCGTGTCCAACTT
YZ148	del_tmIns_4886R	To Sanger sequence the plasmid for deleting MGEs inserted downstream of *tmRNA*	CTGTGCAGTGTACGCCAAAC

YZ149	del_tmIns_5575R	To Sanger sequence the plasmid for deleting MGEs inserted downstream of *tmRNA*	GGTACAGACCCTGTTCGTCG
YZ150	del_tmIns_Final_Seq_1	To confirm the genotype of mutants after allelic replacement downstream of *tmRNA*	TTACCGCTGCGATGTTGACT
YZ151	del_tmIns_Final_Seq_2	To confirm the genotype of mutants after allelic replacement downstream of *tmRNA*	GGAGCGTCCGTGAGAGATTC
YZ161	Tn7R-109	To confirm the insertion of Tn7 cassette in SBW25 genome	CAGCATAACTGGACTGATTTCAG
YZ162	SBW25-glmS	To confirm the insertion of Tn7 cassette in SBW25 genome	CACCAAAGCTTTCACCACCCAA

1027

1028

    Table S4. Annotations of MGEs I23, I44 and I55

Element	ORF name	Consensus annotation	Minimum coordinate	Maximum coordinate	Length	Direction of transcription	Prokka product	Pfam domains	DefenseFinder prediction
I23	orf01 (intY)	tyrosine integrase	186	1442	1257	forward	Prophage integrase IntA	PF13356 (Arm-DNA-bind_3; aa 8-94); PF22022 (Phage_int_M; aa 107-200); PF00589 (Phage_integrase; aa 214-378).	NA
I23	orf02	GajA	1631	3436	1806	reverse	ATP-dependent DNA helicase Rep	PF00580 (UvrD-helicase; aa 5-125, 133-239).	Gabija__GajB_1.
I23	orf03	GajB	3462	5312	1851	reverse	hypothetical protein	PF13175 (AAA_15; aa 1-72, 310-381); PF20469 (OLD-like_TOPRIM; aa 429-500).	Gabija__GajA.
I23	orf04	unannotated	5637	5966	330	forward	hypothetical protein	NA	NA
I23	orf05	DUF932	6062	7030	969	forward	hypothetical protein	PF06067 (DUF932; aa 77-302).	NA
I23	orf06	YqaJ	7106	8110	1005	forward	hypothetical protein	PF09588 (YqaJ; aa 29-172).	NA
I23	orf07	Gp3-like	8201	9151	951	forward	hypothetical protein	PF18897 (Gp3-like; aa 12-206).	NA
I23	orf08	RadC	9123	9620	498	forward	hypothetical protein	PF04002 (RadC; aa 44-163).	NA
I23	orf09	unannotated	9634	9843	210	forward	hypothetical protein	NA	NA
I23	orf10	unannotated	9903	10208	306	forward	hypothetical protein	NA	NA
I23	orf11	HTH-containing protein	10195	10542	348	reverse	hypothetical protein	PF01381 (HTH_3; aa 17-71).	NA
I23	orf12	unannotated	10681	11034	354	forward	hypothetical protein	NA	NA
I23	orf13	unannotated	11155	11973	819	forward	hypothetical protein	NA	NA
I23	orf14	DNA topoisomerase	12150	12920	771	reverse	hypothetical protein	PF08378 (NERD; aa 33-145); PF01396 (zf-C4_Topoisom; aa 217-245).	NA
I23	orf15	T5orf172 helicase	13022	14206	1185	reverse	hypothetical protein	PF10544 (T5orf172; aa 263-374).	NA
I23	orf16	DEAD-like helicase	14199	16265	2067	reverse	hypothetical protein	PF00270 (DEAD; aa 28-193).	NA
I23	orf17	unannotated	16268	16372	105	reverse	hypothetical protein	NA	NA
I23	orf18	DNA methyltransferase	16369	19131	2763	reverse	hypothetical protein	PF20464 (MmeI_N; aa 1-159); PF20465 (MmeI_hel; aa 170-250); PF20473 (MmeI_Mtase; aa 325-606); PF20466 (MmeI_TRD; aa 629-832); PF20467 (MmeI_C; aa 834-911).	RM_Type_IIG__Type_IIG.
I23	orf19	inovirus Gp2	19299	20000	702	reverse	hypothetical protein	PF11726 (Inovirus_Gp2; aa 50-231).	NA
I23	orf20 (alpA)	AlpA	20507	20740	234	reverse	hypothetical protein	PF05930 (Phage_AlpA; aa 1-51).	NA
I23	orf21	unannotated	20810	22114	1305	reverse	hypothetical protein	NA	NA
I23	orf22	unannotated	22346	23020	675	forward	hypothetical protein	NA	NA

I44	orf01 (intY)	tyrosine integrase	185	1441	1257	forward	Prophage integrase IntA	PF13356 (Arm-DNA-bind_3; aa 8-94); PF22022 (Phage_int_M; aa 107-200); PF00589 (Phage_integrase; aa 214-377).	NA
I44	orf02	type II REase	1609	2364	756	forward	hypothetical protein	PF01844 (HNH; aa 180-235).	RM_Type_II__Type_II_REases
I44	orf03	type II Mtase	2467	3531	1065	forward	putative BsuMI modification methylase subunit YdiO	PF00145 (DNA_methylase; aa 4-336).	RM_Type_II__Type_II_MTases
I44	orf04	VSR endonuclease	3482	3898	417	reverse	Very short patch repair protein	PF03852 (Vsr; aa 2-64).	NA
I44	orf05	DrmC	3916	4554	639	reverse	Cardiolipin synthase B	PF13091 (PLDc_2; aa 67-200).	DISARM__drmC
I44	orf06	DrmB	4682	6538	1857	reverse	hypothetical protein	PF09369 (MZB; aa 483-583).	DISARM__drmB
I44	orf07	DrmA	6541	10575	4035	reverse	hypothetical protein	NA	DISARM__drmA
I44	orf08	unannotated	10579	10908	330	reverse	hypothetical protein	NA	NA
I44	orf09	DrmMI	10905	14981	4077	reverse	hypothetical protein	PF07669 (Eco57I; aa 845-931).	DISARM_1__drmMI
I44	orf10	DrmD	14978	18103	3126	reverse	RNA polymerase-associated protein RapA	PF00176 (SNF2-rel_dom; aa 134-317); PF00271 (Helicase_C; aa 514-620).	DISARM_1__drmD
I44	orf11	MFS transporter	18695	19495	801	reverse	Glycerol-3-phosphate transporter	PF07690 (MFS_1; aa 29-257).	NA
I44	orf12	alcohol dehydrogenase	19609	20700	1092	reverse	NAD-dependent methanol dehydrogenase	PF00465 (Fe-ADH; aa 8-351).	NA
I44	orf13	avidin	20847	21623	777	reverse	hypothetical protein	PF01382 (Avidin; aa 3-85).	NA
I44	orf14	MFS transporter	21691	23055	1365	reverse	Putative niacin/nicotinamide transporter NaiP	PF07690 (MFS_1; aa 46-378).	NA
I44	orf15	Asp/Glu/hydantoin racemase	23118	23939	822	reverse	Hydantoin racemase	PF01177 (Asp_Glu_race; aa 20-225).	NA
I44	orf16	unannotated	23960	24850	891	reverse	hypothetical protein	NA	NA
I44	orf17	metallopeptidase	24907	26292	1386	reverse	Xaa-Pro aminopeptidase	PF05195 (AMP_N; aa 7-123); PF00557 (Peptidase_M24; aa 173-438).	NA
I44	orf18	hydantoinase B	26320	27978	1659	reverse	Acetophenone carboxylase delta subunit	PF02538 (Hydantoinase_B; aa 5-522).	NA
I44	orf19	hydantoinase A	27992	30076	2085	reverse	Acetophenone carboxylase gamma subunit	PF05378 (Hydant_A_N; aa 9-185); PF01968 (Hydantoinase_A; aa 208-500); PF19278 (Hydant_A_C; aa 514-689).	NA

I44	orf20	HTH-type transcriptional regulator	30253	32856	2604	reverse	HTH-type transcriptional regulator MalT	PF00196 (GerE; aa 802-856).	NA
I44	orf21	unannotated	33248	33571	324	forward	hypothetical protein	NA	NA
I44	orf22	DUF932	33664	34632	969	forward	hypothetical protein	PF06067 (DUF932; aa 77-302).	NA
I44	orf23	YqaJ	34707	35711	1005	forward	hypothetical protein	PF09588 (YqaJ; aa 29-172).	NA
I44	orf24	Gp3-like	35790	36740	951	forward	hypothetical protein	PF18897 (Gp3-like; aa 12-206).	NA
I44	orf25	RadC	36712	37209	498	forward	hypothetical protein	PF04002 (RadC; aa 44-163).	NA
I44	orf26	unannotated	37223	37432	210	forward	hypothetical protein	NA	NA
I44	orf27	unannotated	37534	37779	246	forward	hypothetical protein	NA	NA
I44	orf28	HTH-containing protein	37787	38134	348	reverse	hypothetical protein	PF01381 (HTH_3; aa 17-71).	NA
I44	orf29	unannotated	38272	38658	387	forward	hypothetical protein	NA	NA
I44	orf30	unannotated	38675	39493	819	forward	hypothetical protein	NA	NA
I44	orf31	inovirus Gp2	39957	40916	960	reverse	hypothetical protein	PF11726 (Inovirus_Gp2; aa 149-232).	NA
I44	orf32 (alpA)	AlpA	41252	41482	231	reverse	hypothetical protein	PF05930 (Phage_AlpA; aa 1-51).	NA
I44	orf33	unannotated	41531	42988	1458	reverse	hypothetical protein	NA	NA
I44	orf34	unannotated	43158	43829	672	forward	hypothetical protein	NA	NA
I55	orf01 (intY)	tyrosine integrase	185	1441	1257	forward	Prophage integrase IntA	PF13356 (Arm-DNA-bind_3; aa 8-94); PF22022 (Phage_int_M; aa 107-200); PF00589 (Phage_integrase; aa 214-378).	NA
I55	orf02	dynamin	2594	4903	2310	forward	hypothetical protein	PF00350 (Dynamin_N; aa 59-226).	NA
I55	orf03	dynamin	4914	7166	2253	forward	hypothetical protein	PF00350 (Dynamin_N; aa 40-177).	NA
I55	orf04	unannotated	7235	7738	504	forward	hypothetical protein	NA	NA
I55	orf05	unannotated	7735	8307	573	forward	hypothetical protein	NA	NA
I55	orf06	unannotated	8349	10991	2643	forward	hypothetical protein	NA	NA
I55	orf07	unannotated	11420	12151	732	forward	hypothetical protein	NA	NA
I55	orf08	unannotated	12148	13686	1539	forward	hypothetical protein	NA	NA
I55	orf09	helicase	14030	16783	2754	reverse	RNA polymerase-associated protein RapA	PF04851 (ResIII; aa 36-245); PF00271 (Helicase_C; aa 755-814).	NA
I55	orf10	unannotated	16770	18677	1908	reverse	hypothetical protein	NA	NA
I55	orf11	unannotated	18674	20056	1383	reverse	hypothetical protein	NA	NA
I55	orf12	unannotated	20681	21103	423	forward	hypothetical protein	NA	NA
I55	orf13	DUF1232	21186	21524	339	forward	hypothetical protein	PF06803 (DUF1232; aa 50-84).	NA
I55	orf14	unannotated	21652	22302	651	forward	hypothetical protein	NA	NA
I55	orf15	unannotated	22316	22447	132	reverse	hypothetical protein	NA	NA

I55	orf16	AAA	22605	24695	2091	forward	ATP-dependent zinc metalloprotease FtsH	PF00004 (AAA; aa 255-377, 491-614).	NA
I55	orf17	WYL	24791	25813	1023	forward	hypothetical protein	PF13280 (WYL; aa 156-330).	NA
I55	orf18	unannotated	25859	26395	537	forward	hypothetical protein	NA	NA
I55	orf19	Sir2	26521	27174	654	reverse	NAD-dependent protein deacylase	PF02146 (SIR2; aa 20-196).	NA
I55	orf20	unannotated	27411	27668	258	reverse	hypothetical protein	NA	NA
I55	orf21	Sel1	27658	28299	642	reverse	hypothetical protein	PF08238 (Sel1; aa 96-127, 133-166, 171-200).	NA
I55	orf22	unannotated	28421	28585	165	forward	hypothetical protein	NA	NA
I55	orf23	RecD	28573	30666	2094	reverse	RecBCD enzyme subunit RecD	PF21185 (RecD_N; aa 40-153); PF13245 (AAA_19; aa 205-357); PF13538 (UvrD_C_2; aa 615-662).	NA
I55	orf24	RecB	30663	34337	3675	reverse	RecBCD enzyme subunit RecB	PF00580 (UvrD-helicase; aa 13-452); PF13361 (UvrD_C; aa 459-660); PF13361 (UvrD_C; aa 710-848); PF12705 (PDDEXK_1; aa 1072-1214).	NA
I55	orf25	RecC	34334	37804	3471	reverse	RecBCD enzyme subunit RecC	PF04257 (Exonuc_V_gamma; aa 11-360); PF17946 (RecC_C; aa 835-1081).	NA
I55	orf26	unannotated	38042	38371	330	forward	hypothetical protein	NA	NA
I55	orf27	DUF932	38462	39430	969	forward	hypothetical protein	PF06067 (DUF932; aa 77-302).	NA
I55	orf28	YqaJ	39513	40514	1002	forward	hypothetical protein	PF09588 (YqaJ; aa 28-171).	NA
I55	orf29	Gp3-like	40605	41498	894	forward	hypothetical protein	PF18897 (Gp3-like; aa 12-206).	NA
I55	orf30	unannotated	41554	41742	189	forward	hypothetical protein	NA	NA
I55	orf31	unannotated	41803	42126	324	forward	hypothetical protein	NA	NA
I55	orf32	HTH-containing protein	42136	42483	348	reverse	hypothetical protein	PF01381 (HTH_3; aa 17-71).	NA
I55	orf33	unannotated	42622	42975	354	forward	hypothetical protein	NA	NA
I55	orf34	unannotated	43189	43812	624	forward	hypothetical protein	NA	NA
I55	orf35	MUG113	43980	45188	1209	reverse	hypothetical protein	PF13455 (MUG113; aa 285-373).	NA
I55	orf36	DEAD-like helicase	45181	47247	2067	reverse	hypothetical protein	PF00270 (DEAD; aa 28-193).	NA
I55	orf37	DNA methyltransferase	47349	50105	2757	reverse	hypothetical protein	PF20464 (MmeI_N; aa 1-159); PF20465 (MmeI_hel; aa 170-250); PF20473 (MmeI_Mtase; aa 325-605); PF20466 (MmeI_TRD; aa 627-826); PF20467 (MmeI_C; aa 827-909).	RM_Type_IIG__Type_IIG
I55	orf38	inovirus Gp2	50350	51321	972	reverse	hypothetical protein	NA	NA
I55	orf39 (alpA)	AlpA	51661	51888	228	reverse	hypothetical protein	PF05930 (Phage_AlpA; aa 1-51).	NA
I55	orf40	unannotated	51950	53260	1311	reverse	hypothetical protein	NA	NA

I55	orf41	unannotated	53417	54289	873	forward	hypothetical protein	NA	NA

- **ORF name**: This is the name given to a gene on an MGE. On each MGE, genes are named serially and consecutively from 5'-end (near tyrosine integrase) to 3'-end in the format *orf#*, where # is a two-digit number. The tyrosine integrase always has #=01, the genes following it have #=02, 03, 04, etc. For some well-recognisable genes, common gene names are indicated inside round brackets following *orf#*.
- **Consensus annotation**: This is the annotation given to a gene after manual checks on Prokka and Pfam predictions. Genes are called "unannotated" when neither predictors find results.
- **Minimum coordinate**: This is the position of the first nucleotide of a gene. For each MGE, the nucleotides are numbered from the 5'-end, with the first nucleotide immediately downstream of the 14 bp repeat (near the 3'-end of *tmRNA*) labelled as 1.
- **Maximum coordinate**: This is the position of the last nucleotide of a gene.
- **Length**: This is the number of nucleotides in a gene, and is equal to Maximum coordinate - Minimum coordinate + 1.
- **Direction of transcription**: This is the direction of transcription of a gene. Genes labelled as "forward" are transcribed from the 5'-end to the 3'-end along the annotated strand (i.e., the strand where the tyrosine integrase is at the 5'-end). Genes labelled as "forward" are transcribed from the 5'-end to the 3'-end along the other stand.
- **Prokka product**: This is the protein product predicted by Prokka in KBase. Proteins without predictions are labelled as "hypothetical protein".
- **Pfam domains**: This is the protein domain(s) predicted by Pfam in InterProScan. For each predicted domain, the Pfam Id is listed, with the corresponding Pfam name and mapped region (given by the interval of amino acids along the amino acid sequence) given in round brackets. For proteins with multiple predicted domains, all domains are listed and separated by semicolons. Proteins with no predicted domains are labelled as "NA".
- **DefenseFinder prediction**: This is the gene name of putative phage defence genes predicted by DenfenseFinder. Genes not predicted to be involved in phage defence activities are labelled as "NA".

1049　Table S5. Mutations detected in isolates with MGEs.

1050

Strain number	Genotype	Gene locus	Genome coordinate	Type of mutation	Change	CDS position	Effect on protein
MPB38392	SBW25 *tmRNA*::I23 *glmS*::Tn*7*-152(GFP, Kan^R) isolate1	*pflu4414*	4878944-4878945	insertion (tandem repeat)	T(2) -> T(3)	1489-1490	frameshift
MPB38393	SBW25 *tmRNA*::I23 *glmS*::Tn*7*-152(GFP, Kan^R) isolate2	*pflu4414 - pflu4415*	4880381-4880570	deletion	del 190 bp	*pflu4414*: 1-53; *pflu4415*: 664-789	*pflu4415*: truncation; *pflu4414*: no translation
MPB38394	SBW25 *tmRNA*::I23 *glmS*::Tn*7*-152(GFP, Kan^R) isolate3	*pflu4414*	4880259	SNP (transition)	G -> A	175	truncation
MPB38395	SBW25 *tmRNA*::I23 *glmS*::Tn*7*-152(GFP, Kan^R) isolate4	*pflu3022*	3295100-3295889	deletion	del 790 bp	387-1176	truncation
		pflu4414	4879083-4879084	insertion (tandem repeat)	T(5) -> T(6)	1350-1351	frameshift
MPB38396	SBW25 *tmRNA*::I23 *glmS*::Tn*7*-152(GFP, Kan^R) isolate5	*pflu4414 - pflu4415*	4880381-4880570	deletion	del 190 bp	*pflu4414*: 1-53; *pflu4415*: 664-789	*pflu4415*: truncation; *pflu4414*: no translation
MPB38895	SBW25 *tmRNA*::I23 *glmS*::Tn*7*-155(RFP, Kan^R) isolate6	None	-	-	-	-	-
MPB38906	SBW25 *tmRNA*::I44 *glmS*::Tn*7*-155(RFP, Kan^R) isolate1	*pflu4744*	5217912	SNP (transition)	G -> A	116	substitution (S39N)
MPB38907	SBW25 *tmRNA*::I44 *glmS*::Tn*7*-155(RFP, Kan^R) isolate2	*pflu4744*	5217912	SNP (transition)	G -> A	116	substitution (S39N)
MPB33357	SBW25 *tmRNA*::I55 *glmS*::Tn*7*-152(GFP, Kan^R) isolate1	*pflu3805*	4199776	SNP (transversion)	A -> T	33	substitution (N11K)
		pflu4414	4880277-4880278	insertion (tandem repeat)	C(5) -> C(6)	157-158	frameshift

1051

1052

Table S6. Rank abundance of genes carried by similar MGEs

Domains	COG category	Description	Counts
Phage_integrase	L	Belongs to the 'phage' integrase family	111
AlpA	K	transcriptional regulator	109
Inovirus_Gp2	S	Inovirus Gp2	101
WYL	K	regulation of single-species biofilm formation	87
DUF932	S	Domain of unknown function (DUF932)	80
YqaJ	L	COG5377 Phage-related protein	78
RadC	L	RadC-like JAB domain	66
DUF1524,DUF262	S	Protein of unknown function DUF262	42
HsdM_N,N6_Mtase	V	Type I restriction-modification system methyltransferase subunit	40
Methylase_S	V	PFAM Type I restriction modification DNA specificity domain	31
N6_Mtase	V	Type II restriction enzyme, methylase	24
DUF1788	S	Domain of unknown function (DUF1788)	23
AAA_5	V	GTPase subunit of restriction endonuclease	20
Dynamin_N	T	ATPase. Has a role at an early stage in the morphogenesis of the spore coat	20
HTH_3,HTH_31	K	sequence-specific DNA binding	20
MMR_HSR1	P	Dynamin family	17
ADP_ribosyl_GH	O	ADP-ribosylglycohydrolase	16
DNA_methylase	H	Belongs to the class I-like SAM-binding methyltransferase superfamily. C5-methyltransferase family	16
DUF262	S	Protein of unknown function DUF262	16
Helicase_C,SNF2_N	KL	Transcription regulator that activates transcription by stimulating RNA polymerase (RNAP) recycling in case of stress conditions such as supercoiled DNA or high salt concentrations. Probably acts by releasing the RNAP, when it is trapped or immobilized on tightly supercoiled DNA. Does not activate transcription on linear DNA. Probably not involved in DNA repair	15
Vsr	L	May nick specific sequences that contain T G mispairs resulting from m5C-deamination	15
EcoR124_C,HSDR_N,ResIII	L	Subunit R is required for both nuclease and ATPase activities, but not for modification	14
Lon_2,Lon_C	O	Putative ATP-dependent Lon protease	14
PDDEXK_1,UvrD-helicase,UvrD_C	L	A helicase nuclease that prepares dsDNA breaks (DSB) for recombinational DNA repair. Binds to DSBs and unwinds DNA via a highly rapid and processive ATP-dependent bidirectional helicase activity. Unwinds dsDNA until it encounters a Chi (crossover hotspot instigator) sequence from the 3' direction. Cuts ssDNA a few nucleotides 3' to the Chi site. The properties and activities of the enzyme are changed at Chi. The Chi-altered holoenzyme produces a long 3'-ssDNA overhang and facilitates RecA-binding to the ssDNA for homologous DNA recombination and repair. Holoenzyme degrades any linearized DNA that is unable to undergo homologous recombination. In the holoenzyme this subunit contributes ATPase, 3'-5' helicase, exonuclease activity and loads RecA onto ssDNA	14
Helicase_C	KL	Helicase conserved C-terminal domain	13
HTH_1,LysR_substrate	K	transcriptional regulator	13

McrBC	V	McrBC 5-methylcytosine restriction system component	13
HSDR_N,ResIII	L	Subunit R is required for both nuclease and ATPase activities, but not for modification	12
HTH_17	K	TIGRFAM DNA binding domain, excisionase family	12
DUF1998	S	Domain of unknown function (DUF1998)	11
PglZ	H	PglZ domain	11
Ank_2,Ank_4,Ank_5	G	response to abiotic stimulus	10
N6_N4_Mtase	H	DNA methylase N-4 N-6	10
ResIII	KL	Type III restriction enzyme, res subunit	10
DEAD,DUF1998,Helicase_C	L	COG1205 Distinct helicase family with a unique C-terminal domain including a metal-binding cysteine cluster	9
DUF1819	S	Putative inner membrane protein (DUF1819)	9
DUF45	S	Metal-dependent hydrolase	9
HTH_Tnp_1	L	Transposase and inactivated derivatives	9
Lon_2,Lon_C,MIT_C	O	ATP-dependent Lon-type protease	9
PLDc_2	I	phospholipase d transphosphatidylase	9
AAA_15	L	AAA ATPase domain	8
BPD_transp_1	P	Binding-protein-dependent transport system inner membrane component	8
DNA_processg_A	LU	DNA recombination-mediator protein A	8
DUF697,MMR_HSR1	D	50S ribosome-binding GTPase	8
Transp_cyt_pur	FH	Permease for cytosine/purines, uracil, thiamine, allantoin	8
AAA_15,AAA_21,AAA_23	T	AAA domain, putative AbiEii toxin, Type IV TA system	7
DEAD,Helicase_C,RecQ_Zn_bind	L	DNA helicase	7
Eco57I,N6_Mtase	V	COG1002 Type II restriction enzyme, methylase subunits	7
AAA_11,AAA_12,AAA_30,PDDEXK_1,RNase_H_2	L	RNase_H superfamily	6
AAA_19,AAA_30,UvrD_C_2	L	A helicase nuclease that prepares dsDNA breaks (DSB) for recombinational DNA repair. Binds to DSBs and unwinds DNA via a highly rapid and processive ATP-dependent bidirectional helicase activity. Unwinds dsDNA until it encounters a Chi (crossover hotspot instigator) sequence from the 3' direction. Cuts ssDNA a few nucleotides 3' to the Chi site. The properties and activities of the enzyme are changed at Chi. The Chi-altered holoenzyme produces a long 3'-ssDNA overhang and facilitates RecA-binding to the ssDNA for homologous DNA recombination and repair. Holoenzyme degrades any linearized DNA that is unable to undergo homologous recombination. In the holoenzyme this subunit has ssDNA-dependent ATPase and 5'-3' helicase activity. When added to pre-assembled RecBC greatly stimulates nuclease activity and augments holoenzyme processivity. Negatively regulates the RecA-loading ability of RecBCD	6
AAA_23	L	ATPase involved in DNA repair	6
ABC_tran,DUF726,Dynamin_N	S	ATPase. Has a role at an early stage in the morphogenesis of the spore coat	6
CxxCxxCC	S	PFAM Uncharacterised protein family (UPF0153)	6
DUF2357,PDDEXK_7	S	PD-(D/E)XK nuclease superfamily	6

DUF445	S	membrane	6
EH_Signature	S	EH_Signature domain	6
Exonuc_V_gamma	L	A helicase nuclease that prepares dsDNA breaks (DSB) for recombinational DNA repair. Binds to DSBs and unwinds DNA via a highly rapid and processive ATP-dependent bidirectional helicase activity. Unwinds dsDNA until it encounters a Chi (crossover hotspot instigator) sequence from the 3' direction. Cuts ssDNA a few nucleotides 3' to the Chi site. The properties and activities of the enzyme are changed at Chi. The Chi-altered holoenzyme produces a long 3'-ssDNA overhang and facilitates RecA-binding to the ssDNA for homologous DNA recombination and repair. Holoenzyme degrades any linearized DNA that is unable to undergo homologous recombination. In the holoenzyme this subunit recognizes the wild- type Chi sequence, and when added to isolated RecB increases its ATP-dependent helicase processivity	6
Helicase_C,Mrr_cat,SNF2_N, SWIM	KL	COG0553 Superfamily II DNA RNA helicases, SNF2 family	6
PDDEXK_1	L	PD-(D/E)XK nuclease superfamily	6
Amidohydro_3	F	cytosine deaminase	5
DDE_Tnp_1,DDE_Tnp_1_6,DUF772	L	Transposase DDE domain	5
DUF1232	S	Protein of unknown function (DUF1232)	5
DUF3987	T	Protein of unknown function (DUF3987)	5
Helicase_C,ResIII,SNF2_N	L	PFAM Helicase conserved C-terminal domain	5
HNH_2	V	restriction endonuclease	5
Hydant_A_N,Hydantoinase_A	EQ	N-methylhydantoinase A acetone carboxylase, beta subunit	5
Hydantoinase_B	EQ	Hydantoinase B/oxoprolinase	5
Isochorismatase	Q	Isochorismatase family	5
MFS_1	EGP	Major Facilitator Superfamily	5
MFS_1,Sugar_tr	EGP	Major facilitator superfamily	5
MotB_plug,OmpA	N	Membrane MotB of proton-channel complex MotA/MotB	5
PDDEXK_7	S	PD-(D/E)XK nuclease superfamily	5
UvrD-helicase,UvrD_C,UvrD_C_2	L	Uvrd rep helicase	5
Virulence_RhuM	S	COG3943 Virulence protein	5
AAA	O	COG0464 ATPases of the AAA class	4
AAA_13	L	AAA domain	4
AAA_5,DUF3578	V	Domain of unknown function (DUF3578)	4
ABC_tran,TOBE_2	P	Belongs to the ABC transporter superfamily	4
DAO	E	FAD dependent oxidoreductase	4
DUF2950	N	Protein of unknown function (DUF2950)	4
DUF3077	S	Protein of unknown function (DUF3077)	4
DUF4357	S	Domain of unknown function (DUF4357)	4

Fer2_4	C	2Fe-2S iron-sulfur cluster binding domain	4
Fer2_BFD,Pyr_redox_2	C	FAD-dependent pyridine nucleotide-disulphide oxidoreductase	4
Helicase_C,ResIII	L	Type III restriction protein res subunit	4
His_Phos_1,NUDIX	L	nUDIX hydrolase	4
HTH_30,PucR	QT	transcriptional regulator	4
Ketoacyl-synt_C,ketoacyl-synt	IQ	catalyzes a condensation reaction in fatty acid biosynthesis addition of an acyl acceptor of two carbons from malonyl-ACP	4
Pkinase	KLT	Protein tyrosine kinase	4
Ribonuc_L-PSP	J	Endoribonuclease L-PSP	4
SBP_bac_6,SBP_bac_8	E	Bacterial extracellular solute-binding protein	4
AAA_11,AAA_12,NERD,Pkinase	L	AAA domain	3
Arm-DNA-bind_3,Phage_integrase	L	Belongs to the 'phage' integrase family	3
DDE_Tnp_IS66,DDE_Tnp_IS66_C,LZ_Tnp_IS66,zf-IS66	S	zinc-finger binding domain of transposase IS66	3
DEAD,Helicase_C	L	helicase superfamily c-terminal domain	3
DJ-1_PfpI,HTH_18	K	COG4977 Transcriptional regulator containing an amidase domain and an AraC-type DNA-binding HTH domain	3
DUF3387,HSDR_N,ResIII	F	Subunit R is required for both nuclease and ATPase activities, but not for modification	3
Helicase_C,PLDc_2,SNF2_N	L	SNF2 family N-terminal domain	3
HTH_21,rve	L	Transposase and inactivated derivatives	3
Mrr_cat	V	Restriction endonuclease	3
NA37	S	37-kD nucleoid-associated bacterial protein	3
RelA_SpoT	S	Region found in RelA / SpoT proteins	3
RHH_1	S	Ribbon-helix-helix protein, copG family	3
Sel1	S	Sel1-like repeats.	3
TnpB_IS66	L	Transposase	3
AAA_15,AAA_21,ABC_tran	L	(ABC) transporter	2
AAA_19,NERD,UvrD_C_2	L	COG0210 Superfamily I DNA and RNA helicases	2
AAA_19,UvrD-helicase	L	UvrD/REP helicase N-terminal domain	2
AAA_19,UvrD-helicase,UvrD_C,UvrD_C_2	L	UvrD/REP helicase N-terminal domain	2
Acyl-CoA_dh_1,Acyl-CoA_dh_M,Acyl-CoA_dh_N	I	Acyl-CoA dehydrogenase, middle domain	2
adh_short_C2	IQ	PFAM Short-chain dehydrogenase reductase SDR	2
AIPR	S	AIPR protein	2
Aldedh	C	Aldehyde dehydrogenase family	2

CoA_transf_3	C	CoA-transferase family III	2
DUF1837	S	Domain of unknown function (DUF1837)	2
DUF3223	S	Protein of unknown function (DUF3223)	2
DUF3883	O	Molecular chaperone. Has ATPase activity	2
DUF4868	S	Domain of unknown function (DUF4868)	2
DUF726	S	Protein of unknown function (DUF726)	2
EcoEI_R_C,HSDR_N,Helicase_C,ResIII	L	EcoEI R protein C-terminal	2
GAF,HATPase_c,HisKA,PAS_2,PHY	T	PAS fold	2
HATPase_c_3	L	Histidine kinase-, DNA gyrase B-, and HSP90-like ATPase	2
HNH	L	HNH nucleases	2
HpcH_HpaI	G	HpcH/HpaI aldolase/citrate lyase family	2
HTH_29,rve,rve_2,rve_3	L	COG2801 Transposase and inactivated derivatives	2
HTH_31	K	Helix-turn-helix XRE-family like proteins	2
HxlR	K	transcriptional regulator	2
MaoC_dehydrat_N,MaoC_dehydratas	I	N-terminal half of MaoC dehydratase	2
MUG113,T5orf172	P	T5orf172	2
NmrA	GM	NmrA family	2
PDDEXK_1,Peptidase_M78,UvrD-helicase,UvrD_C	L	Unwinds DNA duplexes with 3' to 5' polarity with respect to the bound strand and initiates unwinding most effectively when a single-stranded region is present	2
Plug,TonB_dep_Rec	P	TonB dependent receptor	2
PRiA4_ORF3	L	Plasmid pRiA4b ORF-3-like protein	2
RES	S	RES	2
Response_reg	T	response regulator	2
SIR2_2	F	SIR2-like domain	2
UvrD_C_2	L	UvrD-like helicase C-terminal domain	2
3HCDH,3HCDH_N	I	3-hydroxyacyl-coa dehydrogenase	1
AAA_11,AAA_12,DUF3320,DUF4011,DUF559	L	AAA domain	1
AAA_16,GAF,GAF_2,HATPase_c,HisKA,PAS_3,Pkinase	T	Histidine kinase	1
AAA_18,AAA_33,CPT	S	AAA domain	1
AAA_19,UvrD-helicase,UvrD_C	L	DNA helicase	1
AAA_21	S	AAA domain, putative AbiEii toxin, Type IV TA system	1

AAA_30,UvrD_C_2	L	PIF1-like helicase	1
Abhydrolase_1,Abhydrolase_6	I	Alpha beta hydrolase	1
Abhydrolase_1,Hydrolase_4	I	hydrolases or acyltransferases (alpha beta hydrolase superfamily)	1
ABM	S	Antibiotic biosynthesis monooxygenase	1
AcetylCoA_hyd_C,AcetylCoA_hydro	C	acetyl-CoA hydrolase	1
Acetyltransf_1,Acetyltransf_10,tRNA_m1G_MT	K	acetyltransferase	1
ACR_tran	V	TIGRFAM transporter, hydrophobe amphiphile efflux-1 (HAE1) family	1
Acyl-CoA_dh_1,Acyl-CoA_dh_N	I	acyl-CoA dehydrogenase	1
Adenylsucc_synt	F	Adenylosuccinate synthetase	1
ADH_N,ADH_zinc_N	C	Zinc-binding dehydrogenase	1
ADH_N,ADH_zinc_N,ADH_zinc_N_2	C	alcohol dehydrogenase	1
ADH_N_2,ADH_zinc_N	S	N-terminal domain of oxidoreductase	1
adh_short	S	Belongs to the short-chain dehydrogenases reductases (SDR) family	1
adh_short,adh_short_C2	IQ	KR domain	1
Amidohydro_2	S	Amidohydrolase	1
AMP_N,Peptidase_M24	E	Aminopeptidase P, N-terminal domain	1
ApbA,ApbA_C	H	Ketopantoate reductase PanE/ApbA C terminal	1
AraC_binding,HTH_18	K	Transcriptional regulator	1
Arginase	E	Belongs to the arginase family	1
Biotin_lipoyl_2,HlyD_3,HlyD_D23	M	Belongs to the membrane fusion protein (MFP) (TC 8.A.1) family	1
CitMHS	C	TIGRFAM citrate H symporter, CitMHS family	1
Cytochrome_CBB3	C	photosynthesis	1
DDE_3,HTH_23,HTH_32,HTH_33	L	COG3335 Transposase and inactivated derivatives	1
DDE_Tnp_1	L	PFAM Protein	1
DDE_Tnp_1,DUF772	L	COG3039 Transposase and inactivated derivatives, IS5 family	1
DDE_Tnp_1,Methylase_S	V	type I restriction modification DNA specificity domain	1
DEAD,Helicase_C,Pribosyltran,RecQ_Zn_bind,UvrD-helicase,UvrD_C	L	helicase superfamily c-terminal domain	1
DJ-1_PfpI	S	DJ-1/PfpI family	1

DoxX_2	S	DoxX-like family	1
DUF1302	M	Protein of unknown function (DUF1302)	1
DUF2236	S	Uncharacterized protein conserved in bacteria (DUF2236)	1
DUF2867	S	Protein of unknown function (DUF2867)	1
DUF2938	S	Protein of unknown function (DUF2938)	1
DUF2946	S	Protein of unknown function (DUF2946)	1
DUF4113,IMS,IMS_C,IMS_HHH	L	DNA polymerase	1
DUF4420	S	Putative PD-(D/E)XK family member, (DUF4420)	1
DUF928,SIR2_2	F	5-carbamoylmethyl uridine residue modification	1
Dynamin_N,MMR_HSR1,Patatin	M	Esterase of the alpha-beta hydrolase superfamily	1
EcoEI_R_C,Helicase_C,ResIII	L	EcoEI R protein C-terminal	1
EcoRII-C	L	EcoRII C terminal	1
ETF	C	Electron transfer flavoprotein	1
ETF,ETF_alpha	C	Electron transfer flavoprotein	1
ETF_QO,FAD_binding_2,NAD_binding_8,Thi4	C	Electron transfer flavoprotein-ubiquinone oxidoreductase	1
Fe-ADH	C	Iron-containing alcohol dehydrogenase	1
Fer4_12,Fer4_14,Radical_SAM	S	Radical SAM superfamily	1
Fer4_12,Radical_SAM	S	Radical SAM superfamily	1
Flg_new,LRR_5,LRR_6	S	regulation of response to stimulus	1
GAF,GAF_2,GGDEF,PAS_3	T	GAF PAS GGDEF domain-containing protein	1
GerE	K	helix_turn_helix, Lux Regulon	1
GerE,Response_reg,Trans_reg_C	K	response regulator	1
GntR,UTRA	K	UTRA	1
Histone_HNS	-	-	1
HTH_11,WYL	K	HTH domain	1
HTH_18	K	AraC Family	1
HTH_20	K	transcriptional regulator	1
HTH_21,rve,rve_3	L	PFAM Integrase catalytic region	1
HTH_23,HTH_24,rve	L	Transposase and inactivated derivatives	1
HTH_26	K	transcriptional regulator	1
HTH_3,Peptidase_M78	K	IrrE N-terminal-like domain	1

HTH_IclR,IclR	K	transcriptional regulator	1
Integrase_1,Phage_int_SAM_2	L	Belongs to the 'phage' integrase family	1
Lactamase_B	S	Metallo-beta-lactamase superfamily	1
Lumazine_bd_2	S	Putative lumazine-binding	1
LysR_substrate	K	LysR substrate binding domain	1
M20_dimer,Peptidase_M20	S	Peptidase dimerisation domain	1
MaoC_dehydratas	I	MaoC like domain	1
MS_channel,cNMP_binding	MT	mechanosensitive ion channel	1
N6_N4_Mtase,TypeIII_RM_meth	L	DNA methylase	1
NAD_binding_10,NmrA	GM	NmrA-like family	1
NERD,zf-C4_Topoisom	L	COG0551 Zn-finger domain associated with topoisomerase type I	1
NUDIX	L	NUDIX domain	1
OEP	M	RND efflux system, outer membrane lipoprotein	1
OmpA	N	Flagellar Motor Protein	1
OsmC	O	response to oxidative stress	1
Oxidored_FMN	C	PFAM NADH flavin oxidoreductase NADH oxidase	1
PCuAC	S	Copper chaperone PCu(A)C	1
PDDEXK_4	S	PD-(D/E)XK nuclease superfamily	1
PKD,Tannase	C	PFAM blue (type 1) copper domain protein	1
Prok-JAB	S	Prokaryotic homologs of the JAB domain	1
PrpF	P	protein conserved in bacteria	1
Radical_SAM	C	Elongator protein 3, MiaB family, Radical SAM	1
Radical_SAM,TRAM,UPF0004	J	Catalyzes the methylthiolation of an aspartic acid residue of ribosomal protein S12	1
RloB	S	RloB-like protein	1
RraA-like	H	Catalyzes the aldol cleavage of 4-hydroxy-4-methyl-2- oxoglutarate (HMG) into 2 molecules of pyruvate. Also contains a secondary oxaloacetate (OAA) decarboxylase activity due to the common pyruvate enolate transition state formed following C-C bond cleavage in the retro-aldol and decarboxylation reactions	1
SCO1-SenC	S	SCO1/SenC	1
SIR2	K	NAD-dependent lysine deacetylase and desuccinylase that specifically removes acetyl and succinyl groups on target proteins. Modulates the activities of several proteins which are inactive in their acylated form	1
TauD	Q	Taurine catabolism dioxygenase TauD, TfdA family	1
Tautomerase_2	S	tautomerase	1
TetR_C_13,TetR_N	K	transcriptional regulator	1
TetR_N	K	Bacterial regulatory proteins, tetR family	1

ThiF	H	ThiF family	1
Thiolase_C,Thiolase_N	I	Acetyl-CoA acetyltransferase	1
Thioredoxin	O	OST3 / OST6 family, transporter family	1
Thymidylate_kin	F	Phosphorylation of dTMP to form dTDP in both de novo and salvage pathways of dTTP synthesis	1
Transposase_mut	L	Transposase	1
tRNA_anti-like	S	tRNA_anti-like	1
UvrD-helicase,UvrD_C	L	UvrD-like helicase C-terminal domain	1
Z1	L	Z1 domain	1

1055

1056

a

(i) Live filtrate ✓

Compost wash → Inoculation → Incubation → Transfer → Incubation → Transfer → Incubation ······

Filtration → Addition

(ii) Co-culture ✗

Compost wash → *Pseudomonas fluorescens* SBW25 → Inoculation → Incubation → Transfer → Incubation → Transfer → Incubation ······

Inoculation

Addition ← Addition

(iii) Fixed filtrate ✓

Compost wash → Inoculation → Incubation → Filtration & storage

b

SBW25 + community filtrate → Colony isolation → Whole-genome NGS → raw reads → Mapping to reference → SBW25 genome / unmapped reads → *de novo* assembly → foreign element → Ends comparison → SBW25 genome + foreign element → PCR over junction

c

Foreign element	Length / bp	Community filtrate	Regime	Frequency
I23	23,199	Compost B	Fixed filtrate	3 lines in 10 lines (1, 1 and 4 out of 24 isolates)
I44	44,236	Compost B	Fixed filtrate	1 line in 10 lines (2 out of 24 isolates)
I55	54,419	Compost C	Live filtrate	1 line in 8 lines (1 out of 3 isolates)

d

foreign element

Integration

CCCCGGCTCCACCA

SBW25 Genome: *tmRNA* — PFI-12

CCCCGGCTCCACCA *Drawn not to scale*

e

Legend:
- tyrosine integrase
- AlpA
- RadC
- T5orf172 helicase
- YqaJ
- HTH-containing protein
- DUF932
- inovirus Gp2
- DNA methyltransferase
- Gp3-like
- phage defence system
- other annotated protein
- unannotated protein

blastn identity 100% — 69%

5 kb

a Δ*intY* GFP | RFP | Coculturing +/− community filtrate | Count GFP-RFP ratio at the start and end of incubation

b Relative fitness of I55 to WT — Filtrate: +, −

c dilution 10^0 — 10^{-2}; I55Δ*intY*; WT

d PFU per mL — I55-Δ*intY*, WT **

e Relative fitness of I55 to WT — P1, P2, Ctrl

f (i) PFU per mL — I55-Δ*intY*, WT **; (ii) PFU per mL — I55-Δ*intY*, WT **