β-catenin localization in the ctenophore *Mnemiopsis leidyi* suggests an ancestral role in cell adhesion and nuclear function

Authors: Brian M. Walters[1]*, Lucas J. Guttieres[2]*, Mayline Goëb[2], Stanley J. Marjenberg[1], Mark Q. Martindale[2], Athula H. Wikramanayake[1#]
1 Department of Biology, University of Miami, 1301 Memorial Dr., Coral Gables, Fl 33124, USA.
2 The Whitney Laboratory for Marine Bioscience, and the Department of Biology, University of Florida, 9505 N, Ocean Shore Blvd, Saint Augustine, Fl 32080-8610, USA.
* These two authors contributed equally to the work
# Corresponding author: athula@miami.edu

Abstract: The emergence of multicellularity in animals marks a pivotal evolutionary event which was likely enabled by molecular innovations in the way cells adhere and communicate with one another. β-catenin is significant to this transition due to its dual role as both a structural component in the cadherin-catenin complex and as a transcriptional coactivator involved in the Wnt/β-catenin signaling pathway. However, our knowledge of how this protein functions in ctenophores, one of the earliest diverging metazoans, is limited. To study β-catenin function in the ctenophore *Mnemiopsis leidyi,* we generated affinity-purified polyclonal antibodies targeting *Ml*β-catenin. We then used this tool to observe β-catenin protein localization in developing *Mnemiopsis* embryos. In this article we provide evidence of consistent β-catenin protein enrichment at cell-cell interfaces in *Mnemiopsis* embryos, suggesting that *Ml*β-catenin has an ancestral role in cell adhesion. Additionally, we found β-catenin enrichment in some nuclei, particularly restricted to the oral pole around the time of gastrulation, suggesting *Ml*β-catenin may have nuclear function in *Mnemiopsis* embryos. The *Ml*β-catenin affinity-purified antibodies now provide us with a powerful reagent to study the ancestral functions of β-catenin in cell adhesion and transcriptional regulation.

1. Introduction

The origin of multicellularity in animals represents a major evolutionary transition, marking the divergence from single-celled ancestors to complex organisms composed of multiple, specialized cell types. Understanding the mechanisms that evolved to enable this transition will provide insights into the fundamental principles that underpin animal evolution and the diversity of life forms we see today. Critical innovations such as cell-cell adhesion, cell-extracellular matrix interactions, cell-cell communication, and cell fate specification are hypothesized to have facilitated cellular cooperation and the partitioning of roles in the cells of the metazoan last common ancestor (Richter and King 2013; Ros-Rocher et al. 2021). β-catenin is a central molecule to many of these cellular processes and thus stands out as a key protein for understanding this transition (Shapiro 1997; Valenta et al. 2012).

β-catenin is a structurally and functionally conserved protein found to be important for developmental processes and maintaining tissue homeostasis across many metazoan lineages. It is well established that β-catenin plays two important roles: acting as a structural component of the cadherin-catenin complex (CCC) that maintains cell-cell adhesion through adherens junctions (AJs) and as a transcriptional coactivator in the Wnt/β-catenin signaling pathway (reviewed in Valenta et al. 2012). Because of its dual purpose, β-catenin has been identified as a key component involved in cell adhesion, cell fate

specification, neurogenesis, spindle orientation, cell migration, cell polarity, and maintenance of stem cells (Valenta et al. 2012).

In its role in cell-cell adhesion, β-catenin interacts with cadherin, a calcium-dependent transmembrane glycoprotein that links neighboring cells together by forming dimers in the extracellular space (Nollet et al. 2000). Cadherins alone are not sufficient in the establishment of a stable AJ (Ishiyama and Ikura 2012). The intracellular domain of classical cadherins includes binding sites for two catenins: p120 and β-catenin which are essential for stabilizing this complex at the plasma membrane (Fig. 1, Peifer et al. 1992; Orsulic et al. 1999; Huber and Weis 2001; Davis et al. 2003). In the absence of binding, members of the CCC are degraded through multiple mechanisms, destabilizing the AJs and leading to a loss of adhesion between cells (Hinck et al. 1994; Wu and Hirsch, 2009; Kourtidis et al. 2013). This interaction plays a pivotal role in maintaining tissue integrity and organization in multicellular organisms (Meng and Takeichi 2009). Thus, β-catenin is essential in cell-cell adhesion through the formation and maintenance of AJs by interacting with cadherins.

On the cytoplasmic side of the membrane, β-catenin has a role in recruiting α-catenin (α-cat) to the CCC, enabling interaction between the plasma membrane and the actin cytoskeleton. This interaction is necessary for cell adhesion and provides the mechanical force that maintains tissue integrity (reviewed in Lecuit and Yap 2015). In turn, this mechanical force exerted on the CCC is at the heart of the mechano-transduction process, which is characterized by the translation of a mechanical stress into a biochemical message. This mechanically regulated cadherin/β-catenin signaling pathway is notably involved in the specification of the anterior endoderm in *D. melanogaster* (Farge 2003; Desprat et al. 2008) and in the activation of key genes in mesoderm specification by β-catenin in zebrafish and Drosophila (Brunet et al. 2013). Furthermore, there is evidence that selective nuclearization of CCC associated β-catenin can be induced through tension-relaxation events, leading to β-catenin facilitated transcription (Röper et al. 2018; Gayrard et al. 2018).

Fig. 1. The dual role of β-catenin in cell adhesion and transcriptional activation. LRP, Low-density lipoprotein Receptor-related Protein; APC, Adenomatous Polyposis Coli; CK1α, Casein Kinase 1α; GSK-3β, Glycogen Synthase Kinase 3β; TCF, T-Cell Factor; LEF, Lymphoid Enhancer Factor; Dsh, Dishevelled; β-cat, β-catenin; α-cat, α-catenin.

Furthermore, β-catenin plays an important role in the Wnt/β-catenin signaling pathway, which can be maintained in two states: 'off' and 'on' (Fig. 1; Nusse and Clevers 2017). In the absence of the Wnt ligand, the pathway remains 'off'. Here a multi-protein complex, known as the destruction complex, regulates cytoplasmic β-catenin stability. Within this complex, Axin and Adenomatous Polyposis Coli (APC) serve as recruitment and scaffolding proteins while Casein Kinase 1 alpha (CK1α) and Glycogen Synthase Kinase-3β (GSK-3β) sequentially phosphorylate β-catenin, marking it for degradation through the proteasome pathway (Aberle et al. 1997; Stamos and Weis, 2013). On the other hand, when the Wnt ligand is present the pathway shifts to the 'on' state. The Wnt ligand binds to the Frizzled (Fzd) and Low-density lipoprotein Receptor-related Protein (LRP5/6) transmembrane co-receptors which then each respectively recruit Dishevelled (Dsh) and Axin to the cell membrane. Receptor inhibition of GSK-3β and Axin phosphorylation interfere with the destruction complex's enzymatic activity, allowing cytoplasmic β-catenin to accumulate. At a high enough concentration, β-catenin is able to translocate to the nucleus where it binds to T-Cell Factor/Lymphoid Enhancer Factor (TCF/LEF) transcription factors and acts as a transcriptional coactivator. The ultimate outcome of β-catenin nuclearization during the 'on' state of the pathway is the transcription of downstream target gene expression. Mutations and altered expression patterns of this pathway have been linked to human diseases including cancer (Clevers 2006; Klaus and Birchmeier 2008).

Most of our understanding of β-catenin's functional role in metazoan biology comes from bilaterian and cnidarian models (Wikramanayake et al. 2003; Martindale and Hejnol 2009; van der Wal and van Amerongen 2020). While these studies are important for establishing an understanding of these highly

conserved processes, investigating earlier diverging taxa will provide valuable insights into the evolutionary origins of these mechanisms. For instance, the role of β-catenin in either a cell signaling or cell-cell adhesion context in ctenophores remains poorly characterized. Several lines of evidence now indicate that the ctenophores, or comb jellies, were the earliest emerging metazoan clade (Dunn et al. 2008; Hejnol et al. 2009; Whelan et al. 2017; Schultz et al. 2023). Ctenophores have a small genome (for example, the *Mnemiopsis leidyi* genome is roughly 150 megabases) with a reduced molecular repertoire relative to other metazoans, which simplifies the identification of conserved proteins and interactions necessary for molecular function. For example, initial *in silico* analysis suggests that key interacting proteins such as Cadherin and Axin lack the domain responsible for binding to β-catenin (Belahbib et al. 2018; Sun 2020), and there is no evidence for a Wnt/Planar Cell Polarity (PCP) signaling pathway (Ryan et al. 2013; Moroz et al. 2014). This raises questions about what type of information will emerge from studying ancestral protein functions in ctenophores. It could either narrow down essential domains required for structural interactions or uncover alternative methods employed for accomplishing developmental and homeostatic processes. Moreover, studying key proteins in ctenophores can enhance our understanding of their evolutionary origins and shed light on how these essential biological processes evolved across Metazoa. Ctenophores, like *M. leidyi*, occupy a critical phylogenetic position, offering potentially unique insights into the molecular transition from unicellularity to multicellularity in the animal lineage. Therefore, developing tools to study β-catenin and other critical molecules in ctenophores will be instrumental in uncovering these insights. Here we report on the successful generation of affinity-purified rabbit polyclonal antibodies targeting *M. leidyi* β-catenin protein and use it to carry out localization studies to determine the subcellular distribution of the protein during early *M. leidyi* development. Using these antibodies we were able to observe β-catenin protein localize to the cell-cell interface in the earliest stages of development and in some subsets of nuclei of developing *M. leidyi* embryos. This reagent provides a useful tool for future studies on the potential function of β-catenin in cell-cell adhesion and transcriptional activation in this member of an early emerging metazoan taxon.

2. Materials and Methods

2.1 Structural analysis of *Ml*β-catenin protein domains

The protein domains of *Ml*β-catenin (*Ml*β-cat, ML073715a) have been predicted and checked with Interpro 100.0 (https://www.ebi.ac.uk/interpro/, (Paysan-Lafosse et al. 2023)) and SMART 9.0 (http://smart.embl-heidelberg.de/, (Schultz et al. 1998)). Tertiary structure was predicted using AlphaFold v2.3.1 on the COSMIC^2 platform (https://cosmic2.sdsc.edu/, (Cianfrocco et al. 2017)).

2.2 Maintenance and spawning

The *M. leidyi* adults were caught in a marina at Flagler Beach (Florida, United States, 29°30'60.0"N 81°08'47.1"W), and were maintained in 40L kreisels with constant seawater flow, temperature, and salinity at the Whitney Laboratory for Marine Bioscience of the University of Florida (Saint Augustine, Florida, USA). Care and maintenance of adults including spawning, fertilization, and embryo culturing have been carried out as previously described (Salinas-Saavedra and Martindale 2018), however the salinity the adult animals were kept in was adjusted to ⅔ strength filtered sea water (FSW, 25 parts per thousand) to improve spawning efficiency. Spawning was induced by incubating the adults for 3 hours in the dark at RT and then moving them to glass bowls approximately one hour before spawning to collect embryos. Embryos were kept in glass dishes in ⅔ FSW until the desired stage.

2.3 Molecular cloning of *Ml*β-cat

The total RNA was extracted from mixed embryonic stages using Trizol (Sigma, cat.# T9424). A partial length *Ml*β-cat coding sequence was amplified from cDNA using the Advantage RT for PCR kit (Clontech, cat #639506) using the following primers:

*Ml*β-cat pGEX forward: ATAGGATCCATGGAAACGCCAGTATAT

*Ml*β-cat pGEX reverse: TATAAGAATTCTCAGGTCGTTGTGACGGG

The amplified cDNA was then ligated into a pGEX-6P-1 vector for cloning and inducing a GST-tagged *Ml*β-cat protein. The first 200 amino acids of *Ml*β-cat were selected to be used as the antigen for the polyclonal antibodies due to its predicted high immunogenicity. BLAST search against the *M. leidyi* Protein Model 2.2 (*Mnemiopsis* Genome Project Portal, https://research.nhgri.nih.gov/mnemiopsis/) using the antigen sequence only recovered the β-catenin sequence at a significant E-value.

2.4 Antigen preparation and antibody generation

BL21(DE3) competent *E. coli* (New England Biolabs cat.# C2530H) were transformed with the *Ml*β-cat pGEX-6P-1 construct. Five milliliters of inoculated terrific broth (TB) medium were incubated overnight (12-18 hours) in a shaking incubator at 37˚C at 200 rpm. Overnight cultures were used to inoculate 500 mLs of TB medium which were then incubated until they reached an OD_{600} between 0.2-0.3, at which point the bacteria were induced by adding 0.5 mM isopropyl β-d-1-thiogalactopyranoside (IPTG). Cultures were allowed to incubate for an additional 3 hours before bacteria were collected by centrifuging at 5000g for 15 minutes at 4˚C. The induced GST-tagged *Ml*β-cat protein was affinity purified on a glutathione sepharose beads matrix as described (Kielkopf et al. 2020). Approximately 1.5 mg of Mlβ-cat protein collected via PreScission Protease (GenScript cat.# Z02799) digest and 6 mg of *Ml*β-cat-GST protein collected via 10 mM glutathione whole protein elution were sent to Labcorp Early Development Laboratories Inc. (Denver, PA). There, the *Ml*β-cat N-terminal polypeptide was used as the antigen for single rabbit polyclonal antibody generation and the *Ml*β-cat-GST protein was used for affinity purification of the antibody from final bleed serum.

2.5 Western blot analysis

M. leidyi embryo protein samples were collected by centrifuging mixed stage embryos (between 8 and 12 hours post fertilization, hpf) in an Eppendorf tube to pellet, removing as much sea water as possible, and adding minimum volume (between 30-50 μL) of 1X SDS gel-loading buffer (50 mM Tris-Cl pH 6.8, 100 mM dithiothreitol, 2% sodium dodecyl sulfate, 0.1% bromophenol blue, 10% glycerol) to fully lyse the pellet. *M. leidyi* adult protein samples were collected by dissecting the lobes off, using a homogenizer to dissociate cells, centrifuging to pellet, removing as much sea water as possible, and adding minimum volume (between 100-500 μL) of 1X SDS gel-loading buffer to fully lyse the sample. Lysates were boiled for 5 minutes and 30 μL was then separated on 10% SDS-PAGE gels, transferred onto 0.45μm nitrocellulose membranes (Bio-Rad Laboratories, Inc. cat. # 1620115) and blocked overnight in a 5% milk powder 1X tris buffered saline solution. The immunoblot was probed with the rabbit anti-*Ml*β-cat (1:500). Blots were developed using a IRDye 800 (1:10,000) donkey anti-rabbit secondary antibody (LI-COR, Inc. cat.# 926-32213) and imaged using a LI-COR Odyssey Infrared Imaging System.

2.6 Fixing and immunofluorescence analysis of embryos

The vitelline membranes of *M. leidyi* embryos were mechanically removed using forceps before fixation. Embryos were kept in plastic dishes treated with 3% Bovine Serum Albumin (BSA) in ⅔ FSW to prevent them from sticking to the dish. Embryos of different developmental stages were fixed (4% paraformaldehyde, 0.2 % glutaraldehyde in 1X phosphate buffered saline) 5 minutes at RT, before being fixed in a second fixative solution (4% PFA in 1X phosphate buffered saline) 10 minutes at RT. Embryos were washed three times with 1X PBS then permeabilized in 0.2% Triton in PBS. Embryos were rinsed several times in 1X PBT (0.1% BSA, 0.2% Triton in 1X phosphate buffered saline) before the blocking step in 5% Normal Goat Serum (NGS in 1X PBT) for 1 hour at RT on a rocker. The blocking solution was then replaced by the primary antibody (anti-*Ml*β-cat; 1:500 in blocking reagent) and incubated overnight at 4° with gentle rocking. Fixed embryos were washed several times with 1X PBT. A goat anti-rabbit IgG AlexaFluor 647 (Invitrogen,cat.# A-21245) secondary antibody was diluted in 5% NGS (1:250) and embryos were incubated in the dark at 4° overnight on a rocker. Negative controls were incubated with the secondary antibody only. Samples were washed with 1X PBT for a total period of 2 hours. Embryos were finally stained with DAPI (1:1,000 in PBT; Invitrogen, Inc. Cat. #D1306) to allow nuclear visualization. Stained samples were rinsed 2 times with 1X phosphate buffered saline before mounting in the same solution. The embryos were imaged under a Zeiss 710 scanning confocal microscope. All the images were processed through ImageJ (z-stack and maximal intensity projection). More than 30 embryos were examined at each embryonic stage.

3. Results

3.1. *β-catenin protein structure in M. leidyi*

In most metazoans, the core of the β-catenin protein is composed of 12 Armadillo repeat domains (Arm), essential for the interaction with β-catenin binding partners for both Wnt/β-catenin signaling and in AJs (Schneider et al. 2003, van der Wal and van Amerongen 2020). Those binding sites are overlapping, meaning that β-catenin can either bind to cadherin or TCF/LEF, but cannot simultaneously accomplish both functions (Orsulic et al. 1999). The *M. leidyi* genome encodes one single *β-catenin* gene of 2,688 bps coding for 895 amino acids (Ryan et al. 2013). Protein domain prediction softwares were only able to identify 9 out of 12 individual Arm repeats conserved in *Ml*β-cat, but also highlighted a larger Arm repeat superfamily that extended past the final recognized domain (Fig. 2A). Upon further investigation into the tertiary structure, 3 additional complete triple alpha helices were identified (Fig. 2B). Arm domain 7 was predicted with low confidence likely due to only 2 out of the 3 alpha helices being conserved in predicted 3D structure. However a third, short alpha helix is present immediately before the domain but truncated by a low confidence value region of disorder. Furthermore, the alignment of *Ml*β-cat with structurally characterized orthologs in other animals revealed the conservation of the 2 lysine residues (K312 and K435 in mouse epithelial cadherin) in *Ml*β-cat, crucial for the interaction between a classical cadherin and β-catenin (Huber and Weis 2001). Moreover, the phosphorylation sites required by β-catenin destruction complex components, part of the Wnt/β-catenin pathway, are also conserved (Serine 33, Serine 37, and Threonine 41 for GSK3β ; Serine 45 for CK1α). Additionally, 9 out of the 14 essential binding residues of the α-catenin binding motif are conserved (Belahbib et al. 2018).

Fig. 2. The predicted β-catenin architecture in *M. leidyi*. **(A)** *Ml*β-cat protein contains 11 of the 12 Arm domains identified with high confidence flanked by a N-Terminal and a C-terminal region. Asterisk above domain 7 indicates predicted partial (2 out of 3 alpha helices) domain conservation. The amino acids necessary for the interaction with GSK3β and CK1α are conserved, as well as the α-cat binding site. The two crucial lysines (K), essential for the interaction with a classical cadherin, are also conserved. **(B)** Predicted tertiary structure of *Ml*β-cat protein with Arm domains denoted by color. Red through pink domains were identified through domain prediction software; silver, gray, and white domains fall within the Arm repeat superfamily but were identified manually. The blue Arm domain (domain 7) was identified with low confidence likely due to the missing alpha helix. **(C)** pLDDT plot for predicted structure. Dip in confidence values for residues between 472-478 corresponds with a disordered region that truncates an alpha helix in Arm domain 7.

3.2 Affinity-purified rabbit polyclonal antibodies generated to Mlβ-catenin protein specifically targets endogenous Mlβ-catenin

To confirm that the affinity-purified rabbit polyclonal antibodies generated were specifically targeting *Ml*β-catenin, whole cell lysates from both embryo and adult *M. leidyi* were collected and subjected to Western blot analysis. Only a single band was detected around the expected size (~100 kDa) for both samples (Fig. 3A). Next, fixed whole embryos from multiple stages were immunostained using the *Ml*β-cat antibody and nuclei were labeled with DAPI. Fixed embryos incubated with the secondary antibody alone were used as a negative control. Samples imaged using scanning confocal microscopy

demonstrated robust nuclear staining by DAPI, but showed fluorescence only when embryos were incubated with the primary *Ml*β-cat antibody (Fig. 3B).

Fig. 3. *Ml*β-cat antibody specifically recognizes endogenous protein. **(A)** Western blot analysis of *M. leidyi* cell lysates using affinity-purified anti-*Ml*β-cat antibodies. The specificity of the antibody was tested against whole-cell lysates from both *M. leidyi* embryos and adults. A single band at the expected size (~100kDa) was detected for both samples. **(B)** Immunostaining of *M. leidyi* embryos using affinity-purified anti-*Ml*β-cat polyclonal antibodies. The top panels show nuclei stained by DAPI; the bottom panels show results of *Ml*β-cat staining with and without primary antibody (negative control). Images are maximal projection. Scale bar, 50 µM.

3.3 Mlβ-cat localizes at cell-cell contacts

β-catenin is known to play a structural role by its importance in cell-cell adhesion by its association with cadherins at the plasma membrane. To determine if *Ml*β-cat was putatively involved in cell-cell adhesion, we stained *M. leidyi* embryos at different cell-stages from fertilized egg to 5 hpf gastrula. We were not able to detect any membrane staining of *Ml*β-cat at the 1-cell stage, however a cortical localization of *Ml*β-cat at cell-cell contacts at the 2-cell stage was visible (not shown). This localization continued from the 2-cell stage through the late stages of the development in *M. leidyi* (Fig. 4). This localization pattern was only present at cell-cell contacts, while the region of cell membranes that lacked contact displayed a diffuse level of *Ml*β-cat staining that more closely resembled the cytoplasm.

Fig. 4. *Ml*β-catenin localizes at cell-cell contacts during *M. leidyi* embryonic development. Immunostaining of *M. leidyi* embryos at different stages using affinity-purified β-cat antibodies. The top panels exhibit nuclei stained by DAPI; the bottom panels exhibit results of *Ml*β-cat antibody staining. Images are maximal projection. Note that enriched localization of β-catenin only occurs at cell-cell contacts. Scale bar, 50 μM.

3.4 Mlβ-cat translocates into the nucleus during early embryogenesis

The β-catenin function in cell signaling is associated with its accumulation into the nucleus to act as a transcriptional cofactor with TCF/LEF. We therefore looked for nuclear staining using the *Ml*β-cat antibody. We first noticed that *Ml*β-cat was uniformly expressed in the cytoplasm at 1-cell stage, indicating that this protein is a maternal component in *M. leidyi*. However, we did not consistently detect any clear nuclear signal of β-catenin from 1-cell to 32-cell (Fig. 5). The first *Ml*β-cat nuclear translocation event that we observed was at the 60-cell stage in some, but not all, 3E and 2M\ macromeres (Fig. 5). *Ml*β-cat nuclear staining continues to be random until 4 hpf where nuclearization can be observed in cells at the oral pole, but nuclear staining was absent from the aboral pole.

Fig. 5. The localization of β-catenin in *M. leidyi* embryos. Immunostaining of *M. leidyi* embryos at different stages using affinity-purified β-cat antibodies. The left column shows nuclei stained by DAPI, the middle column shows results of *Ml*β-cat antibody staining, and the right column zooms in on an inset of the 4 hpf images (dotted boxes). The 4-cell, 60-cell, and 4 hpf aboral view are z-stack projected images. The 4 hpf oral view is a maximal projection image. The yellow arrows show absence of β-catenin nuclear translocation; while red arrows indicate β-catenin nuclear translocation. Scale bar, 50 μM.

4. Discussion

4.1 β-catenin robustly localizes to cell-cell interfaces in M. leidyi embryos

β-catenin involvement in the CCC is crucial for the adhesion between cells in every bilaterian animal where this process has been experimentally tested (Halbleib and Nelson 2006). Nevertheless, only a few studies focused on the roles of this complex in early branching lineages. Previous studies on the cnidarian *Nematostella vectensis* have shown that the CCC is involved in cell-cell adhesion and germ layer formation (Clarke et al. 2019; Pukhlyakova et al. 2019). Additionally, a functional CCC has been shown to be involved in cell-cell adhesion in sponges as supported by the co-immunoprecipitation of interacting CCC components, β-catenin localizing to cell-cell boundaries in *Ephydatia muelleri* (Schippers and Nichols 2018) and furthermore by *Yeast* 2-Hybrid screening in another sponge, *Oscarella carmela* (Nichols et al. 2012).

As for ctenophores, a prior study using polyclonal *Ml*β-cat antibodies generated against the first 10 amino acids of the protein was unable to capture evidence of the protein localized to the cell-cell interface in *M. leidyi* (Salinas-Saavedra et al. 2019). Thus it was concluded that the CCC is not conserved and hence does not play a role in cell-cell adhesion in ctenophores. However, there is a possibility that the short epitope used to generate the antibodies in the Salines-Saavedra et al. (2019) study were being blocked by β-catenin binding partners at the cell surface, impeding the antibodies' ability to bind and hindering the ability to observe this role. Unfortunately, the peptide antibody generated in the 2019 study appears to have lost its activity and we were not able to confirm its expression in embryos or Western blots (data not shown). Our results show that *Ml*β-catenin clearly localizes at cell-cell contacts during *M. leidyi* embryonic development and therefore likely has a cell-to-cell adhesion function in the earliest metazoan lineage (Fig. 5). Additionally, transcriptomic and genomic data showed that *M. leidyi* has a full set of CCC components (Ryan et al. 2013) and the amino acids essential for the interaction between β-catenin and α-catenin and between β-catenin and cadherin are mostly conserved (Fig. 2) (Belahbib et al. 2018). The conservation of almost all 12 Arm domains on *Ml*β-catenin leads us to predict that future functional studies will demonstrate that *Ml*β-catenin maintains its ability to interact with CCC partners in ctenophores.

However, it is unclear if a functional CCC exists in *M. leidyi* since the β-catenin binding site, essential for recruiting β-catenin at the membrane, is either cryptic or absent from *Mnemiopsis* cadherin sequences (Belahbib et al. 2018; Salinas-Saavedra et al. 2019). An alternative hypothesis is that the CCC emerged from an interaction between β-catenin and a protocadherin instead of a classical cadherin. Indeed, it has been shown that a protocadherin can interact with β-catenin (Chen et al. 2002; Parenti et al. 2010; de Nys et al. 2024), but no study has shown that this interaction is necessary for cell-cell adhesion. A third option is cadherins are entirely unnecessary for cell-cell adhesion in ctenophores. An example of this was shown in the social amoeba *Dictyostelium discoideum* where in response to starving this organism undergoes the

formation of a fruiting body, corresponding to a "multicellular" stage (Urushihara 2008). Analysis of the *D. discoideum* genome indicates that α-catenin and β-catenin genes are present while cadherins are absent. Knocking down the *β-cat* ortholog, called *Aardvark*, demonstrated that the catenins were essential for the establishment of this epithelium-like structure (Dickinson et al. 2011). Clarifying the interacting partners of *Ml*β-cat at the cell membrane will provide insights into the evolutionary origin of the CCC across metazoa.

4.2 β-catenin is nuclearized in cells around the blastopore in M. leidyi embryos
β-catenin can be found in three locations inside cells: in the cytoplasm, at the cell surface on the cytoplasmic side of the cell membrane, and in the nucleus. In each of these locations, it interacts with different proteins and maintains different roles. In order to fulfill its role as a cell signaling molecule, β-catenin must avoid being marked for degradation by the destruction complex in the cytoplasm and then translocate into the nucleus. In the vast majority of organisms where this process has been studied, this begins to occur in the early stages of development and facilitates transcription of genes involved in specifying cell fate (Martindale 2005; Petersen and Reddien 2009). A prior study in *Mnemiopsis leidyi* tracked the spatiotemporal expression of the *Ml*β-catenin protein during development and found nuclearization beginning at the zygote stage (Salinas-Saavedra et al. 2019). Furthermore, this study reported that there was continued nuclear expression of *Ml*β-cat in macromeres at the animal half in the endomesodermal progenitors through 11 hpf, arguing that this is substantial evidence to conclude that *Ml*β-cat has a role in specifying endomesoderm in early *M. leidyi* embryogenesis. When trying to replicate these results using the affinity-purified antibodies we generated to a *Ml*β-catenin polypeptide we were unable to find clear and consistent evidence of nuclear *Ml*β-cat in embryos prior to the 60-cell stage. From the 60-cell to 3 hpf we find nuclear *Ml*β-cat to be patchy and without a clear pattern. This is unexpected due to the highly stereotyped cleavage program and precocious specification of cell fate seen in ctenophores (Martindale & Henry 1999). Early stage ctenophore embryos develop at an accelerated pace making it difficult to capture specific stages for fixation and staining. Improving methods for collecting and staining may make obtaining spatial-temporal protein expression data more consistent in these early stages.

Scanning confocal analysis of *Ml*β-cat antibody stained *Mnemiopsis* embryos at around 4 hpf started to show a clear pattern of nuclear staining restricted blastomeres at the oral pole, while cells at the aboral pole did not show any nuclear staining. This is evidence that *Ml*β-cat may maintain its role as a co-activator of transcription, but does not align with the previously described expression pattern. The fact that this pattern is restricted to one pole does indicate the possibility for differential gene expression or stabilization, although patchy nuclear translocation during early development and a restriction to the ectodermal cells at the oral pole leaves us with questions about this protein's functional role in cell type specification. Future functional experiments are required to determine if β-catenin maintains its role as a transcriptional co-activator in the Wnt/β-catenin pathway in ctenophores.

Furthermore, transportation of β-catenin into the nucleus is not well understood in all metazoans. There are several proposed mechanisms for transport including direct nuclear pore complex interaction, various piggyback or chaperone candidates, and post-translational modification (Anthony et al. 2020). With their

ancestral molecular repertoire and favorable optical properties, ctenophore embryos might be a powerful model to investigate the mechanisms behind β-catenin nuclearization.

4.3 Developing tools to study ctenophores

The combination of Western blot analysis and immunohistochemistry results gives us high confidence that the newly generated affinity-purified *Ml*β-cat antibody specifically targets β-catenin without any detectable non-specific interactions. As such, this antibody can now be used as a tool to ask follow-up questions about the functional roles of β-catenin in the earliest metazoan branch. The evolution and developmental community has come to recognize that ctenophores sit at a crucial point for understanding the early evolutionary history of metazoans and has been working to develop experimental methods and resources like assembled reference genomes, transcriptomes, morpholino oligonucleotides, and CRISPR-Cas9 (Moreland et al. 2014; Davidson et al. 2017; Yamada et al. 2010; Presnell and Browne 2021; Presnell et al. 2022). Hopefully with this expanding toolkit we can begin to answer questions about transition from unicellularity to multicellularity in the animal lineage as well as better understand the molecular components the earliest metazoans maintain that are necessary for cell-cell adhesion and cell fate specification. As we develop new tools and techniques to study ctenophores, we will not only better understand how these animals operate on a molecular level, but also gain insights into how conserved mechanisms have evolved over time.

References

Aberle, H., Bauer, A., Stappert, J., Kispert, A., & Kemler, R. (1997). B-Catenin Is a Target for the Ubiquitin-Proteasome Pathway. *EMBO Journal*, *16*(13), 3797–3804. https://doi.org/10.1093/emboj/16.13.3797

Anthony, C. C., Robbins, D. J., Ahmed, Y., & Lee, E. (2020). Nuclear regulation of Wnt/β-catenin signaling: It's a complex situation. *Genes*, *11*(8), 1–11. https://doi.org/10.3390/genes11080886

Belahbib, H., Renard, E., Santini, S., Jourda, C., Claverie, J. M., Borchiellini, C., & Le Bivic, A. (2018). New genomic data and analyses challenge the traditional vision of animal epithelium evolution. *BMC Genomics*, *19*(1), 1–15. https://doi.org/10.1186/s12864-018-4715-9

Brunet, T., Bouclet, A., Ahmadi, P., Mitrossilis, D., Driquez, B., Brunet, A. C., … Farge, E. (2013). Evolutionary conservation of early mesoderm specification by mechanotransduction in Bilateria. *Nature Communications*, *4*(2821), 1–15. https://doi.org/10.1038/ncomms3821

Chen, M. W., Vacherot, F., De la Taille, A., Gil-Diez-de-Medina, S., Shen, R., Friedman, R. A., … Buttyan, R. (2002). The emergence of protocadherin-PC expression during the acquisition of apoptosis-resistance by prostate cancer cells. *Oncogene*, *21*(51), 7861–7871. https://doi.org/10.1038/sj.onc.1205991

Cianfrocco, M. A., Wong-Barnum, M., Youn, C., Wagner, R., & Leschziner, A. (2017). COSMIC2: A science gateway for cryo-electron microscopy structure determination. *ACM International Conference Proceeding Series*, *Part F1287*, 13–17. https://doi.org/10.1145/3093338.3093390

Clarke, D. N., Lowe, C. J., & Nelson, W. J. (2019). The Cadherin-Catenin Complex is Necessary for Cell Adhesion and Embryogenesis in Nematostella vectensis. *Developmental Biology*, *447*(2), 170–181. https://doi.org/10.1016/j.ydbio.2019.01.007.

Clevers, H. (2006). Wnt/β-Catenin Signaling in Development and Disease. *Cell*, *127*(3), 469–480. https://doi.org/10.1016/j.cell.2006.10.018

Davidson, P. L., Koch, B. J., Schnitzler, C. E., Henry, J. Q., Martindale, M. Q., Baxevanis, A. D., & Browne, W. E. (2017). The maternal-zygotic transition and zygotic activation of the Mnemiopsis leidyi genome occurs within the first three cleavage cycles. *Molecular Reproduction and Development*, *84*(11), 1218–1229. https://doi.org/10.1002/mrd.22926

Davis, M. A., Ireton, R. C., & Reynolds, A. B. (2003). A core function for p120-catenin in cadherin turnover. *Journal of Cell Biology*, *163*(3), 525–534. https://doi.org/10.1083/jcb.200307111

de Nys, R., Gardner, A., van Eyk, C., Mincheva-Tasheva, S., Thomas, P., Bhattacharjee, R., … Gecz, J. (2024). Proteomic analysis of the developing mammalian brain links PCDH19 to the Wnt/β-catenin signalling pathway. *Molecular Psychiatry*, (July 2023), 1–12. https://doi.org/10.1038/s41380-024-02482-z

Desprat, N., Supatto, W., Pouille, P. A., Beaurepaire, E., & Farge, E. (2008). Tissue Deformation Modulates Twist Expression to Determine Anterior Midgut Differentiation in Drosophila Embryos. *Developmental Cell*, *15*(3), 470–477. https://doi.org/10.1016/j.devcel.2008.07.009

Dickinson, D. J., Nelson, W. J., & Weis, W. I. (2011). A Polarized Epithelium Organized by β- and α-Catenin Predates Cadherin and Metazoan Origins. *Science*, *331*(6022), 1336–1339. https://doi.org/10.1126/science.1199633.

Dunn, C. W., Hejnol, A., Matus, D. Q., Pang, K., Browne, W. E., Smith, S. A., … Giribet, G. (2008). Broad phylogenomic sampling improves resolution of the animal tree of life. *Nature*, *452*(7188), 745–749. https://doi.org/10.1038/nature06614

Farge, E. (2003). Mechanical Induction of Twist in the Drosophila Foregut/Stomodeal Primordium. *Current Biology*, *13*, 1365–1377. https://doi.org/10.1016/S0960-9822(03)00576-1

Gayrard, C., Bernaudin, C., Déjardin, T., Seiler, C., & Borghi, N. (2018). Src- and confinement-dependent FAK activation causes E-cadherin relaxation and β-catenin activity. *Journal of Cell Biology*, *217*(3), 1063–1077. https://doi.org/10.1083/jcb.201706013

Halbleib, J. M., & Nelson, W. J. (2006). Cadherins in development: Cell adhesion, sorting, and tissue morphogenesis. *Genes and Development*, *20*(23), 3199–3214. https://doi.org/10.1101/gad.1486806

Hejnol, A., Obst, M., Stamatakis, A., Ott, M., Rouse, G. W., Edgecombe, G. D., … Dunn, C. W. (2009). Assessing the root of bilaterian animals with scalable phylogenomic methods. *Proceedings of the Royal Society B: Biological Sciences*, *276*(1677), 4261–4270. https://doi.org/10.1098/rspb.2009.0896

Hinck, L., Näthke, I. S., Papkoff, J., & Nelson, W. J. (1994). Dynamics of cadherin/catenin complex formation: Novel protein interactions and pathways of complex assembly. *Journal of Cell Biology*, *125*(6), 1327–1340. https://doi.org/10.1083/jcb.125.6.1327

Huber, A. H., & Weis, W. I. (2001). The structure of the β-catenin/E-cadherin complex and the molecular basis of diverse ligand recognition by β-catenin. *Cell*, *105*(3), 391–402. https://doi.org/10.1016/S0092-8674(01)00330-0

Ishiyama, N., & Ikura, M. (2012). The Three-Dimensional Structure of the Cadherin–Catenin Complex. In *Adherens Junctions: From Molecular Mechanisms to Tissue Development and Disease* (pp. 39–62). Springer, Dordrecht. https://doi.org/https://doi.org/10.1007/978-94-007-4186-7_3

Kielkopf, C. L., Bauer, W., & Urbatsch, I. L. (2020). Purification of Fusion Proteins by Affinity Chromatography on Glutathione Resin. *Cold Spring Harbor Protocols*, *2020*(6), 217–224.

Klaus, A., & Birchmeier, W. (2008). Wnt signalling and its impact on development and cancer. *Cancer*, *8*.

Kourtidis, A., Ngok, S. P., & Anastasiadis, P. Z. (2013). P120 catenin: An essential regulator of cadherin stability, adhesion-induced signaling, and cancer progression. *Progress in Molecular Biology and Translational Science*, *116*, 409–432. https://doi.org/10.1016/B978-0-12-394311-8.00018-2

Lecuit, T., & Yap, A. S. (2015). E-cadherin junctions as active mechanical integrators in tissue dynamics. *Nature Cell Biology*, *17*(5), 533–539. https://doi.org/10.1038/ncb3136

Martindale, M. Q., & Henry, J. Q. (1999). Intracellular fate mapping in a basal metazoan, the ctenophore Mnemiopsis leidyi, reveals the origins of mesoderm and the existence of indeterminate cell lineages. *Developmental Biology*, *214*(2), 243–257. https://doi.org/10.1006/dbio.1999.9427

Martindale, M. Q. (2005). The evolution of metazoan axial properties. *Nature Reviews Genetics*, *6*(12), 917–927. https://doi.org/10.1038/nrg1725

Martindale, M. Q., & Hejnol, A. (2009). A Developmental Perspective: Changes in the Position of the Blastopore during Bilaterian Evolution. *Developmental Cell*, *17*(2), 162–174. https://doi.org/10.1016/j.devcel.2009.07.024

Meng, W., & Takeichi, M. (2009). Adherens Junction: Molecular Architecture and Regulation. *Cold Spring Harbor Perspectives in Biology*, *1*, 1–13. https://doi.org/10.1098/rstb.1989.0070

Moreland, R. T., Nguyen, A. D., Ryan, J. F., Schnitzler, C. E., Koch, B. J., Siewert, K., … Baxevanis, A. D. (2014). A customized Web portal for the genome of the ctenophore Mnemiopsis leidyi. *BMC Genomics*, *15*(1), 1–13. https://doi.org/10.1186/1471-2164-15-316

Moroz, L. L., Kocot, K. M., Citarella, M. R., Dosung, S., Norekian, T. P., Povolotskaya, I. S., … Kohn, A. B. (2014). The Ctenophore Genome and the Evolutionary Origins of Neural Systems. *Nature*, *510*(7503), 109–114. https://doi.org/10.1038/nature13400.

Nichols, S. A., Roberts, B. W., Richter, D. J., Fairclough, S. R., & King, N. (2012). Origin of metazoan cadherin diversity and the antiquity of the classical cadherin/β-catenin complex. *Proceedings of the National Academy of Sciences of the United States of America*, *109*(32), 13046–13051. https://doi.org/10.1073/pnas.1120685109

Nollet, F., Kools, P., & Van Roy, F. (2000). Phylogenetic analysis of the cadherin superfamily allows identification of six major subfamilies besides several solitary members. *Journal of Molecular Biology*, *299*(3), 551–572. https://doi.org/10.1006/jmbi.2000.3777

Nusse, R., & Clevers, H. (2017). Wnt/β-Catenin Signaling, Disease, and Emerging Therapeutic Modalities. *Cell*, *169*(6), 985–999. https://doi.org/10.1016/j.cell.2017.05.016

Orsulic, S., Huber, O., Aberle, H., Arnold, S., & Kemler, R. (1999). E-cadherin binding prevents β-catenin nuclear localization and β-catenin/LEF-1-mediated transactivation. *Journal of Cell Science*, *112*(8), 1237–1245. https://doi.org/10.1242/jcs.112.8.1237

Parenti, S., Ferrarini, F., Zini, R., Montanari, M., Losi, L., Canovi, B., … Grande, A. (2010). Mesalazine inhibits the β-catenin signalling pathway acting through the upregulation of μ-protocadherin gene in colo-rectal cancer cells. *Alimentary Pharmacology and Therapeutics*, *31*(1), 108–119. https://doi.org/10.1111/j.1365-2036.2009.04149.x

Paysan-Lafosse, T., Blum, M., Chuguransky, S., Grego, T., Pinto, B. L., Salazar, G. A., … Bateman, A. (2023). InterPro in 2022. *Nucleic Acids Research*, *51*(D1), D418–D427. https://doi.org/10.1093/nar/gkac993

Peifer, M., McCrea, P. D., Green, K. J., Wieschaus, E., & Gumbiner, B. M. (1992). The Vertebrate Adhesive Junction Proteins β-catenin and Plakoglobin and the Drosophila Segment Polarity Gene armadillo Form a Multigene Family with Similar Properties Mark. *The Journal of Cell Biology*, *181*(3), 681–691.

Petersen, C. P., & Reddien, P. W. (2009). Wnt Signaling and the Polarity of the Primary Body Axis. *Cell*, *139*(6), 1056–1068. https://doi.org/10.1016/j.cell.2009.11.035

Presnell, J. S., Bubel, M., Knowles, T., Patry, W., & Browne, W. E. (2022). Multigenerational laboratory culture of pelagic ctenophores and CRISPR–Cas9 genome editing in the lobate Mnemiopsis leidyi. *Nature Protocols*, *17*(8), 1868–1900. https://doi.org/10.1038/s41596-022-00702-w

Presnell, Jason S., & Browne, W. E. (2021). Krüppel-like factor gene function in the ctenophore Mnemiopsis leidyi assessed by CRISPR/Cas9-mediated genome editing. *Development (Cambridge)*, *148*(17). https://doi.org/10.1242/DEV.199771

Pukhlyakova, E. A., Kirillova, A. O., Kraus, Y. A., Zimmermann, B., & Technau, U. (2019). A cadherin switch marks germ layer formation in the diploblastic sea anemone Nematostella vectensis. *Development (Cambridge)*, *146*(20), 1–15. https://doi.org/10.1242/dev.174623

Richter, D. J., & King, N. (2013). The genomic and cellular foundations of animal origins. *Annual Review of Genetics*, *47*, 509–537. https://doi.org/10.1146/annurev-genet-111212-133456

Röper, J.-C., Mitrossilis, D., Stirnemann, G., Waharte, F., Brito, I., Fernandez-Sanchez, M.-E., … Farge, E. (2018). The major β-catenin/E-cadherin junctional binding site is a primary molecular mechano-transductor of differentiation in vivo. *ELife*, *32*(3), 152–160. https://doi.org/10.1016/S1553-7250(06)32020-X

Ros-Rocher, N., Pérez-Posada, A., Leger, M. M., & Ruiz-Trillo, I. (2021). The origin of animals: An ancestral reconstruction of the unicellular-to-multicellular transition. *Open Biol.*, *11(2)*, doi: 10.1098/rsob.200359.

Ryan, J. F., Pang, K., Schnitzler, C. E., Nguyen, A., Moreland, R. T., Simmons, D. K., … Baxevanis, A. D. (2013). The genome of the ctenophore Mnemiopsis leidyi and its implications for cell type evolution. *Science*, *342*(6164). https://doi.org/10.1126/science.1242592.The

Salinas-Saavedra, M., & Martindale, M. Q. (2018). Improved protocol for spawning and immunostaining embryos and juvenile stages of the ctenophore Mnemiopsis leidyi. *Protocol Exchange*, 1–10. https://doi.org/10.1038/protex.2018.092

Salinas-Saavedra, M., Wikramanayake, A., & Martindale, M. (2019). B-Catenin Has an Ancestral Role in Cell Fate Specification But Not Cell Adhesion. *BioRxiv*, 520957. Retrieved from https://doi.org/10.1101/520957

Schippers, K. J., & Nichols, S. A. (2018). Evidence of signaling and adhesion roles for β-Catenin in the sponge ephydatia muelleri. *Molecular Biology and Evolution*, *35*(6), 1407–1421. https://doi.org/10.1093/molbev/msy033

Schneider, S. Q., Finnerty, J. R., & Martindale, M. Q. (2003). Protein Evolution: Structure-Function Relationships of the Oncogene Beta-Catenin in the Evolution of Multicellular Animals. *Journal of Experimental Zoology Part B: Molecular and Developmental Evolution*, *295*(1), 25–44. https://doi.org/10.1002/jez.b.6

Schultz, D. T., Haddock, S. H. D., Bredeson, J. V., Green, R. E., Simakov, O., & Rokhsar, D. S. (2023). Ancient gene linkages support ctenophores as sister to other animals. *Nature*, *618*(7963), 110–117. https://doi.org/10.1038/s41586-023-05936-6

Schultz, J., Milpetz, F., Bork, P., & Ponting, C. P. (1998). SMART, a simple modular architecture research tool: Identification of signaling domains. *Proceedings of the National Academy of Sciences of the United States of America*, *95*(11), 5857–5864. https://doi.org/10.1073/pnas.95.11.5857

Shapiro, L. (1997). The multi-talented β-catenin makes its first appearance. *Structure*, *5*(10), 1265–1268. https://doi.org/10.1016/S0969-2126(97)00278-5

Stamos, J.L. & Weis, W.I. (2013). The β-catenin destruction complex. *Cold Spring Harb Perspect Biol*, *5*(1), doi: 10.1101/cshperspect.a007898

Sun, H. (2020). *Evolution of the Wnt Signaling Pathway : Canonical and Non-Canonical Roles for Wnt Pathway Components in Regulating the Primary Axis in Bilaterian and Non-bilaterian Embryos*. University of Miami.

Urushihara, H. (2008). Developmental biology of the social amoeba: History, current knowledge and prospects. *Development Growth and Differentiation*, *50*(277–281). https://doi.org/10.1111/j.1440-169X.2008.01013.x

Valenta, T., Hausmann, G., & Basler, K. (2012). The many faces and functions of Î 2-catenin. *EMBO Journal*, *31*(12), 2714–2736. https://doi.org/10.1038/emboj.2012.150

van der Wal, T., & van Amerongen, R. (2020). Walking the tight wire between cell adhesion and WNT signalling: A balancing act for β-catenin. *Open Biology*, *10*(12). https://doi.org/10.1098/rsob.200267

Whelan, N. V., Kocot, K. M., Moroz, T. P., Mukherjee, K., Williams, P., Paulay, G., … Halanych, K. M. (2017). Ctenophore relationships and their placement as the sister group to all other animals. *Nature Ecology and Evolution*, *1*(11), 1737–1746. https://doi.org/10.1038/s41559-017-0331-3.

Wikramanayake, A. H., Hong, M., Lee, P. N., Pang, K., Byrum, C. A., Bince, J. M., … Martindale, M. Q. (2003). An ancient role for nuclear β-catenin in the evolution of axial polarity and germ layer segregation. *Nature*, *426*(6965), 446–450. https://doi.org/10.1038/nature02113

Wu, W. J., & Hirsch, D. S. (2009). Mechanism of E-cadherin lysosomal degradation. *Nature Reviews Cancer*, *9*(2), 143. https://doi.org/10.1038/nrc2521-c1

Yamada, A., Martindale, M. Q., Fukui, A., & Tochinai, S. (2010). Highly conserved functions of the Brachyury gene on morphogenetic movements: Insight from the early-diverging phylum Ctenophora. *Developmental Biology*, *339*(1), 212–222. https://doi.org/10.1016/j.ydbio.2009.12.019