Functional characterisation of neuropeptides that act as ligands for both

calcitonin-type and pigment-dispersing factor-type receptors in a deuterostome

Xiao Cong [1, 2, #], Huachen Liu [1, #], Lihua Liu [1], Michaela Egertová [3],

Maurice R. Elphick [3, *] and Muyan Chen [1, *]

1 The Key Laboratory of Mariculture, Ministry of Education, Ocean University of

China, Qingdao, China

2 Yantai Institute of Coastal Zone Research, Chinese Academy of Sciences, Yantai,

China

3 School of Biological & Behavioural Sciences, Queen Mary University of London,

London, United Kingdom

* Correspondence: m.r.elphick@qmul.ac.uk; chenmuyan@ouc.edu.cn;

[#] These authors contributed equally to this work.

Abstract

The calcitonin (CT) family of related peptides exerts diverse physiological effects in mammals via two G-protein coupled receptors: CTR and the CTR-like receptor CLR. Phylogenetic analysis of CT-type signaling has revealed the presence of CT-type peptides and CTR/CLR-type proteins in both deuterostome and protostome invertebrates. Furthermore, experimental studies have demonstrated that in the protostome *Drosophila melanogaster* the CT-like peptide DH_{31} can act as a ligand for a CTR/CLR-type receptor and a pigment-dispersing factor (PDF) receptor. Here we investigated the signaling mechanisms and functions of CT-type neuropeptides in a deuterostome invertebrate, the sea cucumber *Apostichopus japonicus* (phylum Echinodermata). In *A. japonicus,* a single gene encodes two CT-type peptides (AjCT1 and AjCT2) and both peptides act as ligands for a CTR/CLR-type receptor (AjCTR) and two PDF-type receptors (AjPDFR1, AjPDFR2), but with differential activation of downstream cAMP/PKA, $G\alpha q/Ca^{2+}/PKC$ and ERK1/2 signaling pathways. Transcriptic analysis of the AjCT1/AjCT2-encoding gene revealed expression in the central nervous system and a variety of organ systems, which was visualized immunohistochemically using an antiserum to a starfish CT-type peptide (ArCT). *In vitro* pharmacological experiments demonstrated that AjCT1 and/or AjCT2 cause dose-dependent relaxation of longitudinal muscle and intestine preparations. Furthermore, *in vivo* pharmacological experiments, combined with gain- and loss-of-function experiments, revealed a potential physiological role for AjCT2/AjPDFR2 signaling in promoting feeding and growth in *A. japonicus*. To our knowledge, this is the first study to obtain evidence that CT-type peptides can act as ligands for both CTR/CLR-type and PDF-type receptors in a deuterostome. Moreover, it provides the first evidence for appetite-stimulating and growth-promoting effects of CT-type neuropeptides in bilaterians. Given the economic importance of *A. japonicus* as a foodstuff, the discovery of CT-type peptides as potential regulators of feeding and growth in this species may offer novel strategies for aquaculture applications.

Graphical abstract

A schematic showing proposed molecular mechanisms by which CT-type signaling regulates
feeding and growth in *A. japonicus*. AjCT1 and AjCT2 are represented by purple and yellow
circles, respectively. For cell signaling, black arrows indicate that this process is activated, whilst
the pink arrow indicates that ERK1/2 is activated in the AjCT1/AjPDFR1, AjCT2/AjPDFR1 and
AjCT2/AjPDFR2 signaling pathways.

Keywords: calcitonin; neuropeptide; echinoderm; sea cucumber; calcitonin receptor;
pigment-dispersing factor receptor; feeding; growth

1. Introduction

The peptide calcitonin (CT) was first identified as a hormone that regulates blood calcium levels and subsequently a family of CT-related neuropeptides was discovered in mammals, including calcitonin gene related peptides (CGRPs), amylin (AMY), adrenomedullin (AM), and CT receptor-stimulating peptide (CRSP) (Copp et al., 1962; Sekiguchi et al., 2009; Sekiguchi et al., 2016). Members of the CT-type family of peptides typically have a disulfide bond in their N-terminal region (Cai et al., 2018) and exert their physiological effects in mammals by binding to two related G protein-coupled receptors: CTR (CT receptor) and CLR (CTR-like receptor) (Martins et al., 2014). However, receptor activity-modifying proteins (RAMPs) acting as accessory proteins are key determinants of ligand selectivity for CTR and CLR (Mclatchie et al., 1998). Thus, formation of a complex with a RAMP is essential for CLR function, whereas RAMPs are not required for CTR ligand recognition (Kuwasako et al., 2011; Bomberger et al., 2012). Furthermore, binding of CT to CTR in the absence of RAMPs (Katafuchi et al., 2009) can activate signaling via several downstream pathways, including cAMP accumulation, Ca^{2+} mobilization, and ERK activation (Chen et al., 1998; Pondel, 2000; Walker et al., 2010; Andreassen et al., 2014).

Transcriptome/genome sequencing has enabled analysis of the phylogenetic distribution of CT-type signaling systems, indicating that the evolutionary origin of CT-type peptides and CTR/CLR-related receptors can be traced back to the urbilaterian common ancestor of deuterostomes and protostomes (Furuya et al., 2000; Johnson et al., 2005; Jékely, 2013; Mirabeau and Joly, 2013). Furthermore, in protostomes two different types of CT-related peptides have been identified. Firstly, peptides that are structurally similar to vertebrate CT-type peptides in having an N-terminal disulphide bridge and secondly, diuretic hormone 31 (DH_{31})-type peptides that lack an N-terminal disulphide bridge (Furuya et al., 2000; Conzelmann et al., 2013). Furthermore, phylogenetic analysis indicates that gene duplication in a common ancestor of the protostomes gave rise to genes encoding CT-like peptides with a disulphide bridge and DH_{31}-type peptides, which were then differentially retained in protostome taxa (Conzelmann et al.,

2013; Cai et al., 2018). Receptors for CT-like peptides with a disulphide bridge

have been characterised in a molluscan species, the oyster *Crassostrea gigas*,

which has seven CTR/CLR-type receptors. Two of these receptors, named Cg-

CT-R and Cragi-CTR2, were shown to be specifically activated by the CT-type

neuropeptides Cragi-CT1b and Cragi-CT2, respectively, thereby mediating

intracellular Ca^{2+} mobilization and cAMP accumulation (Schwartz *et al.*, 2019).

CT-like peptides with a disulphide bridge are not present in the insect *Drosophila*

melanogaster. However, receptors that are activated by DH_{31} have been identified

in this species and, interestingly, DH_{31} can act as a ligand for two types of

receptors. Thus, DH_{31} acts as a ligand for a CTR/CLR-related protein (CG17415),

but it also acts as a ligand for a protein (CG13758) that is the receptor for

pigment-dispersing factor (PDF) (Johnson et al., 2005; Mertens et al., 2005;

Shafer et al., 2008). Furthermore, the functional significance of the existence of

dual receptor signaling pathways for DH_{31} in *Drosophila* has been investigated

extensively (Johnson et al., 2005; Goda et al., 2019).

Currently, relatively little is known about CT-type signaling systems in

deuterostome invertebrates. Analysis of transcriptome/genome sequence data has

enabled identification of genes/transcripts encoding CT-type peptides and candidate

CTR/CLR-type receptors for these peptides in several deuterostome invertebrate taxa,

including   urochordates   (e.g.   *Ciona   intestinalis*),   cephalochordates   (e.g.

Branchiostoma floridae), hemichordates (e.g. *Saccoglossus kowalevskii*) and

echinoderms (e.g. *Strongylocentrotus purpuratus*) (Burke et al., 2006; Sekiguchi et al.,

2009; Sekiguchi et al., 2016; Rowe and Elphick, 2012; Martins et al., 2014).

Furthermore, analysis of the expression and action of CT-type peptides identified in

the starfish *Asterias rubens* and *Patiria pectinifera* has revealed that they act as

muscle relaxants (Cai et al., 2018; Semmens et al., 2016). However, experimental

confirmation of ligand-receptor partners has thus far only been reported for the

cephalochordate *B. floridae*, where three CT-type peptides were shown to act as

ligands for a CTR/CLR-type receptor when it is co-expressed with RAMP-type

proteins (Sekiguchi et al., 2016). To gain further insights into the mechanisms of CT-

type signaling in invertebrate deuterostomes, the aim of this study was to functionally characterise CT-type peptides and their receptors in an echinoderm – the sea cucumber *Apostichopus japonicus*.

A. japonicus is an economically important species in Asian aquaculture (Pangestuti and Arifin, 2018; Guo et al., 2020) and a species representative of the phylum Echinodermata. As a phylum of deuterostome invertebrates, echinoderms form a sister clade to the chordates together with hemichordates and xenoturbellids (Bromham and Degnan, 1999; Bourlat et al., 2006; Dunn et al., 2008) and as such they occupy a key "intermediate" phylogenetic position relative to the protostome invertebrates and vertebrates (Zandawala et al., 2017), which make them attractive as experimental systems to address evolutionary questions. In our previous studies, we have established this species as a model system to obtain insights into the evolutionary history and comparative physiology of neuropeptide signaling systems in bilaterians (Chen et al., 2019; Li et al., 2022; Zheng et al., 2022; Cong et al., 2023; Zheng et al., 2024). A gene encoding two CT-type peptides (AjCT1 and AjCT2) has been identified in *A. japonicus* (Chen et al., 2019) and a candidate receptor (AjCTR) for these peptides has also been predicted (Huang et al., 2021). Furthermore, relevant to the discovery that the CT-like peptide DH_{31} can act as a ligand for both a CTR/CLR-type receptor and a PDF-type receptor in *Drosophila*, the sequences of two PDF-type receptors have been identified in *A. japonicus* (AjPDFR1 and AjPDFR2) (Huang et al., 2021).

The first aim of this study was to investigate if AjCT1 and/or AjCT2 act as ligands for AjCTR, AjPDFR1 and AjPDFR2 and to characterise the downstream signaling pathways that are activated when AjCT1 and/or AjCT2 bind to their receptor(s). The second aim was to investigate the physiological roles of AjCT1 and AjCT2 in *A. japonicus* by analysing the expression of the gene encoding these peptides, investigating CT-type neuropeptide expression using immunohistochemistry, examining the *in vitro* and *in vivo* pharmacological effects of these peptides and performing loss-of-function tests. The functional characterisation of CT-type signaling in *A. japonicus* reported in this study provides new insights into the evolution and

comparative physiology of neuropeptides and a basis for application of neuropeptide-

based strategies to improve aquaculture methods for this economically important

species.

2. Results

2.1 Characterization and phylogenetic analysis of AjCT1/2

The complete cDNA sequences of the CT-type neuropeptide precursors *AjCTP1*

and *AjCTP2* were confirmed. The coding sequence of *AjCTP2* is shorter than *AjCTP1*

(Figure 1A) but both transcripts encode precursor proteins that comprise the same

predicted 23-residue N-terminal signal peptide (Figure 1-figure supplement 1A).

AjCTP1 encodes both putative CT-type neuropeptides (AjCT1 and AjCT2), while

AjCTP2 only encodes AjCT2 (Figure 1A, Figure 1-figure supplement 1A and 1B).

Exon and intron structure analysis showed that *AjCTP1* comprises four exons, with

AjCT1 and AjCT2 encoded in the third and fourth exons, respectively. *AjCTP2*

comprises three exons, excluding the third exon that encodes AjCT1 (Figure 1A).

Phylogenetic analysis revealed that AjCTP1 and AjCTP2 are closely related to CT-

type precursors from other sea cucumbers (*Holothuria glaberrima* and *Holothuria*

scabra), and are positioned within a clade that also contains CT-type precursors from

other echinoderms (sea urchin *S. purpuratus*, starfish *A. rubens*). Furthermore, the

echinoderm CT-type precursors are positioned within a clade that also comprises CT-

type precursors from chordates, whilst DH_{31}/CT-type precursors from protostomes are

positioned in a separate clade (Figure 1B). Sequence alignment of CT-related

neuropeptides revealed that, with the exception of DH_{31}-type peptides, the N-terminal

region of CT-related peptides has a pair of conserved cysteine residues that are known

or predicted to form a disulfide bond. Another conserved feature is a C-terminal

proline residue, with the exception of CGRPs (Figure 1-figure supplement 2).

Accordingly, both AjCT1 and AjCT2 have these conserved characteristics.

2.2 Characterization and phylogenetic analysis of three candidate receptors for

AjCT1/2

The sequences of three candidate receptors for AjCT1 and AjCT2 were obtained

by RACE and phosphorylation sites were predicted, as shown in Figure 1-figure

supplement 3A, 3B, and 3C. Phylogenetic analysis revealed that CTR/CLR-type

receptors divided into two clades and an *A. japonicus* CTR/CLR-type receptor

(AjCTR) was positioned in a clade with other deuterostome CTR/CLR-type receptors,

whilst the other clade comprised CTR/CLR-type receptors from protostomes.

Similarly, two *A. japonicus* PDF-type receptors (AjPDFR1 and AjPDFR2) were

positioned in a clade comprising PDF-type receptors from other deuterostomes, whilst

another clade comprised PDF-type receptors from protostomes (Figure 1C). Domain

prediction of the candidate receptors showed that AjCTR had HRM (Hormone

receptor) and 7tm_GPCRs (seven-transmembrane G protein-coupled receptor)

superfamily domains (Figure 1-figure supplement 4A), whereas AjPDFR1 and

AjPDFR2 only presented the 7tm_GPCRs superfamily domain (Figure 1-figure

supplement 4B). To further investigate an orthologous relationship of the candidate

receptors with homologous receptors in other echinoderms, we conducted a gene

synteny analysis in *A. japonicus*, *A. rubens* and *L. variegatus*. The neighboring genes

of *AjCTR* included *GOLGA4*, *NeuroD*, *CRF2*, *ppGalNAc-T*, *URB3*, *SLC40A1*, *CMAH*

and *MAGP*. The relative positions of these genes were different from those in *A.*

rubens and *L. variegatus* (Figure 1-figure supplement 4C). The same situation was

also observed for the synteny analysis of PDFRs.

Figure 1. CT-type neuropeptide precursors and candidate receptors for CT-type peptides in *A.*
japonicus. (A) Alternative splicing schematic of *AjCTP1* and *AjCTP2*. Exons are represented by
green rectangles, introns are represented by lines, AjCT1 and AjCT2 are represented by red and

orange rectangles respectively, upstream and downstream non-coding regions are represented by
gray rectangles. The numbers represent the length of exons or introns. (B) Phylogenetic analysis
of AjCTP1 and AjCTP2 and CT-related neuropeptide precursors from other bilaterians. CT-type
precursors from deuterostomes are shown in green (echinoderms) and purple (chordates), CT-type
precursors from protostomes are shown in pink. (C) Phylogenetic analysis of relationships of
AjCTR, AjPDFR1 and AjPDFR2 (red letters) with CTR/CLR (calcitonin receptor/calcitonin
receptor-like receptor), DH31R (diuretic hormone 31 receptor) and PDFR (pigment-dispersing
factor receptor) in other taxa (black letters), using VP/OT-type (vasopressin/oxytocin) receptors as
the outgroup. Considering that VIP (vasoactive intestinal peptide) and PDF both have roles in
regulation of circadian and gut activity, the phylogenetic tree also includes VIPRs. Full species
names and accession numbers are listed in Supplementary table 2.

2.3 AjCT1 and AjCT2 act as ligands for AjCTR, AjPDFR1 and AjPDFR2

To investigate if AjCT1 and/or AjCT2 act as ligands for AjCTR, AjPDFR1 and

AjPDFR2, here, two receptor assay methods were employed. As shown in Figure 2A,

both AjCT1 and AjCT2 (10^{-6} M) caused a significant increase in CRE luciferase

activity in HEK293T cells transfected with AjCTR, AjPDFR1 or AjPDFR2.

Furthermore, this effect was inhibited by pre-incubation of cells with the PKA

inhibitor H89. The intracellular accumulation of cAMP was also measured, and this

showed that AjCT1 and AjCT2 (10^{-9} – 10^{-5} M) caused a dose-dependent increase in

cAMP in HEK293T cells transfected with AjCTR, AjPDFR1 or AjPDFR2 (Figure 2B).

However, when exposed to AjCT1 or AjCT2, the maximum accumulation of

intracellular cAMP was higher in cells transfected with AjCTR than in cells

transfected with AjPDFR1 and AjPDFR2 (Figure 2B). Measurement of SRE

luciferase activity revealed that AjCT1 and AjCT2 (10^{-6} M) administration evoked an

increase in activity in HEK293T cells transfected with AjCTR and AjPDFR1 (Figure

3A). However, in HEK293T cells transfected with AjPDFR2, an increase in SRE

luciferase activity was only observed with AjCT2, not with AjCT1. Furthermore, as

shown in Figure 3A, AjCT1- and AjCT2-induced increases in SRE luciferase activity

were not observed in cells that had been pre-incubated with FR900359 (Gαq protein

inhibitor) and Gö 6983 (PKC inhibitor). As shown in Figure 3-figure supplement 1,

receptors localized in the cell membrane, but after exposure to AjCT1 and AjCT2 (10^{-}

^{6} M) for 15 min, internalization of pEGFP-N1/*AjCTR*, pEGFP-N1/*AjPDFR1* and

pEGFP-N1/*AjPDFR2* was observed in transfected HEK293T cells (Figure 3B). Taken

together, all these findings indicate that AjCT1 and AjCT2 can act as ligands for the

three candidate receptors and then activate cAMP/PKA signaling and Gαq/Ca^{2+}/PKC

signaling in all combinations, with the exception of AjCT1/AjPDFR2 which only

activates the cAMP/PKA cascade.

To investigate whether the activated signaling pathways may exert downstream

effects via the ERK–MAPK pathway, the phosphorylation levels of ERK1/2 were

analysed in transfected HEK293T cells. pERK1/2 levels in cells transfected with

AjCTR or AjPDFR1, but not AjPDFR2, were significantly increased after a 5 min

incubation with AjCT1 (Figure 4A). Furthermore, this effect was blocked by the PKC

inhibitor in AjCTR-expressing cells, and this effect was blocked by both PKA and

PKC inhibitors in AjPDFR1-expressing cells (Figure 4-figure supplement 1A).

pERK1/2 levels in cells transfected with AjCTR or AjPDFR1 or AjPDFR2, were

significantly increased after a 5 min incubation with AjCT2 (Figure 4B). Moreover,

this effect persisted until 15 min after exposure to AjCT in both AjCTR-transfected

cells and AjPDFR1-transfected cells but not in AjPDFR2-transfected cells (Figure 4B,

Table 1). Furthermore, these effects were blocked by PKA and PKC inhibitors in

AjPDFR1 and AjPDFR2 expressing cells, whereas in AjCTR expressing cells,

peptide-induced pERK1/2 activity was only blocked by a PKC inhibitor (Figure 4-

figure supplement 1B). These results indicate that, with the exception of

AjCT1/AjPDFR2, both AjCT1 and AjCT2 can activate the ERK1/2 cascade by

binding to the candidate receptors.

Figure 2. Pharmacological characterisation of AjCT1 and AjCT2 as ligands for the *A. japonicus* receptors AjCTR, AjPDFR1 and AjPDFR2. (A) CRE-driven luciferase activity measured in HEK293T cells transfected with one of the three candidate receptors after exposure to AjCT1 or AjCT2 or neuropeptide + DMSO or neuropeptide + H89 (10 μM) or serum-free DMEM (CO, negative control). The neuropeptide concentration used here was 10^{-6} M. Mean values with standard deviations (n = 9) are shown. * indicates a statistically significant difference with $p <$ 0.05. (B) Measurement of cAMP accumulation after exposure to AjCT1 or AjCT2 (10^{-9} – 10^{-5} M). No cAMP elevation is observed in control experiments where HEK293T cells were transfected with empty pcDNA 3.1(+) (black circle). The estimated EC_{50} of AjCT1 for AjCTR, AjPDFR1 and

AjPDFR2 were 2.04×10^{-7} M (95% CI: 1.06×10^{-7} -8.69×10^{-7} M), 9.12×10^{-7} M (95% CI: 5.56

$\times$ 10^{-7} -4.45 $\times$ 10^{-6} M) and 6.09 $\times$ 10^{-8} M (95% CI: 2.23 $\times$ 10^{-8} -2.84 $\times$ 10^{-7} M) respectively. The

estimated EC$_{50}$ of AjCT2 for AjCTR, AjPDFR1 and AjPDFR2 were 1.50 $\times$ 10^{-8} M (95% CI: 1.03

$\times$ 10^{-9} -3.89 $\times$ 10^{-8} M), 1.44 $\times$ 10^{-6} M (95% CI: 1.04 $\times$ 10^{-6} -1.99 $\times$ 10^{-6} M) and 7.90 $\times$ 10^{-7} M (95%

CI: 5.73 $\times$ 10^{-7} -1.06 $\times$ 10^{-6} M) respectively. Error bars represent SEM for three independent

experiments.

Figure 3. SRE-driven luciferase activity and receptor internalization in cells transfected with
AjCTR, AjPDFR1 or AjPDFR2 and exposed to AjCT1 or AjCT2. (A) SRE-driven luciferase
activity measured after incubation with neuropeptide (10^{-6} M AjCT1 or AjCT2)/DMSO,
neuropeptide (10^{-6} M)/FR900359 (1uM) and neuropeptide (10^{-6} M)/Gö 6983 (1 μM). Mean values
with standard deviations (n = 9) are shown. * indicates a statistically significant difference with *p*

< 0.05. (B) Internalization of AjCTR, AjPDFR1 and AjPDFR2 after 15 min treatment with 10^{-6} M

AjCT1 or AjCT2. Green fluorescence represents the localization of pEGFP-N1/receptors in cells.

The cell nucleus probe (DAPI) was used for cell nucleus staining. Scale bar: 50 µm. All pictures

are representative of three independent experiments.

Figure 4. ERK1/2 activity in cells transfected with AjCTR, AjPDFR1 or AjPDFR2 and exposed to

AjCT1 or AjCT2. (A) and (B) show the immunoblot intensity of representative phosphorylation

bands. The transfected HEK293T cells were incubated with AjCT1 (10^{-6} M) or AjCT2 (10^{-6} M) for

0, 5, 15, 30, 45 and 60 min. The p-ERK1/2 was normalized based on t-ERK1/2. All images are

representative of three independent experiments. Error bars represent SEM for three independent

experiments. Statistically significant differences are indicated as follows: *: $p < 0.05$, **: $p < 0.01$.

Table 1. Activation of signaling pathways

Assays / Receptors	cAMP/PKA/ERK1/2		Gαq/Ca^{2+}/PKC/ERK1/2	
	AjCT1	AjCT2	AjCT1	AjCT2
AjCTR	×*	×*	√	√
AjPDFR1	√	√	√	√
AjPDFR2	×*	√	×	√

A tick indicates cAMP/PKA/ERK1/2 or Gαq/Ca^{2+}/PKC/ERK1/2 activation, the cross indicates no

activation, the cross with * indicates that the cAMP/PKA pathway is activated, while the ERK1/2

pathway is not activated.

2.4 Expression profile of AjCTP1/2 and its receptors in A. japonicus

qRT-PCR analysis revealed that *AjCTP1/2* had the highest expression level in the

circumoral nervous system (CNS) (Figure 5A). The expression levels of three

receptors differed from those of *AjCTP1/2*. *AjCTR* (Figure 5B) and *AjPDFR1* (Figure

5C) showed the highest expression level in CNS and longitudinal muscle, whereas

AjPDFR2 exhibited highest expression in the respiratory tree (Figure 5D).

297
298 Figure 5. The relative expression level of *AjCTP1/2* (A), *AjCTR* (B), *AjPDFR1* (C) and *AjPDFR2*
299 (D) in different tissues. The circumoral nervous system was selected as the control group. Values
300 are means ± SEM (*AjCTP1/2*, n = 6; *AjCTR*, *AjPDFR1* and *AjPDFR2*, n = 5). Asterisks represent
301 significant differences as follows: *: $p < 0.05$, **: $p < 0.01$, ***: $p < 0.001$. CNS, circumoral

nervous system; Gd, gonad; Ie, intestine; LM, longitudinal muscle; RT, respiratory tree.

To investigate the distribution of CT-type neuropeptides in *A. japonicus* using immunohistochemistry we employed an antiserum to the starfish (*Asterias rubens*) CT-type neuropeptide ArCT, which has been used previously for visualization of the expression of ArCT in *A. rubens* (Cai et al., 2018) and a CT-type neuropeptide in the feather star *Antedon mediterranea* (AmCT; Aleotti et al., 2022). The antigen peptide that was used to generate the ArCT antiserum (KYNSPFGASGP-NH$_2$) shares a C-terminal FGxxGP-NH$_2$ (where X is variable) motif with both AjCT1 (FGSAGP-NH$_2$) and AjCT2 (FGSGGP-NH$_2$) and therefore antibodies to ArCT may cross-react with AjCT1 and/or AjCT2. Accordingly, the ArCT antiserum revealed immunostaining in the central nervous system (radial nerve cords and circumoral nerve ring) (Figure 6A–F), which was abolished by pre-incubation of the antiserum with the ArCT antigen peptide (Figure 6-figure supplement 1). More specifically, high magnification images revealed immunostaining was localised in the ectoneural region, but not in the hyponeural region, of the radial nerve cords, with bilaterally symmetrical immunostained cells (neuronal somata) present in the ectoneural epithelium (Figure 6C). Likewise, immunostained cells were revealed in the ectoneural region of the circumoral nerve ring (Figure 6D-F). Furthermore, immunostaining was also revealed in tentacle nerves, which emanate from the circumoral nerve ring (Figure 6D-F), and in several regions of the digestive system, including the pharynx/stomach (Figure 6G), intestine (Figure 6H) and cloaca (Figure 6I). No immunostaining was observed in the longitudinal muscles (Figure 6A, B).

Figure 6. Immunohistochemical localization of CT-type neuropeptide expression in *A. japonicus* using antibodies to the starfish CT-type neuropeptide ArCT. A. Transverse section showing immunostaining in the radial nerve cord. B. Longitudinal section showing immunostaining in the radial nerve cord. C. High magnification image of a transverse section showing immunostained cells (arrowheads) and immunostained fibers in the neuropile (asterisks) of the ectoneural region of the radial nerve cord. Note the absence of immunostaining in the hyponeural region. D, E. Longitudinal sections showing immunostaining in the circumoral nerve ring and in tentacle nerves that emanate from the circumoral nerve ring. F. High magnification image of a transverse section showing immunostained cells (arrowheads) in the circumoral nerve ring. G. Longitudinal section showing immunostained fibers (asterisk) at the junction between the pharynx and the stomach. H. Longitudinal section showing immunostained fibers (arrowheads) in the intestine. I. Longitudinal section showing immunostained fibers (arrowheads) in the cloaca. Abbreviations: An, anus; BW, body wall; Cl, cloaca; CONR, circumoral nerve ring; Ec, Ectoneural region of the radial nerve cord; Hy, hyponeural region of the radial nerve cord; LM, longitudinal muscle; Lu, lumen; Mu, mucosa; RNC, radial nerve cord; TN, tentacle nerve. Scale bar, as shown in panel I, is 100 µm in A, B, D, E, I; 50 µm in F, G, H; 25 µm in C.

2.5 AjCT1 and AjCT2 causes dose-dependent relaxation of longitudinal muscle and intestine preparations from A. japonicus

Both AjCT1 and AjCT2 caused dose-dependent relaxation of longitudinal

muscles from *A. japonicus* at 10^{-10}–10^{-6} M (Figure 7). The maximum relaxant effects

of AjCT2 (10^{-6} M) on longitudinal muscles are illustrated in Figure 7-figure

supplement 1A. At the highest concentration tested (10^{-6} M), AjCT1 and AjCT2

caused 30.07 ± 7.49% and 32.50 ± 14.49% reversal of acetylcholine (ACh)-induced

contraction, respectively (Figure 7A, 7B). AjCT1 had no effect on the contractile state

of intestine preparations, even at concentrations as high as 10^{-5} M (Figure 7C). Unlike

AjCT1, AjCT2 caused dose-dependent relaxation of intestine preparations, with 26.33

± 8.75% reversal of ACh-induced contraction observed at 10^{-5} M (Figure 7D).

Control experiments in which ACh-induced contraction was recorded without

addition of CT-type peptides showed that ACh-induced contraction of longitudinal

muscles and intestinal preparations persisted for ≥15 minutes, confirming that

relaxation effects observed could be attributed to addition of AjCT1 or AjCT2 (Figure

7-figure supplement 1B, 1C).

Figure 7. Pharmacological effects of AjCT1 and AjCT2 on longitudinal muscle and intestine preparations from *A. japonicus*. (A) and (B) show representative recordings of the relaxing effects of AjCT1 and AjCT2 on longitudinal muscle preparations. The graphs show the dose-dependent relaxing effects of AjCT1 and AjCT2 (10^{-10} – 10^{-6} M), calculated as the percentage reversal of the contracting effect of 10^{-6} M ACh. (C) and (D) show representative recordings of experiments in which AjCT1 (10^{-7} – 10^{-5} M) and AjCT2 (10^{-9} – 10^{-5} M) were tested on intestine preparations. AjCT1 had no effect, but AjCT2 caused relaxation and the graph shows the dose-dependent relaxing effect of AjCT2 (10^{-9} – 10^{-5} M), calculated as the percentage reversal of the contracting effect of 10^{-5} M ACh. All recordings shown are representative of three independent experiments.

Mean values ± SEM were determined from three preparations. Lowercase letters a, b and c above

columns indicate the statistical difference at $p < 0.05$, whilst there is no significant difference
between bc and c.

2.6 AjCT2 affects feeding and growth in A. japonicus

To investigate the effects of AjCT1 and AjCT2 *in vivo*, a long-term injection
experiment was carried out at concentrations of 5×10^{-3} mg/mL (low-dose group) and
5×10^{-1} mg/mL (high-dose group), with the injection volume calculated based on the
wet weight of sea cucumbers at 1 μL/g. As shown in Figure 8A, there were no
significant changes in weight gain rate (WGR) and specific growth rate (SGR) in the
AjCT1 low concentration group (CT1L), AjCT1 high concentration group (CT1H)
and AjCT2 low concentration group (CT2L) ($p \geq 0.05$), but in the AjCT2 high
concentration group (CT2H) there was a significant increase in WGR and SGR in
comparison with the control group (CO) ($p < 0.05$). Correspondingly, the mass of
feces increased and the residual bait mass decreased (Figure 8A-figure supplement 1).
Furthermore, the relative intestinal expression levels of several growth factors were
investigated and the results showed that *AjGDF-8*, a growth inhibitory factor (Li et al.,
2016), was significantly down-regulated in all the experimental groups. In contrast,
the relative expression levels of *AjIgf* and *AjMegf6,* growth promoting factors
(Párrizas and LeRoith, 1997; Deribe et al., 2009), were significantly up-regulated in
intestines (Figure 8B).

Subsequently, the relative intestinal expression levels of AjCT1 and AjCT2
receptors were further tested for the CT2H group. The results showed that the relative
expression levels of all three receptors were significantly up-regulated, with the
ranked fold change being *AjCTR > AjPDFR1 > AjPDFR2* (Figure 8C). Conversely,
the relative expression levels of the three receptor genes were significantly down-
regulated in *siAjCTP1/2*-1 treated animals (Figure 9A, 9B and 9C). The interference
rate for si*AjCTP1/2*-1 was 66 % (Figure 9-figure supplement 1).

In long-term AjCT2 loss-of-function studies, the si*AjCTP1/2*-1 group exhibited
significant reductions in WGR and SGR compared to the control group (siNC) ($p <$
0.05, Figure 10A). Correspondingly, except for phase I, the si*AjCTP1/2*-1 group

showed significantly increased remaining bait and decreased excrement across phases

II-VI (Figure 10B). Furthermore, the growth inhibitory factor *AjGDF-8* was

significantly up-regulated and the growth promoting factor *AjMegf6* was significantly

down-regulated in the si*AjCTP1/2*-1 group (Figure 10C).

Figure 8. *In vivo* pharmacological tests revealed that AjCT1 and AjCT2 affect feeding and growth-
related gene expression in *A. japonicus*. (A) Weight gain rate (WGR) and specific growth rate
(SGR) in experimental and control groups. Values are means ± SEM (n = 3, representing three
parallel experiments of each group). (B) The relative intestinal transcript expression levels of
AjGDF-8, *AjIgf* and *AjMegf6* in different groups after injection of AjCT1 or AjCT2 or sterilized
seawater (CO) *in A. japonicus*. Values are means ± SEM (n = 5 biological duplication). (C) The
relative intestinal expression levels of AjCT1/2 receptors (*AjCTR*, *AjPDFR1* and *AjPDFR2*) in
CT2H and CO groups after 24 days injection experiment. Values are means ± SEM (n = 5
biological duplication). CO: control group (sterilized seawater); CT1L/CT2L: AjCT1/AjCT2 low
concentration group (5×10^{-3} mg/mL); CT1H/CT2H: AjCT1/AjCT2 high concentration group (5
$\times 10^{-1}$ mg/mL). * indicates statistically significant differences with $p < 0.05$, ** indicates
statistically significant differences with $p < 0.01$.

Figure 9. The relative intestinal expression levels of AjCT1/2 receptors after *AjCTP1/2*-1
knockdown. (A), (B) and (C) show the relative expression levels of *AjCTR*, *AjPDFR1* and
AjPDFR2 respectively. Values are means ± SEM (n = 5). ** indicates statistically significant
differences with $p < 0.01$.

Figure 10. Effects of 24-day *AjCTP1/2*-1 knockdown *in vivo* on feeding and growth in *A.*
japonicus. (A) Weight gain rate (WGR) and specific growth rate (SGR) in siNC and *siAjCTP1/2*-1
groups. Values are means ± SEM (n = 3, representing three parallel experiments for each group).
(B) Calculation of the remaining bait and excrement mass in different phases. The 24-day
experiment was divided into six phases (I-VI), with each phase comprising four days. Values are
means ± SEM (n = 3, representing three parallel experiments for each group). (C) The relative
intestinal transcript expression levels of *AjGDF-8*, *AjIgf* and *AjMegf6* in siNC and *siAjCTP1/2*-1
groups. Values are means ± SEM (n = 5 biological duplication). siNC: negative control group;
si*AjCTP1/2*-1: effectively experimental groups targeting *AjCTP1/2*. Asterisks represent
statistically significant differences as follows: *: $p < 0.05$, **: $p < 0.01$, ***: $p < 0.001$.

2.7 AjPDFR2 mediates CT-type signaling-induced feeding and growth promotion in A. japonicus

The differential activation of signaling pathways and associated physiological outcomes between AjCT2 and AjCT1 suggested that AjCT2 may promote feeding and growth via AjPDFR2-mediated mechanisms. To investigate this hypothesis, we performed long term knockdown experiments targeting AjPDFR2. As shown in Figure 11A, si*AjPDFR2-1* effectively knocked down the expression of *AjPDFR2*. In the si*AjPDFR2-1* group, both WGR and SGR were significantly decreased compared to siNC group (Figure 11B). Correspondingly, the remaining bait weight was significantly increased in si*AjPDFR2-1* group (except during phase I), and the weight of excrement was significantly decreased in phase V and VI (Figure 11C). Additionally, the growth promoting factor *AjIgf* was significantly down-regulated in the si*AjPDFR2-1* group (Figure 11D).

Figure 11. Effects of *AjPDFR2*-1 knockdown *in vivo* on feeding and growth in *A. japonicus*. (A)
The relative expression level of *AjPDFR2* in siNC, si*AjPDFR2*-1 and si*AjPDFR2*-2. (B) Weight
gain rate (WGR) and specific growth rate (SGR) in siNC and *siAjPDFR2*-1 groups. Values are
means ± SEM (n = 3, representing three parallel experiments for each group). (C) The remaining
bait and excrement weight in different phases. The 24-day experiment was divided into six phases
(I-VI), with each phase comprising four days. Values are means ± SEM (n = 3, representing three
parallel experiments of each group). (D) The relative intestinal transcript expression levels of
AjGDF-8, *AjIgf* and *AjMegf6* in siNC and *siAjPDFR2*-1 groups. Values are means ± SEM (n = 5
biological duplication). siNC: negative control group; si*AjPDFR2*-1 and si*AjPDFR2*-2:

458 experimental groups targeting A*jPDFR2*. Asterisks represent statistically significant differences as
459 follows: *: $p < 0.05$, **: $p < 0.01$, ***: $p < 0.001$. ns: not significant.
460
3. Discussion
Previous studies have revealed that the CT-type neuropeptide DH_{31} can act as a
ligand for a CTR/CLR-type receptor and a pigment-dispersing factor (PDF) receptor
in a protostome – the insect *D. melanogaster* (Mertens et al., 2005; Shafer et al., 2008;
Schwartz et al., 2019). Here we show that two CT-type peptides (AjCT1 and AjCT2)
act as ligands for a CTR/CLR-type receptor and two pigment-dispersing factor (PDF)
type receptors in a deuterostome – the sea cucumber *A. japonicus* (phylum
Echinodermata). This is an important finding because it indicates that the ability of
CT-type neuropeptides to act as ligands for both CTR/CLR-type receptors and PDF-
type receptors may be an evolutionarily ancient property that originated in a common
ancestor of the Bilateria. Alternatively, the ability of CT-type neuropeptides to act as
ligands for PDF-type receptors in *D. melanogaster* and *A. japonicus* may have
evolved independently. Further studies on a wider variety of both protostome (e.g.
molluscs, annelids) and deuterostome taxa (e.g. other echinoderms, hemichordates)
are needed to address this issue. Here, we also investigated the signal transduction
pathways that are activated by AjCT1 and AjCT2 when these peptides bind to their
receptors and then proceeded to investigate the expression and actions of these
peptides to gain insights into their physiological/behavioral roles in *A. japonicus*, as
discussed below.
480
3.1 Alternative RNA splicing gives rise to transcripts encoding one (AjCT2) or two
(AjCT1, AjCT2) CT-type neuropeptides in A. japonicus
In vertebrates, alternative RNA splicing gives rise to transcripts encoding CT or
CGRP (Amara et al., 1982; Morris et al., 1984). Analysis of transcriptome/genome
sequence data from *A. japonicus* has revealed that in this species alternative RNA
splicing gives rise to transcripts encoding one (AjCT2) or two (AjCT1, AjCT2) CT-
type neuropeptides (Rowe et al., 2014; Zandawala et al., 2017; Chen et al., 2019; this
study). A similar pattern of alternative RNA splicing has been reported for CT-type

precursors in other sea cucumbers (*H. scabra* and *H. glaberrima*; class Holothuroidea)

and in brittle stars (*Amphiura filiformis* and *Ophiopsila Aranea*; class Ophiuroidea)

(Zandawala et al., 2017; Suwansa-Ard et al., 2018). In contrast, CT-type precursors

identified in other echinoderm classes (Asteroidea, Echinoidea and Crinoidea)

comprise a single CT-type neuropeptide (Rowe and Elphick, 2012; Mirabeau and Joly,

2013; Rowe et al., 2014; Semmens et al., 2016; Cai et al., 2018; Aleotti et al., 2022).

However, it remains to be determined if this feature of CT-type precursors in sea

cucumbers and brittle stars evolved independently or originated in a common ancestor

of eleutherozoan echinoderms but with subsequent loss in Asteroidea and Echinoidea

(Figure 12).

Analysis of the sequences of AjCT1 and AjCT2 revealed the presence of two

cysteine residues in their N-terminal regions, which is an evolutionarily conserved

feature of CT-type peptides. Structural analysis of CT-type peptides in vertebrates and

in starfish has revealed that these cysteines form a disulphide bridge in the mature

peptides (Cai et al., 2018) and therefore it is inferred that this post-translational

modification also occurs in AjCT1 and AjCT2. Furthermore, the C-terminal residue of

AjCT1 and ACT2 is predicted to be an amidated proline, which is a conserved post-

translational modification of CT/DH31-type peptides in many taxa, but not vertebrate

CGRPs (Cai et al., 2018).

3.2 Identification and characterisation of receptors for CT-type neuropeptides in A.

japonicus

Informed by previous genome-wide analysis of GPCRs in *A. japonicus* (Huang

et al., 2021), we identified three candidate receptors for AjCT1 and/or AjCT2. Firstly,

AjCTR, which is a homolog of CT/DH$_{31}$-type receptors that have been identified in

other taxa, including vertebrates and insects (Garelja et al., 2022; Johnson et al., 2005;

Toribio et al., 2003). Secondly, AjPDFR1 and AjPDF2, which are homologs of the

PDF-type receptor that has been shown to be activated by DH$_{31}$ in *D. melanogaster*

(Mertens et al., 2005; Zandawala, 2012).

In mammals, CTR and CLR mediate effects of all CT-related neuropeptides, but

their ligand specificity relies on the presence and types of receptor activity-modifying proteins (RAMPs) (McLatchie et al., 1998; Purdue et al., 2002; Conner et al., 2004; Hay et al., 2018). Thus, CTR acts as a functional receptor for CT-type peptides without an associated RAMP (Katafuchi et al., 2009), whereas activation of downstream signaling via CLR requires an associated RAMP (Hay et al., 2018). Here, we investigated the occurrence of RAMPs in *A. japonicus* using the tblastn method employed by Sekiguchi *et al.*, (2016), which enabled identification of RAMPs in the cephalochordate *B. floridae*. However, no RAMPs were identified in *A. japonicus*, suggesting that AjCTR may be functionally more similar to CTR than CLR. Accordingly, tblastn analysis of other available echinoderm genomes (*A. rubens*, *S. purpuratus* and *Anneissia japonica*), as well as transcriptome databases for brittle stars (*O. victoriae*, *O. aranea* and *A. filiformis*), also failed to identify RAMPs. Likewise, RAMPs have also not been found in protostomes (Schwartz et al., 2019). Therefore, RAMPs may be unique to chordates, although interestingly RAMPs have also not been identified in the urochordate *C. intestinalis* (Sekiguchi et al., 2009) (Figure 12).

As highlighted above, here we show that the CT-type peptides AjCT1 and AjCT2 both act as ligands for a CTR/CLR-type receptor (AjCTR) and two PDF-type receptors (AjPDFR1, AjPDFR2) in *A. japonicus*. Furthermore, characterization of downstream signaling mechanisms revealed that both AjCT1 and AjCT2 trigger AjCTR-mediated cAMP accumulation, Ca^{2+} mobilization, and ERK signaling. These findings are consistent with previous studies on CT-type receptors in vertebrates and in the mollusc *Crassostrea gigas* (Pondel, 2000; Walker et al., 2010; Andreassen et al., 2014; Schwartz et al., 2019). Both AjCT1 and AjCT2 trigger cAMP accumulation, Ca^{2+} mobilization, and ERK signaling mediated by AjPDFR1. Both AjCT1 and AjCT2 triggered cAMP accumulation mediated by AjPDFR2, but only AjCT2 (not AjCT1) triggered Ca^{2+} mobilization and ERK signaling mediated by AjPDFR2. These differences in the signal transduction mechanisms activated by AjCT1 and AjCT2 may account for differences in the pharmacological activities and physiological roles of these peptides, as discussed below.

3.3 AjCT1 and AjCT2 act as muscle relaxants in A. japonicus

Analysis of the expression of the gene encoding the precursors comprising

AjCT2 or AjCT1 and AjCT2 revealed the occurrence of transcripts in the circumoral

nervous system as well as in peripheral tissues/organs, including the longitudinal

muscle, digestive system and gonads. Immunohistochemical analysis using an

antiserum to the starfish (*A. rubens*) CT-type neuropeptide ArCT enabled visualisation

of CT-type neuropeptide(s) in juvenile specimens of *A. japonicus*. Intense and

extensive immunostaining was revealed in the ectoneural region of the radial nerve

cords and circumoral nerve ring, consistent with the high levels *AjCTP1/2* transcripts

detected in circumoral nervous tissue. Immunostained nerve fibers were also observed

in the digestive system (including the intestine), consistent with the detection

AjCTP1/2 transcripts in intestine. However, no immunostaining was observed in

longitudinal muscles, which is inconsistent with the detection of *AjCTP1/2* transcripts

this tissue. This may reflect differences in the detection sensitivity of the techniques

used. In accordance with the expression pattern of AjCTP1/2 in *A. japonicus*, analysis

of CT-type neuropeptide expression in other echinoderms has revealed a widespread

pattern of expression in larval and adult starfish (*A. rubens*) and in juvenile and adult

featherstars (*Antedon mediterranea*) (Mayorova et al., 2016; Cai et al., 2018; Aleotti

et al., 2022).

Several CT-related peptides act as muscle relaxants in vertebrates (Steiner et al.,

1991; Pinto et al., 1996; Yoshimoto et al., 1998; Edvinsson et al., 2001; Golpon et al.,

2001; Kandilci et al., 2008) and, accordingly, previous studies have revealed that CT-

type neuropeptides cause relaxation of muscle preparations in starfish, including

locomotory organs (tube feet) and the apical muscle, which is functionally similar to

the longitudinal muscle in sea cucumbers (Cai et al., 2018). Therefore, it was of

interest to investigate if AjCT1 and/or AjCT2 act as muscle relaxants in *A. japonicus*.

Both AjCT1 and AjCT2 caused dose-dependent relaxation of longitudinal muscle

preparations, consistent with the relaxing effect of CT-type peptides on starfish apical

muscle preparations. It is worth noting that while AjCT2 at 10^{-6} M exhibited the

maximum efficacy in reversing ACh-induced contraction, its relaxing effect weakened

compared to the $10\square\square$ M treatment when applied at progressively higher

concentrations. This diminished efficacy likely stems from receptor desensitization,

induced by repeated AjCT2 application to longitudinal muscle preparations

(Tsurumaki et al., 2003; Arrowsmith and Wray, 2014). Furthermore, AjCT2, but not

AjCT1, caused dose-dependent relaxation of intestine preparations from *A. japonicus*.

Interestingly, the CT-type peptide ArCT does not to cause relaxation of cardiac

stomach preparations from the starfish *A. rubens* (Cai et al., 2018). Therefore, from a

functional perspective ArCT is more like AjCT1 than AjCT2 and we speculate that

this may reflect loss of an exon encoding a AjCT2-like peptide in the Asteroidea

lineage. Further insights into this issue and the evolution of CT-type neuropeptide

function echinoderms could be obtained by investigation of the actions of CT-type

peptides in brittle stars and sea urchins. Differences in the effects of AjCT1 and

AjCT2 on the intestine may be explained by differences in their signal transduction

mechanisms. Thus, only AjCT2, not AjCT1, triggered AjPDFR2-mediated activation

of Ca^{2+} mobilization and ERK signaling and therefore we speculate that the relaxing

effect of AjCT2 on the intestine may be mediated by activation of AjPDFR2.

3.4 AjCT2/AjPDFR2 stimulates feeding and growth in A. japonicus

In vivo pharmacological assays demonstrated that only high concentrations of

AjCT2 significantly enhanced feeding and growth rates in *A. japonicus*. In contrast,

neither a low concentration of AjCT2 nor any concentration of AjCT1 (low or high)

induced detectable effects. Furthermore, long-term knockdown of *AjCTP1/2* further

validated the essential role of AjCT2 in regulating feeding and growth in this species.

To elucidate the receptor mediating AjCT2's feeding- and growth-promoting effects,

we selected AjPDFR2 based on its distinct activation profile: AjCT2 selectively

activated AjPDFR2, inducing downstream ERK1/2 phosphorylation, whereas AjCT1

exhibited no activity toward this receptor. Given this receptor specificity, we

performed *AjPDFR2* knockdown experiments, which revealed phenotypic changes

consistent with those in *AjCTP1/2* knockdown animals, including significantly

reduced WGR and SGR, alongside increased remaining bait accumulation and diminished excrement output compared to control. Collectively, these results support a model wherein AjCT2 promotes feeding and growth in *A. japonicus* via AjPDFR2-dependent activation of the cAMP/PKA/ERK1/2 and Gαq/Ca^{2+}/PKC/ERK1/2 cascades. Considering the inherent complexity of neuropeptide signaling systems, which involve multiple GPCR subtypes coupled to diverse signaling cascades, ligands bound to the same receptor may activate distinct G protein subforms within a single cell (Møller et al., 2003; Mendel et al., 2020). Receptor activation modes may be modulated by structural polymorphisms or binding site diversity (Wong et al., 2000; Changeux, 2010), as well as by the differential efficacy of peptides in activating receptors *in vivo*.

Collectively, these findings indicate that AjCT2 has a physiological role in stimulating feeding and growth in *A. japonicus*. To the best of our knowledge, this is the first study to report orexigenic and growth-promoting effects of a CT-type neuropeptide in bilaterian animals. Interestingly, previous studies have reported the expression of CT-type neuropeptides in the digestive system of invertebrate chordates (Sekiguchi et al., 2009; Sekiguchi et al., 2016). However, in contrast to the stimulatory effect of AjCT2 on feeding/growth in *A. japonicus*, CT-type neuropeptides have been found to inhibit feeding and growth in vertebrates (Del et al., 2002; Hay et al., 2018; Nakamura et al., 2018). Thus, it appears that opposing roles of CT-type neuropeptide signaling systems as regulators for feeding/growth have evolved in deuterostomes, with a stimulatory role reported here in an echinoderm and an inhibitory role reported previously in vertebrates (Figure 12). Further research is now needed to investigate if the stimulatory effect of AjCT2 on feeding/growth observed in here in *A. japonicus* also applies to other sea cucumber species. Moreover, our discovery of this effect of AjCT2 on *A. japonicus* may provide a basis for development of novel approaches to improve aquaculture of this economically important species.

Figure 12. Phylogenetic summary of the occurrence and properties of CT-type and PDF-type signaling systems and effects of CT-type peptides on feeding/growth regulation in representative species. Echinoderms are boxed in purple with dashed lines and *A. japonicus*, the focus of this study, is framed in red with dashed lines. Pink filled rectangles indicate the occurrence of CT-type precursors/neuropeptides, CTR/CLR-type receptors and RAMPs. Green filled rectangles indicate the occurrence of PDF-type precursors/neuropeptides and receptors. Blue filled rectangles indicate species in which CT-type peptides have been shown to act as ligands for PDF-type receptors. Yellow filled rectangles indicate species in which effects or lack of effects of CT-type peptides on feeding/growth have been reported (I: Inhibition; P: Promotion; N: No effect). Red circles indicate that neuropeptides are generated by alternative splicing of precursor transcripts, with an unfilled circle indicating that transcripts encode one neuropeptide and with a filled circle indicating that transcripts can encode two neuropeptides in tandem. A yellow asterisk marks species in which neuropeptides have been shown to act as ligands for receptors experimentally. Unfilled rectangles indicate the absence of a protein. Unfilled rectanges containing a cross (X) indicate loss of a protein during evolution. Unfilled rectanges containing a question mark (?) indicate the existence of precursors/peptides that have yet to be identified.

 4. Materials and methods

Key Resources Table				
Reagent type (species) or resource	Designation	Source or reference	Identifiers	Additional information
Cell line (*Homo sapiens*)	HEK293 cell line	Procell	CL-0005	
Antibody	phosphorylated ERK1/2 kinases (Thr202/Tyr204) (goat)	Cell Signaling Technology	CAT#4930	IF (1:2000)
Antibody	ERK1/2 kinases (137F5) (goat)	Cell Signaling Technology	CAT#4695T	IF (1:1000)
Antibody	HRP-conjugated goat anti-rabbit IgG (goat)	Absin	Abs20040ss	IF (1:2000)
sequence-based reagent	Primers for pcDNA3.1(+) plasmid construction--AjCTR	This paper	PCR PRIMER	FORWARD CGCGGATCCAAAATGCAGGAGAACGATTC BamH □
sequence-based reagent	Primers for pcDNA3.1(+) plasmid construction--AjCTR	This paper	PCR PRIMER	REVERSE CCGGAATTCTTAAACTACCGTTGTTTTAT EcoR □
sequence-based reagent	Primers for pcDNA3.1(+) plasmid construction—AjPDFR1	This paper	PCR PRIMER	FORWARD CGCGGATCCACAATGAGTGTCGGGAACGT BamH □
sequence-based reagent	Primers for pcDNA3.1(+) plasmid	This paper	PCR PRIMER	REVERSE CCGGAATTCTCATTTCCCACCCCACTCAC

	construction— AjPDFR1			EcoR □
sequence-based reagent	Primers for pcDNA3.1(+) plasmid construction— AjPDFR2	This paper	PCR PRIMER	FORWARD CGCGGATCCAAGATGGCGAACA ATGAAAC BamH □
sequence-based reagent	Primers for pcDNA3.1(+) plasmid construction— AjPDFR2	This paper	PCR PRIMER	REVERSE CCGGAATTCTCATTCATTAAATT TGCCGT EcoR □
sequence-based reagent	Plasmid construction AjCTR–EGFP	This paper	PCR PRIMER	FORWARD CCGGAATTCTGAAAATGCAGGA GAACGATTC EcoR □
sequence-based reagent	Plasmid construction AjCTR–EGFP	This paper	PCR PRIMER	REVERSE CGCGGATCCCGAACTACCGTTG TTTTATCTG BamH □
sequence-based reagent	Plasmid construction AjPDFR1–EGFP			CCGGAATTCTGACAATGAGTGT CGGGAACGT EcoR □
sequence-based reagent	Plasmid construction AjPDFR1–EGFP			CGCGGATCCCGTTTCCCACCCC ACTCACTGT BamH □
sequence-based reagent	Plasmid construction AjPDFR2–EGFP			CCGGAATTCTGAAGATGGCGAA CAATGAAAC EcoR □
sequence-based reagent	Plasmid construction AjPDFR2–EGFP			CGCGGATCCCGTTCATTAAATTT GCCGTTCA BamH □
sequence-based reagent	Primers for qPCR assays β-Actin	This paper	PCR PRIMER	FORWARD AAGGTTATGCTCTTCCTCACGCT
sequence-based reagent	Primers for qPCR assays β-Actin	This paper	PCR PRIMER	REVERSE GATGTCACGGACGATTTCACG

sequence-based reagent	Primers for qPCR assays β-Tubulin	This paper	PCR PRIMER	FORWARD GAAAGCCTTACGACGGAACA
sequence-based reagent	Primers for qPCR assays β-Tubulin	This paper	PCR PRIMER	REVERSE CACCACGTGGACTCAAAATG
sequence-based reagent	Primers for qPCR assays AjCTP1/2	This paper	PCR PRIMER	FORWARD GTTTGGCGGTACCCATTAC
sequence-based reagent	Primers for qPCR assays AjCTP1/2	This paper	PCR PRIMER	REVERSE GGCGTTCGGTCTGTTAATAG
sequence-based reagent	Primers for qPCR assays AjMegf6	This paper	PCR PRIMER	FORWARD CGGTCCTTACTGTGCGGAGA
sequence-based reagent	Primers for qPCR assays AjMegf6	This paper	PCR PRIMER	REVERSE CGTCCTCCCAGCTGGACATC
sequence-based reagent	Primers for qPCR assays AjIgf	This paper	PCR PRIMER	FORWARD TGGAAGGCAGTAAGGCTA
sequence-based reagent	Primers for qPCR assays AjIgf	This paper	PCR PRIMER	REVERSE TAGGCTGATGTGGTTGGA
sequence-based reagent	Primers for qPCR assays AjGDF-8	This paper	PCR PRIMER	FORWARD ATACGCCCATACTCACATCC
sequence-based reagent	Primers for qPCR assays AjGDF-8	This paper	PCR PRIMER	REVERSE TGGCAGGAGTCAACCTTCAT
sequence-based reagent	Primers for qPCR assays AjCTR	This paper	PCR PRIMER	FORWARD GACGAATGGAGGAGAAGG
sequence-based reagent	Primers for qPCR assays AjCTR	This paper	PCR PRIMER	REVERSE CGCCGCAATTAGAGTAGT

sequence-based reagent	Primers for qPCR assays *AjPDFR1*	This paper	PCR PRIMER	FORWARD TTATCAACCGCCGTCTTC
sequence-based reagent	Primers for qPCR assays *AjPDFR1*	This paper	PCR PRIMER	REVERSE CAGGAGGAGGAACACATTC
sequence-based reagent	Primers for qPCR assays *AjPDFR2*	This paper	PCR PRIMER	FORWARD TGTTCTCCTGCCTCTTCT
sequence-based reagent	Primers for qPCR assays *AjPDFR2*	This paper	PCR PRIMER	REVERSE GTCCTTCTCATCGGTGTG
sequence-based reagent	Primers for 3'RACE *AjCTP1* *AjCTP2*	This paper	PCR PRIMER	GATTACGCCAAGCTTCGAGCTG TGTCGATAACATTACGCATGG
sequence-based reagent	Primers for 3'RACE AjPDFR1	This paper	PCR PRIMER	GATTACGCCAAGCTTGAGGGAA AGCCAAACCAGCGAGACG
sequence-based reagent	Primers for 3'RACE AjPDFR2	This paper	PCR PRIMER	GATTACGCCAAGCTTGTGCAAC CGGACGAAGATCCACGTG
sequence-based reagent	Primers for 5'RACE AjPDFR1	This paper	PCR PRIMER	GATTACGCCAAGCTTCGTCTCG CTGGTTTGGCTTTCCCTC
sequence-based reagent	Primers for 5'RACE AjPDFR2	This paper	PCR PRIMER	GATTACGCCAAGCTTGCCTTCC ACAAACATCCATGCGAACGC
sequence-based reagent	Primers for 5'RACE AjCTR	This paper	PCR PRIMER	GATTACGCCAAGCTTACGCCGG GCTCTCAAGAAGTCATTC
sequence	RNAi	This paper	si*AjCTP1*-1	GGAAUUAUCAAGACAGUAU (dT)(dT) AUACUGUCUUGAUAAUUCC (dT)(dT)
sequence	RNAi	This paper	si*AjCTP1*-2	GAAAGUCGCUAAUGCAGUA (dT)(dT) UACUGCAUUAGCGACUUUC

				(dT)(dT)
sequence	RNAi	This paper	si*AjPDFR2*-1	GUAGCGAUUGGUCGAAUAA (dT)(dT) UUAUUCGACCAAUCGCUAC (dT)(dT)
sequence	RNAi	This paper	si*AjPDFR2*-2	CAGUCAUUGUUAUCUACCA (dT)(dT) UGGUAGAUAACAAUGACUG (dT)(dT)
sequence	RNAi	This paper	siNC	UUCUCCGAACGUGUCACGU (dT)(dT) ACGUGACACGUUCGGAGAA (dT)(dT)
commercial assay or kit	SMARTer RACE 5'/3' Kit	Takara	Cat No. 634858	
commercial assay or kit	FastPure Gel DNA Extraction Mini Kit	Vazyme	DC301-01	
commercial assay or kit	Hifair® Ⅲ 1st Strand cDNA Synthesis SuperMix	YEASEN	Cat No. 11141ES60	
commercial assay or kit	Hieff UNICON Universal Blue qPCR SYBR Green Master Mix	YEASEN	Cat No. 11184ES08	
commercial assay or kit	Firefly luciferase reporter gene assay kit	MK	MF4001	
commercial assay or kit	cAMP Assay kit	R&D systems	Cat No. KGE002B	

4.1 Animals

Adult sea cucumbers were collected from an aquaculture breeding farm in

Weihai, Shandong, China. All the animals were acclimated in breeding aquaria (18 □,

32 ppt) for at least seven days and fed daily with a commercially formulated diet.

Tissues used for cDNA cloning and qRT-PCR were obtained from six adult *A.*

japonicus (~120 g) and kept at -80 □ until use. Juvenile specimens of *A. japonicus*

(~15 mm in length) were used for immunohistochemical analysis of CT-type

neuropeptide expression. *In vitro* functional analysis and *in vivo* experiments were

carried out using adult sea cucumbers with body masses of ~120 g and ~20 g,

respectively.

Ethical approval

All methods were carried out in accordance with the ARRIVE guidelines

(https://arriveguidelines.org) for animal experiments. All animal care and use

procedures were approved by the Institutional Animal Care and Use Committee of

Ocean University of China (Permit Number: 20141201). They were performed

according to the Chinese Guidelines for the Care and Use of Laboratory Animals

(GB/T 35892-2018).

4.2 Molecular structure and phylogenetic analysis of CT-type neuropeptides and

precursors

A. japonicus CT-type neuropeptide precursor sequences (*AjCTP1*, GenBank

accession number: MF401985; *AjCTP2*, GenBank accession number: MF401986)

were identified in our previous study (Chen et al., 2019). Based on the published

sequences, transcript-specific primers for *AjCTP1* and *AjCTP2* (Key resources table)

were designed using Primer3Plus (https://www.primer3plus.com/) and sequence

accuracy was confirmed by SMARTer RACE 5'/3' Kit (TaKaRa, Kusatsu, Japan)

according to the manufacturer's instruction. A sequence translation tool

(https://web.expasy.org/translate/) was used to obtain the corresponding amino acid

sequences encoded by the cloned cDNAs. To identify the conserved features of CT-

type neuropeptides in Bilateria, sequences were aligned using Clustal-X with default

settings and then predicted using the online tool Multiple Sequence Comparison

Display Tool (http://www.bio-soft.net/sms/index.html). A neighbor joining tree was constructed to investigate relationships of AjCTP1, AjCTP2 and CT-type precursors from other Bilateria. Amino acid sequences were aligned using MEGA7 (v.7170509) with default settings and bootstrap values were calculated from 1000 replicate analysis and then modified using iTOL (https://itol.embl.de/login.cgi). The accession numbers or citations for all sequences used for analysis are listed in Supplementary table 1.

4.3 Cloning, sequence analysis and phylogenetic tree construction for candidate AjCT1/2 receptors in A. japonicus

Five potential receptors for AjCT1 and AjCT2 in *A. japonicus* (GenBank accession number: PIK49998.1, PIK49999.1, PIK49604.1, PIK34007.1 and PIK62458.1) were identified based on genome-wide analysis (Huang et al., 2021). Sequence comparison revealed that both PIK49998.1 and PIK62458.1 were partial sequences of PIK34007.1. Thus, three candidate receptors belonging to Class B (Secretin-like) GPCRs named as *AjCTR* (Calcitonin receptor, GenBank accession number: PIK49604.1), *AjPDFR1* (Pigment-dispersing factor receptor 1, GenBank accession number: PIK34007.1), and *AjPDFR2* (Pigment-dispersing factor receptor 2, GenBank accession number: PIK49999.1) were predicted and then analyzed. To confirm the accuracy of these sequences, cDNAs were cloned using the SMARTer RACE 5'/3' Kit (Takara, Cat No. 634858) and then sequenced (Beijing Genomics Institute, China). Receptor sequence analysis and phylogenetic analysis were performed as described in 4.2. In addition, the online tool NetPhos-3.1 (https://services.healthtech.dtu.dk/service.php?NetPhos-3.1) was used to predict potential phosphorylation sites. The conserved domains of protein sequences were predicted by the Batch SMART plugin of TBtools (Toolbox for biologists, v.1.098652) software. For analysis of gene synteny, neighboring genes of *AjCTR*, *AjPDFR1* and *AjPDFR2* were found based on an *A. japonicus* chromosome-level genome assembly (Wang et al., 2022) and the corresponding genes in the starfish *Asterias rubens* and the sea urchin *Lytechinus variegatus* were also identified by performing BLASTp in

the NCBI database. Specific primers for cloning of cDNAs encoding the candidate

receptors are shown in the Key resources table and the details of the sequences used

for constructing the phylogenetic tree are listed in Supplementary table 2.

4.4 Testing AjCT1 and AjCT2 as ligands for candidate receptors and exploration of

downstream signaling pathways

AjCT1 and AjCT2 (both with a disulfide bridge between the two cysteine

residues and with C-terminal amidation) were custom synthesized by GL Biochem

Ltd. (Shanghai, China) with >95% purity. To test candidate receptors for CT-type

neuropeptides and investigate their downstream activation mechanisms the following

assays were used to measure cAMP signaling or Ca²☐ signaling: i). direct

measurement of cAMP levels using ELISA ii). indirect measurement of cAMP

signaling using a cAMP response element-Luciferase (CRE-Luc) assay that measures

cAMP-induced gene expression, iii). serum response element (SRE)-Luc assays that

measure Ca^{2+}-induced gene expression, iv). receptor translocation assays and v).

Western blotting.

AjCTR, *AjPDFR1* and *AjPDFR2* incorporating enhanced green fluorescent

protein (EGFP) were constructed respectively according to the method described by

Li *et al.* (2022). Briefly, the complete ORFs of candidate receptors were cloned and

sequenced. Then, the PCR products were digested with EcoRI and BamHI (TaKaRa,

Kusatsu, Japan) and then incorporated into the pEGFP-N1 vector (YouBio, Changsha,

China) that was subject to the same double-digestion. Finally, the constructed plasmid

with correct sequences was transfected into HEK293T cells seeded on glass

coverslips in 12-well plates. After incubation with AjCT1 (10^{-6} M), AjCT2 (10^{-6} M) or

serum-free DMEM (Biosharp, Hefei, China) for 15 min, cells were washed with PBS

(Biosharp, Hefei, China) and fixed with 4% PFA for 10 min. Finally, cell nuclei were

stained with DAPI (G-CLONE, Beijing, China) for 5 min and an Olympus bright field

light microscope was used for image capture. The pcDNA 3.1(+) vectors (YouBio,

Changsha, China) comprising the complete ORF of the three candidate receptors were

also constructed as described by Li *et al.* (2022). Then, they were co-transfected with

the cAMP reporter pCRE-Luc (YouBio, Changsha, China) or Ca^{2+} signaling reporter

pSRE-Luc plasmid including a PKC reporter gene (YouBio, Changsha, China) into

HEK293T cells. Incubation was carried out with AjCT1 (10^{-6} M), AjCT2 (10^{-6} M) or

serum-free DMEM (CO, negative control) for 4 h at 37 °C. Finally, luciferase activity

was tested using the firefly luciferase assay kit (MK, Shanghai, China). When

required, transfected cells described above were co-incubated with H89 (PKA

inhibitor) (Absin, Shanghai, China), FR900359 ($G\alpha q$ protein inhibitor) (Cayman

Chemical, Ann Arbor, Michigan) or Gö 6983 (PKC inhibitor) (Cayman Chemical,

Ann Arbor, Michigan) before the experiment to explore the potential signaling

pathways. To confirm the concentration of intracellular cAMP after stimulation,

HEK293T cells expressing pcDNA 3.1(+)/*AjCTR*, pcDNA 3.1(+)/*AjPDFR2* or

pcDNA 3.1(+)/*AjPDFR1* were treated with AjCT1 or AjCT2 (10^{-9} – 10^{-5} M),

respectively, for 15 min after 2 h starvation in serum-free medium. Subsequently,

cAMP was measured using an ELISA-based cAMP assay kit (R&D systems,

USA&Canada) following the manufacturer's instruction. HEK293T cells transfected

with empty pcDNA3.1(+) were used as a negative control. The specific primers used

for constructing plasmids are listed in Supplemental Table 1.

Western blotting was performed as described previously (Li et al., 2022). Briefly,

HEK293T cells transfected with the constructed pcDNA 3.1(+) plasmid were starved

in serum-free medium for 2 h, and then incubated with AjCT1 or AjCT2 (10^{-6} M).

PVDF membranes containing proteins were blocked with 5% BSA (G-CLONE,

Beijing, China) and incubated overnight with antibodies to phosphorylated ERK1/2

kinases (CST, Boston, China). After that, goat anti-rabbit horseradish peroxidase-

conjugated secondary antibody (Absin, Shanghai, China) was applied and incubated,

followed by the addition of Omni-ECL™ Femto Light Chemiluminescence reagents

(EpiZyme, Shanghai, China). Finally, RVL-100-G (ECHO, America) was applied for

visualization of the PVDF membrane. When required, transfected cells were co-

incubated with PKA or PKC inhibitor before the experiment to identify signaling

pathways that activate downstream ERK1/2 cascade.

4.5 Total RNA extraction and Quantitative Real-Time PCR (qRT-PCR) detection

qRT-PCR was employed to detect *AjCTR*, *AjPDFR1*, *AjPDFR2*, *AjCTP1* and

AjCTP2 transcript expression in five tissues from *A. japonicus*, including the intestine,

longitudinal muscle, respiratory tree, gonad and circumoral nervous system, and the

circumoral nervous system was selected as the control group. Because it was difficult

to design specific primers for *AjCTP1* and *AjCTP2* (Zheng et al., 2022), total

transcript expression levels for *AjCTP1/2* (*AjCTP1* + *AjCTP2*) were investigated.

Total RNA was extracted from the collected tissues using Trizol RNA isolation

reagent (Vazyme, Nanjing, China) and Hifair® □ 1st Strand cDNA Synthesis

SuperMix (YEASEN, Shanghai, China) was used to remove genomic DNA and

synthesize cDNA according to manufacturer's instructions. For qRT-PCR,

amplification was performed in 20 µL volume reactions based on Hieff UNICON

Universal Blue qPCR SYBR Green Master Mix (YEASEN, Shanghai, China) and

expression levels were assessed by a Corbett Rotoe-Gnen Q (Qiagen, Germany). β-

actin (GenBank accession number: PIK61412.1) and β-tubulin (GenBank accession

number: PIK51093) were used as house-keeping genes for normalization (Zhao et al.,

2014). The $2^{-\Delta\Delta CT}$ method was applied to analyze the relative expression levels.

Specific sequences of primers used for qRT-PCR are listed in Supplemental Table 1.

4.6 Localization of CT-type neuropeptide expression in A. japonicus using
immunohistochemistry

799
800

Juvenile specimens of *A. japonicus* (~15 mm in length) were fixed for 24 hours

at room temperature or for 48 hours at 4 ^{o}C in Bouin's fixative (75 mL saturated picric

acid in seawater, 25 mL 37% formaldehyde, 5 mL acetic acid) days and then washed

in water before being transferred to 50% ethanol followed by 70% ethanol. The fixed

specimens were in wrapped in tissue paper soaked in 70% ethanol and then

transported from China to the UK. Following dehydration through 90% and 100%

ethanol and clearing in xylene, specimens were embedded in paraffin wax and then

sectioned (9 or 10 µm) transversely or longitudinally using a Leica RM2145

microtome. Sections were mounted on chrome alum-gelatin coated glass slides and

then after incubation in xylene to remove wax followed by 100% ethanol, slides were

incubated in 1% hydrogen peroxide in phosphate buffered saline (PBS, pH 7.3) to

quench endogenous peroxidases. After using an ethanol series to rehydrate sections,

slides were blocked in 5% goat serum in PBS containing 0.1% Tween-20 (PBST).

Then slides were incubated overnight at 4 $^{\circ}$C with the ArCT antiserum diluted to

1:2000, 1:3000 or 1:4000 in PBST. Following washing in PBST, slides were

incubated with Peroxidase-AffiniPure Goat Anti-Rabbit IgG (H + L) conjugated to

Horseradish Peroxidase (RRID: AB_2313567; Jackson ImmunoResearch, West Grove,

PA) diluted 1:1000 in 2% goat serum/PBST for 3 h. Bound antibodies were revealed

using a solution of 0.05% diaminobenzidine (VWR Chemicals, UK), 0.05% nickel

chloride (Sigma-Aldrich, Gillingham, UK), 0.015% hydrogen peroxide (VWR

Chemicals, UK) diluted in PBS. When staining was observed, slides were washed in

water and dehydrated through an ethanol series into xylene and mounted with

coverslips over DPX mounting medium (Thermo Fisher Scientific). The specificity of

immunstaining was assessed by pre-absorption of the ArCT antiserum (at 1:100

dilution) for 1 h at room temperature with the ArCT antigen peptide (25 µM) prior to

testing on sections at 1:3000 dilution. Images of stained sections were taken using a

QIClich CCD Colour Camera (Qimaging, UK) linked to a DMRAI light microscope

(Leica Microsystems, UK) with Volocity v. 6.3.1 image analysis software (Perkin-

Elmer, USA) running on an iMac computer (27 inch with OS Yosemite, v. 10.10).

Montages of images were prepared and labelled using GIMP (version 3.0.2) and

Adobe Photoshop (version 26.7) running on a MacBook Pro laptop (OS Sequoia 15.5).

4.7 Analysis of the in vitro activity of AjCT1 and AjCT2 on preparations from A.

japonicus

Longitudinal muscles and intestines (20 mm in length) from *A. japonicus* were

dissected and then suspended in a 50 mL glass organ bath containing 20 mL sterilized

seawater. Both ends of the preprations were tied with cotton ligatures and then one

end was tied to a metal hook and the other end was connected with a High Grade

Isotonic Transducer (ADinstruments MLT0015, Oxford, UK). Before testing, the

preparation was stabilized for about 20 min, and the resting tension was adjusted to 1.0 g. Informed by previous findings from Cai et al. (2018), we investigated if AjCT1 and AjCT2 caused relaxation of preparations. First, acetylcholine (ACh) was applied to induce the contraction of preparations and then when a stable state of contraction was achieved, 20 uL AjCT1 or AjCT2 at a range of concentrations was added and relaxation responses were recorded. Output signal (cm) from PowerLab data acquisition hardware (ADinstruments PowerLab 4/26, Oxford, UK) was calculated using LabChart (v8.0.7) software.

4.8 Investigation of the in vivo pharmacological effects of AjCT1 and AjCT2 in A. japonicus

A total of 150 adult sea cucumbers were randomly assigned to five groups (30 individuals per group), including two high concentration groups (CT1H and CT2H, 5×10^{-1} mg/ml), two low concentration groups (CT1L and CT2L, 5×10^{-3} mg/ml) and the control group (CO, sterilized seawater), with each group consisting of three parallel experiments comprising ten animals. Before the experiments, sea cucumbers were starved for a day to normalize their physiological status and their wet weights were measured. AjCT1 and AjCT2 were dissolved in sterilized seawater and then injected through the peristome at noon every other day (1 μl/g wet weight) and the CO group were injected in the same way but with sterilized seawater (1 μl/g wet weight). Feces and residual bait were collected and weighed after drying every day. After 24 days, wet weights of sea cucumbers were measured again. Then, WGR and SGR were calculated to assess their feeding and growth conditions. Formulas are as follows: $WGR = 100*(W_t-W_0)/W_0$, $SGR = 100*(lnW_t-lnW_0)/D$. In the formula, W_0 represents the initial wet body weight (g), W_t represents the final wet body weight (g), D represents the feeding days. Subsequently, animals were dissected and intestines were collected for detecting the expression levels of the three receptor genes (*AjCTR, AjPDFR1, AjPDFR2*) and several growth factors, including multiple epidermal growth factor 6 (*AjMegf6*, GenBank accession number: MG018199.1), insulin-like growth factor (*AjIgf*, GenBank accession number: PIK50518.1) and growth and

differentiation factor-8 (*AjGDF-8*, GenBank accession number: KP100064.1) (Li et

al., 2016; Wang et al., 2018). The specific primers for qRT-PCR analysis are listed in

the Key resources table.

4.9 Loss-of-function experiments by RNA interference (RNAi)

To investigate how receptor expression is affected by knock-down of AjCTP

expression, an RNAi experiment was carried out. It was difficult to design *AjCTP1-*

and *AjCTP2-* specific interference oligonucleotides and therefore, we designed and

customized generic interference oligonucleotides targeting both *AjCTP1* and *AjCTP2*

(si*AjCTP1/2*-1 and si*AjCTP1/2*-2, containing the extra 2′ ome modification) (Beijing

Tsingke Biotech Co., Ltd. China). Meanwhile, a negative control siRNA (siNC) was

also customized for the experiment (Key resources table), which does not target genes

in *A. japonicus*. The si*AjCTP1/2*-1, si*AjCTP1/2*-2 or siNC dry powder was diluted to

20 μM with DEPC water following the manufacturer's instruction, then 60 μl

sterilized seawater was mixed with 20 μl si*AjCTP1/2*-1, si*AjCTP1/2*-2 or siNC, and

20 μl Lipo6000 transfection reagent (Beyotime, Shanghai, China) as the transfection

solution. Eighteen sea cucumbers were randomly divided into three groups and

injected with three different transfection solutions respectively. The mixed solutions

were injected through the peristome once a day (1 μl/g wet weight) and intestines

were dissected and frozen in liquid nitrogen 3 days after injection. The expression

level of *AjCTP1/2* was investigated by qRT-PCR and the interference rate was

calculated using $(1 - 2^{-\Delta\Delta CT})$ to determine the inhibition efficiency of siRNAs. Finally,

the relative expression levels of *AjCTR*, *AjPDFR1* and *AjPDFR2* were investigated

using the selected *A. japonicus* intestines.

To assess the effects of CT-type signaling inhibition on feeding and growth in *A.*

japonicus, a 24-day RNAi experiment was conducted following the methodology

outlined in Section 4.8. Sea cucumbers were randomly allocated into two groups: the

negative control group received siNC injections, while the experimental group was

injected with si*AjCTP1/2*-1 (effectively targeting *AjCTP1/2*). Throughout the

experiment, the dry weight of remaining bait and excrement, along with the wet
weight of individual sea cucumbers, were recorded. Subsequently, WGR and SGR, as
indices of feeding efficiency and growth performance, were calculated. In addition,
the expression levels of the three growth factors (*AjMegf6*, *AjIgf* and *AjGDF-8*) in the
intestinal tissue were tested by qPCR.

To investigate the potential mediation of CT-type signaling effects on feeding
and growth by AjPDFR2 in *A. japonicus*, an RNAi experiment targeting *AjPDFR2*
was performed following the same knockdown protocol established for *AjCTP*.
Briefly, the effective siRNA targeting *AjPDFR2* (si*AjPDFR2*-1 or si*AjPDFR2*-2) was
selected by testing *AjPDFR2* expression in the intestines of injected sea cucumbers
using qRT-PCR and calculating the interference rate efficiency. Subsequently, the
validated siRNA was then employed in a 24-day RNAi experiment. The dry weight of
remaining bait and excrement, as well as the wet weight of the sea cucumbers were
measured, enabling calculation of WGR and SGR. Furthermore, the expression levels
of three key growth factors (*AjMegf6*, *AjIgf* and *AjGDF-8*) in the intestinal tissue were
analyzed via qPCR.

4.10 Statistical analysis

Statistical analyses were carried out using GraphPad Prism 8.0. Significant
differences between groups were determined by one-way analysis of variance
(ANOVA) followed by a Tukey's multiple comparisons test. Statistical significance
thresholds were defined as follows: $p < 0.05$ (*) for significant differences and $p <$
0.01 (**) for highly significant differences, with annotations directly marked on the
figures. All data were presented as mean ± SEM from at least three independent
biological replicates. Non-linear regression analysis was performed using a four-
parameter logistic equation with automatic outlier exclusion to fit the dose-response
curve. The half maximal effective concentration (EC_{50}) was derived from the fitted
curves through built-in algorithms in GraphPad Prism 8.0.

Competing Interests

The authors declare that no competing interests exist.

Funding

This research was supported by National Natural Science Foundation of China [grant

number 42276103] and Biotechnology and Biological Sciences Research Council

(UK; grant number BB/M001644/1).

Author Contributions

Xiao Cong, Data curation, Formal analysis, Investigation, Methodology, Software,
Visualization, Writing—original draft preparation; Huachen Liu, Investigation,
Methodology, Validation; Lihua Liu, Methodology, Validation; Michaela Egertová,
Investigation, Methodology, Visualization; Maurice R. Elphick, Formal analysis,
Validation, Visualization, Writing—review & editing; Muyan Chen,
Conceptualization, Formal analysis, Funding acquisition, Methodology, Project
administration, Resources, Supervision, Validation, Visualization, Writing—review &
editing

References:

Aleotti A, Wilkie I C, Yañez-Guerra L A, et al. Discovery and functional characterization of
946   neuropeptides in crinoid echinoderms. Frontiers in Neuroscience, 2022, 16: 1006594. doi:
947   10.3389/fnins.2022.1006594.

948 Amara S G, Jonas V, Rosenfeld M G, et al. Alternative RNA processing in calcitonin gene
949   expression generates mRNAs encoding different polypeptide products. Nature, 1982,
950   298(5871): 240-244. doi: 10.1038/298240a0.

951 Andreassen K V, Hjuler S T, Furness S G, et al. Prolonged calcitonin receptor signaling by salmon,
952   but not human calcitonin, reveals ligand bias. PloS one, 2014, 9(3): e92042. doi:
953   10.1371/journal.pone.0092042.

954 Arrowsmith S, Wray S. Oxytocin: its mechanism of action and receptor signaling in the
955   myometrium. Journal of neuroendocrinology, 2014, 26(6): 356-369. dio: 10.1111/jne.12154.

956 Bomberger J M, Parameswaran N, Spielman W S. Regulation of GPCR trafficking by RAMPs.
957   Adv. Exp. Med. Biol, 2012, 744: 25-37. doi: 10.1007/978-1-4614-2364-5_3.

958 Bourlat S J, Juliusdottir T, Lowe C J, et al. Deuterostome phylogeny reveals monophyletic
959   chordates and the new phylum Xenoturbellida. Nature, 2006, 444(7115): 85-88. doi:
960   10.1038/nature05241

961 Bromham L D, Degnan B M. Hemichordates and deuterostome evolution: robust molecular
962   phylogenetic support for a hemichordate+ echinoderm clade. Evolution & development, 1999,
963   1(3): 166-171. doi: 10.1046/j.1525-142x.1999.99026.x

Burke R D, Angerer L M, Elphick M R, et al. A genomic view of the sea urchin nervous system. Developmental biology, 2006, 300(1): 434-460. doi: 10.1016/j.ydbio.2006.08.007.

Cai W, Kim C H, Go H J, et al. Biochemical, anatomical, and pharmacological characterization of calcitonin-type neuropeptides in starfish: discovery of an ancient role as muscle relaxants. Frontiers in Neuroscience, 2018, 12: 382. doi: 10.3389/fnins.2018.00382.

Changeux J P. Allosteric receptors: from electric organ to cognition. Annual review of pharmacology and toxicology, 2010, 50(1): 1-38. doi: 10.1146/annurev.pharmtox.010909.105741.

Chen M, Talarovicova A, Zheng Y, et al. Neuropeptide precursors and neuropeptides in the sea cucumber *Apostichopus japonicus*: a genomic, transcriptomic and proteomic analysis. Scientific Reports, 2019, 9(1): 8829. doi: 10.1038/s41598-019-45271-3.

Chen Y, Shyu J F, Santhanagopal A, et al. The calcitonin receptor stimulates Shc tyrosine phosphorylation and Erk1/2 activation: involvement of Gi, protein kinase C, and calcium[J]. Journal of Biological Chemistry, 1998, 273(31): 19809-19816. doi: 10.1074/jbc.273.31.19809.

Cong X, Liu H, Zheng Y, et al. A Putative Role of Vasopressin/Oxytocin-Type Neuropeptide in Osmoregulation and Feeding Inhibition of *Apostichopus japonicus*. International Journal of Molecular Sciences, 2023, 24(18): 14358. doi: 10.3390/ijms241814358.

Conner A C, Simms J, Hay D L, et al. Heterodimers and family-B GPCRs: RAMPs, CGRP and adrenomedullin. Biochemical Society Transactions, 2004, 32(5): 843-846. doi: 10.1042/BST0320843.

Conzelmann M, Williams E A, Krug K, et al. The neuropeptide complement of the marine annelid *Platynereis dumerilii*. BMC genomics, 2013, 14: 1-15. doi:10.1186/1471-2164-14-906.

Copp D H, Cameron E C, Cheney B A, et al. Evidence for calcitonin—a new hormone from the parathyroid that lowers blood calcium. Endocrinology, 1962, 70(5): 638-649. doi: 10.1210/endo-70-5-638.

Del Prete E, Schade B, Riediger T, et al. Effects of amylin and salmon calcitonin on feeding and drinking behavior in pygmy goats. Physiology & behavior, 2002, 75(4): 593-599. doi: 10.1016/s0031-9384(02)00655-8.

Deribe Y L, Wild P, Chandrashaker A, et al. Regulation of epidermal growth factor receptor trafficking by lysine deacetylase HDAC6. Science signaling, 2009, 2(102): ra84-ra84. doi: 10.1126/scisignal.2000576.

Dunn C W, Hejnol A, Matus D Q, et al. Broad phylogenomic sampling improves resolution of the animal tree of life. Nature, 2008, 452(7188): 745-749. doi: 10.1038/nature06614

Edvinsson L, Goadsby P J, Uddman R. Amylin: localization, effects on cerebral arteries and on local cerebral blood flow in the cat. The scientific world journal, 2001, 1: 168-180. doi: 10.1100/tsw.2001.23.

Furuya K, Milchak R J, Schegg K M, et al. Cockroach diuretic hormones: characterization of a calcitonin-like peptide in insects. Proceedings of the National Academy of Sciences, 2000, 97(12): 6469-6474. doi: 10.1073/pnas.97.12.6469.

Garelja M L, Bower R L, Brimble M A, et al. Pharmacological characterisation of mouse calcitonin and calcitonin receptor□like receptors reveals differences compared with human receptors. British journal of pharmacology, 2022, 179(3): 416-434. doi: 10.1111/bph.15628.

Goda T, Umezaki Y, Alwattari F, et al. Neuropeptides PDF and DH31 hierarchically regulate free-running rhythmicity in Drosophila circadian locomotor activity. Scientific reports, 2019, 9(1): 838. doi:10.1038/s41598-018-37107-3.

Golpon H A, Puechner A, Welte T, et al. Vasorelaxant effect of glucagon-like peptide-(7-36) amide and amylin on the pulmonary circulation of the rat. Regulatory peptides, 2001, 102(2-3): 81-86. doi: 10.1016/s0167-0115(01)00300-7.

Guo K, Su L, Wang Y, et al. Antioxidant and anti-aging effects of a sea cucumber protein hydrolyzate and bioinformatic characterization of its composing peptides. Food & Function, 2020, 11(6): 5004-5016. doi: 10.1039/d0fo00560f.

Hay D L, Conner A C, Howitt S G, et al. The pharmacology of adrenomedullin receptors and their relationship to CGRP receptors. Journal of Molecular Neuroscience, 2004, 22: 105-113. doi: 10.1385/JMN:22:1-2:105.

Hay D L, Garelja M L, Poyner D R, et al. Update on the pharmacology of calcitonin/CGRP family of peptides: IUPHAR Review 25. British journal of pharmacology, 2018, 175(1): 3-17. doi: 10.1111/bph.14075.

Huang D, Zhang B, Han T, et al. Genome-wide prediction and comparative transcriptomic analysis reveals the G protein-coupled receptors involved in gonadal development of *Apostichopus japonicus*. Genomics, 2021, 113(1): 967-978. doi: 10.1016/j.ygeno.2020.10.030.

Jékely G. Global view of the evolution and diversity of metazoan neuropeptide signaling. Proceedings of the National Academy of Sciences, 2013, 110(21): 8702-8707. doi: 10.1073/pnas.1221833110.

Johnson E C, Shafer O T, Trigg J S, et al. A novel diuretic hormone receptor in Drosophila: evidence for conservation of CGRP signaling. Journal of Experimental Biology, 2005, 208(7): 1239-1246. doi: 10.1242/jeb.01529.

Kandilci H B, Gumusel B, Lippton H. Intermedin/adrenomedullin-2 (IMD/AM2) relaxes rat main pulmonary arterial rings via cGMP-dependent pathway: role of nitric oxide and large conductance calcium-activated potassium channels (BKCa). Peptides, 2008, 29(8): 1321-1328. doi: 10.1016/j.peptides.2008.04.008.

Katafuchi T, Yasue H, Osaki T, et al. Calcitonin receptor-stimulating peptide: Its evolutionary and functional relationship with calcitonin/calcitonin gene-related peptide based on gene structure. Peptides, 2009, 30(9): 1753-1762. doi: 10.1016/j.peptides.2009.06.012.

Kuwasako K, Kitamura K, Nagata S, et al. Shared and separate functions of the RAMP-based
adrenomedullin receptors. Peptides, 2011, 32(7): 1540-1550. doi:
10.1016/j.peptides.2011.05.022.

Li C, Zheng Y, Cong X, et al. Molecular and functional characterization of the luqin-type
neuropeptide signaling system in the sea cucumber *Apostichopus japonicus*. Peptides, 2022,
155: 170839. doi: 10.1016/j.peptides.2022.170839.

Li S, Zhou Z, Dong Y, et al. Molecular characterization, expression analysis of the myostatin gene
and its association with growth traits in sea cucumber (*Apostichopus japonicus*). Comparative
Biochemistry and Physiology Part B: Biochemistry and Molecular Biology, 2016, 201: 12-20.
doi: 10.1016/j.cbpb.2016.06.005.

Martins R, Vieira F A, Power D M. Calcitonin receptor family evolution and fishing for function
using in silico promoter analysis. General and Comparative Endocrinology, 2014, 209: 61-73.
doi: 10.1016/j.ygcen.2014.04.025.

Mathiesen D S, Lund A, Vilsbøll T, et al. Amylin and calcitonin: potential therapeutic strategies to
reduce body weight and liver fat. Frontiers in Endocrinology, 2021, 11: 617400. doi:
10.3389/fendo.2020.617400.

Mayorova T D, Tian S, Cai W, Semmens D C. Odekunle E A, Zandawala M, Badi Y, Rowe M L,
Egertová M, Elphick M R. (2016). Localization of neuropeptide gene expression in larvae of
an echinoderm, the starfish *Asterias rubens*. Front. Neurosci. 10:553. doi:
10.3389/fnins.2016.00553.

McLatchie L M, Fraser N J, Main M J, et al. RAMPs regulate the transport and ligand specificity
of the calcitonin-receptor-like receptor. Nature, 1998, 393(6683): 333-339. doi:
10.1038/30666.

Mendel H C, Kaas Q, Muttenthaler M. Neuropeptide signalling systems–an underexplored target
for venom drug discovery. Biochemical pharmacology, 2020, 181: 114129. doi:
10.1016/j.bcp.2020.114129.

Mertens I, Vandingenen A, Johnson E C, et al. PDF receptor signaling in Drosophila contributes to
both circadian and geotactic behaviors. Neuron, 2005, 48(2): 213-219. doi:
10.1016/j.neuron.2005.09.009.

Mirabeau O, Joly J S. Molecular evolution of peptidergic signaling systems in bilaterians.
Proceedings of the national academy of sciences, 2013, 110(22): E2028-E2037. doi:
10.1073/pnas.1219956110.

Morris H R, Panico M, Etienne T, et al. Isolation and characterization of human calcitonin gene-
related peptide. Nature, 1984, 308(5961): 746-748. doi: 10.1038/308746a0.

Møller L N, Stidsen C E, Hartmann B, et al. Somatostatin receptors. Biochimica et Biophysica
Acta (BBA)-Biomembranes, 2003, 1616(1): 1-84. doi: 10.1016/s0165-6147(97)89535-x.

Nakamura M, Nomura S, Yamakawa T, et al. Endogenous calcitonin regulates lipid and glucose
metabolism in diet-induced obesity mice. Scientific Reports, 2018, 8(1): 17001. Endogenous
calcitonin regulates lipid and glucose metabolism in diet-induced obesity mice

Pangestuti R, Arifin Z. Medicinal and health benefit effects of functional sea cucumbers. Journal
of   traditional   and   complementary   medicine,   2018,   8(3):   341-351.   doi:
10.1016/j.jtcme.2017.06.007.

Párrizas M, LeRoith D. Insulin-like growth factor-1 inhibition of apoptosis is associated with
increased expression of the bcl-xL gene product. Endocrinology, 1997, 138(3): 1355-1358.
doi: 10.1210/endo.138.3.5103.

Perlow M J, Freed W J, Carman J S, et al. Calcitonin reduces feeding in man, monkey and rat.
Pharmacology Biochemistry and Behavior, 1980, 12(4): 609-612. doi: 10.1016/0091-
3057(80)90196-3.

Pinto A, Sekizawa K, Yamaya M, et al. Effects of adrenomedullin and calcitonin gene‐related
peptide on airway and pulmonary vascular smooth muscle in guinea‐pigs. British journal of
pharmacology, 1996, 119(7): 1477-1483. doi: 10.1111/j.1476-5381.1996.tb16061.x.

Pondel M. Calcitonin and calcitonin receptors: bone and beyond. International journal of
experimental pathology, 2000, 81(6): 405-422. doi: 10.1046/j.1365-2613.2000.00176.x.

Purdue B W, Tilakaratne N, Sexton P M. Molecular pharmacology of the calcitonin receptor.
Receptors and Channels, 2002, 8(3-4): 243-255. doi: 10.3109/10606820213681

Rowe M L, Achhala S, Elphick M R. Neuropeptides and polypeptide hormones in echinoderms:
new insights from analysis of the transcriptome of the sea cucumber *Apostichopus japonicus*.
General   and   Comparative   Endocrinology,   2014,   197:   43-55.   doi:
10.1016/j.ygcen.2013.12.002.

Rowe M L, Elphick M R. The neuropeptide transcriptome of a model echinoderm, the sea urchin
Strongylocentrotus purpuratus. General and comparative endocrinology, 2012, 179(3): 331-
344. doi: 10.1016/j.ygcen.2012.09.009.

Schwartz J, Réalis-Doyelle E, Dubos M P, et al. Characterization of an evolutionarily conserved
calcitonin signaling system in a lophotrochozoan, the Pacific oyster (*Crassostrea gigas*).
Journal of Experimental Biology, 2019, 222(13): jeb201319. doi: 10.1242/jeb.201319.

Sekiguchi T. Evolution of calcitonin/calcitonin gene-related peptide family in chordates:
identification of CT/CGRP family peptides in cartilaginous fish genome. General and
Comparative Endocrinology, 2022, 328: 114123. doi: 10.1016/j.ygcen.2022.114123.

Sekiguchi T, Kuwasako K, Ogasawara M, et al. Evidence for conservation of the calcitonin
superfamily and activity-regulating mechanisms in the basal chordate *Branchiostoma*
floridae: insights into the molecular and functional evolution in chordates. Journal of
Biological Chemistry, 2016, 291(5): 2345-2356. doi: 10.1074/jbc.M115.664003.

Sekiguchi T, Suzuki N, Fujiwara N, et al. Calcitonin in a protochordate, *Ciona intestinalis*–the prototype of the vertebrate calcitonin/calcitonin gene□related peptide superfamily. The FEBS Journal, 2009, 276(16): 4437-4447. doi: 10.1111/j.1742-4658.2009.07151.x.

Semmens D C, Mirabeau O, Moghul I, Pancholi M R, Wurm Y, and Elphick M R (2016). Transcriptomic identification of starfish neuropeptide precursors yields new insights into neuropeptide evolution. Open Biol. 6:150224. doi: 10.1098/rsob.150224.

Shafer O T, Kim D J, Dunbar-Yaffe R, et al. Widespread receptivity to neuropeptide PDF throughout the neuronal circadian clock network of Drosophila revealed by real-time cyclic AMP imaging. Neuron, 2008, 58(2): 223-237. doi: 10.1016/j.neuron.2008.02.018.

Smith M K, Wang T, Suwansa-Ard S, et al. The neuropeptidome of the Crown-of-Thorns Starfish, *Acanthaster planci*. Journal of proteomics, 2017, 165: 61-68. doi: 10.1016/j.jprot.2017.05.026.

Steiner D F, Ohagi S, Nagamatsu S, et al. Is islet amyloid polypeptide a significant factor in pathogenesis or pathophysiology of diabetes?. Diabetes, 1991, 40(3): 305-309. doi: 10.2337/diab.40.3.305.

Suwansa-Ard S, Chaiyamoon A, Talarovicova A, et al. Transcriptomic discovery and comparative analysis of neuropeptide precursors in sea cucumbers (Holothuroidea). Peptides, 2018, 99: 231-240. doi: 10.1016/j.peptides.2017.10.008.

Toribio R E, Kohn C W, Leone G W, et al. Molecular cloning and expression of equine calcitonin, calcitonin gene-related peptide-I, and calcitonin gene-related peptide-II. Molecular and cellular endocrinology, 2003, 199(1-2): 119-128. doi: 10.1016/s0303-7207(02)00289-7.

Tsurumaki T, Muraoka O, Yamaguchi T, et al. Neuropeptide Y-induced contraction and its desensitization through the neuropeptide Y receptor subtype in several rat veins. Journal of Cardiovascular Pharmacology, 2003, 41: S23-S27.

Walker C S, Conner A C, Poyner D R, et al. Regulation of signal transduction by calcitonin gene-related peptide receptors. Trends in pharmacological sciences, 2010, 31(10): 476-483. doi: 10.1016/j.tips.2010.06.006.

Wang Y, Yang Y, Li Y, et al. Identification of sex determination locus in sea cucumber *Apostichopus japonicus* using genome-wide association study. BMC genomics, 2022, 23(1): 391. doi: 10.1186/s12864-022-08632-3.

Wang Z, Cui J, Song J, et al. Comparative transcriptome analysis reveals growth-related genes in juvenile Chinese sea cucumber, Russian sea cucumber, and their hybrids. Marine biotechnology, 2018, 20: 193-205. doi: 10.1007/s10126-018-9796-6.

Wong A H C, Buckle C E, Van Tol H H M. Polymorphisms in dopamine receptors: what do they tell us? European journal of pharmacology, 2000, 410(2-3): 183-203. doi: 10.1016/s0014-2999(00)00815-3.

Yoshimoto R, Mitsui☐Saito M, Ozaki H, et al. Effects of adrenomedullin and calcitonin
gene☐related peptide on contractions of the rat aorta and porcine coronary artery. British
journal of pharmacology, 1998, 123(8): 1645-1654. doi: 10.1038/sj.bjp.0701805.

Zandawala M. Calcitonin-like diuretic hormones in insects. Insect biochemistry and molecular
biology, 2012, 42(10): 816-825. doi: 10.1016/j.ibmb.2012.06.006.

Zandawala M, Moghul I, Yañez Guerra L A, et al. Discovery of novel representatives of bilaterian
neuropeptide families and reconstruction of neuropeptide precursor evolution in ophiuroid
echinoderms. Open biology, 2017, 7(9): 170129. doi: 10.1098/rsob.170129.

Zhao Y, Chen M, Wang T, et al. Selection of reference genes for qRT-PCR analysis of gene
expression in sea cucumber *Apostichopus japonicus* during aestivation. Chinese Journal of
Oceanology and Limnology, 2014, 32(6): 1248-1256. doi: 10.1007/s00343-015-4004-2

Zheng Y, Cong X, Liu H, et al. Nervous system development and neuropeptides characterization
in embryo and larva: insights from a non-chordate deuterostome, the sea cucumber
Apostichopus japonicus. Biology, 2022, 11(10): 1538. doi: 10.3390/biology11101538.

Zheng Y, Liu H, Cong X, et al. A potential feeding regulation strategy during aestivation:
Relaxation of intestine mediated by pedal peptide/orcokinin-type neuropeptides in sea
cucumber *Apostichopus japonicus*. Aquaculture, 2024, 579: 740150.
doi:10.1016/j.aquaculture.2023.740150.

Supplementary figures

1166

1167

Figure 1-figure supplement 1. Calcitonin-type (CT-type) neuropeptide precursors in *A. japonicus*.
(A) *AjCTP1* comprises two putative CT-type neuropeptides, AjCT1 and AjCT2, whereas *AjCTP1*
generated by alternative splicing comprises only AjCT2. The predicted signal peptide is shown in
blue, the predicted cleavage sites (KR or KK) are shown in green, CT-type neuropeptides are
shown in red and C-terminal glycines that are potential substrates for amidation are shown in
orange. Predicted disulfide bridges in AjCT1 and AjCT2 are formed between the two underlined
cysteines. (B) Nucleotide sequences encoding CT-type neuropeptide precursors and the deduced
amino acid sequences in *A. japonicus*. 3'UTR and 5'UTR are shown in lowercase and the ORF is
shown in uppercase. Blue uppercase labels the predicted signal peptide, the green uppercase
shows the predicted cleavage site, AjCT1 and AjCT2 are shown in red and the putative substrate
for amidation is shown in orange.

1179

1180 Figure 1-figure supplement 2. Comparison of the sequences CT-type neuropeptides from Bilateria:

1181 deuterostomes-ambulacraria (green), deuterostomes-chordates (purple) and protostomes (pink).

1182 The conserved residues are shown with white lettering highlighted in gray and black. The

1183 accession numbers or citations are shown in the Supplementary table 1.

1184

A

AjCTR

(Nucleotide and deduced amino acid sequence of AjCTR)

B

AjPDFR1

(Nucleotide and deduced amino acid sequence of AjPDFR1)

1185

1186

1187
1188 Figure 1-figure supplement 3. Amino acid sequences and phosphorylation sites of AjCTR,
1189 AjPDFR1 and AjPDFR2 in *A. japonicus*. (A), (B) and (C) show the cDNA sequences encoding the
1190 receptors and their deduced amino acid sequences. ORFs are shown in uppercase; 3'UTR and
1191 5'UTR are shown in lowercase. The stop codon is shown by the asterisk (*). Predicted potential
1192 phosphorylation sites are shown with squares.

1193
Figure 1-figure supplement 4. Characterization of AjCT1 and AjCT2 receptors (AjCTR, AjPDFR1
and AjPDFR2). (A) Conserved domains of CTR/CLR-type protein sequences. Black line
represents the length of proteins and different domains are represented by rectangles. (B)
Conserved domains of PDFRs in Bilateria. Black line represents the length of proteins and
different domains are represented by rectangles. (C) Synteny analysis of AjCT1 and AjCT2
receptors (AjCTR, AjPDFR1 and AjPDFR2). The chromosomal locations of AjCT1 and AjCT2
receptor genes from *A. rubens*, *L. variegatus* and *A. japonicus* are shown, with different colors
used to mark neighboring genes and with their transcriptional direction shown by arrows.
GOLGA4, golgin subfamily A member 4; *NeuroD*, neurogenic differentiation factor; *CRF2*,
corticotropin-releasing factor receptor 2; *CTR*, calcitonin type receptor; *CLR*, calcitonin gene-
related peptide type 1 receptor; *ppGalNAc-T*, polypeptide N-acetylgalactosaminyltransferase;
URB3, E3 ubiquitin-protein ligase ubr3; *SLC40A1*, solute carrier family 40 member 1; *CMAH*,
cytidine monophosphate-N-acetylneuraminic acid hydroxylase; *MAGP*, microfibril-associated
glycoprotein; *HCN*, hyperpolarization-gated and cyclic nucleotide regulated channel; *ATG9A*,
autophagy-related protein 9A-like; *lysRS*, lysine--tRNA ligase; *FCN*, ficolin; *PDFR1*, pigment-
dispersing factor receptor 1; *PDFR2*, pigment-dispersing factor receptor 2; *PSMD8*, 26S
proteasome regulatory subunit 8; *SARNP*, SAP domain-containing ribonucleoprotein; *HPM1*,
histidine protein methyltransferase 1; *PNPO*, pyridoxine-5'-phosphate oxidase.
1212
1213

1214

1215 Figure 3-figure supplement 1. Negative control for analysis of AjCTR, AjPDFR1 and AjPDFR2

1216 internalization. The images show the localization of AjCTR, AjPDFR1 and AjPDFR2 in

1217 HEK293T cells without AjCT1 or AjCT2 stimulation. Green fluorescence represents the

1218 localization of pEGFP-N1/receptors in cell membrane. Cell nucleus probe (DAPI) was applied for

1219 cell nucleus staining. All pictures are representative of three independent experiments.

1220

Figure 4-figure supplement 1. Effects of PKA or PKC inhibitor on AjCT1 or AjCT2 stimulated ERK1/2 phosphorylation in AjCTR, AjPDFR1 or AjPDFR2 expressing HEK293T cells. (A) ERK1/2 phosphorylation, activated by AjCT1, is blocked by PKC inhibitor Gö 6983 in AjCTR expressing HEK293T cells and is blocked by PKA inhibitor H89 and PKC inhibitor in AjPDFR1 expressing HEK293T cells. (B) ERK1/2 phosphorylation, activated by AjCT2, is blocked by PKC inhibitor in AjCTR expressing HEK293T cells and is blocked by PKA inhibitor and PKC inhibitor in AjPDFR1 and AjPDFR2 expressing HEK293T cells. Serum-starved HEK293 cells were pre-treated with DMSO, PKA inhibitor, or PKC inhibitor before AjCT1 or AjCT2 stimulation. All images are representative of three independent experiments. Error bars represent SEM for three independent experiments. Statistically significant differences are indicated as follows: *: $p < 0.05$, ns: no significant.

1233

1234 Figure 6–figure supplement 1. Immunostaining in the radial nerve cord of *A. japonicus* is abolised

1235 by pre-absorption of ArCT antiserum with the ArCT antigen peptide. (A) Transverse section of a

1236 juvenile specimen of *A. japonicus* showing immunostaining in the radial nerve cord revealed by

1237 an antiserum to the starfish calcitonin-type neuropeptide ArCT, as also shown in Figure 6A. (B) A

1238 transverse section neighbouring the section shown in A was incubated with ArCT antiserum that

1239 had been pre-absorbed with the ArCT antigen peptide; note the absence of immunostaining in the

1240 radial nerve cord, demonstrating the specificity of the immunostaining in A. Abbreviations: LM,

1241 longitudinal muscle; RNC, radial nerve cord. Scale is 100 µm.

1242

Figure 7-figure supplement 1. Positive control on experimental tissues and pharmacological
effects of 10^{-6} M AjCT2 on longitudinal muscle from *A. japonicus*. (A) shows a recording of the
relaxing effect of AjCT2 (10^{-6} M) on a longitudinal muscle preparation. (B) and (C) respectively
are recordings of the contracting effects of ACh on longitudinal muscle and intestine preparations,
showing that the effect of ACh persists for at least 15 minutes. All recordings shown are
representative of three independent experiments.

1249

Figure 8A-figure supplement 1. Measurement of the mass of remaining bait and excrement for
AjCT2-treated and control animals. (A) Remaining bait mass in different phases. (B) The mass of
excrement in different phases. The 24-day experiment was divided into six phases (I-VI), with
each phase comprising four days for statistical analysis. CO: control group (sterilized seawater);
CT2L: AjCT2 low concentration group (5×10^{-3} mg/mL); CT2H: AjCT2 high concentration group

(5×10^{-1} mg/mL). Values are means ± SEM (n = 3, representing three parallel experiments of each

group). Asterisks represent the significant difference as follows: *: $p < 0.05$, **: $p < 0.01$.

1257

1258 Figure 9-figure supplement 1. The relative expression level of *AjCTP1/2* in siNC, si*AjCTP1/2*-1
1259 and si*AjCTP1/2*-2. siNC: negative control group; si*AjCTP1/2*-1 and si*AjCTP1/2*-2: experimental
1260 groups targeting *AjCTP1/2*. Values are means ± SEM (n = 5). Asterisks represent the significant
1261 differences as follows: *: $p < 0.05$, **: $p < 0.01$.

1262

1263 **Supplementary tables**

1264 Supplementary table 1. Accession numbers or citations of calcitonin-type family peptides or
1265 precursors in Bilateria

Abbreviation	Species Names	Accession Numbers And/or Citations
H_sapCTP	*Homo sapiens*	GI:179819
H_sapCGRPP	*Homo sapiens*	NP_001029125.1
T_rubCTP	*Takifugu rubripes*	GenBank: CAC81278.1
T_rubCGRPP	*Takifugu rubripes*	GenBank: CAC81277.1
C_intCTP	*Ciona intestinalis*	GenBank: BAI63095.1
B_floCTP1	*Branchiostome floridae*	GI:260826569
B_floCTP2	*Branchiostome floridae*	GI:260826567
B_floCTP3	*Branchiostome floridae*	GI:260826573
S_kowCTP	*Saccoglossus kowalevskii*	GI:187217193
A_japCTP1	*Apostichopus japonicus*	GenBank: AWU78769.1
A_japCTP2	*Apostichopus japonicus*	GenBank: AWU78770.1
A_rubCTP	*Asterias rubens*	GenBank: KT601715.1
P_pecCTP	*Patiria pectinifera*	GenBank: AWX63418.1
A_plaCTP	*Acanthaster planci*	Smith et al. (2017)
S_purCTP	*Strongylocentrotus purpuratus*	GI:115767208
O_vicCTP1	*Ophionotus. victoriae*	GenBank: ASK86238.1
O_vicCTP2	*Ophionotus. victoriae*	GenBank: ASK86238.1
O_araCTP1	*Ophiopsila aranea*	Zandawala et al. (2017)
O_araCTP2	*Ophiopsila aranea*	
A_filCTP1	*Amphiura filiformis*	
A_filCTP2	*Amphiura filiformis*	
H_scaCTP1	*Holothuria scabra*	Suwansa-Ard et al. (2018)

H_scaCTP2	Holothuria scabra		
H_glaCTP1	Holothuria glaberrima		
H_glaCTP2	Holothuria glaberrima		
A_medCTP	Antedon mediterranea	Aleotti et al., (2022)	
An_japCTP	Anneissia japonica		
L_gigCTP	Lottia gigantea	GI:163526287	
D_melDH31P	Drosophila melanogaster	NP_523514.1	
L_migDH31P	Locusta migratoria	GenBank: AKN21237.1	
C_telCTP	Capitella teleta	GI: 161220966	
C.telDH31P	Capitella teleta	Rowe et al. (2014)	

1266

1267   Supplementary table 2. Accession numbers or citations of CTR/CLR, PDF receptors, VP/OT-type
1268   receptors and VIP receptors in Bilateria

Classification	Abbreviation	Species Names	Accession Numbers And/or Citations
CTR/CLR	B_floCTR	Branchiostoma floridae	XP_035687900.1
	S_purCTR	Strongylocentrotus purpuratus	XP_003725171.1
	A_japCTR	Apostichopus japonicus	PIK49604.1
	A_rubCTR	Asterias rubens	XP_033639437.1
	P_minCTR	Patiria miniata	XP_038079052.1
	A_plaCTR	Acanthaster planci	XP_022079821.1
	A_filCTR	Amphiura filiformis	XP_072028982.1
	L_varCTR	Lytechinus variegatus	XP_041478637.1
	H_sapCTR	Homo sapiens	NP_001158209
	H_sapCLR	Homo sapiens	NP_005786
	B_belCTR	Branchiostoma belcheri	XP_019645944.1
	P_spaCLR	Polyodon spathula	XP_041102166.1
	E_calCLR	Erpetoichthys calabaricus	XP_028672925.1
	S_fasCLR	Stegostoma fasciatum	XP_048390160.1
	X_couCLR	Xiphophorus couchianus	XP_027880277.1
	X_macCLR	Xiphophorus maculatus	XP_005798205.1
	An_japCTR	Anneissia japonica	XP_033104142.1
	B_terDH31R	Bombus terrestris	XP_003398998.1
	B_morDH31R	Bombyx mori	NP_001127732
	C_telCTR	Capitella teleta	ELU10120.1
	C_gigCTR	Crassostrea gigas	AJ551182
	C_gigCTR2	Crassostrea gigas	MK290745
	P_dumDH31R	Platynereis dumerilii	KP293952
	D_melDH31R	Drosophila melanogaster	NP_725278.1
Deuterostomes PDFR	A_japPDFR1	Apostichopus japonicus	PIK34007.1
	A_japPDFR2	Apostichopus japonicus	PIK49999.1
	A_plaPDFR1	Acanthaster planci	XP_022080055.1
	S_purPDFR1	Strongylocentrotus purpuratus	XP_030845952.1
	L_varPDFR1	Lytechinus variegatus	XP_041477899.1
	P_minPDFR1	Patiria miniata	XP_038077382.1
	A_rubPDFR1	Asterias rubens	XP_033639833.1
	S_purPDFR2	Strongylocentrotus purpuratus	XP_030845954.1
	L_varPDFR2	Lytechinus variegatus	XP_041478256.1
	A_plaPDFR2	Acanthaster planci	XP_022080156.1
	P_minPDFR2	Patiria miniata	XP_038077934.1
	A_rubPDFR2	Asterias rubens	XP_033640032.1

	An_japPDFR	Anneissia japonica	XP_033104115.1
	S_kowPDFR	Saccoglossus kowalevskii	XP_006825104.1
	A_fil PDFR1	Amphiura filiformis	XP_072029557.1
	A_fil PDFR2	Amphiura filiformis	XP_072029560.1
VP/OTR	A_japVP/OTR	Apostichopus japonicus	PIK60987
	A_rubVP/OTR	Asterias rubens	MK279533
	S_purVP/OTR	Strongylocentrotus purpuratus	XM_779280.4
VIPR	H_sapVIPR1	Homo sapiens	KAI4029133.1
	H_sapVIPR2	Homo sapiens	KAI4016600.1
	M_musVIPR1	Mus musculus	EDL09145.1
	M_musVIPR2	Mus musculus	AAI38573.1
	G_galVIPR1	Gallus gallus	NP_001090992.2
	G_galVIPR1	Gallus gallus	NP_001014970.1
	C_livVIPR1	Columba livia	PKK32249.1
	C_livVIPR2	Columba livia	PKK32065.1
	X_laeVIPR1	Xenopus laevis	XP_018122563.1
	X_laeVIPR2	Xenopus laevis	XP_041423294.1
	X_troVIPR1	Xenopus tropicalis	XP_031759879.1
	X_troVIPR2	Xenopus tropicalis	NP_001120616.1
	E_orbVIPR1	Emys orbicularis	XP_065254637.1
	E_orbVIPR2	Emys orbicularis	XP_065255130.1
	D_rerVIPR1	Danio rerio	AAI62979.1
	D_rerVIPR2	Danio rerio	AAI62944.1
	N_furVIPR1	Nothobranchius furzeri	KAF7213846.1
	N_furVIPR2	Nothobranchius furzeri	SBP44365.1
Protostomes PDFR	P_chiPDFR	Penaeus chinensis	XP_047472721.1
	O_eduPDF1R	Ostrea edulis	XP_048754972.1
	C_virPDFR	Crassostrea virginica	XP_022310010.1
	C_gigPDFR	Crassostrea gigas	XP_034317823.1
	E_sinPDFR	Eriocheir sinensis	XP_050724174.1
	P_triPDFR	Portunus trituberculatus	XP_045102386.1
	P_maxPDF1R	Pecten maximus	XP_033762294.1
	D_virPDFR	Drosophila virilis	XP_032288784.1
	A_aegPDFR	Aedes aegypti	XP_021711391.1
	Ae_albPDFR	Aedes albopictus	XP_029719191.1
	An_albPDFR	Anopheles albimanus	XP_035773075.1
	H_rufPDFR	Haliotis rufescens	XP_048242666.1
	A_ampPDF1R	Amphibalanus amphitrite	XP_043235105.1
	A_macPDF1R	Anopheles maculipalpis	XP_050069493.1
	A_aquPDFR	Anopheles aquasalis	XP_050085801.1
	A_darPDFR	Anopheles darlingi	XP_049535712.1
	D_novPDFR	Drosophila novamexicana	XP_030566780.1
	A_funPDFR	Anopheles funestus	XP_049279756.1

G_aegPDFR	Gigantopelta aegis	XP_041366859.1
P_monPDFR	Penaeus monodon	XP_037798712.1
M_yesPDFR	Mizuhopecten yessoensis	XP_021374228.1
C_secPDFR	Cryptotermes secundus	XP_023726525.1
A_stePDF1R	Anopheles stephensi	XP_035902082.1
B_glaPDFR	Biomphalaria glabrata	KAI8755269.1
O_bimPDFR	Octopus bimaculoides	XP_014783031.1
A_merPDFR	Anopheles merus	XP_041761546.1
A_araPDFR	Anopheles arabiensis	XP_040175433.1
T_prePDFR	Trichogramma pretiosum	XP_014225639.1
C_quiPDF1R	Culex quinquefasciatus	XP_038122111.1
E_marPDF1R	Elysia marginata	GFS10866.1
E_rowPDFR	Euperipatoides rowelli	QKL20129.1
T_casPDFR	Tribolium castaneum	XP_015835556.1
B_kinPDFR	Belonocnema kinseyi	XP_033223584.1
A_colPDFR	Anopheles coluzzii	XP_040220269.2
D_melPDFR	Drosophila melanogaster	AAT84083.1
S_picPDFR	Schistocerca piceifrons	XP_047108870.1
P_japPDFR	Penaeus japonicus	XP_042879729.1
P_maxPDF2R	Pecten maximus	XP_033762295.1
P_polPDFR	Pollicipes pollicipes	XP_037077774.1
A_ampPDF2R	Amphibalanus amphitrite	XP_043235103.1
E_marPDF2R	Elysia marginata	GFS10866.1
P_ocePDFR	Plakobranchus ocellatus	GFO33290.1
O_eduPDF2R	Ostrea edulis	XP_048755458.1
C_quiPDF2R	Culex quinquefasciatus	XP_038122111.1
H_exePDFR	Hypsibius exemplaris	OQV13188.1
P_vulPDFR	Patella vulgata	XP_050418767.1
A_stePDF2R	Anopheles stephensi	XP_035902080.1
A_macPDF2R	Anopheles maculipalpis	XP_050069491.1
P_claPDFR	Procambarus clarkii	XP_045622346.1
H_vitPDFR	Homalodisca vitripennis	XP_046669177.1
O_tauPDFR	Onthophagus taurus	XP_022919236.1
L_anaPDFR	Lingula anatina	XP_013413106.1
H_rubPDFR	Haliotis rubra	XP_046569917.1
C. elePDFR1a	Caenorhabditis elegans	NP_001021170.1
C. elePDFR1b	Caenorhabditis elegans	EF141317.1
C. elePDFR1c	Caenorhabditis elegans	EF141318
C_telPDFR	Capitella teleta	ELU10117.1

1269

A

B

Bilateria		Representative species	CT-type signaling				PDF-type signaling			CT acting as ligand for PDFR	CT effect on feeding/growth
			precursor	predicted peptides	CTR/CLR	RAMP	precursor	predicted peptides	PDFR		
Deuterostomia	Vertebrata	*H. sapiens*	▢	◯	✳		X	X	X		I
	Urochordata	*C. intestinalis*	▢	▢	▢	X	X	X	X		
	Cephalochordata	*B. floridae*	▢	▢	✳	▢	X	X	X		
	Hemichordata	*S. kowalevskii*	▢	◯	▢	▢	?	?	▢		
	Echinodermata	*A. japonicus*	▢	●	✳	▢	?	?	▢	✳	P
		S. purpuratus	▢	▢	▢	▢	?	?	▢		
		A. rubens	▢	▢	▢	▢	?	?	▢		
		A. filiformis	▢	●	▢	▢	?	?	▢		
		A. japonica	▢	▢	▢	▢	?	?	▢		
Protostomia	Annelida	*C. teleta*	▢	▢	▢	▢	▢	▢	▢		
	Mollusca	*C. gigas*	▢	▢	✳	▢	▢	▢	▢		N
	Arthropoda	*D. melanogaster*	▢	▢	✳	▢	▢	▢	✳	✳	
	Nematoda	*C. elegans*	▢	▢	▢	▢	▢	▢	✳		