Functional characterisation of neuropeptides that act as ligands for both

calcitonin-type and pigment-dispersing factor-type receptors in a deuterostome

Xiao Cong [1, 2, #], Huachen Liu [1, #], Lihua Liu [1], Nayeli Escudero Castelán [3],

Kite G.E. Jones [3], Michaela Egertová [3], Maurice R. Elphick [3, *] and Muyan Chen [1, *]

1   The Key Laboratory of Mariculture, Ministry of Education, Ocean University of

China, Qingdao, China

2   Yantai Institute of Coastal Zone Research, Chinese Academy of Sciences, Yantai,

China

3   School of Biological & Behavioural Sciences, Queen Mary University of London,

London, United Kingdom

*   Correspondence: m.r.elphick@qmul.ac.uk; chenmuyan@ouc.edu.cn;

#   These authors contributed equally to this work.

Abstract

The calcitonin (CT) family of related peptides exerts diverse physiological effects in mammals via two G-protein coupled receptors: CTR and the CTR-like receptor CLR. Phylogenetic analysis of CT-type signaling has revealed the presence of CT-type peptides and CTR/CLR-type proteins in both deuterostome and protostome invertebrates. Furthermore, experimental studies have demonstrated that in the protostome *Drosophila melanogaster* the CT-like peptide DH_{31} can act as a ligand for a CTR/CLR-type receptor and a pigment-dispersing factor (PDF) receptor. Here we investigated the signaling mechanisms and functions of CT-type neuropeptides in a deuterostome invertebrate, the sea cucumber *Apostichopus japonicus* (phylum Echinodermata). In *A. japonicus,* a single gene encodes two CT-type peptides (AjCT1 and AjCT2) and both peptides act as ligands for a CTR/CLR-type receptor (AjCTR) and two PDF-type receptors (AjPDFR1, AjPDFR2), but with differential activation of downstream cAMP/PKA, $G\alpha q/Ca^{2+}/PKC$ and ERK1/2 signaling pathways. AjCT1/AjCT2-encoding transcripts were detected in the central nervous system and a variety of organ systems and neuropeptide expression was visualized immunohistochemically using an antiserum to a starfish CT-type peptide (ArCT). *In vitro* pharmacological experiments demonstrated that AjCT1 and/or AjCT2 cause dose-dependent relaxation of longitudinal muscle and intestine preparations. Furthermore, *in vivo* pharmacological experiments, combined with gain- and loss-of-function experiments, revealed a potential physiological role for AjCT2/AjPDFR2 signaling in promoting feeding and growth in *A. japonicus*. To our knowledge, this is the first study to obtain evidence that CT-type peptides can act as ligands for both CTR/CLR-type and PDF-type receptors in a deuterostome. Moreover, it provides the first evidence for appetite-stimulating and growth-promoting effects of CT-type neuropeptides in bilaterians. Given the economic importance of *A. japonicus* as a foodstuff, the discovery of CT-type peptides as potential regulators of feeding and growth in this species may offer novel strategies for aquaculture applications.

43 **Graphical abstract**

44
A schematic showing proposed molecular mechanisms by which CT-type signaling regulates
feeding and growth in *A. japonicus*. AjCT1 and AjCT2 are represented by purple and yellow
circles, respectively. For cell signaling, black arrows indicate that this process is activated, whilst
the pink arrow indicates that ERK1/2 is activated in the AjCT1/AjPDFR1, AjCT2/AjPDFR1 and
AjCT2/AjPDFR2 signaling pathways.

Keywords: calcitonin; neuropeptide; echinoderm; sea cucumber; calcitonin receptor;
pigment-dispersing factor receptor; feeding; growth

1. Introduction

The peptide calcitonin (CT) was first identified as a hormone that regulates blood calcium levels and subsequently a family of CT-related neuropeptides was discovered in mammals, including calcitonin gene related peptides (CGRPs), amylin (AMY), adrenomedullin (AM), and CT receptor-stimulating peptide (CRSP) (Copp et al., 1962; Sekiguchi et al., 2009; Sekiguchi et al., 2016). Members of the CT-type family of peptides typically have a disulfide bond in their N-terminal region (Cai et al., 2018) and exert their physiological effects in mammals by binding to two related G protein-coupled receptors: CTR (CT receptor) and CLR (CTR-like receptor) (Martins et al., 2014). However, receptor activity-modifying proteins (RAMPs) acting as accessory proteins are key determinants of ligand selectivity for CTR and CLR (Mclatchie et al., 1998). Thus, formation of a complex with a RAMP is essential for CLR function, whereas RAMPs are not required for CTR ligand recognition (Kuwasako et al., 2011; Bomberger et al., 2012). Furthermore, binding of CT to CTR in the absence of RAMPs (Katafuchi et al., 2009) can activate signaling via several downstream pathways, including cAMP accumulation, Ca^{2+} mobilization, and ERK activation (Chen et al., 1998; Pondel, 2000; Walker et al., 2010; Andreassen et al., 2014).

Transcriptome/genome sequencing has enabled analysis of the phylogenetic distribution of CT-type signaling systems, indicating that the evolutionary origin of CT-type peptides and CTR/CLR-related receptors can be traced back to the urbilaterian common ancestor of deuterostomes and protostomes (Furuya et al., 2000; Johnson et al., 2005; Jékely, 2013; Mirabeau and Joly, 2013). Furthermore, in protostomes two different types of CT-related peptides have been identified. Firstly, peptides that are structurally similar to vertebrate CT-type peptides in having an N-terminal disulphide bridge and secondly, CT-like diuretic hormone 31 (DH_{31})-type peptides that lack an N-terminal disulphide bridge (Furuya et al., 2000; Conzelmann et al., 2013). Furthermore, phylogenetic analysis indicates that gene duplication in a common ancestor of the protostomes gave rise to genes encoding CT-type peptides with a disulphide bridge and DH_{31}-type peptides, which were then differentially retained in protostome taxa (Conzelmann et al., 2013; Cai et al., 2018).

Receptors for CT-type peptides with a disulphide bridge have been characterised in

a molluscan species, the oyster *Crassostrea gigas* (now known as *Magallana gigas*),

which has seven CTR/CLR-type receptors. Two of these receptors, named Cg-CT-R

and Cragi-CTR2, were shown to be specifically activated by the CT-type

neuropeptides Cragi-CT1b and Cragi-CT2, respectively, thereby mediating

intracellular Ca^{2+} mobilization and cAMP accumulation (Schwartz *et al.*, 2019).

CT-type peptides with a disulphide bridge are not present in the insect *Drosophila*

melanogaster. However, receptors that are activated by DH_{31} have been identified in

this species and, interestingly, DH_{31} can act as a ligand for two types of receptors.

Thus, DH_{31} acts as a ligand for a CTR/CLR-related protein (CG17415), but it also

acts as a ligand for a protein (CG13758) that is the receptor for pigment-dispersing

factor (PDF) (Johnson et al., 2005; Mertens et al., 2005; Shafer et al., 2008).

Furthermore, the functional significance of the existence of dual receptor signaling

pathways for DH_{31} in *Drosophila* has been investigated extensively (Johnson et al.,

2005; Goda et al., 2019).

Currently, relatively little is known about CT-type signaling systems in

deuterostome invertebrates. Analysis of transcriptome/genome sequence data has

enabled identification of genes/transcripts encoding CT-type peptides and candidate

CTR/CLR-type receptors for these peptides in several deuterostome invertebrate taxa,

including urochordates (e.g. *Ciona intestinalis*), cephalochordates (e.g.

Branchiostoma floridae), hemichordates (e.g. *Saccoglossus kowalevskii*) and

echinoderms (e.g. *Strongylocentrotus purpuratus*) (Burke et al., 2006; Sekiguchi et al.,

2009; Sekiguchi et al., 2016; Rowe and Elphick, 2012; Martins et al., 2014).

Furthermore, analysis of the expression and action of CT-type peptides identified in

the starfish *Asterias rubens* and *Patiria pectinifera* has revealed that they act as

muscle relaxants (Cai et al., 2018; Semmens et al., 2016). However, experimental

confirmation of ligand-receptor partners has thus far only been reported for the

cephalochordate *B. floridae*, where three CT-type peptides were shown to act as

ligands for a CTR/CLR-type receptor when it is co-expressed with RAMP-type

proteins (Sekiguchi et al., 2016). To gain further insights into the mechanisms of CT-

type signaling in invertebrate deuterostomes, the aim of this study was to functionally characterise CT-type peptides and their receptors in an echinoderm – the sea cucumber *Apostichopus japonicus*.

A. japonicus is an economically important species in Asian aquaculture (Pangestuti and Arifin, 2018; Guo et al., 2020) and a species representative of the phylum Echinodermata. As a phylum of deuterostome invertebrates, echinoderms form a sister clade to the chordates together with hemichordates and xenoturbellids (Bromham and Degnan, 1999; Bourlat et al., 2006; Dunn et al., 2008) and as such they occupy a key "intermediate" phylogenetic position relative to the protostome invertebrates and vertebrates (Zandawala et al., 2017), which make them attractive as experimental systems to address evolutionary questions. In our previous studies, we have established this species as a model system to obtain insights into the evolutionary history and comparative physiology of neuropeptide signaling systems in bilaterians (Chen et al., 2019; Li et al., 2022; Zheng et al., 2022; Cong et al., 2023; Zheng et al., 2024). A gene encoding two CT-type peptides (AjCT1 and AjCT2) has been identified in *A. japonicus* (Chen et al., 2019) and a candidate receptor (AjCTR) for these peptides has also been predicted (Huang et al., 2021). Furthermore, relevant to the discovery that the CT-like peptide DH_{31} can act as a ligand for both a CTR/CLR-type receptor and a PDF-type receptor in *Drosophila*, the sequences of two PDF-type receptors have been identified in *A. japonicus* (AjPDFR1 and AjPDFR2) (Huang et al., 2021).

The first aim of this study was to investigate if AjCT1 and/or AjCT2 act as ligands for AjCTR, AjPDFR1 and AjPDFR2 and to characterise the downstream signaling pathways that are activated when AjCT1 and/or AjCT2 bind to their receptor(s). The second aim was to investigate the physiological roles of AjCT1 and AjCT2 in *A. japonicus* by analysing the expression of the gene encoding these peptides, investigating CT-type neuropeptide expression using immunohistochemistry, examining the *in vitro* and *in vivo* pharmacological effects of these peptides and performing loss-of-function tests. The functional characterisation of CT-type signaling in *A. japonicus* reported in this study provides new insights into the evolution and

comparative physiology of neuropeptides and a basis for application of neuropeptide-

based strategies to improve aquaculture methods for this economically important

species.

2. Results

2.1 Characterization and phylogenetic analysis of AjCT1/2

The complete cDNA sequences of the CT-type neuropeptide precursors *AjCTP1*

and *AjCTP2* were confirmed. The coding sequence of *AjCTP2* is shorter than *AjCTP1*

(Figure 1A) but both transcripts encode precursor proteins that comprise the same

predicted 23-residue N-terminal signal peptide (Figure 1-figure supplement 1A).

AjCTP1 encodes both putative CT-type neuropeptides (AjCT1 and AjCT2), while

AjCTP2 only encodes AjCT2 (Figure 1A, Figure 1-figure supplement 1A and 1B).

Exon and intron structure analysis showed that *AjCTP1* comprises four exons, with

AjCT1 and AjCT2 encoded in the third and fourth exons, respectively. *AjCTP2*

comprises three exons, excluding the third exon that encodes AjCT1 (Figure 1A).

Phylogenetic analysis revealed that AjCTP1 and AjCTP2 are closely related to CT-

type precursors from other sea cucumbers (*Holothuria glaberrima* and *Holothuria*

scabra), and are positioned within a clade that also contains CT-type precursors from

other echinoderms (sea urchin *S. purpuratus*, starfish *A. rubens*). Furthermore, the

echinoderm CT-type precursors are positioned within a clade that also comprises CT-

type precursors from chordates, whilst DH_{31}/CT-type precursors from protostomes are

positioned in a separate clade (Figure 1B). Sequence alignment of CT-related

neuropeptides revealed that, with the exception of DH_{31}-type peptides, the N-terminal

region of CT-related peptides has a pair of conserved cysteine residues that are known

or predicted to form a disulfide bond. Another conserved feature is a C-terminal

proline residue, with the exception of CGRPs (Figure 1-figure supplement 2).

Accordingly, both AjCT1 and AjCT2 have these conserved characteristics.

2.2 Characterization and phylogenetic analysis of three candidate receptors for

AjCT1/2

The sequences of three candidate receptors for AjCT1 and AjCT2 were obtained

by RACE and phosphorylation sites were predicted, as shown in Figure 2-figure

supplement 1A, 1B, and 1C. Phylogenetic analysis revealed that CTR/CLR-related

receptors divided into three clades and an *A. japonicus* CTR/CLR-type receptor

(AjCTR) was positioned in a clade with other deuterostome CTR/CLR-type receptors,

including human CTR and CLR. The other two clades comprised protostome DH31-

type receptors, which include the *Drosophila* receptor that has been identified

experimentally as the receptor for DH31, and other protostome CTR/CLR-related

receptors (Cluster A), including two receptors in the mollusc *Magallana gigas* that

have been identified experimentally as receptors for CT-like peptides with a

disulphide bridge. The phylogenetic analysis also revealed that two *A. japonicus* PDF-

type receptors (AjPDFR1 and AjPDFR2) were positioned in a clade comprising PDF-

type receptors, with one branch containing PDF-type receptors from deuterostomes

(including AjPDFR1 and AjPDFR2) and another branch containing PDF-type

receptors from protostomes (including the *Drosophila* PDF receptor) (Figure 2).

Domain prediction of the candidate receptors showed that AjCTR had HRM

(Hormone receptor) and 7tm_GPCRs (seven-transmembrane G protein-coupled

receptor) superfamily domains (Figure 2-figure supplement 2A), whereas AjPDFR1

and AjPDFR2 only presented the 7tm_GPCRs superfamily domain (Figure 2-figure

supplement 2B). To further investigate an orthologous relationship of the candidate

receptors with homologous receptors in other echinoderms, we conducted a gene

synteny analysis in *A. japonicus*, *A. rubens* and *L. variegatus*. The neighboring genes

of *AjCTR* included *GOLGA4*, *NeuroD*, *CRF2*, *ppGalNAc-T*, *URB3*, *SLC40A1*, *CMAH*

and *MAGP*. The relative positions of these genes were different from those in *A.*

rubens and *L. variegatus* (Figure 2-figure supplement 2C). The same situation was

also observed for the synteny analysis of PDFRs.

Figure 1. CT-type neuropeptide precursors and candidate receptors for CT-type peptides in *A. japonicus.* (A) Alternative splicing schematic of *AjCTP1* and *AjCTP2*. Exons are represented by green rectangles, introns are represented by lines, AjCT1 and AjCT2 are represented by red and orange rectangles respectively, upstream and downstream non-coding regions are represented by gray rectangles. The numbers represent the length of exons or introns. (B) Phylogenetic analysis of AjCTP1 and AjCTP2 and CT-related neuropeptide precursors from other bilaterians. CT-type precursors from deuterostomes are shown in green (echinoderms) and purple (chordates), CT-type precursors from protostomes are shown in pink. Full species names and accession numbers are listed in Supplementary table 1.

Figure 2. Maximum-likelihood phylogenetic tree of bilaterian CT-type and PDF-type receptor families, including those found in *A. japonicus,* rooted with CRH-type receptors. The tree comprises distinct CT-type receptor and PDF-type receptor clades with bootstrap support >90. The CT-type receptor clade is subdivided into a deuterostome-specific clade (dCTR, yellow), which includes the *A. japonicus* CTR (AjCTR), and two protostome receptor clades. One comprises DH31 receptors (DH31R, orange), and the other comprises protostome-specific CTRs (p-CTR (Cluster A), green). The name 'Cluster A' is referenced for consistency with Cardoso et al., 2024. The PDF-type receptors are subdivided into protostome (p-PDFR, blue) and deuterostome (d-PDFR, purple) clades, with the latter including the two *A. japonicus* PDFRs (AjPDFR1, AjPDFR2). The stars represent branch support (bootstrap 1000 replicates, see key). The scale bar indicates amino acid substitutions per site. The *A. japonicus* receptors characterised in this paper are highlighted in red, and CT-type family receptors for which ligands have been identified experimentally in other published studies are indicated with an asterisk (Gorn et al., 1992; McLatchie et al., 1998; Johnson et al., 2005; Bauknecht & Jekely, 2015; Sekiguchi et al., 2016; Schwartz et al., 2019; Beets et al., 2023; Huang et al., 2024). Species names are abbreviated as follows: Acal (*Aplysia californica*), Afil (*Amphiura filiformis*), Ajap (*Apostichopus japonicus*), Amel (*Apis mellifera*), Anja (*Anneissia japonica*), Arub (*Asterias rubens*), Bflo (*Branchiostoma*

floridae), Bimp (*Bombus impatiens*), Cele (*Caenorhabditis elegans*), Cint (*Ciona intestinalis*),
Dmel (*Drosophila melanogaster*), Ggal (*Gallus gallus*), Ggec (*Gekko gecko*), Hexe (*Hypsibius*
exemplaris), Hsap (*Homo sapiens*), Llon (*Lineus longissimus*), Locu (*Lepisosteus oculatus*), Lpol
(*Limulus polyphemus*), Lter (*Lumbricus terrestris*), Mgig (*Magallana gigas*), Ofus (*Owenia*
fusiformis), Pcau (*Priapulus caudatus*), Pdum (*Platynereis dumerilii*), Pmax (*Pecten maximus*),
Pmet (*Paramacrobiotus metropolita*), Rsoc (*Rotaria socialis*), Scla (*Styela clava*), Skow
(*Saccoglossus kowalevskii*), Sles (*Sepioteuthis laessoniana*), Smed (*Schmidtea mediterranea*),
Spur (*Strongylocentrotus purpuratus*), Tcas (*Tribolium castaneum*), Xtro (*Xenopus tropicalis*).
Sequence accession numbers are listed in supplementary table 2.
236

2.3 AjCT1 and AjCT2 act as ligands for AjCTR, AjPDFR1 and AjPDFR2

To investigate if AjCT1 and/or AjCT2 act as ligands for AjCTR, AjPDFR1 and

AjPDFR2, here, two receptor assay methods were employed. As shown in Figure 3A,

both AjCT1 and AjCT2 (10^{-6} M) caused a significant increase in CRE luciferase

activity in HEK293T cells transfected with AjCTR, AjPDFR1 or AjPDFR2.

Furthermore, this effect was inhibited by pre-incubation of cells with the PKA

inhibitor H89. The intracellular accumulation of cAMP was also measured, and this

showed that AjCT1 and AjCT2 (10^{-9} – 10^{-5} M) caused a dose-dependent increase in

cAMP in HEK293T cells transfected with AjCTR, AjPDFR1 or AjPDFR2 (Figure 3B).

However, when exposed to AjCT1 or AjCT2, the maximum accumulation of

intracellular cAMP was higher in cells transfected with AjCTR than in cells

transfected with AjPDFR1 and AjPDFR2 (Figure 3B). Measurement of SRE

luciferase activity revealed that AjCT1 and AjCT2 (10^{-6} M) administration evoked an

increase in activity in HEK293T cells transfected with AjCTR and AjPDFR1 (Figure

4A). However, in HEK293T cells transfected with AjPDFR2, an increase in SRE

luciferase activity was only observed with AjCT2, not with AjCT1. Furthermore, as

shown in Figure 4A, AjCT1- and AjCT2-induced increases in SRE luciferase activity

were not observed in cells that had been pre-incubated with FR900359 (Gαq protein

inhibitor) and Gö 6983 (PKC inhibitor). As shown in Figure 4-figure supplement 1,

receptors localized in the cell membrane, but after exposure to AjCT1 and AjCT2 (10^{-6}

^{6} M) for 15 min, internalization of pEGFP-N1/*AjCTR*, pEGFP-N1/*AjPDFR1* and

pEGFP-N1/*AjPDFR2* was observed in transfected HEK293T cells (Figure 4B). Taken

together, all these findings indicate that AjCT1 and AjCT2 can act as ligands for the

three candidate receptors and then activate cAMP/PKA signaling and $G\alpha q/Ca^{2+}/PKC$

signaling in all combinations, with the exception of AjCT1/AjPDFR2 which only

activates the cAMP/PKA cascade.

To investigate whether the activated signaling pathways may exert downstream

effects via the ERK–MAPK pathway, the phosphorylation levels of ERK1/2 were

analysed in transfected HEK293T cells. pERK1/2 levels in cells transfected with

AjCTR or AjPDFR1, but not AjPDFR2, were significantly increased after a 5 min

incubation with AjCT1 (Figure 5A). Furthermore, this effect was blocked by the PKC

inhibitor in AjCTR-expressing cells, and this effect was blocked by both PKA and

PKC inhibitors in AjPDFR1-expressing cells (Figure 5-figure supplement 1A).

pERK1/2 levels in cells transfected with AjCTR or AjPDFR1 or AjPDFR2, were

significantly increased after a 5 min incubation with AjCT2 (Figure 5B). Moreover,

this effect persisted until 15 min after exposure to AjCT in both AjCTR-transfected

cells and AjPDFR1-transfected cells but not in AjPDFR2-transfected cells (Figure 5B,

Table 1). Furthermore, these effects were blocked by PKA and PKC inhibitors in

AjPDFR1 and AjPDFR2 expressing cells, whereas in AjCTR expressing cells,

peptide-induced pERK1/2 activity was only blocked by a PKC inhibitor (Figure 5-

figure supplement 1B). These results indicate that, with the exception of

AjCT1/AjPDFR2, both AjCT1 and AjCT2 can activate the ERK1/2 cascade by

binding to the candidate receptors.

Figure 3. Pharmacological characterisation of AjCT1 and AjCT2 as ligands for the *A. japonicus* receptors AjCTR, AjPDFR1 and AjPDFR2. (A) CRE-driven luciferase activity measured in HEK293T cells transfected with one of the three candidate receptors after exposure to AjCT1 or AjCT2 or neuropeptide + DMSO or neuropeptide + H89 (10 μM) or serum-free DMEM (CO, negative control). The neuropeptide concentration used here was 10^{-6} M. Mean values with standard deviations (n = 9) are shown. * indicates a statistically significant difference with $p <$ 0.05. (B) Measurement of cAMP accumulation after exposure to AjCT1 or AjCT2 ($10^{-9} - 10^{-5}$ M). No cAMP elevation is observed in control experiments where HEK293T cells were transfected with empty pcDNA 3.1(+) (black circle). The estimated EC_{50} of AjCT1 for AjCTR, AjPDFR1 and

AjPDFR2 were 2.04×10^{-7} M (95% CI: 1.06×10^{-7} -8.69×10^{-7} M), 9.12×10^{-7} M (95% CI: 5.56

$\times 10^{-7}$ -4.45 $\times 10^{-6}$ M) and 6.09 $\times 10^{-8}$ M (95% CI: 2.23 $\times 10^{-8}$ -2.84 $\times 10^{-7}$ M) respectively. The

estimated EC_{50} of AjCT2 for AjCTR, AjPDFR1 and AjPDFR2 were 1.50 $\times 10^{-8}$ M (95% CI: 1.03

$\times 10^{-9}$ -3.89 $\times 10^{-8}$ M), 1.44 $\times 10^{-6}$ M (95% CI: 1.04 $\times 10^{-6}$ -1.99 $\times 10^{-6}$ M) and 7.90 $\times 10^{-7}$ M (95%

CI: 5.73 $\times 10^{-7}$ -1.06 $\times 10^{-6}$ M) respectively. Error bars represent SEM for three independent

experiments.

Figure 4. SRE-driven luciferase activity and receptor internalization in cells transfected with
AjCTR, AjPDFR1 or AjPDFR2 and exposed to AjCT1 or AjCT2. (A) SRE-driven luciferase
activity measured after incubation with neuropeptide (10^{-6} M AjCT1 or AjCT2)/DMSO,
neuropeptide (10^{-6} M)/FR900359 (1uM) and neuropeptide (10^{-6} M)/Gö 6983 (1 μM). Mean values
with standard deviations (n = 9) are shown. * indicates a statistically significant difference with *p*

< 0.05. (B) Internalization of AjCTR, AjPDFR1 and AjPDFR2 after 15 min treatment with 10^{-6} M
AjCT1 or AjCT2. Green fluorescence represents the localization of pEGFP-N1/receptors in cells.
The cell nucleus probe (DAPI) was used for cell nucleus staining. Scale bar: 50 µm. All pictures
are representative of three independent experiments.

Figure 5. ERK1/2 activity in cells transfected with AjCTR, AjPDFR1 or AjPDFR2 and exposed to
AjCT1 or AjCT2. (A) and (B) show the immunoblot intensity of representative phosphorylation
bands. The transfected HEK293T cells were incubated with AjCT1 (10^{-6} M) or AjCT2 (10^{-6} M) for
0, 5, 15, 30, 45 and 60 min. The p-ERK1/2 was normalized based on t-ERK1/2. All images are
representative of three independent experiments. Error bars represent SEM for three independent
experiments. Statistically significant differences are indicated as follows: *: $p < 0.05$, **: $p < 0.01$.

<center>Table 1. Activation of signaling pathways</center>

Assays / Receivers	cAMP/PKA/ERK1/2		Gαq/Ca^{2+}/PKC/ERK1/2	
	AjCT1	AjCT2	AjCT1	AjCT2
AjCTR	×*	×*	√	√
AjPDFR1	√	√	√	√
AjPDFR2	×*	√	×	√

A tick indicates cAMP/PKA/ERK1/2 or Gαq/Ca^{2+}/PKC/ERK1/2 activation, the cross indicates no

activation, the cross with * indicates that the cAMP/PKA pathway is activated, while the ERK1/2

pathway is not activated.

2.4 Expression profile of AjCTP1/2 and its receptors in A. japonicus

qRT-PCR analysis revealed that *AjCTP1/2* had the highest expression level in the

circumoral nervous system (CNS) (Figure 6A). The expression levels of three

receptors differed from those of *AjCTP1/2*. *AjCTR* (Figure 6B) and *AjPDFR1* (Figure

6C) showed the highest expression level in CNS and longitudinal muscle, whereas

AjPDFR2 exhibited highest expression in the respiratory tree (Figure 6D).

326
327   Figure 6. The relative expression level of *AjCTP1/2* (A), *AjCTR* (B), *AjPDFR1* (C) and *AjPDFR2*
328   (D) in different tissues. The circumoral nervous system was selected as the control group. Values
329   are means ± SEM (*AjCTP1/2*, n = 6; *AjCTR*, *AjPDFR1* and *AjPDFR2*, n = 5). Asterisks represent
330   significant differences as follows: *: $p < 0.05$, **: $p < 0.01$, ***: $p < 0.001$. CNS, circumoral

331 nervous system; Gd, gonad; Ie, intestine; LM, longitudinal muscle; RT, respiratory tree.
332

To investigate the distribution of CT-type neuropeptides in *A. japonicus* using

immunohistochemistry we employed an antiserum to the starfish (*Asterias rubens*)

CT-type neuropeptide ArCT, which has been used previously for visualization of the

expression of ArCT in *A. rubens* (Cai et al., 2018) and a CT-type neuropeptide in the

feather star *Antedon mediterranea* (AmCT; Aleotti et al., 2022). The antigen peptide

that was used to generate the ArCT antiserum (KYNSPFGASGP-NH$_2$) shares a C-

terminal FGxxGP-NH$_2$ (where X is variable) motif with both AjCT1 (FGSAGP-NH$_2$)

and AjCT2 (FGSGGP-NH$_2$) and therefore antibodies to ArCT may cross-react with

AjCT1 and/or AjCT2. Accordingly, the ArCT antiserum revealed immunostaining in

the central nervous system (radial nerve cords and circumoral nerve ring) (Figure 7A–

F), which was abolished by pre-incubation of the antiserum with the ArCT antigen

peptide (Figure 7-figure supplement 1). More specifically, high magnification images

revealed immunostaining was localised in the ectoneural region, but not in the

hyponeural region, of the radial nerve cords, with bilaterally symmetrical

immunostained cells (neuronal somata) present in the ectoneural epithelium (Figure

7C). Likewise, immunostained cells were revealed in the ectoneural region of the

circumoral nerve ring (Figure 7D-F). Furthermore, immunostaining was also revealed

in tentacle nerves, which emanate from the circumoral nerve ring (Figure 7D-F), and

in several regions of the digestive system, including the pharynx/stomach (Figure 7G),

intestine (Figure 7H) and cloaca (Figure 7I). No immunostaining was observed in the

longitudinal muscles (Figure 7A, B).

Figure 7. Immunohistochemical localization of CT-type neuropeptide expression in *A. japonicus*
using antibodies to the starfish CT-type neuropeptide ArCT. A. Transverse section showing
immunostaining in the radial nerve cord. B. Longitudinal section showing immunostaining in the
radial nerve cord. C. High magnification image of a transverse section showing immunostained
cells (arrowheads) and immunostained fibers in the neuropile (asterisks) of the ectoneural region
of the radial nerve cord. Note the absence of immunostaining in the hyponeural region. D, E.
Longitudinal sections showing immunostaining in the circumoral nerve ring and in tentacle nerves
that emanate from the circumoral nerve ring. F. High magnification image of a transverse section
showing immunostained cells (arrowheads) in the circumoral nerve ring. G. Longitudinal section
showing immunostained fibers (asterisk) at the junction between the pharynx and the stomach. H.
Longitudinal section showing immunostained fibers (arrowheads) in the intestine. I. Longitudinal
section showing immunostained fibers (arrowheads) in the cloaca. Abbreviations: An, anus; BW,
body wall; Cl, cloaca; CONR, circumoral nerve ring; Ec, Ectoneural region of the radial nerve
cord; Hy, hyponeural region of the radial nerve cord; LM, longitudinal muscle; Lu, lumen; Mu,
mucosa; RNC, radial nerve cord; TN, tentacle nerve. Scale bar, as shown in panel I, is 100 µm in
A, B, D, E, I; 50 µm in F, G, H; 25 µm in C.

2.5 AjCT1 and AjCT2 causes dose-dependent relaxation of longitudinal muscle and

intestine preparations from A. japonicus

Both AjCT1 and AjCT2 caused dose-dependent relaxation of longitudinal

muscles from *A. japonicus* at 10^{-10}–10^{-6} M (Figure 8). The maximum relaxant effects

of AjCT2 (10^{-6} M) on longitudinal muscles are illustrated in Figure 8-figure

supplement 1A. At the highest concentration tested (10^{-6} M), AjCT1 and AjCT2

caused 30.07 ± 7.49% and 32.50 ± 14.49% reversal of acetylcholine (ACh)-induced

contraction, respectively (Figure 8A, 8B). AjCT1 had no effect on the contractile state

of intestine preparations, even at concentrations as high as 10^{-5} M (Figure 8C). Unlike

AjCT1, AjCT2 caused dose-dependent relaxation of intestine preparations, with 26.33

± 8.75% reversal of ACh-induced contraction observed at 10^{-5} M (Figure 8D).

Control experiments in which ACh-induced contraction was recorded without

addition of CT-type peptides showed that ACh-induced contraction of longitudinal

muscles and intestinal preparations persisted for ≥15 minutes, confirming that

relaxation effects observed could be attributed to addition of AjCT1 or AjCT2 (Figure

8-figure supplement 1B, 1C).

Figure 8. Pharmacological effects of AjCT1 and AjCT2 on longitudinal muscle and intestine preparations from *A. japonicus*. (A) and (B) show representative recordings of the relaxing effects of AjCT1 and AjCT2 on longitudinal muscle preparations. The graphs show the dose-dependent relaxing effects of AjCT1 and AjCT2 (10^{-10} – 10^{-6} M), calculated as the percentage reversal of the contracting effect of 10^{-6} M ACh. (C) and (D) show representative recordings of experiments in which AjCT1 (10^{-7} – 10^{-5} M) and AjCT2 (10^{-9} – 10^{-5} M) were tested on intestine preparations. AjCT1 had no effect, but AjCT2 caused relaxation and the graph shows the dose-dependent relaxing effect of AjCT2 (10^{-9} – 10^{-5} M), calculated as the percentage reversal of the contracting effect of 10^{-5} M ACh. All recordings shown are representative of three independent experiments.

Mean values ± SEM were determined from three preparations. Lowercase letters a, b and c above

399 columns indicate the statistical difference at $p < 0.05$, whilst there is no significant difference
400 between bc and c.

2.6 AjCT2 affects feeding and growth in A. japonicus

To investigate the effects of AjCT1 and AjCT2 *in vivo*, a long-term injection

experiment was carried out at concentrations of 5×10^{-3} mg/mL (low-dose group) and

5×10^{-1} mg/mL (high-dose group), with the injection volume calculated based on the

wet weight of sea cucumbers at 1 µL/g. As shown in Figure 9A, there were no

significant changes in weight gain rate (WGR) and specific growth rate (SGR) in the

AjCT1 low concentration group (CT1L), AjCT1 high concentration group (CT1H)

and AjCT2 low concentration group (CT2L) ($p \geq 0.05$), but in the AjCT2 high

concentration group (CT2H) there was a significant increase in WGR and SGR in

comparison with the control group (CO) ($p < 0.05$). Correspondingly, the mass of

feces increased and the residual bait mass decreased (Figure 9A-figure supplement 1).

Furthermore, the relative intestinal expression levels of several growth factors were

investigated and the results showed that *AjGDF-8*, a growth inhibitory factor (Li et al.,

2016), was significantly down-regulated in all the experimental groups. In contrast,

the relative expression levels of *AjIgf* and *AjMegf6,* growth promoting factors

(Párrizas and LeRoith, 1997; Deribe et al., 2009), were significantly up-regulated in

intestines (Figure 9B).

Subsequently, the relative intestinal expression levels of AjCT1 and AjCT2

receptors were further tested for the CT2H group. The results showed that the relative

expression levels of all three receptors were significantly up-regulated, with the

ranked fold change being *AjCTR > AjPDFR1 > AjPDFR2* (Figure 9C). Conversely,

the relative expression levels of the three receptor genes were significantly down-

regulated in *siAjCTP1/2*-1 treated animals (Figure 10A, 10B and 10C). The

interference rate for si*AjCTP1/2*-1 was 66 % (Figure 10-figure supplement 1).

In long-term AjCT2 loss-of-function studies, the si*AjCTP1/2*-1 group exhibited

significant reductions in WGR and SGR compared to the control group (siNC) ($p <$

0.05, Figure 11A). Correspondingly, except for phase I, the si*AjCTP1/2*-1 group

showed significantly increased remaining bait and decreased excrement across phases

II-VI (Figure 11B). Furthermore, the growth inhibitory factor *AjGDF-8* was

significantly up-regulated and the growth promoting factor *AjMegf6* was significantly

down-regulated in the si*AjCTP1/2*-1 group (Figure 11C).

Figure 9. *In vivo* pharmacological tests revealed that AjCT1 and AjCT2 affect feeding and growth-
related gene expression in *A. japonicus*. (A) Weight gain rate (WGR) and specific growth rate
(SGR) in experimental and control groups. Values are means ± SEM (n = 3, representing three
parallel experiments of each group). (B) The relative intestinal transcript expression levels of
AjGDF-8, *AjIgf* and *AjMegf6* in different groups after injection of AjCT1 or AjCT2 or sterilized
seawater (CO) *in A. japonicus*. Values are means ± SEM (n = 5 biological duplication). (C) The
relative intestinal expression levels of AjCT1/2 receptors (*AjCTR*, *AjPDFR1* and *AjPDFR2*) in
CT2H and CO groups after 24 days injection experiment. Values are means ± SEM (n = 5
biological duplication). CO: control group (sterilized seawater); CT1L/CT2L: AjCT1/AjCT2 low
concentration group (5×10^{-3} mg/mL); CT1H/CT2H: AjCT1/AjCT2 high concentration group (5
$\times 10^{-1}$ mg/mL). * indicates statistically significant differences with $p < 0.05$, ** indicates
statistically significant differences with $p < 0.01$.

Figure 10. The relative intestinal expression levels of AjCT1/2 receptors after *AjCTP1/2*-1
knockdown. (A), (B) and (C) show the relative expression levels of *AjCTR*, *AjPDFR1* and
AjPDFR2 respectively. Values are means ± SEM (n = 5). ** indicates statistically significant
differences with $p < 0.01$.

Figure 11. Effects of 24-day *AjCTP1/2*-1 knockdown *in vivo* on feeding and growth in *A. japonicus*. (A) Weight gain rate (WGR) and specific growth rate (SGR) in siNC and *siAjCTP1/2*-1 groups. Values are means ± SEM (n = 3, representing three parallel experiments for each group). (B) Calculation of the remaining bait and excrement mass in different phases. The 24-day experiment was divided into six phases (I-VI), with each phase comprising four days. Values are means ± SEM (n = 3, representing three parallel experiments for each group). (C) The relative intestinal transcript expression levels of *AjGDF-8*, *AjIgf* and *AjMegf6* in siNC and *siAjCTP1/2*-1 groups. Values are means ± SEM (n = 5 biological duplication). siNC: negative control group; si*AjCTP1/2*-1: effectively experimental groups targeting *AjCTP1/2*. Asterisks represent statistically significant differences as follows: *: $p < 0.05$, **: $p < 0.01$, ***: $p < 0.001$.

2.7 AjPDFR2 mediates CT-type signaling-induced feeding and growth promotion in A. japonicus

The differential activation of signaling pathways and associated physiological outcomes between AjCT2 and AjCT1 suggested that AjCT2 may promote feeding and growth via AjPDFR2-mediated mechanisms. To investigate this hypothesis, we performed long term knockdown experiments targeting AjPDFR2. As shown in Figure 12A, si*AjPDFR2-1* effectively knocked down the expression of *AjPDFR2*. In the si*AjPDFR2-1* group, both WGR and SGR were significantly decreased compared to siNC group (Figure 12B). Correspondingly, the remaining bait weight was significantly increased in si*AjPDFR2-1* group (except during phase I), and the weight of excrement was significantly decreased in phase V and VI (Figure 12C). Additionally, the growth promoting factor *AjIgf* was significantly down-regulated in the si*AjPDFR2-1* group (Figure 12D).

Figure 12. Effects of *AjPDFR2*-1 knockdown *in vivo* on feeding and growth in *A. japonicus*. (A)
The relative expression level of *AjPDFR2* in siNC, si*AjPDFR2*-1 and si*AjPDFR2*-2. (B) Weight
gain rate (WGR) and specific growth rate (SGR) in siNC and *siAjPDFR2*-1 groups. Values are
means ± SEM (n = 3, representing three parallel experiments for each group). (C) The remaining
bait and excrement weight in different phases. The 24-day experiment was divided into six phases
(I-VI), with each phase comprising four days. Values are means ± SEM (n = 3, representing three
parallel experiments of each group). (D) The relative intestinal transcript expression levels of
AjGDF-8, *AjIgf* and *AjMegf6* in siNC and *siAjPDFR2*-1 groups. Values are means ± SEM (n = 5
biological duplication). siNC: negative control group; siA*jPDFR2*-1 and siA*jPDFR2*-2:

experimental groups targeting A*jPDFR2*. Asterisks represent statistically significant differences as

follows: *: $p < 0.05$, **: $p < 0.01$, ***: $p < 0.001$. ns: not significant.

3. Discussion

Previous studies have revealed that the CT-type neuropeptide DH_{31} can act as a

ligand for a CTR/CLR-type receptor and a pigment-dispersing factor (PDF) receptor

in a protostome – the insect *D. melanogaster* (Mertens et al., 2005; Shafer et al., 2008;

Schwartz et al., 2019). Here we show that two CT-type peptides (AjCT1 and AjCT2)

act as ligands for a CTR/CLR-type receptor and two pigment-dispersing factor (PDF)

type receptors in a deuterostome – the sea cucumber *A. japonicus* (phylum

Echinodermata). This is an important finding because it indicates that the ability of

CT-type neuropeptides to act as ligands for both CTR/CLR-type receptors and PDF-

type receptors may be an evolutionarily ancient property that originated in a common

ancestor of the Bilateria. Alternatively, the ability of CT-type neuropeptides to act as

ligands for PDF-type receptors in *D. melanogaster* and *A. japonicus* may have

evolved independently. Further studies on a wider variety of both protostome (e.g.

molluscs, annelids) and deuterostome taxa (e.g. other echinoderms, hemichordates)

are needed to address this issue. Here, we also investigated the signal transduction

pathways that are activated by AjCT1 and AjCT2 when these peptides bind to their

receptors and then proceeded to investigate the expression and actions of these

peptides to gain insights into their physiological/behavioral roles in *A. japonicus*, as

discussed below.

3.1 Alternative RNA splicing gives rise to transcripts encoding one (AjCT2) or two

(AjCT1, AjCT2) CT-type neuropeptides in A. japonicus

In vertebrates, alternative RNA splicing gives rise to transcripts encoding CT or

CGRP (Amara et al., 1982; Morris et al., 1984). Analysis of transcriptome/genome

sequence data from *A. japonicus* has revealed that in this species alternative RNA

splicing gives rise to transcripts encoding one (AjCT2) or two (AjCT1, AjCT2) CT-

type neuropeptides (Rowe et al., 2014; Zandawala et al., 2017; Chen et al., 2019; this

study). A similar pattern of alternative RNA splicing has been reported for CT-type

precursors in other sea cucumbers (*H. scabra* and *H. glaberrima*; class Holothuroidea) and in brittle stars (*Amphiura filiformis* and *Ophiopsila Aranea*; class Ophiuroidea) (Zandawala et al., 2017; Suwansa-Ard et al., 2018). In contrast, CT-type precursors identified in other echinoderm classes (Asteroidea, Echinoidea and Crinoidea) comprise a single CT-type neuropeptide (Rowe and Elphick, 2012; Mirabeau and Joly, 2013; Rowe et al., 2014; Semmens et al., 2016; Cai et al., 2018; Aleotti et al., 2022). However, it remains to be determined if this feature of CT-type precursors in sea cucumbers and brittle stars evolved independently or originated in a common ancestor of eleutherozoan echinoderms but with subsequent loss in Asteroidea and Echinoidea (Figure 13).

Analysis of the sequences of AjCT1 and AjCT2 revealed the presence of two cysteine residues in their N-terminal regions, which is an evolutionarily conserved feature of CT-type peptides. Structural analysis of CT-type peptides in vertebrates and in starfish has revealed that these cysteines form a disulphide bridge in the mature peptides (Cai et al., 2018) and therefore it is inferred that this post-translational modification also occurs in AjCT1 and AjCT2. Furthermore, the C-terminal residue of AjCT1 and ACT2 is predicted to be an amidated proline, which is a conserved post-translational modification of CT/DH31-type peptides in many taxa, but not vertebrate CGRPs (Cai et al., 2018).

3.2 Identification and characterisation of receptors for CT-type neuropeptides in A. japonicus

Informed by previous genome-wide analysis of GPCRs in *A. japonicus* (Huang et al., 2021), we identified three candidate receptors for AjCT1 and/or AjCT2. Firstly, AjCTR, which is a homolog of CT/DH$_{31}$-type receptors that have been identified in other taxa, including vertebrates and insects (Garelja et al., 2022; Johnson et al., 2005; Toribio et al., 2003). Secondly, AjPDFR1 and AjPDF2, which are homologs of the PDF-type receptor that has been shown to be activated by DH$_{31}$ in *D. melanogaster* (Mertens et al., 2005; Zandawala, 2012).

In mammals, CTR and CLR mediate effects of all CT-related neuropeptides, but

their ligand specificity relies on the presence and types of receptor activity-modifying proteins (RAMPs) (McLatchie et al., 1998; Purdue et al., 2002; Conner et al., 2004; Hay et al., 2018). Thus, CTR acts as a functional receptor for CT-type peptides without an associated RAMP (Katafuchi et al., 2009), whereas activation of downstream signaling via CLR requires an associated RAMP (Hay et al., 2018). Here, we investigated the occurrence of RAMPs in *A. japonicus* using the tblastn method employed by Sekiguchi *et al.*, (2016), which enabled identification of RAMPs in the cephalochordate *B. floridae*. However, no RAMPs were identified in *A. japonicus*, suggesting that AjCTR may be functionally more similar to CTR than CLR. Accordingly, tblastn analysis of other available echinoderm genomes (*A. rubens*, *S. purpuratus* and *Anneissia japonica*), as well as transcriptome databases for brittle stars (*O. victoriae*, *O. aranea* and *A. filiformis*), also failed to identify RAMPs. Likewise, RAMPs have also not been found in protostomes (Schwartz et al., 2019). Therefore, RAMPs may be unique to chordates, although interestingly RAMPs have also not been identified in the urochordate *C. intestinalis* (Sekiguchi et al., 2009) (Figure 13).

As highlighted above, here we show that the CT-type peptides AjCT1 and AjCT2 both act as ligands for a CTR/CLR-type receptor (AjCTR) and two PDF-type receptors (AjPDFR1, AjPDFR2) in *A. japonicus*. Furthermore, characterization of downstream signaling mechanisms revealed that both AjCT1 and AjCT2 trigger AjCTR-mediated cAMP accumulation, Ca^{2+} mobilization, and ERK signaling. These findings are consistent with previous studies on CT-type receptors in vertebrates and in the mollusc *Magallana gigas* (Pondel, 2000; Walker et al., 2010; Andreassen et al., 2014; Schwartz et al., 2019). Both AjCT1 and AjCT2 trigger cAMP accumulation, Ca^{2+} mobilization, and ERK signaling mediated by AjPDFR1. Both AjCT1 and AjCT2 triggered cAMP accumulation mediated by AjPDFR2, but only AjCT2 (not AjCT1) triggered Ca^{2+} mobilization and ERK signaling mediated by AjPDFR2. These differences in the signal transduction mechanisms activated by AjCT1 and AjCT2 may account for differences in the pharmacological activities and physiological roles of these peptides, as discussed below.

3.3 AjCT1 and AjCT2 act as muscle relaxants in A. japonicus

Analysis of the expression of the gene encoding the precursors comprising
AjCT2 or AjCT1 and AjCT2 revealed the occurrence of transcripts in the circumoral
nervous system as well as in peripheral tissues/organs, including the longitudinal
muscle, digestive system and gonads. Immunohistochemical analysis using an
antiserum to the starfish (*A. rubens*) CT-type neuropeptide ArCT enabled visualisation
of CT-type neuropeptide(s) in juvenile specimens of *A. japonicus*. Intense and
extensive immunostaining was revealed in the ectoneural region of the radial nerve
cords and circumoral nerve ring, consistent with the high levels *AjCTP1/2* transcripts
detected in circumoral nervous tissue. Immunostained nerve fibers were also observed
in the digestive system (including the intestine), consistent with the detection
AjCTP1/2 transcripts in intestine. However, no immunostaining was observed in
longitudinal muscles, which is inconsistent with the detection of *AjCTP1/2* transcripts
this tissue. This may reflect differences in the detection sensitivity of the techniques
used. In accordance with the expression pattern of AjCTP1/2 in *A. japonicus*, analysis
of CT-type neuropeptide expression in other echinoderms has revealed a widespread
pattern of expression in larval and adult starfish (*A. rubens*) and in juvenile and adult
featherstars (*Antedon mediterranea*) (Mayorova et al., 2016; Cai et al., 2018; Aleotti
et al., 2022).

Several CT-related peptides act as muscle relaxants in vertebrates (Steiner et al.,
1991; Pinto et al., 1996; Yoshimoto et al., 1998; Edvinsson et al., 2001; Golpon et al.,
2001; Kandilci et al., 2008) and, accordingly, previous studies have revealed that CT-
type neuropeptides cause relaxation of muscle preparations in starfish, including
locomotory organs (tube feet) and the apical muscle, which is functionally similar to
the longitudinal muscle in sea cucumbers (Cai et al., 2018). Therefore, it was of
interest to investigate if AjCT1 and/or AjCT2 act as muscle relaxants in *A. japonicus*.
Both AjCT1 and AjCT2 caused dose-dependent relaxation of longitudinal muscle
preparations, consistent with the relaxing effect of CT-type peptides on starfish apical
muscle preparations. It is worth noting that while AjCT2 at 10^{-6} M exhibited the

maximum efficacy in reversing ACh-induced contraction, its relaxing effect weakened

compared to the $10\square\square$ M treatment when applied at progressively higher

concentrations. This diminished efficacy likely stems from receptor desensitization,

induced by repeated AjCT2 application to longitudinal muscle preparations

(Tsurumaki et al., 2003; Arrowsmith and Wray, 2014). Furthermore, AjCT2, but not

AjCT1, caused dose-dependent relaxation of intestine preparations from *A. japonicus*.

Interestingly, the CT-type peptide ArCT does not to cause relaxation of cardiac

stomach preparations from the starfish *A. rubens* (Cai et al., 2018). Therefore, from a

functional perspective ArCT is more like AjCT1 than AjCT2 and we speculate that

this may reflect loss of an exon encoding a AjCT2-like peptide in the Asteroidea

lineage. Further insights into this issue and the evolution of CT-type neuropeptide

function echinoderms could be obtained by investigation of the actions of CT-type

peptides in brittle stars and sea urchins. Differences in the effects of AjCT1 and

AjCT2 on the intestine may be explained by differences in their signal transduction

mechanisms. Thus, only AjCT2, not AjCT1, triggered AjPDFR2-mediated activation

of Ca^{2+} mobilization and ERK signaling and therefore we speculate that the relaxing

effect of AjCT2 on the intestine may be mediated by activation of AjPDFR2.

3.4 AjCT2/AjPDFR2 stimulates feeding and growth in A. japonicus

In vivo pharmacological assays demonstrated that only high concentrations of

AjCT2 significantly enhanced feeding and growth rates in *A. japonicus*. In contrast,

neither a low concentration of AjCT2 nor any concentration of AjCT1 (low or high)

induced detectable effects. Furthermore, long-term knockdown of *AjCTP1/2* further

validated the essential role of AjCT2 in regulating feeding and growth in this species.

To elucidate the receptor mediating AjCT2's feeding- and growth-promoting effects,

we selected AjPDFR2 based on its distinct activation profile: AjCT2 selectively

activated AjPDFR2, inducing downstream ERK1/2 phosphorylation, whereas AjCT1

exhibited no activity toward this receptor. Given this receptor specificity, we

performed *AjPDFR2* knockdown experiments, which revealed phenotypic changes

consistent with those in *AjCTP1/2* knockdown animals, including significantly

reduced WGR and SGR, alongside increased remaining bait accumulation and

diminished excrement output compared to control. Collectively, these results support

a model wherein AjCT2 promotes feeding and growth in *A. japonicus* via AjPDFR2-

dependent activation of the cAMP/PKA/ERK1/2 and $G\alpha q/Ca^2\square/PKC/ERK1/2$

cascades. Considering the inherent complexity of neuropeptide signaling systems,

which involve multiple GPCR subtypes coupled to diverse signaling cascades, ligands

bound to the same receptor may activate distinct G protein subforms within a single

cell (Møller et al., 2003; Mendel et al., 2020). Receptor activation modes may be

modulated by structural polymorphisms or binding site diversity (Wong et al., 2000;

Changeux, 2010), as well as by the differential efficacy of peptides in activating

receptors *in vivo*.

Collectively, these findings indicate that AjCT2 has a physiological role in

stimulating feeding and growth in *A. japonicus*. To the best of our knowledge, this is

the first study to report orexigenic and growth-promoting effects of a CT-type

neuropeptide in bilaterian animals. Interestingly, previous studies have reported the

expression of CT-type neuropeptides in the digestive system of invertebrate chordates

(Sekiguchi et al., 2009; Sekiguchi et al., 2016). However, in contrast to the

stimulatory effect of AjCT2 on feeding/growth in *A. japonicus*, CT-type neuropeptides

have been found to inhibit feeding and growth in vertebrates (Del et al., 2002; Hay et

al., 2018; Nakamura et al., 2018). Thus, it appears that opposing roles of CT-type

neuropeptide signaling systems as regulators for feeding/growth have evolved in

deuterostomes, with a stimulatory role reported here in an echinoderm and an

inhibitory role reported previously in vertebrates (Figure 13). Further research is now

needed to investigate if the stimulatory effect of AjCT2 on feeding/growth observed

here in *A. japonicus* also applies to other sea cucumber species. Moreover, our

discovery of this effect of AjCT2 on *A. japonicus* may provide a basis for

development of novel approaches to improve aquaculture of this economically

important species.

Figure 13. Phylogenetic summary of the occurrence and properties of CT-type and PDF-type signaling systems and effects of CT-type peptides on feeding/growth regulation in representative species. Echinoderms are boxed in purple with dashed lines and *A. japonicus*, the focus of this study, is framed in red with dashed lines. Pink filled rectangles indicate the occurrence of CT-type precursors/neuropeptides, CTR/CLR-type receptors and RAMPs. Green filled rectangles indicate the occurrence of PDF-type precursors/neuropeptides and receptors. Blue filled rectangles indicate species in which CT-type peptides have been shown to act as ligands for PDF-type receptors. Yellow filled rectangles indicate species in which effects or lack of effects of CT-type peptides on feeding/growth have been reported (I: Inhibition; P: Promotion; N: No effect). Red circles indicate that neuropeptides are generated by alternative splicing of precursor transcripts, with an unfilled circle indicating that transcripts encode one neuropeptide and with a filled circle indicating that transcripts can encode two neuropeptides in tandem. A yellow asterisk marks species in which neuropeptides have been shown to act as ligands for receptors experimentally. Unfilled rectangles indicate the absence of a protein. Unfilled rectanges containing a cross (X) indicate loss of a protein during evolution. Unfilled rectangles containing a question mark (?) indicate the existence of precursors/peptides that have yet to be identified.

685 **4. Materials and methods**

Key Resources Table				
Reagent type (species) or resource	Designation	Source or reference	Identifiers	Additional information
Cell line (*Homo sapiens*)	HEK293 cell line	Procell	CL-0005	
Antibody	phosphorylated ERK1/2 kinases (Thr202/Tyr204) (goat)	Cell Signaling Technology	CAT#4930	IF (1:2000)
Antibody	ERK1/2 kinases (137F5) (goat)	Cell Signaling Technology	CAT#4695T	IF (1:1000)
Antibody	HRP-conjugated goat anti-rabbit IgG (goat)	Absin	Abs20040ss	IF (1:2000)
sequence-based reagent	Primers for pcDNA3.1(+) plasmid construction--AjCTR	This paper	PCR PRIMER	FORWARD CGCGGATCCAAAATGCAGGAGAACGATTC BamH □
sequence-based reagent	Primers for pcDNA3.1(+) plasmid construction--AjCTR	This paper	PCR PRIMER	REVERSE CCGGAATTCTTAAACTACCGTTGTTTTAT EcoR □
sequence-based reagent	Primers for pcDNA3.1(+) plasmid construction—AjPDFR1	This paper	PCR PRIMER	FORWARD CGCGGATCCACAATGAGTGTCGGGAACGT BamH □
sequence-based reagent	Primers for pcDNA3.1(+) plasmid	This paper	PCR PRIMER	REVERSE CCGGAATTCTCATTTCCCACCCCACTCAC

	construction—AjPDFR1			EcoR □
sequence-based reagent	Primers for pcDNA3.1(+) plasmid construction—AjPDFR2	This paper	PCR PRIMER	FORWARD CGCGGATCCAAGATGGCGAACA ATGAAAC BamH □
sequence-based reagent	Primers for pcDNA3.1(+) plasmid construction—AjPDFR2	This paper	PCR PRIMER	REVERSE CCGGAATTCTCATTCATTAAATT TGCCGT EcoR □
sequence-based reagent	Plasmid construction AjCTR–EGFP	This paper	PCR PRIMER	FORWARD CCGGAATTCTGAAAATGCAGGA GAACGATTC EcoR □
sequence-based reagent	Plasmid construction AjCTR–EGFP	This paper	PCR PRIMER	REVERSE CGCGGATCCCGAACTACCGTTG TTTTATCTG BamH □
sequence-based reagent	Plasmid construction AjPDFR1–EGFP			CCGGAATTCTGACAATGAGTGT CGGGAACGT EcoR □
sequence-based reagent	Plasmid construction AjPDFR1–EGFP			CGCGGATCCCGTTTCCCACCCC ACTCACTGT BamH □
sequence-based reagent	Plasmid construction AjPDFR2–EGFP			CCGGAATTCTGAAGATGGCGAA CAATGAAAC EcoR □
sequence-based reagent	Plasmid construction AjPDFR2–EGFP			CGCGGATCCCGTTCATTAAATTT GCCGTTCA BamH □
sequence-based reagent	Primers for qPCR assays β-Actin	This paper	PCR PRIMER	FORWARD AAGGTTATGCTCTTCCTCACGCT
sequence-based reagent	Primers for qPCR assays β-Actin	This paper	PCR PRIMER	REVERSE GATGTCACGGACGATTTCACG

sequence-based reagent	Primers for qPCR assays β-Tubulin	This paper	PCR PRIMER	FORWARD GAAAGCCTTACGACGGAACA
sequence-based reagent	Primers for qPCR assays β-Tubulin	This paper	PCR PRIMER	REVERSE CACCACGTGGACTCAAAATG
sequence-based reagent	Primers for qPCR assays AjCTP1/2	This paper	PCR PRIMER	FORWARD GTTTGGCGGTACCCATTAC
sequence-based reagent	Primers for qPCR assays AjCTP1/2	This paper	PCR PRIMER	REVERSE GGCGTTCGGTCTGTTAATAG
sequence-based reagent	Primers for qPCR assays AjMegf6	This paper	PCR PRIMER	FORWARD CGGTCCTTACTGTGCGGAGA
sequence-based reagent	Primers for qPCR assays AjMegf6	This paper	PCR PRIMER	REVERSE CGTCCTCCCAGCTGGACATC
sequence-based reagent	Primers for qPCR assays AjIgf	This paper	PCR PRIMER	FORWARD TGGAAGGCAGTAAGGCTA
sequence-based reagent	Primers for qPCR assays AjIgf	This paper	PCR PRIMER	REVERSE TAGGCTGATGTGGTTGGA
sequence-based reagent	Primers for qPCR assays AjGDF-8	This paper	PCR PRIMER	FORWARD ATACGCCCATACTCACATCC
sequence-based reagent	Primers for qPCR assays AjGDF-8	This paper	PCR PRIMER	REVERSE TGGCAGGAGTCAACCTTCAT
sequence-based reagent	Primers for qPCR assays AjCTR	This paper	PCR PRIMER	FORWARD GACGAATGGAGGAGAAGG
sequence-based reagent	Primers for qPCR assays AjCTR	This paper	PCR PRIMER	REVERSE CGCCGCAATTAGAGTAGT

sequence-based reagent	Primers for qPCR assays *AjPDFR1*	This paper	PCR PRIMER	FORWARD TTATCAACCGCCGTCTTC
sequence-based reagent	Primers for qPCR assays *AjPDFR1*	This paper	PCR PRIMER	REVERSE CAGGAGGAGGAACACATTC
sequence-based reagent	Primers for qPCR assays *AjPDFR2*	This paper	PCR PRIMER	FORWARD TGTTCTCCTGCCTCTTCT
sequence-based reagent	Primers for qPCR assays *AjPDFR2*	This paper	PCR PRIMER	REVERSE GTCCTTCTCATCGGTGTG
sequence-based reagent	Primers for 3'RACE *AjCTP1* *AjCTP2*	This paper	PCR PRIMER	GATTACGCCAAGCTTCGAGCTG TGTCGATAACATTACGCATGG
sequence-based reagent	Primers for 3'RACE AjPDFR1	This paper	PCR PRIMER	GATTACGCCAAGCTTGAGGGAA AGCCAAACCAGCGAGACG
sequence-based reagent	Primers for 3'RACE AjPDFR2	This paper	PCR PRIMER	GATTACGCCAAGCTTGTGCAAC CGGACGAAGATCCACGTG
sequence-based reagent	Primers for 5'RACE AjPDFR1	This paper	PCR PRIMER	GATTACGCCAAGCTTCGTCTCG CTGGTTTGGCTTTCCCTC
sequence-based reagent	Primers for 5'RACE AjPDFR2	This paper	PCR PRIMER	GATTACGCCAAGCTTGCCTTCC ACAAACATCCATGCGAACGC
sequence-based reagent	Primers for 5'RACE AjCTR	This paper	PCR PRIMER	GATTACGCCAAGCTTACGCCGG GCTCTCAAGAAGTCATTC
sequence	RNAi	This paper	si*AjCTP1*-1	GGAAUUAUCAAGACAGUAU (dT)(dT) AUACUGUCUUGAUAAUUCC (dT)(dT)
sequence	RNAi	This paper	si*AjCTP1*-2	GAAAGUCGCUAAUGCAGUA (dT)(dT) UACUGCAUUAGCGACUUUC

				(dT)(dT)
sequence	RNAi	This paper	si*AjPDFR2*-1	GUAGCGAUUGGUCGAAUAA (dT)(dT) UUAUUCGACCAAUCGCUAC (dT)(dT)
sequence	RNAi	This paper	si*AjPDFR2*-2	CAGUCAUUGUUAUCUACCA (dT)(dT) UGGUAGAUAACAAUGACUG (dT)(dT)
sequence	RNAi	This paper	siNC	UUCUCCGAACGUGUCACGU (dT)(dT) ACGUGACACGUUCGGAGAA (dT)(dT)
commercial assay or kit	SMARTer RACE 5'/3' Kit	Takara	Cat No. 634858	
commercial assay or kit	FastPure Gel DNA Extraction Mini Kit	Vazyme	DC301-01	
commercial assay or kit	Hifair® □ 1st Strand cDNA Synthesis SuperMix	YEASEN	Cat No. 11141ES60	
commercial assay or kit	Hieff UNICON Universal Blue qPCR SYBR Green Master Mix	YEASEN	Cat No. 11184ES08	
commercial assay or kit	Firefly luciferase reporter gene assay kit	MK	MF4001	
commercial assay or kit	cAMP Assay kit	R&D systems	Cat No. KGE002B	

4.1 Animals

Adult sea cucumbers were collected from an aquaculture breeding farm in

Weihai, Shandong, China. All the animals were acclimated in breeding aquaria (18 □,

32 ppt) for at least seven days and fed daily with a commercially formulated diet.

Tissues used for cDNA cloning and qRT-PCR were obtained from six adult *A.*

japonicus (~120 g) and kept at -80 □ until use. Juvenile specimens of *A. japonicus*

(~15 mm in length) were used for immunohistochemical analysis of CT-type

neuropeptide expression. *In vitro* functional analysis and *in vivo* experiments were

carried out using adult sea cucumbers with body masses of ~120 g and ~20 g,

respectively.

Ethical approval

All methods were carried out in accordance with the ARRIVE guidelines

(https://arriveguidelines.org) for animal experiments. All animal care and use

procedures were approved by the Institutional Animal Care and Use Committee of

Ocean University of China (Permit Number: 20141201). They were performed

according to the Chinese Guidelines for the Care and Use of Laboratory Animals

(GB/T 35892-2018).

4.2 Molecular structure and phylogenetic analysis of CT-type neuropeptides and

precursors

A. japonicus CT-type neuropeptide precursor sequences (*AjCTP1*, GenBank

accession number: MF401985; *AjCTP2*, GenBank accession number: MF401986)

were identified in our previous study (Chen et al., 2019). Based on the published

sequences, transcript-specific primers for *AjCTP1* and *AjCTP2* (Key resources table)

were designed using Primer3Plus (https://www.primer3plus.com/) and sequence

accuracy was confirmed by SMARTer RACE 5'/3' Kit (TaKaRa, Kusatsu, Japan)

according to the manufacturer's instruction. A sequence translation tool

(https://web.expasy.org/translate/) was used to obtain the corresponding amino acid

sequences encoded by the cloned cDNAs. To identify the conserved features of CT-

type neuropeptides in Bilateria, sequences were aligned using Clustal-X with default

settings and then predicted using the online tool Multiple Sequence Comparison

Display Tool (http://www.bio-soft.net/sms/index.html). To investigate relationships of AjCTP1, AjCTP2 and CT-type precursors from other Bilateria, phylogenetic analyses were performed using the Neighbor joining method. The precursor sequences were aligned using the MUSCLE plugin (iterative, 10 iterations, UPGMB as clustering method), a phylogenetic tree was generated using the neighbor joining method with MEGA7 (v.7170509) and bootstrap values were calculated from 1000 replicate analysis and then modified using iTOL (https://itol.embl.de/login.cgi). The accession numbers or citations for all sequences used for analysis are listed in Supplementary table 1.

4.3 Cloning, sequence analysis and phylogenetic tree construction for candidate AjCT1/2 receptors in A. japonicus

Five potential receptors for AjCT1 and AjCT2 in *A. japonicus* (GenBank accession number: PIK49998.1, PIK49999.1, PIK49604.1, PIK34007.1 and PIK62458.1) were identified based on genome-wide analysis (Huang et al., 2021). Sequence comparison revealed that both PIK49998.1 and PIK62458.1 were partial sequences of PIK34007.1. Thus, three candidate receptors belonging to Class B (Secretin-like) GPCRs named as *AjCTR* (Calcitonin receptor, GenBank accession number: PIK49604.1), *AjPDFR1* (Pigment-dispersing factor receptor 1, GenBank accession number: PIK34007.1), and *AjPDFR2* (Pigment-dispersing factor receptor 2, GenBank accession number: PIK49999.1) were predicted and then analyzed. To confirm the accuracy of these sequences, cDNAs were cloned using the SMARTer RACE 5'/3' Kit (Takara, Cat No. 634858) and then sequenced (Beijing Genomics Institute, China).

Phylogenetic relationships of AjCTR, AjPDFR1 and AjPDFR2 with GPCRs belonging to the CT-type (including DH31 and Cluster A) and PDF-type receptor families in other bilaterian animals were investigated, with CRH-type receptors used as an outgroup. Receptor sequences (see supplementary table 2) were aligned using MUSCLE 5.1 with the PPP algorithm (Edgar, 2022). The alignment was trimmed with the gappyout option using trimAl 1.2rev59 (Capella-Gutiérrez et al., 2009), and

maximum-likelihood phylogenetic trees were generated with IQ-tree 2.0.7 for Linux

64-bit using the SH-aLRT and ultrafast bootstrap (UFBoot) methods with nearest

neighbor interchange (NNI) correction tests (Guindon and Gascuel, 2003; Guindon et

al., 2010; Hoang et al., 2018; Minh et al., 2020; Nguyen et al., 2015). The

VT+F+I+G4 amino acid substitution model was selected, with branch support of 1000

bootstrap replicates. The tree was rooted using CRH-type receptors as the outgroup,

and edited using Treeviewer version 2.2.0 (Bianchini and Sanchez-Baracaldo, 2024).

In addition, the online tool NetPhos-3.1

(https://services.healthtech.dtu.dk/service.php?NetPhos-3.1) was used to predict

potential phosphorylation sites. The conserved domains of protein sequences were

predicted by the Batch SMART plugin of TBtools (Toolbox for biologists, v.1.098652)

software. For analysis of gene synteny, neighboring genes of *AjCTR*, *AjPDFR1* and

AjPDFR2 were found based on an *A. japonicus* chromosome-level genome assembly

(Wang et al., 2022) and the corresponding genes in the starfish *Asterias rubens* and

the sea urchin *Lytechinus variegatus* were also identified by performing BLASTp in

the NCBI database. Specific primers for cloning of cDNAs encoding the candidate

receptors are shown in the Key resources table and the details of the sequences used

for constructing the phylogenetic tree are listed in Supplementary table 2.

4.4 Testing AjCT1 and AjCT2 as ligands for candidate receptors and exploration of

downstream signaling pathways

AjCT1 and AjCT2 (both with a disulfide bridge between the two cysteine

residues and with C-terminal amidation) were custom synthesized by GL Biochem

Ltd. (Shanghai, China) with >95% purity. To test candidate receptors for CT-type

neuropeptides and investigate their downstream activation mechanisms the following

assays were used to measure cAMP signaling or Ca²□ signaling: i). direct

measurement of cAMP levels using ELISA ii). indirect measurement of cAMP

signaling using a cAMP response element-Luciferase (CRE-Luc) assay that measures

cAMP-induced gene expression, iii). serum response element (SRE)-Luc assays that

measure Ca^{2+}-induced gene expression, iv). receptor translocation assays and v).

Western blotting.

AjCTR, *AjPDFR1* and *AjPDFR2* incorporating enhanced green fluorescent

protein (EGFP) were constructed respectively according to the method described by

Li *et al.* (2022). Briefly, the complete ORFs of candidate receptors were cloned and

sequenced. Then, the PCR products were digested with EcoRI and BamHI (TaKaRa,

Kusatsu, Japan) and then incorporated into the pEGFP-N1 vector (YouBio, Changsha,

China) that was subject to the same double-digestion. Finally, the constructed plasmid

with correct sequences was transfected into HEK293T cells seeded on glass

coverslips in 12-well plates. After incubation with AjCT1 (10^{-6} M), AjCT2 (10^{-6} M) or

serum-free DMEM (Biosharp, Hefei, China) for 15 min, cells were washed with PBS

(Biosharp, Hefei, China) and fixed with 4% PFA for 10 min. Finally, cell nuclei were

stained with DAPI (G-CLONE, Beijing, China) for 5 min and an Olympus bright field

light microscope was used for image capture. The pcDNA 3.1(+) vectors (YouBio,

Changsha, China) comprising the complete ORF of the three candidate receptors were

also constructed as described by Li *et al.* (2022). Then, they were co-transfected with

the cAMP reporter pCRE-Luc (YouBio, Changsha, China) or Ca^{2+} signaling reporter

pSRE-Luc plasmid including a PKC reporter gene (YouBio, Changsha, China) into

HEK293T cells. Incubation was carried out with AjCT1 (10^{-6} M), AjCT2 (10^{-6} M) or

serum-free DMEM (CO, negative control) for 4 h at 37 °C. Finally, luciferase activity

was tested using the firefly luciferase assay kit (MK, Shanghai, China). When

required, transfected cells described above were co-incubated with H89 (PKA

inhibitor) (Absin, Shanghai, China), FR900359 (Gαq protein inhibitor) (Cayman

Chemical, Ann Arbor, Michigan) or Gö 6983 (PKC inhibitor) (Cayman Chemical,

Ann Arbor, Michigan) before the experiment to explore the potential signaling

pathways. To confirm the concentration of intracellular cAMP after stimulation,

HEK293T cells expressing pcDNA 3.1(+)/*AjCTR*, pcDNA 3.1(+)/*AjPDFR2* or

pcDNA 3.1(+)/*AjPDFR1* were treated with AjCT1 or AjCT2 (10^{-9} – 10^{-5} M),

respectively, for 15 min after 2 h starvation in serum-free medium. Subsequently,

cAMP was measured using an ELISA-based cAMP assay kit (R&D systems,

USA&Canada) following the manufacturer's instruction. HEK293T cells transfected

with empty pcDNA3.1(+) were used as a negative control. The specific primers used

for constructing plasmids are listed in Key resources table.

Western blotting was performed as described previously (Li et al., 2022). Briefly,

HEK293T cells transfected with the constructed pcDNA 3.1(+) plasmid were starved

in serum-free medium for 2 h, and then incubated with AjCT1 or AjCT2 (10^{-6} M).

PVDF membranes containing proteins were blocked with 5% BSA (G-CLONE,

Beijing, China) and incubated overnight with antibodies to phosphorylated ERK1/2

kinases (CST, Boston, China). After that, goat anti-rabbit horseradish peroxidase-

conjugated secondary antibody (Absin, Shanghai, China) was applied and incubated,

followed by the addition of Omni-ECL™ Femto Light Chemiluminescence reagents

(EpiZyme, Shanghai, China). Finally, RVL-100-G (ECHO, America) was applied for

visualization of the PVDF membrane. When required, transfected cells were co-

incubated with PKA or PKC inhibitor before the experiment to identify signaling

pathways that activate downstream ERK1/2 cascade.

4.5 Total RNA extraction and Quantitative Real-Time PCR (qRT-PCR) detection

qRT-PCR was employed to detect *AjCTR*, *AjPDFR1*, *AjPDFR2*, *AjCTP1* and

AjCTP2 transcript expression in five tissues from *A. japonicus*, including the intestine,

longitudinal muscle, respiratory tree, gonad and circumoral nervous system, and the

circumoral nervous system was selected as the control group. Because it was difficult

to design specific primers for *AjCTP1* and *AjCTP2* (Zheng et al., 2022), total

transcript expression levels for *AjCTP1/2* (*AjCTP1* + *AjCTP2*) were investigated.

Total RNA was extracted from the collected tissues using Trizol RNA isolation

reagent (Vazyme, Nanjing, China) and Hifair® Ⅲ 1st Strand cDNA Synthesis

SuperMix (YEASEN, Shanghai, China) was used to remove genomic DNA and

synthesize cDNA according to manufacturer's instructions. For qRT-PCR,

amplification was performed in 20 µL volume reactions based on Hieff UNICON

Universal Blue qPCR SYBR Green Master Mix (YEASEN, Shanghai, China) and

expression levels were assessed by a Corbett Rotoe-Gnen Q (Qiagen, Germany). *β*-

actin (GenBank accession number: PIK61412.1) and *β*-tubulin (GenBank accession

number: PIK51093) were used as house-keeping genes for normalization (Zhao et al.,

2014). The $2^{-\Delta\Delta CT}$ method was applied to analyze the relative expression levels.

Specific sequences of primers used for qRT-PCR are listed in Key resources table.

4.6 Localization of CT-type neuropeptide expression in A. japonicus using

immunohistochemistry

Juvenile specimens of *A. japonicus* (~15 mm in length) were fixed for 24 hours

at room temperature or for 48 hours at 4 °C in Bouin's fixative (75 mL saturated picric

acid in seawater, 25 mL 37% formaldehyde, 5 mL acetic acid) days and then washed

in water before being transferred to 50% ethanol followed by 70% ethanol. The fixed

specimens were in wrapped in tissue paper soaked in 70% ethanol and then

transported from China to the UK. Following dehydration through 90% and 100%

ethanol and clearing in xylene, specimens were embedded in paraffin wax and then

sectioned (9 or 10 µm) transversely or longitudinally using a Leica RM2145

microtome. Sections were mounted on chrome alum-gelatin coated glass slides and

then after incubation in xylene to remove wax followed by 100% ethanol, slides were

incubated in 1% hydrogen peroxide in phosphate buffered saline (PBS, pH 7.3) to

quench endogenous peroxidases. After using an ethanol series to rehydrate sections,

slides were blocked in 5% goat serum in PBS containing 0.1% Tween-20 (PBST).

Then slides were incubated overnight at 4 °C with the ArCT antiserum diluted to

1:2000, 1:3000 or 1:4000 in PBST. Following washing in PBST, slides were

incubated with Peroxidase-AffiniPure Goat Anti-Rabbit IgG (H + L) conjugated to

Horseradish Peroxidase (RRID: AB_2313567; Jackson ImmunoResearch, West Grove,

PA) diluted 1:1000 in 2% goat serum/PBST for 3 h. Bound antibodies were revealed

using a solution of 0.05% diaminobenzidine (VWR Chemicals, UK), 0.05% nickel

chloride (Sigma-Aldrich, Gillingham, UK), 0.015% hydrogen peroxide (VWR

Chemicals, UK) diluted in PBS. When staining was observed, slides were washed in

water and dehydrated through an ethanol series into xylene and mounted with

coverslips over DPX mounting medium (Thermo Fisher Scientific). The specificity of

immunstaining was assessed by pre-absorption of the ArCT antiserum (at 1:100

dilution) for 1 h at room temperature with the ArCT antigen peptide (25 µM) prior to

testing on sections at 1:3000 dilution. Images of stained sections were taken using a

QIClich CCD Colour Camera (Qimaging, UK) linked to a DMRAI light microscope

(Leica Microsystems, UK) with Volocity v. 6.3.1 image analysis software (Perkin-

Elmer, USA) running on an iMac computer (27 inch with OS Yosemite, v. 10.10).

Montages of images were prepared and labelled using GIMP (version 3.0.2) and

Adobe Photoshop (version 26.7) running on a MacBook Pro laptop (OS Sequoia 15.5).

4.7 Analysis of the in vitro activity of AjCT1 and AjCT2 on preparations from A.

japonicus

Longitudinal muscles and intestines (20 mm in length) from *A. japonicus* were

dissected and then suspended in a 50 mL glass organ bath containing 20 mL sterilized

seawater. Both ends of the preprations were tied with cotton ligatures and then one

end was tied to a metal hook and the other end was connected with a High Grade

Isotonic Transducer (ADinstruments MLT0015, Oxford, UK). Before testing, the

preparation was stabilized for about 20 min, and the resting tension was adjusted to

1.0 g. Informed by previous findings from Cai et al. (2018), we investigated if AjCT1

and AjCT2 caused relaxation of preparations. First, acetylcholine (ACh) was applied

to induce the contraction of preparations and then when a stable state of contraction

was achieved, 20 uL AjCT1 or AjCT2 at a range of concentrations was added and

relaxation responses were recorded. Output signal (cm) from PowerLab data

acquisition hardware (ADinstruments PowerLab 4/26, Oxford, UK) was calculated

using LabChart (v8.0.7) software.

4.8 Investigation of the in vivo pharmacological effects of AjCT1 and AjCT2 in A.

japonicus

A total of 150 adult sea cucumbers were randomly assigned to five groups (30

individuals per group), including two high concentration groups (CT1H and CT2H, 5

$\times 10^{-1}$ mg/ml), two low concentration groups (CT1L and CT2L, 5×10^{-3} mg/ml) and

the control group (CO, sterilized seawater), with each group consisting of three

parallel experiments comprising ten animals. Before the experiments, sea cucumbers were starved for a day to normalize their physiological status and their wet weights were measured. AjCT1 and AjCT2 were dissolved in sterilized seawater and then injected through the peristome at noon every other day (1 µl/g wet weight) and the CO group were injected in the same way but with sterilized seawater (1 µl/g wet weight). Feces and residual bait were collected and weighed after drying every day. After 24 days, wet weights of sea cucumbers were measured again. Then, WGR and SGR were calculated to assess their feeding and growth conditions. Formulas are as follows: $WGR = 100*(W_t-W_0)/W_0$, $SGR = 100*(lnW_t-lnW_0)/D$. In the formula, W_0 represents the initial wet body weight (g), W_t represents the final wet body weight (g), D represents the feeding days. Subsequently, animals were dissected and intestines were collected for detecting the expression levels of the three receptor genes (*AjCTR, AjPDFR1, AjPDFR2*) and several growth factors, including multiple epidermal growth factor 6 (*AjMegf6*, GenBank accession number: MG018199.1)*,* insulin-like growth factor (*AjIgf*, GenBank accession number: PIK50518.1) and growth and differentiation factor-8 (*AjGDF-8*, GenBank accession number: KP100064.1) (Li et al., 2016; Wang et al., 2018). The specific primers for qRT-PCR analysis are listed in the Key resources table.

4.9 Loss-of-function experiments by RNA interference (RNAi)

To investigate how receptor expression is affected by knock-down of AjCTP expression, an RNAi experiment was carried out. It was difficult to design *AjCTP1-* and *AjCTP2-* specific interference oligonucleotides and therefore, we designed and customized generic interference oligonucleotides targeting both *AjCTP1* and *AjCTP2* (si*AjCTP1/2*-1 and si*AjCTP1/2*-2, containing the extra 2′ ome modification) (Beijing Tsingke Biotech Co., Ltd. China). Meanwhile, a negative control siRNA (siNC) was also customized for the experiment (Key resources table), which does not target genes in *A. japonicus*. The si*AjCTP1/2*-1, si*AjCTP1/2*-2 or siNC dry powder was diluted to 20 µM with DEPC water following the manufacturer's instruction, then 60 µl sterilized seawater was mixed with 20 µl si*AjCTP1/2*-1, si*AjCTP1/2*-2 or siNC, and

20 μl Lipo6000 transfection reagent (Beyotime, Shanghai, China) as the transfection

solution. Eighteen sea cucumbers were randomly divided into three groups and

injected with three different transfection solutions respectively. The mixed solutions

were injected through the peristome once a day (1 μl/g wet weight) and intestines

were dissected and frozen in liquid nitrogen 3 days after injection. The expression

level of *AjCTP1/2* was investigated by qRT-PCR and the interference rate was

calculated using $(1 - 2^{-\Delta\Delta^{CT}})$ to determine the inhibition efficiency of siRNAs. Finally,

the relative expression levels of *AjCTR*, *AjPDFR1* and *AjPDFR2* were investigated

using the selected *A. japonicus* intestines.

To assess the effects of CT-type signaling inhibition on feeding and growth in *A.*

japonicus, a 24-day RNAi experiment was conducted following the methodology

outlined in Section 4.8. Sea cucumbers were randomly allocated into two groups: the

negative control group received siNC injections, while the experimental group was

injected with si*AjCTP1/2*-1 (effectively targeting *AjCTP1/2*). Throughout the

experiment, the dry weight of remaining bait and excrement, along with the wet

weight of individual sea cucumbers, were recorded. Subsequently, WGR and SGR, as

indices of feeding efficiency and growth performance, were calculated. In addition,

the expression levels of the three growth factors (*AjMegf6*, *AjIgf* and *AjGDF-8*) in the

intestinal tissue were tested by qPCR.

To investigate the potential mediation of CT-type signaling effects on feeding

and growth by AjPDFR2 in *A. japonicus*, an RNAi experiment targeting *AjPDFR2*

was performed following the same knockdown protocol established for *AjCTP*.

Briefly, the effective siRNA targeting *AjPDFR2* (si*AjPDFR2*-1 or si*AjPDFR2*-2) was

selected by testing *AjPDFR2* expression in the intestines of injected sea cucumbers

using qRT-PCR and calculating the interference rate efficiency. Subsequently, the

validated siRNA was then employed in a 24-day RNAi experiment. The dry weight of

remaining bait and excrement, as well as the wet weight of the sea cucumbers were

measured, enabling calculation of WGR and SGR. Furthermore, the expression levels

of three key growth factors (*AjMegf6*, *AjIgf* and *AjGDF-8*) in the intestinal tissue were

analyzed via qPCR.

4.10 Statistical analysis

Statistical analyses were carried out using GraphPad Prism 8.0. Significant

differences between groups were determined by one-way analysis of variance

(ANOVA) followed by a Tukey's multiple comparisons test. Statistical significance

thresholds were defined as follows: $p < 0.05$ (*) for significant differences and $p <$

0.01 (**) for highly significant differences, with annotations directly marked on the

figures. All data were presented as mean ± SEM from at least three independent

biological replicates. Non-linear regression analysis was performed using a four-

parameter logistic equation with automatic outlier exclusion to fit the dose-response

curve. The half maximal effective concentration (EC_{50}) was derived from the fitted

curves through built-in algorithms in GraphPad Prism 8.0.

Competing Interests

The authors declare that no competing interests exist.

Funding

This research was supported by National Natural Science Foundation of China [grant number 42276103] and Biotechnology and Biological Sciences Research Council (UK; grant numbers BB/M001644/1 and BB/X001024/1).

Author Contributions

Xiao Cong, Data curation, Formal analysis, Investigation, Methodology, Software, Visualization, Writing—original draft preparation; Huachen Liu, Investigation, Methodology, Validation; Lihua Liu, Methodology, Validation; Nayeli Escudero-Castelán, Investigation, Methodology, Visualization, Writing—review & editing; Kite Jones, Investigation, Methodology, Visualization, Writing—review & editing; Michaela Egertová, Investigation, Methodology, Visualization; Maurice R. Elphick, Formal analysis, Funding acquisition, Resources, Supervision, Validation, Visualization, Writing—review & editing; Muyan Chen, Conceptualization, Formal analysis, Funding acquisition, Methodology, Project administration, Resources, Supervision, Validation, Visualization, Writing—review & editing

References:

Aleotti A, Wilkie I C, Yañez-Guerra L A, et al. Discovery and functional characterization of neuropeptides in crinoid echinoderms. Frontiers in Neuroscience, 2022, 16: 1006594. doi: 10.3389/fnins.2022.1006594.

Amara S G, Jonas V, Rosenfeld M G, et al. Alternative RNA processing in calcitonin gene expression generates mRNAs encoding different polypeptide products. Nature, 1982, 298(5871): 240-244. doi: 10.1038/298240a0.

Andreassen K V, Hjuler S T, Furness S G, et al. Prolonged calcitonin receptor signaling by salmon, but not human calcitonin, reveals ligand bias. PloS one, 2014, 9(3): e92042. doi: 10.1371/journal.pone.0092042.

Arrowsmith S, Wray S. Oxytocin: its mechanism of action and receptor signaling in the myometrium. Journal of neuroendocrinology, 2014, 26(6): 356-369. dio: 10.1111/jne.12154.

Bauknecht P, Jékely G. Large-Scale Combinatorial Deorphanization of Platynereis Neuropeptide GPCRs. Cell reports, 2015, 12(4), 684–693. doi: 10.1016/j.celrep.2015.06.052.

Beets I, Zels S, Vandewyer E, Demeulemeester J, et al. System-wide mapping of peptide-GPCR interactions in *C. elegans*. Cell reports, 2023, 42(9), 113058. doi: 10.1016/j.celrep.2023.113058.

Bianchini G, Sánchez-Baracaldo P. TreeViewer: Flexible, modular software to visualise and
manipulate phylogenetic trees. Ecology and evolution, 2024, 14(2), e10873. doi:
10.1002/ece3.10873.

Bomberger J M, Parameswaran N, Spielman W S. Regulation of GPCR trafficking by RAMPs.
Adv. Exp. Med. Biol, 2012, 744: 25-37. doi: 10.1007/978-1-4614-2364-5_3.

Bourlat S J, Juliusdottir T, Lowe C J, et al. Deuterostome phylogeny reveals monophyletic
chordates and the new phylum Xenoturbellida. Nature, 2006, 444(7115): 85-88. doi:
10.1038/nature05241

Bromham L D, Degnan B M. Hemichordates and deuterostome evolution: robust molecular
phylogenetic support for a hemichordate+ echinoderm clade. Evolution & development, 1999,
1(3): 166-171. doi: 10.1046/j.1525-142x.1999.99026.x

Burke R D, Angerer L M, Elphick M R, et al. A genomic view of the sea urchin nervous system.
Developmental biology, 2006, 300(1): 434-460. doi: 10.1016/j.ydbio.2006.08.007.

Cai W, Kim C H, Go H J, et al. Biochemical, anatomical, and pharmacological characterization of
calcitonin-type neuropeptides in starfish: discovery of an ancient role as muscle relaxants.
Frontiers in Neuroscience, 2018, 12: 382. doi: 10.3389/fnins.2018.00382.

Capella-Gutiérrez S, Silla-Martínez J M, Gabaldón T. trimAl: a tool for automated alignment
trimming in large-scale phylogenetic analyses. Bioinformatics (Oxford, England), 2009,
25(15), 1972–1973. doi: 10.1093/bioinformatics/btp348.

Cardoso J C, Mc Shane J C, Li Z, et al. Revisiting the evolution of Family B1 GPCRs and ligands:
Insights from mollusca. Molecular and cellular endocrinology, 2024, 586, 112192. doi:
10.1016/j.mce.2024.112192.

Changeux J P. Allosteric receptors: from electric organ to cognition. Annual review of
pharmacology and toxicology, 2010, 50(1): 1-38. doi:
10.1146/annurev.pharmtox.010909.105741.

Chen M, Talarovicova A, Zheng Y, et al. Neuropeptide precursors and neuropeptides in the sea
cucumber *Apostichopus japonicus*: a genomic, transcriptomic and proteomic analysis.
Scientific Reports, 2019, 9(1): 8829. doi: 10.1038/s41598-019-45271-3.

Chen Y, Shyu J F, Santhanagopal A, et al. The calcitonin receptor stimulates Shc tyrosine
phosphorylation and Erk1/2 activation: involvement of Gi, protein kinase C, and calcium[J].
Journal of Biological Chemistry, 1998, 273(31): 19809-19816. doi:
10.1074/jbc.273.31.19809.

Cong X, Liu H, Zheng Y, et al. A Putative Role of Vasopressin/Oxytocin-Type Neuropeptide in
Osmoregulation and Feeding Inhibition of *Apostichopus japonicus*. International Journal of
Molecular Sciences, 2023, 24(18): 14358. doi: 10.3390/ijms241814358.

Conner A C, Simms J, Hay D L, et al. Heterodimers and family-B GPCRs: RAMPs, CGRP and
adrenomedullin. Biochemical Society Transactions, 2004, 32(5): 843-846. doi:
10.1042/BST0320843.

Conzelmann M, Williams E A, Krug K, et al. The neuropeptide complement of the marine annelid
Platynereis dumerilii. BMC genomics, 2013, 14: 1-15. doi:10.1186/1471-2164-14-906.

Copp D H, Cameron E C, Cheney B A, et al. Evidence for calcitonin—a new hormone from the
parathyroid that lowers blood calcium. Endocrinology, 1962, 70(5): 638-649. doi:
10.1210/endo-70-5-638.

Del Prete E, Schade B, Riediger T, et al. Effects of amylin and salmon calcitonin on feeding and
drinking behavior in pygmy goats. Physiology & behavior, 2002, 75(4): 593-599. doi:
10.1016/s0031-9384(02)00655-8.

Deribe Y L, Wild P, Chandrashaker A, et al. Regulation of epidermal growth factor receptor
trafficking by lysine deacetylase HDAC6. Science signaling, 2009, 2(102): ra84-ra84. doi:
10.1126/scisignal.2000576.

Dunn C W, Hejnol A, Matus D Q, et al. Broad phylogenomic sampling improves resolution of the
animal tree of life. Nature, 2008, 452(7188): 745-749. doi: 10.1038/nature06614

Edgar R C. Muscle5: High-accuracy alignment ensembles enable unbiased assessments of
sequence homology and phylogeny. Nature communications, 2022, 13(1), 6968. doi:
10.1038/s41467-022-34630-w.

Edvinsson L, Goadsby P J, Uddman R. Amylin: localization, effects on cerebral arteries and on
local cerebral blood flow in the cat. The scientific world journal, 2001, 1: 168-180. doi:
10.1100/tsw.2001.23.

Furuya K, Milchak R J, Schegg K M, et al. Cockroach diuretic hormones: characterization of a
calcitonin-like peptide in insects. Proceedings of the National Academy of Sciences, 2000,
97(12): 6469-6474. doi: 10.1073/pnas.97.12.6469.

Garelja M L, Bower R L, Brimble M A, et al. Pharmacological characterisation of mouse
calcitonin and calcitonin receptor□like receptors reveals differences compared with human
receptors. British journal of pharmacology, 2022, 179(3): 416-434. doi: 10.1111/bph.15628.

Goda T, Umezaki Y, Alwattari F, et al. Neuropeptides PDF and DH31 hierarchically regulate free-
running rhythmicity in Drosophila circadian locomotor activity. Scientific reports, 2019, 9(1):
838. doi:10.1038/s41598-018-37107-3.

Golpon H A, Puechner A, Welte T, et al. Vasorelaxant effect of glucagon-like peptide-(7-36) amide
and amylin on the pulmonary circulation of the rat. Regulatory peptides, 2001, 102(2-3): 81-
86. doi: 10.1016/s0167-0115(01)00300-7.

Gorn A H, Lin H Y, Yamin M, et al. Cloning, characterization, and expression of a human
calcitonin receptor from an ovarian carcinoma cell line. The Journal of clinical investigation,

1992, 90(5), 1726–1735. doi: 10.1172/JCI116046.

Guindon S, Dufayard J F, Lefort V, et al. New algorithms and methods to estimate maximum-
likelihood phylogenies: assessing the performance of PhyML 3.0. Systematic biology, 2010,
59(3), 307–321. doi: 10.1093/sysbio/syq010.

Guindon S, Gascuel O. A simple, fast, and accurate algorithm to estimate large phylogenies by maximum likelihood. Systematic biology, 2003, 52(5), 696–704. doi: 10.1080/10635150390235520.

Guo K, Su L, Wang Y, et al. Antioxidant and anti-aging effects of a sea cucumber protein hydrolyzate and bioinformatic characterization of its composing peptides. Food & Function, 2020, 11(6): 5004-5016. doi: 10.1039/d0fo00560f.

Hay D L, Conner A C, Howitt S G, et al. The pharmacology of adrenomedullin receptors and their relationship to CGRP receptors. Journal of Molecular Neuroscience, 2004, 22: 105-113. doi: 10.1385/JMN:22:1-2:105.

Hay D L, Garelja M L, Poyner D R, et al. Update on the pharmacology of calcitonin/CGRP family of peptides: IUPHAR Review 25. British journal of pharmacology, 2018, 175(1): 3-17. doi: 10.1111/bph.14075.

Hoang D T, Chernomor O, von Haeseler A, et al. UFBoot2: Improving the Ultrafast Bootstrap Approximation. Molecular biology and evolution, 2018, 35(2), 518–522. doi: 10.1093/molbev/msx281.

Huang D, Zhang B, Han T, et al. Genome-wide prediction and comparative transcriptomic analysis reveals the G protein-coupled receptors involved in gonadal development of *Apostichopus japonicus*. Genomics, 2021, 113(1): 967-978. doi: 10.1016/j.ygeno.2020.10.030.

Huang T, Su J, Wang X, et al. Functional Analysis and Tissue-Specific Expression of Calcitonin and CGRP with RAMP-Modulated Receptors CTR and CLR in Chickens. Animals: an open access journal from MDPI, 2024, 14(7), 1058. doi: 10.3390/ani14071058.

Jékely G. Global view of the evolution and diversity of metazoan neuropeptide signaling. Proceedings of the National Academy of Sciences, 2013, 110(21): 8702-8707. doi: 10.1073/pnas.1221833110.

Johnson E C, Shafer O T, Trigg J S, et al. A novel diuretic hormone receptor in Drosophila: evidence for conservation of CGRP signaling. Journal of Experimental Biology, 2005, 208(7): 1239-1246. doi: 10.1242/jeb.01529.

Kandilci H B, Gumusel B, Lippton H. Intermedin/adrenomedullin-2 (IMD/AM2) relaxes rat main pulmonary arterial rings via cGMP-dependent pathway: role of nitric oxide and large conductance calcium-activated potassium channels (BKCa). Peptides, 2008, 29(8): 1321-1328. doi: 10.1016/j.peptides.2008.04.008.

Katafuchi T, Yasue H, Osaki T, et al. Calcitonin receptor-stimulating peptide: Its evolutionary and functional relationship with calcitonin/calcitonin gene-related peptide based on gene structure. Peptides, 2009, 30(9): 1753-1762. doi: 10.1016/j.peptides.2009.06.012.

Kuwasako K, Kitamura K, Nagata S, et al. Shared and separate functions of the RAMP-based adrenomedullin receptors. Peptides, 2011, 32(7): 1540-1550. doi: 10.1016/j.peptides.2011.05.022.

Li C, Zheng Y, Cong X, et al. Molecular and functional characterization of the luqin-type neuropeptide signaling system in the sea cucumber *Apostichopus japonicus*. Peptides, 2022, 155: 170839. doi: 10.1016/j.peptides.2022.170839.

Li S, Zhou Z, Dong Y, et al. Molecular characterization, expression analysis of the myostatin gene and its association with growth traits in sea cucumber (*Apostichopus japonicus*). Comparative Biochemistry and Physiology Part B: Biochemistry and Molecular Biology, 2016, 201: 12-20. doi: 10.1016/j.cbpb.2016.06.005.

Martins R, Vieira F A, Power D M. Calcitonin receptor family evolution and fishing for function using in silico promoter analysis. General and Comparative Endocrinology, 2014, 209: 61-73. doi: 10.1016/j.ygcen.2014.04.025.

Mathiesen D S, Lund A, Vilsbøll T, et al. Amylin and calcitonin: potential therapeutic strategies to reduce body weight and liver fat. Frontiers in Endocrinology, 2021, 11: 617400. doi: 10.3389/fendo.2020.617400.

Mayorova T D, Tian S, Cai W, Semmens D C. Odekunle E A, Zandawala M, Badi Y, Rowe M L, Egertová M, Elphick M R. (2016). Localization of neuropeptide gene expression in larvae of an echinoderm, the starfish *Asterias rubens*. Front. Neurosci. 10:553. doi: 10.3389/fnins.2016.00553.

McLatchie L M, Fraser N J, Main M J, et al. RAMPs regulate the transport and ligand specificity of the calcitonin-receptor-like receptor. Nature, 1998, 393(6683): 333-339. doi: 10.1038/30666.

Mendel H C, Kaas Q, Muttenthaler M. Neuropeptide signalling systems–an underexplored target for venom drug discovery. Biochemical pharmacology, 2020, 181: 114129. doi: 10.1016/j.bcp.2020.114129.

Mertens I, Vandingenen A, Johnson E C, et al. PDF receptor signaling in Drosophila contributes to both circadian and geotactic behaviors. Neuron, 2005, 48(2): 213-219. doi: 10.1016/j.neuron.2005.09.009.

Minh B Q, Schmidt H A, Chernomor O, et al. IQ-TREE 2: New Models and Efficient Methods for Phylogenetic Inference in the Genomic Era. Molecular biology and evolution, 2020, 37(5), 1530–1534. doi: 10.1093/molbev/msaa015.

Mirabeau O, Joly J S. Molecular evolution of peptidergic signaling systems in bilaterians. Proceedings of the national academy of sciences, 2013, 110(22): E2028-E2037. doi: 10.1073/pnas.1219956110.

Møller L N, Stidsen C E, Hartmann B, et al. Somatostatin receptors. Biochimica et Biophysica Acta (BBA)-Biomembranes, 2003, 1616(1): 1-84. doi: 10.1016/s0165-6147(97)89535-x.

Morris H R, Panico M, Etienne T, et al. Isolation and characterization of human calcitonin gene-related peptide. Nature, 1984, 308(5961): 746-748. doi: 10.1038/308746a0.

Nakamura M, Nomura S, Yamakawa T, et al. Endogenous calcitonin regulates lipid and glucose metabolism in diet-induced obesity mice. Scientific Reports, 2018, 8(1): 17001. Endogenous calcitonin regulates lipid and glucose metabolism in diet-induced obesity mice

Nguyen L T, Schmidt H A, von Haeseler A, et al. IQ-TREE: a fast and effective stochastic algorithm for estimating maximum-likelihood phylogenies. Molecular biology and evolution, 2015, 32(1), 268–274. doi: 10.1093/molbev/msu300.

Pangestuti R, Arifin Z. Medicinal and health benefit effects of functional sea cucumbers. Journal of traditional and complementary medicine, 2018, 8(3): 341-351. doi: 10.1016/j.jtcme.2017.06.007.

Párrizas M, LeRoith D. Insulin-like growth factor-1 inhibition of apoptosis is associated with increased expression of the bcl-xL gene product. Endocrinology, 1997, 138(3): 1355-1358. doi: 10.1210/endo.138.3.5103.

Perlow M J, Freed W J, Carman J S, et al. Calcitonin reduces feeding in man, monkey and rat. Pharmacology Biochemistry and Behavior, 1980, 12(4): 609-612. doi: 10.1016/0091-3057(80)90196-3.

Pinto A, Sekizawa K, Yamaya M, et al. Effects of adrenomedullin and calcitonin gene□related peptide on airway and pulmonary vascular smooth muscle in guinea□pigs. British journal of pharmacology, 1996, 119(7): 1477-1483. doi: 10.1111/j.1476-5381.1996.tb16061.x.

Pondel M. Calcitonin and calcitonin receptors: bone and beyond. International journal of experimental pathology, 2000, 81(6): 405-422. doi: 10.1046/j.1365-2613.2000.00176.x.

Purdue B W, Tilakaratne N, Sexton P M. Molecular pharmacology of the calcitonin receptor. Receptors and Channels, 2002, 8(3-4): 243-255. doi: 10.3109/10606820213681

Rowe M L, Achhala S, Elphick M R. Neuropeptides and polypeptide hormones in echinoderms: new insights from analysis of the transcriptome of the sea cucumber *Apostichopus japonicus*. General and Comparative Endocrinology, 2014, 197: 43-55. doi: 10.1016/j.ygcen.2013.12.002.

Rowe M L, Elphick M R. The neuropeptide transcriptome of a model echinoderm, the sea urchin *Strongylocentrotus purpuratus*. General and comparative endocrinology, 2012, 179(3): 331-344. doi: 10.1016/j.ygcen.2012.09.009.

Schwartz J, Réalis-Doyelle E, Dubos M P, et al. Characterization of an evolutionarily conserved calcitonin signaling system in a lophotrochozoan, the Pacific oyster (*Crassostrea gigas*). Journal of Experimental Biology, 2019, 222(13): jeb201319. doi: 10.1242/jeb.201319.

Sekiguchi T, Kuwasako K, Ogasawara M, et al. Evidence for conservation of the calcitonin superfamily and activity-regulating mechanisms in the basal chordate *Branchiostoma floridae*: insights into the molecular and functional evolution in chordates. Journal of Biological Chemistry, 2016, 291(5): 2345-2356. doi: 10.1074/jbc.M115.664003.

Sekiguchi T, Suzuki N, Fujiwara N, et al. Calcitonin in a protochordate, *Ciona intestinalis*–the
prototype of the vertebrate calcitonin/calcitonin gene☐related peptide superfamily. The FEBS
Journal, 2009, 276(16): 4437-4447. doi: 10.1111/j.1742-4658.2009.07151.x.

Sekiguchi T. Evolution of calcitonin/calcitonin gene-related peptide family in chordates:
identification of CT/CGRP family peptides in cartilaginous fish genome. General and
Comparative Endocrinology, 2022, 328: 114123. doi: 10.1016/j.ygcen.2022.114123.

Semmens D C, Mirabeau O, Moghul I, Pancholi M R, Wurm Y, and Elphick M R (2016).
Transcriptomic identification of starfish neuropeptide precursors yields new insights into
neuropeptide evolution. Open Biol. 6:150224. doi: 10.1098/rsob.150224.

Shafer O T, Kim D J, Dunbar-Yaffe R, et al. Widespread receptivity to neuropeptide PDF
throughout the neuronal circadian clock network of Drosophila revealed by real-time cyclic
AMP imaging. Neuron, 2008, 58(2): 223-237. doi: 10.1016/j.neuron.2008.02.018.

Smith M K, Wang T, Suwansa-Ard S, et al. The neuropeptidome of the Crown-of-Thorns Starfish,
Acanthaster planci. Journal of proteomics, 2017, 165: 61-68. doi:
10.1016/j.jprot.2017.05.026.

Steiner D F, Ohagi S, Nagamatsu S, et al. Is islet amyloid polypeptide a significant factor in
pathogenesis or pathophysiology of diabetes?. Diabetes, 1991, 40(3): 305-309. doi:
10.2337/diab.40.3.305.

Suwansa-Ard S, Chaiyamoon A, Talarovicova A, et al. Transcriptomic discovery and comparative
analysis of neuropeptide precursors in sea cucumbers (Holothuroidea). Peptides, 2018, 99:
231-240. doi: 10.1016/j.peptides.2017.10.008.

Toribio R E, Kohn C W, Leone G W, et al. Molecular cloning and expression of equine calcitonin,
calcitonin gene-related peptide-I, and calcitonin gene-related peptide-II. Molecular and
cellular endocrinology, 2003, 199(1-2): 119-128. doi: 10.1016/s0303-7207(02)00289-7.

Tsurumaki T, Muraoka O, Yamaguchi T, et al. Neuropeptide Y-induced contraction and its
desensitization through the neuropeptide Y receptor subtype in several rat veins. Journal of
Cardiovascular Pharmacology, 2003, 41: S23-S27.

Walker C S, Conner A C, Poyner D R, et al. Regulation of signal transduction by calcitonin gene-
related peptide receptors. Trends in pharmacological sciences, 2010, 31(10): 476-483. doi:
10.1016/j.tips.2010.06.006.

Wang Y, Yang Y, Li Y, et al. Identification of sex determination locus in sea cucumber
Apostichopus japonicus using genome-wide association study. BMC genomics, 2022, 23(1):
391. doi: 10.1186/s12864-022-08632-3.

Wang Z, Cui J, Song J, et al. Comparative transcriptome analysis reveals growth-related genes in
juvenile Chinese sea cucumber, Russian sea cucumber, and their hybrids. Marine
biotechnology, 2018, 20: 193-205. doi: 10.1007/s10126-018-9796-6.

Wong A H C, Buckle C E, Van Tol H H M. Polymorphisms in dopamine receptors: what do they
tell us? European journal of pharmacology, 2000, 410(2-3): 183-203. doi: 10.1016/s0014-
2999(00)00815-3.

Yoshimoto R, Mitsui□Saito M, Ozaki H, et al. Effects of adrenomedullin and calcitonin
gene□related peptide on contractions of the rat aorta and porcine coronary artery. British
journal of pharmacology, 1998, 123(8): 1645-1654. doi: 10.1038/sj.bjp.0701805.

Zandawala M, Moghul I, Yañez Guerra L A, et al. Discovery of novel representatives of bilaterian
neuropeptide families and reconstruction of neuropeptide precursor evolution in ophiuroid
echinoderms. Open biology, 2017, 7(9): 170129. doi: 10.1098/rsob.170129.

Zandawala M. Calcitonin-like diuretic hormones in insects. Insect biochemistry and molecular
biology, 2012, 42(10): 816-825. doi: 10.1016/j.ibmb.2012.06.006.

Zhao Y, Chen M, Wang T, et al. Selection of reference genes for qRT-PCR analysis of gene
expression in sea cucumber *Apostichopus japonicus* during aestivation. Chinese Journal of
Oceanology and Limnology, 2014, 32(6): 1248-1256. doi: 10.1007/s00343-015-4004-2

Zheng Y, Cong X, Liu H, et al. Nervous system development and neuropeptides characterization
in embryo and larva: insights from a non-chordate deuterostome, the sea cucumber
Apostichopus japonicus. Biology, 2022, 11(10): 1538. doi: 10.3390/biology11101538.

Zheng Y, Liu H, Cong X, et al. A potential feeding regulation strategy during aestivation:
Relaxation of intestine mediated by pedal peptide/orcokinin-type neuropeptides in sea
cucumber *Apostichopus japonicus*. Aquaculture, 2024, 579: 740150.
doi:10.1016/j.aquaculture.2023.740150.

1249

1250 **Supplementary figures**

1251

1252

Figure 1-figure supplement 1. Calcitonin-type (CT-type) neuropeptide precursors in *A. japonicus*.
(A) *AjCTP1* comprises two putative CT-type neuropeptides, AjCT1 and AjCT2, whereas *AjCTP1*
generated by alternative splicing comprises only AjCT2. The predicted signal peptide is shown in
blue, the predicted cleavage sites (KR or KK) are shown in green, CT-type neuropeptides are
shown in red and C-terminal glycines that are potential substrates for amidation are shown in
orange. Predicted disulfide bridges in AjCT1 and AjCT2 are formed between the two underlined
cysteines. (B) Nucleotide sequences encoding CT-type neuropeptide precursors and the deduced
amino acid sequences in *A. japonicus*. 3'UTR and 5'UTR are shown in lowercase and the ORF is
shown in uppercase. Blue uppercase labels the predicted signal peptide, the green uppercase
shows the predicted cleavage site, AjCT1 and AjCT2 are shown in red and the putative substrate
for amidation is shown in orange.

1264

1265 Figure 1-figure supplement 2. Comparison of the sequences CT-type neuropeptides from Bilateria:

1266 deuterostomes-ambulacraria (green), deuterostomes-chordates (purple) and protostomes (pink).

1267 The conserved residues are shown with white lettering highlighted in gray and black. The

1268 accession numbers or citations are shown in the Supplementary table 1.

1269

A

AjCTR

B

AjPDFR1

1270

1271

1272
1273 Figure 2-figure supplement 1. Amino acid sequences and phosphorylation sites of AjCTR,
1274 AjPDFR1 and AjPDFR2 in *A. japonicus*. (A), (B) and (C) show the cDNA sequences encoding the
1275 receptors and their deduced amino acid sequences. ORFs are shown in uppercase; 3'UTR and
1276 5'UTR are shown in lowercase. The stop codon is shown by the asterisk (*). Predicted potential
1277 phosphorylation sites are shown with squares.

1278
Figure 2-figure supplement 2. Characterization of AjCT1 and AjCT2 receptors (AjCTR, AjPDFR1
and AjPDFR2). (A) Conserved domains of CTR/CLR-type protein sequences. Black line
represents the length of proteins and different domains are represented by rectangles. (B)
Conserved domains of PDFRs in Bilateria. Black line represents the length of proteins and
different domains are represented by rectangles. (C) Synteny analysis of AjCT1 and AjCT2
receptors (AjCTR, AjPDFR1 and AjPDFR2). The chromosomal locations of AjCT1 and AjCT2
receptor genes from *A. rubens*, *L. variegatus* and *A. japonicus* are shown, with different colors
used to mark neighboring genes and with their transcriptional direction shown by arrows.
GOLGA4, golgin subfamily A member 4; *NeuroD*, neurogenic differentiation factor; *CRF2*,
corticotropin-releasing factor receptor 2; *CTR*, calcitonin type receptor; *CLR*, calcitonin gene-
related peptide type 1 receptor; *ppGalNAc-T*, polypeptide N-acetylgalactosaminyltransferase;
URB3, E3 ubiquitin-protein ligase ubr3; *SLC40A1*, solute carrier family 40 member 1; *CMAH*,
cytidine monophosphate-N-acetylneuraminic acid hydroxylase; *MAGP*, microfibril-associated
glycoprotein; *HCN*, hyperpolarization-gated and cyclic nucleotide regulated channel; *ATG9A*,
autophagy-related protein 9A-like; *lysRS*, lysine--tRNA ligase; *FCN*, ficolin; *PDFR1*, pigment-
dispersing factor receptor 1; *PDFR2*, pigment-dispersing factor receptor 2; *PSMD8*, 26S
proteasome regulatory subunit 8; *SARNP*, SAP domain-containing ribonucleoprotein; *HPM1*,
histidine protein methyltransferase 1; *PNPO*, pyridoxine-5'-phosphate oxidase.
1297
1298

1299

1300 Figure 4-figure supplement 1. Negative control for analysis of AjCTR, AjPDFR1 and AjPDFR2

1301 internalization. The images show the localization of AjCTR, AjPDFR1 and AjPDFR2 in

1302 HEK293T cells without AjCT1 or AjCT2 stimulation. Green fluorescence represents the

1303 localization of pEGFP-N1/receptors in cell membrane. Cell nucleus probe (DAPI) was applied for

1304 cell nucleus staining. All pictures are representative of three independent experiments.

1305

1306

Figure 5-figure supplement 1. Effects of PKA or PKC inhibitor on AjCT1 or AjCT2 stimulated ERK1/2 phosphorylation in AjCTR, AjPDFR1 or AjPDFR2 expressing HEK293T cells. (A) ERK1/2 phosphorylation, activated by AjCT1, is blocked by PKC inhibitor Gö 6983 in AjCTR expressing HEK293T cells and is blocked by PKA inhibitor H89 and PKC inhibitor in AjPDFR1 expressing HEK293T cells. (B) ERK1/2 phosphorylation, activated by AjCT2, is blocked by PKC inhibitor in AjCTR expressing HEK293T cells and is blocked by PKA inhibitor and PKC inhibitor in AjPDFR1 and AjPDFR2 expressing HEK293T cells. Serum-starved HEK293 cells were pre-treated with DMSO, PKA inhibitor, or PKC inhibitor before AjCT1 or AjCT2 stimulation. All images are representative of three independent experiments. Error bars represent SEM for three independent experiments. Statistically significant differences are indicated as follows: *: $p < 0.05$, ns: no significant.

1318

Figure 7–figure supplement 1. Immunostaining in the radial nerve cord of *A. japonicus* is abolised

by pre-absorption of ArCT antiserum with the ArCT antigen peptide. (A) Transverse section of a

juvenile specimen of *A. japonicus* showing immunostaining in the radial nerve cord revealed by

an antiserum to the starfish calcitonin-type neuropeptide ArCT, as also shown in Figure 7A. (B) A

transverse section neighbouring the section shown in A was incubated with ArCT antiserum that

had been pre-absorbed with the ArCT antigen peptide; note the absence of immunostaining in the

radial nerve cord, demonstrating the specificity of the immunostaining in A. Abbreviations: LM,

longitudinal muscle; RNC, radial nerve cord. Scale is 100 µm.

1327

Figure 8-figure supplement 1. Positive control on experimental tissues and pharmacological
effects of 10^{-6} M AjCT2 on longitudinal muscle from *A. japonicus*. (A) shows a recording of the
relaxing effect of AjCT2 (10^{-6} M) on a longitudinal muscle preparation. (B) and (C) respectively
are recordings of the contracting effects of ACh on longitudinal muscle and intestine preparations,
showing that the effect of ACh persists for at least 15 minutes. All recordings shown are
representative of three independent experiments.

1334

Figure 9A-figure supplement 1. Measurement of the mass of remaining bait and excrement for
AjCT2-treated and control animals. (A) Remaining bait mass in different phases. (B) The mass of
excrement in different phases. The 24-day experiment was divided into six phases (I-VI), with
each phase comprising four days for statistical analysis. CO: control group (sterilized seawater);
CT2L: AjCT2 low concentration group (5×10^{-3} mg/mL); CT2H: AjCT2 high concentration group

(5×10^{-1} mg/mL). Values are means ± SEM (n = 3, representing three parallel experiments of each

group). Asterisks represent the significant difference as follows: *: $p < 0.05$, **: $p < 0.01$.

1342

Figure 10-figure supplement 1. The relative expression level of *AjCTP1/2* in siNC, si*AjCTP1/2*-1 and si*AjCTP1/2*-2. siNC: negative control group; si*AjCTP1/2*-1 and si*AjCTP1/2*-2: experimental groups targeting *AjCTP1/2*. Values are means ± SEM (n = 5). Asterisks represent the significant differences as follows: *: $p < 0.05$, **: $p < 0.01$.

Supplementary tables

Supplementary table 1. Accession numbers or citations of calcitonin-type family peptides or precursors in Bilateria

Abbreviation	Species Names	Accession Numbers And/or Citations
H_sapCTP	*Homo sapiens*	GI:179819
H_sapCGRPP	*Homo sapiens*	NP_001029125.1
T_rubCTP	*Takifugu rubripes*	GenBank: CAC81278.1
T_rubCGRPP	*Takifugu rubripes*	GenBank: CAC81277.1
C_intCTP	*Ciona intestinalis*	GenBank: BAI63095.1
B_floCTP1	*Branchiostome floridae*	GI:260826569
B_floCTP2	*Branchiostome floridae*	GI:260826567
B_floCTP3	*Branchiostome floridae*	GI:260826573
S_kowCTP	*Saccoglossus kowalevskii*	GI:187217193
A_japCTP1	*Apostichopus japonicus*	GenBank: AWU78769.1
A_japCTP2	*Apostichopus japonicus*	GenBank: AWU78770.1
A_rubCTP	*Asterias rubens*	GenBank: KT601715.1
P_pecCTP	*Patiria pectinifera*	GenBank: AWX63418.1
A_plaCTP	*Acanthaster planci*	Smith et al. (2017)
S_purCTP	*Strongylocentrotus purpuratus*	GI:115767208
O_vicCTP1	*Ophionotus. victoriae*	GenBank: ASK86238.1
O_vicCTP2	*Ophionotus. victoriae*	GenBank: ASK86238.1
O_araCTP1	*Ophiopsila aranea*	
O_araCTP2	*Ophiopsila aranea*	Zandawala et al. (2017)
A_filCTP1	*Amphiura filiformis*	
A_filCTP2	*Amphiura filiformis*	
H_scaCTP1	*Holothuria scabra*	Suwansa-Ard et al. (2018)

H_scaCTP2	Holothuria scabra	
H_glaCTP1	Holothuria glaberrima	
H_glaCTP2	Holothuria glaberrima	
A_medCTP	Antedon mediterranea	Aleotti et al., (2022)
An_japCTP	Anneissia japonica	
L_gigCTP	Lottia gigantea	GI:163526287
D_melDH31P	Drosophila melanogaster	NP_523514.1
L_migDH31P	Locusta migratoria	GenBank: AKN21237.1
C_telCTP	Capitella teleta	GI: 161220966
C.telDH31P	Capitella teleta	Rowe et al. (2014)

1351

1352   Supplementary table 2. Accession numbers of CT-type, PDF-type and CRH-type receptor families
1353   in Bilateria

Species	Protein ID	Gene code	Species code
Amphiura filiformis	XP_072029112.1	CRHR	Afil
Amphiura filiformis	XP_072021544.1	CTR	Afil
Amphiura filiformis	XP_072029557.1	PDFR1	Afil
Amphiura filiformis	XP_072028436.1	PDFR2	Afil
Anneissia japonica	XP_033108182.1	CRHR	Anja
Anneissia japonica	XP_033104142.1	CTR	Anja
Anneissia japonica	XP_033104113.1	PDFR	Anja
Apis mellifera	XP_006564466.1	DH31R	Amel
Apis mellifera	XP_016768833.2	PDFR	Amel
Aplysia californica	XP_012938918.2	CRHR	Acal
Aplysia californica	XP_005089839.2	CTR	Acal
Aplysia californica	XP_005093780.3	DH31R	Acal
Apostichopus japonicus	XP_071811976.1	CRHR	Ajap
Apostichopus japonicus	PIK36177.1	CRHR	Ajap
Apostichopus japonicus	XP_071811994.1	CRHR	Ajap
Apostichopus japonicus	PIK41405.1	CRHR	Ajap
Apostichopus japonicus	XP_071812618.1	CTR	Ajap
Apostichopus japonicus	XP_071811731.1	PDFR1	Ajap
Apostichopus japonicus	XP_071812243.1	PDFR2	Ajap
Asterias rubens	ENSASRP00000008640.1	CRHR	Arub
Asterias rubens	ENSASRP00000013963.1	CRHR	Arub
Asterias rubens	ENSASRP00000014061.1	CRHR	Arub
Asterias rubens	ENSASRP00000011186.1	CTR	Arub
Asterias rubens	ENSASRP00000010309.1	PDFR1	Arub
Asterias rubens	ENSASRP00000010166.1	PDFR2	Arub
Bombus impatiens	ENSBIMP00000009704.1	CRHR	Bimp
Bombus impatiens	ENSBIMP00000023653.1	DH31R	Bimp
Bombus impatiens	ENSBIMP00000026387.1	PDFR	Bimp
Branchiostoma floridae	XP_035687893.1	CRHR	Bflo
Branchiostoma floridae	XP_035687894.1	CRHR	Bflo
Branchiostoma floridae	XP_035689670.1	CRHR	Bflo
Branchiostoma floridae	XP_035689671.1	CRHR	Bflo
Branchiostoma floridae	XP_035687895.1	CTR*	Bflo
Caenorhabditis elegans	NP_001122749.1	SEB2*	Cele
Ciona intestinalis	XP_002122381.1	CRHR	Cint
Ciona intestinalis	XP_009858716.1	CRHR	Cint
Drosophila melanogaster	NP_725278.1	DH31R*	Dmel

Drosophila melanogaster	NP_001285174.1	HECTOR	Dmel
Drosophila melanogaster	CAB72288.1	PDFR	Dmel
Gallus gallus	ENSGALP00010018855.1	CLR*	Ggal
Gallus gallus	ENSGALP00010039362.1	CRHR	Ggal
Gallus gallus	ENSGALP00010039367.1	CRHR	Ggal
Gallus gallus	ENSGALP00010033414.1	CRHR	Ggal
Gallus gallus	ENSGALP00010004560.1	CTR*	Ggal
Gekko gecko	ENSEGEP00000030211.1	CLR	Ggec
Gekko gecko	ENSEGEP00000005175.1	CRHR	Ggec
Gekko gecko	ENSEGEP00000009950.1	CRHR	Ggec
Gekko gecko	ENSEGEP00000001898.1	CTR	Ggec
Homo sapiens	ENST04850112052.1	CLR*	Hsap
Homo sapiens	ENST04850047995.1	CRHR	Hsap
Homo sapiens	ENST04850093643.1	CRHR	Hsap
Homo sapiens	ENST04850217190.1	CTR*	Hsap
Hypsibius exemplaris	OQV19025.1	CRHR	Hexe
Hypsibius exemplaris	OQV18662.1	CRHR	Hexe
Hypsibius exemplaris	OWA52582.1	CTR	Hexe
Hypsibius exemplaris	OWA52579.1	DH31Ra	Hexe
Hypsibius exemplaris	OQV24416.1	DH31Rb	Hexe
Hypsibius exemplaris	OQV13187.1	PDFR	Hexe
Lepisosteus oculatus	XP_015199664.1	CLRa	Locu
Lepisosteus oculatus	XP_015214520.1	CLRb	Locu
Lepisosteus oculatus	XP_015209811.1	CRHR	Locu
Lepisosteus oculatus	XP_015217722.1	CRHR	Locu
Lepisosteus oculatus	XP_006636078.2	CTR	Locu
Limulus polyphemus	XP_013771881.1	CRHR	Lpol
Limulus polyphemus	XP_022240360.1	CRHR	Lpol
Limulus polyphemus	XP_013775447.1	CRHR	Lpol
Limulus polyphemus	XP_013791145.1	CRHR	Lpol
Limulus polyphemus	XP_013791446.1	CRHR	Lpol
Limulus polyphemus	XP_022240480.1	CTRa	Lpol
Limulus polyphemus	XP_022240470.1	CTRb	Lpol
Limulus polyphemus	XP_022254242.1	CTRc	Lpol
Limulus polyphemus	XP_022237602.1	CTRd	Lpol
Limulus polyphemus	XP_022246293.1	DH31Ra	Lpol
Limulus polyphemus	XP_022235154.1	DH31Rb	Lpol
Limulus polyphemus	XP_022235431.1	DH31Rc	Lpol
Limulus polyphemus	XP_022239928.1	DH31Rd	Lpol
Limulus polyphemus	XP_022248891.1	DH31Re	Lpol
Limulus polyphemus	XP_022255301.1	PDFR	Lpol
Lineus longissimus	ENSLLNP00015009405.1	CRHR	Llon
Lineus longissimus	ENSLLNP00015009422.1	CRHR	Llon
Lineus longissimus	ENSLLNP00015009434.1	CRHR	Llon
Lineus longissimus	ENSLLNP00015009638.1	CTRa	Llon
Lineus longissimus	ENSLLNP00015008640.1	CTRb	Llon
Lineus longissimus	ENSLLNP00015012550.1	CTRc	Llon
Lineus longissimus	ENSLLNP00015013093.1	DH31R	Llon
Lineus longissimus	ENSLLNP00015009635.1	PDFR	Llon
Lumbricus terrestris	ENSVPPP00000022107.1	CTRa	Lter

Lumbricus terrestris	ENSVPPP00000011858.1	CTRb	Lter
Lumbricus terrestris	ENSVPPP00000018690.1	CTRc	Lter
Lumbricus terrestris	ENSVPPP00000033159.1	CTRd	Lter
Lumbricus terrestris	ENSVPPP00000030225.1	DH31Ra	Lter
Lumbricus terrestris	ENSVPPP00000022149.1	DH31Rb	Lter
Lumbricus terrestris	ENSVPPP00000003533.1	PDFR	Lter
Magallana gigas	NP_001292279.1	CTR1*	Mgig
Magallana gigas	XP_034310675.2	CTR2*	Mgig
Magallana gigas	XP_011442634.2	CTR3	Mgig
Magallana gigas	XP_011442636.3	CTR4	Mgig
Magallana gigas	XP_065929766.1	CTR5	Mgig
Magallana gigas	XP_065923133.1	CTR6	Mgig
Magallana gigas	XP_052684037.1	CTR7	Mgig
Magallana gigas	XP_034317823.1	PDFR	Mgig
Magallana gigas	XP_034310400.1		Mgig
Owenia fusiformis	CAC9666549.1	CRHR	Ofus
Owenia fusiformis	CAC9666335.1	CTRa	Ofus
Owenia fusiformis	CAC9673671.1	CTRb	Ofus
Owenia fusiformis	CAC9666125.1	DH31R	Ofus
Owenia fusiformis	CAC9578850.1	PDFR	Ofus
Paramacrobiotus metropolita	XP_055352663.1	CRHR	Pmet
Paramacrobiotus metropolita	XP_055352664.1	CRHR	Pmet
Paramacrobiotus metropolita	XP_055329231.1	CTR	Pmet
Paramacrobiotus metropolita	XP_055329233.1	DH31R	Pmet
Paramacrobiotus metropolita	XP_055331530.1	PDFR	Pmet
Pecten maximus	XP_033762144.1	CRHR	Pmax
Pecten maximus	XP_033762006.1	CTRa	Pmax
Pecten maximus	XP_033762916.1	CTRb	Pmax
Pecten maximus	XP_033762874.1	CTRc	Pmax
Pecten maximus	XP_033762263.1	CTRd	Pmax
Pecten maximus	XP_033762155.1	DH31Ra	Pmax
Pecten maximus	XP_033762946.1	DH31Rb	Pmax
Pecten maximus	XP_033729115.1	DH31Rc	Pmax
Pecten maximus	XP_033762294.1	PDFR	Pmax
Platynereis dumerilii	AKQ63006.1	DH31R-1*	Pdum
Platynereis dumerilii	WOE99345.1	PDFR	Pdum
Priapulus caudatus	XP_014664111.1	DH31Ra	Pcau
Priapulus caudatus	XP_014661438.1	DH31Rb	Pcau
Rotaria socialis	CAF3356201.1	PDFRa	Rsoc
Rotaria socialis	CAF3546535.1	PDFRb	Rsoc
Saccoglossus kowalevskii	NP_001161520.1	CRHR	Skow
Saccoglossus kowalevskii	XP_006818813.1	CTR	Skow
Saccoglossus kowalevskii	XP_006825104.1	PDFRa	Skow
Saccoglossus kowalevskii	XP_006825103.1	PDFRb	Skow
Schmidtea mediterranea	SMEST051050001	CRHR	Smed
Schmidtea mediterranea	SMEST064388001	CRHR	Smed
Schmidtea mediterranea	SMEST049048003	CRHR	Smed
Schmidtea mediterranea	SMEST080399003	CTRa	Smed
Schmidtea mediterranea	SMEST042095002	CTRb	Smed
Schmidtea mediterranea	SMEST006966001	DH31Ra	Smed

Schmidtea mediterranea	SMEST027375002	DH31Rb	Smed
Schmidtea mediterranea	SMEST023013001	DH31Rc	Smed
Schmidtea mediterranea	SMEST035956002	PDFR	Smed
Sepioteuthis lessoniana	ENSUKOP00000023564.1	CRHR	Sles
Sepioteuthis lessoniana	ENSUKOP00000009609.1	CTRa	Sles
Sepioteuthis lessoniana	ENSUKOP00000023956.1	CTRc	Sles
Sepioteuthis lessoniana	ENSUKOP00000023961.1	CTRd	Sles
Sepioteuthis lessoniana	ENSUKOP00000005365.1	DH31Ra	Sles
Sepioteuthis lessoniana	ENSUKOP00000007960.1	DH31Rb	Sles
Sepioteuthis lessoniana	ENSUKOP00000030254.1	PDFR	Sles
Strongylocentrotus purpuratus	XP_003725174.2	CRHR	Spur
Strongylocentrotus purpuratus	XP_030845739.1	CRHR	Spur
Strongylocentrotus purpuratus	XP_030846612.1	CRHR	Spur
Strongylocentrotus purpuratus	XP_030846621.1	CTR	Spur
Strongylocentrotus purpuratus	XP_030845952.1	PDFR1	Spur
Strongylocentrotus purpuratus	XP_030845953.1	PDFR2	Spur
Styela clava	XP_039255657.1	CRHR	Scla
Tribolium castaneum	XP_008198347.1	CRHR	Tcas
Tribolium castaneum	XP_008192711.1	CRHR	Tcas
Tribolium castaneum	XP_064212169.1	CTR	Tcas
Tribolium castaneum	XP_008193920.1	DH31Ra	Tcas
Tribolium castaneum	XP_064210831.1	DH31Rb	Tcas
Tribolium castaneum	XP_064212793.1	PDFR	Tcas
Xenopus tropicalis	ENSXETP00000114756.1	CLR	Xtro
Xenopus tropicalis	ENSXETP00000109962.1	CRHR	Xtro
Xenopus tropicalis	ENSXETP00000101014.1	CRHR	Xtro
Xenopus tropicalis	ENSXETP00000060302.3	CTR	Xtro

1354

A

B

A

B