Determining the off-target activity of antibiotics and novel translation initiation sites in mitochondria

James Marks[1,#,*], Emma Young[1,**], Markus Hafner[1,#]

[1]RNA Molecular Biology Laboratory, National Institute for Arthritis and Musculoskeletal and Skin Diseases, NIH, Bethesda, MD 20892

Current address:

*Department of Biochemistry, Vanderbilt University School of Medicine, Nashville, TN 37237

**Department of Biological Anthropology, University of California, Los Angeles, CA 90095

[#]Corresponding authors: james.marks@vanderbilt.edu, markus.hafner@nih.gov

Abstract

The 13 mtDNA-encoded proteins are synthesized using a dedicated translation system that is more similar to bacterial systems than the cytoplasmic system. Consequently, many bacterial protein synthesis inhibitors, used as antibiotics, exhibit mitochondrial toxicity as off-target effects. However, whether these antibiotics act through the same mechanisms in mitochondria as in bacteria remains unclear. To address this, we characterized the impact of a panel of bacterial translation initiation and elongation inhibitors on mitochondrial translation through mitoribosome profiling. We found that the mechanism of action for every antibiotic, except telithromycin, is the same in both bacteria and mitochondria. Additionally, mitoribosome profiling data showed that *MT-ND1* and *MT-ND5* have incorrectly annotated translation initiation sites and suggested the presence of several translation initiation sites on ncRNAs that produce mitoribosome footprints, as indicated by the detection of mitoribosome footprints at these locations. This work demonstrates how antibiotics can inhibit mitochondrial translation by mechanisms identically or very similar to those found in bacteria and the utility of mitoribosome profiling for annotating mitochondrial genes.

Introduction

Mitochondria contain their own genome, which canonically encodes thirteen proteins. These proteins are essential building blocks of complexes I, III, and IV of the electron transport chain, as well as ATP synthetase. They are synthesized at the mitochondrial matrix by a dedicated translation system that is conserved from the ancestral bacteria that evolved into the present-day organelles through an endosymbiotic event[1]. As a result, mitochondrial translation is wholly distinct from cytoplasmic translation and utilizes its own mitochondrial ribosomes (mitoribosomes), tRNAs, and translation factors. The RNA components of the machinery are mitochondrially encoded, while its protein components are encoded in the nucleus, translated by the cytoplasmic ribosome, and subsequently imported into the mitochondria[2].

Chronic impairment of mitochondrial translation, most commonly through mutations in translation factors, results in disrupted mitochondrial function and causes diseases such as MERRF (Myoclonic Epilepsy with Ragged-Red Fibers)[3], MELAS (Mitochondrial Encephalopathy, Lactic Acidosis, and Stroke-like episodes)[4], and maternally inherited deafness[5,6]. Furthermore, acute inhibition of mitochondrial translation caused by the off-target action of antibiotics targeting the bacterial ribosome can also cause mitochondrial toxicity[7-11]. This toxicity can manifest as lactic acidosis, ototoxicity[12], nephrotoxicity[13], and peripheral neuropathy[14], depending on the specific antibiotic and the genetic background of the recipient[15]. While the United States has worked to regulate antibiotics with severe side effects, they remain a global issue. For example, chloramphenicol, whose use is discontinued in the United States, causes fatal aplastic anemia at a rate of 1 in 60,000[7] but remains widely prescribed in low-income countries due to its low production cost[16,17]. Surveys suggest that approximately 14.3 doses of antibiotics are administered per 1,000 people every day worldwide[18]. The toxic effects of antibiotics can be exacerbated when taken without medical supervision. In addition, safety concerns surrounding drugs like chloramphenicol and telithromycin, which are linked to liver failure[19], limit our pool of available antibiotics in the face of a rise in multidrug-resistant bacteria. Studying the off-target effects of these drugs can guide the development of safer therapies.

Multiple classes of antibiotics inhibit bacterial translation by binding functionally important sites on the ribosome and disrupting a specific process in translation, resulting in arrested translation on mRNAs at sites characteristic of the mechanism of inhibition[20,21]. In some instances, this relationship is straightforward. For example, the pleuromutilin antibiotic retapamulin binds to the peptidyl-transferase center (PTC) of the bacterial ribosome and impedes translation initiation

by interfering with the positioning of initiator tRNA[22], resulting in bacterial ribosomes accumulating

at start codons[23]. Inhibition by other antibiotics is more complicated, only disrupting translation

when specific sequences are being translated, called context-dependent translation arrest. This

is exemplified by chloramphenicol, which simultaneously interacts with the nascent peptide and

the peptidyl-transferase center at its binding site and disrupts accommodation of aminoacyl-tRNA,

thereby acting as an elongation inhibitor[24]. Due to chloramphenicol's interaction with the nascent

peptide, it selectively causes ribosomes to arrest translation when an alanine, serine, or threonine

is in the −1 (penultimate) position of the nascent peptide[25]. Additionally, chloramphenicol does not

interfere with the accommodation of glycine due to its minimal size. The mechanisms of these

and other antibiotics in bacteria have been determined by ribosome profiling and structural

studies. As well, structural studies have demonstrated the binding characteristics of antibiotics on

the mitochondrial ribosome[11,26,27]. However, it remains unclear whether chloramphenicol or other

antibiotics inhibit mitochondrial translation by identical mechanisms.

To determine the sites of translation arrest, we utilized mitoribosome profiling in HEK293

cells treated with a diverse panel of antibiotics[19,28,29]. Ribosome profiling allows for the mapping

of translating ribosomes with codon-level accuracy by identifying mRNA sequences occupied by

translating ribosomes. To do this, RNase treatment degrades mRNA unprotected by the ribosome

to generate ribosome-protected fragments (RPFs). Because cytoplasmic ribosomes significantly

outnumber mitoribosomes, mitoribosomes containing the mitochondrial RPFs (MRPFs) are

isolated through sucrose gradient fractionation, where the cytoplasmic ribosome and

mitoribosome sediment in the 80S and 55S fractions, respectively (Figure 1A). RPFs are isolated

and quantified by next-generation sequencing. Increases in MRPFs reveal the specific codons

and sequence contexts prevalent at sites at which mitochondrial translation arrests due to

antibiotic treatment.

Understanding the mechanisms of translation inhibitors in mitochondria can guide the

development of therapeutics that are more selective to the bacterial ribosome and lay the

groundwork for determining how these therapeutics disrupt mitochondrial function. In this study,

we utilized mitoribosome profiling in the presence of the bacterial protein synthesis inhibitors

retapamulin, tiamulin, josamycin, erythromycin, chloramphenicol, and linezolid to determine their

mechanism of action. We found that the mechanism of mitochondrial inhibition is the same as

bacterial inhibition for all inhibitors except telithromycin. While telithromycin selectively arrests

bacterial ribosomes when either arginine (R) or lysine (K) is in the -1 position of the nascent chain

and A-site, called an R/K-X-R/K motif[30], it instead arrests mitochondrial translation at an R/K/A-

X-K motif, suggesting an underlying difference in the mechanism of the two systems. Additionally,

nucleotide-level precision of mitoribosome profiling also allowed us to precisely annotate the

mitochondrial gene structure. We found that *MT-ND1* and *MT-ND5* genes use alternative

translation initiation sites and identified a putative open reading frame on the *MT-RNR1* gene.

This work demonstrates that mitoribosome profiling is an effective tool to study the fundamentals

of mitochondrial translation and the mechanisms of translation impairment affecting human

health.

Results

RNase I generates highly phased mitoribosome footprints

To optimize the mitoribosome profiling method, we tested two different RNases typically used for ribosome footprinting, RNase I and MNase, followed by the isolation of the 55S mitoribosome by sucrose gradient fractionation. We mapped the sequence reads to the mitochondrial genome and its thirteen open reading frames (ORFs). For each mapped sequence read, we inferred the position of the ribosomal P-site using a 13-nucleotide (nt) offset from the 5' end, determined by the analysis of footprint distribution surrounding the initiation sites of *MT-ND4*, *MT-ND6*, and the newly reported *MT-ND5-dORF*. We observed coverage across mitochondrial ORFs, suggesting we captured translating mitoribosomes (Figure 1B). Furthermore, analysis of MRPF lengths revealed a distinct difference between the RNase I and MNase-digested footprint. While both libraries are enriched for reads ~30 nt in length, MNase footprints showed a bimodal distribution, with a clear secondary peak at ~22 nts (Figure 1C), which may indicate that MNase-treatment may have captured two distinct conformations of the ribosome. To determine the quality of the ribosome profiling data, we calculated read phasing, which occurs when a nuclease creates uniform MRPFs and the resulting ends of the MRPFs reflect the 3-nt periodicity of elongating ribosomes. While both datasets do show clear phasing, the read phasing is more biased toward position 1 of the codon in the RNase I-treated sample, indicating slightly more consistent digestion (Figure 1D, Table S1). Nevertheless, we observed similar levels of footprint abundance per gene, with the only notable exceptions being that RNase I-digested footprints were enriched on *MT-ND3* and depleted on *MT-ND4L* and *MT-ND6* (Figure 1E). While the cause of the difference is unclear, we did not expect it to interfere when comparing an antibiotic-treated sample to a control. Read coverage also appeared similar across the two methods, with 85%-87% codons having between 2 and 50 reads mapping to them (Figure 1F). Overall, these results indicated that both MNase and RNase I digestion capture mitoribosome footprints with only minor differences. Considering the improved read phasing, we selected RNase I for our subsequent mitoribosome profiling experiments.

143
Figure 1. Mitoribosome profiling captures elongating ribosomes.

(A) Schematic representation of mitoribosome footprinting and separation from cytoplasmic ribosomes by differential centrifugation. (B) Coverage of MRPFs on *MT-CYB* from mitoribosome profiling generated by either RNase I or MNase footprinting. (C) Distribution of read lengths of MRPFs mapping to the mitochondrial genome. (D) Read phasing analysis of MRPFs, where the 5' ends of reads are grouped by their subcodon position. Each dot represents an individual mitochondrial gene. (E) Average coverage of mitochondrial genes, calculated by dividing the total number of reads mapping to an ORF and dividing by the length of the ORF. (F) Distribution of mitochondrial codons grouped by the number of mitoribosome P-sites assigned to them.

Retapamulin, tiamulin, and josamycin inhibit translation initiation

First, we examined the impact of three antibiotics – retapamulin, josamycin, and tiamulin – on translation initiation or mitochondrial translation. Retapamulin and tiamulin are members of the pleuromutilin class of antibiotics. They work by binding to the peptidyl transferase center (PTC) of the bacterial ribosome and disrupting initiator tRNA and A site tRNA positioning (Figure 2A, left)[22,23,31-33]. Josamycin is a macrolide antibiotic that binds in the nascent peptide exit tunnel near the PTC and arrests bacterial ribosomes at initiation by interfering with A-site tRNA positioning (Figure 2A, right)[34]. We sought to understand whether these antibiotics also target translation initiation in mitochondria. To this end, we treated HEK293 cells with high concentrations of these antibiotics (100 µg/ml of tiamulin and josamycin, 10 µg/mL of retapamulin) for 30 minutes, performed mitoribosome profiling, and compared the treated experiments to a DMSO-treated control.

We mapped MRPFs to the 13 mitochondrial genes. On *MT-CYB* and all other genes, we observed that after treatment with antibiotics, reads accumulated at the 5' end of the transcript and MRPFs across the rest of the coding region were lost or decreased (Figures 2B, S1). We quantified the percentage of MRPFs mapping at the 5' end relative to the total ORF, excluding polycistronic mRNAs. Treatment with translation initiation inhibitors increased the percentage of those reads from 2.4 to 5.2-fold relative to the DMSO-treated cells (Figure 2C). Additionally, we quantified the cumulative footprint abundance moving from the start to the stop codon (Figure 2D). While ribosome footprints are generally evenly spread across the open reading frame in the DMSO control, MRPFs from cells treated with the initiation inhibitors showed a sharp increase in the percentage of reads found at the 5' end. Together, these results indicate that retapamulin, tiamulin, and josamycin inhibit mitochondrial translation initiation, likely by the same mechanism as in prokaryotes.

Figure 2. Inhibition of translation initiation in mitochondria.

(A) (Left) Schematic representation of retapamulin (orange) or tiamulin (yellow) binding to the PTC and disrupting the first round of elongation. (Right) Josamycin (blue) disrupts accommodation of the first aminoacyl-tRNA. All three inhibit initiation in bacteria. (B) MRPF coverage of *MT-CYB* treated with either retapamulin, tiamulin (note y-axis is segmented), or josamycin. (C) Average fold change increase in the percentage of reads that map to the 5' of mitochondrial transcripts. (D) Cumulative distribution of ribosome density normalized across all mitochondrial ORFs.

Chloramphenicol, linezolid, and telithromycin arrest mitochondrial translation in a context-dependent manner

Chloramphenicol, linezolid, and telithromycin selectively inhibit bacterial translation when ribosomes are translating specific sequences, in a process called context-dependent translation arrest[25,30]. Linezolid, like chloramphenicol, induces context-dependent arrest when alanine, serine, or threonine is at the penultimate position of the nascent peptide (Figure 3A, left). Unlike chloramphenicol and linezolid, telithromycin binds in the nascent peptide exit tunnel near the PTC and adjacent to the nascent peptide[35]. During elongation, telithromycin causes ribosomes to arrest translation at an R/K-X-R/K motif, when the identity of both the penultimate position of the nascent peptide and the incoming aminoacyl-moiety is either arginine or lysine (Figure 3A, right).

We carried out mitoribosome profiling on HEK293 cells treated with either antibiotic to understand whether these context-dependent translation inhibitors arrest mitoribosomes in the same manner as they arrest bacterial ribosomes. Unlike the cells treated with initiation inhibitors, cells treated with chloramphenicol or linezolid carried ribosome density throughout the entirety of the ORF, suggesting that, as expected, they were not acting as inhibitors of initiation (Figures 3B, S1). Cells treated with telithromycin did show a slight increase in MRPFs abundance at the 5' end of *MT-CYB.* However, across all genes, the levels of MRPFs at the 5' end were not near the levels seen when using the initiation inhibitor retapamulin and were more similar to chloramphenicol, linezolid, and the DMSO control (Figure 3C). The cumulative distribution of MRPFs suggested that ribosome density was biased towards the 5' end of the gene for chloramphenicol and telithromycin, but not significantly for linezolid (Figure 3D). These results suggest that these three antibiotics are not acting as initiation inhibitors and that linezolid and telithromycin redistribute MRPFs to the 5' end of the ORF.

Next, we determined whether this observed redistribution of elongating ribosomes was dependent on the context of the sequence being translated. We calculated the average change of MRPFs depending on the identity of the amino acid at either the -1, P-site, or A-site positions. For both chloramphenicol and linezolid, we observed a specific increase in ribosome footprints when mitoribosomes encountered codons for alanine, serine, or threonine in the -1 position, with a 2-, 2.2-, and 1.3-fold change for chloramphenicol and a 2.3-, 1.4-, and 1.6-fold change for linezolid, respectively (Figure 3E). Additionally, we also observed context-dependent translation arrest induced by telithromycin. Unlike the R/K-X-R/K motif observed in bacteria, we found that telithromycin caused mitoribosomes to arrest at sites with a lysine positioned in the A-site, with an average 2.5-fold increase in MRPFs around those sites, and that neither residue was enriched

in the -1 position. Due to the limited size of the mitochondrial genome, there are only nine

instances of the R/K-X-R/K motif; nevertheless, when we observed the positions with the greatest

fold change, 3 of the top 4 contained the motif with lysine in the A-site (Figure 3F), suggesting

that the motif was at least partially recapitulated. To specifically look for the motif, we calculated

the average fold change when we fixed arginine or lysine in the A-site and varied the identity of

the -1 amino acid (Figure 3G). We found that when lysine was in the A-site while either arginine,

lysine, or alanine was in the -1 position, there was strong arrest (> 5-fold change). However, when

arginine was in the A-site, we did not observe a strong stalling event with any -1 position residue.

Therefore, in mitochondria, telithromycin induces stalling at R/K/A-X-K motifs. This motif appears

in only 6 of the 13 mitochondrially encoded genes, indicating that mitochondrial protein synthesis

is unequally inhibited (Figure S2A, B). We concluded that the mechanism underlying context-

dependent translation arrest for telithromycin is different in mitochondria compared to bacteria.

Figure 3. Context-dependent translation arrest by elongation inhibitors.

(A) Schematic representation of chloramphenicol (purple), linezolid (cyan), and telithromycin (green) inducing context-dependent translation arrest. (B) MRPF coverage of *MT-CYB* from cells treated with either chloramphenicol, linezolid, or telithromycin. (C) Average fold change increase in the percentage of reads that map to the 5' of mitochondrial transcripts. (D) Cumulative distribution of ribosome density normalized across all mitochondrial ORFs. (E) Volcano plots of changes in MRPF fold change from cells treated with translation inhibitors. Each dot is the mean $\log_2$-fold change grouped by the identity of the -1 amino acid (top), P-site amino acid (middle), or A-site amino acid (right). (F) The top 4 strongest stalling sites induced by telithromycin treatment. Red amino acids indicate a classical R/K-X-R/K stalling motif. (G) Fold change in ribosome density, calculated as $2^{\text{mean}(\log_2(\text{Telithromycin/Control}))}$. Bars differ in the identity of the P-site amino acid and contain either a lysine in the A-site (top) or arginine (bottom).

Translation initiation on *MT-ND1 and MT-ND5* occurs at an alternative start codon

During the process of assigning the P-site positions of MRPFs, we observed that no ribosomes
were assigned in the first 13 nucleotides of the 5' end of the transcript. This is not surprising,
considering that mitochondrial transcripts often have no or very short 5' untranslated regions -
typically less than 4 nts long - and we applied a 13-nt offset from the 5' end of MRPFs. It reasons
that as mitoribosomes begin translation on a leaderless mRNA, the 5' end of the resulting MRPF
would have a full-length region 3' of the P-site and no sequence 5' of the P-site. As a ribosome
moves down the transcript, the 5' end of the MRPF would increase until it is accessible for
digestion, at which point it would generate 31 nt-long MRPFs (Figure 4A). As accurate mapping
of mitoribosomes at initiation could be a useful tool to understand mitochondrial translation, we
attempted to map the location of initiating mitoribosomes at the 5' end using footprint length, rather
than the 5' offset. Because mitoribosome profiling with MNase generated a population of
ribosomes with short MRPFs, we hypothesized that these footprints could be indicative of
capturing initiating ribosomes (Figure 1C). We plotted the distribution of MRPFs from 20 to 34 nts
in length that mapped to the 5' end of the 11 leaderless or short leader ORFs and compared them
to reads mapping to the whole ORF. We observed that MRPFs 22-23 nts in length were 6-fold
more abundant and MRPFs >28 nts in length were depleted compared to the whole ORF (Figure
4B). As mitochondrial ORFs vary in the length of their 5' UTR and we expect most ribosomes to
be positioned at the start codon, we tested whether leaderless ORFs were enriched in shorter
reads compared to ORFs preceded by a short leader. Leaderless ORFs, such as *MT-CO2*, *MT-
CO3*, and *MT-ND4L,* contained MRPFs mainly 20-23 nucleotides in length at the 5' end (Figure
4C, top). *MT-ATP8*, which contains a start codon beginning at the third nt, contained footprints
that were 22-24 nt long, and *MT-CO1,* with a start codon at the fourth nt, contained MRPFs 23-
24 nt long (Figure 4C, middle). Not only did we observe longer MRPFs in short leader ORFs, but
the size of the MRPFs also appeared proportional to the location of the start codon. Together,
these results indicate that the size of the footprint is directly related to the start codon position at
the 5' end of mitochondrial mRNAs.

Two genes, *MT-ND*1 and *MT-ND5,* deviated from this trend. The annotated start site for
MT-ND5 is a leaderless AUA codon (a start codon in mitochondria), but we observed few 20-22
nt MRPFs (Figure 4C, bottom). Similarly, the annotated start site for *MT-ND1* is also an AUA
codon that begins at the third nt, and it is depleted in reads for MRPFs that are smaller than 25
nts. Instead, MRPFs derived from the 5' end of both genes are enriched in longer reads: 27 nts-
long for *MT-ND5* and 28-29 nts-long for *MT-ND1*. Curiously, the third codon for both genes is an

in-frame AUG, and the read lengths proportionally correspond to mitoribosomes positioned at

these codons, suggesting mitoribosomes initiate translation at these sites. MRPFs generated from

cells treated with initiation inhibitors and footprinted by MNase yield similarly sized reads at the 5'

end of *MT-ND1* and *MT-ND5* (Figure S3). Using protein mass spectrometry, we tested whether

peptides matching the N-terminus of these proteins would correspond to initiation at the canonical

start codons or alternative start codons. We identified a peptide that mapped to the N-terminal

region of MT-ND1 but lacked the first two amino acids, accordant with translation initiating from

the alternative start codon (Figure 4D). While we did not identify N-terminal peptides mapping to

MT-ND5 from our mass spectrometry data, the protein mass spectrometry database MassIVE[34]

covered the N-terminal region of MT-ND5 (Figure 4E). Consistent with our mitoribosome profiling

data, the majority of N-terminal fragments did not include the amino acids from the annotated start

or second codon but began with methionine at codon 3. We also observed that MRPFs on *MT-*

ND6, which initiates at the first of two AUG codons immediately at the 5' end, correlated to

ribosomes positioned at both AUG codons, indicating that this gene may initiate translation from

the second AUG codon as well. However, we did not observe any peptides mapping to the N-

terminus of *MT-ND6*. Together, these data show that *MT-ND1* and *MT-ND5* may not initiate from

their canonical AUA codons but may use alternative in-frame AUG codons. Additionally, our

findings indicate that the mapping of mitochondrial translation initiation events by size, rather than

5' offset, is an accurate strategy when used in conjunction with MNase footprinting.

304
Figure 4. MRPF sizes indicate use of alternative initiation sites.

(A) Schematic representation of changing MRPF sizes as a function of the mitoribosome's
position near the 3' end of mRNAs. (B) Read length measurement of reads mapping to the 5'
ends of mitochondrial transcripts (red) compared to the entire ORF (black). (C) Read lengths of
MRPFs mapping to the 5' ends of mitochondrial transcripts. Each row contains the nucleotide
sequence of each transcript starting from the 5' end. The coloring of the box is proportional
abundance of MRPFs with read lengths from 20-34 nts. Nucleotide sequence and read
abundance are overlayed to compare the abundance of reads to potential ribosome start sites.
Transcripts are grouped by being leaderless, having a leader, or having a potential alternative
start site. Black boxes indicate annotated start codons and green boxes indicate alternative
initiation sites. (D) Visual representation of the peptide coverage (green bars or highlights) from
mass spectrometry data covering the N-terminus of *MT-ND1* or (E) *MT-ND5.*
317

Mitoribosome profiling may reveal novel translation events

During our analysis of 5' end mapping, we did not remove reads mapping to non-coding RNAs (ncRNAs) before mapping to the mitochondrial genome and performing P-site assignment. Consequently, we observed that some tRNAs and rRNAs had an abundance of reads mapping to 13 nts away from their 5' ends, similarly to what we observe on mRNAs for translation initiation. In addition, some of these ncRNAs contained a potential start codon near the 5' end. These reads could either be products of random digestion of the abundant background of ncRNAs or be genuine MRPFs. We hypothesized that if these reads were the result of mitoribosomes initiating or attempting to initiate translation at the 5' end of the ncRNAs, then we could use the same analysis of read length distribution at 5' ends of mRNAs we performed above and determine of the sizes correlated to an AUG, AUU, GUG, or UUG as potential start codons. While GUG and UUG are not used in canonical mitochondrial initiation, they were included in the search because noncanonical initiation has been observed at GUG[36] , and both are frequently utilized in bacterial initiation. We identified sites on four tRNAs that contained MRPFs with a size that correlated to either AUG or UUG codons (Figure 5A). The predicted ORFs and thus resulting potential translation events varied widely. On *MT-TF*, the start codon is followed by an immediate stop codon. The ORF on *MT-TI* is three codons long and terminates with the non-canonical mitochondrial termination codon AGA. On *MT-TY, a* 7-codon ORF is terminated by a UAA codon. *MT-TS1* contains the largest open reading frame identified on tRNAs, at 20 codons, and it terminates with the U AGA sequence, as used in *MT-CO1*[37]. Additionally, we identified a putative start site at the 5' end of *MT-RNR1* (Figure 5A), and mitoribosome profiling of cells treated with initiation inhibitors and digested with MNase shows similar footprint sizes (Figure S3)*.* However, this site did not occur at a canonical AUG or AUU codon, but instead at a UUG codon. This putative start codon is followed by a 58-codon ORF and terminates at a UAG stop codon.

To determine whether translation elongation occurs at these putative ORFs, we determined the read phasing characteristic for MRPFs derived from translating ribosomes at these ORFs. However, the sites on *MT-TF*, *MT-TI*, and *MT-TY* are short and close to the 5' end, hindering phasing analysis and thus do not allow us to conclude whether our sequence reads are indicative of translation. While the ORF on *MT-TS1* is longer, MRPF density was low, and we did not observe read phasing and thus it is likely not translated (not shown). The reads mapping to the ORF on *MT-RNR1* had a phasing profile similar to those mapping to the 13 canonical mitochondrial genes (Figure 5B). In contrast, reads mapping to the remainder of *MT-RNR1* or mapping to the region of *MT*-RNR2 that codes for the putative mitochondrial micropeptide

Humanin show no or conflicting read phasing. When compared to the amino acid composition of

the mitochondrially-encoded proteome, the amino acid composition of the resulting peptide shows

a stronger positive correlation (R^2=0.534) (Figure 5C) to the remaining non-coding portion of *MT-*

RNR1 (R^2=0.241) (Figure 5D). Although mitoribosome profiling suggests that this putative ORF

is translated, we did not detect any corresponding peptides in the protein mass spectrometry of

isolated mitochondria. Nevertheless, our mass spectrometry dataset, or the MASSIVE database,

does not cover many regions of the mitochondrial proteome, and thus, the absence of a spectrum

is not necessarily evidence that the peptide is not translated. The read phasing analysis suggests

this region of *MT-RNR1* is translated and could encode a 15[th] mitochondrial protein. Overall, this

data shows that our mitoribosome profiling can be used as a tool to study non-canonical

translation events resulting from ncRNAs.

363
364
Figure 5. MRPFs on ncRNAs indicate potential novel ORFs

(A) Read lengths of MRPFs mapping to the 5' end of ncRNAs. Each row contains the nucleotide
sequence beginning from the 5' end of listed ncRNAs. The coloring of the box is proportional the
abundance of MRPFs with read lengths 20-34 nts. The nucleotide sequence and read MRPF
abundance are overlayed to identify potential new start sites. Putative start sites are indicated by
green boxes. (B) Read phasing analysis of all canonical ORFs (grey), a mutative ORF located on
MT-RNR1, the non-coding region of *MT-RNR1,* and Humanin. (C) A correlation plot comparing
the frequence of amino acid usage in canonical ORFs compared to the putative ORF on *MT-*
RNR1 or (D) the non-coding region of *MT-RNR1.*

Discussion

In this work, we sought to understand the mechanisms of bacterial protein synthesis inhibitors on their primary off-target, the mitoribosome. We found that inhibitors of bacterial translation initiation, retapamulin, tiamulin, and josamycin, also act as initiation inhibitors in mitochondria. Likewise, chloramphenicol and linezolid selectively inhibit elongation when the mitoribosome carries either an alanine, serine, or threonine in the penultimate position of the nascent peptide, matching their mechanism in bacteria. In contrast, the context-dependence of telithromycin differed when arresting mitoribosomes, wherein it preferentially arrested translation at the R/K/A-X-K motif.

Context-dependent translation arrest by chloramphenicol and linezolid suggests similarities between the binding of the bacterial ribosome and the mitoribosome. Structural and biochemical studies have demonstrated that binding of chloramphenicol and linezolid is stabilized by alanine, serine, or threonine in the penultimate position of nascent peptide[38,39]. Our findings suggest a similar or identical interaction is occurring in the mitoribosome. In contrast, these inhibitors are unable to inhibit peptide bond formation in bacteria when glycine is positioned in the A-site, due to the minimal size of the side chain, allowing the inhibitor and the aminoacyl-moiety to coexist in the PTC. We did not observe a lack of mitoribosome translation arrest when glycine was positioned in the A-site. This may have technical reasons, e.g., it may be caused by an insufficient depth in sequencing or indicate that the relatively small mitochondrial genome limits the variety of stalling and non-stalling events we can observe. But more likely it shows that chloramphenicol and linezolid are capable of arresting translation when glycine is in the A-site. Recently, mitoribosome profiling analysis showed that chloramphenicol added during lysis did not induce context-dependent translation arrest[40]. This suggests that during lysis, there is insufficient mitochondrial translation to induce a redistribution of mitoribosomes rather than an inability of the chloramphenicol to induce context-dependent arrest.

During the preparation of this manuscript, two other groups studied the effect of translation inhibitors on mitochondrial translation. One group's mitoribosome profiling in the presence of chloramphenicol and linezolid agreed with our observed context-dependent translation arrest[41]. The structure of linezolid in the mitochondrial peptidyl-transferase center also suggested some nuanced differences in binding between the bacterial and the mitochondrial ribosome. Another group carried out mitoribosome profiling in the presence of retapamulin[40]. However, they observed additional ribosome density throughout the coding region and potentially some associated with putative start codons. This differs significantly from our observed accumulation at

only annotated start sites and may be due to differences in mitoribosome profiling methods.

Future work may reveal how each strategy contributes to our understanding of mitochondrial

translation.

Inhibition of mitochondrial translation through either small molecules or mutations

localized in mitochondrial translation factors can induce a wide range of pathologies[42-44]. However,

it is unclear how or whether the relationships between the mechanism of translation inhibition

relate to the resulting disease. Recently, the proteins mtRF-R and MTRES1 were shown to act

as quality control factors, rescuing stalled mitoribosomes by removing the nascent peptide and

peptidyl-tRNA[45]. In addition, treatment with antibiotics can reduce the presence of polyamines in

the mitoribosome, potentially reducing translation[46]. As the mitoribosome can arrest translation

through a variety of mechanisms and small molecules, it remains unclear whether these factors

can rescue all stalled mitoribosomes or decrease the translation rate to reduce the total amount

of translation arrest. Further work to dissect the relationship between mechanisms of inhibition,

activation of rescue factors, and stress response can complement our understanding of

pathologies resulting from mitoribosomopathies and antibiotic treatment.

In this study, we also reported that the translation start codon of two mitochondrial genes,

MT-ND1 and *MT-ND5*, does not use the annotated AUA codon, but instead begins translation

from a nearby in-frame AUG codon. Others have noted that haplogroups L1b and F2 contain a

single nucleotide polymorphism at either position T3308C or T12338C, respectively, which

disrupts the annotated AUA start codons of *MT-ND1* and *MT-ND5* yet does not result in any

discernible pathogenicity[47-49]. The authors noted that the same AUG codons we identified as start

codons could be used in lieu of the canonical AUA start codons. Because we observe MRPF

lengths primarily correlating with the AUG codons, we suggest that these downstream start

codons are the primary start site for these genes, though this does not preclude the use of both

sites. Given the limited examples of translation initiation sites inherent to the mitochondrial

genome, the reannotation of 2 sites is a significant revision and can guide future work to

understand how start codon selection occurs in mitochondria.

Recently, mitoribosome profiling was used to identify an open reading frame in region

immediately 3' of the gene *MT-ND5,* which we confirmed observing in our profiling experiments[50].

However, this uses an internal initiation mechanism for polycistronic genes, similar to *MT-ND4*

and *MT-ATP6.* Until now, accurate mapping of mitochondrial translation events at the 5' ends of

transcripts has been unavailable. We identified MRPFs that map to putative start codons

occurring on mitochondrial MT-tRNAs and *MT-RNR1,* based on the size of the reads occurring at

the 5' end of the RNA and their relationship to a start codon localized in the sequence. The site

of *MT-RNR1* was further supported by analysis of the resulting ORF, suggesting translation

initiation and elongation occurs. Curiously, tRNA^{Tyr}, encoded by *MT-TY*, is polyadenylated during

processing, which may be related to our identification of a ribosome binding event on this tRNA[51].

Due to both the short ORFs, minimal read coverage, and lack of a detectable peptide we could

not determine if translation elongation occurred on the mitochondrial tRNAs. These sites may be

unproductive mitoribosome binding events or simply from tRNAs partially digesting during MNase

treatment. Future work is needed to understand whether these are actual translation events and

to determine the function of these potential novel peptides. Together, our findings improve our

understanding of mitochondrial translation initiation, gene expression, and the mechanisms of

translation inhibition.

Materials and Methods

Isolation of mitoribosomes for profiling and western blotting

Mitoribosome profiling was carried out in HEK293 Flp-In T-REx cells. 15 cm plates were treated with either DMSO as a control or 100 μg/mL of protein synthesis inhibitor for 1 hour, washed with 4°C PBS, and flash frozen by floating the plate on liquid nitrogen. Cells were collected by scraping on ice with 600 μL of lysis buffer (1.5x Lysis buffer (30 mM Tris–HCl pH 7.8, 150 mM KCl, 15 mM $MgCl_2$, 1.5 mM DTT, 1.5% Triton X-100, 0.15% NP40, 1 × complete phosphatase and protease inhibitors)) and lysed by passing through a 27-G needle 10 times. Lysate was centrifuged for 10 minutes at 21,000 g at 4°C, and the supernatant was collected. 300 μL of lysate was footprinted with either digested with 1,5000 U of MNase, 10 μL of SUPERase-In, and 5 mM of $CaCl_2$ at 22°C for 1 hour and stopped by adding EGTA to a concentration of 6 mM or with 8 U/μL of RNase I at 22°C for 1 hour and stopped by adding 10 μL of SUPERase-In. Titration of the digestion enzymes was not performed and is based on previous published work[29,50]. The digestion was clarified by centrifugation at 21,000 g for 5 minutes. Lysates were loaded onto a 5%-30% sucrose gradient (20 mM Tris-HCl, pH 7.8, 100 mM KCl, 10mM $MgCl_2$, 1 mM DTT) and spun for 2.5 hours at 40,000 rpm in a SW40 rotor. 55S mitoribosome fractions were collected using a BioComp Fractionator and MRPFs were isolated by SDS/hot phenol/chloroform extraction.

Mitoribosome profiling library construction

Samples were run on a 15% TBE-Urea polyacrylamide gel and footprint sizes from 15-34 nts were excised and footprints were eluted in 0.3 M NaOAc, 1 mM EDTA, and 0.25% SDS followed by purification by Oligo Clean & Concentrator. Samples were treated with Quick CIP. 3' adapters (5'-rAppNN-6 nt UMI-TGGAATTCTCGGGTGCCAAGG-L) were ligated using Rnl2(1-249)K227Q ligase for 2 hours at 16°C and pooled. Pooled samples were treated with T4 PNK for 30 minutes at 37°C. A 5' chimeric DNA-RNA adapter (GTTCAGAGTTCTACAGTCCGACGATCrNrNrNrN) was ligated to the samples with T4 RNL1. Reverse transcription was carried out using SuperScript IV followed by PCR amplification by Taq DNA polymerase. Libraries were size selected using a 3% agarose Pippin prep and final libraries were PCR amplified. Samples were sequenced as either single- or paired-end 50 nt reads on either a NextSeq550 or NovaSeq.

Mitoribosome profiling analysis

Adapter sequences were removed with cutadapt and aligned with bowtie2 to a fasta containing rRNA and tRNA sequences. To identify stalling sites, reads were mapped to a custom

mitochondrial transcriptome containing all 5' UTRs, coding sequences, and 3' UTRs for all
processed mRNAs. Reads were aligned with bowtie2. A 13-nt P-site offset was assigned using
Plastid and mapping reads relative to the start codons of *ND4*, *ATP6*, *MT-ND5-dORF.* We
determined the relative ribosome density by normalizing the number of reads at each codon by
the total number of reads mapping to its respective gene for each sample. Fold change was then
calculated by dividing the relative ribosome density in a treated sample by that in an untreated
sample, excluding sites that carried less than 0.1% of the total ribosome density of each gene. To
identify putative novel translation initiation sites, we mapped our footprints to the entire
mitochondrial genome and visually inspected the 3' ends of mitochondrial tRNAs and rRNAs.
495

Mass spectrometry analysis of mitochondrial proteins
Mitochondria were isolated from 2×10^7 HEK293 cells using the Mitochondria Isolation Kit for
Cultured Cells (Thermo Scientific). Mitochondrial proteins were processed for trypsin/LysC
digestion, and the digested peptide mixture was then concentrated and desalted using C18
column. Reconstituted desalted peptides in 25 µl of 0.1% formic acid. 12 µl of peptides from each
sample was analyzed by a 110 min LC/MS/MS run. The LC/MS/MS analysis of tryptic peptides
for each sample was performed sequentially with a blank run between each two sample runs
using a Thermo Scientific Orbitrap Exploris 240 Mass Spectrometer and a Thermo Dionex
UltiMate 3000 RSLCnano System. Peptides from trypsin digestion were loaded onto a peptide
trap cartridge at a flow rate of 5 µL/min. The trapped peptides were eluted onto a reversed-phase
Easy-Spray Column PepMap RSLC, C18, 2 µM, 100A, 75 µm × 250 mm (Thermo Scientific) using
a linear gradient of acetonitrile (3-36%) in 0.1% formic acid. The elution duration was 60 min at a
flow rate of 0.3 µl/min. Eluted peptides from the Easy-Spray column were ionized and sprayed
into the mass spectrometer, using a Nano Easy-Spray Ion Source (Thermo) under the following
settings: spray voltage, 1.6 kV, Capillary temperature, 275°C. Other settings were empirically
determined. Raw data files were searched against the human protein sequences database and
the client target protein sequences database using the Proteome Discoverer 2.5 software
(Thermo, San Jose, CA) based on the SEQUEST algorithm. Carbamidomethylation (+57.021 Da)
of cysteines was set as a fixed modification, and Oxidation / +15.995 Da (M), Phospho / +79.966
Da (S, T, Y), and Deamidated / +0.984 Da (N, Q) were set as dynamic modifications. The
minimum peptide length was specified to be five amino acids. The precursor mass tolerance was
set to 15 ppm, whereas the fragment mass tolerance was set to 0.05 Da. The maximum false
peptide discovery rate was specified as 0.01. The resulting Proteome Discoverer Report contains

all assembled proteins with peptides sequences and peptide spectrum match counts (PSM#) and
MS1 peak area relative abundance.

Data sharing

522 Mitoribosome fastq files, wig files mapping to mitochondrial mRNAs, and read length files for
523 MNase-treated samples mapping to chrM are available through GEO Series GSE277563.
524

Author contributions

526 J.M. conceived the project, designed and executed experiments, analyzed data, and wrote the
527 manuscript. E.Y. analyzed mitoribosome profiling data. M.H. supervised the project, advised on
528 data interpretation and experimental design, and edited the manuscript.
529

Acknowledgments

We'd like to thank Nicholas Guydosh (NIDDK), Sezen Meydan (Vanderbilt University), Astrid
Haase (NIDDK), the Haase lab members, and the Hafner lab members for their feedback on this
work. We thank Daniel Benhalvey (Tel Aviv University) for his assistance in carrying out ribosome
profiling and Sarah Young-Baird (Uniformed Services University) for assistance with sucrose
gradient fractionation. We thank Stefania Dell'Orso and Faiza Naz (NIH/NIAMS Genomic
Technology Section) for sequencing experiments. This research was supported by the Intramural
Research Program of the National Institutes of Health (NIH). The contributions of the NIH authors
are considered Works of the United States Government. The findings and conclusions presented
in this paper are those of the authors and do not necessarily reflect the views of the NIH or the
U.S. Department of Health and Human Services.
541

Funding

543 Intramural Research Program of the National Institutes of Health, National Institute of Arthritis and
544 Musculoskeletal and Skin Diseases (ZIA-AR041205 to M.H.).
545

Supplementary Figures and Tables

	Nucleotide 1	Nucleotide 2	Nucleotide 3
MT-ND1	0.407	0.321	0.272
MT-ND2	0.381	0.332	0.287
MT-CO1	0.406	0.297	0.297
MT-CO2	0.396	0.314	0.290
MT-ATP8	0.386	0.303	0.311
MT-ATP6	0.382	0.321	0.297
MT-CO3	0.387	0.304	0.308
MT-ND3	0.404	0.302	0.293
MT-ND4L	0.359	0.347	0.294
MT-ND4	0.418	0.302	0.280
MT-ND5	0.385	0.337	0.278
MT-ND6	0.369	0.294	0.338
MT-CYB	0.408	0.304	0.287
MT-RNR1_ORF	0.365	0.347	0.288
MT-RNR1_ncRNA	0.338	0.331	0.331
MT-Humanin	0.345	0.376	0.279
MT-TI	0.030	0.685	0.284
MT-TS1	0.245	0.460	0.296

Table S1. Read phasing values for mitoribosomes footprinted with RNaseI.
The read phasing values for the canonical and putative ORFs.

553
Figure S1. Mitoribosome profiling traces of *MT-CYB*.
Individual traces from mitoribosome profiling from each untreated control labeled with its'
corresponding antibiotic-treated sample.

557
558
559

A

B

Gene	Stall site (codon)	Log Fold Change	Stall Motif	Length of Gene (codons)
MT-ND4	80	55.66929134	RKK	459
MT-ND2	104	34.63291139	AMK	347
MT-ATP6	96	13.34866237	KIK	226
MT-CO1	480	8.44860196	KRK	513
MT-ATP6	94	7.213050223	RSK	226
MT-ND1	53	6.980821918	AMK	318
MT-ND5	511	6.752314298	KLK	603
MT-ND2	57	4.222222222	AIK	347
MT-CO1	478	4.049649208	ASK	513
MT-ND2	313	2.738072055	KMK	347
MT-ND5	513	1.706953542	KMK	603
MT-ND5	222	0.9491412	AGK	603

560
561 **Figure S2. Distribution of A/R/K-X-K telithromycin stalling site.**
562 (A) The frequency of each telithromycin stalling sites across canonical mtDNA-encoded genes
563 and (B) the measured log fold change at those sites as compared to an untreated control.
564

Figure S3. MRPF lengths after treatment with initiation inhibitors correlate to alternative initiation sites.

(A) Read lengths of MNase-generated MRPFs from cells treated with josamycin or (B) retapamulin.

Supplementary Files

Supplemental File 1

Peptides identified by protein mass spectrometry of isolated mitochondria.

References

1. Gray, M.W., Burger, G., and Lang, B.F. (1999). Mitochondrial evolution. Science *283*, 1476-1481. 10.1126/science.283.5407.1476.
2. D'Souza, A.R., and Minczuk, M. (2018). Mitochondrial transcription and translation: overview. Essays Biochem *62*, 309-320. 10.1042/EBC20170102.
3. Enriquez, J.A., Chomyn, A., and Attardi, G. (1995). MtDNA mutation in MERRF syndrome causes defective aminoacylation of tRNA(Lys) and premature translation termination. Nat Genet *10*, 47-55. 10.1038/ng0595-47.
4. Webb, B.D., Diaz, G.A., and Prasun, P. (2020). Mitochondrial translation defects and human disease. J Transl Genet Genom *4*, 71-80. 10.20517/jtgg.2020.11.
5. Prezant, T.R., Agapian, J.V., Bohlman, M.C., Bu, X., Oztas, S., Qiu, W.Q., Arnos, K.S., Cortopassi, G.A., Jaber, L., Rotter, J.I., and et al. (1993). Mitochondrial ribosomal RNA mutation associated with both antibiotic-induced and non-syndromic deafness. Nat Genet *4*, 289-294. 10.1038/ng0793-289.
6. Zheng, J., Ji, Y., and Guan, M.X. (2012). Mitochondrial tRNA mutations associated with deafness. Mitochondrion *12*, 406-413. 10.1016/j.mito.2012.04.001.
7. Smick, K.M., Condit, P.K., Proctor, R.L., and Sutcher, V. (1964). Fatal Aplastic Anemia. An Epidemiological Study of Its Relationship to the Drug Chloramphenicol. J Chronic Dis *17*, 899-914. 10.1016/0021-9681(64)90161-4.
8. Woodhead, J.L., Yang, K., Oldach, D., MacLauchlin, C., Fernandes, P., Watkins, P.B., Siler, S.Q., and Howell, B.A. (2019). Analyzing the Mechanisms Behind Macrolide Antibiotic-Induced Liver Injury Using Quantitative Systems Toxicology Modeling. Pharm Res *36*, 48. 10.1007/s11095-019-2582-y.
9. De Vriese, A.S., Coster, R.V., Smet, J., Seneca, S., Lovering, A., Van Haute, L.L., Vanopdenbosch, L.J., Martin, J.J., Groote, C.C., Vandecasteele, S., and Boelaert, J.R. (2006). Linezolid-induced inhibition of mitochondrial protein synthesis. Clin Infect Dis *42*, 1111-1117. 10.1086/501356.
10. Gupta, A., Okesli-Armlovich, A., Morgens, D., Bassik, M.C., and Khosla, C. (2020). A genome-wide analysis of targets of macrolide antibiotics in mammalian cells. J Biol Chem *295*, 2057-2067. 10.1074/jbc.RA119.010770.
11. Amunts, A., Brown, A., Toots, J., Scheres, S.H.W., and Ramakrishnan, V. (2015). Ribosome. The structure of the human mitochondrial ribosome. Science *348*, 95-98. 10.1126/science.aaa1193.
12. Fu, X., Wan, P., Li, P., Wang, J., Guo, S., Zhang, Y., An, Y., Ye, C., Liu, Z., Gao, J., et al. (2021). Mechanism and Prevention of Ototoxicity Induced by Aminoglycosides. Front Cell Neurosci *15*, 692762. 10.3389/fncel.2021.692762.
13. Morales-Alvarez, M.C. (2020). Nephrotoxicity of Antimicrobials and Antibiotics. Adv Chronic Kidney Dis *27*, 31-37. 10.1053/j.ackd.2019.08.001.
14. Soriano, A., Miro, O., and Mensa, J. (2005). Mitochondrial toxicity associated with linezolid. N Engl J Med *353*, 2305-2306. 10.1056/NEJM200511243532123.
15. Palenzuela, L., Hahn, N.M., Nelson, R.P., Jr., Arno, J.N., Schobert, C., Bethel, R., Ostrowski, L.A., Sharma, M.R., Datta, P.P., Agrawal, R.K., et al. (2005). Does linezolid cause lactic acidosis by inhibiting mitochondrial protein synthesis? Clin Infect Dis *40*, e113-116. 10.1086/430441.
16. Bale, B.I., Elebesunu, E.E., Manikavasagar, P., Agwuna, F.O., Ogunkola, I.O., Sow, A.U., and Lucero-Prisno, D.E., 3rd (2023). Antibiotic resistance in ocular bacterial infections: an

integrative review of ophthalmic chloramphenicol. Trop Med Health *51*, 15.
10.1186/s41182-023-00496-x.

17. Lorenzo, D. (2019). Chloramphenicol Resurrected: A Journey from Antibiotic Resistance
in Eye Infections to Biofilm and Ocular Microbiota. Microorganisms *7*.
10.3390/microorganisms7090278.

18. Browne, A.J., Chipeta, M.G., Haines-Woodhouse, G., Kumaran, E.P.A., Hamadani,
B.H.K., Zaraa, S., Henry, N.J., Deshpande, A., Reiner, R.C., Jr., Day, N.P.J., et al. (2021).
Global antibiotic consumption and usage in humans, 2000-18: a spatial modelling study.
Lancet Planet Health *5*, e893-e904. 10.1016/S2542-5196(21)00280-1.

19. Graham, D.J. (2006). Telithromycin and acute liver failure. N Engl J Med *355*, 2260-2261.
10.1056/NEJMc066372.

20. Paternoga, H., Crowe-McAuliffe, C., Bock, L.V., Koller, T.O., Morici, M., Beckert, B.,
Myasnikov, A.G., Grubmuller, H., Novacek, J., and Wilson, D.N. (2023). Structural
conservation of antibiotic interaction with ribosomes. Nat Struct Mol Biol *30*, 1380-1392.
10.1038/s41594-023-01047-y.

21. Lin, J., Zhou, D., Steitz, T.A., Polikanov, Y.S., and Gagnon, M.G. (2018). Ribosome-
Targeting Antibiotics: Modes of Action, Mechanisms of Resistance, and Implications for
Drug Design. Annu Rev Biochem *87*, 451-478. 10.1146/annurev-biochem-062917-
011942.

22. Yan, K., Madden, L., Choudhry, A.E., Voigt, C.S., Copeland, R.A., and Gontarek, R.R.
(2006). Biochemical characterization of the interactions of the novel pleuromutilin
derivative retapamulin with bacterial ribosomes. Antimicrob Agents Chemother *50*, 3875-
3881. 10.1128/AAC.00184-06.

23. Meydan, S., Marks, J., Klepacki, D., Sharma, V., Baranov, P.V., Firth, A.E., Margus, T.,
Kefi, A., Vazquez-Laslop, N., and Mankin, A.S. (2019). Retapamulin-Assisted Ribosome
Profiling Reveals the Alternative Bacterial Proteome. Mol Cell *74*, 481-493 e486.
10.1016/j.molcel.2019.02.017.

24. Choi, J., Marks, J., Zhang, J., Chen, D.H., Wang, J., Vazquez-Laslop, N., Mankin, A.S.,
and Puglisi, J.D. (2020). Dynamics of the context-specific translation arrest by
chloramphenicol and linezolid. Nat Chem Biol *16*, 310-317. 10.1038/s41589-019-0423-2.

25. Marks, J., Kannan, K., Roncase, E.J., Klepacki, D., Kefi, A., Orelle, C., Vazquez-Laslop,
N., and Mankin, A. (2016). Context-specific inhibition of translation by ribosomal
antibiotics targeting the peptidyl transferase center. Proc Natl Acad Sci U S A *113*, 12150-
12155. 10.1073/pnas.1613055113.

26. Itoh, Y., Singh, V., Khawaja, A., Naschberger, A., Nguyen, M.D., Rorbach, J., and Amunts,
A. (2022). Structure of the mitoribosomal small subunit with streptomycin reveals Fe-S
clusters and physiological molecules. Elife *11*. 10.7554/eLife.77460.

27. Amunts, A., Brown, A., Bai, X.C., Llacer, J.L., Hussain, T., Emsley, P., Long, F.,
Murshudov, G., Scheres, S.H.W., and Ramakrishnan, V. (2014). Structure of the yeast
mitochondrial large ribosomal subunit. Science *343*, 1485-1489.
10.1126/science.1249410.

28. Li, S.H., Nofal, M., Parsons, L.R., Rabinowitz, J.D., and Gitai, Z. (2021). Monitoring
mammalian mitochondrial translation with MitoRiboSeq. Nat Protoc *16*, 2802-2825.
10.1038/s41596-021-00517-1.

29. Rooijers, K., Loayza-Puch, F., Nijtmans, L.G., and Agami, R. (2013). Ribosome profiling
reveals features of normal and disease-associated mitochondrial translation. Nat
Commun *4*, 2886. 10.1038/ncomms3886.

30. Sothiselvam, S., Neuner, S., Rigger, L., Klepacki, D., Micura, R., Vazquez-Laslop, N., and
Mankin, A.S. (2016). Binding of Macrolide Antibiotics Leads to Ribosomal Selection

against Specific Substrates Based on Their Charge and Size. Cell Rep *16*, 1789-1799.
10.1016/j.celrep.2016.07.018.

31. Schlunzen, F., Pyetan, E., Fucini, P., Yonath, A., and Harms, J.M. (2004). Inhibition of
peptide bond formation by pleuromutilins: the structure of the 50S ribosomal subunit from
Deinococcus radiodurans in complex with tiamulin. Mol Microbiol *54*, 1287-1294.
10.1111/j.1365-2958.2004.04346.x.

32. Poulsen, S.M., Karlsson, M., Johansson, L.B., and Vester, B. (2001). The pleuromutilin
drugs tiamulin and valnemulin bind to the RNA at the peptidyl transferase centre on the
ribosome. Mol Microbiol *41*, 1091-1099. 10.1046/j.1365-2958.2001.02595.x.

33. Davidovich, C., Bashan, A., Auerbach-Nevo, T., Yaggie, R.D., Gontarek, R.R., and
Yonath, A. (2007). Induced-fit tightens pleuromutilins binding to ribosomes and remote
interactions enable their selectivity. Proc Natl Acad Sci U S A *104*, 4291-4296.
10.1073/pnas.0700041104.

34. Arenz, S., Ramu, H., Gupta, P., Berninghausen, O., Beckmann, R., Vazquez-Laslop, N.,
Mankin, A.S., and Wilson, D.N. (2014). Molecular basis for erythromycin-dependent
ribosome stalling during translation of the ErmBL leader peptide. Nat Commun *5*, 3501.
10.1038/ncomms4501.

35. Beckert, B., Leroy, E.C., Sothiselvam, S., Bock, L.V., Svetlov, M.S., Graf, M., Arenz, S.,
Abdelshahid, M., Seip, B., Grubmuller, H., et al. (2021). Structural and mechanistic basis
for translation inhibition by macrolide and ketolide antibiotics. Nat Commun *12*, 4466.
10.1038/s41467-021-24674-9.

36. Dubot, A., Godinot, C., Dumur, V., Sablonniere, B., Stojkovic, T., Cuisset, J.M., Vojtiskova,
A., Pecina, P., Jesina, P., and Houstek, J. (2004). GUG is an efficient initiation codon to
translate the human mitochondrial ATP6 gene. Biochem Biophys Res Commun *313*, 687-
693. 10.1016/j.bbrc.2003.12.013.

37. Kruger, A., Remes, C., Shiriaev, D.I., Liu, Y., Spahr, H., Wibom, R., Atanassov, I., Nguyen,
M.D., Cooperman, B.S., and Rorbach, J. (2023). Human mitochondria require mtRF1 for
translation termination at non-canonical stop codons. Nat Commun *14*, 30.
10.1038/s41467-022-35684-6.

38. Tsai, K., Stojkovic, V., Lee, D.J., Young, I.D., Szal, T., Klepacki, D., Vazquez-Laslop, N.,
Mankin, A.S., Fraser, J.S., and Fujimori, D.G. (2022). Structural basis for context-specific
inhibition of translation by oxazolidinone antibiotics. Nat Struct Mol Biol *29*, 162-171.
10.1038/s41594-022-00723-9.

39. Syroegin, E.A., Flemmich, L., Klepacki, D., Vazquez-Laslop, N., Micura, R., and
Polikanov, Y.S. (2022). Structural basis for the context-specific action of the classic
peptidyl transferase inhibitor chloramphenicol. Nat Struct Mol Biol *29*, 152-161.
10.1038/s41594-022-00720-y.

40. Wakigawa, T., Mito, M., Yamashiro, H., Tomuro, K., Tani, H., Tomizawa, K., Chujo, T.,
Nagao, A., Suzuki, T., Wei, F.-Y., et al. (2023). Complexity and dynamics of in
organello translation landscape assessed by high-resolution mitochondrial
ribosome profiling. bioRxiv, 2023.2007.2019.549812. 10.1101/2023.07.19.549812.

41. Bibel, B., Raskar, T., Couvillion, M., Lee, M., Kleinman, J.I., Takeuchi-Tomita, N.,
Churchman, L.S., Fraser, J.S., and Fujimori, D.G. (2024). Context-specific inhibition of
mitochondrial ribosomes by phenicol and oxazolidinone antibiotics. bioRxiv,
2024.2008.2021.609012. 10.1101/2024.08.21.609012.

42. Brandon, M.C., Lott, M.T., Nguyen, K.C., Spolim, S., Navathe, S.B., Baldi, P., and
Wallace, D.C. (2005). MITOMAP: a human mitochondrial genome database--2004
update. Nucleic Acids Res *33*, D611-613. 10.1093/nar/gki079.

43. Kalghatgi, S., Spina, C.S., Costello, J.C., Liesa, M., Morones-Ramirez, J.R., Slomovic, S.,
Molina, A., Shirihai, O.S., and Collins, J.J. (2013). Bactericidal antibiotics induce

mitochondrial dysfunction and oxidative damage in Mammalian cells. Sci Transl Med *5*,
192ra185. 10.1126/scitranslmed.3006055.

44.   D'Achille, G., and Morroni, G. (2023). Side effects of antibiotics and perturbations of
mitochondria functions. Int Rev Cell Mol Biol *377*, 121-139.
10.1016/bs.ircmb.2023.03.009.

45.   Desai, N., Yang, H., Chandrasekaran, V., Kazi, R., Minczuk, M., and Ramakrishnan, V.
(2020). Elongational stalling activates mitoribosome-associated quality control. Science
370, 1105-1110. 10.1126/science.abc7782.

46.   Singh, V., Itoh, Y., Del'Olio, S., Hassan, A., Naschberger, A., Flygaard, R.K., Nobe, Y.,
Izumikawa, K., Aibara, S., Andrell, J., et al. (2024). Mitoribosome structure with cofactors
and modifications reveals mechanism of ligand binding and interactions with L1 stalk. Nat
Commun *15*, 4272. 10.1038/s41467-024-48163-x.

47.   Kong, Q.P., Yao, Y.G., Sun, C., Zhu, C.L., Zhong, L., Wang, C.Y., Cai, W.W., Xu, X.M.,
Xu, A.L., and Zhang, Y.P. (2004). Phylogeographic analysis of mitochondrial DNA
haplogroup F2 in China reveals T12338C in the initiation codon of the ND5 gene not to be
pathogenic. J Hum Genet *49*, 414-423. 10.1007/s10038-004-0170-3.

48.   Vilarinho, L., Chorao, R., Cardoso, M.L., Rocha, H., Nogueira, C., and Santorelli, F.M.
(1999). The ND1 T3308C mutation may be a mtDNA polymorphism. Report of two
Portuguese patients. J Inherit Metab Dis *22*, 90-91. 10.1023/a:1005471904710.

49.   Rocha, H., Flores, C., Campos, Y., Arenas, J., Vilarinho, L., Santorelli, F.M., and Torroni,
A. (1999). About the "Pathological" role of the mtDNA T3308C mutationellipsis. Am J Hum
Genet *65*, 1457-1459. 10.1086/302641.

50.   Soto, I., Couvillion, M., Hansen, K.G., McShane, E., Moran, J.C., Barrientos, A., and
Churchman, L.S. (2022). Balanced mitochondrial and cytosolic translatomes underlie the
biogenesis of human respiratory complexes. Genome Biol *23*, 170. 10.1186/s13059-022-
02732-9.

51.   Fiedler, M., Rossmanith, W., Wahle, E., and Rammelt, C. (2015). Mitochondrial poly(A)
polymerase is involved in tRNA repair. Nucleic Acids Res *43*, 9937-9949.
10.1093/nar/gkv891.