Genomic insights into the adaptive evolution of the memorial rose (*Rosa lucieae*; Rosaceae) in saline environments

Ji-Hyeon Jeon[1], Myunghee Jung[2], Younhee Shin[2*] & Seung-Chul Kim[1*]

[1]Department of Biological Sciences, Sungkyunkwan University, Suwon 16419, Republic of Korea;

[2]Research and Development Center, Insilicogen Inc., Yongin 16954, Republic of Korea

*Corresponding authors:

Seung-Chul Kim, sonchus96@skku.edu;

Younhee Shin, yhshin@insilicogen.com

ABSTRACT

Rosa lucieae (syn. *Rosa wichurana* or *R. wichuraiana*; Rosaceae) is one of the major wild

progenitors of modern rose cultivars. In breeding roses, *R. lucieae* has contributed not only to

white blooms and shiny leaves, but most of all, to succeed in various harsh conditions and

high resistance to diseases and pests. The wild *R. lucieae* inhabits littoral sand banks or rocks,

where drought stress and high salinity inhibit plant growth. In this study, we assembled the

chromosome-level genome sequence of a wild accession of *R. lucieae* and identified potential

genes that may responsible for the evolution of salt tolerance. Within the 544.7 Mb-long

genome of *R. lucieae*, recent duplications and divergence of the genes responsible to stress

tolerance were identified, including stress response, signaling, homeostasis, and

detoxification. Additionally, the gene family evolution analyses elucidated rapid evolution of

stress tolerant functions in *R. lucieae*. The first high-quality genome assembly of wild *R.*

lucieae contributes significantly to our understanding of the genomic evolution and adaptive

mechanism of *R. lucieae*, which promises the enhancement of our efforts in rose breeding

programs.

Keywords: *Rosa lucieae*; *Rosa wichurana*; rose genome; salt tolerance; salinity resistance;

adaptive evolution

INTRODUCTION

The roses are the flowering plants belonging to the genus *Rosa* in the family Rosaceae, encompassing 100–200 wild species and more than 30,000 cultivars (Rehder, 1940; Young et al., 2007). As renowned for their beauty and fragrance, roses have been beloved for millennia along with the history of human civilization, and cultivated for ornamental, aromatizing, therapeutic and culinary purposes in global (Rehder, 1940; Yu and Ku, 1985; Gudin, 2003; Nybom, 2009). Despite importance of roses, rose cultivation has faced various hurdles to overcome, including environmental changes, diseases and pests, hybrid incompatibility, and productivity (Gerard, 1897; Van Fleet, 1908; Gudin, 2003; Nybom, 2009). However, breeding hardy wild roses developed many robust hybrid cultivars which can cope with those problems (Van Fleet, 1908; Gudin, 2003). *Rosa lucieae*, also well known as its synonym, *Rosa wichurana* or *R. wichuraiana,* or the memorial rose, has been cherished for centuries as one of the major progenitors contributed to developing many modern rose cultivars (Gerard, 1897; Van Fleet, 1908; Yu and Ku, 1985; Crespel and Mouchotte, 2003; Marriott, 2003; Nybom, 2009). *Rosa lucieae* is a rambler rose vigorously growing even in the bad climate and on bare soil or rocks (Gerard, 1897; Rehder, 1940; Yu and Ku, 1985), highly resistant to the prevalent rose diseases such as black spot (*Diplocarpon rosae*; Noack, 2003; Byrne et al., 2005), and easy to cross (Gerard, 1897; Van Fleet, 1908). Such characteristics of *R. lucieae* have led the successful development of thousands of robust *R. lucieae* hybrid cultivars (Gerard, 1897; Van Fleet, 1908; Rehder, 1940; Yu and Ku, 1985; Byrne et al., 2005). Consequently, in the classification scheme of modern roses, the Hybrid Wichurana is now identified as one of the cultivar groups, bred by crossing *R. lucieae* and other roses and characterized by its rambling habit and wide variety of flower shapes and colors (Cairns, 2003).

Rosa lucieae has been classified as a member of the section *Synstylae* within genus

Rosa, where the species are closely related but diversified with unique characteristics to adapt

to various ecological habitats (Rehder, 1940; Wissemann, 2003; Jeon et al., 2024). Unlike

most of other *Rosa* sect. *Synstylae* species, the natural habitats of *R. lucieae* are open and

exposed rocky slopes or banks, especially along the sea shores and riverside at low elevation

(Rehder, 1940; Ohba, 1993; Ku and Robertson, 2003). In these habitats, both abiotic stresses

such as high light, high salinity and heavy metals, and biotic stresses caused by various

herbivores and pathogens restrict plant growths and introduction. Nevertheless, *R. lucieae* has

adapted to harsh environments of those habitats, evolving to resist and tolerate to various

abiotic and biotic stresses. In addition to their ornamental characteristics, stress resistance and

resilience in various environmental conditions are pivotal in breeding roses, efficiently

increasing productivity and durability (Van Fleet, 1908; Gudin, 2003; Debener and Linde,

2009; Leus et al., 2018). Utilizing wild roses including *R. lucieae* in the rose breeding

programs will pave the way for introducing the stress tolerant traits to cultivated roses (Van

Fleet, 1908; Noack, 2003; De Cock et al., 2007). Understanding genetic and genomic

architectures of roses also promises improvement in breeding roses for the resistance by

allowing the applications such as marker-assisted selection and genetic engineering (Debener

and Hibrand-Saint Oyant, 2009; Debener and Linde, 2009; Gahlaut et al., 2021). Recently,

several transcriptomic approaches illustrated genetic functions and mechanisms of the stress

responses in hardy wild roses, including disease resistance (Xiang et al., 2019) and cold

tolerance (Zhang et al., 2016; Zhuang et al., 2021). Additionally, the genome of *Rosa rugosa*

provided insights into the evolution of stress response in harsh environments (Chen et al.,

2021). Nevertheless, the genomic intricacies of roses have impeded the extensive

understanding on the genomic bases of key characteristics in the rose cultivation (Gahlaut et

al., 2021).

The salt stress is one of the major abiotic stresses in plants, which induces osmotic

stress, ion toxicity, oxidative stress, and decreasing in photosynthesis (Zhu, 2001; Munns and

Tester, 2008). In agriculture, nearly 20 % of global irrigated lands have been salinized, and

recurrent irrigation with brackish groundwater or return flow caused from limited sources of

fresh water accumulates salts in agricultural lands (Ghassemi et al., 1995; Tilman et al., 2002;

Munns, 2005). Meanwhile, the increased level of soil salinity has severely affected the

decline in productivity and quality of crops, resulting in consistent and complex economic

losses (Munns and Gilliham, 2015). Not only can the high concentration of salts directly

affect to the cytological and physiological dysfunction, but also involve other secondary

cytotoxic effects caused from generation of reactive oxygen species (ROS), imbalance of

other essential ions such as K^+, and malnutrition (Zhu, 2001; Botella et al., 2005). In plants,

three major mechanisms of salt tolerance are known: homeostasis, detoxification, and growth

control (Zhu, 2001; Munns, 2005; Munns and Tester, 2008). Homeostasis can be established

by compartmentalization of Na^+ and Cl^- into the vacuoles, modulation Na^+ influx and efflux,

and accumulating compatible osmolytes, which are regulated by root endoderm and xylem, or

stress signaling pathway including Ca^{2+}-mediated cascades (Zhu, 2001; Botella et al., 2005;

Munns, 2005; Van Zelm et al., 2020). Ionic and oxidative toxicity can be detoxified or

prevented by protein protection and repair, antioxidant metabolites, and mitogen-activated

protein kinase (MAPK) cascades-induced detoxifying enzymes (Zhu, 2001; Chinnusamy and

Zhu. 2003; Botella et al., 2005; Munns, 2005; Van Zelm et al., 2020). Salt stress also can

inhibit the plant growth with stomal closure induced by abscisic acid (ABA), restricting the

photosynthesis (Botella et al., 2005; Van Zelm et al., 2020). However, the growth-promoting

hormones such as jasmonic acid (JA), brassinosteroid (BR), and gibberellic acid (GA) are

responsible for the growth recovery after salt stress (Van Zelm et al., 2020).

Roses are generally salt sensitive with lower yield, fewer flowers, and poor growths

even in relatively low salt concentrations (Cabrera and Perdomo, 2003; Cabrera et al., 2009;

Niu and Sun, 2017). Nevertheless, the genetic or genomic studies of roses not only lag behind

other rosaceous crops (e.g., apples, peaches, and strawberries; Shulaev et al., 2008; Foucher,

2009; Gahlaut et al., 2021), but the salt tolerance of roses is also poorly understood compared

to the aesthetic features (e.g., flower forms, colors, and fragrance), which have been priorly

considered in breeding roses (Gudin, 2003; Bao et al., 2021). Given the deteriorating soil

salinity stemming from recurrent irrigation, urbanization and climate changes, the salt

tolerance becomes pivotal in the rose cultivation (Cai et al., 2014; Corwin, 2021).

Consequently, the advent of high-throughput next-generation sequencing technique has led

recent multi-omics investigation into the salt tolerance in various rose cultivars of *Rosa*

chinensis (Tian et al., 2018; Bao et al., 2021), *R. rugosa* (Chen et al., 2021; Qi et al., 2023), *R.*

× damascena and *R. × hybrida* (Ren et al., 2024).

In this study, a chromosome-level genome of a wild accession of *R. lucieae*, inhabiting

the saline sea shore, was assembled in order to understand the genomic evolution and nature

of salt tolerance in *R. lucieae*. The chromosome-level genome of *R. lucieae* and its evolution

in salt tolerance will contribute to our understanding of genomic features and adaptive

mechanisms of wild roses, and promise to enhance our efforts in rose breeding programs.

20   **RESULTS**

Molecular sequencing, genome assembly, and annotation of Rosa lucieae

A total of 374.6 million paired-end short-sequence reads with 56.6 Gbp (123.4×

coverage depths of the estimated haploid genome size) were generated through Illumina

HiSeq platform, and 2.7 million long-sequence reads with 28.7 Gbp (62.6× coverage depths

of the estimated genome size) were generated through PacBio Sequel platform

(Supplementary Table S1). Each of two transcriptome samples from the young leaves and

flower buds was sequenced into 9.0-Gbp data with 89 million paired-end short-sequence

reads through Illumina NovaSeq platform (Supplementary Table S1). The *k*-mer analysis

using genomic short reads estimated the haploid genome size of *R. lucieae* at 458.3 Mbp, of

which repeat content was 59.61 % (273.2 Mbp) and the heterozygosity rate was 1.35% (Table

1; Supplementary Fig. S1).

The error-corrected PacBio subreads were *de novo* assembled into 2067 primary

contigs with a total of 619.5 Mbp lengths, and this heterozygous primary assembly was

curated into the 1356 contigs with a total of 544.9 Mbp lengths. Finally, error sequences of

the curated assembly were corrected with short sequence reads, establishing the 544.5 Mbp-

long draft *de novo* genome assembly with 1356 contigs (Supplementary Table S2). The

BUSCO assessment identified more than 96 % of BUSCO eudicots-, embryophyte-, and

viridiplantae-ortholog datasets within the draft assembly of *R. lucieae* respectively (Table 1).

The *de novo* assembly was subsequently scaffolded into seven chromosomal sequences and

187 unplaced contigs, finalizing the 544.7 Mb-long chromosomal genomic sequences of *R.*

lucieae (Fig. 1; Table 1; Supplementary Table S3). The genome collinearity analysis with the

reference chromosomes indicated no genomic gaps, missing data, or possible mis-assembly

within the draft genome assembly of *R. lucieae* in the present study (Supplementary Fig. S2).

Within the genome of *R. lucieae*, 49.67 % of sequences (270.5 Mbp) were identified as

repeat sequences. Among them, most repeat sequences were interspersed repeats (97.4 %), of

which 54.1 % were identified as long terminal repeat (LTR) elements, 15.5 % were identified

as DNA elements, and 24.5 % were unclassified (Supplementary Table S4). A total of 36,876

protein-coding genes were identified and annotated within the genome of *R. lucieae* (Table 1),

the number of which was close to the numbers of annotated protein-coding gene of other

Rosa genomes (*R. lucieae*: 32,674; *R. chinensis*: 30,924–39,669; *R. rugosa*: 29,146–39,704

genes; Hibrand Saint-Oyant et al., 2018; Raymond et al., 2018; Chen et al., 2021; Zang et al.,

2021; Zhang et al., 2024). Among the protein-coding genes, all the genes except three

(36,873 genes; 99.99 %) were functionally annotated from the public databases, including nr,

gene ontology (GO), KEGG, and Pfam databases (Supplementary Table S5).

Structural genomic evolution of Rosa lucieae

The genome of *R. lucieae* was assembled into a total of 544.7 Mbp-long scaffolds and

contigs, of which the size was almost similar to the genomes of *R. chinensis* (488.6–541.3

Mbp; sect. *Chinenses*; Raymond et al., 2018; Hibrand Saint-Oyant et al., 2018; Zhang et al.,

2024), and slightly to moderately larger than those of *R. rugosa* (401.2–450.0 Mbp; sect.

Rosa; GCF_958449725.1; Chen et al., 2021; Zang et al., 2021), *R. roxburghii* (504.0–529.9

Mbp; *sect. Microphyllae*; Yang et al., 2024; Zong et al., 2024), and *R. persica* (366.1 Mbp;

subg. *Hulthemia*; GCA_025331145.1). The genome size profile of these *Rosa* species was

corroborated with the nucleic DNA amount investigation, estimating 2C at 1.13 pg in *R.*

lucieae, 1.16 pg in *R. chinensis*, 0.98 pg in *R. rugosa*, 0.95 pg in *R. roxburghii*, and 0.84 pg in

R. persica (Yokoya et al., 2000). Despite the increased genome size, the repeat content of

genome of *R. lucieae* (49.7 %; Table 1; Supplementary Table S4) was nearly similar to *R.*

rugosa (50.3–54.1 %; Chen et al., 2021; Zang et al., 2021), whereas it was notably increased

in the genomes of *R. chinensis* (61.8–67.9 %; Raymond et al., 2018; Hibrand Saint-Oyant et

al., 2018; Zhang et al., 2024).

Within 36,876 protein-coding genes identified within the genome of *R. lucieae*, the

ortholog group inference analysis classified 32,666 genes into 19,667 gene family groups, of

which 14,436 groups were shared by Rosoideae species (i.e., *R. lucieae*, *R. chinensis*, *R.*

rugosa and *Fragaria vesca*), 15,790 groups were shared within the genus *Rosa*, and 586

groups were unique to *R. lucieae* (Fig. 2A). Among 32,666 classified genes, 31,264 were

interspecific orthologous and the other 1402 were species-specific paralogous genes (Fig. 2B).

Based on the gene annotations of Rosoideae genomes and their homologies, the interspecific

collinear blocks were identified to infer the genome synteny. Within the genus *Rosa*, the

genome synteny was highly conserved among species, especially between *R. chinensis* and *R.*

rugosa. The genome synteny was also moderately conserved between the genera *Rosa* and

Fragaria, even though several structural variations were identified (Supplementary Fig. S3).

Despite the increased repeat content in the genomes of *R. chinensis*, the genome synteny was

highly conserved between *R. chinensis* and *R. rugosa*. Moreover, despite the closer

phylogenetic relationship of *R. chinensis* to *R. lucieae* than *R. rugosa* (Wissemann and Ritz,

2005; Fougère-Danezan et al., 2015; Debray et al., 2019; Jeon et al., 2024), more inter-

chromosomal collinear blocks were identified between *R. chinensis* and *R. lucieae*, which

also prevailed in the genomic study of *R. lucieae* 'Basye's thornless' (Zhong et al., 2021),

compared to between *R. chinensis* and *R. rugosa* (Fig. 2C; Supplementary Fig. S3). Given the

intra-genomic collinear blocks and increased gene content within the genome of *R. lucieae*

(Fig. 1; Table 1), putative gene duplication may have influenced the genomic evolution of *R.*

lucieae, which aligned with the pronounced expansion of gene families in *R. lucieae* lineage

detected from the gene family evolution analysis (Fig. 2C).

Rapid evolution of salt stress-related gene families in Rosa lucieae

The ortholog group inference analysis identified 2228 single-copy orthologous gene

families shared among the genomes of *R. lucieae*, five other rosaceous species, and the

outgroup taxa *Populus trichocarpa*. The Bayesian phylogenetic inference using the single-

copy orthologous gene set corroborated the taxonomic classification of the species, resolving

robust monophyly of the family Rosaceae, subfamilies Rosoideae and Amygdaloideae, and

genus *Rosa* respectively (Fig. 2C; Supplementary Table S6). Within the genus *Rosa*, the

closer relationship between *R. lucieae* and *R. chinensis* was inferred compared to with *R.*

rugosa, which is congruent with the previous phylogenetic inferences of the genus *Rosa*

using conventional molecular markers (Wissemann and Ritz, 2005; Fougère-Danezan et al.,

2015). In pairwise comparisons of Rosoideae species on distributions of the synonymous

substitution rates (Ks) of orthologous genes, orthologous Ks values between *R. lucieae* and *R.*

chinensis were lower (mode Ks = 0.021) than those between *R. rugosa* and other *Rosa*

species (mode Ks = 0.033–0.035), and between *F. vesca* and *Rosa* species (mode Ks = 0.187–

0.199; Supplementary Fig. S4). This result suggested more recent divergence of *R. lucieae*

and *R. chinensis* compared to their divergence with *R. rugosa*, aligning with the divergence

time estimation (Fig. 2C).

Based on the inferred orthologous gene families and species phylogeny, the gene family

evolution was inferred within each phylogenetic lineage. While contractions of gene families

were predominantly involved in the evolution of most other rosaceous lineages, the evolution

of *R. lucieae* involved the pronounced expansion of gene families, from the stem lineage of

genus *Rosa* (Fig. 2C). After the divergence with *R. chinensis*, 1875 gene families were

expanded with 2579 more genes, whereas 385 gene families were contracted with 609 less

genes within the lineage of *R. lucieae*. Among the expanded or contracted gene families

involved in the *R. lucieae* lineage, 283 gene families were inferred to be rapidly evolving in

this lineage. For the *R. lucieae* genes of rapidly evolving families, GOs related to plant

signaling, secondary metabolites, salinity stress, root system, homeostasis, calcium ion,

protein repair, or defense mechanism were significantly enriched (Fig. 3; Supplementary

Table S7). Likewise, based on their KEGG orthology (KO) annotations, KEGG pathways

related to the steroid hormones including BRs, secondary metabolites, and stress responses

were significantly enriched (Supplementary Fig. S5).

Within the genus *Rosa*, paralogous Ks values were greater (mode Ks = 0.109–0.122)

than Ks values of orthologous genes, and their distributions were mostly overlapped with

each other (Fig. 4A). This implied that paralogous genes in most gene families were likely

duplicated in their common ancestral lineage after the divergence with *F. vesca*, while the

orthologous genes have been accumulating unique genetic variation after their divergence by

duplications. Particularly with paralogous genes annotated with the GO term 'response to salt

stress' (GO:0009651) or its child terms, Ks values in *R. lucieae* were remarkably decreased

(mode Ks = 0.017) compared to Ks values of all paralogous genes (mode Ks = 0.114),

whereas Ks values in other *Rosa* species and *F. vesca* were barely changed or slightly

increased (mode Ks = 0.102–0.169; Fig. 4B). The unique Ks distribution of paralogous salt

stress-related genes in *R. lucieae* with lower values indicated the recent duplications and

divergence of these genes specifically within the evolution of *R. lucieae*.

Duplication and divergence of salt tolerance-related genes in Rosa lucieae

Within the rapidly evolving gene families of *R. lucieae*, GOs related to responses to salt

and osmotic stresses were significantly enriched (Fig. 3; Supplementary Table S7),

suggesting possible influence of gene duplications or divergence of *R. lucieae* on adaptation

to salt stress. Moreover, the unique Ks distribution with lower values of paralogous *R. lucieae*

genes annotated with GOs related to responses to salt stress corroborated their recent

divergence compared to those genes in other close species or other genes in *R. lucieae* (Fig.

4). Given that gene duplication and copy-number variation can contribute to adaptive

evolution by increasing the genetic diversity or product dosage (Ohno, 1970; Kondrashov,

2012; Magadum et al., 2013), these results substantiated the influence of duplications and

divergence of salinity response-related genes on the unique adaptation to salinity in *R. lucieae*.

Similarly, the duplications of various salinity response-related genes were identified in the

evolution of salt tolerance of the saltwater cress, *Eutrema salsugineum* (syn. *Thellungiella*

salsuginea), which is a close halophytic relative of *Arabidopsis thaliana* (Wu et al., 2012;

Monihan et al., 2019).

In addition to GOs related to responses to salt and osmotic stresses, GOs related to

ABA signaling, Ca^{2+} signaling and ROS signaling, and KEGG pathways related to MAPK

signaling were also significantly enriched in the rapidly evolving gene families of *R. lucieae*,

which are principal signaling pathways of salinity response in plants (Fig. 3; Supplementary

Fig. S5; Supplementary Table S7; Zhu, 2001; Botella et al., 2005; Van Zelm et al., 2020). The

genetic evolution related to early signaling under salt stress possibly influenced the

adaptation to salinity of *R. lucieae* by enabling a rapid and/or enhanced response to salt stress.

Within the genome of *R. lucieae*, the duplications and divergence of various salt stress

signaling-related genes were identified, including genes of Ca^{2+} channels such as cyclic

nucleotide-gated ion channel 1 (CNGC1) (Table 2; Supplementary Figs. S6 and S7;

Supplementary Table S8).

CNGC1 is a member of cyclic nucleotide-gated channels (CNGCs), which are Ca^{2+}-

permeable cation channels involved in signal transduction related to stress tolerance in plants

(Sanders et al., 2002; Jha et al., 2016). While some other members of CNGCs in *A. thaliana*,

such as AtCNGC19 and AtCNGC20, are known to be involved in salt stress (Kugler et al.,

2009), the function of *Arabidopsis* CNGC1 in stress tolerance is less understood but Pb^{2+}

tolerance (Sunkar et al., 2000). Even though the direct relationship of AtCNGC1 to salt stress

is barely known, several recent studies suggested the role of AtCNGC1 in Ca^{2+} uptakes at

roots, accumulation of Ca^{2+}, and maintaining proper Ca^{2+} levels in plants (Ma et al., 2006;

Urquhart et al., 2011). In the genome of *R. lucieae*, the copy number of CNGC1 genes was

increased compared to other close Rosoideae species (Supplementary Fig. S6). Gene

duplication of *R. lucieae CNGC1* (*RlCNGC1*) was not likely to originate from a specific

homologous gene cluster, but has occurred in diverse homologous gene clusters

independently (Supplementary Fig. S6A). The synteny of orthologous *CNGC1* mostly

conflicted with the synteny of genomic collinear blocks, and several copies of *RlCNGC1*

were inserted in multiple unplaced contigs of *R. lucieae* (Supplementary Fig. S6B, C).

The increased number of *RlCNGC1* may have affected the accumulation of Ca^{2+} in *R.*

lucieae by regulating Ca^{2+} uptake at roots, which can induce a rapid increase in Ca^{2+} levels

after salinity stress (Ma et al., 2006; Kader and Lindberg, 2010). *Rosa lucieae* can be more

Ca^{2+}-sensitive than other roses with the increased gene copy numbers of Ca^{2+} sensors such as

Calcineurin B-like protein 1 (CBL1), of which overexpression in transgenic *A. thaliana*

plants enhanced salt tolerance (Albrecht et al., 2003; Cheong et al., 2003). After the elevation

of Ca^{2+} levels induced by salt stress, the increased gene copy numbers of salt stress-related

signaling proteins including ion channels, receptors and kinases possibly influenced the

subsequent ABA, Ca^{2+} and ROS signaling in *R. lucieae* (Table 2). For example, transgenic *A.*

thaliana plants overexpressing an ABA receptor, pyrabactin resistance 1-like 9 (PYL9),

exhibited improved osmoregulation (Zhao et al., 2016). Likewise, transgenic *A. thaliana*

plants overexpressing zeaxanthin epoxidase (ZEP), which is pivotal in ABA biosynthesis,

also exhibited enhanced salt tolerance (Park et al., 2008). Thus, overexpression of stress

response- and signaling-related genes by duplication may have contributed to the adaptive

evolution of *R. lucieae* in saline environments. Given the increased genetic diversity after

duplication (Ohno, 1970; Kondrashov, 2012), the duplicated gene pairs were also possibly led

to asymmetric evolution, resulting in functional and conditional differences in stress signaling

(VanderSluis et al., 2010).

Because the early responses of salt stress including generation of ABA and ROS can

damage and impair plant systems, the downstream detoxification and recovery are another

pivotal mechanism in salt tolerance (Van Zelm et al., 2020). The significant enrichment of

metabolite-, ionic and osmotic homeostasis-, and protein repair-related GOs, and secondary

metabolite- and protein processing-related KEGG pathways implied the putative evolution of

ion- or ROS-detoxification strategies in *R. lucieae* (Fig. 3; Supplementary Fig. S5;

Supplementary Table S7). Within the genome of *R. lucieae*, the duplications and divergence

of various detoxification- and homeostasis-related genes were also identified, including genes

of ion transporters such as auto-inhibited Ca^{2+}-ATPases (ACAs) (Fig. 5; Table 2;

Supplementary Fig. S7; Supplementary Table S8).

ACAs and endoplasmic reticulum-type Ca^{2+}-ATPases (ECAs) are P-type Ca^{2+}-ATPases

found in plants, of which a primary role is terminating Ca^{2+}-mediated signaling by Ca^{2+}

efflux from the cytosol (Bose et al., 2011; Park and Shin, 2022). Thus, ACAs and ECAs are

pivotal in the removal of excessive Ca^{2+} and restoration after stress responses. ACAs are

generally classified into four conserved clusters in flowering plants: □) ACA1, ACA2 and

ACA7 in cluster 1, □) ACA4 and ACA11 in cluster2, □) ACA12 and ACA13 in cluster 3,

and □) ACA8, ACA9 and ACA10 in cluster 4 (Yu et al., 2018). The unique gene functions of

each cluster in stress responses were investigated in several model plants including *A.*

thaliana and *Oryza sativa* (Bose et al., 2011; Huda et al., 2013; Yu et al., 2018; Park and Shin,

2022). Within the genome of *R. lucieae*, genes of various members of ACA family were

duplicated, including genes of ACA10, ACA12 and ACA13 (Fig. 5). Similar to *RlCNGC1*,

the duplications of *R. lucieae ACA10*, *ACA12* and *ACA13* have occurred in diverse

homologous gene clusters independently (Fig. 5A, B). The copy numbers of *ACA13* of *Rosa*

species were notably increased compared to their relative *F. vesca*, and the copy number of

ACA13 of *R. lucieae* was increased again within the genus *Rosa* (Fig. 5A, C).

The excessive Na^+, Cl^- and Ca^{2+} resulting from high salinity can cause the ion toxicity

by themselves, and also induce severe stress responses such as overaccumulation of ROS

(Zhu, 2001; Munns and Tester, 2008). In addition to *RlACA13*, the duplications and

divergence of *R. lucieae* genes of calmodulin-like protein 18 (CML18), high-affinity K^+

transporter 1 (HKT1), inositol transporter 4 (INT4), SHORT ROOT IN SALT MEDIUM 1

(RSA1), and slow anion channel 1 homolog 3 (SLAH3) have possibly affected alleviation

and detoxification of excessive Na^+, Cl^- and Ca^{2+}, rescuing *R. lucieae* from salt stress. For

examples, the point mutation in *A. thaliana HKT1* increased the Na^+ tolerance with reduced

Na^+ uptake (Rubio et al., 1995), and three paralogous copies of *MhINT4* in the Chinese crab

apple (*Malus hupehensis*; Rosaceae) were differently involved in salt tolerance, in which

MhINT4.2 was particularly up-regulated under salt stress (Hu et al., 2020). Furthermore,

transgenic *M. hupehensis* plants overexpressing SLAH3 exhibited increased Cl^- efflux rate

and reduced ROS accumulation after ABA signaling. Likewise, transgenic *A. thaliana* plants

overexpressing RSA1 exhibited enhanced expression of salt overly sensitive 1 (SOS1), a

Na^+/H^+ antiporter crucial for Na^+ homeostasis under salt stress, and genes encoding

oxidoreductase and peroxidase, resulting in Na^+ and ROS detoxification (Guan et al., 2013).

In addition to *RlRSA1* and *RlSLAH3*, the duplications and divergence of *R. lucieae* genes of

leucine aminopeptidase2 (LAP2), L-type lectin-domain containing receptor kinase IX.1

(LRK91), salt tolerance receptor-like cytoplasmic kinase 1 (STRK1), and suppressor of Var 3

(SUV3) are possibly responsible for scavenging and detoxification of ABA-induced ROS.

The various receptor-like kinases such as LRK91 and STRK1 are well known to be involved

in responses to plant abiotic stresses (Ye et al., 2017; Gandhi and Oelmüller, 2023). For

instance, transgenic sweet cherry (*Prunus avium*; Rosaceae) plants overexpressing LRK91

homolog (PaLectinL16) and transgenic *O. sativa* plants over-expressing STRK1 exhibited

enhanced antioxidant ability under salt stress (Zhou et al., 2018; Sun et al., 2022). Likewise,

transgenic *O. sativa* plants overexpressing SUV3 also exhibited reduced ROS production

with increased antioxidant activities, and increased growth rates with higher content of

phytohormones under salt stress (Tuteja et al., 2013; Sahoo et al., 2014). Furthermore, given

the diverse roles of 70 kD heat shock proteins (HSP70s) as molecular chaperones in abiotic

stress tolerance, the increased copy number and genetic diversity of *RlHSP70* possibly

influenced various abiotic stress tolerance in *R. lucieae*, including salt tolerance (Vu et al.,

2023; Pan et al., 2024). Thus, increased gene copy numbers and diversity after duplications of

homeostasis- and detoxification-related genes of *R. lucieae* may have contributed to the

evolution of increased viability and rapid recovery from severe salt stress, resulting in the

vigorous growth of *R. lucieae* in saline environments.

BRs are a group of natural steroid hormones broadly found in plants, which are

considered to be related to plant growth, development and stress response. Although their

roles in plant growth and development are well established, the mechanism of stress

resistance induced by BRs is poorly understood (Krishna, 2003). Nevertheless, several recent

studies suggested BRs enhance the stress resistance with the complex interaction with other

hormones, including cross-talk between BRs and ABA (Krishna, 2003; Choudhary et al.,

2012). Under salt stress, BRs modulate the gene transcription and ROS levels by acting as

signaling molecules and interacting other phytohormones (Anwar et al., 2018; Kumar et al.,

2022). Although not many BR-related genes affecting salinity resistance are known in plants,

pronounced enrichment of BR-related GOs and KEGG pathways suggested the putative

influence of the rapid evolution of BR-related genes on salt tolerance of *R. lucieae* (Fig. 3;

Supplementary Fig. S5; Supplementary Table S7).

Modes of gene duplication in adaptive evolution of Rosa lucieae

There are four different modes of gene duplication known so far: tandem duplication,

whole-genome duplication, chromosomal segmental duplication, and single gene

transposition-duplication (Freeling, 2009). Among those, whole-genome duplication is less

likely to affect the gene duplications and divergence in *R. lucieae*, considering its similar

genome size and nucleic DNA amount compared to its close relatives (Yokoya et al., 2000).

Except for whole-genome duplication, salt tolerant-related genes of *R. lucieae* have been

possibly duplicated through any other modes of duplication. Especially, all the three modes of

duplication are likely to influence the divergence of *RlCNGC1*. The clusters of adjacent

CNGC1 paralogs located on chromosome 1 of *R. lucieae*, which are conserved in Rosoideae

species, may have originated from the ancestral tandem duplication within the Rosoideae

lineage (Supplementary Fig. S6B; Supplementary Table S8). One of these *CNGC1* clusters in

R. lucieae was duplicated into unplaced contig 62, hypothesizing a recent segmental

duplication event in *R. lucieae* probably increased the copy number of *RCNGC1*

(Supplementary Fig. S6B, C). The close genetic distances of *RlCNGC1* genes between the

two clusters on chromosome 1 and unplaced contig 62 corroborated their recent divergence

(Supplementary Fig. S6A). On the other hand, many *CNGC1* genes of which synteny was not

aligned with the synteny of genomic collinear blocks were dispersed on various loci of whole

genomes (Supplementary Fig. S6B). This distribution can stem from transposition-

duplication, which relocates genes to new and arbitrary loci and blurs the syntenic evidence

(Freeling, 2009). The dense distributions of LTR elements and DNA repeat elements near the

duplicated *RlCNGC1* genes on novel positions supported the possible transposition-

duplication of *RlCNGC1* (Supplementary Fig. S6C). The various modes of duplication of

RlCNGC1 suggested that all duplication modes but whole-genome duplication may have

influenced the recent and rapid adaptive evolution of *R. lucieae* in saline environments.

Evolution of other stress tolerance in Rosa lucieae

In addition to salt stress in their natural habitat, wild *R. lucieae* plants are exposed to

various biotic and abiotic stresses, and become adapted and tolerate to these stresses. The

Ca^{2+} and ABA signaling, and ROS accumulation are common processes of responses to

diverse biotic and abiotic stresses (Mauch-Mani and Mauch, 2005; Fujita et al., 2006;

Boudsocq and Sheen, 2010; Brosché et al., 2010; Rock et al., 2010). Moreover, some genes

responsible to salt tolerance are also related to other abiotic stress, especially drought stress,

which induces many common responses with salt stress by causing osmotic stress (Bartels

and Sunkar, 2005). For example, the overexpression of CBL1 enhanced drought tolerance in

transgenic *A. thaliana* plants as well, in addition to salt tolerance (Albrecht et al., 2003;

Cheong et al., 2003). Similarly, closely related genes to putative salt tolerance-responsible

genes in *R. lucieae* may be related to other stress tolerance of *R. lucieae*. For example, similar

to *RlACA13*, the copy numbers of *RlACA10* and *RlACA12* were increased compared to close

relatives of *R. lucieae* (Fig. 5C). Despite the genetic structures and function of Ca^{2+} efflux

similar to ACA13, ACA10 and ACA12 are responsible to plant immune response and defense

to pathogens and insects in *A. thaliana* but not salt tolerance (Frei dit Frey et al., 2012; Yu et

al., 2018; Fotouhi et al., 2022). This result suggested that gene duplications and divergence

also enhanced stress response, signaling and restoration of *R. lucieae* under various biotic and

abiotic stresses.

Aligning with the contribution of *R. lucieae* to the improvement in resistance to

diseases and pathogens in rose breeding (Noack, 2003; Byrne et al., 2005), GOs and KEGG

pathways related to plant immunity and defense mechanism were significantly enriched in the

rapidly evolving gene families of *R. lucieae* (Fig. 3; Supplementary Fig. S5; Supplementary

Table S7). Likewise, the significant enrichment of GOs and KEGG pathways related to

phytohormones and secondary metabolites responsible for plant immunity and defense

supported the enhanced resistance to biotic stresses in *R. lucieae*. The phytohormones

salicylic acid (SA), JA and ethylene are renowned for their roles in biotic stress responses

(Fujita et al., 2006), and GOs related to all these hormones were significantly enriched in the

rapidly evolving gene families of *R. lucieae* (Fig. 3; Supplementary Table S7). Given that SA

and ethylene can interact with BRs to enhance salt tolerance (Divi et al., 2010; Choudhary et

al., 2012), the evolution of salt tolerance may be closely associated with the evolution of

enhanced defense to pathogens and insects in *R. lucieae*. The secondary metabolites including

terpenoids, phenolics, alkaloids and flavonoids are pivotal for plants in defense and

adaptation to environmental stresses (Akula and Ravishankar, 2011; Erb and Kliebenstein,

2020). Particularly, the importance of secondary metabolites in salinity tolerance in plants

was highlighted with their crucial role as antioxidants alleviating oxidative stress induced by

high salinity (Isah, 2019). The recent multi-omics study also emphasized the importance of

flavonoid and flavonol metabolism in response to salt stress in roses (Ren et al., 2024).

Within secondary metabolites, plant volatiles particularly act on attracting pollinators as

floral scent components or deterring herbivory as insect repellents, and non-volatiles are

involved in various defense mechanism distinct from volatiles (Pichersky and Gershenzon,

2002; Unsicker et al., 2009; Tholl, 2015). The earlier comparative genomic investigations of

rose species suggested that the duplication and divergence of volatile terpenoid metabolism-

involved Nudix hydrolase 1 (NUDX1) genes modulate the content of scent compounds of

rose flowers (Conart et al., 2022; Shang et al., 2024). However, the copy numbers of *NUDX1*

were not only decreased in *R. lucieae* (four copies in *R. lucieae*, nine in *R. chinensis*, and 15

in *R. rugosa*), but the amount of volatile compounds was also reduced in flower petals of *R.*

lucieae (Feng et al., 2022; Shang et al., 2024). Moreover, different profiles of paralogous

NUDX1 in involvement in floral scent were identified between *R. chinensis* and hybrid

cultivars of *R. lucieae* (Conart et al., 2023). In rose flowers, the chemical composition of

volatile compounds was notably different among rose species, and the interspecific

relationships based on characteristics of floral scent were inconsistent with the species

phylogeny (Debray et al., 2019; Feng et al., 2022; Noh et al., 2024; Jeon et al., 2024; Shang

et al., 2024), which implied the independent and unique evolution of secondary metabolites in

different rose species. This diversity and complexity of secondary metabolites among closely

related species can be easily affected by environmental factors (Borghi et al. 2019; Paul et al.,

2022). Moreover, the biosynthetic pathways of volatiles are also responsible for production of

various non-volatile secondary metabolites involved in stress resistance (Paul et al., 2022).

Given the enhanced resistance to biotic and abiotic stresses in *R. lucieae*, the unique

biosynthetic mechanism of secondary metabolites in *R. lucieae* may have alternatively

evolved to complement resistance to environmental stresses. For example, the prominently

enriched GOs of saponin and glycoside metabolism in *R. lucieae* may be responsible for

defense to herbivory, diseases and pathogens (Supplementary Table S7; Tholl, 2015). Further

biochemical investigation of *R. lucieae* will contribute to understanding protective

compounds of roses and their mechanisms of defense to diseases, pathogens and other

environmental stresses, promising our understanding of hardiness in rose breeding programs.

18   **DISCUSSION**

Rosa lucieae inhabits open and exposed rocky slopes or banks, especially along the sea

shores and riverside at low elevation (Rehder, 1940; Ohba, 1993; Ku and Robertson, 2003),

where the salts are constantly accumulated in their soils. In these natural habitats, *R. lucieae*

has evolved to adapt to saline environments, resisting to and growing vigorously under salt

stress. Although *R. rugosa* is another rose species renowned for salt tolerance, its earlier

genomic study could not find any difference in genetic component related to salt tolerance,

but only find the duplications of two putative genes (*DRIP2* and *PTR3*) related to water

deprivation (Chen et al., 2021). However, the first chromosome-level genome of wild *R.*

lucieae from the present study shed light on the adaptive evolution to salt stress in roses.

Utilizing high-throughput genome sequencing techniques, the genome of wild *R.*

lucieae was sequenced and assembled into 544.7 Mb-long chromosome-level genomic

sequences. This assembly presents the genomic basis of a wild rose, which provides insights

into the nature of roses. The chromosome-level genome assembly of *R. lucieae* indicated the

conserved genomic structure with other comprehensive rose genomic assembly of *R.*

chinensis and *R. rugosa* (Supplementary Fig. S3). Nevertheless, coupled with the increased

genome size, the number of protein-coding genes within the genome of *R. lucieae* was

increased with gene duplications (Table 1). This remarkable gene duplication was identified

as a major driving force behind the adaptive evolution in salt tolerance of *R. lucieae*. The

diverse genes responsible for various processes of salt tolerance in plants were duplicated in

R. lucieae, including Ca^{2+} (*BON*, *CBL1*, *CNGC1* and *CSC1*), abscisic acid (*BON*, *CIP1*,

LRK10L1.2, *PYL9*, *SD25* and *ZEP*), and ROS signaling (*IP5P7*, *LRX4* and *RBOH*), ion

homeostasis (*ACA13*, *CML18*, *HKT1*, *INT4*, *RSA1* and *SLAH3*), ROS detoxification (*LAP2*,

LRK91, *SLAH3*, *STRK1* and *SUV3*), protein repair (*HSP70*), and growth recovery (*LRX4* and

SUV3) (Table 2; Supplementary Fig. S7). Moreover, the rapid evolution of gene families

related to phytohormones including BRs, SA and ethylene also possibly enhanced tolerance

to various stresses including salinity. The unique evolution in composition of secondary

metabolites may have allowed the adaptation to environmental stresses and defense to

diseases and pathogens. These results substantiate an important role of gene duplication in the

adaptive evolution of plants, and shed light on the nature of adaptive evolution of *R. lucieae*

in saline environments. The genomic basis of salt tolerance of *R. lucieae* will pave the way

for the advancement in rose breeding programs by contributing to enhanced stress resistance

in rose cultivation. Particularly, understanding genomic mechanism of resistance to

environmental stresses in roses and its utilization will significantly contribute to future

horticultural and agricultural industries with recovered or enhanced productivity and quality

of rose flowers or other rosaceous crops in ongoing global urbanization and climate changes.

A further comparative analysis of expression profiles in *R. lucieae* will corroborate influence

of the putative salt tolerance-responsible genes on the adaptation of *R. lucieae* to saline

environments. Additionally, further genome-wide association studies and comparative

genomic studies including diverse *R. lucieae* accessions from various sources will unveil the

influence of genetic mutations affecting salt tolerance in roses.

11   **MATERIALS AND METHODS**

Sample collection and preparation

To develop a genomic sequence of *R. lucieae*, a wild individual of *R. lucieae* (accession

ID: SG001; NCBI BioSample accession: SAMN39614440) was collected from a coastal field

in Jeju, Korea, and its voucher specimen was deposited into the Ha Eun Herbarium (SKK) of

Sungkyunkwan University. Genomic DNA (gDNA) was isolated from fresh young leaves of

the accession using the Exgene Plant SV mini kit (GeneAll Biotechnology Co., Ltd., Seoul,

Korea), following the manufacturer's instructions. For the quality control, the isolated gDNA

was quantified and assessed using QuantiFluor dsDNA System (Promega Corp., WI, USA)

and the 2100 Bioanalyzer System (Agilent Technologies, Inc., CA, USA). Additionally,

flower buds (BioSample accession: SAMN40548981) and young leaves (BioSample

accession: SAMN40548982) were collected from the same individual to provide evidence for

the splice site prediction. The samples were stored in RNAlater solution (Thermo Fisher

Scientific Inc., MA, USA) immediately after the collection to stabilize their RNAs. Total

RNAs were isolated using Maxwell RSC Plant RNA Kit and Instrument (Promega) by

Macrogen, Inc. (Seoul, Korea). For the quality control, the isolated RNAs were quantified

and assessed using Quant-iT RiboGreen RNA Assay Kit (Thermo Fisher Scientific) and the

2100 Bioanalyzer System.

Molecular sequencing and genome assembly

The gDNA was prepared into respective libraries for Illumina HiSeq short-read and

PacBio SMRT long-read sequencing platforms. The short-read sequencing library was

prepared by random fragmentation, followed by 5′- and 3′-adapter ligations, using the TruSeq

Nano DNA kit (Illumina, Inc., CA, USA). The fragmented library was sequenced via

Illumina HiSeq platform (Illumina) by Macrogen after the size selection. The PacBio

SMRTbell library was prepared using SMRTbell Express Template Prep Kit (Pacific

Biosciences of California, Inc., CA, USA), following the manufacturer's instructions. The

large fragments longer than 20 kb were selected using BluePippin system (Sage Science, Inc.,

MA, USA), and the large-insert library was sequenced via PacBio Sequel platform (Pacific

Biosciences) by DNALink, Inc. (Seoul, Korea). The total RNAs were prepared into

sequencing libraries by purifying poly-A containing mRNAs, random fragmentation, cDNA

synthesis, and 5′- and 3′-adapter ligations, using the TruSeq Stranded mRNA kit (Illumina).

The fragmented libraries were sequenced via Illumina NovaSeq platform (Illumina) by

Macrogen.

The genome size, heterozygosity and repeat content of *R. lucieae* were estimated with

k-mer analysis using Jellyfish v2.1.3 (Marçais and Kingsford, 2011) and GenomeScope v2.0

(Ranallo-Benavidez et al., 2020), after preprocessing short reads by trimming the sequencing

adapters, PCR dimers, and low-quality sequences using BBDuk v38.26 (Bushnell, 2014) and

removing contaminant reads by mapping reads to NCBI RefSeq complete bacterial, viral and

organellar genomes using bowtie2 v2.3.4 (Langmead and Salzberg, 2012). PacBio subreads

were error-corrected into consensus reads using SMRT Link v2.3 (Pacific Biosciences), and

preprocessed short reads were mapped to the consensus reads to remove contaminant reads

using bwa v0.7 (Li, 2013). The preprocessed PacBio consensus reads were *de novo*

assembled into primary contigs using a diploid-aware assembler FALCON, and the primary

contigs were phased and updated using a diploid assembler FALCON-Unzip (Chin et al.,

2016). The phased primary contigs of the diploid genome assembly were curated to remove

haplotigs to complete the draft genome assembly using Purge haplotigs v1.1.1 (Roach et al.,

2018). Error sequences in the draft genome assembly were corrected using Pilon v1.23, after

mapping preprocessed short reads to the draft genome assembly using bwa aligner (Li, 2013;

Walker et al., 2014). The *de novo* genome assembly was assessed for the completeness based

on the universal single-copy orthologs using BUSCO v5.3.0 and eudicots_odb10,

embryophyta_odb10, and viridiplantae_odb10 datasets (Manni et al., 2021). The *de novo*

genome assembly was scaffolded into chromosomes using RagTag v2.1.0 (Alonge et al.,

2022), referenced from the chromosome-level assembly of *R. lucieae* 'Basye's Thornless'

(NCBI GenBank accession: GCA_024704745.1; Zhong et al., 2021). The scaffolded

chromosomes were assessed with the genome collinearity analysis with the reference

assembly using MUMmer4 v4.0.0rc1 (Marçais et al., 2018), examining any genomic gaps or

putative misassembled contigs.

Genome annotation

The repeat sequences of genome of *R. lucieae* were *de novo* predicted and developed

into the *de novo* repeat library using RepeatModeler v1.0.8 (Smit and Hubley, 2008),

incorporating complementary repeat annotation methods like RECON v1.08 (Bao and Eddy,

2002), RepeatScout v1.0.5 (Price et al., 2005), and TRF v4.04 (Benson, 1999). The repeat

regions of draft genome were identified, classified into subclasses based on the Dfam

database v2.0 (Hubley et al., 2016) and RepBase database v23.10 (Bao et al., 2015), and

masked using RepeatMasker v4.0.7 (Smit et al., 2013).

Within the genome of *R. lucieae*, protein-coding genes were predicted and annotated by

integrating *ab initio*, RNA-seq evidenced, and homology-based gene predictions. For *ab*

initio gene prediction, the protein-coding gene models of rosaceous reference genomes of *R.*

chinensis 'Old Blush' (RefSeq accession: GCF_002994745.2; Raymond et al., 2018), *R.*

rugosa (GCF_958449725.1; Ruhsam et al., 2024), *F. vesca* (GCF_000184155.1; Shulaev et

al., 2011), *Prunus persica* 'Lovell' (GCF_000346465.2; IPGI, 2013), and *Malus domestica*

'Golden Delicious' (GCF_002114115.1; Daccord et al., 2017) were trained and optimized

respectively using AUGUSTUS v3.1.0 (Stanke and Waack, 2003). With meta-parameters of

rosaceous reference gene models, the *ab initio* gene model of *R. lucieae* was predicted within

the draft genome using AUGUSTUS v3.1 (Stanke and Waack, 2003). For homology-based

prediction, protein-coding genes from reference genomes (Supplementary Table S6) were

aligned to the draft genome of *R. lucieae* to identify homologous regions using Exonerate

v2.4.0 (Slater and Birney, 2005). For RNA-seq evidenced prediction, each RNA-seq

accession was aligned to the draft genome of *R. lucieae* using STAR v2.5.1a (Dobin et al.,

2013). Leveraging the RNA-seq alignment, reference gene models of *R. chinensis*, *R. rugosa*,

and *F. vesca*, and evidence from *ab initio* and homology-based gene predictions, an integrated

consensus gene model was predicted using GeMoMa v1.9 (Keilwagen et al., 2018) based on

conserved coding-sequence and intron positions.

A set of protein-coding genes of *R. lucieae* were annotated in their functions. The GOs

(Ashburner et al., 2000) and definitions were annotated using blastp v2.15.0 (Camacho et al.,

2009), Blast2GO v6.0.3 (Conesa et al., 2005), InterProScan v5.52 (Jones et al., 2014), and

Trinotate v4.0.2 (Bryant et al., 2017), based on public databases including NCBI non-

redundant protein sequence (nr; accessed on 12 Feb 2024), Gene Ontology Annotation

(GOA; v2023.08; Camon et al., 2004), Pfam (v36.0; El-Gebali et al., 2019), KEGG

(Kanehisa et al., 2012), UniProtKB/Swiss-Prot (v2024_01; Boutet et al., 2007), and EggNOG

(v5.0; Huerta-Cepas et al., 2019).

Comparative genomic and evolutionary analyses

To infer the genetic evolution of *R. lucieae*, the orthologous gene groups were

identified from the gene models of *R. lucieae* and other reference species including *R.*

chinensis, *R. rugosa*, *F. vesca*, *P. persica*, *M. domestica*, and *P. trichocarpa* (RefSeq

accession: GCF_000002775.4; Tuskan et al., 2006) using OrthoMCL v2.0.9 (Li et al., 2003).

The phylogenetic inference and divergence time estimation were analyzed with conserved

single-copy orthologs. Divergence times of *R. lucieae* and reference species were estimated

by running two Markov chain Monte Carlo chains for 10,000,000 generations using BEAST2

v2.6.3 (Bouckaert et al., 2014) with the partitioned alignment and JTT amino-acid exchange

rate matrix from IQ-TREE v2.2.0 (Chernomor et al., 2016) and ModelFinder

(Kalyaanamoorthy et al., 2017), discarding the first 10 % of trees as post-burn-in. The

divergence time estimation was calibrated using the oldest fossil records within Rosaceae

(113.0–100.5 Mya; Peppe et al., 2008) and Eurosids (145.0–100.5 Mya; Ward and Jenney,

1899; Mellon et al., 1963). Based on identified orthologous groups and estimated divergence

time, gene family evolution in *R. lucieae* and other rosaceous species was inferred using

CAFE v5.1.0 (Mendes et al., 2020), identifying gene families with significant family size

change ($p < 0.01$) as rapidly evolving gene families. To infer the significantly evolved gene

functions in *R. lucieae*, GO and KO enrichment analyses on expanded or contracted gene

families were performed using the R package clusterProfiler v4.4.4 (Wu et al., 2021).

The genomic synteny among the genome of *R. lucieae* and the reference genomes of its

closely related species in Rosoideae (i.e., *R. chinensis*, *R. rugosa*, and *F. vesca*) was analyzed

to infer the structural genomic variation within their evolution. The pairwise genetic

homologies between species were assessed using blastp, and homology-based collinear

blocks were identified using MCScanX (Wang et al., 2012). The collinearity blocks between

species were visualized using SynVisio (Bandi and Gutwin, 2020). To detect the gene

duplications within the genome of *R. lucieae*, the paralogous collinear blocks were also

identified using blastp and MCscanX.

To understand the sequence divergences of homologous genes and their duplications

within the Rosoideae lineage, Ks within paralogs of a species or orthologs between a pair of

species were calculated using ParaAT v2.0 (Zhang et al., 2012), incorporating sequence

alignments from MAFFT and Ks calculation from KaKs_Calculator v3.0 (Zhang, 2022). For

Ks calculation, pairs of homologous genes were identified with percent identity scores over

70 % and percent positive scores over 80 % in all-to-all pairwise blastp comparisons of genes

in each orthologous group. Particularly, the Ks of homologous pairs of genes annotated with

the GO term for 'response to salt stress' or its child terms were calculated to understand the

evolution of salt stress-related genes in *R. lucieae*.

Gene duplication and divergence analysis

The duplications and divergence of genes in rapidly evolving gene families or

annotated with gene functions involved in salt tolerance were explored to elucidate the

influence of gene duplication and diversification in the evolution of salt tolerance of *R.*

lucieae. The genes responsible for salt tolerance were compiled based on the broad

investigation of plant salt-stress resistance (Botella et al., 2005; Munns, 2005; Gandhi and

Oelmüller, 2023). The candidate genes were identified by their functional annotation based

on the UniProtKB/Swiss-Prot and NCBI nr databases (E-value ≤ 1E–5), followed by the

protein family and domain searches using HMMsearch v3.4 (Eddy, 2011) based on the Pfam

database (E-value ≤ 1E–3). For example, ion transport protein family (PF00520) and cyclic

nucleotide-binding domain (PF00027) were searched for the identification of CNGC1 genes,

and E1-E2 ATPase (PF00122) or cation transport ATPase (P-type) (PF13246) families and

cation transporting ATPase C-terminus (PF00689) or N-terminus (PF00690) domains were

searched for the identification of ACA and ECA genes.

Among Rosoideae species, the genealogies of CNGC1 and ACA genes, of which the

numbers of copies were notably increased in *R. lucieae*, were respectively inferred using IQ-

TREE based on maximum-likelihood approaches with the JTT+F+R5 (CNGC1) and

JTT+I+G4 (ACAs) substitution models selected by ModelFinder, after aligning amino-acid

sequences using MAFFT. The robustness of each tree was assessed by 5000 replicates of

bootstrap approximation using UFBoot2 (Hoang et al., 2018). The structural motifs enriched

in Rosoideae CNGC1 or ACAs were identified using STREME v5.5.7 (Bailey, 2021), with

the minimum motif width of 6 aa and the *p*-value cutoff of 0.01. The pairwise synteny of

orthologous genes was identified using blastp by finding the best match between a pair of

species.

22   **ACKNOWLEDGEMENTS**

23   This study was supported by the National Research Foundation of Korea [grant number

24   NRF-2017H1A2A1045787] to the first author.

AUTHOR CONTRIBUTIONS

3   J.-H.J., Y.S. and S.-C.K. designed and coordinated the study. J.-H.J. and S.-C.K.

4 collected the materials. J.-H.J., M.J. and Y.S. analyzed molecular data. J.-H.J. and S.-C.K.

5 wrote the manuscript. The manuscript was reviewed and accepted by all the authors.

DATA AVAILABILITY

8   Molecular sequencing data and genome assembly generated in this study were

9 deposited into NCBI under BioProject accession PRJNA1069305.

CONFLICT OF INTERESTS

12   The authors declare that there are no conflicts of interest.

SUPPLEMENTARY DATA

Supplementary data is available at online in the Supplemental Material section.

Fig. S1. *K*-mer distribution in the short-read sequencing data of *Rosa lucieae*.

Fig. S2. Syntenic dotplot of two *R. lucieae* assemblies.

Fig. S3. Pairwise genome synteny of *R. lucieae* and its close relatives

Fig. S4. Comparison of pairwise Ks distribution of orthologous genes among the Rosoideae

species.

Fig. S5. KEGG pathway enrichment analysis on *Rosa lucieae* genes within the rapidly

evolving gene families based on the gene family evolution analysis.

Fig. S6. Duplication and divergence of CNGC1 genes within Rosoideae.

Fig. S7. Duplication of candidates of salt tolerance-related genes in *R. lucieae*.

Table S1. Summary of whole-genome and transcriptome sequencing of *R. lucieae* and

subsequent read preprocessing.

Table S2. Statistics of the de novo genome assembly of *R. lucieae*.

Table S3. Statistics of the genome scaffolding of *R. lucieae* assembly.

Table S4. Repeat sequences within the genome of *R. lucieae*.

Table S5. Statistics of functional annotation of protein-coding genes from the genome of

R. lucieae.

Table S6. List of the reference genomes and their gene models used for the homology-based

gene prediction of *R. lucieae*.

Table S7. List of gene ontologies enriched in rapidly evolving gene families of *R. lucieae*.

Table S8. List of candidate genes related to salt tolerance in *R. lucieae*.

REFERENCES

Akula R, Ravishankar GA. 2011. Influence of abiotic stress signals on secondary metabolites in plants. *Plant Signaling & Behavior* 6(11): 1720–1731.

Albrecht V, Weinl S, Blazevic D, et al. 2003. The calcium sensor CBL1 integrates plant responses to abiotic stresses. *The Plant Journal* 36(4): 457–470.

Alonge M, Lebeigle L, Kirsche M, et al. 2022. Automated assembly scaffolding using RagTag elevates a new tomato system for high-throughput genome editing. *Genome Biology* 23(1): 258.

Anwar A, Liu Y, Dong R, Bai L, Yu X, Li Y. 2018. The physiological and molecular mechanism of brassinosteroid in response to stress: a review. *Biological Research* 51: 46.

Ashburner M, Ball CA, Blake JA, et al. 2000. Gene Ontology: tool for the unification of biology. *Nature Genetics* 25(1): 25–29.

Bailey TL. 2021. STREME: accurate and versatile sequence motif discovery. *Bioinformatics* 37(18): 2834–2840.

Bandi V, Gutwin C. 2020. Interactive exploration of genomic conservation. In: *Proceedings of the 46th Graphics Interface Conference on Proceedings of Graphics Interface 2020 (GI'20)*. Waterloo: Canadian Human-Computer Communications Society.

Bao W, Kojima KK, Kohany O. 2015. Repbase Update, a database of repetitive elements in eukaryotic genomes. *Mobile DNA* 6: 1–6.

Bao Y, Chen C, Fu L, Chen Y. 2021. Comparative transcriptome analysis of *Rosa chinensis* 'Old Blush' provides insights into the crucial factors and signaling pathways in salt stress response. *Agronomy Journal* 113(4): 3031–3050.

Bao Z, Eddy SR. 2002. Automated de novo identification of repeat sequence families in sequenced genomes. *Genome Research* 12(8): 1269–1276.

Bartels D, Sunkar R. 2005. Drought and salt tolerance in plants. *Critical Reviews in Plant Sciences* 24(1): 23–58.

Begum K, Das A, Ahmed R, Akhtar S, Kulkarni R, Banu S. 2024. Genome-wide analysis of respiratory burst oxidase homolog (*Rboh*) genes in *Aquilaria* species and insight into ROS-mediated metabolites biosynthesis and resin deposition. *Frontiers in Plant Science* 14: 1326080.

Benson G. 1999. Tandem repeats finder: a program to analyze DNA sequences. *Nucleic Acids Research* 27(2): 573–580.

Berthomieu P, Conéjéro G, Nublat A, et al. 2003. Functional analysis of *AtHKT1* in Arabidopsis shows that Na$^+$ recirculation by the phloem is crucial for salt tolerance. *The EMBO Journal* 22(9): 2004–2014.

Borghi M, Perez de Souza L, Yoshida T, Fernie AR. 2019. Flowers and climate change: a metabolic perspective. *New Phytologist* 224(4): 1425–1441.

Bose J, Pottosin II, Shabala SS, Palmgren MG, Shabala S. 2011. Calcium efflux systems in stress signaling and adaptation in plants. *Frontiers in Plant Science* 2: 85.

Botella MA, Rosado A, Bressan RA, Hasegawa PM. 2005. Plant adaptive responses to salinity stress. In: Jenks MA, Hasegawa PM (eds.). *Plant Abiotic Stress*. Oxford: Blackwell Publishing, 37–70.

Bouckaert R, Heled J, Kühnert D, et al. 2014. BEAST 2: a software platform for Bayesian evolutionary analysis. *PLoS Computational Biology* 10(4): e1003537.

Boudsocq M, Sheen J. 2010. Stress signaling □: calcium sensing and signaling. In: Pareek A,

Sopory SK, Bohnert HJ, Govindjee (eds.). *Abiotic Stress Adaptation in Plants: physiological, molecular and genomic foundation*. Dordrecht: Springer, 75–90.

Boutet E, Lieberherr D, Tognolli M, Schneider M, Bairoch A. 2007. UniProtKB/Swiss-Prot: the manually annotated section of the UniProt KnowledgeBase. In: Edwards D (ed.). *Plant Bioinformatics: Methods and Protocols*. Totowa: Humana Press, 89–112.

Brosché M, Overmyer K, Wrzaczek M, Kangasjärvi J, Kangasjärvi S. 2010. Stress signaling □: reactive oxygen species (ROS). In: Pareek A, Sopory SK, Bohnert HJ, Govindjee (eds.). *Abiotic Stress Adaptation in Plants: physiological, molecular and genomic foundation*. Dordrecht: Springer, 91–102.

Bryant DM, Johnson K, DiTommaso T, et al. 2017. A tissue-mapped axolotl de novo transcriptome enables identification of limb regeneration factors. *Cell Reports* 18(3): 762–776.

Bushnell B. 2014. *BBMap: a fast, accurate, splice-aware aligner*. Berkeley: Lawrence Berkeley National Lab., United States.

Cabrera RI, Perdomo P. 2003. Reassessing the salinity tolerance of greenhouse roses under soilless production conditions. *HortScience* 38(4): 533–536.

Cabrera RI, Solís-Pérez AR, Sloan JJ. 2009. Greenhouse rose yield and ion accumulation responses to salt stress as modulated by rootstock selection. *HortScience* 44(7): 2000–2008.

Cai X, Niu G, Starman T, Hall C. 2014. Response of six garden roses (*Rosa × hybrida* L.) to salt stress. *Scientia Horticulturae* 168: 27–32.

Cairns T. 2003. Classification: Horticultural Classification Schemes. In: Roberts AV, Debener T, Gudin S (eds.). *Encyclopedia of Rose Science*. Oxford: Academic Press, 117–124.

Camacho C, Coulouris G, Avagyan V, et al. 2009. BLAST+: architecture and applications. *BMC Bioinformatics* 10: 1–9.

Camon E, Magrane M, Barrell D, et al. 2004. The Gene Ontology Annotation (GOA) Database: sharing knowledge in UniProt with Gene Ontology. *Nucleic Acids Research* 32(Suppl_1): D262–D266.

Chen F, Su L, Hu S, et al. 2021. A chromosome-level genome assembly of rugged rose (*Rosa rugosa*) provides insights into its evolution, ecology, and floral characteristics. *Horticulture Research* 8: 141.

Chen K, Gao J, Sun S, et al. 2020. BONZAI proteins control global osmotic stress responses in plants. *Current Biology* 30(24): 4815–4825.

Cheong YH, Kim KN, Pandey GK, Gupta R, Grant JJ, Luan S. 2003. CBL1, a calcium sensor that differentially regulates salt, drought, and cold responses in Arabidopsis. *The Plant Cell* 15(8): 1833–1845.

Chernomor O, Von Haeseler A, Minh BQ. 2016. Terrace aware data structure for phylogenomic inference from supermatrices. *Systematic Biology* 65(6): 997–1008.

Chin CS, Peluso P, Sedlazeck FJ. 2016. Phased diploid genome assembly with single-molecule real-time sequencing. *Nature Methods* 13(12): 1050–1054.

Chinnusamy V, Zhu JK. 2003. Plant salt tolerance. In: Hirt H, Shinozaki K (eds.). *Plant Responses to Abiotic Stress*. Berlin: Springer, 241–270.

Choudhary SP, Yu JQ, Yamaguchi-Shinozaki K, Shinozaki K, Tran LSP. 2012. Benefits of brassinosteroid crosstalk. *Trends in Plant Science* 17(10): 594–605.

Conart C, Saclier N, Foucher F, et al. 2022. Duplication and specialization of *NUDX1* in Rosaceae led to geraniol production in rose petals. *Molecular Biology and Evolution*

39(2): msac002.

Conart C, Saclier N, Foucher F, et al. 2023. Evolution of scent genes in roses. *Acta Horticulturae* 1368: 361–370.

Conesa A, Götz S, García-Gómez JM, Terol J, Talón M, Robles M. 2005. Blast2GO: a universal tool for annotation, visualization and analysis in functional genomics research. *Bioinformatics* 21(18): 3674–3676.

Corwin DL. 2021. Climate change impacts on soil salinity in agricultural areas. *European Journal of Soil Science* 72(2): 842–862.

Crespel L, Mouchotte J. 2003. Breeding: Methods of Cross-Breeding. In: Roberts AV, Debener T, Gudin S (eds.). *Encyclopedia of Rose Science*. Oxford: Academic Press, 30–33.

Daccord N, Celton JM, Linsmith G, et al. 2017. High-quality *de novo* assembly of the apple genome and methylome dynamics of early fruit development. *Nature Genetics* 49(7): 1099–1106.

Debener T, Hibrand-Saint Oyant L. 2009. Genetic engineering and tissue culture of roses. In: Folta KM, Gardiner SE (eds.). *Genetics and Genomics of Rosaceae*. New York: Springer, 393–409.

Debener T, Linde M. 2009. Exploring complex ornamental genomes: the rose as a model plant. *Critical Reviews in Plant Sciences* 28(4): 267–280.

Debray K, Marie-Magdelaine J, Ruttink T, Clotault J, Foucher F, Malécot V. 2019. Identification and assessment of variable single-copy orthologous (SCO) nuclear loci for low-level phylogenomics: a case study in the genus *Rosa* (Rosaceae). *BMC Evolutionary Biology* 19: 152.

Divi UK, Rahman T, Krishna P. 2010. Brassinosteroid-mediated stress tolerance in Arabidopsis shows interactions with abscisic acid, ethylene and salicylic acid pathways. *BMC Plant Biology* 10: 1–14.

Dobin A, Davis CA, Schlesinger F, et al. 2013. STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* 29(1): 15–21.

Eddy SR. 2011. Accelerated profile HMM searches. *PLoS Computational Biology* 7(10): e1002195.

Erb M, Kliebenstein DJ. 2020. Plant secondary metabolites as defenses, regulators, and primary metabolites: the blurred functional trichotomy. *Plant Physiology* 184(1): 39–52.

Feng D, Jian H, Zhang H, et al. 2022. Comparison of volatile compounds between wild and cultivated roses. *HortScience* 57(5): 657–663.

Fotouhi N, Fischer-Stettler M, Lenzoni G, et al. 2022. ACA pumps maintain leaf excitability during herbivore onslaught. *Current Biology* 32(11): 2517–2528.

Foucher F. 2009. Functional genomics in rose. In: Folta KM, Gardiner SE (eds). *Genetics and Genomics of Rosaceae*. New York: Springer, 381–392.

Fougère-Danezan M, Joly S, Bruneau A, Gao XF, Zhang LB. 2015. Phylogeny and biogeography of wild roses with specific attention to polyploids. *Annals of Botany* 115(2): 275–291.

Freeling M. 2009. Bias in plant gene content following different sorts of duplication: tandem, whole-genome, segmental, or by transposition. *Annual Review of Plant Biology* 60(1): 433–453.

Frei dit Frey N, Mbengue M, Kwaaitaal M, et al. 2012. Plasma membrane calcium ATPases are important components of receptor-mediated signaling in plant immune responses and

development. *Plant Physiology* 159(2): 798–809.

Fujita M, Fujita Y, Noutoshi Y, et al. 2006. Crosstalk between abiotic and biotic stress responses: a current view from the points of convergence in the stress signaling networks. *Current Opinion in Plant Biology* 9(4): 436–442.

Gahlaut V, Kumari P, Jaiswal V, Kumar S. 2021. Genetics, genomics and breeding in *Rosa* species. *The Journal of Horticultural Science and Biotechnology* 96(5): 545–559.

Gandhi A, Oelmüller R. 2023. Emerging roles of receptor-like protein kinases in plant response to abiotic stresses. *International Journal of Molecular Sciences* 24(19): 14762.

Gerard JN. 1897. Rose Hybrids. In: Sargent CS (ed.). *Garden and Forest: A Journal of Horticulture, Landscape Art and Forestry*, Vol. 10. New York: The Garden and Forest Publishing Co., 147.

Ghassemi F, Jakeman AJ, Nix HA. 1995. *Salinisation of Land and Water Resources: human causes, extent, management and case studies*. Wallingford: CAB International.

Guan Q, Wu J, Yue X, Zhang Y, Zhu J. 2013. A nuclear calcium-sensing pathway is critical for gene regulation and salt stress tolerance in *Arabidopsis*. *PLoS Genetics* 9(8): e1003755.

Gudin S. 2003. Breeding: Overview. In: Roberts AV, Debener T, Gudin S (eds.). *Encyclopedia of Rose Science*. Oxford: Academic Press, 25–30.

Hibrand Saint-Oyant L, Ruttink T, Hamama L, et al. 2018. A high-quality genome sequence of *Rosa chinensis* to elucidate ornamental traits. *Nature Plants* 4(7): 473–484.

Hoang DT, Chernomor O, Von Haeseler A, Minh BQ, Vinh LS. 2018. UFBoot2: improving the ultrafast bootstrap approximation. *Molecular Biology and Evolution* 35(2): 518–522.

Hou C, Tian W, Kleist T, et al. 2014. DUF221 proteins are a family of osmosensitive calcium-permeable cation channels conserved across eukaryotes. *Cell Research* 24(5): 632–635.

Hu L, Zhou K, Yang S, et al. 2020. MdINT1 enhances apple salinity tolerance by regulating the antioxidant system, homeostasis of ions, and osmosis. *Plant Physiology and Biochemistry* 154: 689–698.

Hubley R, Finn RD, Clements J, et al. 2016. The Dfam database of repetitive DNA families. *Nucleic Acids Research* 44(D1): D81–D89.

Huda KMK, Banu MSA, Tuteja R, Tuteja N. 2013. Global calcium transducer P-type Ca^{2+}-ATPases open new avenues for agriculture by regulating stress signalling. *Journal of Experimental Botany* 64(11): 3099–3109.

Huerta-Cepas J, Szklarczyk D, Heller D, et al. 2019. eggNOG 5.0: a hierarchical, functionally and phylogenetically annotated orthology resource based on 5090 organisms and 2502 viruses. *Nucleic Acids Research* 47(D1): D309–D314.

International Peach Genome Initiative (IPGI). 2013. The high-quality draft genome of peach (*Prunus persica*) identifies unique patterns of genetic diversity, domestication and genome evolution. *Nature Genetics* 45(5): 487–494.

Isah T. 2019. Stress and defense responses in plant secondary metabolites production. *Biological Research* 52: 39.

Jeon JH, Maki M, Chiang YC, Kim SC. 2024. Inferring complex evolutionary history of the closely related East Asian wild roses in *Rosa* sect. *Synstylae* (Rosaceae) based on genomic evidence from conserved orthologues. *Annals of Botany*: mcae170. (in press)

Jha SK, Sharma M, Pandey GK. 2016. Role of cyclic nucleotide gated channels in stress management in plants. *Current Genomics* 17(4): 315–329.

Jones P, Binns D, Chang HY, et al. 2014. InterProScan 5: genome-scale protein function classification. *Bioinformatics* 30(9): 1236–1240.

Kader MA, Lindberg S. 2010. Cytosolic calcium and pH signaling in plants under salinity stress. *Plant Signaling & Behavior* 5(3): 233–238.

Kalyaanamoorthy S, Minh BQ, Wong TK, Von Haeseler A, Jermiin LS. 2017. ModelFinder: fast model selection for accurate phylogenetic estimates. *Nature Methods* 14(6): 587–589.

Kanehisa M, Goto S, Sato Y, Furumichi M, Tanabe M. 2012. KEGG for integration and interpretation of large-scale molecular data sets. *Nucleic Acids Research* 40(D1): D109–D114.

Kaye Y, Golani Y, Singer Y, et al. 2011. Inositol polyphosphate 5-phosphatase7 regulates the production of reactive oxygen species and salt tolerance in Arabidopsis. *Plant Physiology* 157(1): 229–241.

Keilwagen J, Hartung F, Paulini M, Twardziok SO, Grau J. 2018. Combining RNA-seq data and homology-based gene prediction for plants, animals and fungi. *BMC Bioinformatics* 19: 1–12.

Kondrashov FA. 2012. Gene duplication as a mechanism of genomic adaptation to a changing environment. *Proceedings of the Royal Society B: Biological Sciences* 279(1749): 5048–5057.

Krishna P. 2003. Brassinosteroid-mediated stress responses. *Journal of Plant Growth Regulation* 22: 289–297.

Ku TC, Robertson KR. 2003. *Rosa*. In: Wu Z, Raven PH, Hong D (eds.). *Flora of China. Volume 9: Pittosporaceae through Connaraceae*. Beijing: Science Press, 339–381.

Kugler A, Köhler B, Palme K, Wolff P, Dietrich P. 2009. Salt-dependent regulation of a CNG

channel subfamily in Arabidopsis. *BMC Plant Biology* 9: 1–11.

Kumar V, Singh P, Sharma J, Sharma A. 2022. Role of Brassinosteroids in plants responses to salinity stress: a review. *Journal of Applied and Natural Science* 14(2): 582–599.

Kwak JM, Mori IC, Pei ZM, et al. 2003. NADPH oxidase *AtrbohD* and *AtrbohF* genes function in ROS☐dependent ABA signaling in *Arabidopsis*. *The EMBO Journal* 22(11): 2623–2633.

Langmead B, Salzberg SL. 2012. Fast gapped-read alignment with Bowtie 2. *Nature Methods* 9(4): 357–359.

Leus L, Van Laere K, De Riek J, Van Huylenbroeck J. 2018. Rose. In: Van Huylenbroeck J (ed.). *Ornamental Crops*. Cham: Springer International Publishing, 719–767.

Li CH, Tien HJ, Wen MF, Yen HE. 2021. *Myo*☐inositol transport and metabolism participate in salt tolerance of halophyte ice plant seedlings. *Physiologia Plantarum* 172(3): 1619–1629.

Li H. 2013. Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM. *arXiv*: 1303.3997. (Preprint: not peer reviewed)

Li L, Stoeckert CJ, Roos DS. 2003. OrthoMCL: identification of ortholog groups for eukaryotic genomes. *Genome Research* 13(9): 2178–2189.

Lim CW, Yang SH, Shin KH, Lee SC, Kim SH. 2015. The AtLRK10L1.2, *Arabidopsis* ortholog of wheat LRK10, is involved in ABA-mediated signaling and drought resistance. *Plant Cell Reports* 34: 447–455.

Ma W, Ali R, Berkowitz GA. 2006. Characterization of plant phenotypes associated with loss-of-function of AtCNGC1, a plant cyclic nucleotide gated cation channel. *Plant Physiology and Biochemistry* 44(7–9): 494–505.

Maathuis FJM, Filatov V, Herzyk P, et al. 2003. Transcriptome analysis of root transporters reveals participation of multiple gene families in the response to cation stress. *The Plant Journal* 35(6): 675–692.

Magadum S, Banerjee U, Murugan P, Gangapur D, Ravikesavan R. 2013. Gene duplication as a major force in evolution. *Journal of Genetics* 92(1): 155–161.

Manni M, Berkeley MR, Seppey M, Simão FA, Zdobnov EM. 2021. BUSCO update: novel and streamlined workflows along with broader and deeper phylogenetic coverage for scoring of eukaryotic, prokaryotic, and viral genomes. *Molecular Biology and Evolution* 38(10): 4647–4654.

Marçais G, Delcher AL, Phillippy AM, et al. 2018. MUMmer4: A fast and versatile genome alignment system. *PLoS Computational Biology* 14(1): e1005944.

Marçais G, Kingsford C. 2011. A fast, lock-free approach for efficient parallel counting of occurrences of *k*-mers. *Bioinformatics* 27(6): 764–770.

Marriott M. 2003. History of roses in cultivation: Modern (Post–1800). In: Roberts AV, Debener T, Gudin S (eds.). *Encyclopedia of Rose Science*. Oxford: Academic Press, 402–409.

Mauch-Mani B, Mauch F. 2005. The role of abscisic acid in plant–pathogen interactions. *Current Opinion in Plant Biology* 8(4): 409–414.

Mellon GB, Wall JH, Stelck CR. 1963. Lower Cretaceous section, Belcourt Ridge, northeastern British Columbia. *Bulletin of Canadian Petroleum Geology* 11(1): 64–72.

Mendes FK, Vanderpool D, Fulton B, Hahn MW. 2020. CAFE 5 models variation in evolutionary rates among gene families. *Bioinformatics* 36(22-23): 5516–5518.

Monihan SM, Ryu CH, Magness CA, Schumaker KS. 2019. Linking duplication of a calcium

sensor to salt tolerance in *Eutrema salsugineum*. *Plant Physiology* 179(3): 1176–1192.

Munns R. 2005. Genes and salt tolerance: bringing them together. *New Phytologist* 167(3): 645–663.

Munns R, Gilliham M. 2015. Salinity tolerance of crops – what is the cost? *New Phytologist* 208(3): 668–673.

Munns R, Tester M. 2008. Mechanisms of salinity tolerance. *Annual Review of Plant Biology* 59: 651–681.

Niu G, Sun Y. 2019. Salt tolerance in roses: a review. *Acta Horticulturae* 1232: 143–150.

Noack R. 2003. Breeding: Selection Strategies for Disease and Pest Resistance. In: Roberts AV, Debener T, Gudin S (eds.). *Encyclopedia of Rose Science*. Oxford: Academic Press, 49–55.

Noh YM, Ait Hida A, Raymond O, Comte G, Bendahmane M. 2024. The scent of roses, a bouquet of fragrance diversity. *Journal of Experimental Botany* 75(5): 1252–1264.

Nybom H. 2009. Introduction to *Rosa*. In: Folta KM, Gardiner SE (eds.). *Genetics and Genomics of Rosaceae*. New York: Springer, 339–351.

Ohba H. 1993. *Rosa* L. In: Iwatsuki K, Boufford DE, Ohba H (eds.). *Flora of Japan*, Vol. 3. Tokyo: Kodansha, 169–177.

Ohno S. 1970. *Evolution by Gene Duplication*. Berlin: Springer.

Pan X, Zheng Y, Lei K, Tao W, Zhou N. 2024. Systematic analysis of Heat Shock Protein 70 (*HSP70*) gene family in radish and potential roles in stress tolerance. *BMC Plant Biology* 24(1): 2.

Park CJ, Shin R. 2022. Calcium channels and transporters: roles in response to biotic and

abiotic stresses. *Frontiers in Plant Science* 13: 964059.

Park HY, Seok HY, Park BK, et al. 2008. Overexpression of Arabidopsis *ZEP* enhances tolerance to osmotic stress. *Biochemical and Biophysical Research Communications* 375(1): 80–85.

Paul I, Poddar Sarkar M, Bhadoria PBS. 2022. Floral secondary metabolites in context of biotic and abiotic stress factors. *Chemoecology* 32(2): 49–68.

Peppe DJ, Hickey LJ, Miller IM, Green WA. 2008. A morphotype catalogue, floristic analysis and stratigraphic description of the Aspen Shale flora (Cretaceous–Albian) of southwestern Wyoming. *Bulletin of the Peabody Museum of Natural History* 49(2): 181–208.

Pichersky E, Gershenzon J. 2002. The formation and function of plant volatiles: perfumes for pollinator attraction and defense. *Current Opinion in Plant Biology* 5(3): 237–243.

Price AL, Jones NC, Pevzner PA. 2005. *De novo* identification of repeat families in large genomes. *Bioinformatics* 21(Suppl_1): i351–i358.

Qi S, Wang X, Wu Q, et al. 2023. Morphological, physiological and transcriptomic analyses reveal potential candidate genes responsible for salt stress in *Rosa rugosa*. *Ornamental Plant Research* 3(1): 21.

Ranallo-Benavidez TR, Jaron KS, Schatz MC. 2020. GenomeScope 2.0 and Smudgeplot for reference-free profiling of polyploid genomes. *Nature Communications* 11(1): 1432.

Raymond O, Gouzy J, Just J, et al. 2018. The *Rosa* genome provides new insights into the domestication of modern roses. *Nature Genetics* 50(6): 772–777.

Rehder A. 1940. *Manual of Cultivated Trees and Shrubs Hardy in North America: exclusive of the subtropical and warmer temperate regions*, 2nd edn. New York: Macmillan. Pp.

426–451.

Ren C, Zhu X, Zhang P, Gong Q. 2016. Arabidopsis COP1-interacting protein 1 is a positive regulator of ABA response. *Biochemical and Biophysical Research Communications* 477(4): 847–853.

Ren H, Yang W, Jing W, et al. 2024. Multi-omics analysis reveals key regulatory defense pathways and genes involved in salt tolerance of rose plants. *Horticulture Research* 11(5): uhae068.

Roach MJ, Schmidt SA, Borneman AR. 2018. Purge Haplotigs: allelic contig reassignment for third-gen diploid genome assemblies. *BMC Bioinformatics* 19: 1–10.

Rock CD, Sakata Y, Quatrano RS. 2010. Stress signaling □: the role of abscisic acid (ABA). In: Pareek A, Sopory SK, Bohnert HJ, Govindjee (eds.). *Abiotic Stress Adaptation in Plants: physiological, molecular and genomic foundation*. Dordrecht: Springer, 33–73.

Rubio F, Gassmann W, Schroeder JI. 1995. Sodium-driven potassium uptake by the plant potassium transporter HKT1 and mutations conferring salt tolerance. *Science* 270(5242): 1660–1663.

Ruhsam M, Hollingsworth PM, Darwin Tree of Life Consortium. 2024. The genome sequence of the Japanese rose, *Rosa rugosa* Thunb., 1784 (Rosaceae). *Wellcome Open Research* 9: 513.

Rus A, Yokoi S, Sharkhuu A, et al. 2001. AtHKT1 is a salt tolerance determinant that controls Na$^+$ entry into plant roots. *Proceedings of the National Academy of Sciences USA* 98(24): 14150–14155.

Sahoo RK, Ansari MW, Tuteja R, Tuteja N. 2014. *OsSUV3* transgenic rice maintains higher

endogenous levels of plant hormones that mitigates adverse effects of salinity and sustains crop productivity. *Rice* 7: 1–3.

Sanders D, Pelloux J, Brownlee C, Harper JF. 2002. Calcium at the crossroads of signaling. *The Plant Cell* 14(suppl_1): S401–S417.

Shang J, Feng D, Liu H, et al. 2024. Evolution of the biosynthetic pathways of terpene scent compounds in roses. *Current Biology* 34(15): 3550–3563.

Shulaev V, Korban SS, Sosinski B, et al. 2008. Multiple models for Rosaceae genomics. *Plant Physiology* 147(3): 985–1003.

Shulaev V, Sargent DJ, Crowhurst RN, et al. 2011. The genome of woodland strawberry (*Fragaria vesca*). *Nature Genetics* 43(2): 109–116.

Slater GS, Birney E. 2005. Automated generation of heuristics for biological sequence comparison. *BMC Bioinformatics* 6(1): 1–11.

Smit AFA, Hubley R. 2008–2015. RepeatModeler Open-1.0. Available at https://www.repeatmasker.org/RepeatModeler/

Smit AFA, Hubley R, Green P. 2013–2015. RepeatMasker Open-4.0. Available at https://www.repeatmasker.org/RepeatMasker/

Song J, Yan J, Sun B, et al. 2024. Abscisic acid regulates Cl⁻ efflux via the ABI5-ZAT10-SLAH3 module in chloride-stressed *Malus hupehensis*. *Horticulture Research* 11(9): uhae200.

Stanke M, Waack S. 2003. Gene prediction with a hidden Markov model and a new intron submodel. *Bioinformatics* 19(Suppl_2): ii215–ii225.

Sun Y, Zhao X, Gao Y, et al. 2022. Genome-wide analysis of lectin receptor-like kinases

(LecRLKs) in sweet cherry (*Prunus avium*) and reveals PaLectinL16 enhances sweet cherry resistance with salt stress. *Environmental and Experimental Botany* 194: 104751.

Sunkar R, Kaplan B, Bouche N, et al. 2000. Expression of a truncated tobacco *NtCBP4* channel in transgenic plants and disruption of the homologous *Arabidopsis CNGC1* gene confer Pb^{2+} tolerance. *The Plant Journal* 24(4): 533–542.

Tholl D. 2015. Biosynthesis and biological functions of terpenoids in plants. In: Schrader J, Bohlmann J (eds.). *Biotechnology of Isoprenoids*. Cham: Springer, 63–106.

Tian X, Wang Z, Zhang Q, et al. 2018. Genome-wide transcriptome analysis of the salt stress tolerance mechanism in *Rosa chinensis*. *PloS One* 13(7): e0200938.

Tian Y, Zeng H, Wu JC, et al. 2024. The zinc finger protein DHHC09 S-acylates the kinase STRK1 to regulate H_2O_2 homeostasis and promote salt tolerance in rice. *The Plant Cell* 36(4): 919–940.

Tilman D, Cassman KG, Matson PA, Naylor R, Polasky S. 2002. Agricultural sustainability and intensive production practices. *Nature* 418(6898): 671–677.

Tuskan GA, Difazio S, Jansson S, et al. 2006. The genome of black cottonwood, *Populus trichocarpa* (Torr. & Gray). *Science* 313(5793): 1596–1604.

Tuteja N, Sahoo RK, Garg B, Tuteja R. 2013. OsSUV3 dual helicase functions in salinity stress tolerance by maintaining photosynthesis and antioxidant machinery in rice (*Oryza sativa* L. cv. IR64). *The Plant Journal* 76(1): 115–127.

Unsicker SB, Kunert G, Gershenzon J. 2009. Protective perfumes: the role of vegetative volatiles in plant defense against herbivores. *Current Opinion in Plant Biology* 12(4): 479–485.

Urquhart W, Chin K, Ung H, Moeder W, Yoshioka K. 2011. The cyclic nucleotide-gated

channels AtCNGC11 and 12 are involved in multiple Ca^{2+}-dependent physiological responses and act in a synergistic manner. *Journal of Experimental Botany* 62(10): 3671–3682.

Van Fleet W. 1908. Breeding wild roses. *Journal of Heredity* os-4(1): 240–244.

Van Zelm E, Zhang Y, Testerink C. 2020. Salt tolerance mechanisms of plants. *Annual Review of Plant Biology* 71: 403–433.

VanderSluis B, Bellay J, Musso G, et al. 2010. Genetic interactions reveal the evolutionary trajectories of duplicate genes. *Molecular Systems Biology* 6(1): 429.

Vu NT, Nguyen NBT, Ha HH, et al. 2023. Evolutionary analysis and expression profiling of the *HSP70* gene family in response to abiotic stresses in tomato (*Solanum lycopersicum*). *Science Progress* 106(1): 00368504221148843.

Walker BJ, Abeel T, Shea T, et al. 2014. Pilon: an integrated tool for comprehensive microbial variant detection and genome assembly improvement. *PloS One* 9(11): e112963.

Wang Y, Tang H, DeBarry JD, et al. 2012. MCScanX: a toolkit for detection and evolutionary analysis of gene synteny and collinearity. *Nucleic Acids Research* 40(7): e49.

Ward LF, Jenney WP. 1899. *The Cretaceous Formation of the Black Hills as Indicated by the Fossil Plants*. Washington: US Government Printing Office.

Wissemann V. 2003. Classification: Conventional Taxonomy (Wild Roses). In: Roberts AV, Debener T, Gudin S (eds.). *Encyclopedia of Rose Science*. Oxford: Academic Press, 111–117.

Wissemann V, Ritz CM. 2005. The genus *Rosa* (Rosoideae, Rosaceae) revisited: molecular analysis of nrITS-1 and *atpB–rbcL* intergenic spacer (IGS) versus conventional taxonomy. *Botanical Journal of the Linnean Society* 147(3): 275–290.

Wu HJ, Zhang Z, Wang JY, et al. 2012. Insights into salt tolerance from the genome of *Thellungiella salsuginea*. *Proceedings of the National Academy of Sciences USA* 109(30): 12219–12224.

Wu T, Hu E, Xu S, et al. 2021. clusterProfiler 4.0: A universal enrichment tool for interpreting omics data. *The Innovation* 2(3): 100141.

Xiang G, Zhang H, Jian H, et al. 2019. De Novo assembly and characterization of the transcriptome of susceptible and resistant rose species in response to powdery mildew. *Scientia Horticulturae* 257: 108653.

Xiong L, Lee H, Ishitani M, Zhu JK. 2002. Regulation of osmotic stress-responsive gene expression by the *LOS6/ABA1* locus in *Arabidopsis*. *Journal of Biological Chemistry* 277(10): 8588–8596.

Yamaguchi T, Aharon GS, Sottosanto JB, Blumwald E. 2005. Vacuolar Na^+/H^+ antiporter cation selectivity is regulated by calmodulin from within the vacuole in a Ca^{2+}-and pH-dependent manner. *Proceedings of the National Academy of Sciences USA* 102(44): 16107–16112.

Yang J, Zhang J, Yan H, et al. 2024. The chromosome-level genome and functional database accelerate research about biosynthesis of secondary metabolites in *Rosa roxburghii*. *BMC Plant Biology* 24(1): 410.

Ye Y, Ding Y, Jiang Q, Wang F, Sun J, Zhu C. 2017. The role of receptor-like protein kinases (RLKs) in abiotic stress response in plants. *Plant Cell Reports* 36: 235–242.

Yokoya K, Roberts AV, Mottley J, Lewis R, Brandham PE. 2000. Nuclear DNA amounts in roses. *Annals of Botany* 85(4): 557–561.

Young MA, Schorr P, Baer R. 2007. *Modern Roses 12: the comprehensive list of roses in*

cultivation or of historical or botanical importance. Shreveport: The American Rose Society.

Yu H, Yan J, Du X, Hua J. 2018. Overlapping and differential roles of plasma membrane calcium ATPases in Arabidopsis growth and environmental responses. *Journal of Experimental Botany* 69(10): 2693–2703.

Yu TT, Ku TC. 1985. *Rosa* L. In: Yu TT (ed.). *Flora Reipublicae Popularis Sinicae*, Vol. 37. Beijing: Science Press, 360–455.

Zang F, Ma Y, Tu X, et al. 2021. A high-quality chromosome-level genome of wild *Rosa rugosa*. *DNA Research* 28(5): dsab017.

Zhang X, Wu Q, Lan L, et al. 2024. Haplotype-resolved genome assembly of the diploid *Rosa chinensis* provides insight into the mechanisms underlying key ornamental traits. *Molecular Horticulture* 4(1): 14.

Zhang X, Zhang J, Zhang W, Yang T, Xiong Y, Che D. 2016. Transcriptome sequencing and de novo analysis of *Rosa multiflora* under cold stress. *Acta Physiologiae Plantarum* 38: 1–13.

Zhang Y, Wang LF, Li TT, Liu WC. 2021. Mutual promotion of *LAP2* and *CAT2* synergistically regulates plant salt and osmotic stress tolerance. *Frontiers in Plant Science* 12: 672672.

Zhang Z. 2022. KaKs_Calculator 3.0: calculating selective pressure on coding and non-coding sequences. *Genomics, Proteomics and Bioinformatics* 20(3): 536–540.

Zhang Z, Xiao J, Wu J, et al. 2012. ParaAT: a parallel tool for constructing multiple protein-coding DNA alignments. *Biochemical and Biophysical Research Communications* 419(4): 779–781.

Zhao C, Jiang W, Zayed O, et al. 2021. The LRXs-RALFs-FER module controls plant growth and salt stress responses by modulating multiple plant hormones. *National Science Review* 8(1): nwaa149.

Zhao C, Zayed O, Yu Z, et al. 2018. Leucine-rich repeat extensin proteins regulate plant salt tolerance in *Arabidopsis*. *Proceedings of the National Academy of Sciences USA* 115(51): 13123–13128.

Zhao Y, Chan Z, Gao J, et al. 2016. ABA receptor PYL9 promotes drought resistance and leaf senescence. *Proceedings of the National Academy of Sciences USA* 113(7): 1949–1954.

Zhong MC, Jiang XD, Yang GQ, et al. 2021. Rose without prickle: genomic insights linked to moisture adaptation. *National Science Review* 8(12): nwab092.

Zhou J, Ju P, Zhang F, et al. 2020. OsSRK1, an atypical S-receptor-like kinase positively regulates leaf width and salt tolerance in rice. *Rice Science* 27(2): 133–142.

Zhou YB, Liu C, Tang DY, et al. 2018. The receptor-like cytoplasmic kinase STRK1 phosphorylates and activates CatC, thereby regulating H_2O_2 homeostasis and improving salt tolerance in rice. *The Plant Cell* 30(5): 1100–1118.

Zhu JK. 2001. Plant salt tolerance. *Trends in Plant Science* 6(2): 66–71.

Zhuang D, Ma C, Xue L, et al. 2021. Transcriptome and de novo analysis of *Rosa xanthina* f. *spontanea* in response to cold stress. *BMC Plant Biology* 21: 1–11.

Zong D, Liu H, Gan P, et al. 2024. Chromosomal□scale genomes of two *Rosa* species provide insights into genome evolution and ascorbate accumulation. *The Plant Journal* 117(4): 1264–1280.

1   **Table 1**. Statistics of the genome assembly and annotation of *Rosa lucieae*.

Feature	Value
Genome assembly	
Estimated haploid size (bp)	458,297,472
Estimated repeat sequences (bp)	273,209,581
Estimated heterozygosity (%)	1.35
Assembled genome size (bp)	544,655,161
Number of chromosomes	7
Total size of chromosomes (bp)	520,416,292
Number of unplaced contigs	187
Total size of unplaced contigs (bp)	24,238,869
Contig N50 (bp)	180,204
GC content (%)	38.71
Gap length (bp)	116,200
Complete eudicots_odb10 BUSCOs (%)	96.17
Complete embryophyte_odb10 BUSCOs (%)	96.78
Complete viridiplantae_odb10 BUSCOs (%)	97.17
Genome annotation	
Repeat sequences (bp)	270,481,675
Number of protein-coding genes	36,876
Average gene length (bp)	3651.95
Average CDS length (bp)	1263.19
Average number of exons per gene	5.16
Average exon length (bp)	244.86
Average intron length (bp)	574.38
Gene content (%)	24.12
CDS content (%)	8.55

Table 2. List of candidates of salt tolerance-related genes duplicated in *Rosa lucieae*.

Genes	Product	Function	Reference species	Reference
Stress Response & Signaling				
BON	Protein BONZAI	Ca^{2+} signaling & ABA accumulation	*Arabidopsis thaliana*	Chen et al. (2020)
CIP1	COP1-interactive protein 1	ABA signaling	*Arabidopsis thaliana*	Ren et al. (2016)
CBL1	Calcineurin B-like protein 1	Ca^{2+} sensor	*Arabidopsis thaliana*	Cheong et al. (2003)
CNGC1	Cyclic nucleotide-gated ion channel 1	Ca^{2+} uptake & homeostasis	*Arabidopsis thaliana*	Ma et al. (2006); Urquhart et al. (2011)
CSC1	Calcium permeable stress-gated cation channel 1	Ca^{2+} signaling	*Arabidopsis thaliana*	Hou et al. (2014)
IP5P7	Type IV inositol polyphosphate 5-phosphatase 7	ROS signaling	*Arabidopsis thaliana*	Kaye et al. (2011)
LRK10L1.2	Leaf rust 10 disease-resistance locus receptor- like protein kinase-like 1.2	ABA signaling	*Arabidopsis thaliana*	Lim et al. (2015)
LRX4	Leucine-rich repeat extensin-like protein 4	hormonal homeostasis & ROS accumulation	*Arabidopsis thaliana*	Zhao et al. (2018, 2021)
PYL9	Abscisic acid receptor PYL9	ABA signaling	*Arabidopsis thaliana*	Zhao et al. (2016)
RBOH	Respiratory burst oxidase homolog protein	ROS signaling	*Arabidopsis thaliana*	Kwak et al. (2003); Begum et al. (2024)
SD25	G-type lectin S-receptor-like serine/threonine-protein kinase SD2-5	ABA signaling	*Oryza sativa*	Zhou et al. (2020)
ZEP	Zeaxanthin epoxidase	ABA biosynthesis	*Arabidopsis thaliana*	Xiong et al. (2002); Park et al. (2008)
Homeostasis & Detoxification				
ACA13	Calcium-transporting ATPase 13	Ca^{2+} efflux	*Arabidopsis thaliana*	Maathuis et al. (2003); Yu et al. (2018)
CML18	Probable calcium-binding protein CML18	NHX1 binding	*Arabidopsis thaliana*	Yamaguchi et al. (2005)
HKT1	Sodium transporter HKT1	Na^+/K^+ homeostatis & Na^+ recirculation	*Arabidopsis thaliana*	Rubio et al. (1995); Rus et al. (2001); Berthomieu et al. (2003)
HSP70	Heat shock cognate 70 kDa protein	Chaperone	*Raphanus sativus*, *Solanum lycopersicum*	Vu et al. (2023); Pan et al. (2024)
INT4	Inositol transporter 4	Na^+/K^+ homeostatis	*Malus hupehensis*, *Mesembryanthemum crystallinum*	Hu et al. (2020); Li et al. (2021)
LAP2	Leucine aminopeptidase 2	ROS scavenging	*Arabidopsis thaliana*	Zhang et al. (2021)
LRK91	L-type lectin-domain containing receptor kinase IX.1	ROS scavenging	*Prunus avium*	Sun et al. (2022)
RSA1	SHORT ROOT IN SALT MEDIUM	ROS detoxification & Na^+ homeostasis	*Arabidopsis thaliana*	Guan et al. (2013)
SLAH3	S-type anion channel SLAH3	Cl^- efflux	*Malus hupehensis*	Song et al. (2024)
STRK1	Salt tolerance receptor-like cytoplasmic kinase 1	ROS scavenging	*Oryza sativa*	Zhou et al. (2018); Tian et al. (2024)
SUV3	DExH-box ATP-dependent RNA helicase DExH16	antioxidant & hormonal homeostasis	*Oryza sativa*	Tuteja et al. (2013); Sahoo et al. (2014)

2   **Fig. 1.** Genome map of *Rosa lucieae*. **A)** GC content, **B)** density of *Gypsy*, **C)** density of *Copia*,

3   **D)** density of protein-coding genes, and **E)** collinear relationships within the genome. The GC

4   content and densities of genetic elements were calculated using sliding windows of 500kb

5   without overlaps.

Fig. 2. Comparative genomic and evolutionary analyses of *Rosa lucieae*. **A)** Venn diagram of orthologous gene families among *Rosa lucieae*, *R. chinensis*, *R. rugosa* and *Fragaria vesca*. **B)** Gene family classification profile of protein-coding genes of *R. lucieae* and other close reference species. **C)** Chronogram and gene family evolution of *R. lucieae* and other close reference species. In the Bayesian phylogenetic tree, the posterior probability for every branch was 1.0.

2 **Fig. 3.** Gene ontology (GO) enrichment analysis on *Rosa lucieae* genes within the rapidly

3 evolving gene families based on the gene family evolution analysis. BP: biological process;

4 MF: molecular function; CC: cellular component; FDR: false discovery rate.

Fig. 4. Density plots of the synonymous substitution rates (Ks) of pairs of homologous genes within the four Rosoideae species. **A)** Pairwise comparisons of Ks distribution of paralogous and orthologous genes within pairs of the Rosoideae species. **B)** Pairwise comparisons of Ks distribution of paralogous and orthologous genes annotated with the gene ontology (GO) term 'response to salt stress' (GO:0009651) or its child terms (left), of which the ancestor chart is shown on the right side.

Fig. 5. Duplications and divergence of ACA genes within Rosoideae. **A)** Unrooted gene tree of ACA10, ACA12 and ACA13 genes (left) and the structural motifs in each ACA gene (right). **B)** Ideogram of the four Rosoideae species showing synteny of ACA and ECA genes between species. Colored lines represent the synteny relationships of orthologous ACA and ECA gene pairs, and gray lines represent the synteny relationships of genomic collinear blocks. The colors of lines represent the clusters within the ACA and ECA gene family (see subfigure C). The cladogram on the right side represents the species relationships. **C)** Copy number variation of ACA and ECA genes within the genomes of four Rosoideae species. The numbers in blocks and the color shades of blocks represent the gene copy numbers.