Non-diffusive slow heat dissipation induces high local temperature in living cells

Masaharu Takarada[1], Ryo Shirakashi[2], Masahiro Takinoue[3,4], Motohiko Ishida[1], Masamune Morita[3,5], Hiroyuki Noji[6], Kazuhito V. Tabata[4,6], Takashi Funatsu[1,7], and Kohki Okabe[1,4*]

[1]Graduate School of Pharmaceutical Sciences, The University of Tokyo, 7-3-1 Hongo, Bunkyo-ku, Tokyo 113-0033, Japan

[2]Institute of Industrial Science, the University of Tokyo, 4-6-1 Komaba Meguro-ku, Tokyo 153-8505, Japan

[3]Department of Computer Science, Institute of Science Tokyo, 4259 Nagatsuta-cho, Midori-ku, Yokohama 226-8502, Japan

[4]JST, PRESTO, 4-8-1 Honcho, Kawaguchi, Saitama 332-0012, Japan

[5]Biomedical Research Institute, National Institute of Advanced Industrial Science and Technology (AIST), Center 6, 1-1-1 Higashi, Tsukuba, Ibaraki, 305-8566, Japan

[6]Graduate School of Engineering, The University of Tokyo, 7-3-1 Hongo, Bunkyo-ku, Tokyo 113-8656, Japan

[7]Graduate School of Integrated Sciences for Life, Hiroshima University, 1-4-1 Kagamiyama, Higashi-Hiroshima 739-8528, Japan

*Correspondence should be addressed to: K.O. (okabe@mol.f.u-tokyo.ac.jp)

Abstract

Recently, intracellular thermometry has revealed temperature variations within cells. Although the biological significance of intracellular temperature change is recognized, the physical principles of intracellular temperature change remain a mystery. Here, we investigate intracellular heat transfer through intracellular temperature mapping using a fluorescent polymeric thermometer and high-speed fluorescence lifetime imaging microscopy. Through infrared laser irradiation-assisted heating, we track changes in temperature distribution to examine the mechanism of intracellular heat dissipation in comparison with heat conduction. Continuous heating induces the significantly slower relaxation of the average temperature of single cells compared with that of liposomes containing homogeneous aqueous solutions of comparable size; to the scale of seconds. We additionally elucidate that these phenomena are impacted by intracellular structures and molecules. Finally, we discover that this slow intracellular temperature relaxation originates from non-diffusive heat dissipation distinct from the conventional heat conduction model. Our results provide insights into the mechanisms of temperature variation in cells that are unresolved based on our current understanding, establishing a framework for understanding intracellular thermodynamics under non-equilibrium conditions.

Introduction

Temperature has a great influence on biological activity at various hierarchical levels. It is well established that physiological functions include adapting to and utilizing changes in ambient temperature. The study of the control of biological functions through heating has a long history, under topics such as hyperthermia therapy in medicine and

the heat shock response in biology. In recent years, intracellular temperature has attracted increasing attention. Fluorescent molecular thermometers (fluorescent polymeric thermometers [FPT][1,2], fluorescent proteins[3-5], fluorescent small molecules[6], and nanoparticles such as upconversion nanoparticles [UCNP][7,8] and fluorescent nanodiamonds [FND][9-11]) and non-optical methods (thermocouples[12,13] and electrochemical methods[14]) have revealed temporal and spatial variations of temperature within cells. These intracellular temperature changes are characterized by spontaneous heat generation within cells and are associated with cellular functions[15-18].

Very recently, it has been reported that heat endogenously generated within cells in response to physiological stimuli drives cellular functions such as neuronal differentiation[9], brain edema[19], and heat shock response[20]. In these phenomena, the driving force of the functions is not a change in external environmental temperature, but a temperature variation of about 1−2 °C induced by spontaneous intracellular heat. This emerging phenomenon produced by intracellular heat propagation (termed thermal signaling) is considered a potential universal intracellular signaling mechanism as all molecular dynamics and chemical reactions are governed by heat. Thus, intrinsic temperature changes in cells are a promising unexplored property in biology and medicine that drives cellular activity and determines the cellular state[15-18].

Although the biological significance of intracellular temperature variations is being revealed, the mechanism by which heat generated within the cell produces this temperature change of approximately 1–2 °C remains unknown. It has been pointed out that, according to the heat conduction equation, temperature changes due to spontaneous heat generation within cells can only reach approximately 10^{-5} °C (the 10^5 gap issue)[21]. While the values calculated from the heat conduction equation and experimental observations show a significant discrepancy, the observed result of intracellular temperature change by about 1 °C due to spontaneous heat generation has been verified through robust reproducibility, across different methodological principles[15]. Via attempts to explain the physical mechanism behind this discrepancy, it has been proposed that heat generated within intracellular regions where heat conduction is restricted could lead to elevated temperatures[5,22,23]. Recently, various studies have estimated cellular thermal conductivity (0.1–0.6 W m^{-1} K^{-1})[23-26] and thermal diffusivity (0.27–1.34×10^{-7} m^2 s^{-1})[5,27] to be between one-sixth and one-fold of those in water. It should be noted that the actual thermal conductivity and diffusivity reported in these studies showed significant variation. Even assuming conditions in which heat conduction is severely restricted (thermal conductivity: 0.1 W m^{-1} K^{-1}, thermal diffusivity: 0.27×10^{-7} m^2 s^{-1}), the discrepancy between the experimental results and the calculated values remains substantial (over 10^4-fold). This gap cannot be fully explained by considering additional parameters, such as the intensity of and the distance from the heat source. Thus, this issue concerning intracellular temperature variation is not a matter of the cellular thermal properties, but rather a fundamental mystery of the heat dissipation process itself[18]. This unresolved paradox of intracellular thermodynamics hinders the physical and biochemical understanding of the mechanisms of all chemical reactions and thermal signaling phenomena in cells. Despite the potential benefits of addressing this challenge, which could open new fields in cell biology and biophysics, no experimental investigation of the fate of intracellular heat has been performed to date.

In this study, in order to perform a detailed analysis of the dissipation process of heat generated in cells, we observed the heat transfer using a controllable heat source and a method for tracking the changes in the temperature distribution within the cells. First, we developed a real-time temperature mapping method using FPT by utilizing a

time-correlated single photon counting (TCSPC)-based FLIM instrument, which has a very high photon acquisition

capacity, in addition to employing a method for highly efficient fluorescence lifetime determination. Next, we used

the developed real-time temperature mapping method to track the transient temperature change caused by an artificial

heat source in living cells and liposomes, revealing that the temperature relaxation in cells is slow and, moreover,

that it is location- and molecule-dependent inside of the cells. The detailed analysis of the intracellular temperature

relaxation performed in comparison with the heat conduction equation revealed that the intracellular temperature

changes during continuous heating, similar to physiological thermogenesis, include temperature changes that do not

depend on heat conduction. These results suggested that thermal energy remains localized without immediate

dissipation, and the resultant local heat controls thermal signaling in cellular functions.

Results

High-speed temperature mapping in living cells

In this study, to explore the process of the change in temperature distribution within cells we adopted FPT[2] (Figure

1a), which responds sensitively to temperature changes above the lower critical solution temperature (LCST).

Intracellularly introduced FPT exists in a dispersed state within cells and can capture the temperature distribution[2].

This ability is advantageous over other intracellular temperature measurement methods, such as fixed-point

temperature measurement using fluorescent nanoparticles or temperature tracking within specific organelles using

fluorescent small molecules and proteins. We previously demonstrated that FPT selectively detects intracellular

temperature changes due to its functional independence; FPT did not respond to physicochemical factors other than

temperature, such as pH, ionic strength, biomolecular concentration, and viscosity, when they were quantitatively

altered in solution[2]. In this study, we used a control copolymer (CP)[2] with a chemical composition almost identical

to that of FPT, but lacking temperature sensitivity, to confirm the temperature selectivity of FPT. In complex

environments like the intracellular space, various physicochemical changes can occur due to heating. Therefore,

rigorous verification using this CP is essential to achieving our objectives. The inclusion of a control probe that

enables the investigation of temperature selectivity is a unique feature of FPT that is not found in other

nanothermometers.

To achieve the spatio-temporal resolution of temperature measurements that allows tracking of intracellular heat,

we changed the method for calculating fluorescence lifetime in TCSPC. Conventional fluorescence lifetime

determination via exponential approximation requires a large number of photons, such that a 60-second signal

integration barely provides a snapshot of the temperature distribution with organelle-sized spatial resolution[2].

Because we expected linear approximation to reduce variability in fluorescence lifetime determination, we adopted

the fast fluorescence lifetime[28] defined by the average photon arrival time ("fast lifetime"), which requires fewer

photons than the conventional method (Figure 1b and Supplementary Note 1).

First, we compared the fast lifetime and conventional fluorescence lifetime of FPT (Figures 1a, 1c, and

Supplementary Figure 1). Figure 1c shows the relationship between the temperature and the fast lifetime of FPT in

the solution of COS7 cell extracts. The temperature response curve of FPT by fast lifetime was almost the same as

that of the conventional method[2]. This difference in the fluorescence lifetime determination method also did not affect

the temperature response curve using living cells (Supplementary Figure 1). We confirmed that CP does not exhibit

temperature dependence even in fast lifetime measurements (Figure 1d).

We investigated whether reliable fast lifetime measurements could be achieved using fewer photons than with

conventional methods. We experimentally evaluated the impact of different fluorescence lifetime estimation

algorithms on the trueness and precision of the resulting values. (Figures 1e, 1f, and Supplementary Figure 2). The

results showed that fast lifetime-based fluorescence lifetime determination outperformed conventional methods in

terms of trueness (Figure 1e) and precision (Figure 1f and Supplementary Figure 2). Furthermore, it was confirmed

that the errors and variability associated with FPT fast lifetime determination using a limited number of photons did

not significantly affect the measurement values in this study (Supplementary Note 1).

Here, we performed intracellular temperature mapping with fast lifetime of FPT in living COS7 cells using the

state-of-the-art FLIM having high photon acquisition capability (photon detection efficiency per unit time is 25 times

higher than that of our previous system[2]). Comparing the fluorescence lifetime images at different measurement times,

we found that the intracellular temperature mapping became more detailed in a time-dependent manner; the spatial

heterogeneity of the intracellular temperature could be observed even at a measurement time of only 0.66 s, while a

previous study using FPT required a measurement time of 60 s (Figure 1g)[2]. The highly efficient fluorescence lifetime

determination of this method enabled fluorescence lifetime imaging at high compartmentalization (e.g., 512×512

pixels). The results of high-resolution temperature mapping in a single COS7 cell (Figure 1h) clearly revealed a

heterogeneous temperature distribution not only at the organelle level, such as a high temperature in the nucleus, but

also in micro-spaces in the cytoplasm and nucleoplasm. When the heterogeneous temperature distribution visualized

in an intracellular micro-space was compared with that in a homogeneous aqueous solution of the same size, the

spatial fluctuation of the intracellular temperature was notably large (Figure 1i). Next, mapping of the temperature

distribution was performed in cells in which mitochondria, the main heat-generating organelles, were visualized

(Figure 1j). As in our previous study[2], the mitochondria were clearly shown to have a higher temperature ($\Delta T = 1.29$

°C) than the surrounding region in steady-state cells.

Local heating using IR laser in living cells

Next, to observe the change in intracellular temperature with this mapping method, we heated an intracellular local

area via infrared (IR) laser (1480 nm) irradiation as an artificial heat source. By directly stimulating the O–H vibration

of water molecules, an irradiating IR laser focused on a glass surface (Figure 2a) can heat up an aqueous solution in

a local area (about 1 μm in diameter) in a transient and power-dependent manner[29,30]. The near-infrared spectrum of

water showed absorption of light at the wavelength of the IR laser used in this study (1,480 nm), while no absorption

of this light by FPT itself was observed (Supplementary Figure 3). Considering that the fluorescence lifetime values

and the components of the fluorophore (7-[*N*,*N*-dimethylaminosulfonyl]-2,1,3-benzoxadiazol-4-yl, DBD) in FPT are

sensitive to the chemical environment (e.g., hydrophobicity)[31], we analyzed the details of the fluorescence decay

curve of FPT at the heating center in the nucleus and the cytoplasm. As shown in Supplementary Figure 4, the

concordance of the fluorescence decay curves (i.e., the fluorescence lifetimes and their compositions of the two

components) during IR laser heating and medium heating was confirmed. Furthermore, the analysis of the

fluorescence decay curve of CP confirmed that factors other than temperature (e.g., viscosity) induced by IR laser

irradiation do not directly affect the FPT response (Supplementary Figure 5). Figure 2b shows the results of fluorescence lifetime imaging in the cells under IR laser irradiation, demonstrating the concentric distribution of temperature around the point of IR laser irradiation. On the other hand, IR laser irradiation did not change the fluorescence lifetime distribution of CP (Figure 2c). These results clearly demonstrate that the spatial distribution of the fluorescence lifetime of FPT created by IR laser irradiation is solely due to heat, independent of the chemical environment. When the nucleus was irradiated with the IR laser, the temperature gradient formed in the nucleus was steeper than that in the cytoplasm, even though the amount of IR laser power applied to the nucleus and the cytoplasm was the same (Supplementary Figure 6). Next, the increase in intracellular averaged temperature upon IR laser irradiation was dependent on the laser intensity (Figure 2d). To investigate the reliability of intracellular temperature change recorded during this artificial heating, we examined the results using another fluorescent thermometer with a completely different temperature response principle. Rhodamine B has long been known for the high temperature dependence of its fluorescence quantum yield[32] and has been used to measure solution temperature in nanoscale space[33]. The fluorescence intensity-based measurement of temperature using fluorescent small molecules is not suitable for intracellular temperature mapping, observation of local temperature where local fluorophore concentration changes, or temperature measurement in systems where environmental factors may change. However, it can be used to track the average temperature of the entire cell because this study confirmed that the effects of IR laser irradiation on factors other than heat are negligible. Here, we quantitatively investigated the intracellular temperature change during IR laser irradiation using the temperature-dependent change in fluorescence intensity of intracellular Rhodamine B-labeled dextran (molecular weight 10,000) (Supplementary Figure 7). The results showed that the change in averaged temperature during IR laser heating was the same between the two methods, which confirmed the validity of the temperature change measurement. In addition, by repeating the temperature mapping upon transient intracellular heating with an IR laser, we examined the ability to track the intracellular temperature distribution using this method. As shown in Figure 2e, the increase in fluorescence lifetime upon IR laser irradiation returned to the pre-irradiation value when the irradiation was stopped, and the temperature gradient could be repeatedly formed. Lastly, we quantitatively verified the temporal and spatial resolution of this temperature measurement using an artificially temperature-manipulatable IR laser, and determined that they are 9 ms and at the diffraction limited level, respectively (Supplementary Figures 8, 9, and Supplementary Note 2). These results clearly demonstrate that the spatial distribution of the fluorescence lifetime of FPT created by IR laser irradiation is solely due to heat, and that the resulting changes in intracellular temperature distribution can be tracked.

Characteristics of intracellular temperature relaxation

By taking advantage of the high temporal resolution of this high-speed temperature mapping method (9 ms), we observed the time-course of temperature changes in cells during IR laser irradiation (Figure 3a). Using CP, we confirmed that FPT responds only to temperature changes induced by IR laser irradiation (Figure 3b). Here, the average temperature of a finite region containing the heat source was analyzed. This average temperature change represents the change in the thermal energy of this region, divided by the heat capacity. First, the changes in averaged temperature of whole cells during pulsed and continuous heating are shown in Figure 3c. The results of fast rise and fast relaxation of temperature change during pulsed heating confirm that our high-speed temperature mapping method

is useful for tracking fast intracellular temperature change of several tens of milliseconds. The temperature relaxation

during pulsed heating was consistent with the time resolution limit of the present temperature tracking method

(Supplementary Figure 8 and Supplementary Note 2). In the case of continuous heating, the temperature rapidly

increased followed by a gradual increase upon IR laser irradiation, and then gradually decreased once irradiation was

terminated (Figure 3c). Because the relaxation rate did not depend on the magnitude of the temperature rise

(Supplementary Figure 10), we normalized the temperature change and compared the curves of relaxation, finding

that the relaxation after stopping continuous heating was slower than that in the case of pulse heating. As shown in

Supplementary Note 3, the relaxation of the average whole-cell temperature after this continuous heating exhibited

two-component time constants on the order of seconds. Further investigation of the heating duration dependence of

the intracellular relaxation rate revealed that it slows down in a heating duration-dependent manner and gradually

saturates after heating for more than 5–10 seconds (Supplementary Figure 11). We also tracked the average

temperature of whole cells using Rhodamine B. As shown in Supplementary Figure 12, the slow intracellular

temperature relaxation after continuous heating shown by Rhodamine B was in excellent agreement with the results

obtained by FPT. In addition, we confirmed the slow intracellular temperature relaxation using Cy3 tagged by

Streptavidin (Supplementary Figure 13). These results show that this slow temperature relaxation upon continuous

heating was an intrinsic cellular property.

Tracking the temperature of local areas in cells can help us understand the principles that determine the organelle-

specific temperature variation within them. Thus, we conducted temperature tracking of micro-spaces of cells. By

following the changes in fluorescence lifetime in local regions of cells, we were able to track the changes in average

temperature of a finite region (a square area of 10 μm per side with vertical thickness of z-resolution in this imaging

method) with a time resolution of several tens of milliseconds, and we observed the continuous heating-dependent

slow temperature relaxation even in the local cytoplasm (Figure 3d). Therefore, we compared the temperature

relaxation in different locations in cells. We monitored the temperature in the nucleus when heated; as shown in

Figure 3e, the relaxation of averaged temperature in the nucleus was slower than that in the cytoplasm. This slow

nuclear temperature relaxation may be partly due to nuclear-specific transcription and RNA content, because

transcription inhibition accelerated nuclear temperature relaxation (Supplementary Figure 14). Furthermore, the rate

of temperature relaxation in small spaces can be measured in giant plasma membrane vesicles (GPMVs) of 10 μm

diameter directly isolated from living cells, which was significantly faster than a region of similar size in the

cytoplasm (Supplementary Figure 15).

Cell-intrinsic temperature relaxation is significantly slower than heat conduction

To compare the results of slow temperature relaxation obtained in cells with a volume of water of the same size,

we next attempted to track the temperature variations of liposome, a micrometer-sized artificial compartment with a

lipid bilayer (Supplementary Figure 16 and Supplementary Movie 1). The comparison of the relaxation of averaged

temperature after continuous heating of liposomes with that in an intracellular local space of the same size clearly

showed that the temperature relaxation observed in single living cells was slower than that in water (Figure 3f). Next,

through several control experiments (Supplementary Note 4 and Supplementary Figure 17), we confirmed the

reliability of the FPT response (its insensitivity to factors other than temperature, such as viscosity) and the validity

of IR laser irradiation regarding the aforementioned slow intracellular heat dissipation, ensuring that structural
changes, distribution changes, or reactions—including physiological functions—of intracellular molecules do not
induce artifacts. These results suggest that the slow temperature relaxation observed in cells is an intracellular
component-dependent phenomenon.

Here, the time of temperature relaxation observed in cells after continuous heating (Figures 3c−3f) was compared
with numerical calculations based on heat conduction. Numerical calculations based on the unsteady heat conduction
equation, which consists of only a simple thermal diffusion term, using the thermal conductivity and specific heat of
cells showed that the time required for the relaxation of averaged temperature in cells by heat conduction is on the
order of ms (Supplementary Figure 18 and Supplementary Note 5). This comparison with the numerical calculation
shows that the relaxation of averaged temperature upon continuous heating observed in cells (which occurs on the
order of seconds) is much slower than that by heat conduction.

Conduction-independent, non-diffusive heat dissipation in living cells
Owing to the non-negligible difference in the relaxation time, we examined whether the intracellular temperature
variation originates from heat conduction. Given the diffusive nature of heat, the temperature distribution
immediately before the cessation of heating is essential for understanding the mechanism of temperature relaxation.
For this purpose, we constructed a method to track the intracellular temperature distribution during heating and
relaxation. Because the temperature mapping with high temporal resolution severely limits the number of photons
per pixel, we employed repeated measurements to achieve temperature distribution tracking in single cells (Figure
4a, Supplementary Figure 19, and Supplementary Movie 2), by taking advantage of the long-time measurement
capability of this method (photobleaching-independence of fast lifetime) and the property of our thermal
measurements where the temperature returned to its initial state.

Using this method, we investigated whether the characteristic dependency of the relaxation of intracellular
averaged temperature on the heating durations shown in Figures 3c, 3d, and Supplementary Figure 11 are attributable
to heat conduction. According to the heat conduction equation where thermal properties remain constant and there
are no exothermic or endothermic sources, the rate at which the average temperature relaxes after heating stops is
determined by the temperature distribution in the initial condition (Supplementary Note 6). Therefore, we
investigated whether the average intracellular temperature relaxation rate, as observed in Figure 3, depends on the
initial temperature distribution. As shown in Figure 4b, the average temperature relaxation rate was directly compared
under the condition that the temperature distribution was kept constant immediately before heating was stopped (i.e.,
the initial condition), regardless of the heating time. This was achieved by adjusting the IR laser intensity. The results
showed that the average temperature relaxation rate of the whole cell was slower with longer heating times, despite
almost identical initial temperature distributions (Figure 4c and Supplementary Movies 3–5). These results suggest
that the rate of intracellular temperature relaxation is not determined by the initial temperature distribution; however,
it is influenced by the heating duration, indicating that the intracellular temperature relaxation rate cannot be
explained by heat conduction alone.

The results were then also verified with respect to the diffusive nature of heat transfer in heat conduction. If heat
dissipation only occurs via conduction, the heat quantity decreases due to conduction from the region's boundary;

therefore, for the same temperature distribution, the time constant for the average temperature relaxation rate depends

on the size of the region. By using the equation (S3 and Supplementary Note 5) of heat conduction, we first

numerically calculated and compared the average temperature relaxation rate of various areas (D = 3, 5, and 10 μm,

where D is diameter). The calculated relaxation time of the average temperature in the finite region containing the

heat source was found to increase with an increase in the region size (Figure 4d and Supplementary Figure 20). Next,

we investigated this in living cells, examining the intracellular relaxation rate of the area-averaged temperature for

observation areas of varying sizes centered on the heat source (L = 3, 5, and 10 μm, where L is length of one side of

the square area). In contrast to the numerical calculation based on heat conduction, the relaxation rate of the area-

averaged temperature observed in the cells exhibited negligible dependence on the area size (Figure 4e), exhibiting

the distinctive characteristic of non-diffusive dissipation. The clear difference in the size dependence of the

temperature relaxation rate between simulations based on the heat conduction equation and observations in living

COS7 cells (Figure 4f) indicates that intracellular heat dissipation does not depend solely on heat conduction.

Therefore, it is suggested that, in addition to heat conduction, exothermic and endothermic processes contribute to

the intracellular heat transfer pathway that governs in situ temperature change.

Direct observation of high local temperature remaining in living cells during heat dissipation

Finally, to investigate the cause of the slow temperature relaxation in cells, we attempted to directly observe the slow

relaxation of temperature gradient in the intracellular micro-space. Figures 5a and 5b show the tracking of the

temperature distribution of local areas when stopping IR laser irradiation. In both the cytoplasm (Figure 5a and

Supplementary Movie 6) and the nucleus (Figure 5b and Supplementary Movie 7), the temperature gradient was

maintained around the heating point for approximately 200 ms (cytoplasm) and over 300 ms (nucleus) without

immediate diffusion after the heat source disappeared. In addition to this, the temperature relaxation of the entire

field of view was observed on the order of seconds. The temperature relaxation in the localized regions in the nucleus

was slower than that in the cytoplasm (Figures 5a and 5b), which is consistent with the findings on the tracking of

cellular average temperature shown in Figure 3. Furthermore, the difference in temperature relaxation in these two

areas was clearly confirmed by the difference in the relaxation rate of the temperature gradient when heating was

performed on the nuclear membrane (Supplementary Figure 21). These results indicate that slow heat dissipation in

the intracellular micro-space contributes to the slow temperature relaxation in cells.

Discussion

In this study, we revealed that the measured average intracellular temperature slowly relaxes by using a high-speed

temperature mapping method in cells through FPT and ultrasensitive FLIM, in contrast to the rapid relaxation of the

average temperature in liposomes containing homogeneous aqueous solutions. Furthermore, we found that

temperature relaxation depends on the intracellular location and its related molecules. Based on the temperature

selectivity and fast response time of FPT, the results of control experiments using CP, the independence of the

measurement methodology, and the fast relaxation at specific heating durations (e.g., 100 ms), we concluded that this

significantly slower temperature relaxation measured inside cells compared to a homogeneous aqueous solution when

cells were continuously heated is an intrinsic property of cells, and is not due to undesirable side effects of FPT.

Additionally, by exploiting the high temporal and spatial resolution of temperature mapping, we observed transient changes in intracellular temperature distribution after stopping heating and discovered that slow and non-diffusive heat dissipation within cells does not depend on heat conduction. The slow intracellular temperature relaxation upon continuous heating was confirmed in the investigations in another mammalian cell line using HeLa cells (Supplementary Figure 22).

By observing in situ heat transfer locally inside cells, we revealed that continuous heating induced a slow temperature relaxation and this temperature relaxation process does not follow the heat conduction equation with only a diffusion term (Figure 4). Furthermore, based on the discussion in Supplementary Note 7 regarding the possibility that thermal resistance at multiphase interfaces within the cell slows heat dissipation, the slow thermal relaxation occurring on the order of seconds within the intracellular space cannot be explained by heat conduction alone, even when considering thermal resistance. This finding suggests that intracellular temperature changes are accompanied by the process of non-diffusive heat dissipation involving energy conversion in addition to heat conduction. Therefore, we reconsidered the energy dissipation process in cells during continuous heating. The average temperature relaxation rate in the intracellular region depends on intracellular organelles (such as the nucleus), RNA (Supplementary Figure 14), and structures and large molecules that do not enter the GPMV (Supplementary Figure 15). In addition, we observed that both heat dissipation from the inside to the outside of the cell and heat transfer in the intracellular space were slow (Figures 5a and 5b). From these results, we speculate that the thermal energy generated by the absorption of IR laser light by intracellular water and biomolecules is propagated through energy absorption associated with state changes of specific molecules (related to the above structures) in the vicinity, in addition to being dissipated by heat conduction. This phenomenon may be attributable to the longer and thus non-negligible molecular relaxation times (compared with the relaxation times caused by heat conduction) of high energy states of molecules (e.g., higher-order structures) induced by continuous heating. For instance, the intrinsic relaxation times of RNAs depend on their interactions[34]. When these relaxation times are longer than the relaxation time caused by heat conduction (on the order of ms, see Supplementary Note 5), we can assume that the molecular relaxation of RNAs is a kinetic rate-limiting step in heat dissipation. As a result, this molecular relaxation dominates the heat dissipation of both RNAs and molecules that can share energy with RNAs. The thermal excitation of these biopolymers and the subsequent energy conversion keeps them in a higher energy state, preventing them from diffusive heat dissipation. However, considering that measurements with different temperature probe molecules—which are expected to have distinct affinities to biomolecules, varying temperature-sensing chemistry, and different molecular relaxation times—recorded the same intracellular heat dissipation rates (Supplementary Figures 12 and 13) and intracellular temperature changes on the same order[1-16,18-20], it is unlikely that molecular relaxation of the probe molecules affects the measurement of intracellular temperature. As similar temperature measurements can be made regardless of the type of thermometer, the implication is that energy is shared among biopolymers, each of which possesses unique dynamics.

On the other hand, an equilibrium system also exists within the cell in which heat is dissipated by conduction. In fact, our pulsed heating experiments and the recently developed label-free temperature measurement by refractive index change showed a fast temperature relaxation and a thermal conductivity not much different from that of water[27]. These facts suggest that intracellular temperature change includes both the temperature defined by a group of

biomolecules under non-equilibrium conditions and the temperature governed by the heat conduction equation which relies solely on thermal diffusivity. Examples of such non-equilibrium relationships between systems with different mechanisms of heat transfer (i.e., different temperature-determining populations) can be found in plasmas (electrons and cations)[35] and metal crystals (free electrons and phonons)[36]. In these examples, as the rate of energy conversion between systems in non-equilibrium states is not determined by the thermal conductivity of each system, it is possible to define different temperatures for the two systems within a time shorter than the relaxation time of that energy conversion. In cells, because of the energy conversion of certain biomolecules through state changes and their subsequent relaxation, resulting in low heat transfer rates, it is possible to define different temperatures for the two systems within the time required for energy dissipation. The elucidation of the specific mechanisms of energy conversion and molecular relaxation of biomolecules that generate this hypothetical temperature difference between the two systems may lead to an accurate definition of intracellular temperature, encompassing non-equilibrium conditions—an area that requires further exploration in future thermodynamic investigations.

To understand heat transfer in biological samples of various sizes, such as cells, tissues, and organs, it is necessary to consider the spatial scale of the observations. The time required for the temperature variations within the entire system to disappear via heat conduction strongly depends on the size of the space. On the other hand, as biological macromolecules have unique conformational change timescales that depend on their structure[34], the relaxation times of thermally excited molecules are independent of the size of the space. Therefore, the time required for heat dissipation via heat conduction including the effects of interfacial thermal resistance in spaces of certain sizes (i.e., in large spaces) is non-negligible compared with the relaxation time of the molecules. The relaxation time of subcutaneous tissue temperature observed in mice using a luminescent nanothermometer showed little deviation from the heat conduction model[37]. The phenomenon revealed in this study, i.e., that the intracellular temperature change is not completely governed by heat conduction, is observed only in the single-cell space. This suggests that, especially in cells, it is inappropriate to assume that all thermal energy is dissipated via heat conduction when determining the thermal conductivity and other thermal properties from temperature changes, given the substantial effects of energy conversion and molecular relaxation of biomolecules. The cellular thermal conductivity recently estimated from the local temperature change of molecular thermometers[5,23,24] differed significantly from the cellular thermal conductivity determined from a measurement in bulk (cell suspension) at a relatively large spatial scale[25] and equilibrium thermal measurements[26,27]. The low thermal conductivities and the thermal diffusivity estimated in cells using the former methods should reflect the significant influence of molecular relaxation on temperature changes at the single-cell scale, as discovered in this study.

Until now, the physical mechanism causing the discrepancy (the 10^5 gap) between the temperature change inside cells and the temperature change calculated by the heat conduction equation of a homogeneous system was unknown. The discovery of the existence of a very slow heat dissipation process that is independent of conduction highlights a critical flaw in attempts to estimate the temperature change during spontaneous intracellular heat generation from thermal conductivity and the heat conduction equation. In particular, regarding the large discrepancy of the 10^5 gap issue, our findings challenge the basic assumption that intracellular temperature changes are caused solely by simple heat conduction. Slow intracellular non-diffusive heat dissipation discovered in this study (Figure 4), which cannot be described by the simple heat conduction equation, can explain the temperature change during intracellular heat

generation that lasts for more than 1 s. In fact, cells with slower heat dissipation had a greater temperature increase

during endogenous heat generation (i.e., mitochondrial stimulation) (Supplementary Figure 23). In addition, we

estimate an intracellular temperature increase of 1.1 °C when heat dissipates independent of conduction (i.e., the cell

can be assumed to be adiabatic homogeneous during the relaxation time) upon mitochondrial stimulation

(Supplementary Note 8). This temperature increase is in excellent agreement with the intracellular temperature

increase measured in both the present (Supplementary Figure 23) and previous studies[15,16]. This consistency

demonstrates the importance of considering the thermal excitation of intracellular biopolymers and their energy

dissipation processes, rather than relying solely on simple heat conduction equations when attempting to understand

the mechanisms of intracellular temperature change.

Furthermore, the temperature elevation of biomolecules (which can be sensed by molecular thermometers) induced

via slow intracellular heat dissipation has physiological significance. The slow heat transfer during continuous

artificial heating observed in this study is also expected to occur during relatively long-term (on the order of minutes)

physiological heating within cells, such as mitochondrial heating (Supplementary Figure 23). Our study has reported

that spontaneous heat generation, which was dependent on the activity of neurons, caused an intracellular temperature

rise ($\Delta T = 2.12$ °C), activating its own thermosensitive Transient Receptor Potential Vanilloid 4 (TRPV4)[38]. More

recently, a thermal signaling mechanism has been discovered in neuronal cells where spontaneous temperature

increase ($\Delta T = 0.9$ °C) in the nucleus has a positive effect on important developmental processes during neuronal

differentiation[9]. The slow intracellular non-diffusive heat dissipation could explain how heat generated at the single-

cell level contributes to these thermal signaling mechanisms, which are activated by the aforementioned high

temperatures. Thus, slow heat dissipation in cells may play a pivotal role in facilitating the temperature rises that

trigger various physiological cellular functions[9,18-20]. This new concept should help to elucidate the intracellular

signaling mechanisms involved in dynamic medical phenomena (e.g., epilepsy[38,39], cancer hyperthermia[40], and

inflammation[41]) and cell biological events (body temperature control by brown adipose tissue[15,16], embryogenesis[42,43],

neuronal activity[9,19,44], apoptosis[1], and liquid–liquid phase separation[45]) that are closely related to temperature

changes. Through these investigations, the concepts discovered in this study should provide a revolutionary

perspective on life science research that has long been solely based on molecular communication.

Methods

Cell culture

COS7 cells (No. RCB0539, RIKEN BRC, Tsukuba, Japan) and HeLa cells (No. RCB0007, RIKEN BRC) were

cultured in Dulbecco's Modified Eagle's Medium (DMEM) with 10% fetal bovine serum supplemented with

penicillin–streptomycin, L-glutamine, sodium pyruvate, and nonessential amino acids at 37 °C in 5% CO_2. For live-

cell imaging, cells were cultured in 35-mm glass-bottomed dishes (AGC Techno Glass, Shizuoka, Japan) and the

medium was replaced by phenol red-free culture medium containing HEPES buffer (2 mL) before live-cell imaging.

All solutions were from Thermo Fisher Scientific (MA, USA).

Microinjection into living cells and culturing on the microscope

Microinjection of FPT, CP, Rhodamine B-labeled dextran (molecular weight 10,000; Thermo Fisher Scientific), and

Cy3 tagged Streptavidin (FUJIFILM Wako Pure Chemical Corporation, Osaka, Japan) into living cells was

performed with Femtojet (Eppendorf, Hamburg, Germany), controlled by a micromanipulator (Eppendorf). FPT (0.5

w/v%), CP (0.5 w/v%), Rhodamine B-labeled dextran (20 µM), and Cy3 tagged Streptavidin (3 µM) were dissolved

in an aqueous solution containing 140 mM KCl, 10 mM KH_2PO_4–K_2HPO_4, and 4 mM NaCl. The solution was filtered

using an Ultrafree-MC filter (Merck Millipore, MA, USA) and microinjected into the cytoplasm with a glass capillary

needle (Femtotips II; Eppendorf). The volume of injected solution was estimated to be 2 fL.

The temperature of the culture medium was maintained at 26 °C for monitoring temperature change and at 24 °C

for imaging temperature distribution. We chose these temperatures because they are the temperatures associated with

the highest sensitivity for measuring temperature with FPT (see Figure 1c) in each experiment. The temperature of

the culture medium was controlled and monitored with a stage-top incubator and a microscope objective lens heater

(INUBSF-ZILCS; Tokai Hit, Fujinomiya, Japan).

Preparation of cell extract

The COS7 cell pellets were collected and resuspended in a buffer (30 mM HEPES (pH 7.4), 75 mM Sucrose,

225 mM Mannitol, and 0.1 mM EGTA). After the addition of protease inhibitor cocktail (EDTA-free, Nacalai

Tesque, Inc, Kyoto, Japan), the cell suspend was lysed using Dounce Tissue Grinder (Wheaton, Millville, NJ),

followed by the three times of centrifugation (600 g, 10 min, 4 °C). Following further centrifugation (10,000 g,

10 min, 4 °C) of the supernatant, KCl was added to make the ionic strength 0.3 M. The concentration of proteins

in the obtained cell extract was quantified with BCA Protein Assay Kit (Thermo Fisher Scientific) after treated

with 1% Triton X-100 (Sigma-Aldrich, St. Louis, MO) to be 2.9 g L^{-1}.

Fluorescence lifetime imaging of living cells, cell extracts and GPMVs

Time-correlated single-photon counting (TCSPC)-based fluorescence lifetime imaging microscopy (FLIM) was

performed on a TCS SP8-FALCON confocal laser-scanning microscope (Leica Microsystems) with a pulsed diode

laser (PDL 800-B, 470 nm; PicoQuant, Berlin, Germany) at a repetition rate of 20 MHz, and the emission from 500

to 640 nm was obtained through the described objective with 1–18.2 zoom factors and a binning procedure (factor:

1 or 2) in a 64×64–1024×1024-pixel format for 0.7–257.5 s. To obtain the conventional fluorescence lifetime (τ_f),

the fluorescence decay curves of FPT in single cells and cell extract were fitted with a double exponential function:

$$I(t) = A_1 \exp(- t/\tau_1) + A_2 \exp(- t/\tau_2) \qquad (1)$$

Then, the conventional fluorescence lifetime (τ_f) was calculated using following equation:

$$\tau_f = (A_1 \tau_1^2 + A_2 \tau_2^2)/(A_1 \tau_1 + A_2 \tau_2) \qquad (2)$$

In determining the intracellular temperature change (ΔT) during heating by an artificial heat source, we used the

temperature response curve of τ_{FAST} of FPT (Supplementary Figure 1) for calibration when the entire cell was heated

to ignore the temperature increase due to the steady-state spontaneous heat generation of cells. For intracellular
temperature mapping, we used a calibration using cell extracts[2] (Figure 1c). The temperature response curves for the
temperature mapping with FPT in single cells and cell extract were obtained by approximating the relationship
between the averaged fast fluorescence lifetime (τ_{FAST}) of FPT (in triplicate) and the temperature using fourth- to
sixth-degree polynomials (coefficients of determination for both $R^2 = 0.99$):

$$T = A_1 \tau_{FAST}^6 + A_2 \tau_{FAST}^5 + A_3 \tau_{FAST}^4 + A_4 \tau_{FAST}^3 + A_5 \tau_{FAST}^2 + A_6 \tau_{FAST} + A_7 \qquad (3)$$

465 where T and τ_{FAST} represent the temperature (°C) and the fast fluorescence lifetime (ns), respectively. The coefficients
466 (A_1–A_7) in Equation (3) are listed in Supplementary Table 1.

Fluorescence imaging of the cells

Confocal fluorescence imaging was performed on a TCS SP8-FALCON confocal laser-scanning microscope (Leica
Microsystems, Wetzler, Germany) with a laser (638 nm) and an HC PL APO 63×/1.40 Oil CS2 objective (Leica
Microsystems). Mitochondria and the nucleus were labeled with MitoTracker Deep Red FM (Thermo Fisher
Scientific) and SiR-DNA (Cytoskeleton, Inc., CO, USA), respectively, and the fluorescence from 650 to 750 nm was
obtained.

Confocal fluorescence imaging of Rhodamine B and Cy3 were performed on a TCS SP8-FALCON confocal laser-
scanning microscope (Leica Microsystems) with a laser (552 nm and 488 nm), and the fluorescence from 560 to 700
nm and 500 to 700 nm were obtained, respectively.

Monitoring transient heating with IR laser irradiation

An IR laser (FRL-DC 1480 nm, maximum power, 3 W; IRE-Polus, MA, USA) was installed in a TCS SP8-FALCON
confocal laser-scanning microscope (Leica Microsystems) through the IR-LEGO system (Sigma Koki, Tokyo, Japan)
for the local heating of living cells and liposome, as shown in Figure 2a. An HC PL APO 63× 1.40 N.A. OIL CS2
objective (Leica Microsystems) was used to focus the IR laser and image the fluorescence lifetime of specimens; the
diameter of the focused area in the IR laser irradiation was estimated to be 1.29 μm. An electronically controlled
shutter (SSH-C4B; Sigma Koki) was used to start and stop the irradiation of the IR laser. The durations of irradiation
were 0.1–5 s. The timing of stopping heating was analytically determined by the rapid drop in FPT fluorescence
intensity. Considering that the heating operation using the shutter and the timing of the scanning in imaging are not
perfectly synchronized, the first frame after the fluorescence lifetime drop was excluded because it may not reflect
the temperature.

Absorption spectra of FPT

491 Absorption spectra of FPT (0.1 w/v% in KCl aqueous solution [150 mM]) were obtained using an IRT-5200 Infrared
492 Microscope (JASCO, Tokyo, Japan). A sample fixture was installed in a handmade cryostage for the microscope, and
493 the sample temperature was stably controlled at 25 °C using liquid nitrogen and an electric heater integrated into the
494 cryostage[46].

Tracking structural transition of FPT

The response time (τ) of the thermometers was determined by approximating with a single exponential function:

$$F(t) = a \exp(-t/\tau) + b \qquad (4)$$

where $F(t)$ and t represent the fluorescence intensity and time (s), respectively, and a and b are constants.

Fluorescence recovery after photobleaching (FRAP) analysis of FPT in living COS7 cells

The mobility of FPT in living cells was investigated by fluorescence recovery after photobleaching (FRAP). In the FRAP experiments, a small round area in living COS7 cells microinjected with FPT was photobleached for 2 s with 488 nm laser light focused on the area in the cytoplasm, followed by time-lapse confocal fluorescence imaging (described in Methods) over 200 s. In the image analysis, the fluorescence recovery was determined using the ratio of the averaged fluorescence intensity of the photobleached spot area to that of the whole cytoplasm calculated for each time point. The time course of fluorescence recovery for FRAP employing a uniform circular beam was fitted using the formula of Soumpasis et al.[47].

Inhibition of endogenous metabolic thermogenesis

Endogenous thermogenesis via glycolysis and oxidative phosphorylation was inhibited by adding 10 mM 2-deoxy-D-glucose (Sigma-Aldrich), 1 µM rotenone, and 1 µM antimycin A (Sigma-Aldrich) to the cell observation medium immediately before the experiment.

Preparation of giant plasma membrane vesicles (GPMVs)

COS7 cells injected with FPT were washed twice, then incubated with PFA (25 mM) and DTT (2 mM) dissolved in buffer (10 mM HEPES [pH 7.4], 150 mM NaCl, and 2 mM CaCl$_2$) for 3 hours at room temperature.

Preparation and fluorescence imaging of liposome

Briefly, 1,2-Dioleoyl-sn-glycero-3-phosphocholine (DOPC) was dissolved in chloroform/methanol mixtures (2/1, v/v). Then, 20 µL of DOPC solution (5 mM) was poured into a glass test tube and gently dried using a flow of air to produce a thin DOPC film at the bottom of the glass tube. This DOPC film was subsequently dried under a vacuum for 2 h. It was then hydrated with 190 µL of Rhodamine B-labeled dextran (10 µM) in a 200 mM sucrose solution at 50 °C overnight after 5 min of prehydration (50 °C) using 10 µL of the solution. The final DOPC concentration was 0.5 mM.

To prevent liposomes from moving out of the field of view upon IR laser irradiation due to convection, a microsized chamber array[48] of 20 µm in diameter and 50 µm in depth was introduced and a single liposome was placed in the well. First, a cover slip, on which microchambers were fabricated with SU-8, was treated with 200 µL of 200 mM glucose solution and deaeration, followed by the application of 2 µL of the solution containing liposomes and 18 µL of 200 mM glucose solution. Then, this cover slip with microchambers was sealed with a cover glass to allow

fluorescence lifetime imaging.

Confocal fluorescence imaging of liposomes was performed on a TCS SP8-FALCON confocal laser scanning

microscope (Leica Microsystems), as described above. The fluorescence signal observed inside the liposome using

confocal microscopy confirmed the inclusion of Rhodamine B-labeled dextran. Temperature changes were

determined from the temperature-dependent fluorescence intensity of Rhodamine B in aqueous solution

(Fluorescence intensity = $1.8294e^{-0.0229T}$, where T indicates the temperature; $R^2 = 0.99$).

Comparison of the time-course of temperature relaxation

The time-course of the temperature relaxation normalized by the temperature increase just before cessation of the

heating by IR laser irradiation was analyzed. The line of fitting with an exponential function is shown for reference.

The area under the curve (AUC) of normalized temperature relaxation for 3.5 s was also calculated, followed by

statistical analysis. The first two points were excluded in the calculation of AUC because the time required for the

structural relaxation of FPT does not reflect the temperature (Supplementary Figure 8).

Treatment with Actinomycin D

The chemical stimulus was conducted by adding 2 mg L^{-1} Actinomycin D (Nacalai Tesque, Inc., Kyoto, Japan) to the

culture medium 6 h prior to the observation.

Tracking of intracellular temperature distribution upon IR laser heating

Because the temperature mapping with high temporal resolution severely limits the number of photons per pixel, we

tracked the intracellular temperature distribution by repeating the measurements on the same cells (Supplementary

Figure 19). Fluorescence lifetime movies lasting 55 s (heating duration: 500 ms) and 105 s (heating duration: 1 and

5 s), including the duration of IR laser irradiation, were recorded five times in the same cells (Figures 4a and 4b) and

once in 10 cells (Figures 5a and 5b) or 20 cells (Supplementary Figure 22d), and averaged at the timing when IR

laser irradiation was stopped.

Measurement of temperature change during mitochondrial thermogenesis

The mitochondrial stimulus was conducted by quickly adding 1 mL of 200 µM FCCP (Sigma-Aldrich) in fetal bovine

serum (FBS)-free culture medium to the studied cells in 1 mL of FBS-free culture medium in a glass base dish. The

temperature of the culture medium was maintained at 26.0 °C.

Statistical analysis and reproducibility

Statistical significance was determined by Welch's two-sided t test. No statistical method was used to predetermine

the sample sizes. The experiments were not randomized. The investigators were not blinded to allocation during

experiments and outcome assessment. Cells displaying significant morphological alterations following

microinjection, laser heating, or treatment with inhibitors or cells in which the signal (fluorescence intensity or

lifetime) from the fluorescent probes did not return to preheating levels were excluded from the analysis. Data

indicating fluorescence lifetime values outside the range of the temperature response curve with FPT were excluded.

Data availability

All data that support the findings of this study are included in the paper. The raw data acquired by FLIM exceeds 200 GB and cannot be deposited in a public repository due to size limitations. All data are preserved by the authors and can be accessed by contacting the corresponding author. Requests will receive a response within 1 month. Source data are provided with this paper.

Code availability

The scripts used in this study are available online at:

https://github.com/tkrdmshr/Non-diffusive_slow_heat_dissipation_induces_high_local_temperature_in_living_cells.git. The scripts are also available at https://doi.org/10.5281/zenodo.18934454.

References

1. Gota, C., Okabe, K., Funatsu, T., Harada, Y., & Uchiyama, S. Hydrophilic fluorescent nanogel thermometer for intracellular thermometry. *J. Am. Chem. Soc.* **131**, 2766–2767 (2009).

2. Okabe, K. *et al.* Intracellular temperature mapping with a fluorescent polymeric thermometer and fluorescence lifetime imaging microscopy. *Nat. Commun.* **3**, 705 (2012).

3. Kiyonaka, S. *et al.* Genetically encoded fluorescent thermosensors visualize subcellular thermoregulation in living cells. *Nat. Methods* **10**, 1232–1238 (2013).

4. Nakano, M. *et al.* Genetically encoded ratiometric fluorescent thermometer with wide range and rapid response. *PLoS One* **12**, e0172344 (2017).

5. Lu, K. *et al.* Intracellular heat transfer and thermal property revealed by kilohertz temperature imaging with a genetically encoded nanothermometer. *Nano Letters* **22**, 5698–5707 (2022).

6. Kriszt, R. *et al.* Optical visualisation of thermogenesis in stimulated single-cell brown adipocytes. *Sci. Rep.* **7**, 1–14 (2017).

7. Di, X. *et al.* Quantitatively Monitoring In Situ Mitochondrial Thermal Dynamics by Upconversion Nanoparticles, *Nano Letters* **21**, 1651–1658 (2021).

8. Di, X. *et al.* Spatiotemporally mapping temperature dynamics of lysosomes and mitochondria using cascade organelle-targeting upconversion nanoparticles. *Proc. Natl. Acad. Sci. USA* **119**, e2207402119 (2022).

9. Chuma, S. *et al.* Implication of thermal signaling in neuronal differentiation revealed by manipulation and measurement of intracellular temperature. *Nat. Commun.* **15**, 3473 (2024).

10. Petrini, G. *et al.* Nanodiamond–quantum sensors reveal temperature variation associated to hippocampal neurons firing. *Adv. Sci.* **9**, 2202014. (2022).

11. Lee, Y., Kim, K., Kim, D., & Lee, J. S. Organelle-Specific Quantum Thermometry Using Fluorescent Nanodiamonds: Insights into Cellular Metabolic Thermodynamics. *J. Am. Chem. Soc.* **147**, 13180–13189 (2025).

12. Rajagopal, M. C. *et al.* Transient heat release during induced mitochondrial proton uncoupling. *Commun. Biol.* **2**, 1–6 (2019).

13. Wu, L. *et al.* Combined cellular thermometry reveals that salmonella typhimurium warms macrophages by
inducing a pyroptosis-like phenotype. *J. Am. Chem. Soc.* **144**, 19396–19409 (2022).

14. Ding, H., Liu, K., Zhao, X., Su, B., & Jiang, D. Thermoelectric nanofluidics probing thermal heterogeneity inside
single cells. *J. Am. Chem. Soc.* **145**, 22433–22441 (2023).

15. Zhou, J., Del Rosal, B., Jaque, D., Uchiyama, S. & Jin, D. Advances and challenges for fluorescence
nanothermometry. *Nat. Methods* **17**, 967–980 (2020).

16. Okabe, K., Sakaguchi, R., Shi, B., & Kiyonaka, S. Intracellular thermometry with fluorescent sensors for thermal
biology. *Pflügers Archiv Eur. J. Physiol.* **470**, 717–731 (2018).

17. Evanko, D. A thermometer for cells. *Nat. Methods* **9**, 328 (2012).

18. Okabe, K. & Uchiyama, S. Intracellular thermometry uncovers spontaneous thermogenesis and associated
thermal signaling, *Commun Biol* **4**, 1377 (2021).

19. Hoshi, Y. *et al.* Ischemic brain injury leads to brain edema via hyperthermia-induced TRPV4 activation. *J.*
Neurosci. **38**, 5700–5709 (2018).

20. Kang, M. G. *et al.* Mitochondrial Thermogenesis Can Trigger Heat Shock Response in the Nucleus. *ACS Central*
Science **10**, 1231–1241 (2024).

21. Baffou, G., Rigneault, H., Marguet, D., & Jullien, L. A critique of methods for temperature imaging in single
cells. *Nat. Methods* **11**, 899–901 (2014).

22. Suzuki, M. & Plakhotnik, T. The challenge of intracellular temperature. *Biophys. Rev.* **12**, 593–600 (2020).

23. Sotoma, S. *et al.* In situ measurements of intracellular thermal conductivity using heater-thermometer hybrid
diamond nanosensors. *Sci. Adv.* **7**, eabd7888 (2021).

24. Song, P. *et al.* Heat transfer and thermoregulation within single cells revealed by transient plasmonic imaging.
Chem **7**, 1569–1587 (2021).

25. Park, B. K. *et al.* Thermal conductivity of biological cells at cellular level and correlation with disease state. *J.*
Appl. Phys. **119** (2016).

26. Shrestha, R., Atluri, R., Simmons, D. P., Kim, D. S., & Choi, T. Y. Thermal conductivity of a Jurkat cell measured
by a transient laser point heating method. *Int. J. Heat Mass Transf.* **160**, 120161 (2020).

27. Toda, K. *et al.* Label-free mid-infrared photothermal microscopy revisits intracellular thermal dynamics: what do
fluorescent nanothermometers measure? *arXiv*:2406.16265.

28. Becker, W. Fluorescence lifetime imaging–techniques and applications. *J. Microscopy* **247**, 119–136 (2012).

29. Kamei, Y. *et al.* Infrared laser-mediated gene induction in targeted single cells in vivo. *Nat. Methods* **6**, 79–81
(2009).

30. Ebbinghaus, S., Dhar, A., McDonald, J. D., & Gruebele, M. Protein folding stability and dynamics imaged in a
living cell. *Nat. Methods* **7**, 319–323 (2010).

31. Gota, C., Uchiyama, S., Yoshihara, T., Tobita, S., & Ohwada, T. Temperature-dependent fluorescence lifetime of
a fluorescent polymeric thermometer, poly(*N*-isopropylacrylamide), labeled by polarity and hydrogen bonding
sensitive 4-sulfamoyl-7-aminobenzofurazan. *J. Phys. Chem. B* **112**, 2829–2836 (2008).

32. Kubin, R. F., & Fletcher, A. N. Fluorescence quantum yields of some rhodamine dyes. *J. Luminescence* **27**, 455–
462 (1982).

33. Moreau, D., Lefort, C., Burke, R., Leveque, P., & O'Connor, R. P. Rhodamine B as an optical thermometer in cells focally exposed to infrared laser light or nanosecond pulsed electric fields. *Biomed. Opt. Exp.* **6**, 4105–4117 (2015).

34. Bothe, J. R. *et al.* Characterizing RNA dynamics at atomic resolution using solution-state NMR spectroscopy. *Nat. Methods* **8**, 919–931 (2011).

35. Boulos, M. I., Fauchais, P., & Pfender, E. *Thermal plasmas* (Vol. 1). 5−8. Springer Science & Business Media. (1994).

36. Novko, D., Caruso, F., Draxl, C., & Cappelluti, E. Ultrafast Hot Phonon Dynamics in MgB_2 Driven by Anisotropic Electron-Phonon Coupling. *Phys. Rev. Lett.* **124**, 077001 (2020).

37. Ximendes, E. C. *et al.* Unveiling in vivo subcutaneous thermal dynamics by infrared luminescent nanothermometers. *Nano Lett.* **16**, 1695–1703 (2016).

38. Shibasaki, K. *et al.* Temperature elevation in epileptogenic foci exacerbates epileptic discharge through TRPV4 activation. *Lab. Invest.* **100**, 274–284 (2020).

39. Yang, X. F., Chang, J. H., & Rothman, S. M. Intracerebral temperature alterations associated with focal seizures. *Epilepsy Res.* **52**, 97–105 (2002).

40. Zhu, X., Li, J., Qiu, X., Liu, Y., Feng, W., & Li, F. Upconversion nanocomposite for programming combination cancer therapy by precise control of microscopic temperature. *Nat. Commun.* **9**, 2176 (2018).

41. Evans, S. S., Repasky, E. A., & Fisher, D. T. Fever and the thermal regulation of immunity: the immune system feels the heat. *Nat. Rev. Immunol.* **15**, 335–349 (2015).

42. Rodenfels, J., Neugebauer, K. M., & Howard, J. Heat oscillations driven by the embryonic cell cycle reveal the energetic costs of signaling. *Dev. Cell* **48**, 646–658 (2019).

43. Choi, J. *et al.* Probing and manipulating embryogenesis via nanoscale thermometry and temperature control. *Proc. Natl. Acad. Sci. USA* **117**, 14636–14641 (2020).

44. Ermakova, Y. G. *et al.* Thermogenetic neurostimulation with single-cell resolution. *Nat. Commun.* **8**, 15362 (2017).

45. Falahati, H. & Wieschaus, E. Independent active and thermodynamic processes govern the nucleolus assembly *in vivo*. *Proc. Natl. Acad. Sci. USA* **114**, 1335–1340 (2017).

46. Kawai, K., & Shirakashi, R. Water rotational relaxation time measurement by shortwave infrared micro spectroscopy (SWIR) at sub-zero temperatures. *Spectrochim. Acta A* **322**, 124707 (2024).

47. Soumpasis, D. M. Theoretical analysis of fluorescence photobleaching recovery experiments. *Biophys. J.* **41**, 95–97 (1983).

48. Watanabe, R. *et al.* Arrayed lipid bilayer chambers allow single-molecule analysis of membrane transporter activity. *Nat. Commun.* **5**, 4519 (2014).

Acknowledgment

We thank Professor Katsunori Hanamura, Professor Takeharu Nagai, Professor Takuro Ideguchi, and Dr. Keiichiro Toda for critical comments on this manuscript. We are also grateful for financial support from PRESTO and CREST

Author contributions

689 K.O. organized this work. K.O., M.Takarada, M.Takinoue, and T.F. designed this work. M.Takarada and K.O.
690 performed all imaging experiments and analyzed the data. M.Takarada., M.Takinoue, M.I., and M.M. prepared and
691 performed experiments using liposomes. K.V.T. and H.N. designed and fabricated the microarray chambers. R.S.
692 obtained absorption spectra, performed numerical calculations, and contributed to the interpretation of the results.
693 K.O. and M.Takarada wrote the paper.

Competing interest statement

696 The authors declare no competing interests.

Figure legends

Figure 1 | High-resolution mapping of temperature in living COS7 cell with τ_{FAST}-based FLIM.

a Functional diagram of FPT responding above the lower critical solution temperature (LCST) and its temperature
response curve using conventional fluorescence lifetime (τ_{f}). **b** Schematic diagram of fluorescence lifetime
determination in FLIM. In the conventional analysis, the fluorescence lifetime (τ_{f}) is calculated from the parameters
obtained by approximating the fluorescence decay curve with a two-component exponential function, while the fast
fluorescence lifetime (τ_{FAST}) is obtained by just averaging the arrival time (τ_{arr}) of photons in TCSPC. A and N
represent amplitude and number of photons, respectively. **c** The temperature response curve of FPT using fast
fluorescence lifetime (τ_{FAST}). **d** Relationship between temperature and the fast fluorescence lifetime (τ_{FAST}) of control
copolymer (CP). **e** Relationship between the accumulated photon count and fluorescence lifetime at 24, 27, and 30
°C (τ_{f}: left and τ_{FAST}: right) **f** The relationship between photon counts in TCSPC-FLIM and the standard deviation of
fluorescence lifetime (τ_{f}: black and τ_{FAST}: red). **g** Comparison of fluorescence lifetime images of FPT using τ_{FAST} with
various accumulation times. **h** Fluorescence lifetime (τ_{FAST}) images of FPT. The region of interest, as shown in the
square in the left panel, is enlarged in the right panel. **i** Comparison of temperature distribution in aqueous solution
and in the cytoplasm of a cell. Scale bar represents 2 μm. **j** The colocalization of the intracellular area with a higher
temperature than the surroundings (middle) with mitochondria visualized using MitoTracker Deep Red FM (left).
Histogram of fluorescence lifetime (τ_{FAST}) is shown on the right. Dotted line indicates the outline of the cell. ΔT was
calculated by subtracting the average temperature of the cytoplasm from that around mitochondria. $<\Delta T>$ represents
an average of the histogram. The experiments (**g**–**j**) were performed three times with similar results. Scale bars in **g**,
h, and **j** represent 10 μm. Source data are provided as a Source Data file.

Figure 2 | Local heating using IR laser in living COS7 cell.

a Optical set-up. **b,c** Fluorescence lifetime images of FPT (**b**) or control copolymer (CP, **c**) without (Off) and with

(On) IR laser irradiation (0.55 mW). The right graph indicates the profile of ΔT (**b**) or $\Delta\tau_{\mathrm{FAST}}$ (**c**) of the cell. **d** The

relationship between IR laser power and the increase in averaged temperature of whole cells (ΔT_{ave}). Data are

presented as mean ± s.e.m. ($n = 30$ cells). **e** Reversibility of the fluorescence lifetime response of FPT to local heating

(IR laser: 0.67 mW). The right graph shows the fluorescence lifetime of a 10 μm diameter area indicated by a dotted

circle in the left image. The experiments (**b,c,e**) were performed three times with similar results. Source data are

provided as a Source Data file.

Figure 3 | Monitoring temperature variation upon heating with IR laser in living COS7 cell and liposome.

a Schematic diagram of transient heating. Heating was applied by an IR laser for X s during 20 s of intracellular

temperature tracking. **b** Tracking of fast fluorescence lifetime changes of FPT (red) and control copolymer (CP, black)

in cells when heated by continuous (0.55 mW for 5 s) irradiation with an IR laser ($n = 29$ [FPT] and 19 [CP] cells).

c–f Tracking of the average temperature changes under IR laser heating and normalized temperature relaxation after

heating stops. The averaged temperature of whole cells (**c**) and square area with a 10 μm side centered on the heating

point (**d–f**) were analyzed. (**c,d**) The results of continuous heating (red, 0.55 mW, 5 s) are compared with those of

pulsed heating (black, 1.1 mW, 0.1 s). (**e**) The results of continuous heating (0.55 mW, 5 s) of the nucleus (red) are

compared with those of the cytoplasm (black). (**f**) The results of continuous heating (0.83 mW for 5 s) of the

cytoplasm (red) are compared with those of water within liposomes. Data including the inset in the right graph are

mean ± s.e.m. (Welch's two-sided *t* test: **c**, $p = 1.13 \times 10^{-19}$; $n = 29$ [continuous] and 15 [pulsed] cells; **d**, $p = 2.26 \times 10^{-}$

^{21}; $n = 30$ [continuous] and 15 [pulsed] cells; **e**, $p = 1.71 \times 10^{-7}$; $n = 30$ cells; **f**, $p = 1.07 \times 10^{-6}$; $n = 30$ cells and $n = 10$

liposomes). Insets in the right panels of **c–f** indicate areas under the curves (AUC) of normalized temperature

relaxation. The unit a.u. in AUC means arbitrary units. $^{***}P < 0.001$ (Welch's two-sided *t* test). Vertical scale bars in

the right panels of **c–f** represent indicated temperature changes. ΔT was calculated by subtracting the average

temperature from that of before heating. Source data are provided as a Source Data file.

Figure 4 | Independence of intracellular temperature relaxation after continuous heating from the heat conduction

model.

a The temperature mapping of a whole cell with local heating (0.55 mW for 1 s). The tracking of average temperature

change (left) and representative temperature distributions in cells at various time points. The cell is indicated by a

dotted line (right). Scale bar represents 10 μm. The experiment was performed four times with similar results. **b** The

intracellular temperature distribution with different heating conditions (top [0.89 mW for 500 ms], middle [0.83 mW

for 1 s], and bottom [0.55 mW for 5 s]). The line profiles of ΔT of cells and the temperature mapping during relaxation

are also shown. Scale bars represent 10 μm. The experiment was performed two (top), four (middle), and five

(bottom) times respectively with similar results. **c** The tracking of average temperature change of the whole cell,

normalized temperature relaxation, and its area under the curve (AUC, inset). The unit a.u. in AUC means arbitrary

units. Data in the inset are mean ± s.e.m. ($n = 10$ [500 ms], 9 [1 s], and 10 [5 s] cells). Vertical scale bars represent

temperature change. **d** The average temperature relaxation for various spatial sizes was analyzed from the time course

of the temperature distribution numerically calculated by the heat conduction equation (HCE) in Supplementary Note

5. Areas of diameter (D) = 3, 5, and 10 µm are shown. **e** The relaxation of average temperature of various sized

regions observed in the cytoplasm of living COS7 cells after continuous heating (5 s) (Figure 3**d**). Data in cells are

presented as mean ± s.e.m. ($n = 30$ cells). Areas of a square with a side (L) = 3, 5, and 10 µm are shown. **f** Comparison

of the size dependence of temperature relaxation times evaluated using fold change in area under the curve (AUC)

between the simulation based on the HCE and the observation in living COS7 cells. **a–c** ΔT was calculated by

subtracting the temperature from that of after the temperature relaxation. Source data are provided as a Source Data

file.

Figure 5 | Observation of slow temperature relaxation in intracellular local area.

High-speed mapping of intracellular temperature in the cytoplasm (**a**) and in the nucleus (**b**) immediately after

stopping local heating (1.1 mW [cytoplasm] and 0.83 mW [nucleus] for 5 s). Fluorescence lifetime images were taken

every 41 ms. The right graphs show the relaxation of average temperature in the heating areas indicated in the left

images. ΔT was calculated by subtracting the temperature from that of after the temperature relaxation. Scale bars

represent 2 µm. The experiments were performed three times with similar results. Source data are provided as a

Source Data file.

a

T < LSCT
Weakly fluorescent

T > LSCT
Strongly fluorescent

Heating / Cooling

| Thermo-sensitive sequence
⊖ Hydrophilic unit
☆,★ Fluorescent unit
Water molecule

b

Excitation
Photon (emission)

50 ns

τ_{arr} τ_{arr}

Arrival time

Time

Curve fitting

A_1, τ_1
A_2, τ_2

(Conventional)

Averaging

$\tau_f = \dfrac{A_1\tau_1^2 + A_2\tau_2^2}{A_1\tau_1 + A_2\tau_2}$

Accumulation

Averaging

$\tau_{FAST} = (1/N)\Sigma\tau_{arr}$

g Accumulation time

0.7 sec — Binning: 4
5 sec — Binning: 1
15 sec — Binning: 1
60 sec — Binning: 1

Fluorescence lifetime (τ_{FAST}, ns)
7.5 (29°C) — 4.9 (22°C)

h

Fluorescence lifetime (τ_{FAST}, ns)
6.6 (27°C) — 4.9 (22°C)

i Solution   Cytoplasm

Difference from average temperature (°C)
3.0 — -3.0

j Mitochondria   Temperature

Fluorescence lifetime (τ_{FAST}, ns)
7.0 (28°C) — 4.9 (22°C)

$\langle\Delta T\rangle$ = 1.29 °C

Cytoplasm
Mitochondria

a

Point of irradiation

IR Laser (1480 nm)

b

off   on

4.9 (22 ˚C)   8.9 (33 ˚C)
Fluorescence lifetime (τ_{FAST}, ns)

c

off   on

1.5   4.5
Fluorescence lifetime (τ_{FAST}, ns)

CP

d

e

1 2 3 4 5
6 7 8 9 10

5.4   9.5
(24 ˚C)   (35 ˚C)
Fluorescence lifetime (τ_{FAST}, ns)