Sec18 side-loading is essential for universal SNARE recycling across cellular contexts

Yousuf A. Khan[1,2,3,4,9], K. Ian White[1,2,3,4,5], Richard A. Pfuetzner[1,2,3,4,5], Bharti Singal[6], Luis

Esquivies[1,2,3,4,5], Garvey Mckenzie[7], Fang Liu[7], Katherine DeLong[1],

Uchoer B. Choi[1,2,3,4,5], Elizabeth Montabana[6], Theresa Mclaughlin[7], William T. Wickner[8],

Axel T. Brunger[1,2,3,4,5,9*]

[1]Department of Molecular and Cellular Physiology, Stanford University, Stanford, CA, USA
[2]Department of Neurology and Neurological Sciences, Stanford University, Stanford, CA, USA
[3]Department of Structural Biology, Stanford University, Stanford, CA, USA
[4]Department of Photon Science, Stanford University, Stanford, CA, USA
[5]Howard Hughes Medical Institute, Stanford University, Stanford, CA, USA
[6]Stanford Cryo-EM microscopy center, Stanford University, Palo Alto, CA, USA
[7]Stanford University Mass Spectrometry, Stanford University, Palo Alto, CA, USA
[8]Department of Biochemistry and Cell Biology, Geisel School of Medicine at Dartmouth,
Dartmouth College, Hanover, NH 03755.
[9]Correspondence: yousuf@stanford.edu (YAK), brunger@stanford.edu (ATB)

Summary

SNARE proteins drive membrane fusion as their core domains zipper into a parallel four-helix
bundle[1,2]. After fusion, these bundles are disassembled by the AAA+ protein Sec18/NSF and its
adaptor Sec17/ α-SNAP[3,4] to make them available for subsequent rounds of membrane fusion.
SNARE domains are often flanked by C-terminal transmembrane or N-terminal domains[5].
Previous structures of the NSF–α-SNAP–SNARE complex revealed SNARE domain threaded
through the D1 ATPase ring[6], posing a topological constraint as SNARE transmembrane
domains would prevent complete substrate threading as suggested for other AAA+ systems[7].
Here, *in vivo* mass-spectrometry reveals N-terminal SNARE domain interactions with Sec18,
exacerbating this topological issue. Cryo-EM structures of a yeast SNARE complex, Sec18, and
Sec17 in a non-hydrolyzing condition shows SNARE Sso1 threaded through the D1 and D2
ATPase rings of Sec18, with its folded, N-terminal Habc domain interacting with the D2 ring.
This domain does not unfold during Sec18/NSF activity. Cryo-EM structures under hydrolyzing
conditions revealed substrate-released and substrate-free states of Sec18 with a coordinated
opening in the side of the ATPase rings. Thus, Sec18/NSF operates by substrate side-loading and
unloading topologically constrained SNARE substrates.

Keywords:

ATPases associated with diverse cellular activities (AAA+);
N-ethylmaleimide-sensitive factor (NSF);
Sec18;
Soluble N-ethylmaleimide sensitive factor Attachment protein Receptor (SNARE);
Exocytosis;
Quality control;
cryo-electron microscopy;
single-molecule fluorescence resonance energy transfer;

Main

Cellular compartmentalization, growth, hormone secretion, transport, neurotransmission, and many other pathways depend on precise, rapid, and regulated membrane fusion[8,9]. Membrane fusion in eukaryotic cells is mediated by a highly conserved superfamily of SNAREs (Soluble N-ethylmaleimide sensitive factor Attachment protein Receptors)[5]. All SNAREs share a characteristic 60–70 amino acid SNARE domain often flanked by a C-terminal transmembrane domain, membrane anchors, and a folded N-terminal variable domain specific to the SNARE's function and intracellular pathway[10]. SNAREs on opposing membranes interact primarily through their SNARE domains to form a parallel *trans*-SNARE complex, juxtaposing two different membranes[11,12]. Membrane fusion commences when these SNARE domains zipper together in a directed fashion[1,2]. After fusion, the SNAREs form a highly stable parallel helical bundle, the so-called *cis*-SNARE complex[1]. This highly stable four-helix *cis*-SNARE complex is disassembled by Sec18 (N-ethylmaleimide-sensitive factor, NSF, in higher eukaryotes)[4,13,14] to provide the energy for subsequent membrane fusion events[15].

Sec18/NSF is a universally conserved AAA+ (ATPases associated with diverse cellular activities) protein translocase[16–18]. Sec18/NSF consists of an N-domain, an active D1 AAA+ domain, and a catalytically inactive D2 AAA+ oligomerization domain[19]. It was initially discovered as a critical complementation group required by the yeast secretory pathway, and it is now known for its role in recycling SNAREs[20] and SNARE assembly quality control for proper SNARE complex assembly[21–24]. Sec18/NSF, with the adaptor protein Sec17 (α-SNAP in higher eukaryotes), disassembles *cis*-SNARE complexes. The disassembly requires multiple ATP hydrolysis events in the presence of Mg^{2+} ; as few as 6 ATP hydrolysis events are sufficient[25,26].

Cryo-EM structures of the mammalian 20S complex (NSF, α-SNAP, and neuronal SNAREs ternary complex) in a non-hydrolyzing condition (*i.e.*, in the absence of divalent cations) revealed a supramolecular architecture in which the NSF N-domains bind α-SNAP molecules, three or four of which in turn bind the four-helix SNARE complex consisting of the SNARE domains of syntaxin-1A, SNAP-25, and synaptobrevin[3,13]. The N-terminal residues of the SNARE domain of SNAP-25 was bound to the D1 ring pore without apparent ATP hydrolysis, where it interacts with several conserved tyrosine amino acids in a spiral staircase-

like pattern[13,16,27]. In these EM maps, no ordered density was observed in the D2 ring pore,
consistent with the notion that it is catalytically inactive and primarily responsible for NSF
oligomerization rather than substrate engagement[28]. The interaction between the SNARE
substrate and the D1 pore is like that observed for other AAA+ translocases and suggests a
conserved mechanism for substrate threading through the D1 pore. However, the membrane
anchors and domains of the SNAREs would seemingly prevent complete threading of the type
suggested for other AAA+ systems[7,29,30], posing a topological challenge. Furthermore, SNAREs
contain globular N-terminal domain(s) of variable length and structure; for example, the N-
terminal domain of syntaxin consists of a three-helix bundle (Habc domain) involved in
regulating its function[31], and the N-terminal domains of Use1 and Sec20 in part form a stable
255 kDa tethering complex[32]. Complete threading would imply that such N-terminal domains are
somehow unfolded. These topological constraints surrounding SNARE loading, processing, and
release through Sec18/NSF are further compounded by the observation that Sec18/NSF
disassembles all SNAREs in all cellular contexts[5,10,33], which all contain a variety of different N-
and C-terminal domains and membrane arrangements/linkages.
89
***In vivo* cross-linking mass-spectrometry with Sec18**

These questions led us to investigate the space of Sec18/NSF—SNARE interactions
through *in vivo* protein crosslinking mass-spectrometry (XL-MS) in yeast. Due to its power as a
model system and relatively simple SNARE proteome, *S. cerevisiae* is an excellent model for
investigating Sec18 interactions with different substrates in live cells. As such, we developed a
protocol for *in vivo* crosslinking mass-spectrometry (XL-MS) in yeast to identify binding
partners of Sec18 (**Extended Data Figure 1A, Methods**).
In total, 35 identified proteins are shared between disuccinimidyl glutarate crosslinker
(DSG) treated and untreated conditions (*i.e.*, they are non-specific proteins) (**Figure 1A**). The
nine proteins observed only in the untreated condition had few unique spectra mapping to their
identified proteins, suggesting that they were not detected in the treated condition due to their
low abundance and difficulty of consistent detection. The 205 proteins unique to the 5 mM DSG
treated condition were processed using spatial analysis of functional enrichment (SAFE) analysis
to visualize the various cellular processes to which the identified proteins contribute[34]. At
increasing levels of significance thresholds ($p = 0.01$, 0.001, and 0.0001), the identified proteins
only enriched the vesicle trafficking processes. Gene ontology (GO) analysis also revealed
lesser-known processes, such as vacuolar acidification and ergosterol biosynthesis
(**Supplementary Figure 1**), consistent with Sec18's functions in these contexts[35–37]. The
enrichment of these processes validated our crosslinking protocol for specifically targeting Sec18
and its binding partners inside the cell.
Within our enriched protein dataset, we found many yeast SNAREs involved in different
pathways and compartments of the cell. We then mapped crosslinked residues of these SNAREs
to those of Sec18 (**Figure 1B**). Using a series of empirical constraints (**Methods**), we classified a
crosslink as high confidence if it met all constraints, medium confidence if it met some but not
all, and low confidence if it did not. Since crosslinking was performed in live cells, we expect

these crosslinks to represent interactions between SNAREs and Sec18 during substrate loading,

disassembly, and substrate release.

Previous cryo-EM structures of the complex of NSF, α-SNAP, and neuronal SNAREs in

a non-hydrolyzing condition revealed that the four-helix SNARE bundle interacts with between

two and four α-SNAP molecules, which in turn are bound by the N-domains of NSF; the N-

terminal end of one of the SNAREs is bound to the pore of the D1 ring[3,13,27]. Consistent with

these structures, we found crosslinks between SNARE domains and the N-domain or D1 domain

(examples in **Figure 1C, E, F**). Additionally, we observed several high-confidence crosslinks of

yeast SNARE proteins to the D2 domain, an unexpected result given an absence of interactions

between the SNAREs and the D2 ring in the cryo-EM structures and that the D2 domain is

catalytically inactive[19] and has had no previously reported role in SNARE recycling. Considering

this unexpected result, we thus focused on SNAREs with at least one high-confidence crosslink

to the D2 domain of Sec18 (**Figures 1C-F**).

Specifically, seven of ten high-confidence D2 crosslinks connect to regions N-terminal to

a SNARE domain (**Figures 1C-F**). For example, for Sso1, the high-confidence crosslink to D2

involves its Habc domain (**Figure 1D, Extended Data Figure 1B**). The Habc domain is a stable

three-helix bundle[31]; thus, assuming complete threading through the D1 and D2 pores, the Habc

domain (~30 Å diameter) would have to be unfolded transiently. The remaining three high-

confidence D2 crosslinks connect to C-terminal regions within SNARE domains. At first glance,

this suggests complete substrate threading through both the D1 and D2 rings, akin to other

AAA+ protein translocases[38]. However, considering the topology imposed by membrane

domains, anchors, and folded N-terminal domains, how is a SNARE substrate loaded and

released? To answer these vexing topological questions, we next determined cryo-EM structures

of this orthologous yeast complex together with Sec18 and Sec17.

The Sec18—Sec17—Sso1—Snc1—Sec9 (y20S) complex

To corroborate the surprising LC-MS/MS results, we determined cryo-EM structures of a

yeast SNARE complex together with Sec18 and Sec17. We chose the Sso1—Sec9 —Snc1/Snc2

complex (referred to as ySNARE complex) since each component of this complex crosslinked

with high confidence to the D2 domain of Sec18 (Snc1 is highly homologous to Snc2). We

prepared the Sec18—Sec17—Sso1—Snc1—Sec9 (y20S) complex (**Extended Data Figure 2,**

Methods), and after single particle cryo-EM data collection and processing, we obtained

381,591 high-quality particles that yielded eight 3D classes into which models were built

(**Figure 2A–E, Supplementary Figure 2A-B, Supplementary Table 1**).

In the structures of all eight classes, Sso1 is threaded through the D1 pore of Sec18

(**Figure 2A**). A characteristic phenylalanine side chain density in Sso1 is present at the same

position relative to Sec18 in all classes, allowing for reliable indexing of Sso1. The nucleotide

states and the arrangements of D1 around the substrate were also the same among all eight

classes, with the ADP-engaged E protomer forming the top of a spiral staircase of tyrosine

residues. Protomers D, B, and C were all ATP bound and formed the middle of the staircase. The

bottom protomer, A, forms the base of the staircase. Protomer F, with no apparent density for
nucleotide in D1, was not associated with Sso1 (**Figure 2B**). This arrangement of the protomers
is driven, in part, by the angle between the more mobile D1 domain and rigid D2 domain. From
protomers E to A, the angle between the D1 and D2 decreases monotonically (**Figure 2C**). The
protomer at the top of the staircase maintains the largest angle between D1 and D2, while the one
at the bottom has the smallest. The eight classes from the non-hydrolyzing y20S dataset also
show differences in the ySNARE-Sec17-Sec18-N-domain subcomplex arrangement ("spire")
and Sec17 stoichiometry above the D1 ring of the y20S complex before disassembly (**Figure
2D-E, Extended Data Figure 3B-C**; **Supplementary Discussion**).
The high-confidence crosslinks for Sec18 intra- and inter-protomer interactions are
consistent with the cryo-EM structure, specifically with the arrangements of the N-domain and
D2-domain (**Figure 3A, Extended Data Figure 3A**). Moreover, inter-protomer crosslinks were
found between pore loop regions, and between the inter-protomer loop between residues 476 and
490 and N-domain linker regions of D1. These high-confidence crosslinks were within the
theoretical maximum crosslinking distance (N_ζ–N_ζ: ~40 Å)[39].
Remarkably, the structures reveal that Sso1 is also threaded through the D2 pore and that
the Habc domain of Sso1 interacts with the outside of the D2 ring, with class 1 having the most
well-defined, discrete density (**Figure 3A–E**). In class 1, the density for Sso1 within Sec18 is
continuous from the D1 pore entrance to the D2 pore exit, where the Habc domain begins
(**Figure 3A, D**). The three-helix bundle of the Habc domain is then positioned outside the D2
pore, where it packs between two small D2 subdomains (protomers A and B) in an interaction
driven by charge complementarity. Indeed, the electrostatic potential surface of the Habc domain
forms a roughly opposite that of the bottom of the D2 surface against which it packs (**Figure 3B-
C**). Given that this charge distribution is replicated between each pair of D2 subdomains around
the D2 surface, it is unsurprising that the Habc domain adopts multiple corresponding rotational
states over the eight classes, presumably due to the flexible linker that follows it (**Figure 3E**).
This structural observation is consistent with XL-MS data showing a crosslink between the Habc
and the D2 domain of Sec18 (**Figure 3A, Extended Data Figure 1B**). Although the putative
distance between the crosslinked residues (K570 in Sec18 and K111 of Sso1) observed in the
model for class 1 (42 Å) is at the limit for a crosslinking distance, the conformational flexibility
of the Habc domain to rotate and exist in multiple conformations about the D2 ring (**Figure 3E**)
likely produces conformations well within the crosslinking distance threshold.
187

Sec18 does not unfold the Sso1/syntaxin Habc domain

Given that this complex was assembled in non-hydrolyzing conditions, we next asked
how Sso1 is loaded into Sec18 given the diameter of the Habc domain of Sso1 is ~30 Å whereas
the D1 pore of Sec18 has a diameter of ~11 Å. In other AAA+ translocases, such as ClpX, ATP
hydrolysis drives complete substrate threading through the AAA+ pore[7]. However, in the case of

Sec18 in a non-hydrolyzing condition, such complete threading would require the Habc domain
to unfold without any energetic input from Sec18.
We employed two orthogonal approaches to test if the Sso1 Habc domain unfolds during
SNARE disassembly. In our first approach, we turned to a single-molecule Fluorescence
Resonance Energy Transfer (smFRET) system developed previously for NSF, α-SNAP, and
neuronal SNAREs[14] in which individual SNARE domains were synthetically linked and
stochastically labeled with FRET pairs. This system allows one to observe multiple rounds of
SNARE disassembly as the disassembled SNARE domains readily reassemble into a *cis*-SNARE
complex due to the covalent linkages. We chose this system since it is well-established.
Moreover, it is relevant for the Sec18 system studied here since the primary sequence is
conserved in the core regions of Sec18/NSF responsible for ATP hydrolysis and substrate
processing (**Extended Data Figure 4A**). In addition, the D1 and D2 rings are structurally
conserved when comparing the D1 and D2 rings of Sec18/NSF with engaged substrates, with an
average Cα RMSD of 1.64 Å (**Extended Data Figure 4B-C**). The only protomer that exhibits
much difference is the F protomer, which is likely due to its mobility and not due to inherent
differences between the structures. Furthermore, we tested the ability of Sec18 and Sec17 to
disassemble fluorophore-labeled neuronal SNARE and, *vice versa*, NSF and α-SNAP to
disassemble fluorophore-labeled exocytic ySNARE complex (**Extended Data Figure 5A-E,**
Supplementary Discussion). We found that both could process the species-ortholog SNARE
complex, albeit at different rates, further corroborating the interchangeability of NSF and Sec18.
We thus used the smFRET assay to study the effect of NSF on the dynamics of the three-
helix bundle Habc domain of syntaxin during disassembly. We either stochastically labeled the
SNARE domains or stochastically labeled the Habc domain at two distinct residue positions to
monitor either SNARE disassembly or the folded state of the Habc domain (**Figure 4A**). As in
our previous work[14], we observed repeated rounds of disassembly and re-assembly for the
SNARE domains as indicated by the changes in single FRET intensity over tens of seconds
(**Figure 4B-C**). In stark contrast, we did not observe a change in single-molecule FRET intensity
for the labeled Habc domain in non-disassembly and disassembly conditions, producing a FRET
intensity distribution consisting only of a high-FRET state (the small peak at 0 is due to traces
where acceptor photobleaching occurred since the particular traces never transitioned back to
high FRET and no change in the peak intensity between non-disassembly and disassembly
conditions was observed) (**Figure 4D-F**). This result suggests that the Habc domain does not
unfold during NSF activity.
To corroborate this finding with an orthogonal approach for the yeast system, we double-
crosslinked the Habc domain of Sso1 by incorporating an unnatural amino acid, 4-azido-L-
phenylalanine[40] (**Extended Data Figure 5F-G**). This double crosslink was introduced to prevent
the Habc domain from unfolding during any Sec18-driven threading. After crosslinking, we
performed a disassembly assay on fluorescently labeled un-crosslinked and crosslinked
complexes and monitored progress by native gel electrophoresis (**Figure 4G**). The top band,
representing the fully assembled exocytic SNARE complex, is present in both conditions but

disappears when adding Mg^{2+} to initiate hydrolysis and the disassembly reaction. There was no
qualitative difference in the disassembly between uncrosslinked and crosslinked conditions after
normalizing for labeling efficiency (**Figure 4H**). Thus, the folded state of the Habc domain does
not change during yeast SNARE disassembly, and crosslinking the Habc domain does not affect
the disassembly kinetics. Taken together, these results for both the yeast and neuronal systems
further argue against a threading model of Sso1 engagement since the Habc domain would be
unable to pass through the D1 and D2 domains.
240

Substrate-released Sec18 structures reveal coordinated ring opening

Together, these observations beg the question of how Sso1 enters the Sec18 ATPase
rings. To address this topological challenge, we next determined structures of the y20S complex
after initiating Sec18 hydrolysis by adding Mg^{2+} (referred to as "hydrolyzing condition").
Informed by a fluorescent protein disassembly assay (**Extended Data Figure 5H**), we initiated
the disassembly reaction and waited 7 seconds before sample vitrification.
In the resulting cryo-EM dataset, we found a small number of particles (6,356 or 2.26%
of final particles, class 1) that consist of entire y20S assemblies (**Figure 5A**), largely similar to
those observed in the non-hydrolyzing condition (**Figure 2**). While the resolution of this
reconstruction is low (10.88 Å), the D1 ring is flattened relative to reconstructions from the non-
hydrolyzing condition (**Figure 5B**). Considering that our sample consists of purified y20S
complexes before initiating disassembly by adding Mg^{2+}, this D1 ring flattening likely occurs
due to Mg^{2+} binding or ATP-hydrolysis at one or more subunits.
The remaining particles (274,883 or 97.74 % of final particles) were substrate-free
(classes 2–4 in **Figure 5C–E**). The D1 ring is also flat in all these substrate-free classes, with all
D1 protomers bound to ADP. This uniform binding to ADP, as opposed to the spiral staircase of
apo, ADP, and ATP-bound protomers, likely explains the flattened nature of the protomers. In
the D2 ring all protomers are still ATP bound, consistent with its role in oligomerization. Class 4
consists of a configuration where both the D1 and D2 rings are heptameric (**Figure 5C**), and
class 3 is similar to this heptameric class, except the 7th protomers is only partially occupied
(**Figure 5D).**
Class 2 has a well-resolved hexameric configuration with a coordinated split in class 2 in
the D1 and D2 rings (**Figure 5E**). This split spans ~20 Å between protomers and is large enough
to accommodate a polypeptide chain entering or leaving the rings. This conformational change is
likely induced by the nucleotide state, and specifically coupled to the presence of ADP
throughout the ring following hydrolysis and SNARE substrate processing; this opening allows
the processed SNARE substrate to be released from the side. This side-release mechanism thus
solves one part of the topological challenge state above; threading occurs, but the substrate exits
from the side, avoiding the membrane domains and linkages in SNARE proteins.
270

Substrate-free Sec18 structures reveal coordinated ring opening

Next, we asked if this same side-opening of Sec18 occurs in the hydrolyzing condition without SNARE substrate. We prepared purified wild-type Sec18 in the presence of ATP and Mg^{2+} in a three-stage procedure that yielded a pure sample used for cryo-EM studies (**Extended Data Figure 6A-C**). Two bands were visible in native gels, indicating two oligomeric states (**Extended Data Figure 6D**). Processing and classification led to three classes, two of which appeared to be heptameric and identical (280,935 particles and 73,253 particles, respectively; **Figure 6A**) and the third class, which contained indeterminate density in the D1 ring (70,513 particles). Density for the N-domains was not well resolved in any of the classes, presumably due to the conformational flexibility of the N-domains without substrate or adaptor present.

3D variability analysis (3DVA)[41] of the third class suggests that it consists of a mixture of heptameric and hexameric Sec18 (**Extended Figure 6E),** consistent with the two bands seen on a native gel of the sample**.** The hexameric class reconstructed from 3DVA particle slices from the first mode (**Figure 6B**) is like class 2 observed for the y20S complex under hydrolyzing conditions (**Figure 5C**), wherein hexameric Sec18 has a large, coordinated opening in its D1 and D2 domain rings without substrate. This similarity in Sec18 conformations provides further evidence that the substrate is both loaded and released through the side of the rings, explaining why the Habc domain is not unfolded during Sec18/NSF activity (**Figure 3**) and that crosslinking the Habc domain does substantially affect the disassembly kinetics (**Figure 4H**). This solves the second part of the topological question stated above.

The heptameric class (**Figure 6A**) is nearly identical to class 4 observed in the y20S complex in a hydrolyzing condition (**Figure 4A**). Each protomer in the heptamer is similar to the others, with an average RMSD of <1Å between them, with the angle between the D1 and D2 rings varying only with a small range of 61.9°–67.1°, in contrast to hexamer, with a range of 60.8°–92.3° (**Figure 2C, Extended Data Figure 6F**). The D1 pore diameter is 30 Å, larger than substrate-bound hexamer, with a diameter of ~11 Å. The substrate-free structures (hexamer and heptamer) are nearly identical (RMSD = 0.48 Å) to the substrate-released structures.

To test if these D1 ADP-bound, ring-flattened states of Sec18 observed by single particle cryo-EM convert to a disassembly-competent state, we tested the same protein preparation used for cryo-EM with a vacuolar/lysosome proteoliposome fusion assay[42,43] (**Extended Figure 6H-J, Supplementary Discussion**); this assay confirmed that the Sec18 preparation used for cryo-EM was active and processed vacuolar SNAREs. Moreover, NSF also adopts this flattened ring state and a heptameric or hexameric arrangement when exchanged into a buffer containing ATP and Mg^{2+} without substrate (**Extended Data Figure 7, Supplementary Discussion**), consistent with our observations that Sec18 and NSF share the same functional mechanism.

Implications for ATPase activity

The observed structures of Sec18 in substrate-free, substrate-loaded, and substrate-released states suggest large conformational change upon ATP hydrolysis, consistent with

previous observations[3,27,44]. In the substrate-free and substrate-released states, examination of the
D1 nucleotide binding pockets reveals critical catalytic residues far from the nucleotide (**Figure**
6C). However, in the substrate-engaged state, these conserved residues are tightly bound to
nucleotide in a state preceding the binding of divalent cation required to drive hydrolysis.
Employing the same statistical workflow used to cluster the different y20S classes (**Methods**),
we identified the most significant residues that vary between the no substrate and substrate-
bound state by generating an ensemble of models for each class via ensemble refinement
(residues colored red in **Figure 6D,** variance contribution per component in **Extended Figure**
5I). For both the arginine finger loop and Walker B motif, we observed substantial changes to
side chain and backbone conformations (**Figure 6E-F**), wherein coordinating residues are
positioned closer to the nucleotide-binding pocket in the presence of substrate, presumably
establishing a hydrolysis-ready state. In the case of the arginine fingers, the side chain
conformational change is likely coupled to nucleotide identity, as hydrolysis disrupts phosphate
coordination with ATP, weakens the interprotomer interface, and leads to rigid body motion of
the D1 large subdomain as the arginine fingers disengage. The case of the shift in the Walker B
element, on the other hand, appears to be the flexing of the loop without substantial changes to
the sidechains of D349 and E350, induced by the motions of pore loop 2 (GVG motif) that bind
to the substrate in the pore.

Discussion

Based on our LC-MS/MS, structural, and functional results, we propose the following
model for SNARE recycling (**Figure 6G**), consisting of four phases. Sec18/NSF transiently
forms split-open conformation of both D1 and D2 rings in the resting phase with no substrate.
Once the *cis*-SNARE complex has been coated with adaptor Sec17/α-SNAP molecules, the
Sec18 N-domains can bind the subcomplex and position the unstructured linker connecting the
N-terminal Sso1 Habc domain to the SNARE domain proximal to the D1/D2 split, thus
promoting the loading of substrate into the catalytic core of the enzyme. More generally, this
side-loading mechanism would allow Sec18/NSF to process complexes incorporating all
SNAREs in the cell regardless of the presence of folded N-terminal domains, provided that, in
the latter case, a long enough linker connects them to the following SNARE domain. Upon the
formation of the 20S particle, ATP-driven hydrolysis likely fuels the disassembly of the SNARE
complex by some degree of threading, accompanied by D1 ring flattening. In the case of Sso1,
since full threading cannot occur due to the transmembrane domain, partial melting of the N-
terminal end of the SNARE domain could be sufficient to destabilize the SNARE complex.
Finally, the pore-bound SNARE is released through the side-split of the D1 and D2 rings,
regardless of membrane anchors or transmembrane domains in SNARE proteins. This model
explains how Sec18/NSF can disassemble SNAREs with constrained topologies.
346
As observed in our cross-linking LC-MS/MS experiments (**Figure 1**), the Sec18 D2 ring
can interact with not just with the Habc domain of Sso1 but also with other N-terminal accessory

domains (GOS1, TLG1, TLG2, etc). In turn, N-terminal domains often interact with other
factors. For example, the trimeric Dsl1 complex binds to the SNAREs Use1, Ufe1, and Sec20
through interactions with N-terminal SNARE domains. This 255 kDa complex tethers
endoplasmic reticulum membranes together before forming the *trans*-SNARE complex and
likely remains complexed before, during, and after fusion[32]. Another example is HOPS, a
hexameric complex that tethers lysosomal/vacuolar membranes and orchestrates their fusion[45].
HOPS has two Rab-binding subunits for tethering and a SNARE binding subunit that catalyzes
initial SNARE complex assembly[46]. Sec18/NSF and Sec17/α-SNAP cooperate with HOPS to
ensure rapid and efficient SNARE-mediated fusion[47,48]. For the case of Dsl1, the N-terminal
domains of SNAREs remain in a supramolecular arrangement that must be bypassed. For
vacuolar/lysosomal fusion, SNAREs must be preprocessed and handed off to the HOPS
complex. These complexes pose topological constraints on both ends of the SNAREs that can be
resolved by side-loading and side-release.
362
Finally, we note that distantly related AAA+ clamp loaders and their clamp load their
nucleic acid substrate from the side as well[49]. DNA is loaded into a split opened clamp and
AAA+ clamp loader. Once the clamp loader recognizes the substrate, the ATPase is triggered,
and the clamp loader closes the complex around the DNA. Once the clamp is fully secured, the
clamp-loader is then released. This shared mechanism suggests the possibility of side-loading
and release across the entire AAA+ protein family.
369
In summary, Sec18/NSF can load and release SNARE substrates via a coordinated
opening in the D1 and D2 rings. This mechanism does not preclude the more conventional notion
of complete substrate threading through the pores of Sec18/NSF if the SNARE substrate is not
topologically restricted by membrane domains or anchors or by N-terminal domains or
complexes. However, considering these topological considerations are ubiquitous for SNAREs,
the side-loading/release mechanism is likely conserved for the entire Sec18/NSF class of AAA+
proteins.
377

Figure Legends

Figure 1: *In vivo* crosslinking mass-spectrometry of Sec18
a, Top left: Venn diagram of proteins found in the condition without crosslinking agent (red)
added and with crosslinking agent added (green). Bottom right: SAFE-analysis[34] of yeast
proteins that specifically crosslink to Sec18. Blobs represent enrichments in the pathway. Blobs
of different colors represent enrichments that are specific to different p-value thresholds (brown
< 0.01, blue < 0.001, red < 0.0001). **b,** Heatmap of yeast SNARE proteins crosslinking to
specific Sec18 positions. The Sec18 primary sequence along the x-axis is colored coded by
domain, and yeast SNARE proteins with at least one medium confidence crosslink are shown
along the y-axis. Red squares represent positions where one or more manually verified, high-

confidence crosslinks were found (**Methods**). Orange squares represent where one or more
manually verified medium-confidence crosslinks were found. White squares represent positions
in which manually verified crosslinks were not found or were of low confidence. **c-f,**
Crosslinking schematic of Sec18 and SNAREs that contain at least one high-confidence
crosslink to the Sec18 D2 domain. Red dashed lines represent high-confidence crosslinks, and
orange dashed lines represent medium-confidence crosslinks.
394
Figure 2: The y20S supramolecular complex
a, Domain diagrams of the components used to prepare the y20S complex; coloring follows this
throughout (top left). The structure of class 1 of y20S, with ATP and ADP nucleotides (red and
blue, respectively; bottom right). Cryo-EM density for class 1 and the Sso1 atomic model around
residue F173 (inset, bottom left). **b,** Sso1 substrate is engaged by Y315 in all D1 protomers with
bound nucleotide. **c,** Conformations of the protomers of Sec18 when engaged to substrate. D,
Top-down views of class 1 and class 8 cryo-EM maps. **e,** Cartoon top-down views of eight y20S
classes, which differ primarily by spire configuration, *i.e.*, the pattern of N-domain and α-SNAP
engagement.
404
Figure 3: Sso1 is threaded through the D1 and D2 domains
a, y20S class 1 atomic model emphasizing the position of Sso1. Stars represent the crosslink
identified by crosslinking mass spectrometry (**Figure 1B**). **b,** Bottom view of y20S class 1 D1
ring; the Habc domain is packed against the D2 ring surface, between protrusions corresponding
to the C-terminal helical regions of two D2 small subdomains. All surfaces are colored grey,
except for protomers A and B, which are colored by their electrostatic potential **c,** Rotated view
of y20S class 1 relative to **b** showing the Habc tucked between protomers A and B with now the
Habc domain colored by its electrostatic potential **d,** Composite cryoEM map (blue, Relion-
unsharpened map: Habc domain and D1 density and CryoSPARC 3DFlex map: D2 density) of
y20S class 1. Nucleotides are shown as red (ATP) or blue (ADP) spheres. **e,** Cryo-EM maps of
all eight y20S classes reveal the Habc domain associated with the Sec18 D2 ring in all classes
(red). In 6/8 classes, the Habc density is radially averaged into multiple discrete positions around
the D2 ring, leading to more diffuse density.
418
Figure 4: Sec18/NSF does not unfold the Habc domain
a, Diagram of the smFRET disassembly/re-assembly assay with linked neuronal SNAREs. The
linked SNAREs are either stochastically labeled within the SNARE domains or the syntaxin
Habc domain. **b,** Representative traces of labeled neuronal SNARE complex in the non-
hydrolyzing condition. Violet represents the acceptor, and orange represents the donor-dye
fluorescence-intensity time traces. n = 53. **c,** Representative time trace of labeled SNARE
complex in disassembly conditions, the black line represents fitting by a vbGMM[50]. n = 114. **d,**
Representative time-trace of labeled Habc domain in the non-hydrolyzing condition. Green
represents the donor, and red is the acceptor dye fluorescence intensity time-traces. n = 27. **e,**

Representative time-trace of labeled Habc domain in disassembly conditions. The black line
represents fitting a vbGMM. n = 22. **f,** Histograms of E-FRET for all four conditions. The blue
dotted line represents a Gaussian fit of the data. The y-axis is the normalized frequency of
occurrence of the E-FRET states in the traces. Top left: SNARE E-FRET for the non-
hydrolyzing condition, top right: SNARE E-FRET for the hydrolyzing condition. Habc E-FRET
for the non-hydrolyzing condition (bottom left). Habc E-FRET for the hydrolyzing condition
(bottom right). **g,** Native gel of labeled uncrosslinked or crosslinked yeast SNARE complex that
has been labeled with Oregon Green Maleimide 488. **h,** Quantification of gel densitometry of n =
3 independent disassembly experiments. Error bars represent the standard deviation of the mean.
Densitometries were all normalized to 0 min values.
438
Figure 5: Post-disassembly of y20S reveals substrate-released states
a, Cryo-EM map of y20S in the hydrolyzing condition, colored by domain. **b,** The distance
between the D1 and D2 domains decreases for some protomers under hydrolyzing conditions,
leading to D1 ring flattening. The orange model corresponds to protomer E from the merged
D1/D2 focused class in the non-hydrolyzing condition, which was rigid-body fit into the density
of the hydrolyzing condition to generate the blue model. **c,** Cryo-EM map and model of class 4
of Sec18, showing a heptamer. **d,** Cryo-EM map and atomic model of class 3 of Sec18,
suggesting a transition state. **e, Cryo-**EM map and atomic model of class 2 of Sec18, showing a
split hexamer. In panels **c-e**, the top is the EM-map, while the middle and bottom show atomic
models in top and side views, respectively.
449
Figure 6: Structures of substrate-free Sec18 in the hydrolyzing condition
a, Atomic model of the substrate-free heptameric state of Sec18 (class 1). **b,** Atomic model of
the split substrate-free hexameric state of Sec18 (derived from 3DVA of class 3). **c,** The D1
nucleotide binding pocket is remodeled as a function of SNARE substrate binding and nucleotide
state. The arginine fingers R406 and R409 are contributed by the neighboring protomer. The
substrate-free and substrate-released states are nearly identical in conformation. Water molecules
are shown as red spheres. **d**, Atomic model of Sec18 with protomer A colored blue if not a
significant residue from conformational analysis and colored red if significant (p-value < 0.05).
e, Atomic models showing the difference of the arginine finger loop in the no substrate
(transparent) and substrate condition. **f,** Atomic models of the loop proximal to the Walker B
motif in the no substrate (transparent) and substrate condition. For **d-f**, residues are colored red if
they are in the top 5% of residues that significantly vary between conditions. **g.** We propose a
general model of SNARE recycling, regardless of the cellular context. First, a *cis*-SNARE
complex is coated with at least 1 Sec17/α-SNAP adaptor molecule, allowing Sec18/NSF to
recognize it. The SNARE substrate is then loaded into a split hexamer with a coordinated
opening in both D1 and D2 rings through the side, bypassing whatever N-terminal domains may
be present. The substrate is then threaded coaxially through the Sec18/NSF pore. Upon
completion of processing, the SNARE substrate is released through the side, bypassing the

topological constraints of the membrane. Finally, Sec18 returns to its 'resting' state until more

substrate is encountered. Yellow arrows indicate the direction of loading and release.

Extended Data Figure Legends

Extended Data Figure 1: In vivo mass-spec workflow and additional analysis

a, Workflow of how samples were processed before mass-spec. **b,** Representative mass-

spectrometry raw data of high confidence **(Methods)**, suggesting that the Habc domain is

crosslinked to the D2 domain. In the sequence diagram, the cross-linked lysine residues are

shown in orange.

Extended Data Figure 2: Purification and assembly of y20S complex

a, Sec18 SDS-PAGE gel after the final SEC run. Pooled fractions were frozen and stored. **b,**

Sec17 SDS-PAGE gel after final SEC run. Pooled fractions were frozen and stored. **c,** Schematic

of y20S assembly. Snc1, Sso1, and Sec9 were purified individually before being co-folded for

complex formation (**Methods**). The resulting SNARE complex was added to purified Sec17,

EDTA was added, and then purified Sec18 was added. The resulting y20S complex was purified

by SEC, and fractions were pooled. The final gel shows reference inputs for Sec18, Sec17, and

yeast SNARE complex, the final concentrated y20S complex, and the fractions from SEC that

were pooled (yellow lines).

Extended Data Figure 3: Intra and inter-protomer crosslinks mapped on the y20S complex

structure

a, Two Sec18 protomers of the y20S complex are shown. The black dashed lines represent

crosslinks designated as excellent (passing all manual verifications) and superimposed over the

diagram. The two insets show inter-protomer crosslinks. **b,** Electrostatic surface potentials of the

interaction between SNAREs and Sec17. Negatively charged regions are colored red, and

positive ones are colored blue. **c,** Electrostatic surface potentials of the interaction between

Sec17 and the N-domains of Sec18. Alternating views of these interactions are shown to

demonstrate the complementarity of the electrostatic interactions.

Extended Data Figure 4: Sequence and structural conservation between NSF and Sec18

a, Primary sequence alignment between NSF *c. griseus* and Sec18 *s. cerevisiae*. **b,** cartoon

representation of an atomic model of Sec18 D1/D2 focused model colored by Cα root-mean-

square-difference (RMSD) to the structure of the binary SNARE complex with NSF and α-

SNAP[27]. **c,** Focused view of substrate in D1. The substrate bound to the D1 pore is represented

as spheres.

Extended Data Figure 5: Cross-complementation disassembly assays

a, Scheme of fluorescent-protein labeling. Snc1 was labeled with mTurq, and SSO1 was labeled with mVenus (**Methods**, referred to as fpySNARE). **b**, SDS Page gel of the final SEC purification of y20S complex. **c,** Representative traces from the disassembly assay (**Methods**). The mTurq emission is monitored, and an increase in emission as the complex suggests that SNARE disassembly occurs (loss of fluorescence resonance energy transfer to mVenus upon disassembly). **d-e**, Disassembly plots of SNAREs and NSF/Sec18 combinations. **f,** Residues that were replaced with p-azido-l-phenylalanine are shown as yellow spheres in a black box **g,** MS raw data of double crosslinked yeast exocytic SNARE complex. **h,** Michaelis-Menten diagram of the y20S disassembly reaction. The red line shows the fitted line used to calculate M-M metrics, whereas the black points and error bars show the actual data. i, Variance contributed by each principal component.

Extended Data Figure 6: Functional studies of Sec18.

a, Schematic of purification of Sec18. **b,** SDS-PAGE gel of final SEC fractions and the fractions that were pooled for CryoEM studies of the y20S complex and the vacuolar fusion assay. **c,** Corresponding SEC trace showing the Sec18 oligomer eluting at expected volume. **d,** Native gel of final sample showing two oligomeric species. **e,** Representative slices of class 3 of the y20S complex under the hydrolyzing condition, subjected to 3DVA revealing split hexamer class. **f,** Alignment of all seven protomers, where each protomer is colored uniquely. **g,** Cartoon depicting the large pore in the heptameric state of the y20S complex under the hydrolyzing condition. **h,** Cartoon of the vacuolar fusion assay [42,43] (**Methods**) used to determine the function of the same Sec18 sample used for CryoEM studies of the y20S complex. Fusion only occurs when Sec18 prepares SNAREs for fusion, leading to fluorescence. **i,** Representative traces of fusion activity. **j,** Dot-plots of all activity assays.

Extended Data Figure 7: CryoEM structures of substrate-free NSF in the hydrolyzing condition

a-d, CryoEM maps (sharpened cryoSPARC) in three orientations (top to bottom rows) of the four classes of NSF in the hydrolyzing condition. The NSF heptamer (class 1) is designated as class 1. Classes 2-4 showed hexameric NSF in similar orientations with varying resolution for the substrate, so the best one, class 3 (initial) (c), was chosen for 3DVA analysis in CryoSPARC. **e,** The main principal component of the 3DVA analysis was used to select a subset of particles clearly showing an NSF N-terminal domain that is engaged with the D2 ring and then re-refined against this subset of particles to yield the final map for class 3. **f-g,** insets showing that an N-domain acts as a bound substrate in class 3 (final).

Methods

HA-Sec18 *S. cerevisiae* **strain preparation and treatment**

Briefly, we created an *S. cerevisiae* strain containing HA-tagged Sec18 and treated it with
zymolyase and lyticase to dissolve its cell wall but maintain its integrity as a living cell
(spheroplast). We then incubated the spheroplasts with disuccinimidyl glutarate (DSG), a
membrane permeable protein crosslinking agent, for 30 minutes before cellular lysis and anti-HA
co-immunoprecipitation (coIP) to isolate Sec18 and any bound proteins to it. This sample was
then subjected to SDS-PAGE gel electrophoresis, and the bands corresponding to crosslinked
Sec18 were excised and sent for liquid chromatography-mass spectrometry (LC-MS/MS)
analysis. More specifically, tagged Sec18 (pYK175, 3x-Flag N-terminal, HA-tag C-terminal)
plasmid was transformed into *S. cerevisiae* S288C using the Frozen-EZ Yeast Transformation
Kit (Zymo Research T2001). Expression of tagged Sec18 was accomplished by growing yeast in
1% raffinose and 1% galactose media. Cells were spun down and treated with 100 units of
zymolyase (Zymo Research #1004) at 37°C for 1 hour. After incubation, spheroplasts were
pelleted with a 1000 xg spin. DSG diluted in 1x PBS was used to resuspend the pellet, and the
sample was incubated for 30 minutes. The excess DSG was then quenched with 1M Tris and
incubated for another 15 minutes. Cells were washed with cold PBS and were resuspended in
lysis buffer (5mM EDTA, 1mM ATP, 1mM TCEP, 150 mM NaCl, 10% glycerol, 0.5% NP40,
and 50 mM HEPES pH 7.4) and glass beads (BioSpec 11079105). This bead-lysis-cell solution
was vortexed for 5 minutes at 4°C three times with ice incubations between each vortexing. The
solution was then spun down at 10,000 xg for 5 minutes at 4°C. The supernatant was taken and
then spun down again at 10,000 xg for 15 minutes at 4°C. The supernatant was then taken and
subjected to magnetic-bead anti-HA coIP (Thermo Scientific 88838). The eluant of the beads
was run on an SDS-PAGE gel, and the bands corresponding to Sec18 were excised.
568
569

XL-MS sample preparation

Protein samples were embedded in Coomassie-stained gel bands, fixed in 1% acetic acid.
The gels were washed several times in 50 mM ammonium bicarbonate, followed by reduction
with 10 mM of dithriothreitol (DTT) and incubation for twenty minutes at room temperature.
They were then alkylated using 30 mM acrylamide for 30 minutes. This was proceeded by
overnight digestion at 37 degrees Celsius with 500ng of mass spectrometry grade trypsin/LysC
mix (Promega). Post-digestion, samples were quenched with formic acid (adjusted to a pH ~3)
and desalted using MonoSpin C18 Solid-Phase Extraction (SPE) columns (GL Sciences).
Finally, the samples were dried via SpeedVac (Thermo Scientific, San Jose, CA) and exchanged
into LC-MS reconstitution buffer (2% acetonitrile with 0.1% formic acid in water) for
instrumental analysis.
581
582

LC-MS/MS analysis

Mass spectrometry experiments were performed using an Orbitrap Eclipse Tribrid mass
spectrometer (Thermo Scientific, San Jose, CA) attached to an Acquity M-Class UPLC system

(Waters Corporation, Milford, MA). The UPLC system was set to a flow rate of 300 nL/min,
where mobile phase A was 0.2% formic acid in water and mobile phase B was 0.2% formic acid
in acetonitrile. The analytical column was prepared in-house with an I.D. of 100 microns pulled
to a nanospray emitter using a P2000 laser puller (Sutter Instrument, Novato, CA). The column
was packed with Dr. Maisch 1.9 micron C18 stationary phase to a length of approximately 25
cm. Peptides were directly injected into the column with a gradient of 3-45% mobile phase B,
followed by a high-B wash over a total of 80 minutes. The mass spectrometer was operated in a
data-dependent mode using CID fragmentation for MS/MS spectra generation.
The RAW data were analyzed using Byonic v5.2.5 (Protein Metrics, Cupertino, CA) to
identify peptides and infer proteins. Initial Byonic analyses used a concatenated FASTA file
containing the Uniprot *Saccharomyces cerevisiae* proteins and other likely contaminants and
impurities. Once sample complexity was determined, a second round of Byonic analyses was
completed using a targeted FASTA file, which included only the sequences present in the samples
and allowances for crosslinked peptides with the appropriate linker. Proteolysis with Trypsin/LysC
was assumed to be fully specific with up to two missed cleavage sites. The precursor ion tolerance
was set to 12 ppm. The fragment ion tolerance was set to 0.4 Da. Cysteine modified with
propionamide was set as a fixed modification in the search. Variable modifications included
oxidation on methionine and acetylation on protein N-terminus. Proteins were held to a false
discovery rate of 1% using the standard reverse-decoy technique (Elias & Gygi Nat. Meth. 2007).
Potential crosslinked peptides were then validated and graded as described previously
using Byologic v5.2.31 (Protein Metrics) (Kobilka & Benovic, Cell, 2017). Briefly, a series of
empirical constraints were used for inspection, including minimum peptide and crosslink partner
length, chromatographic profile, MS1 coelution and MS/MS spectrum fragmentation quality.
Following this, identified crosslinked peptides were further categorized into three groups: high
confidence, medium confidence, and low confidence, where "high confidence" succeeded on all
these rules and "low confidence" failed at least in three of these rules.
612
613

Preparation of y20S in the non-hydrolyzing condition
The y20S complex consists of the Sso1–Snc1–Sec9 SNARE (ySNARE) complex, the
Sec18 oligomeric complex, and Sec17 adaptor proteins, and it was assembled in a multi-stage
workflow. Starting with the yeast exocytic complex, individual SNARE components Sec9 (401-
651), Sso1 (1-265), and Snc1 (1-93) were purified individually. All three proteins were cloned
into the pET28b plasmid vector with a 6x-His tag and a TEV cleavage sequence N-terminal to
the SNARE protein sequence.
For Sec9, we used a lysis buffer of pH 7.5, 50 mM NaPi, 300 mM NaCl, 10 mM
Imidazole, 0.5 mM TCEP, 1% Triton X-100 and SIGMAFAST Protease Inhibitor Cocktail
Tablet (Sigma-Aldrich S8830) which was used to resuspend bacterial cell pellets from an 8L
culture of auto-induced One Shot BL21 (DE3) *E*. coli (Thermofisher C600003). This
resuspended lysate was subjected to sonication for 20 minutes (3 seconds on, 9 seconds off, 60%

amplitude), followed by a 30-minute 4000 xg spin and a 1-hour 40,000 xg spin. The supernatant
was bound (1 hour, 4°C), run through a Ni-NTA column, and washed with pH 7.5, 50 mM NaPi,
30 mM Imidazole, 300 mM NaCl, and 0.5 mM TCEP buffer. The sample is then eluted with a
pH 8 50 mM NaPi, 400 mM Imidazole, 300 mM NaCl, and 0.5 mM TCEP buffer. This sample is
then digested overnight with TEV protease in a dialysis buffer (pH 8 20 mM Tris, 250 mM
NaCl, 0.5 mM TCEP, and 1 mM EDTA). The cleaved sample was then subjected to a MonoS
5/50 GL with buffer A consisting of pH 8 20 mM Tris, 50 mM NaCl, 0.5 mM TCEP, 1mM
EDTA and with buffer B consisting of pH 8 20 mM Tris, 500 mM NaCl, 0.5 mM TCEP, and
1mM EDTA. Taking the majority peak corresponding to Sec9, we then subjected this peak to
size exclusion chromatography (SEC) with the HiLoad 16/60 Superdex 200. The peak
corresponding to Sec9 was concentrated and flash-frozen. We followed the same protocol for
individually purifying Snc1 and Sso1.
Once all three individual SNAREs were purified, a 1:1:1 molar ratio of these proteins
was added to a 6M GdHCl solution. This solution was then slowly dialyzed overnight at 4°C into
a solution of 250 mM NaCl, 50 mM HEPES pH 7.6. This sample was then diluted to a NaCl
concentration of 75 mM and then subjected to MonoQ with the low salt buffer at 75 mM and the
high salt buffer at 500 mM with both buffers in 50 mM HEPES pH 7.6. The peak corresponding
to a 1:1:1 ratio of all three bands, indicative of the yeast exocytic complex, was then
concentrated for y20S formation without freezing.
The complete Sec18 protein coding sequence was cloned into the pMZ0002/pYK103
backbone with a 6x-His tag and TEV cleavage sequence N-terminal to Sec18. The 8L culture of
One Shot BL21 (DE3) *E.* coli (Thermofisher C600003) was spun down and resuspended in lysis
buffer of pH 7.5 100 mM HEPES, 500 mM KCl, 5 mM ATP, 5mM MgCl$_2$, protease inhibitor
tablets, and benzonase. This lysate was sonicated for 20 minutes (3 seconds on, 9 seconds off,
60% amplitude) and clarified with a 45,000 xg spin for 30 minutes. The lysate was supplemented
with 20 mM Imidazole bound to Ni-NTA beads (4°C) and washed with pH 7.5 20 mM HEPES,
480 mM KCl, 0.5 mM ATP, 0.5 mM TCEP, 20 mM Imidazole, 1 mM MgCl$_2$, and 10% glycerol.
The sample was then eluted with pH 7.5 20 mM HEPES, 480 KCl, 0.5 mM ATP, 0.5 mM TCEP,
500 mM Imidazole, 1 mM MgCl$_2$, and 10% glycerol. The protein-containing peak was then
equilibrated in SEC buffer pH 6.8 20 mM PIPES, 125 mM KCl, 0.2M sorbitol, 5 mM MgCl$_2$, 2
mM ATP, 2 mM DTT, and 10% glycerol and injected a HiLoad 16/60 Superdex 200. The
fractions were pooled, concentrated, and snap-frozen for later usage. The band right below Sec18
is an oligomeric *E. coli* contaminant, which is removed in the final steps of y20S complex
purification (**Figure S2A, C**). Moreover, this contaminant is absent in the protocol for preparing
high-purity Sec18 (see below and **Figure S6B**).
The full Sec17 coding sequence was cloned with a 10x-His tag and a TEV cleavage
sequence fused N-terminally. The 8L culture of One Shot BL21 (DE3) *E. coli* was spun down
and resuspended in a lysis buffer of pH 8 50 mM NaPi, 300 mM NaCl, 20 mM Imidazole, and
0.5 mM TCEP. This mixture was then sonicated (3 seconds on, 9 seconds off, 60% amplitude)
and clarified at 45,000 xg for 30 minutes. This supernatant was then incubated with Ni-NTA

beads for an hour and then washed with lysis buffer. The protein was eluted with lysis buffer
supplemented with 500 mM Imidazole and 5 mM EDTA. TEV protease was added to pooled
fractions and slowly dialyzed overnight at 4°C against a pH 8 50 mM Tris, 100 mM NaCl, 1 mM
EDTA, 0.5 mM TCEP buffer. This sample was then diluted to a NaCl concentration of 50 mM
and run on a MonoQ 10/100 from a 50 mM to 500 mM salt gradient. The major peak
corresponding to Sec17 was then pooled and injected into a HiLoad 16/60 Superdex 200 SEC.
The major peak from this run was taken, pooled, and snap-frozen for later usage.
To assemble the entire y20S complex, Sec17 was added to the freshly prepared yeast
exocytic SNARE complex, followed by EDTA quenched Sec18 with a final ratio of
Sec18:SNARE:Sec17 at 1:1.67:10, where a total of 5580 picomoles of Sec18 was used, in the
following buffer: pH 8 50mM Tris, 150 mM NaCl, 1 mM TCEP, 1 mM ATP, and 1 mM EDTA.
Following this assembly, this mixture was injected onto a Superose 6 10/300 increase column,
and the peak corresponding to the y20S complex, as assessed by elution volume and
corresponding SDS-PAGE gel, was concentrated to ~40 mg/mL.
680
Single particle Cryo-EM grid preparation of y20S in the non-hydrolyzing condition
Quantifoil R1.2/1.3 200 mesh gold grids were treated with chloroform and dried
overnight. Grids were glow discharged, and 5 microliters of the sample (at a final concentration
of 20 mg/mL with 0.05% v/v Nonidet P-40) was blotted onto the grids and further blotted and
vitrified in liquid ethane using an FEI Vitrobot (ThermoFisher Scientific). Grids were blotted for
4 seconds with a 5-second wait time (reduced to 3 seconds in hydrolyzing condition).
687
688 **Sample preparation and single particle Cryo-EM of y20S in the hydrolyzing condition**
689 Samples were prepared identically to y20S in the non-hydrolyzing condition, except that excess
690 $MgCl_2$ was blotted onto grids immediately before vitrification, and the total blot and wait time
691 was reduced to 7 seconds before freezing.
692
Sample preparation and single particle Cryo-EM of substrate-free Sec18 in the hydrolyzing
condition
To prepare high-purity, substrate-free Sec18, an alternative protocol was required since
low-purity Sec18, which can form y20S, was not amenable to high-quality cryo-EM studies. We
used a protocol similar NSF[18], but at the final reassembly step, we used $MgCl_2$ instead of EDTA.
The cryo-EM studies were performed similarly to the studies of Y20S in the non-hydrolyzing
condition.
700
701 **Sample preparation and single particle Cryo-EM of NSF in the hydrolyzing condition**
702   A sample of high-purity NSF was prepared, as described in the previous section. This
703 sample was exchanged into a buffer supplemented with 1mM magnesium chloride instead of
704 EDTA and incubated before a final SEC and freezing.
705

Single particle cryo-EM data collection, processing, and model building
Single particle cryo-EM data collection and processing workflow are described
thoroughly in the **Extended Table 1** and **Supplementary Figure 2**. An FEI Titan Krios
(ThermoFisher Scientific) cryo-electron microscope with either a K3 Gatan or a Falcon F4i
camera was used for data collection. Micrographs were analyzed using a combination of initial
Relion 3.1[51] processing followed by additional rounds of 3D classification or heterogenous
refinement, and final refinements in CryoSPARC v4[52]. A general workflow description is
provided in **Supplementary Figure 2,** and specific dataset information is provided in
Supplementary Figures 2-6 as well.
y20S models were constructed by first docking the crystal structure of the exocytic
SNARE complex (PDB ID: 3B5N) and AlphaFold2[53] models for Sec17 into the unsharpened
density first obtained in Relion 3.1[51]. Next, sharpened cryoSPARC-v4[52] maps were then used to
model the Sec18 complex and the region of Sso1 that extended from the spire into the D1 pore.
For class 1, 3DFlex analysis in cryoSPARC-v4 was performed. A mixture of the Relion 3.1 and
cryoSPARC-v4 unsharpened and sharpened maps, and the 3Dflex map were used to model the
Habc domain outside of the D2 ring. The resulting atomic models were iteratively refined with a
combination of COOT[54], Phenix real-space refinement[55], and ISOLDE[56].
Models of substrate-free Sec18 and NSF were built *de novo* in COOT using cryoSPARC
sharpened maps, and iteratively refined with ISOLDE and Phenix.
725
smFRET disassembly/re-assembly assay
smFRET assays with linked neuronal SNAREs, NSF, α-SNAP, and reagents were
prepared as previously reported[14]. Briefly, linked SNARE complex (L-SNARE) composed of
SNAP25A, synaptobrevin-2, and syntaxin-1A were biotinylated and flowed onto the chamber
that was passivated by 10 mg/mL egg phosphatidylcholine 50 nm liposomes coated with 1
mg/mL of biotinylated BSA to mimic the lipid environment inside the cell. 0.1 mg/mL
streptavidin was added to surface-tether the biotinylated and labeled L-SNARE complexes. L-
20S was then assembled by first adding αSNAP and then later NSF (if in disassembly
conditions). L-SNARE was diluted such that about 500 molecules of L-SNARE per 45 x 90 μm^2
field of view was visible. The raw data was imported into tMaven[44] for analysis. The standard,
recommended tMaven workflow was employed. Traces were not subject to any preprocessing
workflow. Automatic photobleach detection was used to determine the locations of
photobleaching, and a vbGMM mixture model was used to detect the different states.
739
Crosslinking assays
Cysteines were introduced into Sec9 and Sso1 for labeling with Oregon Green Maleimide
488 (O6034). UAG stop codons were introduced at V57 and F83 in Sso1 for crosslinking. Cys-
Sec9, Cys-Sso1 (with or without stop codon depending on if used for crosslinked), and 6x-His-
TEV-Snc1 were co-expressed (if crosslinked, also transformed with pEVOL-pAzF AddGene
#31186, 2mM AzF and 0.2% v/v arabinose) and induced with IPTG. Cell pellets were lysed with

pH 7 50 mM Tris, 200 mM NaCl, 1% v/v Triton X-100, 20 mM Imidazole, 0.5 mM TCEP and
protease inhibitor tablets. Lysate was sonicated for 20 minutes (3 seconds on, 9 seconds off, 60%
amplitude) and then clarified at 40,000 xg for 30 minutes. Following a 1-hour incubation at 4°C,
the beads were washed with two different buffers. Buffer 1 is simply a lysis buffer without
Triton X-100, and Buffer 2 is a high salt wash of pH 7 50 mM Tris, 1M NaCl, 50 mM Imidazole,
0.5 mM TCEP. The complex was eluted with pH 7 50 mM Tris, 200 mM NaCl, 350 mM
Imidazole, and 0.5 mM TCEP. The sample was then digested overnight with TEV protease and
dialyzed into pH 7 20 mM Tris, 200 mM NaCl, 0.5 mM TCEP. This sample was then injected
onto a HiLoad 16/60 Superdex 200. The fractions containing all three in equimolar ratios were
taken and concentrated. Samples were labeled with Oregon Green Maleimide 488 in molar
excess overnight before cleanup with a Zeba spin column (ThermoFisher 89889). If the sample
was meant to be crosslinked, it was subjected to UV irradiation before Oregon Green Maleimide
labeling.
For native gel disassembly assays, reactions were assembled in test tubes, initiated with
$MgCl_2$, and quenched with EDTA. Samples were then loaded into Any kD Mini-Protean gels
(Bio-Rad 4569033) and used Tris-glycine running buffer (Bio-Rad) supplemented with 1 mM
ATP to prevent Sec18 from falling apart in the gel. Gels were first visualized in an iBright 1500
(ThermoFisher Scientific) to visualize ySNARE bands only before coomassie staining. Gel
densitometry was assessed in ImageJ.
765
FRET and fluorescent dequenching disassembly assays
Neuronal disassembly assays were carried out as previously reported[10,45], leveraging
fluorescence dequenching of the neuronal SNARE complex labeled with Oregon Green
Maleimide 488. For yeast SNARE disassembly assays, a strategy was devised to tag Snc1 with
mTurq and Sso1 with mVenus. When assembled, excitation by a 434 nm laser leads to mVenus
emission. 434 nm laser excitation leads to mTurq emission. Disassembly is followed by
monitoring excitation/emission at 434/474 nm such that as the SNARE complex is disassembled,
mTurq emission increases. We used the same instrument (FlexStation II 384, Molecular Devices)
and a 384-well format to perform both the dequenching and FRET assays that were initiated by
titrating $MgCl_2$ in excess of the reactions.
776
Structural analysis
Statistical structural analysis was performed to identify the most significant backbone
motions between the many different structural states observed. Using Python 3, principle
component analysis (PCA) on backbone ϕ/ψ angles was performed, and the single mode that
contributed most to the variance (>70%) was analyzed. y20S models were then clustered in this
reduced 5-dimensional space. When comparing the substrate-free and substrate-engaged states,
ensembles consistent with a given map were generated to avoid model-building bias using
Phenix's simple molecular dynamics implementation for 2000 steps followed by a round of
phenix.refine[55]. These ensembles were then subjected to backbone ϕ/ψ PCA, and the mode that

contributed to the largest variance between the two clusters for the two classes was used to
determine which residues contributed most to the difference in structures.
788
Vacuolar fusion assay
Assays were performed as previously reported[42,43] with the same substrate-free Sec18
protein sample/batch that was purified and used for CryoEM. Briefly, proteoliposomes with
vacuolar SNAREs were prepared with either Phycoerythirin-biotin (PhycoE) or Cy5 and
incubated in a fluorescence plate reader for 40 minutes. FRET between PhycoE and Cy5 was
measured in intervals of 5-15s. Reaction mixtures contained HOPS complex, Sec17, and variable
amounts of Sec18.
796
Data availability statement
The cryo-EM maps will be deposited at the Electron Microscopy Data Bank (EMDB) and
the coordinates will be deposited in the Protein Data Bank. Motion-corrected micrographs will
be deposited in the Electron Microscopy Public Image Archive (EMPIAR). The data will be
made publicly available as of the date of publication. Original code reported in this paper will be
deposited on a publicly available GitHub page. The mass spectroscopy data will be deposited in
a publicly available repository.
804

Acknowledgments
We thank William I. Weis and Joseph D. Puglisi for stimulating discussions. We thank
the National Institutes of Mental Health (NIMH) for support (RO1MH63105 to A.T.B.,
1F31MH134477 to Y.A.K.). Y.A.K was additionally supported in part by an NSF GRFP and the
Knight-Hennessy Scholarship. K.I.W. was supported by a postdoctoral fellowship from the
Helen Hay Whitney Foundation, supported by the Howard Hughes Medical Institute. W.T.W. is
supported by a grant from the NIH (2R35GM118037). We thank the Vincent Coates Foundation
Mass Spectrometry Laboratory, Stanford University Mass Spectrometry (RRID:SCR_017801)
for utilizing the Thermo Orbitrap Eclipse nanoLC/MS system (RRID:SCR_022212) that was
purchased with funding from the National Institutes of Health Shared Instrumentation Grant
1S10OD030473, the Stanford Cancer Institute Proteomics/Mass Spectrometry Shared Resource
(NIH P30 CA124435).

816
Author contributions
Y.A.K and A.T.B conceived and designed experiments. K.I.W, B.S., and E.M. assisted
Y.A.K. with cryo-EM data collection. Y.A.K. performed and analyzed all experimental data.
K.I.W. assisted Y.A.K. with cryo-EM data processing and model building. R.A.F. and L.E.
assisted with protein purification. G.M., F.L., and T.M. assisted with high-resolution MS
proteomics data collection and analysis. K.D. assisted with bulk disassembly assays. U.B.C.
collected and assisted with smFRET data. W.T.W. performed and analyzed vacuolar fusion
assays. Y.A.K., A.T.B., and K.I.W. wrote the manuscript.

References

1. Sutton, R. B., Fasshauer, D., Jahn, R. & Brunger, A. T. Crystal structure of a SNARE

complex involved in synaptic exocytosis at 2.4 Å resolution. *Nature* **395**, 347–353 (1998).

2. Weber, T. *et al.* SNAREpins: Minimal Machinery for Membrane Fusion. *Cell* **92**, 759–772

(1998).

3. Zhao, M. *et al.* Mechanistic insights into the recycling machine of the SNARE complex.

Nature **518**, 61–67 (2015).

4. Söllner, T., Bennett, M. K., Whiteheart, S. W., Scheller, R. H. & Rothman, J. E. A protein

assembly-disassembly pathway in vitro that may correspond to sequential steps of synaptic

vesicle docking, activation, and fusion. *Cell* **75**, 409–418 (1993).

5. Jahn, R., Cafiso, D. C. & Tamm, L. K. Mechanisms of SNARE proteins in membrane fusion.

Nat. Rev. Mol. Cell Biol. 1–18 (2023) doi:10.1038/s41580-023-00668-x.

6. White, K. I., Zhao, M., Choi, U. B., Pfuetzner, R. A. & Brunger, A. T. Structural principles of

SNARE complex recognition by the AAA+ protein NSF. *eLife* **7**, e38888 (2018).

7. Maillard, R. A. *et al.* ClpX(P) Generates Mechanical Force to Unfold and Translocate Its

Protein Substrates. *Cell* **145**, 459–469 (2011).

8. Wickner, W. & Schekman, R. Membrane fusion. *Nat. Struct. Mol. Biol.* **15**, 658–664 (2008).

9. Südhof, T. C. Neurotransmitter Release: The Last Millisecond in the Life of a Synaptic

Vesicle. *Neuron* **80**, 675–690 (2013).

10. Ungar, D. & Hughson, F. M. SNARE Protein Structure and Function. *Annu. Rev. Cell Dev.*

Biol. **19**, 493–517 (2003).

11. Rizo, J. Molecular Mechanisms Underlying Neurotransmitter Release. *Annu. Rev. Biophys.*

51, 377–408 (2022).

12. Liang, B., Kiessling, V. & Tamm, L. K. Prefusion structure of syntaxin-1A suggests pathway

for folding into neuronal trans-SNARE complex fusion intermediate. *Proc. Natl. Acad. Sci.*

110, 19384–19389 (2013).

13. White, K. I., Zhao, M., Choi, U. B., Pfuetzner, R. A. & Brunger, A. T. Structural principles of

SNARE complex recognition by the AAA+ protein NSF. *eLife* **7**, e38888 (2018).

14. Choi, U. B. *et al.* NSF-mediated disassembly of on- and off-pathway SNARE complexes and

inhibition by complexin. *eLife* **7**, e36497 (2018).

15. Yoon, T.-Y. & Munson, M. SNARE complex assembly and disassembly. *Curr. Biol.* **28**,

R397–R401 (2018).

16. Khan, Y. A., White, K. I. & Brunger, A. T. The AAA+ superfamily: a review of the structural

and mechanistic principles of these molecular machines. *Crit. Rev. Biochem. Mol. Biol.* **57**,

156–187 (2022).

17. Hanson, P. I. & Whiteheart, S. W. AAA+ proteins: have engine, will work. *Nat. Rev. Mol. Cell*

Biol. **6**, 519–529 (2005).

18. Puchades, C., Sandate, C. R. & Lander, G. C. The molecular principles governing the

activity and functional diversity of AAA+ proteins. *Nat. Rev. Mol. Cell Biol.* **21**, 43–58 (2020).

19. Nagiec, E. E., Bernstein, A. & Whiteheart, S. W. Each Domain of the N-Ethylmaleimide-

sensitive Fusion Protein Contributes to Its Transport Activity (*). *J. Biol. Chem.* **270**, 29182–

29188 (1995).

20. Novick, P., Field, C. & Schekman, R. Identification of 23 complementation groups required

for post-translational events in the yeast secretory pathway. *Cell* **21**, 205–215 (1980).

21. Ma, C., Su, L., Seven, A. B., Xu, Y. & Rizo, J. Reconstitution of the vital functions of Munc18

and Munc13 in neurotransmitter release. *Science* **339**, 421–425 (2013).

22. Lai, Y. *et al.* Molecular Mechanisms of Synaptic Vesicle Priming by Munc13 and Munc18.

Neuron **95**, 591-607.e10 (2017).

23. Gao, Y. *et al.* Sensory deficit screen identifies nsf mutation that differentially affects SNARE

recycling and quality control. *Cell Rep.* **42**, 112345 (2023).

24. Leitz, J. *et al.* Beyond the MUN domain, Munc13 controls priming and depriming of synaptic

vesicles. *Cell Rep.* **43**, 114026 (2024).

25. Ryu, J.-K. *et al.* Spring-loaded unraveling of a single SNARE complex by NSF in one round

of ATP turnover. *Science* **347**, 1485–1489 (2015).

26. Ryu, J.-K., Jahn, R. & Yoon, T.-Y. Review: Progresses in understanding N-ethylmaleimide

sensitive factor (NSF) mediated disassembly of SNARE complexes. *Biopolymers* **105**, 518–

531 (2016).

27. White, K. I. *et al.* NSF converts syntaxin clusters to a priming-ready state. *Unpublished*.

28. Whiteheart, S. W. *et al.* N-ethylmaleimide-sensitive fusion protein: a trimeric ATPase whose

hydrolysis of ATP is required for membrane fusion. *J. Cell Biol.* **126**, 945–954 (1994).

29. de la Peña, A. H., Goodall, E. A., Gates, S. N., Lander, G. C. & Martin, A. Substrate-

engaged 26S proteasome structures reveal mechanisms for ATP-hydrolysis–driven

translocation. *Science* **362**, eaav0725 (2018).

30. Rizo, A. N. *et al.* Structural basis for substrate gripping and translocation by the ClpB AAA+

disaggregase. *Nat. Commun.* **10**, 2393 (2019).

31. Fernandez, I. *et al.* Three-dimensional structure of an evolutionarily conserved N-terminal

domain of syntaxin 1A. *Cell* **94**, 841–849 (1998).

32. DAmico, K. A. *et al.* Structure of a membrane tethering complex incorporating multiple

SNAREs. *Nat. Struct. Mol. Biol.* 1–9 (2024) doi:10.1038/s41594-023-01164-8.

33. Wang, T., Li, L. & Hong, W. SNARE proteins in membrane trafficking. *Traffic* **18**, 767–775

(2017).

34. Baryshnikova, A. Systematic Functional Annotation and Visualization of Biological

Networks. *Cell Syst.* **2**, 412–421 (2016).

35. Thomas, P. D. *et al.* PANTHER: Making genome-scale phylogenetics accessible to all.

Protein Sci. **31**, 8–22 (2022).

36. Kato, M. & Wickner, W. Ergosterol is required for the Sec18/ATP-dependent priming step of

homotypic vacuole fusion. *EMBO J.* **20**, 4035–4040 (2001).

37. Ungermann, C., Wickner, W. & Xu, Z. Vacuole acidification is required for trans-SNARE

pairing, LMA1 release, and homotypic fusion. *Proc. Natl. Acad. Sci.* **96**, 11194–11199

(1999).

38. Baker, T. A. & Sauer, R. T. ClpXP, an ATP-powered unfolding and protein-degradation

machine. *Biochim. Biophys. Acta BBA - Mol. Cell Res.* **1823**, 15–28 (2012).

39. Merkley, E. D. *et al.* Distance restraints from crosslinking mass spectrometry: Mining a

molecular dynamics simulation database to evaluate lysine–lysine distances. *Protein Sci.*

Publ. Protein Soc. **23**, 747–759 (2014).

40. Chin, J. W. *et al.* Addition of p-Azido-l-phenylalanine to the Genetic Code of Escherichia

coli. *J. Am. Chem. Soc.* **124**, 9026–9027 (2002).

41. Punjani, A. & Fleet, D. J. 3D variability analysis: Resolving continuous flexibility and discrete

heterogeneity from single particle cryo-EM. *J. Struct. Biol.* **213**, 107702 (2021).

42. Zucchi, P. C. & Zick, M. Membrane fusion catalyzed by a Rab, SNAREs, and SNARE

chaperones is accompanied by enhanced permeability to small molecules and by lysis. *Mol.*

Biol. Cell **22**, 4635–4646 (2011).

43. Zick, M., Stroupe, C., Orr, A., Douville, D. & Wickner, W. T. Membranes linked by trans-

SNARE complexes require lipids prone to non-bilayer structure for progression to fusion.

eLife **3**, e01879.

44. Shein, M. *et al.* Characterizing ATP processing by the AAA+ protein p97 at the atomic level.

Nat. Chem. **16**, 363–372 (2024).

45. Shvarev, D. *et al.* Structure of the HOPS tethering complex, a lysosomal membrane fusion

machinery. *eLife* **11**, e80901 (2022).

46. Song, H., Orr, A. S., Lee, M., Harner, M. E. & Wickner, W. T. HOPS recognizes each

SNARE, assembling ternary trans-complexes for rapid fusion upon engagement with the 4th

SNARE. *eLife* **9**, e53559 (2020).

47. Song, H., Torng, T. L., Orr, A. S., Brunger, A. T. & Wickner, W. T. Sec17/Sec18 can support

membrane fusion without help from completion of SNARE zippering. *eLife*

https://elifesciences.org/articles/67578 (2021) doi:10.7554/eLife.67578.

48. Orr, A. & Wickner, W. Sec18 supports membrane fusion by promoting Sec17 membrane

association. *Mol. Biol. Cell* **33**, ar127 (2022).

49. Kelch, B. A., Makino, D. L., O'Donnell, M. & Kuriyan, J. Clamp loader ATPases and the

evolution of DNA replication machinery. *BMC Biol.* **10**, 34 (2012).

50. Verma, A. R., Ray, K. K., Bodick, M., Kinz-Thompson, C. D. & Gonzalez, R. L. Increasing

the accuracy of single-molecule data analysis using tMAVEN. *Biophys. J.* (2024)

doi:10.1016/j.bpj.2024.01.022.

51. Scheres, S. H. W. A Bayesian view on cryo-EM structure determination. *J. Mol. Biol.* **415**,

406–418 (2012).

52. cryoSPARC: algorithms for rapid unsupervised cryo-EM structure determination - PubMed.

https://pubmed.ncbi.nlm.nih.gov/28165473/.

53. Jumper, J. *et al.* Highly accurate protein structure prediction with AlphaFold. *Nature* **596**,

583–589 (2021).

54. Emsley, P. & Cowtan, K. Coot: model-building tools for molecular graphics. *Acta Crystallogr.*

D Biol. Crystallogr. **60**, 2126–2132 (2004).

55. Adams, P. D. *et al.* The Phenix Software for Automated Determination of Macromolecular

Structures. *Methods San Diego Calif* **55**, 94–106 (2011).

56. Croll, T. I. ISOLDE: a physically realistic environment for model building into low-resolution

electron-density maps. *Acta Crystallogr. Sect. Struct. Biol.* **74**, 519–530 (2018).

Figure 1

Figure 2

A
Sso1

D1

D2

Habc

180°

B
Habc

Protomer A   Protomer B

C
Habc

Protomer B   Protomer A

180°

Electrostatic
potential (kT/e)
-5   0   5

D
Composite
EM map

D2

D1

E   Class averages

①   ②

③   ④   ⑤

⑥   ⑦   ⑧

Figure 3

Figure 4

Figure 5

Figure 6

Extended Data Table 1 | Cryo-EM data collection, refinement, and validation statistics

	Y20S, non-hydrolyzing condition									Y20S, hydrolyzing condition				Sec18, hydrolyzing condition		NSF, hydrolyzing condition	
	#1 Class 1 (EMDB-) (PDB)	#2 Class 2 (EMDB-) (PDB)	#3 Class 3 (EMDB-) (PDB)	#4 Class 4 (EMDB-) (PDB)	#5 Class 5 (EMDB-) (PDB)	#6 Class 6 (EMDB-) (PDB)	#7 Class 7 (EMDB-) (PDB)	#8 Class 8 (EMDB-) (PDB)	#9 All Merge D1/D2 (EMDB-) (PDB)	#10 Class 1 (no model deposited)	#11 Class 2 (EMDB-) (PDB)	#12 Class 3 (EMDB-) (PDB)	#13 Class 4 (EMDB-) (PDB)	#14 Class 1 (EMDB-) (PDB)	#15 Class 2 (EMDB-) (PDB)	#16 Class 1 (EMDB-) (PDB)	#17 Class 2 (EMDB-) (PDB)
Data collection and processing																	
Magnification	130,000×	130,000×	130,000×	130,000×	130,000×	130,000×	130,000×	130,000×	130,000x	130,000x	130,000x	130,000x	130,000x	130,000x	130,000x	130,000x	130,000x
Camera	Gatan K3	Gatan K3	Gatan K3	Gatan K3	Gatan K3	Gatan K3	Gatan K3	Gatan K3	Gatan K3	Falcon 4i	Falcon 4i	Falcon 4i	Falcon 4i	Gatan K3	Gatan K3	Gatan K3	Gatan K3
Voltage (kV)	300	300	300	300	300	300	300	300	300	300	300	300	300	300	300	300	300
Electron exposure (e–/Å^2)	26.8	26.8	26.8	26.8	26.8	26.8	26.8	26.8	26.8	51.48	51.48	51.48	51.48	25.5	25.5	39.04	39.04
Defocus range (μm)	-1.0 to -2.0	-1.0 to -2.0	-1.0 to -2.0	-1.0 to -2.0	-1.0 to -2.0	-1.0 to -2.0	-1.0 to -2.0	-1.0 to -2.0	-1.0 to -2.0	-1.0 to -2.0	-1.0 to -2.0	-1.0 to -2.0	-1.0 to -2.0	-1.0 to -2.0	-1.0 to -2.0	-1.0 to -2.0	-1.0 to -2.0
Pixel size (Å, super-res)	0.548	0.548	0.548	0.548	0.548	0.548	0.548	0.548	0.548	0.49	0.49	0.49	0.49	0.548	0.548	0.548	0.548
Exposure time (seconds)	4	4	4	4	4	4	4	4	4	6.93	6.93	6.93	6.93	4	4	4	4
Number of frames per exposure	80	80	80	80	80	80	80	80	80	2133 (EER fractionation of 42)	2133 (EER fractionation of 42)	2133 (EER fractionation of 42)	2133 (EER fractionation of 42)	80	80	80	80
Number of movies	12,439	12,439	12,439	12,439	12,439	12,439	12,439	12,439	12,439	14455	14455	14455	14455	7821	7821	13375	13375
Symmetry imposed	C1	C1	C1	C1	C1	C1	C1	C1	C1	C1	C1	C1	C1	C1	C1	C1	C1
Final particle images (no.)	41,159	35,723	79,060	26,603	33,084	64,857	69,608	31,497	381,591	6,356	86,542	91,962	96,291	280,935	16,074	72,069	11,844
Map resolution (Å)	3.89	3.91	3.73	4.61	4.29	3.73	3.75	4.13	3.4	10.88	3.38	3.36	3.18	2.99	6.09	3.60	4.70
FSC threshold	0.143	0.143	0.143	0.143	0.143	0.143	0.143	0.143	0.143	0.143	0.143	0.143	0.143	0.143	0.143	0.143	0.143
Model information																	
Initial model used (PDB code)	3B5N, Sec17/18 from AlphaFold2	3B5N, Sec17/18 from AlphaFold2	3B5N, Sec17/18 from AlphaFold2	3B5N, Sec17/18 from AlphaFold2	3B5N, Sec17/18 from AlphaFold2	3B5N, Sec17/18 from AlphaFold2	3B5N, Sec17/18 from AlphaFold2	3B5N, Sec17/18 from AlphaFold2	Sec18 from AlphaFold2	Y20S EDTA Class 1	*De novo*	*De novo*	*De novo*	*De novo*	*De novo*	3J94	3J94
Bond RMSD (Å)	0.12	0.012	0.007	0.007	0.010	0.005	0.005	0.010	0.006	No model deposited	0.004	0.013	0.007	0.007	0.006	0.006	0.007
Angle RMSD (°)	1.461	1.473	1.186	1.168	1.206	1.023	1.027	1.243	0.920	No model deposited	1.203	1.444	1.186	1.160	1.090	0.715	1.313
Molprobity score	1.68	1.79	1.58	1.79	1.72	1.73	1.62	1.76	1.55	No model deposited	1.23	1.78	1.78	1.51	1.80	2.00	1.72
Clashscore, all atoms	5.77	8.55	5.80	8.26	8.37	9.22	6.68	7.71	4.01	No model deposited	2.67	8.27	4.49	4.52	9.44	5.37	9.81
Ramachandran. Favored/Allowed/Disfacored (%)	94.77 4.85 0.38	95.35 4.49 0.16	96.15 3.75 0.10	95.12 4.54 0.34	96.11 3.69 0.20	96.41 3.45 0.14	96.24 3.67 0.08	95.81 3.88 0.31	94.80 5.01 0.19	No model deposited	96.95 2.92 0.14	95.16 4.65 0.18	95.13 4.58 0.29	95.95 4.05 0.00	95.73 4.17 0.00	95.64 4.36 0.00	97.17 2.26 0.57
Rotamer outliers (%)	0.61	0.44	0.43	0.69	0.88	0.55	0.52	1.14	0.55	No model deposited	0.08	0.00	1.96	0.93	0.97	3.45	1.19
Cβ outliers (%)	0.02	0.04	0.00	0.04	0.02	0.02	0.00	0.02	0.00	No model deposited	0.04	0.10	0.00	0.00	0.00	0.00	0.00
CaBLAM outliers (%)	2.36	2.08	2.06	2.22	2.09	1.99	2.15	2.14	3.06	No model deposited	0.93	1.36	1.20	0.99	0.99	1.44	1.46

A

B

Extended Data Figure 1

A

Sec18 SEC pooled fractions

B

Sec17 SEC pooled fractions

C

Purified individually

Snc1   1   v-SNARE   93

Sso1   1   t-SNARE   hABC   265

Sec9   401   t-SNARE   t-SNARE   651

Mixed in 1:1:1 molar ratios in 6M GdHCl

Cofold ternary complex with dialysis

SEC and MonoQ

Input   Peak 1   Peak 2   Peak 3

◄ Sso1
◄ Sec9
◄ Snc1

Peak 2
Peak 1
Peak 3

A280 (arbitrary units)

Buffer (mL)

Concentrate, mix with purified Sec17, & incubate

Add purified Sec18 complex

SEC

Y20S complex

Void

A290 (arbitrary units)

Buffer (mL)

Void   Y20S complex

Sec18
Sso1
Sec17
Sec9
Snc1

Sec18
Sec17
Yeast Exocytic Complex
Concentrated Y20S

Extended Data Figure 2

A

Protomer C

Protomer D

Protomer C
K235

Protomer D
K482

Finger loop

Protomer D
K314

Protomer C
K314

Pore loop

B   C

-5   0   5
Electrostatic
potential (kT/e)

Extended Data Figure 3

A

Sec18/1-758	1 MFKIPGFGKAAANHTPPDMTNMDTRTRHLKVSNCPNNSYALANVAAVSPNDFPNNIYIII----DNLFVFTTRHSNDIPPGTIGFNGNDRTWGGWSL	93
NSF/1-744	1 - MAGRSMQAARCPTDELSLSNCAVVSEKDYQSGQHVIVRTSPNHKYIFTLRHTHPSVVPGSVAFSLPQRKWAGLSI	74
Sec18/1-758	94 NQDVQAKAFDLFKYSGKQSYLGSIDIDISFRARGKAVSTVFDQDELAKQFVRCYESQIFSPTQYLIMEFQGHFFDLKIRNVQAIDLGDIEPTSA--V	188
NSF/1-744	75 GQEIEVA--LYSFDKAKQCIGTMTIEIDFLQKKNIDSNPYDTDKMAAEFIQQFNNQAFSVGQQLVFSFNDKLFGLLVKDIEAMDPSILKGEPASGK	168
Sec18/1-758	189 ATGIETKGILTKQTQINFFKGRDGLVNLKSSNSLRPRSNAVIRPQFKFEDLGVGGLDKEFTKIFRRAFASRIFPPSVIEKLGISHVKGLLLYGPPGT	285
NSF/1-744	169 RQKII-EVGLVVGNSQVAFEKAENSSLNLIGKAKTKENRQSIINPDWNFEKMGIGGGLDKEFSDIFRRAFASRVFPPEIVEQMGCKHVKGILLYGPPGC	264
Sec18/1-758	286 GKTLIARKIGTMLNAKEPKIVNGPEILSKYVGSSEENIRNLFKDAEAEYRAKGEESSLHIIIFDELDSVFKQRGSRGDGTGVGDNVVNQLLAKMDGV	382
NSF/1-744	265 GKTLLARQIGKMLNAREPKVVNGPEILNKYVGESEANIRKLFADAEEEQRRLGANSGLHIIIFDEIDAICKQRGSMAGSTGVHDTVVNQLLSKIDGV	361
Sec18/1-758	383 DQLNNILVIGMTNRKDLIDSALLRPGRFEVQVEIHLPDEKGRLQIFDIQTKKMRENNMMSDDVNLAELAALTKNFSGAEIEGLVKSASSFAINKTVN	479
NSF/1-744	362 EQLNNILVIGMTNRPDLIDEALLRPGRLEVKMEIGLPDEKGRLQILHIHTARMRGHQLLSADVDIKELAVETKNFSGAELEGLVRAAQSTAMNRHIK	458
Sec18/1-758	480 IGKGATKLNTKDIAKLKVTREDFLNAL-NDVTPAFGISEEDLKTCVEGGMMLYSERVNSILKNGARYVRQVRESDKSRLVSLLIHGPAGSGKTALAA	575
NSF/1-744	459 ASTK-VEVDMEKAESLQVTRGDFLASLENDIKPAFGTNQEDYASYIMNGIIKWGDPVTRVLDDGELLVQQTKNSDRTPLVSVLLEGPPHSGKTALAA	554
Sec18/1-758	576 EIALKSGFPFIRLISPNELSGMSESAKIAYIDNTFRDAYKSPLNILVIDSLETLVDWVPIGPRFSNNILQMLKVALKRKPPQDRRLLIMTTTSAYSV	672
NSF/1-744	555 KIAEESNFPFIKICSPDKMIGFSETAKCQAMKKIFDDAYKSQLSCVVVVDDIERLLDYVPIGPRFSNLVLQALLVLLKKAPPQGRKLLIIGTTSRKDV	651
Sec18/1-758	673 LQQMDILSCFDNEIAVPNMTNLDELNNVMI-ESNFLDDAGRVKVINELSRSCPNFNVGIKKTLTNIETARH--DEDPVNELVELMTQSA------	758
NSF/1-744	652 LQEMEMLNAFSTTIHVPNIATGEQLLEALELLGNFKD-KERTTIAQQV--KGKKVWIGIKKLLMLIEMSLQMDPEYRVRKFLALLREEGASPLDFD	744

B

Sec18
NSF
Cα RMSD
0 Å   5 Å

C

Extended Data Figure 4

Extended Data Figure 5

Extended Data Figure 6

A Class 1

B Class 2

C Class 3 (initial)

D Class 4

E 3D variability analysis

Class 3 (final)

F N-domain central pore

G R205

Cluster A

Extended Data Figure 7