Rapid synthesis and screening of natively paired antibodies against influenza hemagglutinin stem via oPool+ display

Wenhao O. Ouyang[1], Huibin Lv[1,2], Wenkan Liu[1], Zongjun Mou[3], Ruipeng Lei[1], Tossapol Pholcharee[1], Yiquan Wang[1], Katrine E. Dailey[1], Akshita B. Gopal[1], Danbi Choi[1], Madison R. Ardagh[1], Logan Talmage[1], Lucia A. Rodriguez[1], Xinghong Dai[3], Nicholas C. Wu[1,2,4,5,§]

[1] Department of Biochemistry, University of Illinois at Urbana-Champaign, Urbana, IL 61801, USA

[2] Carl R. Woese Institute for Genomic Biology, University of Illinois at Urbana-Champaign, Urbana, IL 61801, USA

[3] Department of Physiology and Biophysics, Case Western Reserve University, School of Medicine, Cleveland, OH 44106, USA

[4] Center for Biophysics and Quantitative Biology, University of Illinois at Urbana-Champaign, Urbana, IL 61801, USA

[5] Carle Illinois College of Medicine, University of Illinois at Urbana-Champaign, Urbana, IL 61801, USA

[§] To whom correspondence should be addressed. Email: nicwu@illinois.edu (N.C.W.)

ABSTRACT

Antibody discovery is crucial for developing therapeutics and vaccines as well as understanding adaptive immunity. However, the lack of approaches to synthesize antibodies with defined sequences in a high-throughput manner represents a major bottleneck in antibody discovery. Here, we presented oPool$^+$ display, which combines oligo pool synthesis and mRNA display to construct and characterize many natively paired antibodies in parallel. As a proof-of-concept, we applied oPool$^+$ display to rapidly screen the binding activity of >300 natively paired influenza hemagglutinin (HA) antibodies against the conserved HA stem domain. Structural analysis of 16.ND.92, one of the identified HA stem antibodies, revealed a unique binding mode distinct from other known broadly neutralizing HA stem antibodies with convergent sequence features. Yet, despite such differences, 16.ND.92 remained broadly reactive and conferred *in vivo* protection. Overall, this study not only established an experimental platform that can be applied in both research and therapeutics to accelerate antibody discovery, but also provides molecular insights into antibody responses to the influenza HA stem, which is a major target for universal influenza vaccine development.

Keywords: High-throughput screening, influenza, hemagglutinin, antibody, broadly neutralizing, oligo pools, mRNA display, cryo-EM

INTRODUCTION

Antibodies are central to the immune system for protection against pathogen infection. Therefore, identification of antibodies that target pathogens of interest is key to the understanding of adaptive immunity as well as the development of effective therapeutics and vaccines. In recent years, advances in single-cell B-cell receptor sequencing (scBCR-seq) have greatly improved the capacity to discover novel antibodies[1]. Thousands of natively paired antibody sequences can be obtained from a single scBCR-seq experiment. By contrast, downstream characterization of these antibody sequences remains costly, labor intensive, and time consuming, involving cloning, expression, purification, and testing the binding activities of different antibodies individually. At the same time, protein display technologies offer a high-throughput solution for characterizing antibody binding activity[2], with antibody library construction being an essential first step. Methods for constructing antibody libraries with random heavy-light chain pairing from B cell repertoires are well-established[3,4]. However, there is a lack of approaches to synthesize custom-made antibody libraries with precise heavy-light chain pairing from a defined list of antibody sequences. This technical barrier has restricted the application of protein display technologies in antibody research.

Hemagglutinin (HA) is the major antigen of Influenza A virus. HA is a homotrimeric glycoprotein consisting of a hypervariable globular head domain atop a conserved stem domain[5,6]. The functions of HA are critical for viral entry. The globular head domain engages the sialylated receptor, whereas the stem domain possesses the membrane fusion machinery. In the past two decades, many human antibodies to the HA stem have been discovered and characterized[7-11]. In contrast to HA head antibodies, which are usually strain-specific, HA stem antibodies often cross-neutralize multiple influenza subtypes. Several recurring sequence features have been observed among HA stem antibodies isolated from different individuals, such as IGHV1-69, IGHV1-18, IGHD6-1, and IGHD3-9[7,11-14]. However, many HA stem antibodies do not contain any known recurring sequence features, suggesting there are additional ones have yet to be discovered. Comprehending the recurring sequence features

of HA stem antibodies is critical for the molecular understanding of broadly neutralizing

antibody responses at the population scale, which will in turn benefit the development of

universal influenza vaccines.

In this study, we developed oPool⁺ display, a rapid and cost-effective *in vitro* platform that

combined oligo pool synthesis with mRNA display to assemble and screen natively paired

antibodies in a highly parallel manner. As a proof-of-concept, we first synthesized a library of

325 natively paired HA antibodies, then screened them for specificity to the HA stem. Our

screen identified 12 HA stem antibody candidates that were not previously known to target the

HA stem. Cryo-electron microscopy (cryo-EM) analysis revealed that the binding mode of one

of these HA stem antibodies, 16.ND.92, was dominated by immunoglobulin heavy diversity 3-

3 (IGHD3-3) gene. 16.ND.92 also exhibited a unique CDR H3 conformation compared to other

known HA stem antibodies that relied on IGHD3-3 for binding. Yet, 16.ND.92 remained broadly

reactive and protected against lethal influenza challenge. This observation substantiated that

IGHD3-3 is a recurring sequence feature of HA stem antibodies and demonstrated its

versatility in binding to HA stem.

RESULTS

High throughput assembly of natively paired antibody library via oligo pools

We previously curated a dataset containing 5,561 human monoclonal HA antibodies, where

1,082 antibodies (19.5%) are known to bind to either the head domain or the stem domain[15].

Of the remaining 4,479 (80.5%) HA antibodies which lack epitope information, 292 have

complete heavy chain variable (V_H) and light chain variable (V_L) sequences available and are

not encoded by well-characterized sequence features of HA stem antibodies, namely IGHV1-

69, IGHV1-18, IGHV6-1, and IGHD3-9[7,12–14] (**Figure S1A and Table S1**). These 292 HA

antibodies were included in the synthesis of our natively paired antibody library. In addition,

three known stem antibodies, namely 31.a.55, AG11-2F01, and 042-100809-2F04[7,16], as well

as 30 known HA head antibodies[6,17–25] were included as positive and negative controls,

respectively, bringing the total library size to 325 antibodies **(Figure 1A)**.

To synthesize the natively paired antibody library in a high-throughput manner, we aimed to

leverage recent advances in oligo pool synthesis. The maximum length of each oligo in

commercial oligo pool synthesis is around 300 to 350 nucleotides. By contrast, the length of

a single-chain variable fragment (scFv), which is the smallest format of a human antibody, is

around 800 to 900 nucleotides. As a result, each given scFv sequence was split into 4 oligos

with overlaps at the diverse complementary-determining regions (CDRs), namely the CDRs

H1, H3, and L3 (**Figure 1B-C and Table S2**). We then performed codon randomization to

ensure that the overlaps among oligos for the same scFv are unique at the nucleic acid level.

This would help prevent mis-annealing between oligos from different scFvs, especially if they

shared similar amino acid sequences (**Figure 1B, Material and Methods**). Through an

overlap PCR, full-length scFv sequences could then be generated with the intended native V_H

and V_L pairing. Subsequently, this strategy was applied to synthesize our library of 325

antibodies using 13 oligo pools with 25 antibodies per pool (**Figure 1C-F, Figure S1B-C, and**

Table S2).

The length of the assembled product was consistent with that of full-length scFvs (**Figure 1D**

and Figure S2). PacBio sequencing revealed that 319 of 325 (98%) natively paired antibodies

were successfully assembled (**Table S4**). Over 88% of the assemblies represented natively

paired scFvs, in which half had no mutations (**Figure 1E**). Only around 10% of the assemblies

were non-natively paired scFvs with V_H and V_L from different antibodies. Mutations in natively

paired scFvs were likely introduced during oligo pool synthesis, which is known to be error-

prone[26,27]. Moreover, a Pearson correlation coefficient of 0.9 was observed between replicates

(**Figure 1F**), demonstrating the reproducibility of our oligo pool-based strategy for antibody

library synthesis. While we opted to assemble this antibody library by 25 antibodies per pool,

we have also tested the assembly accuracy of 50 antibodies per pool. Among the 50

antibodies, 48 (96%) were successfully assembled with a Pearson correlation coefficient of 0.98 between replicates **(Figure S2A-B)**, demonstrating the potential to further increase the throughput of our antibody library synthesis strategy.

Rapid screening of the natively paired antibody library by mRNA display

To identify HA stem antibodies from our natively paired antibody library, we utilized mRNA display[28–30], a well-established *in vitro* screening approach that allows rapid screening for protein binders (**Figure 2A-D**). Briefly, each scFv was covalently linked to the RNA molecule that encoded it, thus providing a phenotype-genotype linkage. The mRNA-displayed antibody library was then selected against H1 and H3 stem constructs that were designed based on H1N1 A/Brisbane/59/2007 HA and H3N2 A/Finland/486/2004 HA, respectively[31,32]. Both pre- and post-selection libraries were analyzed by PacBio sequencing to quantify the enrichment of each scFv during selection. Pearson correlation coefficients of 0.95 and 0.93 were observed between biological replicates of selections against H1 stem and H3 stem, respectively (**Figure 2B-C**), demonstrating the reproducibility of the selections. The positive controls were highly enriched, whereas the negative controls were not (**Figure 2B-C and Table S3**), further validating that selection had taken place. Besides, we identified one antibody candidate targeting H1 stem and 11 antibody candidates targeting H3 stem (**Figure 2D**). Of note, one of the H3 stem antibody candidates, AG2-G02, was concurrently identified as an H3 stem antibody using a machine learning approach and experimentally confirmed in another study of ours[15].

To validate our screening results, we used biolayer interferometry (BLI) to measure the binding activities of selected antibody candidates (**Figure 3A**), including one targeting H1 stem and five targeting H3 stem. We also selected two antibodies that had low enrichment in both H1 stem and H3 stem screens as negative controls. The binding activities of these antibody candidates were successfully confirmed using both fragment antigen-binding (Fab) and scFv

formats (**Figure 3B-C, Figure S3 and S4**). Together, our results showed that oPool⁺ display

enabled rapid discovery of natively paired HA stem antibodies.

AG11-2F01 and 16.ND.92 have similar sequence features but distinct binding modes

The H1 stem antibody identified from our screen was 16.ND.92, which was originally isolated

from a young individual in an H5N1 influenza vaccine trial[7]. 16.ND.92 was encoded by IGHV3-

74/IGHD3-3/IGKV1-5 (**Table S3**). Coincidentally, both IGHD3-3 and IGKV1-5 were utilized by

AG11-2F01, which was one of the two positive controls against H1 stem in our screen (**Figure**

2B and 3B). Moreover, 16.ND.92 and AG11-2F01 shared a similar FG[V/L] motif encoded by

the reading frame +3 of IGHD3-3 (**Figure S5**). This observation led us to hypothesize that

16.ND.92 and AG11-2F01 engaged the HA stem via similar binding modes. Consequently, we

determined the cryo-EM structures of H1N1 A/Solomon Islands/03/2006 (SI/06) HA in complex

with AG11-2F01 and 16.ND.92 to resolutions of 2.89 Å and 2.82 Å, respectively (**Figure 4A-**

E and Table S4). Contrary to our hypothesis, the cryo-EM structures revealed very different

binding modes between the two antibodies. While AG11-2F01 bound to HA stem horizontally,

16.ND.92 had a downward approaching angle towards HA (**Figure 4A**). Relatedly, the epitope

of 16.ND.92 shifted slight upward compared to that of AG11-2F01 (**Figure 4B**).

Although both 16.ND.92 and AG11-2F01 were encoded by IGKV1-5, their light chains

interacted with the HA stem differently. For example, V_L S30 of AG11-2F01 H-bonded with

HA2 Q38, whereas that of 16.ND.92 H-bonded with HA1 K32 (**Figure 4C**). Similarly, despite

sharing an FG[V/L] motif in their IGHD3-3-encoded regions, 16.ND.92 and AG11-2F01 used

this motif to interact with different parts of the HA stem (**Figure 4D**). For the FGL motif in AG11-

2F01, V_H F100 inserted into a hydrophobic pocket in the HA stem centering at HA2 I48,

whereas V_H L100b inserted into a lower pocket centering at HA2 W21. By contrast, this lower

pocket was occupied by the V_H F100a of the FGV motif in 16.ND.92. The paratope of 16.ND.92

also involved IGHD3-3-encoded V_H V100, I100e, and I100f, allowing its CDR H3 to bind to

upper pockets in the HA stem that were not engaged by AG11-2F01 (**Figure 4D**). Together,

our structural analyses showed that AG11-2F01 and 16.ND.92 formed distinct molecular

interactions with HA stem.

16.ND.92 utilizes IGHD3-3 in a unique manner for binding to HA stem

Previous studies have determined the structures of several HA stem antibodies with an

FG[V/L/I] motif in the CDR H3 that was encoded by reading frame +3 of IGHD3-3, including

MEDI8852, 56.a.09, 54-1G05, 39.29, PN-SIA28, and 429 B01[7,9,33–36]. These six HA stem

antibodies used either IGHV6-1 or IGHV3-30, unlike AG11-2F01 and 16.ND.92, which used

IGHV4-38-2 and IGHV3-74, respectively. Nevertheless, the CDR H3 conformation of AG11-

2F01 resembled that of MEDI8852, 56.a.09, 54-1G05, 39.29, PN-SIA28, and 429 B01 (**Figure**

4E). Moreover, the IGHD3-3-encoded FG[V/L/I] motifs of these seven antibodies bound to the

HA stem in a similar fashion (**Figure 4E**). In comparison, the CDR H3 conformation of

16.ND.92 was different due to more extensive involvement of IGHD3-3 in binding (**Figure 4D-**

E).

The unique usage of IGHD3-3 for binding enables 16.ND.92 V_H to interact with the HA stem

exclusively through CDR H3, whereas the V_H paratopes of other IGHD3-3 HA stem antibodies

involved non-CDR H3 regions (**Figure 4F and Table S5**). Similarly, IGHD3-3 accounted for

98.6% of the buried surface area of the 16.ND.92 V_H paratope, but 38% to 63% of the V_H

paratopes of other IGHD3-3 HA stem antibodies (**Figure 4G and Table S5**). These

observations not only substantiated that reading frame +3 of IGHD3-3 was a recurring

sequence feature of HA stem antibodies, but also demonstrated that it could pair with diverse

IGHV genes and interact with HA stem via different binding modes.

AG11-2F01 and 16.ND.92 are neutralizing antibodies with *in vivo* protection activity

Given the different binding modes of AG11-2F01 and 16.ND.92, we further aimed to compare

their binding breath, *in vitro* neutralization, and *in vivo* protection activity. ELISA showed that

both AG11-2F01 and 16.ND.92 bound to all H1 and H5 HAs tested (**Figure 5A and Figure**

S6), Microneutralization assay against six H1N1 strains further revealed their neutralizing

activity (**Figure 5B**). Both AG11-2F01 and 16.ND.92 also protected mice against a lethal

challenge of H1N1 A/Puerto Rico/8/1934 (PR8), based on the weight loss profiles (**Figure 5C-**

D), survival analyses (**Figure 5E-F**), and lung viral titers at day 3 post-infection (**Figure 5G-**

H). Nonetheless, our results indicated that the *in vivo* therapeutic protection activity of

16.ND.92 was stronger than AG11-2F01. While only 20% (1/5) of the mice therapeutically

treated with AG11-2F01 survived (**Figure 5E**), 80% (4/5) of the mice therapeutically treated

with 16.ND.92 survived (**Figure 5F**). Additionally, at day 3 post-infection, lung viral titers of

mice therapeutically treated with 16.ND.92 were ~15-fold lower than those treated with AG11-

2F01 (**Figure 5G-H**). Notably, 16.ND.92 had comparable, if not weaker, *in vitro* neutralizing

activity than AG11-2F01 against PR8 (**Figure 5B**). Consequently, the stronger *in vivo*

protection activity of 16.ND.92 against PR8 may at least be partly attributed to its more

downward approaching angle to the HA stem (**Figure 4A**), which could help position the Fc

region closer to effector cells.

DISCUSSION

Antibody discovery has led to significant advancements on many fronts, including antibody-

based therapeutics as well as vaccine designs[11,37–40]. Discovery of natively paired antibody

sequences has been hugely accelerated by scBCR-seq in the past few years[1]. However, going

from sequence information to specificity characterization remains a major bottleneck for

antibody discovery. In this study, we established oPool[+] display, an experimental platform that

allows specificity characterization of antibodies with defined sequences in a highly parallel

fashion. Importantly, oPool[+] display was more cost-efficient (~$30 per antibody) and faster

(~48-72 hours) than the conventional methods that require cloning and recombinant

expression of individual antibodies (~$175-275 per antibody, weeks to months) (**Table S6**). As

a proof-of-concept, we applied oPool[+] display to identify multiple HA stem antibodies from

hundreds of HA antibodies that were left uncharacterized in the literature. Structural analysis

of one of the identified HA stem antibodies, 16.ND.92, further revealed the importance of

IGHD3-3 in targeting the HA stem. Overall, these results provide not only a proof-of-concept

for oPool[+] display, but also molecular insights into our antibody responses against influenza

virus.

Previous studies have shown that IGHV6-1 and IGHV3-30 HA stem antibodies often utilize

IGHD3-3-encoded FG[V/L/I] motif for binding to HA stem[7,9,33–36]. As demonstrated by our work

here, IGHD3-3-encoded FG[V/L/I] motif can also pair with other IGHV genes to target HA stem,

substantiating that it is an IGHV-independent recurring sequence feature of HA stem

antibodies. Our results further revealed that IGHD3-3 can engage the HA stem via different

binding modes. These observations are comparable to those of IGHD3-9, which is utilized by

HA stem antibodies with various IGHV genes and can bind to HA stem in two different reading

frames[14]. Similarly, recent studies have identified IGHD3-22 as an IGHV-independent

recurring sequence feature of antibodies that target a conserved site on SARS-CoV-2

spike[41,42]. Although antibody sequence analysis typically focuses on IGHV genes, the

contribution of IGHD genes to antibody responses should not be overlooked since emerging

evidence suggests that it might be more important than previously thought.

A key feature of oPool[+] display is its relatively simple protocol. A previous study has shown

that the throughput for screening antibodies with defined sequences can be increased by

using liquid handlers to express individual antibodies one by one[43]. In comparison, oPool[+]

display uses a near one-pot strategy for antibody library synthesis and screening, which has

higher potential for scaling. It also only requires standard benchtop equipment commonly

found in a regular molecular biology lab. Therefore, oPool[+] display can be easily integrated

into antibody discovery pipelines that rely on scBCR-seq[1]. After the natively paired antibody

sequences are obtained from scBCR-seq, oPool[+] display can be applied to validate and

characterize the specificity of a large panel of antibody candidates. Thus, this integration can

streamline the existing antibody discovery pipelines. Although this study focused on influenza

HA stem as a proof-of-concept, oPool[+] display can be generalized to any antigens of interest

as long as they can be recombinantly purified. Furthermore, the capability of constructing

custom-made antibody libraries means that oPool[+] display has the potential to benefit the

development of machine learning models for antibody engineering, specificity prediction and

de novo design in the future[15,44,45]. We envision that prediction results from these models can

be rapidly validated by oPool[+] display, which will in turn facilitate iterative refinement of the

models.

We acknowledge that oPool[+] display has some limitations. First, oPool[+] display requires

antibodies to be presented as scFv, which may lose functionality compared to its Fab

counterpart[46–48]. Second, antibodies with a fast off-rate may result in false negatives in mRNA

display selection, since it depends on monovalent binding. Nonetheless, this caveat can be

addressed by replacing mRNA display with other protein display technologies that support

multivalent binding, such as yeast display and phage display[49,50]. As the length of oligo pool

synthesis continues to improve, the cost and complexity of oPool[+] display will further decrease.

Besides, by including more antibodies in a given oligo pool, which is feasible without

compromising the assembly accuracy as demonstrated in this study, we can increase the

throughput of the antibody library construction. Moreover, the scale of antibody library

construction can potentially be increased by leveraging multiplexed gene synthesis

strategies[26,51]. Overall, we believe that oPool[+] display represents a starting point for the future

development of high-throughput approaches to screen antibodies with defined sequences.

MATERIALS AND METHODS

Selection of HA antibodies for paired antibody library synthesis

Members of the natively paired antibody library were selected from a previously curated

dataset containing 5,561 human monoclonal antibodies to influenza HA from 60 research

publications and three patents[15]. Filters were applied to exclude antibodies that 1) had

incomplete sequence information, 2) utilized germline genes that were regarded as recurring

sequence features of HA stem antibodies, namely IGHV1-69, IGHV6-1, IGHV1-18 and

IGHD3-9[7,12–14], and 3) were members of known HA stem antibody clonotypes. This resulted in

292 antibody sequences from 7 publications[7,16–18,23,52,53]. Three HA stem antibodies, namely

31.a.55, AG11-2F01, and 042-100809_2F04, as well as 30 HA head antibodies were randomly

selected as positive and negative controls[6,7,16–25]. Of note, both AG11-2F01 and AG2-G02

were not previously labeled as an HA stem antibody in the curated dataset[15]. Through

literature search, AG11-2F01 was found to compete with CR9114[10], which is an HA stem

antibody, for binding to H1[16], while 042-100809-2F04 was found to bind to group 2 HA stem

domain[54].

Computational design of the oligo pool sequences for assembly

A summary of the computational design pipeline is described below and summarized in **Figure**

S1.

Sequence preparation

Selected antibody sequences were first annotated using abYsis[55]. Any missing nucleotides at

the 5' and/or 3' ends were then filled in using the sequences from the IGHV and IGHJ genes

that had the highest identity with the given antibody.

Codon randomization and pool assignment of the antibodies

To decrease undesired assembly during antibody library synthesis, codon randomization of

each selected antibody sequence was first performed to reduce nucleotide sequence similarity

among different antibodies. For a given amino acid, codon usages <15% in *Escherichia coli*

were removed from consideration to prevent low translation efficiency during RNA display. For

325 antibodies, total of 2 million randomized sequences were generated to maximize

downstream sequence differentiation. All antibody sequences were then split into 8 segments,

followed by clustering using CD-HIT[56]. The clusters were generated using the criterion of 70%

sequence identity (at least 30% differences between each cluster). The antibody sequences

were then reconstructed by selecting necessary segments from different clusters, followed by

the deletion of the used clusters. Such reconstruction was repeated until a complete pool (total

of 25 antibody sequences) had been reassembled. The deleted clusters were then added

back with the reassembled antibody sequences removed. The process was then iterated to

generate the remaining pools.

Selection of overlap regions and generation of the final oligo pools

Upon the pool assignment of antibodies, the ideal overlap regions were searched over CDR

L3, H1, and H3 of each antibody sequence. Six 30-nucleotide long sequences were extracted

from each region through frame shifting, then aligned to the complete antibody sequences in

the corresponding pool using BLAST+[57]. For each antibody, the overlap sequences that are

least similar to other antibodies in the pool were selected. The antibody sequences were then

split at the overlap region, followed by the addition of universal adaptor regions at fragments

encoding the N-terminal and C-terminal of the antibodies, leading to the generation of the final

oligo pools.

Overlap PCR assembly of the natively paired antibody library

A total of 13 oligo pools were synthesized (Integrated DNA Technologies). The lengths of

oligos ranged from around 180 to 330 nucleotides. Each oligo pool contained 100 oligos

resuspended in 200 µL water. An assembly PCR was set up for each oligo pool using 1,600

ng of oligos as input. The assembly was performed using KAPA HiFi HotStart ReadyMix

(Roche) and a Mastercycler nexus GX2 (Eppendorf). PCR was set up in the absence of

primers. From cycles 1-40, PCR was performed with minimal ramp rate (0.1°C/s) in between

the denaturing (98°C, 20 s) and annealing steps (62°C, 15 s) to reduce erroneous annealing

events. After cycle 40, a universal forward primer 5'-TTC <u>TAA TAC GAC TCA CTA TAG</u> GGA

CAA TTA CTA AAG GAG TAT CC-3' and a universal reverse primer 5'-GGA GCC GCT ACC

<u>CTT ATC GTC GTC ATC CTT GTA ATC</u> GGA TCC T-3' were added to the PCR. The underlined

region in the forward primer sequence is the T7 promoter, whereas the underlined region in

the reverse primer sequence encodes a FLAG tag. Subsequently, another 15 cycles of PCR

were performed to amplify the assembled product. The final PCR product was purified using

a Monarch Gel Extraction Kit (New England Biolabs). Two replicates of the assembly were

performed.

Preparation of the biotinylated H1 stem and H3 stem constructs

The H1 stem (mini-HA #4900)[31] and H3 stem (H3ssF)[32] constructs were cloned into a

customized baculovirus transfer vector. Both constructs contained a N-terminal gp67 signal

peptide at the N-terminus as well as a BirA biotinylation site, a thrombin cleavage site, a

trimerization domain and a $6 \times$His-tag at the C-terminus. Recombinant bacmid DNA that

carried the H1 stem construct or H3 stem construct was generated using the Bac-to-Bac

system (Thermo Fisher Scientific) per manufacturer's instructions. Baculovirus was generated

by transfecting the purified bacmid DNA into adherent Sf9 cells using Cellfectin reagent

(Thermo Fisher Scientific) per manufacturer's instructions. The baculovirus was further

amplified by passaging in adherent Sf9 cells at a multiplicity of infection (MOI) of 1.

Recombinant H1 stem and H3 stem constructs were then expressed using 1L of suspension

Sf9 cells at an MOI of 1. At day 3 post-infection, Sf9 cells were pelleted by centrifugation at

4,000 $\times$g for 25 min. Soluble recombinant H1 stem and H3 stem constructs were purified from

the supernatant by affinity chromatography using Ni Sepharose excel resin (Cytiva) and then

size exclusion chromatography using a HiLoad 16/100 Superdex 200 prep grade column

(Cytiva) in 20 mM Tris-HCl at pH 8.0, and 100 mM NaCl. The purified protein was concentrated

by an Amicon spin filter (Millipore Sigma) and filtered by a 0.22 mm centrifuge Tube Filter

(Costar). The purified H1 stem and H3 stem constructs were then biotinylated using a Biotin-

Protein Ligase-BIRA kit (Avidity) according to the manufacturer's instructions. The biotinylated

proteins were then purified again through size exclusion chromatography as described above.

The A280 absorbance values were measured using a Nanodrop One (Thermo Fisher

Scientific) to quantify the protein concentration.

Antibody screening using mRNA display

The mRNA display was performed based on the protocols provided by previous studies[29,30,58] with slight modifications.

Generation of the puromycin-conjugated mRNA templates

The DNA library was first transcribed by a MEGAscript T7 Transcription Kit (Thermo Fisher Scientific) and purified by a MEGAclear Transcription Clean-Up Kit (Thermo Fisher Scientific) according to manufacturer's instructions. Ligation was performed using 1 nmol of the mRNA product, 1.1 nmol of the splint oligo (5'-TTT TTT TTT TTT GGA GCC GCT ACC-3'), and 1.2 nmol of the puromycin linker (5'-/5Phos/-d(A)21-(C$_9$)3-d(ACC)-puromycin-3') by the T4 DNA ligase (New England Biolabs) in a 100 µL reaction for 1 hour at room temperature, followed by Lambda exonuclease (New England Biolabs) digestion for 30 mins at 37°C. The puromycin-conjugated mRNA product was purified using a Dynabeads mRNA DIRECT Purification Kit (Thermo Fisher Scientific), aliquoted, and stored at -20°C until used.

Preparation of the mRNA-scFv fusion library

The puromycin-conjugated mRNA templates were translated using a PURExpress In Vitro Protein Synthesis Kit (New England Biolabs) with the addition of PURExpress Disulfide Bond Enhancer (New England Biolabs) for 1 hour at 37°C. The reaction was then incubated with 500 mM KCl and 60 mM MgCl$_2$ for at least 30 mins at room temperature to promote fusion between the translated scFv and puromycin. EDTA was then added to dissociate ribosomes. The full-length mRNA-scFv product was then purified using Anti-FLAG M2 Magnetic Beads (Millipore Sigma) followed by elution using 3×FLAG peptides (GlpBio). Subsequently, the purified mRNA-scFv product was reverse transcribed using SuperScript IV reverse transcriptase (Thermo Fisher Scientific). The cDNA/mRNA-scFv product was referred as the "pre-selection library".

Preparation of the magnetic beads coated with H1 stem or H3 stem

Biotinylated H1 stem construct or H3 stem construct was coated onto the Dynabeads M280-

straptavidin (Thermo Fisher Scientific) according to the manufacturer's instruction. Briefly, 150

pmol of biotinylated proteins were incubated with 50 µL of the beads for 30 mins to 1 hour at

room temperature with gentle rotation. The beads were then washed with TBST (20 mM Tris-

HCl at pH 7.5, 100mM NaCl, and 0.025% Tween-20) five times using the DynaMag-2 magnetic

holder (Thermo Fisher Scientific) and then resuspended to the original volume.

Antibody selection against H1 stem or H3 stem

Selection of antibodies against H1 stem construct and H3 stem construct were carried out in

parallel. Briefly, the pre-selection library was mixed with 25 µL of beads coated with H1 stem

construct or H3 stem construct and incubated for 1 hour at room temperature with gentle

rotation. After incubation, the beads were washed thrice with 400 µL TBST. The beads were

then resuspended in water. These samples were referred as the "post-selection libraries".

Next-generation sequencing of the scFv library

The pre-selection libraries, post-selection libraries, and the small pool assemblies selected for

quality control were amplified using PrimeSTAR Max DNA Polymerase (Takara Bio) per

manufacturer's instruction with the following primers (5'-GTA AAA CGA CGG CCA GTT TCA

GGG GAC AAT TAC TAA AGG AGT ATC C-3' and 5'- CAG GAA ACA GCT ATG ACC CAC

TCG TCA TCC TTG TAA TCG GAT CCT CCG GA-3'. The PCR product was purified using a

Monarch Gel Extraction Kit (New England Biolabs). A second round of PCR was carried out

to add the adapter sequence and index to the amplicons (**Table S7**). The final PCR products

were sequenced on one SMRTcell 8M on a PacBio Revio system using the CCS sequencing

mode and a 30-hour movie time.

Analysis of next-generation sequencing data

Circular consensus sequences (CCSs) were generated from the raw subreads using

SMRTLink v13.0, setting the parameters to require 99.9% accuracy and a minimum of 3

passes. CSSs in FASTQ format were parsed using the SeqIO module in BioPython[59] and

filtered based on the base calling quality score, where any read with more than five nucleotides

of phred quality score <40 were removed. The adapter sequences were then identified on

each read and trimmed from the scFv sequences. Reads that did not have the complete

adapter sequences were also removed. The filtered reads were then aligned to the reference

scFv sequences and classified into three categories: 1) natively paired scFvs with no mutation,

2) natively paired scFvs with mutation, 3) others (non-natively paired scFvs and incomplete

assemblies). Only the reads encoding natively paired scFvs with no mutation were used for

downstream analysis. Frequency (*F*) of a scFv *i* of a given sample *s* was computed for each

replicate as follows:

439
$$F_{i,s} = \frac{readcount_{i,s}+1}{\sum_s(readcount_{i,s}+1)} \quad (1)$$

440   A pseudocount of 1 was added to each mutant to avoid division by zero in subsequent steps.

441   We then calculate the enrichment (*E*) of a scFv *i* of a given replicate *k* after the mRNA display

442   selection as follows:

443
$$E_{i,k} = \frac{F_{post-selection,i,k}}{F_{pre-selection,i,k}} \quad (2)$$

Expression and purification of CR9114-FLAG scFv

CR9114 scFv nucleotide sequence with a pelB secretion peptide at the N-terminal and a FLAG

(DYKDDDK) tag followed by a stop codon at the C-terminal were synthesized (Integrated DNA

Technologies) and ligated into a pET-28a plasmid vector backbone using NEBuilder HiFi DNA

Assembly Master Mix (New England Biolabs). The ligated product was then transformed into

DH5a competent cells (Thermo Fisher Scientific), and the plasmids were extracted using a

QIAprep Spin Miniprep Kit (Qiagen). The plasmids were then transformed into BL21(DE3)

competent cells (Thermo Fisher Scientific), followed by overnight growth at 37°C, 225 RPM

shaking. 100mL LB culture (Fisher Scientific) were inoculated at 1:500 ratio using the

overnight culture and shaked at 37°C, 225 RPM. Once the OD_{600} reached approximately 0.6,

IPTG (1mM final concentration) was added to induce protein expression for 16hrs at 30°C,

200 RPM.

The overnight culture was centrifuged at 4,500 ×g, 4°C for 1 hour to remove the supernatant.

The pellet was resuspended using 2 mL of ice-cold 1× TES buffer (200 mM Tris-HCl at pH 8.0,

0.65 mM EDTA, and 0.5 M sucrose). The resuspended mixture was incubated at 4°C for 2

hours with gentle shaking. 5 mL of ice-cold 0.25× TES buffer was then added, followed by

incubation at 4°C overnight. The mixture was centrifuged at 4,500 ×g, 4°C for 1 hour to

remove the pellet. The supernatant was subsequently clarified using a polyethersulfone

membrane filter with a 0.22 mm pore size (Millipore) and purified using ANTI-FLAG M2 Affinity

Gel (Millipore-Sigma) per manufacturer's instructions. The A280 was measured using the

Nanodrop One (Thermo Fisher) to calculate the sample concentration.

Expression, purification, and quantitation of selected scFvs for validation

Nucleotide sequences of selected scFvs with a T7 promoter at the N-terminal and a FLAG

(DYKDDDK) tag followed by a stop codon at the C-terminal were synthesized (Integrated DNA

Technologies) and amplified by PCR using Prime STAR Max DNA polymerase (Takara Bio).

The PCR product was purified using a PureLink PCR purification kit (Thermo Fisher Scientific)

and used as the template for *in vitro* translation using a PURExpress In Vitro Protein Synthesis

Kit (New England Biolabs) with the addition of PURExpress Disulfide Bond Enhancer (New

England Biolabs). The translated scFvs were then reverse purified using Pierce High-Capacity

Ni-IMAC magnetic beads (Thermo Fisher Scientific) per manufacturer's instructions to remove

all His-tagged translation kit components.

To measure the concentration of the purified scFv, quantitation assays were performed by

biolayer interferometry (BLI) using an Octet Red instrument (ForteBio). Briefly, rat anti-FLAG

tag monoclonal antibody (L5) (Thermo Fisher Scientific) at 5 µg/mL in 1× kinetics buffer (1×

PBS at pH 7.4, and 0.002% v/v Tween 20) were loaded onto ProG biosensors (Sartorius),

then incubated with the 40× diluted scFv sample (5 µL of sample added to 195 µL of 1×

kinetics buffer). The standard curve was generated via 2-fold serial dilutions using CR9114-

FLAG scFv. The assay consisted of five steps: (1) baseline: 60 s with $1\times$ kinetics buffer; (2)

antibody capture: 180 s with rat anti-FLAG antibody; (3) baseline: 60 s with $1\times$ kinetics buffer;

(4) binding rate measurement: 120 s with standard and scFv samples; and (5) regeneration:

5s in regeneration buffer (0.1 M Glycine at pH 3.0) followed by 5 s in neutralization buffer (1M

Tris-HCl at pH 7.5), repeated 3 times. The data were analyzed using Octet analysis software

9.0, where the first 30 s of the binding rate measurement were used for final quantitation.

Expression and purification of Fabs and IgGs

Heavy and light chains of the antibodies were cloned into a phCMV3 vector with a mouse

immunoglobulin kappa signal peptide in human IgG1 Fc or Fab format. Plasmids encoding the

heavy and light chains of antibodies were transfected into Expi293F cells using an

ExpiFectamine 293 transfection kit (Gibco) in a 2:1 mass ratio for IgG or a 1:1 mass ratio for

Fab following the manufacturer's protocol. Supernatant was harvested at 6 days post-

transfection and centrifuged at 4,000 $\times$g for 30 mins at 4°C to remove cells and debris. The

supernatant was subsequently clarified using a polyethersulfone membrane filter with a 0.22

mm pore size (Millipore). Antibodies were first purified by CaptureSelect CH1-XL beads

(Thermo Fisher Scientific). Then, the antibodies were further purified by size exclusion

chromatography using a HiLoad 16/100 Superdex 200 prep grade column (Cytiva) in $1\times$ PBS.

The A280 were measured using the Nanodrop One (Thermo Fisher) to calculate the sample

concentration. Antibodies were stored at 4°C until used.

Measurement of binding kinetics using biolayer interferometry

Fab K_D measurement

The binding assay was performed by biolayer interferometry (BLI) using an Octet Red

instrument (ForteBio). Briefly, biotinylated H1 or H3 stem construct at 0.5 µM in $1\times$ kinetics

buffer ($1\times$ PBS at pH 7.4, and 0.002% v/v Tween 20) were loaded onto SA biosensors

(Sartorius) and incubated with 33 nM, 100 nM, and 300 nM of purified Fabs. The assay

consisted of five steps: (1) baseline: 60 s with 1× kinetics buffer; (2) loading: 120 s with

biotinylated HA stem domains; (3) baseline: 60 s with 1× kinetics buffer; (4) association: 120

s with Fab samples; and (5) dissociation: 120 s with 1× kinetics buffer. For estimating the

exact K_D, a 1:1 binding model was used.

scFv K_D measurement

The binding assay was performed by biolayer interferometry (BLI) using an Octet Red

instrument (ForteBio). Briefly, biotinylated H1 or H3 stem construct at 0.5 µM in 1× kinetics

buffer (1× PBS at pH 7.4, and 0.002% v/v Tween 20) were loaded onto SA biosensors

(Sartorius) and incubated with 9× dilution (20 µL of sample added to 160uL of 1× kinetics

buffer) and 18× dilution (10 µL of sample added to 170 µL of 1× kinetics buffer) of the purified

scFv sample. The assay consisted of five steps: (1) baseline: 60 s with 1× kinetics buffer; (2)

loading: 180 s with biotinylated HA stem domains; (3) baseline: 60 s with 1× kinetics buffer;

(4) association: 120 s with Fab samples; and (5) dissociation: 120 s with 1× kinetics buffer.

For estimating the exact K_D, a 1:1 binding model was used.

Expression and purification of HA ectodomains

The HA ectodomains of H1N1 A/Puerto Rico/8/1934, H1N1 A/Beijing/262/1995, H1N1

A/Solomon Island/3/2006 (SI/06), H1N1 A/Michigan/45/2015, H1N1 A/Brisbane/02/2018 were

cloned, expressed, and purified as mentioned above for the H1 and H3 stem constructs. The

HA ectodomains of H1N1 A/California/04/2009 (NR-15749), H5N1 A/bald eagle/Florida/W22-

134-OP/2022 (NR-59476), H5N2 A/snow goose/Missouri/CC15-84A/2015 (NR-50651), and

H5N8 A/northern pintail/WA/40964/2014 (NR-50174) were obtained from BEI Resources

(https://www.beiresources.org/). The HA ectodomains of H1N1 A/USSR/90/1977 and H1N1

A/Taiwan/01/1986 were purchased from SinoBiological.

Cryo-EM sample preparation, data collection, and data processing

AG11-2F01 Fab in complex with SI/06 HA

The AG11-2F01 Fab was incubated with SI/06 HA on ice overnight followed by size exclusion

chromatography. The peak fraction of the Fab-HA complex was concentrated to around 1

mg/mL for cryo-EM sample preparation. Cryo-EM grids were prepared using a Vitrobot Mark

IV (Thermo Fisher Scientific). 3.5 µL of the sample was applied to a 300-mesh Quantifoil

R1.2/1.3 Cu grid pretreated with glow-discharge. Excess liquid was blotted away using filter

paper with blotting force -5 and blotting time 3 s. The grid was then flash frozen in liquid ethane.

Movies were then collected on a Titan Krios microscope equipped with Gatan BioQuantum K3

imaging filter and camera (Thermo Fisher Scientific). Images were recorded at 130,000×

magnification, corresponding to a pixel size of 0.33 Å/pixel at super-resolution mode of the

camera. A defocus range of -0.8 µm to -1.5 µm was set. A total dose of 50 e$^-$/Å^2 of each

exposure was fractionated into 50 frames. Both untilted and 30-degree-tilted data were

collected and combined to alleviate the preferred orientation problem of the sample.

CryoSPRAC[60] was used to process the cryo-EM data. For model building, ABodyBuilder[61] was

used to generate an initial model for AG11-2F01 Fab. This model, together with the model of

SI/06 HA (PDB 6FYT)[62], was fitted into the cryo-EM density map using UCSF Chimera[63]. The

model was manually adjusted in Coot[64] and refined with Phenix real-space refinement

program[65]. This process was iterated for several cycles until no significant improvement of the

model was observed.

16.ND.92 Fab in complex with SI/06 HA

The 16.ND.92 Fab was incubated with SI/06 HA and FISW84 Fab, a known HA anchor

antibody[66], on ice overnight followed by size exclusion chromatography. The peak fraction of

the Fab-HA complex was concentrated to around 3 mg/mL for cryo-EM sample preparation.

0.1% (w/v) of n-octyl-ß-D-glucoside was added to reduce orientation bias. Cryo-EM grids were

prepared using a Vitrobot Mark IV (Thermo Fisher Scientific). 3 µL of the sample was applied

to a 400-mesh Quantifoil R1.2/1.3 Cu grid pretreated with glow-discharge. Excess liquid was

blotted away using filter paper with blotting force 0 and blotting time 3 s. The grid was then

flash frozen in liquid ethane. Movies were then collected on a Titan Krios microscope equipped

with Gatan BioQuantum K3 imaging filter and camera (Thermo Fisher Scientific). Images were

recorded at 81,000× magnification, corresponding to a pixel size of 0.53 Å/pixel at super-

resolution mode of the camera. A defocus range of -0.5 μm to -5 μm was set. A total dose of

57.35 e^-/Å^2 of each exposure was fractionated into 40 frames.

CryoSPRAC[60] was used to process the cryo-EM data. DeepEMhancer[67] was used to generate

the sharpened density map for downstream model building. For model building, IgFold[68] was

used to generate an initial model for 16.ND.92 Fab. This model, together with the model of

SI/06 HA (PDB 6FYT)[62], was fitted into the cryo-EM density map using Phenix DockinMap

module[69]. The models were manually adjusted in Coot[64] and refined with Phenix real-space

refinement program[65]. This process was iterated for several cycles until no significant

improvement of the model was observed.

Structural analysis of HA-antibody complexes

Buried surface areas upon binding and paratope residues of AG11-2F01, 16.ND.92,

MEDI8852[9] (PDB 5JW4), 56.a.09[7] (PDB 5K9J), 54-1G05[33] (PDB 6WIZ), PN-SIA28[35] (PDB

8GV5), 39.29[36] (PDB 4KVN), and 429 B01[34] (PDB 6NZ7) were analyzed using PDBePISA[70].

The CDRH3 region and IGHD 3-3 usage of each antibody was annotated using IgBLAST[71].

The molecular interactions of AG11-2F01 and 16.ND.92 in complex with SI/06 HA were

analyzed and visualized using PyMOL (Schrödinger).

Enzyme-linked immunosorbent assay (ELISA)

Nunc MaxiSorp ELISA plates (Thermo Fisher Scientific) were coated overnight at 4°C with

100 μL of recombinant proteins at 1 μg/mL in 1× PBS. The next day, plates were washed

thrice with 1× PBS containing 0.05% Tween 20 and blocked with 200 μl of 5% non-fat milk in

1× PBS for 2 hours at room temperature. Monoclonal antibodies were serially diluted 10-fold

starting from 100 μg/mL, added to the plates, and incubated for 2 hours at 37°C. Plates were

then washed thrice and incubated with horseradish peroxidase (HRP)-conjugated goat anti-

human IgG antibody (Thermo Fisher Scientific) at 1:5,000 dilution for 1 hour at 37°C. After six

washes with 1× PBS containing 0.05% Tween 20, 100 μL of 1-Step TMB ELISA Substrate

Solution (Thermo Fisher Scientific) was added to each well. After incubation for 10 mins, the

reaction was stopped with 50 μL of 2 M H_2SO_4 solution, and absorbance values were

measured at 450 nm using a BioTek Synergy HTX Multimode Reader (Agilent).

Recombinant virus construction and purification

Recombinant viruses with HA and NA segments from the indicated H1N1 strains and six

internal segments from H1N1 A/Puerto Rico/8/1934 (PR8) were obtained from BEI Resources

(https://www.beiresources.org/). Recombinant viruses were rescued using the eight-plasmid

reverse genetics system[72]. Briefly, plasmids encoding the HA segments from H1N1

A/California/07/2009 and H1N1 A/Michigan/45/2015 along with seven plasmids encoding the

other seven segments from PR8 were transfected into a co-culture of HEK293T (human

embryonic kidney) cells and MDCK-SIAT1 (Madin-Darby Canine Kidney) cells at a 6:1 ratio.

Supernatants were injected into 8-10 days old embryonated chicken eggs and incubated at

37°C for 48 hours. Viruses in the allantoic fluid were plaque-purified on MDCK-SIAT1 cells

grown in Dulbecco's Modified Eagles Medium (Gibco) containing 10% fetal bovine serum

(Gibco) and a penicillin-streptomycin mix (100 U/mL penicillin and 100 μg/mL streptomycin,

Gibco). The HA sequence of each virus was confirmed by Sanger sequencing.

Microneutralization assay

For the microneutralization assay, MDCK-SIAT1 cells were seeded in 96-well plates. After

reaching 100% confluency, MDCK-SIAT1 cells were washed once with 1× PBS. Minimal

essential media (Gibco) containing 25 mM HEPES (Gibco) was then added to the cells.

Monoclonal antibodies were serially diluted 10-fold starting from 100 μg/ml and mixed with

100 $TCID_{50}$ (median tissue culture infectious dose) of viruses at equal volume and incubated

at 37°C for 1 hour. Subsequently, the mixture was inoculated into cells and incubated at 37°C

for another hour. Cell supernatants were discarded and replaced with minimal essential media

containing 25 mM HEPES, and 1 µg/mL TPCK-trypsin (Sigma). Plates were incubated at 37°C

for 72 hours, and virus presence was detected by hemagglutination assay to determine the

MN_{50} titers.

Mice

The animal experiments were performed in accordance with protocols approved by UIUC

Institutional Animal Care and Use Committee (IACUC). Six-week-old female BALB/c mice

(Jackson Laboratory) were used for all animal experiments.

Prophylactic and therapeutic protection experiments

Female BALB/c mice at 6 weeks old (n = 5 per group) were anesthetized with isoflurane and

intranasally infected with 5× lethal dose (LD_{50}) of recombinant PR8 virus. Mice were given the

indicated antibody at a dose of 5 mg/kg intraperitoneally at 4 hours before infection

(prophylaxis) or 4 hours after infection (therapeutics). Weight loss was monitored daily for 14

days. The humane endpoint was defined as a weight loss of 25% from initial weight at day 0.

Of note, while our BALB/c mice were not modified to facilitate interaction with human IgG1,

human IgG1 could interact with mouse Fc gamma receptors[73–75].To determine the lung viral

titers at day 3 post-infection, lungs of infected mice were harvested and homogenized in 1 mL

of minimal essential media with 10% bovine serum albumin using a gentleMACS C Tube

(Miltenyi Biotec). Subsequently, virus titers were measured by $TCID_{50}$ assay.

ACKNOWLEDGEMENTS

We thank Anders Olson, Chris Brooke, David Kranz, and Steve Sligar for helpful discussion

and the Roy J. Carver Biotechnology Center at the University of Illinois at Urbana-Champaign

for assistance with PacBio sequencing. We thank Kristen Flatt and the Materials Research

Laboratory Central Research Facilities at the University of Illinois at Urbana-Champaign for

access to cryo-EM instrumentation during the screening of the Fab-HA complexes. We are

grateful to the Cryo-EM Core and the Core Facility for Advanced Research Computing at Case

Western Reserve University for the access of cryo-EM instrumentation and data processing.

We also thank Frank Vago and the Cryo-EM Facility at Purdue University for the access of

cryo-EM instrumentation and data collection. This work is supported by National Institutes of

Health R01 AI167910 (N.C.W.), DP2 AT011966 (N.C.W.), the Searle Scholars Program

(N.C.W.), Howard Hughes Medical Institute Emerging Pathogens Initiative (N.C.W.), UIUC

William T. and Lynn Jackson Graduate Student Fellowship (W.O.O.), and Carl R. Woese

Institute for Genomic Biology Postdoctoral Fellowship (H.L.).

AUTHOR CONTRIBUTIONS

W.O.O. and N.C.W. conceived and designed the study. W.O.O., H.L., W.L., Y.W., K.E.D. and

N.C.W. developed the methodology. W.O.O. and L.T. assembled the library and performed

mRNA display experiment. W.O.O., W.L. and N.C.W. analyzed the PacBio sequencing data.

W.O.O., H.L., R.L., A.B.G. and L.A.R expressed and purified recombinant proteins. W.O.O.,

Z.M., R.L., T.P., X.D., and N.C.W. performed structural analysis of the antibodies. W.O.O., H.L.,

D.C., and M.R.A performed functional characterization of the antibodies. X.D. and N.C.W.

provided resources and support. W.O.O. H.L., W.L. and N.C.W. wrote the paper, and all

authors reviewed and edited the paper.

DECLARATION OF INTERESTS

N.C.W. consults for HeliXon. The authors declare no other competing interests.

DATA AVAILABILITY

Raw sequencing data have been submitted to the NIH Short Read Archive under accession

number: BioProject PRJNA1150188. Cryo-EM density maps and coordinates have been

deposited to EMDB and PDB with accession numbers: EMD-46727 and EMD-45930; PDB

9DBX and 9CU7.

CODE AVAILABILITY

Custom scripts for all analyses have been deposited to:

https://github.com/nicwulab/oPool_display.

MAIN FIGURES

Figure 1. Curation and synthesis of the natively paired HA antibody library. (A) The

overall breakdown of the HA antibody library. **(B)** Design of oligos for scFv assembly. Each

given scFv construct contains a T7 promoter and a start codon at the N-terminal as well as a

FLAG tag at the C-terminal. The scFv sequences were then split into 4 fragments at the

selected CDR regions, with overlap between adjacent fragments. Through an overlap PCR,

oligos of the same construct would preferably anneal to each other, ensuring the assembly of

natively paired scFvs. **(C)** Synthesis of the natively paired HA antibody library. A total of 13

separate oligo pools (25 antibodies per pool) were synthesized, followed by the assembly,

which consisted of a two-stage PCR (**see Materials and Methods**). **(D)** Comparison between

the unassembled and assembled oligo pools were analyzed by agarose gel electrophoresis.

"U": unassembled oligo pool. "Rep1": replicate 1. "Rep2": replicate 2. The red arrow indicates

the target size (800-900 bp) for full length scFvs. **(E-F)** The composition **(E)** and reproducibility

(F) were assessed. The Pearson correlation coefficient (R) of the occurrence frequencies of

individual scFvs between the two replicates were indicated. Micro-tube icon by Servier

https://smart.servier.com/ is licensed under CC-BY 3.0 Unported, available via Bio Icons.

Figure 2. Rapid screening for HA stem antibodies via mRNA display. (A) Schematic overview of the mRNA display selection. Different colors represent different scFvs. The purple dot represents the puromycin. **(B-C)** Correlation between replicates of mRNA display selection against **(B)** H1 stem and **(C)** H3 stem. The green dots represent the positive controls (known HA stem antibodies), whereas the red dots represent negative controls (known HA head antibodies). The three positive controls in this study are 31.a.55 (both H1 and H3 stem-binding), AG11-2F01 (H1 stem-binding), and 042-100809-2F04 (H3 stem-binding)[7,16]. For each replicate, the average enrichment of negative controls is shown as a dotted line. Pearson correlation coefficient (R) between replicates is indicated. **(D)** The average enrichments against H1 stem and H3 stem are compared. The dotted line represents the cutoff for the identification of HA stem antibody candidates. H1 stem antibody candidates are in lower right quadrant, whereas H3 stem antibody candidates are in the upper left quadrant.

Figure 3. Validation of HA stem antibody candidates. (A) Schematic overview of the validation experiment. H1 and H3 stem antibody candidates in both scFv and Fab formats were validated using biolayer interferometry (BLI). **(B-C)** The dissociation constants (K_D) of antibodies against **(B)** H1 stem and **(C)** H3 stem are shown as heatmaps. The selected antibodies were listed from left to right based on their average enrichments. The enrichment rank of each antibody in the library were shown. Of note, 31.a.55 is a positive control for binding to both H1 stem and H3 stem[7], AG11-2F01 is a positive control for binding to H1 stem[16], and 042-100809_2F04 is a positive control for binding to H3 stem[54].

Figure 4. Structural analysis of AG11-2F01 and 16.ND.92. (A) Cryo-EM structures of AG11-

2F01 and 16.ND.92 in complex with SI/06 HA. HA1 is in light grey. HA2 is in dark grey. Heavy

chain variable domain (V_H) is in orange. Light chain variable domain (V_L) is in pink. **(B)**

Epitopes of AG11-2F01 and 16.ND.92. V_H contacts are in orange. V_L contacts are in pink.

Contacts shared by both V_H and V_L are in purple. **(C)** Interactions between light chain and HA

are shown. H-bonds are represented by black dashed lines. **(D)** Interactions between CDR

729   H3 and HA are shown. **(E)** Overlay of the CDR H3 loops from IGHD3-3 HA-stem antibodies.

730   HA is in surface representation. **(F)** Contributions of CDR H3 (orange), non-CDR H3 V_H

731   (yellow), and V_L (pink) to the paratope buried surface areas (BSA) of the indicated antibodies.

732   **(G)** Contributions of IGHD3-3-encoded residues to V_H paratope BSA of the indicated

733   antibodies.

Figure 5. *in vitro* and *in vivo* protection of AG11-2F01 and 16.ND.92. (A) The binding

activities of AG11-2F01 and 16.ND.92 against recombinant HA proteins from the indicated H1

and H5 strains were measured by ELISA. The EC_{50} values are shown as a heatmap. **(B)** The

neutralization activity of AG11-2F01 and 16.ND.92 against different recombinant H1N1 viruses

was measured by a microneutralization assay. The IC_{50} values are shown as a heatmap. **(A-**

B) Strain names are abbreviated as follows: H1N1 A/Puerto Rico/8/1934 (H1 PR8), H1N1

A/USSR/90/1977 (H1 USSR/77), H1N1 A/Chile/1/1983 (H1 Chile/83), H1N1

A/Taiwan/01/1986 (H1 Taiwan/86), H1N1 A/Beijing/262/1995 (H1 Beijing/95), H1N1

A/Solomon Island/3/2006 (H1 SI/06), H1N1 A/Brisbane/59/2007 (H1 Bris/07), H1N1

A/California/04/2009 (H1 CA/09), H1N1 A/Michigan/45/2015 (H1 Mich/15), H1N1

A/Brisbane/02/2018 (H1 Bris/18), H5N8 A/northern pintail/WA/40964/2014 (H5 Wash/14),

H5N2 A/snow goose/Missouri/CC15-84A/2015 (H5 Missouri/15), and H5N1 A/bald

eagle/Florida/W22-134-OP/2022 (H5 Florida/22). H1 stem and H3 stem represents the HA

stem constructs designed based on H1N1 A/Brisbane/59/2007 HA and H3N2

A/Finland/486/2004 HA, respectively[31,32]. **(C-H)** The *in vivo* protection activity of AG11-2F01

and 16.ND.92 against lethal challenge of PR8 virus was assessed by **(C-D)** weight loss profiles,

(E-F) Kaplan-Meier survival curves, and **(G-H)** lung viral titers at day 3 post-infection. *P* values

were computed by two-tailed student's t-test. ***: $P < 0.001$; **: $P < 0.01$; n.s.: not significant.

SUPPLEMENTAL FIGURES

Figure S1. Curation and design of the natively paired HA antibody library.

Figure S2. Quality assessment of oligo pool-based antibody library assembly.

Figure S3. BLI sensorgrams for antibody validation in Fab format.

Figure S4. BLI sensorgrams for antibody validation in scFv format.

Figure S5. CDR H3 sequence analysis of AG11-2F01 and 16.ND.92.

Figure S6. ELISA titration curves.

SUPPLEMENTAL TABLES

Table S1: Selected HA antibodies.

Table S2: Oligo pool sequences.

Table S3: Enrichment results of oPool[+] display.

Table S4: Cryo-EM data collection, refinement and validation statistics.

Table S5: Buried surface areas upon binding of IGHD3-3 antibodies.

Table S6: Cost and time comparison between traditional approaches for antibody

specificity characterization and oPool[+] display.

Table S7: Sequences of primers used in this study.

REFERENCES

1. Pedrioli, A. & Oxenius, A. Single B cell technologies for monoclonal antibody discovery. *Trends Immunol.* **42**, 1143–1158 (2021).

2. Valldorf, B. *et al.* Antibody display technologies: selecting the cream of the crop. *Biol. Chem.* **403**, 455–477 (2022).

3. Persson, M. A., Caothien, R. H. & Burton, D. R. Generation of diverse high-affinity human monoclonal antibodies by repertoire cloning. *Proc. Natl. Acad. Sci. U. S. A.* **88**, 2432–2436 (1991).

4. Sheets, M. D. *et al.* Efficient construction of a large nonimmune phage antibody library: the production of high-affinity human single-chain antibodies to protein antigens. *Proc. Natl. Acad. Sci. U. S. A.* **95**, 6157–6162 (1998).

5. Kirkpatrick, E., Qiu, X., Wilson, P. C., Bahl, J. & Krammer, F. The influenza virus hemagglutinin head evolves faster than the stalk domain. *Sci. Rep.* **8**, 10432 (2018).

6. Zost, S. J., Wu, N. C., Hensley, S. E. & Wilson, I. A. Immunodominance and antigenic variation of Influenza virus hemagglutinin: implications for design of universal vaccine immunogens. *J. Infect. Dis.* **219**, S38–S45 (2019).

7. Joyce, M. G. *et al.* Vaccine-induced antibodies that neutralize group 1 and group 2 Influenza A viruses. *Cell* **166**, 609–623 (2016).

8. Corti, D. *et al.* A neutralizing antibody selected from plasma cells that binds to group 1 and group 2 Influenza A hemagglutinins. *Science* **333**, 850–856 (2011).

9. Kallewaard, N. L. *et al.* Structure and function analysis of an antibody aecognizing all Influenza A subtypes. *Cell* **166**, 596–608 (2016).

10. Dreyfus, C. *et al.* Highly conserved protective epitopes on Influenza B viruses. *Science* **337**, 1343–1348 (2012).

11. Wu, N. C. & Wilson, I. A. Influenza hemagglutinin structures and antibody recognition. *Cold Spring Harb. Perspect. Med.* **10**, a038778 (2020).

12. Andrews, S. F., Graham, B. S., Mascola, J. R. & McDermott, A. B. Is it possible to develop a "universal" influenza virus vaccine? Immunogenetic considerations underlying B-cell biology in the development of a pan-subtype Influenza A vaccine targeting the

hemagglutinin stem. *Cold Spring Harb. Perspect. Biol.* **10**, a029413 (2018).

13. Pappas, L. *et al.* Rapid development of broadly influenza neutralizing antibodies through

redundant mutations. *Nature* **516**, 418–422 (2014).

14. Wu, N. C. *et al.* Recurring and adaptable binding motifs in broadly neutralizing antibodies

to Influenza virus are encoded on the D3-9 segment of the Ig gene. *Cell Host Microbe* **24**,

569-578.e4 (2018).

15. Wang, Y. *et al.* An explainable language model for antibody specificity prediction using

curated influenza hemagglutinin antibodies. *Immunity* (2024)

doi:10.1016/j.immuni.2024.07.022.

16. Henry, C. *et al.* Influenza virus vaccination elicits poorly adapted B cell responses in elderly

individuals. *Cell Host Microbe* **25**, 357-366.e6 (2019).

17. Andrews, S. F. *et al.* Immune history profoundly affects broadly protective B cell responses

to influenza. *Sci. Transl. Med.* **7**, 316ra192 (2015).

18. Dugan, H. L. *et al.* Preexisting immunity shapes distinct antibody landscapes after

influenza virus infection and vaccination in humans. *Sci. Transl. Med.* **12**, eabd3601 (2020).

19. Cheung, C. S.-F. *et al.* Identification and structure of a multidonor class of head-directed

Influenza-neutralizing antibodies reveal the mechanism for its recurrent elicitation. *Cell*

Rep. **32**, 108088 (2020).

20. Ekiert, D. C. *et al.* Cross-neutralization of influenza A viruses mediated by a single antibody

loop. *Nature* **489**, 526–532 (2012).

21. Henry, C. *et al.* Monoclonal antibody responses after recombinant hemagglutinin vaccine

versus subunit inactivated Influenza Virus vaccine: a comparative study. *J. Virol.* **93**,

e01150-19 (2019).

22. Hu, H. *et al.* A human antibody recognizing a conserved epitope of H5 hemagglutinin

broadly neutralizes highly pathogenic avian influenza H5N1 viruses. *J. Virol.* **86**, 2978–

2989 (2012).

23. Liu, Y. *et al.* Cross-lineage protection by human antibodies binding the influenza B

hemagglutinin. *Nat. Commun.* **10**, 324 (2019).

24. Watanabe, A. *et al.* Antibodies to a conserved Influenza head interface epitope protect by

an IgG subtype-dependent mechanism. *Cell* **177**, 1124-1135.e16 (2019).

25. Lee, J. *et al.* Molecular-level analysis of the serum antibody repertoire in young adults

before and after seasonal influenza vaccination. *Nat. Med.* **22**, 1456–1464 (2016).

26. Plesa, C., Sidore, A. M., Lubock, N. B., Zhang, D. & Kosuri, S. Multiplexed gene synthesis

in emulsions for exploring protein functional landscapes. *Science* **359**, 343–347 (2018).

27. Moravec, Z. *et al.* Discovery of tumor-reactive T cell receptors by massively parallel library

synthesis and screening. *Nat. Biotechnol.* 1–9 (2024) doi:10.1038/s41587-024-02210-6.

28. Roberts, R. W. & Szostak, J. W. RNA-peptide fusions for the in vitro selection of peptides

and proteins. *Proc. Natl. Acad. Sci. U. S. A.* **94**, 12297–12302 (1997).

29. Wu, N. C., Dai, L., Olson, C. A., Lloyd-Smith, J. O. & Sun, R. Adaptation in protein fitness

landscapes is facilitated by indirect paths. *eLife* **5**, e16965 (2016).

30. Tanaka, S. *et al.* Rapid identification of neutralizing antibodies against SARS-CoV-2

variants by mRNA display. *Cell Rep.* **38**, 110348 (2022).

31. Impagliazzo, A. *et al.* A stable trimeric influenza hemagglutinin stem as a broadly protective

immunogen. *Science* **349**, 1301–1306 (2015).

32. Corbett, K. S. *et al.* Design of nanoparticulate group 2 influenza virus hemagglutinin stem

antigens that activate unmutated ancestor B cell receptors of broadly neutralizing antibody

lineages. *mBio* **10**, e02810-18 (2019).

33. Wu, N. C. *et al.* Convergent evolution in breadth of two VH6-1-encoded Influenza antibody

clonotypes from a single donor. *Cell Host Microbe* **28**, 434-444.e4 (2020).

34. Matsuda, K. *et al.* Prolonged evolution of the memory B cell response induced by a

replicating adenovirus-influenza H5 vaccine. *Sci. Immunol.* **4**, eaau2710 (2019).

35. Chen, Y. *et al.* Structural basis for a human broadly neutralizing influenza A hemagglutinin

stem-specific antibody including H17/18 subtypes. *Nat. Commun.* **13**, 7603 (2022).

36. Nakamura, G. *et al.* An in vivo human-plasmablast enrichment technique allows rapid

identification of therapeutic Influenza A antibodies. *Cell Host Microbe* **14**, 93–103 (2013).

37. Lanzavecchia, A., Frühwirth, A., Perez, L. & Corti, D. Antibody-guided vaccine design:

identification of protective epitopes. *Curr. Opin. Immunol.* **41**, 62–67 (2016).

38. Haynes, B. F. *et al.* Strategies for HIV-1 vaccines that induce broadly neutralizing

antibodies. *Nat. Rev. Immunol.* **23**, 142–158 (2023).

39. Toussi, S. S., Hammond, J. L., Gerstenberger, B. S. & Anderson, A. S. Therapeutics for

COVID-19. *Nat. Microbiol.* **8**, 771–786 (2023).

40. Schoeder, C. T. *et al.* Epitope-focused immunogen design based on the ebolavirus

glycoprotein HR2-MPER region. *PLOS Pathog.* **18**, e1010518 (2022).

41. Liu, L. *et al.* An antibody class with a common CDRH3 motif broadly neutralizes

sarbecoviruses. *Sci. Transl. Med.* **14**, eabn6859 (2022).

42. Liu, H. *et al.* Human antibodies to SARS-CoV-2 with a recurring YYDRxG motif retain

binding and neutralization to variants of concern including Omicron. *Commun. Biol.* **5**, 766

(2022).

43. Hunt, A. C. *et al.* A rapid cell-free expression and screening platform for antibody discovery.

Nat. Commun. **14**, 3897 (2023).

44. Joubbi, S. *et al.* Antibody design using deep learning: from sequence and structure design

to affinity maturation. *Brief. Bioinform.* **25**, bbae307 (2024).

45. Bennett, N. R. *et al.* Atomically accurate de novo design of single-domain antibodies.

BioRxiv Prepr. Serv. Biol. 2024.03.14.585103 (2024) doi:10.1101/2024.03.14.585103.

46. Wu, N. C. *et al.* In vitro evolution of an influenza broadly neutralizing antibody is modulated

by hemagglutinin receptor specificity. *Nat. Commun.* **8**, 15371 (2017).

47. Quintero-Hernández, V. *et al.* The change of the scFv into the Fab format improves the

stability and in vivo toxin neutralization capacity of recombinant antibodies. *Mol. Immunol.*

44, 1307–1315 (2007).

48. Kang, T. H. & Seong, B. L. Solubility, stability, and avidity of recombinant antibody

fragments expressed in microorganisms. *Front. Microbiol.* **11**, (2020).

49. Rondot, S., Koch, J., Breitling, F. & Dübel, S. A helper phage to improve single-chain

antibody presentation in phage display. *Nat. Biotechnol.* **19**, 75–78 (2001).

50. Chao, G. *et al.* Isolating and engineering human antibodies using yeast surface display.

Nat. Protoc. **1**, 755–768 (2006).

51. Sidore, A. M., Plesa, C., Samson, J. A., Lubock, N. B. & Kosuri, S. DropSynth 2.0: high-

fidelity multiplexed gene synthesis in emulsions. *Nucleic Acids Res.* **48**, e95 (2020).

52. Huang, K.-Y. A. *et al.* Focused antibody response to influenza linked to antigenic drift. *J.*

Clin. Invest. **125**, 2631–2645 (2015).

53. Li, G.-M. *et al.* Pandemic H1N1 influenza vaccine induces a recall response in humans

that favors broadly cross-reactive memory B cells. *Proc. Natl. Acad. Sci. U. S. A.* **109**,

9047–9052 (2012).

54. Henry Dunand, C. J. *et al.* Preexisting human antibodies neutralize recently emerged

H7N9 influenza strains. *J. Clin. Invest.* **125**, 1255–1268 (2015).

55. Swindells, M. B. *et al.* abYsis: Integrated antibody sequence and structure-management,

analysis, and prediction. *J. Mol. Biol.* **429**, 356–364 (2017).

56. Fu, L., Niu, B., Zhu, Z., Wu, S. & Li, W. CD-HIT: accelerated for clustering the next-

generation sequencing data. *Bioinformatics* **28**, 3150–3152 (2012).

57. Camacho, C. *et al.* BLAST+: architecture and applications. *BMC Bioinformatics* **10**, 421

(2009).

58. Olson, C. A., Wu, N. C. & Sun, R. A comprehensive biophysical description of pairwise

epistasis throughout an entire protein domain. *Curr. Biol.* **24**, 2643–2651 (2014).

59. Cock, P. J. A. *et al.* Biopython: freely available Python tools for computational molecular

biology and bioinformatics. *Bioinformatics* **25**, 1422–1423 (2009).

60. Punjani, A., Rubinstein, J. L., Fleet, D. J. & Brubaker, M. A. cryoSPARC: algorithms for

rapid unsupervised cryo-EM structure determination. *Nat. Methods* **14**, 290–296 (2017).

61. Leem, J., Dunbar, J., Georges, G., Shi, J. & Deane, C. M. ABodyBuilder: Automated

antibody structure prediction with data–driven accuracy estimation. *mAbs* **8**, 1259–1268

(2016).

62. Laursen, N. S. *et al.* Universal protection against influenza infection by a multidomain

antibody to influenza hemagglutinin. *Science* **362**, 598–602 (2018).

63. Pettersen, E. F. *et al.* UCSF Chimera--a visualization system for exploratory research and

analysis. *J. Comput. Chem.* **25**, 1605–1612 (2004).

64. Emsley, P. & Cowtan, K. Coot: model-building tools for molecular graphics. *Acta*
Crystallogr. D Biol. Crystallogr. **60**, 2126–2132 (2004).

65. Afonine, P. V. *et al.* Real-space refinement in PHENIX for cryo-EM and crystallography.
Acta Crystallogr. Sect. Struct. Biol. **74**, 531–544 (2018).

66. Benton, D. J. *et al.* Influenza hemagglutinin membrane anchor. *Proc. Natl. Acad. Sci. U.*
S. A. **115**, 10112–10117 (2018).

67. Sanchez-Garcia, R. *et al.* DeepEMhancer: a deep learning solution for cryo-EM volume
post-processing. *Commun. Biol.* **4**, 874 (2021).

68. Ruffolo, J. A., Chu, L.-S., Mahajan, S. P. & Gray, J. J. Fast, accurate antibody structure
prediction from deep learning on massive set of natural antibodies. *Nat. Commun.* **14**,
2389 (2023).

69. Millán, C., McCoy, A. J., Terwilliger, T. C. & Read, R. J. Likelihood-based docking of models
into cryo-EM maps. *Acta Crystallogr. Sect. Struct. Biol.* **79**, 281–289 (2023).

70. Krissinel, E. & Henrick, K. Inference of macromolecular assemblies from crystalline state.
J. Mol. Biol. **372**, 774–797 (2007).

71. Ye, J., Ma, N., Madden, T. L. & Ostell, J. M. IgBLAST: an immunoglobulin variable domain
sequence analysis tool. *Nucleic Acids Res.* **41**, W34-40 (2013).

72. Neumann, G., Fujii, K., Kino, Y. & Kawaoka, Y. An improved reverse genetics system for
influenza A virus generation and its implications for vaccine production. *Proc. Natl. Acad.*
Sci. U. S. A. **102**, 16825–16829 (2005).

73. Lei, R. *et al.* Leveraging vaccination-induced protective antibodies to define conserved
epitopes on influenza N2 neuraminidase. *Immunity* **56**, 2621-2634.e6 (2023).

74. Overdijk, M. B. *et al.* Crosstalk between human IgG isotypes and murine effector cells. *J.*
Immunol. Baltim. Md 1950 **189**, 3430–3438 (2012).

75. Dekkers, G. *et al.* Affinity of human IgG subclasses to mouse Fc gamma receptors. *mAbs*
9, 767–773 (2017).

A Library breakdown

Stem: 3
Head: 30

325 total
HA abs

Unknown: 292

B scFv construct design

Start codon | CDR L1 | CDR L2 | CDR L3 | Gly-Ser linker | CDR H1 | CDR H2 | CDR H3 | FLAG tag

T7

Variable light chain (V_L)   Variable heavy chain (V_H)

Split to 4 ssDNA fragments

Overlap PCR assembly

scFv with intended V_H-V_L pairing

D Assembled product

Ladder   U   Rep 1   Rep 2

1000 bp
500 bp
300 bp
100 bp

C Parallel assembly of natively paired scFvs

ssDNA (oligo) pool
(25 scFvs per pool)

Low ramp rate
(0.1C/s) annealing

Without
primers

PCR
Extension

PacBio
sequencing

Add universal
primers to amplify

E Composition

Rep 2
Rep 1

0%   25%   50%   75%   100%

■ Native pairing w/ no mutations   ■ Native pairing w/ mutations   ■ Non-native pairing

F Reproducibility

$R = 0.9$

log₁₀ scFv frequency (Rep 2) vs log₁₀ scFv frequency (Rep 1)

A DNA library → *in-vitro* transcription / puromycin ligation → *in-vitro* translation / FLAG purification → reverse transcription → cDNA/mRNA-scFv Library

Beads coated with HA stem — H1 stem or H3 stem

mix → incubate at room temp 1 hour → wash → elute → Post-selection library

PacBio sequencing → Calculate post/pre enrichment → Enrichment data

Antibody	Enrichment
A	0.33
B	3
C	0.5
...	...

B H1 stem screen

log₁₀ enrichment (Rep 2) vs log₁₀ enrichment (Rep 1)

31.a.55, AG11-2F01, *R* = 0.95

C H3 stem screen

log₁₀ enrichment (Rep 2) vs log₁₀ enrichment (Rep 1)

042-100809-2F04, 31.a.55, *R* = 0.93

D H1 vs H3

log₁₀ average enrichment (H3) vs log₁₀ average enrichment (H1)

042-100809-2F04, 31.a.55, AG11-2F01

● Positive control ● Negative control ● Unknown

A

Traditional Fab validation

Gene fragments — V_L | V_H + Expression vector

cloning ~2-3 days
expression ~3-6 days
purification ~1-2 days

Purified Fab

mix with

BLI binding measurement

H1 stem-coated probe or H3 stem-coated probe

Rapid scFv validation

T7 — Start codon | V_L | Gly-Ser linker | V_H | FLAG tag — T7-term

in vitro translation ~2-4 hours
reverse His-tag purification ~0.5-1 hour
BLI quantitation via FLAG tag ~0.5-1 hour

mix with

Purified scFv

B — H1 stem screen validation

Rank:	1	2	3	286	317
Fab K_D					
scFv K_D					

K_D (nM): >10000, 1000, 100, 10, 1, <1

31.a.55, AG11-2F01, 16.ND.92, C3-2.3F02, AG11-3C06

C — H3 stem screen validation

Rank:	1	2	3	4	9	12	216	316
Fab K_D								
scFv K_D								

K_D (nM): >10000, 1000, 100, 10, 1, <1

240-14 IgA 2F02, 56.j.01, 01.ad.01, 31.a.55, 042-100809_2F04, 01.h.02, C3-2.3F02, AG11-3C06

A | Structure overview

AG11-2F01
IGHD3-3 + IGKV1-5

16.ND.92
IGHD3-3 + IGKV1-5

HA1 ☐ HA2 ▦ V_H ▮ V_L ▮

B | Epitope

AG11-2F01 epitope

16.ND.92 epitope

V_H contacts ▮ V_L contacts ▮ Shared contacts ▮

C | Light chain interactions

AG11-2F01

Q42_HA2
Q38_HA2
D19_HA2
V_L W32
V_L S30
V_L Y94

16.ND.92

K32_HA1
N46_HA2
Q42_HA2
V_L S30
V_L W32
V_L N92
V_L Y94
Q38_HA2
D19_HA2

D | CDR H3 interactions

AG11-2F01

V40_HA1
T318_HA1
V52_HA2
V_H F100
T49_HA2
I48_HA2
I45_HA2
W21_HA2
V_H L100b
T41_HA2

16.ND.92

L292_HA1
L42_HA1
V40_HA1
I56_HA2
V_H I100f
V52_HA2
V_H I100e
T318_HA1
I48_HA2
V_H V100
W21_HA2
I45_HA2
V_H F100a
T41_HA2

E | Comparison of CDR H3 conformations

AG11-2F01 ▮
16.ND.92 ▮
MEDI8852 ▮
56.a.09 ▮
54-1G05 ▮
PN-SIA28 ▮
39.29 ▯
429 B01 ▮

F | Paratope buried surface area

CDRH3 ▮ Non-CDRH3 V_H ▮ V_L ▮

Paratope BSA

AG11-2F01, 16.ND.92, MEDI-8852, 56.a.09, 54-1G05, PN-SIA28, 39.29, 429 B01

G | Contribution of IGHD3-3 to V_H paratope

V_H BSA

AG11-2F01, 16.ND.92, MEDI-8852, 56.a.09, 54-1G05, PN-SIA28, 39.29, 429 B01

AG11-2F01 & 16.ND.92 *in vitro* activity

A ELISA — EC$_{50}$ (µg/mL)

AG11-2F01 / 16.ND.92 across: H1 USSR/77, H1 Taiwan/86, H1 Beijing/95, H1 SI/06, H1 CA/09, H1 Mich/15, H1 Bris/18, H5 Wash/14, H5 Missouri/15, H5 Florida/22, H1 stem, H3 stem

B Neutralization — IC$_{50}$ (µg/mL)

AG11-2F01 / 16.ND.92 across: H1 PR8, H1 Chile/83, H1 Beijing/95, H1 Bris/07, CA/09, H1 Mich/15

AG11-2F01 *in vivo* protection

C Weight loss — % of initial weight vs dpi (Protection, Treatment, Control)

E Survival — survival (%) vs dpi (Protection, Treatment, Control)

G Day 3 lung titer — log$_{10}$ TCID$_{50}$/mL (Protection, Treatment, Control); *** , *** , n.s.

16.ND.92 *in vivo* protection

D Weight loss — % of initial weight vs dpi (Protection, Treatment, Control)

F Survival — survival (%) vs dpi (Protection, Treatment, Control)

H Day 3 lung titer — log$_{10}$ TCID$_{50}$/mL (Protection, Treatment, Control); *** , *** , **