High levels of mitotic gene conversion are needed to effectively purge deleterious mutations in asexual organisms

Dominik Kopčak[1,2,3], Matthew Hartfield[1,*]

1) Institute of Ecology and Evolution, The University of Edinburgh, Edinburgh EH9 3FL, UK.
2) Institute of Science and Technology Austria, 3400 Klosterneuburg, Austria.
3) Okinawa Institute of Science and Technology Graduate University, Onna-son, Okinawa, Japan.
* Corresponding author: m.hartfield@ed.ac.uk

Key words: Self-fertilisation, facultative sex, gene conversion, deleterious mutation, dominance, mutation load.

Data availability statement: Simulation code and Conda environments with all used packages, together with all scripts have been archived on GitHub (https://github.com/dominik-kopcak/SelfClonGC).

Author contributions: MH designed the project with input from DK. DK wrote and ran the simulations, and analysed the resulting outputs with feedback from MH. MH developed and analysed the single-locus model. MH and DK wrote and edited the manuscript.

Funding: Matthew Hartfield is supported by a NERC Independent Research Fellowship (NE/R015686/1) and a UKRI Frontier Research Guarantee Grant (EP/X027570/1).

Conflict of interest statement: Nothing to declare.

Acknowledgements. We would like to thank Anders Poulsen Charmouh for advising on how to implement the non-constant s, h model, and members of the Hartfield lab and two anonymous reviewers for providing feedback on the work.

ORCID links:
Dominik Kopčak: https://orcid.org/0009-0005-3758-4390
Matthew Hartfield: https://orcid.org/0000-0001-6941-0647

Abstract

Self-fertilisation and asexual reproduction are both hypothesised to lead to long-term extinction due to inefficient selection acting against deleterious mutations. Self-fertilisation can counter these effects through creating homozygous genotypes and purging deleterious mutations. Although complete asexuality lacks meiotic gene exchange, mitotic gene conversion is a mechanism for creating homozygous regions that could limit deleterious mutation accumulation in a similar manner to self-fertilisation. Here, we compare mutation accumulation in self-fertilising and facultative sexual populations subject to mitotic gene conversion, and quantify the relative efficacy of purging caused by the latter. We first show analytically that this purging is most effective with high levels of asexuality and gene conversion, and when deleterious mutations are recessive. We further show using simulations that, when mitotic gene conversion becomes sufficiently high in obligate asexuals, there is a sudden reduction in the mutation count and a jump in homozygosity, reflecting mutation purging. However, this mechanism is not necessarily as efficient at purging compared to high levels of self-fertilisation, and elevated rates of mitotic gene conversion seem to be needed compared to empirical estimates. Hence, while mitotic gene conversion can remove recessive deleterious mutations, self-fertilisation seems to be a more effective mechanism for doing so.

58 # Teaser Text

59
Asexuality has been argued to be an evolutionary 'dead end', due to a lack of gene

exchange causing inefficient selection acting against deleterious mutations. It has been

proposed that asexuals can counter these negative effects through mitotic gene conversion,

which exposes mutations to selection within individual lineages. Here, we theoretically

investigate how effective this mechanism is. We compare results to those obtained when

individuals reproduce by self-fertilisation, which has similar effects on exposing deleterious

variants. While mitotic gene conversion can be effective in removing recessive deleterious

mutations, high rates are required and it is not always as effective as selfing.

68 Introduction

Mutations are the ultimate source of genetic novelty in all organisms and are essential

for biological evolution (Loewe and Hill 2010). However, the generation of variation is a double-

edged sword; although favourable mutations may emerge that are crucial for long-term

survival, it may also result in a burden of deleterious mutations, which are much more common

than favourable ones (Eyre-Walker and Keightley 2007).

The reproductive system, which determines how offspring are generated in a

population, has a profound impact on the prevalence of deleterious variation. Reproduction

can either be with or without sex. Sex usually involves two phenomena (Hartfield and Keightley

2012): recombination (the exchange of genetic information between homologous DNA

sequences, most commonly as meiotic crossing-over), and fusion of gametes (specialised

cells with halved ploidy and DNA content). In the simplest case when asexuality is equivalent

to clonal reproduction, the genotype of an offspring is identical to that of a parent, except for

de novo mutations. Clonal reproduction encompasses diverse mechanisms, including fission

or budding in unicellular eukaryotes, or budding (e.g., hydra, jellyfishes), fragmentation (e.g.,

corals) and whole-body fission (e.g., flatworms, starfish) in animals, or many forms of

vegetative propagation in plants (De Meeûs et al. 2007). Nevertheless, most eukaryotic

organisms undergo some variant of sexual reproduction, at least occasionally (Speijer et al.

2015). Facultatively sexual organisms reproduce via a mixture of sexual and asexual

reproduction, so that genetic evolution is affected by the frequency of both recombination and

clonal reproduction (Hartfield 2016).

Gametes from which offspring originate may also be derived from the same individual,

leading to self-fertilisation or selfing. The opposite of selfing is panmictic outcrossing, with

anyone in the population having an equal chance of being a mate of anyone else, with self-

mating being rare or even prohibited (if, e.g., self-incompatibility mechanisms are present;

Nettancourt 2001). Cloning and selfing are two types of uniparental reproduction, which forms

a basis of their similarity (e.g., the lack of mate finding) and potential ecological advantages

(e.g., reproductive assurance; Baker 1955, 1967; Theologidis et al. 2014; Mráz and Mrázová

2021).

Despite these apparent similarities, cloning and selfing differ in the manner in which

they affect selection acting against deleterious mutations. Muller's ratchet operates in asexual

populations leading to stochastic mutation accumulation, due to their inability to generate

offspring that have fewer deleterious alleles than those of their ancestors (Muller 1964; Haigh

1978). This mechanism is a form of selection interference, which refers to the effect of strong

genetic linkage on the dynamics of variants under selection (Felsenstein 1974). In completely

clonal populations all sites are in absolute genetic linkage, therefore the presence of multiple

deleterious alleles selectively interferes with one another having a strong impact on mutation

accumulation (Hill and Robertson 1966; Roze and Barton 2006; Hartfield and Keightley 2012).

Recombination breaks this linkage and allows the dynamics of variants to respond to selection

independently, i.e., more efficient removal of deleterious mutations and restoration of the

most-fit genotype. Another important consequence due to the lack of recombination and

gamete fusion in diploid asexuals is the divergence of homologous chromosomes found within

one individual. The two genome copies are always transmitted together across the lineage,

gradually and independently accumulating mutations leading to a high level of heterozygosity;

a process known as 'allelic sequence divergence' or the Meselson effect (Mark Welch and

Meselson 2000; Butlin 2002; Hartfield 2016).

In contrast to cloning, the main genetic consequence of selfing is genome

homogenization. Under complete selfing, parents with heterozygous genotypes *Aa* have a

50% probability of losing heterozygosity, independent of the allele frequencies in the rest of

the population. The probability that heterozygosity is retained decreases exponentially every

generation, rapidly leading to overall genome homogenisation. This homogenisation also

occurs, albeit at a lower rate, under partial selfing (Wright 1951). As a consequence, the

effective recombination rate is reduced since recombination creates variability only when

heterozygosity is present (Nordborg et al. 1996; Nordborg 2000). An additional consequence

of increased homozygosity is the exposure of recessive deleterious alleles to selection, giving

rise to inbreeding depression (Charlesworth and Willis 2009). While this leads to an instant

reduction in the fitness of selfers, the subsequent purging and removal of deleterious

mutations from the population could eventually lead to elevated long-term fitness and

persistence of the selfing population (Glémin 2003; Abu Awad and Billiard 2017). An outcome

of selfing broadly shared with cloning is the removal of neutral variation through extensive

linked selection acting against deleterious mutations, manifesting itself as a reduction in

effective population size (Agrawal and Hartfield 2016).

Both reproductive systems have been called 'dead-end' strategies, meaning that

species using them have an increased extinction probability. In asexual and selfing

populations, this phenomenon has been often ascribed to mutational meltdown, i.e., the

accumulation of deleterious mutations leading to extinction (Lynch et al. 1993, 1995; Olofsson

et al. 2023), but other mechanisms have been proposed for the phylogenetically short-lived

appearance of asexual species (Engelstädter 2008; Janko et al. 2008; Schwander and Crespi

2009). While the 'dead-end' hypothesis has also been invoked for selfing species (Wright et

al. 2013), they can increase their fitness over time through purging deleterious mutations

(Glémin 2003; Abu Awad and Billiard 2017), and this purging could also stop Muller's ratchet

from operating (Charlesworth et al. 1993). Furthermore, there exist ancient asexuals that have

seemingly existed for long periods of time in the absence of sexual reproduction, including

species of bdelloid rotifers (Flot et al. 2013), oribatid mites (Brandt et al. 2021), ostracods

(Tran Van et al. 2021), and trypanosomes (Weir et al. 2016). The persistence of these ancient

asexuals begs the question of whether there are underexplored mechanisms at play that

prevent these species from undergoing excessive accumulation of deleterious variants, similar

to what occurs in self-fertilising species.

Although genome exchange is an intrinsic part of sexual reproduction, gene conversion

is a mechanism of genome exchange that is also available to asexuals in the form of mitotic

recombination (Lee et al. 2009). Gene conversion results in homozygous tracts, akin to the

effect of selfing, and it has been shown to have an important impact on neutral diversity when

sex is rare (Hartfield et al. 2016; Hartfield 2021). Mitotic gene conversion has been argued to

affect deleterious mutation accumulation in asexual species, including the bdelloid rotifer

Adenata vaga (Flot et al. 2013; although see Vakhrusheva et al. 2020 who argue that these

species show evidence of inter-individual gene exchange), the crustacean *Daphnia pulex*

(Tucker et al. 2013) and the planarian worm *Schmidtea mediterranea* (Kershenbaum et al.

2024). Khakhlova and Bock (2006) also show experimental evidence of gene conversion

eliminating deleterious mutations in plastids of tobacco plants, which are uniparentally

inherited and are effectively asexual. Most of these studies argue that this gene conversion is

beneficial for asexual species by removing deleterious mutations (an exception being Tucker

et al. (2013) that argued that mitotic gene conversion reduces fitness by creating homozygous

mutations). However, most of these arguments are verbal, and there exist little theoretical

predictions to determine whether realistic levels of gene conversion can purge deleterious

mutations, and hence maintain high fitnesses in asexual populations, in an analogous manner

to self-fertilisation. One exception is a model in Flot et al. (2013), showing how gene

conversion slows down Muller's ratchet in asexual populations.

The overall goal of this study is to investigate the efficacy of mitotic gene conversion

in facultative sexual and asexual species in purging deleterious mutations, and compare its

effects to that caused by self-fertilisation. We will use two complementary approaches: first,

we develop single-locus analytical models to quantify the frequency of deleterious mutations

under facultative sex and gene conversion, and compare results to equivalent models under

self-fertilisation. We will then use multi-locus simulations to determine how both reproductive

mechanisms influence the genome-wide prevalence of deleterious mutations.

Our principal questions are: (i) Under facultative sex, how does the frequency of

deleterious variants (and hence also the population's mean fitness) scale with the prevalence

of sex and gene conversion, both at individual loci and genome-wide? (ii) What values of

sexual reproduction and gene conversion produce a level of mutation purging similar to a given

level of self-fertilisation? (iii) Can realistic levels of mitotic gene conversion hence be as

effective as self-fertilisation at purging deleterious mutations? The findings of this study will

provide important quantification of a mechanism for maintaining the fitness of facultative

sexuals and asexuals that has been often discussed but seldom investigated theoretically. It

also provides some insight as to which mode of uniparental reproduction might be favoured

by evolution due to its effects on removing deleterious variation.

Single-Locus Mathematical Model

To gain some insight into how mitotic gene conversion affects the purging of

deleterious mutations and how it compares to self-fertilisation, we investigate a single-locus

model of deleterious allele frequencies in a deterministic diploid Wright-Fisher population

subject to either facultative sex with sexual reproduction occurring at frequency σ (i.e., every

generation, a fraction σ of offspring are generated sexually, and the remaining 1–σ are

produced asexually), or self-fertilisation at rate *S* (using similar definitions as for facultative

sex). We consider the frequency of a deleterious allele (denoted *A*) that arose from mutation

from a wild-type allele *a* with probability μ; heterozygous mutant genotypes have fitness 1–*hs*

and homozygotes 1–*s*, for *h* the dominance coefficient and *s* the selection coefficient. Mitotic

gene conversion occurs with frequency γ, which converts heterozygotes to either wild-type or

derived homozygote with equal frequency.

We can track the prevalence of deleterious mutations by creating numerical recursions

of how the frequency is affected by selection, reproduction, mutation and gene conversion in

non-overlapping generations (following the methods of Agrawal and Hartfield 2016, Hartfield

2021; see Supplementary *Mathematica* file for the full recursion equations). Although these

recursions calculate the change in genotype frequencies over time, we can convert these

genotype frequencies to allele frequencies; specifically, if the frequency of the mutant allele is

p_A, then the frequency of the mutant homozygote is $p_A^2 + F p_A (1 - p_A)$ (with similar equations

for the wild-type homozygote), and the heterozygote has frequency $2 p_A (1 - p_A) (1 - F)$.

Here, *F* denotes the reduction in homozygosity compared to Hardy-Weinberg expectations

(Wright 1951), and shows to what extent either gene conversion under facultative sex or self-

fertilisation affects genotype composition by reducing heterozygote frequencies. After making

these transformations, we linearise the equations assuming $\mu, p_A \ll 1$ to obtain steady-state

values of p_A and F (see Otto 2003 for a similar approach).

After these algebraic calculations, we obtain the general equation for the deleterious

allele frequency:

$\hat{p}_A = \dfrac{\mu}{s(F+h-Fh)}$ (1)

This is a known form for mutation-selection balance under non-random mating (Ohta

and Cockerham 1974; Glémin 2003). The inbreeding coefficient F in Equation 1 differs

depending on the form of uniparental reproduction. Under selfing:

$\hat{F}_{self} = \dfrac{((1-hs)\,S\,(1-\gamma)+2\gamma)}{(2-(1-(2-h)s)\,S\,(1-\gamma))}$ (2)

While under facultative sex:

$\hat{F}_{asex} = \dfrac{\gamma}{\gamma+(1-\gamma)(s(1-\sigma)+\sigma)}$ (3)

We can use heuristic arguments to compare F in each case to understand which

reproductive mode has the greatest effect on homozygosity. In the absence of gene

conversion and selection, so we only focus on the effects of selfing, then $\hat{F}_{self} \approx S/(2-S)$ in

line with classic results on how selfing affects homozygosity (e.g., Nordborg and Donnelly

1997). Under facultative sex (Equation 3), positive $\hat{F}_{asex}$ is only formed when gene conversion

is present; interestingly, dominance does not appear in Equation 3 and instead affects allele

frequencies through the mutation-selection balance term (Equation 1). In the absence of

selection and assuming that $\gamma << \sigma$ so the denominator approximates to σ, then $\hat{F}_{asex} \approx \dfrac{\gamma}{\sigma}$ (as

also found by Hartfield 2021). By comparing these two approximations, we can show that the

inbreeding values are equivalent if S = 2/(1+ϕ) for $\phi = \sigma/\gamma$ (Hartfield et al. 2016; Hartfield

2021). As sex is much more common than gene conversion under these assumptions (so ϕ

>> 1) then we see that gene conversion acting under frequent sex is only equivalent to a small

selfing rate. Gene conversion becomes more effective once sex becomes rarer and hence is

more equivalent to elevated selfing rates, but note that this approximation for *S* will break down

once gene conversion and sex act at the same rate.

Figure 1 illustrates these derived equations for different values of the rate of sex and

mitotic gene conversion. If mutations are recessive (Figure 1a) then even small amounts of

gene conversion can reduce the prevalence of deleterious mutations compared to outcrossing

expectations, even for intermediate frequencies of sex. Conversely, the frequency of additive

alleles are close to outcrossing expectations (Figure 1b). These results illustrate that mitotic

gene conversion has the strongest effect on purging deleterious mutations if they are

recessive, with the purging effect magnified as sex becomes rarer. Under self-fertilisation

(Figures 1c, d), the same general results arise, but we note that only a small amount of selfing

is needed to remove recessive deleterious mutations, reflecting the stronger effect of selfing

on homozygosity. Mitotic gene conversion also only has a minor effect on allele frequencies

(note that the contour lines are nearly vertical). Plots of $\hat{F}$ (Figure 2a, b) show that while mitotic

gene conversion under facultative sex leads to elevated homozygosity, the increase is mostly

modest unless sex is rare and deleterious mutations have a weaker selection effect. In

contrast, under self-fertilisation then $\hat{F}$ values tend to be much higher (Figure 2c, d).

To test the accuracy of these approximate analytical solutions (Equations 1-3), we

compared them to exact values obtained from numerical recursions over 50,000 generations,

where the deleterious heterozygous had an initial frequency of 0.001 (Supplementary Figures

S1 and S2, with code in Supplementary *Mathematica* file). We see that they match exact

numerical recursions over a wide range of parameters. One caveat of these models is that

they do not consider drift effects, and hence processes like Muller's ratchet (Muller 1964;

Haigh 1978) that can arise when sex is rare.

253 Multi-Locus Simulation Methods

254 Simulation Outline

While the single-locus model provides insight into how rare sex and gene conversion

affect deleterious mutation frequencies, it does not show how gene conversion affects longer

genome regions (e.g., through longer gene-conversion events purging mutations at multiple

loci, and its effects on the resulting disequilibrium between loci) and hence how effective it is

in purging mutations across larger stretches of the genome. Hence, we next use simulations

to compare how gene conversion leads to mutation removal in asexuals, compared to selfing

species.

Simulations were written using SLiM v4.0.1 (Haller and Messer 2023), and were

implemented in either the Wright-Fisher framework (where the population size is fixed and the

probability of choosing parents are proportional to their fitness; hereafter 'WF model') or in the

non-Wright-Fisher framework (where the population size can change depending on which

parents reproduce or not; hereafter 'non-WF model'). Simulated populations consist of 5,000

diploid individuals, each with a genome size of $L = 25$ Mb that approximates the length of

Arabidopsis thaliana chromosome 1 (The Arabidopsis Genome Initiative 2000). Individuals do

not carry mutations at the start of the simulations. We let populations evolve for 60,000

generations, which is sufficient for the population variance to reach a steady-state

(Supplementary Figure S3). After this time, 50 individuals are sampled and their genotypes

were saved in a VCF file format for downstream analysis. The mutation rate was set to

4×10^{-9}, which is on the same order as that inferred from *Arabidopsis thaliana* (Ossowski et al.

2010). Per-base-position recombination rate was set to 4×10^{-8}, reflecting estimates obtained

from crosses of *Arabidopsis thaliana* (Salome et al. 2012).

Simulated mutations were deleterious and mutation stacking is allowed by default. Two

types of fitness models were investigated. They differ in whether selection and dominance

coefficients were constant or drawn from a distribution (Table 1). In the first type, fixed values

of deleterious mutations and dominance coefficients were chosen. In the second type, the

selection and dominance coefficients were drawn from a distribution based on Spigler et al.

(2017) (see Table 1). Selection coefficient is modelled by a gamma distribution with shape

parameter α and mean $\bar{s}$. The dominance coefficient of a mutation with selection coefficient s

is then drawn from a uniform distribution bounded by 0 and e^{-ks} , where k is a constant (see

Table 1 for exact values). Free parameters in the model, k, $\bar{s}$, and α, were defined so $\bar{s}$ and $\bar{h}$

match with parameters in our main simulations (s = 0.01, h = 0.2) and the shape parameter

was based on Chen *et al.* (2017). The general rationale behind this model is to reflect empirical

observations that strongly deleterious mutations tend to be more recessive (Simmons and

Crow 1977; Agrawal and Whitlock 2011).

Implementing mitotic gene conversion in asexual reproduction

By default, recombination in SLiM (including gene conversion) only acts exclusively

when meiosis is involved. Therefore, the reproduction cycle was manually modified through a

reproduction call-back (i.e., a SLiM function for governing reproduction), so that mitotic gene

conversion in asexuals can be simulated; this can be done only in non-WF model. For

simplicity, the model applies only to a completely asexual population. The development of the

code was facilitated by a post on the SLiM discussion board (Dussert and Haller 2019) and

was extended based on the gene conversion model under SLiM Wright-Fisher simulations

(Haller and Messer 2016).

When an offspring is generated, then the number of gene conversion events that acts

within it is drawn from a Poisson distribution with a mean $L\gamma$ where γ denotes the rate of gene

conversion initiating at a site. For each gene conversion event, the central position is first

drawn uniformly through the chromosome. Then, the tract is extended to the left and right

independently by drawing each length from a geometric distribution with parameter *p:*

$$p = \frac{-\lambda + \sqrt{\lambda}\,\sqrt{8+\lambda}}{2\lambda} \qquad\qquad (2)$$

where λ is the total mean tract length we want to use in simulations. This parameterization is

used to account for the fact that gene conversion tracts with length zero are discarded. This

redefined value of *p* means that the mean tract length is λ after accounting for discarding

these tracts (see Supplementary *Mathematica* notebook for further information). Gene

conversion events that overlap with the chromosome ends are truncated so they stop at the

chromosome edges.

After the gene conversion start- and end-points are drawn, they are checked and

potentially modified as follows: tracts of length 0 are removed; gene conversion tracts may

have the same start or end positions, in which case, the start (or end) of one of them is shifted

to the right (left), so that gene conversion tracts overlap; or two gene conversion tracts overlap,

in which case two interior gene conversion positions are removed so that the resulting gene

conversion tract is a union of the two smaller ones. Since multiple gene conversion tracts can

overlap, this refinement process is repeated until all start and end positions are unique and

alternate after sorting.

Gene conversion can occur on either of the homologous chromosomes with equal

probability and results in the selected part of one chromosome being copied onto the same

position at the other chromosome. Note that for each offspring-production event, all gene

conversion tracts are in the same direction, i.e., material is transferred solely from one strand

to another during each reproduction event. Since the 'initiation' strand is randomly chosen for

each reproduction event, then this effect should not lead to a major bias in results. The mean

length of a gene conversion event is set to 4,000 base pairs; this value accounts for the fact

that empirical studies of mitotic gene conversion tends to find that gene conversion tracts are

bimodally distributed, with a combination of short (on the order of 1kb) and long tracts (on the

order of 10kb) (Yim et al. 2014; Chumki et al. 2016; Jia et al. 2021). Using a mean length of

4,000bp will thus enable a wide distribution of tract lengths to be drawn that cover both these

possibilities. We report simulation results in terms of the 'mean gene conversion rate per site',

which is equal to the initiation rate times the mean tract length. This value is approximately

equivalent to the gene conversion parameter γ in the single-locus model (Frisse et al. 2001).

To enable fair comparison between simulations, further changes to the non-WF model

were made so that it behaves as a standard Wright-Fisher model with a fixed population size

over time, based on section 16.15 in the SLiM Manual (Haller and Messer 2016). Briefly,

survival callback limits the age of an individual to a single generation, and reproduction

callback samples a constant number of parents with probability weighted by their fitness. By

comparing measurements of population fitness (Supplementary Figure S4) and mutation

count (Supplementary Figure S5) between the two simulations, we see that this new routine

yields similar behaviour under low gene conversion rates to the original WF model with

complete asexuality.

100 replicate simulations were run if *s*, *h* were fixed; 20 replicate simulations were run

for WF simulations (i.e., with partial selfing and cloning) where *s*, *h* are drawn from

distributions; and 10 replicate simulations were run for non-WF simulations (i.e., with

asexuality and gene conversion) where *s*, *h* are drawn from distributions. Fewer simulations

replicates were used in the latter two cases due to the much longer runtime of these

simulations.

Mutational Load Measurement

Deleterious mutation load was assessed based on two measurements to reflect those

used in empirical genomic studies (note these terms and measurements are also used by

Laenen et al. (2018) in an analysis of *Arabis alpina*). We calculate (i) the the number of

homozygous mutant genotypes (also called 'recessive load'), which is proportional to the

genetic load expressed by recessive mutations; and (ii) the total number of mutations present

in the population ('additive load'), which is proportional to the load caused by mutations with *h*

= 0.5 (Bertorelle et al. 2022). We also calculated the relative homozygosity for each simulation,

defined as 2*(number of homozygous mutations)/(total number of mutations). This

measurement can be interpreted as the proportion of deleterious mutations that are in

homozygous state, across all sites and individuals. Simulation outputs were analysed using R

(R Core Team 2024).

361　Measuring linkage and identity disequilibrium

Linkage disequilibrium was measured from simulations using similar methods to those

in Hartfield et al. (2018) and Hartfield and Glémin (2024). Briefly, r^2 was measured using

VCFtools (Danecek et al. 2011) after thinning SNPs so measurements were taken between

sites that are 10kb apart and had a minor allele frequency of 5%. LD decay was plotted for

each simulation replicate in bins of 0.5Mb up to 12.5Mb apart, while subsampling the results

per bin so that they all contain the same number as in the lowest bin, but no more than 200

(in case of simulations with fixed fitness effects) or 100 (for simulations with variable fitness

effects). Only replicates with the minimum number of points per bin larger or equal to 5 were

retained.

Identity disequilibrium was measured using the g_2 statistic (David et al. 2007), as

implemented in the 'inbreedR' package (Stoffel et al. 2016). This statistic measures the extent

heterozygosity across a genome is correlated, relative to an expectation under independent

sites. For all simulations, the output VCF file was converted to inbreedR format as described

in the package's manual. The 'g2_snps' function was then used with 100 bootstraps for

generating the mean g_2 estimate along with 95% confidence intervals.

377　Results

378　Comparing self-fertilisation and facultative sex

We first compared the mean fitness and mutation loads with different levels of self-

fertilisation or asexual reproduction, to determine how mutation accumulation differs between

the two mating systems. These simulations were run using the WF model. All results in the

main text assume a fixed *s* and *h*; we will present results with varying *s*, *h* values at the end of

the results section.

Figure 3 shows how the mean fitness changes over time. Under self-fertilisation

(Figure 3a), while the mean fitness is fairly high overall, it is lowest when there is complete

outcrossing. It generally increases with the selfing rate, reaching a maximum for around 95%

self-fertilisation, but then drops for even higher rates. In contrast, the mean fitness drops

dramatically for highly asexual populations (Figure 3b).

Figure 4 (and Supplementary Figure S6) shows mutation counts under these

reproductive regimes. Selfing populations undergo mutation purging, as evidenced by a

decreasing number of mutations in the population (approximately linearly with the selfing rate)

and an increase in homozygous load reflecting higher inbreeding. In contrast, asexual

populations have approximately the same mutation counts (both total and homozygote) as

outcrossing populations, until the population becomes around 95% asexual. At this point, there

is a spike in homozygosity that is then reduced with even higher levels of asexuality, while the

total mutation count rapidly increases. The high mutation count reflects inefficient selection

acting once asexuality becomes prevalent, as reflected in the drop in mean fitness (Figure 3).

The spike in homozygosity likely reflects the fact that for high-but-incomplete asexuality,

mutation accumulation starts acting but there is still sufficient segregation to create

homozygous genotypes. In populations that become completely asexual (i.e., the level of

clonal reproduction approaches 1), there is a lack of genetic segregation so genotypes only

form as heterozygotes (the signature of 'allelic sequence divergence'). The relative

homozygosity (Supplementary Figure S7) also reflects these outcomes; it increases towards

1 with higher selfing, reflecting greater homozygosity due to inbreeding. In contrast, it stays

constant with increased asexuality up to 95% asexual reproduction, but then decreases with

higher rates. No apparent spike at 95% asexuality is observed using this measure.

407 Impact of mitotic gene conversion

Given this increased mutation count under low rates of sex, we next investigated to

what extent mitotic gene conversion can reduce it. Here, we used the non-WF model with the

new gene conversion routine implemented. Figure 5 plots the mean fitness for different per-

base-pair initiation rates of gene conversion. For most values the mean fitness remains low,

similar to results in the absence of gene conversion (Figure 3b). However, once mean gene

conversion per site becomes sufficiently high (in this case, 4×10^{-4} and above) the mean fitness

increases and starts reflecting that of outcrossing populations. These results imply that purging

is indeed occurring, and this mechanism is validated when looking at mutation counts (Figure

6). Here, the mutation count drops while the homozygosity increases (as reflected with

heightened relative homozygosity; Supplementary Figure S8), in line with purging reducing

the total mutation prevalence and increasing the mean fitness.

419 Comparing mitotic gene conversion to self-fertilisation

We next compare the effects of mitotic gene conversion in asexuals to self-fertilisation,

to determine to what extent the two effects are comparable. Figure 7 compares gene

conversion in asexuals against a range of selfing rates. When looking at homozygosity, it

seems that asexuality with the highest gene conversion rates (4×10^{-2}) yield the same

homozygous counts as complete selfing, implying that the two are equivalent. However, the

total mutation count is lower under any level of self-fertilisation, compared to the highest gene

conversion rate used. Hence, it seems that while gene conversion in asexuals can purge

mutations, it does not produce the same benefits in terms of reducing mutation load as

observed under self-fertilisation.

Effects on Linkage and Identity Disequilibrium

As the level of uniparental reproduction increases, so does linkage disequilibrium

across the genome (Supplementary Figures S9, S10). Linkage disequilibrium under self-

fertilisation does not exceed 0.25 on average, while for asexual reproduction it jumps as the

level of clonal reproduction exceeds 95% and stays at high values (r^2 of nearly 1) across the

genome. These results reflect those previously obtained under investigations of LD under

different reproductive modes (Hartfield et al. 2018; Hartfield and Glémin 2024). When gene

conversion is included under clonal reproduction (Supplementary Figure S11), r^2 still stays

constant over the length of genome investigated, but decreases from mean rates of 4×10^{-4}

onwards and dropping below 0.25 and at a similar level to high selfing for the highest value

investigated of 4×10^{-2}.

Identity disequilibrium (ID), as measured by g_2, is mostly zero across simulations, only

becoming non-zero for some high values of selfing or asexual reproduction. Mean ID is also

zero in clonal populations even when gene conversion is present, although there is some

variance around this value under very high gene conversion rates (Supplementary Figure

S12).

Results where *s*, *h* are drawn from distributions

447
To test the generality of these results, we also investigate simulations where *s*, *h* are

drawn from probability distributions, using the same mutation rate. In these cases the fitness

variance also reaches a steady-state within 60,000 generations (Supplementary Figure S13).

Mean fitness results are generally similar to the fixed *s*, *h* case, although if looking just at

selfing populations, the highest fitness is realised for complete selfing (Supplementary Figure

S14). Mutation accumulation in the absence of gene conversion acts slightly differently; while

the highest mutation count occurs under high levels of sexual reproduction, the number of

homozygous genotypes is reduced under both high levels of selfing and asexual reproduction

(Supplementary Figures S15, S16). The relative homozygosity is higher than that for fixed *s*,

h cases under low rates of sex, but drastically drops once the rate of asexuality reaches 95%

and higher (Supplementary Figure S17). Gene conversion in asexuals restores population

fitness in a similar manner to the fixed *s, h* simulations (Supplementary Figure S18), with

slightly lower levels of gene conversion needed to purge deleterious mutations

(Supplementary Figure S19 – S21). LD and ID measurements are also qualitatively similar to

before (Supplementary Figure S22 – S25). Hence, it seems that the results found above do

not strongly depend on the fitness model used.

464   Discussion

465   Summary of results

In this study, we theoretically explored to what extent mitotic gene conversion in

facultative sexual species purge deleterious mutations. We additionally compared this process

to self-fertilisation, as the latter is known to purge deleterious mutations so we can compare

how effective the two are. Using a single-locus model (Figure 1), we first demonstrated that

mitotic gene conversion has the greatest effect in reducing mutation frequency when

asexuality is frequent, but also when mutations are recessive. We also show that, while the

process leads to elevated homozygosity (as measured by the inbreeding coefficient *F;* Figure

2), it will not be substantially elevated and match the effects of selfing unless populations are

fully asexual and the gene conversion rate is high. We subsequently used multi-locus

simulations to explore the effect of mitotic gene conversion on a genome-wide level. In the

absence of gene conversion (Figures 3, 4), highly-asexual species accumulate many

mutations as heterozygotes, leading to a drastic drop in fitness. When gene conversion is

present in fully asexual populations (Figure 5), then we do see a purging effect: once the gene

conversion rate exceeds a critical value (generally an mean rate of 4×10^{-4} per site) the total

mutation count drops and more mutations are present as homozygotes. These levels of gene

conversion also dramatically increase the fitness of asexual populations (Figure 5, Figure

S18). Despite these benefits of gene conversion, the total mutation count is lowest under high

levels of self-fertilisation (Figure 7). Overall, both the single-locus and multi-locus results

indicate that while high rates of gene conversion can purge deleterious mutations under

asexual reproduction, these benefits do not always counter the costs of inefficient selection

under clonality and self-fertilisation can act a stronger force at purging deleterious mutations.

That said, lower levels of gene conversion (rate of 10^{-5} and above) can remove some

deleterious mutations and increase the fitness of asexual organisms, and hence still be

evolutionary favoured.

Comparing mutation accumulation in self-fertilising and asexual populations

491
492

One notable feature of the asexual-selfing comparison is the threshold behaviour of

the mutation load. In the absence of gene conversion, neutral genetic diversity in asexual

populations is similar to outcrossers unless the rate of sex is of the order $1/N$ (Bengtsson 2003;

Ceplitis 2003; Hartfield et al. 2016; Hartfield 2021); background selection also incurs

noticeable reductions in diversity in species with infrequent sex (Agrawal and Hartfield 2016).

We see similar results with regards to the mutation count in Figure 4; it stays fairly stable under

facultative sex unless populations become highly asexual. Conversely, under high selfing we

generally see a drop in the mutation count and heightened homozygosity, in line with mutation

purging. With very high levels of selfing, we see an elevation in both the homozygous mutation

count and in the total mutation number. These increases likely arise due to an additional

increase in mutation number due to genetic drift, as N_e can be dramatically reduced in selfing

populations due to background selection effects (Kamran-Disfani and Agrawal 2014; Roze

2016). Note that this behaviour can change with the fitness model assumptions; under a model

where s and h are drawn from distributions, we instead see a dramatic reduction in the

mutation count under high selfing, in line with a model where genetic variation is purged

(Charlesworth and Charlesworth 1995; Lande and Porcher 2015; Abu Awad and Roze 2018).

Effects of mitotic gene conversion, and comparison with

empirical estimates

Throughout this study, we compared purging via mitotic gene conversion in asexuals

to that via self-fertilisation. We showed that despite the benefits of mitotic gene conversion,

lower mutation counts were observed under high levels of self-fertilisation (see, e.g., Figure

7). These results indicate that self-fertilisation can be a more efficient means of purging

deleterious mutations, likely because it affects the whole genome and homozygous regions

can be formed more rapidly. In contrast, mitotic gene conversion as employed here only

affects a fraction of the genome every generation, so purging acts in a piecemeal manner.

This comparison seems to hold over different mutation models, selfing and gene conversion

rates; one caveat is that there could be a specific, untested combination of mutation and

recombination rates in selfers that might lead to different results.

Our simulation model shows that a mean mitotic gene conversion rate above 4×10^{-4}

per site can start facilitating the purging of deleterious mutations. This is in line with previous

theoretical work showing that, once gene conversion becomes sufficiently high, then it can

substantially reduce genetic variation in asexual populations (Hartfield et al. 2016; Hartfield

2021). Although empirical estimates are sparse, they are usually much lower than this value,

ranging between 10^{-7} and 10^{-5} depending on the study organism (Mandegar and Otto 2007;

Xu et al. 2011; Flot et al. 2013; Sharp and Agrawal 2016). Potentially higher rates were

observed in rice; Jia et al. (2021) inferred a mitotic recombination rate of 10^{-6} with mean tract

lengths ranging between 30-50kb, leading to a per-locus rate on the order of 10^{-2}. Hence, it

seems that while some larger empirical estimates are sufficiently high to cause extensive

deleterious mutation purging, on the whole mitotic gene conversion is usually too weak a force

to lead to extensive purging in nature. However, note that the estimates of Jia et al. (2021)

consider all mitotic recombination events, both crossover and non-crossover events. Our

simulation procedure only considers non-crossover events, which converts more localised

genomic regions. Mitotic crossover events, in contrast, can convert much larger fractions of

the genome (Jia et al. 2021). Even though the net conversion rate per site seems the most

important parameter, longer conversion regions could have a stronger effect through affecting

multiple sites at once, in a similar manner to self-fertilisation. Other forms of asexual

reproduction that also create extensive homozygous regions, such as terminal-fusion

automixis (Mirzaghaderi and Hörandl 2016), could also affect enough of the genome to cause

large levels of mutation purging. Future research can determine whether these different factors

have a stronger effect on asexual fitness.

Another factor not considered in our study is the effect of small and/or changing

population sizes. In small populations, homozygotes can form more frequently due to genetic

drift, which can also lead to the purging of deleterious mutations. However, this process is only

effective at purging highly recessive mutations (Glémin 2003). Gene conversion and drift also

interact to affect the evolution of sex (Roze and Michod 2010). Given the large diploid

population size of 5,000 used in simulations, then drift should have a minimal influence on the

actual purging process under the multi-locus results, but could affect the baseline level of

deleterious mutations in the absence of gene conversion. It will be a rewarding avenue of

future research to determine how drift would affect these processes. Similarly, we also did not

consider demographic feedbacks that are induced during the purging period, where mutation

purging also causes a reduction in population size (Abu Awad and Billiard 2017). The

presence of these feedbacks could increase the extinction probability of populations during

gene-conversion induced purging. Another avenue for the potential further investigation is the

role of adaptive evolution and/or epistasis on the relative advantage of different reproduction

strategies, as they could impact the relative fitness of heterozygotes and homozygotes and

hence the outcomes of the reproductives modes studied here (Kondrashov and Kondrashov

2001; Abu Awad and Roze 2020).

An interesting case where gene conversion may be important under clonal

reproduction is tumour evolution. Here, cells reproduce asexually by mitosis and can

potentially accumulate deleterious mutations due to a lack of recombination (Ní Leathlobhair

and Lenski 2022). Hence, the gene conversion mechanisms investigated here may prevent

transmissible cancers from experiencing mutation meltdown (Ní Leathlobhair and Lenski

2022), and also activate linked mutations that disable tumour-suppressing genes and enable

carcinogenesis (Takahashi and Innan 2022).

Table 1. The summary of model parameters and values used for simulations

Parameter	Values for simulations with f. e.*	
	constant	non-constant
Genome	Diploid, L = 25 Mbp	
Per-base mutation rate	4×10^{-9}	
Per-position recombination rate	4×10^{-8}	
Population size	5 000	
Sample size	50	
Selection	s = 0.01	s ~ Gamma($\bar{s}$, α) mean $\bar{s}$ = 0.01 shape α = 0.2
Dominance	h = 0.2	h ~ U(0, e^{-ks}) k = 91.63
Mean gene conversion tract length	λ = 4 000 bp	

569   * f. e. - fitness effects; selection and dominance coefficients

571   ***Alt text:*** *Table outlining parameter values used in computer simulations.*

Frequency of deleterious mutation $\hat{p}_A$, scaled by μ/hs

Figure 1: Plots of single-locus analytical results for the steady-state frequency of a deleterious mutation. *Results show Equation 1 for different dominance values, scaled to the expected result under outcrossing (μ/hs). Note that μ is not present in this scaled equation. Panels (a), (b) are for different frequencies of sex (with $\hat{F}$ given by Equation 3), while (c), (d) are for different frequencies of self-fertilisation (with $\hat{F}$ given by Equation 2).*

Alt text: *Contour plots showing deleterious mutation frequency with subfigures labelled a to d, illustrating effects of reproductive modes and mitotic gene conversion.*

Reduction in Heterozygosity $\hat{F}$

Figure 2: Plots of single-locus analytical results for the reduction in heterozygosity, $\hat{F}$.

Results show either Equation 2 under facultative sex (panels a, b), or Equation 3 under self-

fertilisation (panels c, d).

Alt text: *Contour plots showing reduction in heterozygosity with subfigures labelled a to d,*

illustrating effects of reproductive modes and mitotic gene conversion.

Figure 3: Mean fitness for different modes of uniparental reproduction. Results are shown as a function of time, for (a) different rates of self-fertilisation or (b) asexual reproduction. Thick lines are mean values, while thinner shading around lines show 95% confidence intervals. Notice the change of scale on the y-axis between (a) and (b).

Alt text: Graphs showing the mean fitness over time for either self-fertilisation or partial asexual reproduction, with different colour lines showing different rates.

Figure 4: Deleterious mutation counts under different uniparental reproductive modes.
As a function of the rate of uniparental reproduction (either selfing or asexual reproduction), these plots show (top) the total mutation count present in the sample of 50 individuals, and (bottom) the number of homozygous genotypes present in the sample. Error bars represent 95% confidence intervals; if they are not shown then they lie within a point. See Supplementary Figure S6 for the total count on a linear scale.

Alt text: *Graphs showing total mutation count and number of homozygote genotypes, with different points indicating different rates of either self-fertilisation or partial asexual reproduction.*

Figure 5: Mean fitness for obligately asexual populations when mitotic gene conversion

is present at different rates. *Results are shown as a function of time. Thick lines are mean*

values, while thinner shading around lines show 95% confidence intervals.

Alt text: *Graph showing the mean fitness over time for complete asexual reproduction, with*

different colour lines showing different rates of mitotic gene conversion.

Figure 6: Deleterious mutation counts for obligate asexual populations with mitotic

gene conversion present. *Results are shown as a function of the rate of gene conversion*

initiation at a single site. Plots show (top) the total mutation count present in the sample of 50

individuals, and (bottom) the number of homozygous genotypes present in the sample. Error

bars represent 95% confidence intervals; if they are not shown then they lie within a point.

Alt text: *Graph showing total mutation count and number of homozygote genotypes, with*

different points indicating different rates of gene conversion under complete asexual

reproduction.

Figure 7: Comparing mutation counts under different levels of self-fertilisation to asexual populations subject to mitotic gene conversion*. Plots show (top) the total mutation count present in the sample of 50 individuals, and (bottom) the number of homozygous genotypes present in the sample. Note that only* gene conversion *rates of $4x10^{-2}$ and $4x10^{-3}$ are shown in the top panel. The points show results under different levels of self-fertilisation; error bars represent 95% confidence intervals, and if they are not shown then they lie within a point. Coloured lines are results for asexuality with different levels of mitotic* gene conversion*, as indicated by the legend; shaded bands around lines indicate 95% confidence intervals.*

***Alt text:** Graph comparing total mutation count and number of homozygote genotypes, with points indicating different selfing rates, and horizontal lines are results for complete asexuality with different gene conversion rates.*

References

Abu Awad, D., and S. Billiard. 2017. The double edged sword: The demographic

consequences of the evolution of self-fertilization. Evolution 71:1178–1190.

Abu Awad, D., and D. Roze. 2018. Effects of partial selfing on the equilibrium genetic

variance, mutation load, and inbreeding depression under stabilizing selection.

Evolution 72:751–769.

Abu Awad, D., and D. Roze. 2020. Epistasis, inbreeding depression, and the evolution of

self-fertilization. Evolution 74:1301–1320.

Agrawal, A. F., and M. Hartfield. 2016. Coalescence with Background and Balancing

Selection in Systems with Bi- and Uniparental Reproduction: Contrasting Partial

Asexuality and Selfing. Genetics 202:313–326.

Agrawal, A. F., and M. C. Whitlock. 2011. Inferences About the Distribution of Dominance

Drawn From Yeast Gene Knockout Data. Genetics 187:553–566.

Baker, H. G. 1955. Self-Compatibility and Establishment After `Long-Distance' Dispersal.

Evolution 9:347–349.

Baker, H. G. 1967. Support for Baker's Law–As a Rule. Evolution 21:853–856.

Bengtsson, B. O. 2003. Genetic variation in organisms with sexual and asexual

reproduction. J. Evol. Biol. 16:189–199.

Bertorelle, G., F. Raffini, M. Bosse, C. Bortoluzzi, A. Iannucci, E. Trucchi, H. E. Morales, and

C. Van Oosterhout. 2022. Genetic load: genomic estimates and applications in non-

model animals. Nat Rev Genet 23:492–503.

Brandt, A., P. Tran Van, C. Bluhm, Y. Anselmetti, Z. Dumas, E. Figuet, C. M. François, N.

Galtier, B. Heimburger, K. S. Jaron, M. Labédan, M. Maraun, D. J. Parker, M.

Robinson-Rechavi, I. Schaefer, P. Simion, S. Scheu, T. Schwander, and J. Bast.

2021. Haplotype divergence supports long-term asexuality in the oribatid mite

Oppiella nova. Proc. Natl. Acad. Sci. U.S.A. 118:e2101485118.

Butlin, R. 2002. The costs and benefits of sex: new insights from old asexual lineages. Nat.

665    Rev. Genet. 3:311–317.

666 Ceplitis, A. 2003. Coalescence times and the Meselson effect in asexual eukaryotes. Genet.

667    Res. 82:183–190.

668 Charlesworth, D., and B. Charlesworth. 1995. Quantitative Genetics in Plants: The Effect of

669    the Breeding System on Genetic Variability. Evolution 49:911–920.

670 Charlesworth, D., M. T. Morgan, and B. Charlesworth. 1993. Mutation accumulation in finite

671    outbreeding and inbreeding populations. Genet. Res. 61:39–56.

672 Charlesworth, D., and J. H. Willis. 2009. The genetics of inbreeding depression. Nat. Rev.

673    Genet. 10:783–796.

674 Chen, J., S. Glémin, and M. Lascoux. 2017. Genetic Diversity and the Efficacy of Purifying

675    Selection across Plant and Animal Species. Mol. Biol. Evol. 34:1417–1428.

676 Chumki, S. A., M. K. Dunn, T. F. Coates, J. D. Mishler, E. M. Younkin, and A. M. Casper.

677    2016. Remarkably Long-Tract Gene Conversion Induced by Fragile Site Instability in

678    *Saccharomyces cerevisiae*. Genetics 204:115–128.

679 Danecek, P., A. Auton, G. Abecasis, C. A. Albers, E. Banks, M. A. DePristo, R. E.

680    Handsaker, G. Lunter, G. T. Marth, S. T. Sherry, G. McVean, and R. Durbin. 2011.

681    The variant call format and VCFtools. Bioinformatics 27:2156–2158.

682 David, P., B. Pujol, F. Viard, V. Castella, and J. Goudet. 2007. Reliable selfing rate estimates

683    from imperfect population genetic data. Molecular Ecology 16:2474–2487.

684 De Meeûs, T., F. Prugnolle, and P. Agnew. 2007. Asexual reproduction: Genetics and

685    evolutionary aspects. Cell. Mol. Life Sci. 64:1355–1372.

686    Dussert, Y., and B. C. Haller. 2019. Mitotic recombination in clonal population. URL:

687    https://groups.google.com/g/slim-discuss/c/5bcUrRLA1cQ/m/kHg3nk4WAgAJ.

688    Accessed July 2024.

689 Engelstädter, J. 2008. Constraints on the evolution of asexual reproduction. BioEssays

690    30:1138–1150.

691 Eyre-Walker, A., and P. D. Keightley. 2007. The distribution of fitness effects of new

692    mutations. Nat. Rev. Genet. 8:610–618.

Felsenstein, J. 1974. The Evolutionary Advantage of Recombination. Genetics 78:737–756.

Flot, J.-F., B. Hespeels, X. Li, B. Noel, I. Arkhipova, E. G. J. Danchin, A. Hejnol, B.

Henrissat, R. Koszul, J.-M. Aury, V. Barbe, R.-M. Barthelemy, J. Bast, G. A. Bazykin,

O. Chabrol, A. Couloux, M. Da Rocha, C. Da Silva, E. Gladyshev, P. Gouret, O.

Hallatschek, B. Hecox-Lea, K. Labadie, B. Lejeune, O. Piskurek, J. Poulain, F.

Rodriguez, J. F. Ryan, O. A. Vakhrusheva, E. Wajnberg, B. Wirth, I. Yushenova, M.

Kellis, A. S. Kondrashov, D. B. Mark Welch, P. Pontarotti, J. Weissenbach, P.

Wincker, O. Jaillon, and K. Van Doninck. 2013. Genomic evidence for ameiotic

evolution in the bdelloid rotifer *Adineta vaga*. Nature 500:453–457.

Frisse, L., R. R. Hudson, A. Bartoszewicz, J. D. Wall, J. Donfack, and A. Di Rienzo. 2001.

Gene Conversion and Different Population Histories May Explain the Contrast

between Polymorphism and Linkage Disequilibrium Levels. Am. J. Hum. Genet.

69:831–843.

Glémin, S. 2003. How are deleterious mutations purged? Drift versus nonrandom mating.

Evolution 57:2678–2687.

Haigh, J. 1978. The accumulation of deleterious genes in a population–Muller's Ratchet.

Theor. Popul. Biol. 14:251–267.

Haller, B. C., and P. W. Messer. 2023. SLiM 4: Multispecies Eco-Evolutionary Modeling. The

American Naturalist 201:E127–E139.

Haller, B. C., and P. W. Messer. 2016. SLiM: An Evolutionary Simulation Framework. URL:

http://benhaller.com/slim/SLiM_Manual.pdf. Accessed July 2024.

Hartfield, M. 2021. Approximating the Coalescent Under Facultative Sex. Journal of Heredity

esaa036.

Hartfield, M. 2016. Evolutionary genetic consequences of facultative sex and outcrossing. J.

Evol. Biol. 29:5–22.

Hartfield, M., and S. Glémin. 2024. Polygenic selection to a changing optimum under self–

fertilisation. PLoS Genet 20:e1011312.

Hartfield, M., and P. D. Keightley. 2012. Current hypotheses for the evolution of sex and

recombination. Integr. Zool. 7:192–209.

Hartfield, M., S. I. Wright, and A. F. Agrawal. 2018. Coalescence and Linkage Disequilibrium

in Facultatively Sexual Diploids. Genetics 210:683–701.

Hartfield, M., S. I. Wright, and A. F. Agrawal. 2016. Coalescent Times and Patterns of

Genetic Diversity in Species with Facultative Sex: Effects of Gene Conversion,

Population Structure, and Heterogeneity. Genetics 202:297–312.

Hill, W. G., and A. Robertson. 1966. The effect of linkage on limits to artificial selection.

Genet. Res. 8:269–294.

Janko, K., P. Drozd, J. Flegr, and J. R. Pannell. 2008. Clonal Turnover Versus Clonal Decay:

A Null Model for Observed Patterns of Asexual Longevity, Diversity and Distribution.

Evolution 62:1264–1270.

Jia, X., Q. Zhang, M. Jiang, J. Huang, L. Yu, M. B. Traw, D. Tian, L. D. Hurst, and S. Yang.

2021. Mitotic gene conversion can be as important as meiotic conversion in driving

genetic variability in plants and other species without early germline segregation.

PLoS Biol 19:e3001164.

Kamran-Disfani, A., and A. F. Agrawal. 2014. Selfing, adaptation and background selection

in finite populations. J. Evol. Biol. 27:1360–1371.

Kershenbaum, S., M. J. Espejo, A. Griffin, and A. Aboobaker. 2024. Gene conversion limits

the cost of asexuality in somatically immortal worms. *bioRxiv*, doi:

10.1101/2023.03.20.533553.

Khakhlova, O., and R. Bock. 2006. Elimination of deleterious mutations in plastid genomes

by gene conversion. Plant J. 46:85–94.

Kondrashov, F. A., and A. S. Kondrashov. 2001. Multidimensional epistasis and the

disadvantage of sex. Proc. Natl. Acad. Sci. USA 98:12089–12092.

Laenen, B., A. Tedder, M. D. Nowak, P. Toräng, J. Wunder, S. Wötzel, K. A. Steige, Y.

Kourmpetis, T. Odong, A. D. Drouzas, M. C. A. M. Bink, J. \AAgren, G. Coupland,

and T. Slotte. 2018. Demography and mating system shape the genome-wide impact

of purifying selection in *Arabis alpina*. Proc. Natl. Acad. Sci. USA 115:816–821.

Lande, R., and E. Porcher. 2015. Maintenance of Quantitative Genetic Variance Under

Partial Self-Fertilization, with Implications for Evolution of Selfing. Genetics 200:891–

906.

Lee, P. S., P. W. Greenwell, M. Dominska, M. Gawel, M. Hamilton, and T. D. Petes. 2009. A

Fine-Structure Map of Spontaneous Mitotic Crossovers in the Yeast Saccharomyces

cerevisiae. PLoS Genet 5:e1000410.

Loewe, L., and W. G. Hill. 2010. The population genetics of mutations: good, bad and

indifferent. Philos. Trans. R. Soc. Lond. B Biol. Sci. 365:1153–1167.

Lynch, M., R. Burger, D. Butcher, and W. Gabriel. 1993. The Mutational Meltdown in

Asexual Populations. J. Hered. 84:339–344.

Lynch, M., J. Conery, and R. Bürger. 1995. Mutational Meltdowns In Sexual Populations.

Evolution 49:1067–1080.

Mandegar, M. A., and S. P. Otto. 2007. Mitotic recombination counteracts the benefits of

genetic segregation. Proc. R. Soc. B 274:1301–1307.

Mark Welch, D. B., and M. Meselson. 2000. Evidence for the Evolution of Bdelloid Rotifers

Without Sexual Reproduction or Genetic Exchange. Science 288:1211–1215.

Mirzaghaderi, G., and E. Hörandl. 2016. The evolution of meiotic sex and its alternatives.

Proc. R. Soc. B 283.

Mráz, P., and V. Mrázová. 2021. Greater reproductive assurance of asexual plant compared

with sexual relative in a low-density sympatric population: Experimental evidence for

pollen limitation. J of Evolutionary Biology 34:1503–1509.

Muller, H. J. 1964. The relation of recombination to mutational advance. Mutat. Res. 1:2–9.

Nettancourt, D. de. 2001. Incompatibility and incongruity in wild and cultivated plants. 2.,

totally rev. and enl. ed. Springer, Berlin Heidelberg New York Barcelona Hong Kong

London Milan Paris Singapore Tokyo.

Ní Leathlobhair, M., and R. E. Lenski. 2022. Population genetics of clonally transmissible

cancers. Nat Ecol Evol 6:1077–1089.

Nordborg, M. 2000. Linkage Disequilibrium, Gene Trees and Selfing: An Ancestral

Recombination Graph With Partial Self-Fertilization. Genetics 154:923–929.

Nordborg, M., B. Charlesworth, and D. Charlesworth. 1996. Increased Levels of

Polymorphism Surrounding Selectively Maintained Sites in Highly Selfing Species.

Proc. R. Soc. B 263:1033–1039.

Nordborg, M., and P. Donnelly. 1997. The Coalescent Process With Selfing. Genetics

146:1185–1195.

Ohta, T., and C. C. Cockerham. 1974. Detrimental genes with partial selfing and effects on a

neutral locus. Genet. Res. 23:191–200.

Olofsson, P., L. Chipkin, R. C. Daileda, and R. B. R. Azevedo. 2023. Mutational meltdown in

asexual populations doomed to extinction. J. Math. Biol. 87:88.

Ossowski, S., K. Schneeberger, J. I. Lucas-Lledo, N. Warthmann, R. M. Clark, R. G. Shaw,

D. Weigel, and M. Lynch. 2010. The Rate and Molecular Spectrum of Spontaneous

Mutations in *Arabidopsis thaliana*. Science 327:92–94.

Otto, S. P. 2003. The Advantages of Segregation and the Evolution of Sex. Genetics

164:1099–1118.

R Core Team. 2024HarRoze LandeFlor. R: A Language and Environment for Statistical

Computing. R Foundation for Statistical Computing, Vienna, Austria.

Roze, D. 2016. Background Selection in Partially Selfing Populations. Genetics 203:937–

957.

Roze, D., and N. H. Barton. 2006. The Hill-Robertson Effect and the Evolution of

Recombination. Genetics 173:1793–1811.

Roze, D., and R. E. Michod. 2010. Deleterious Mutations and Selection for Sex in Finite

Diploid Populations. Genetics 184:1095–1112.

Salome, P. A., K. Bomblies, J. Fitz, R. A. E. Laitinen, N. Warthmann, L. Yant, and D. Weigel.

2012. The recombination landscape in *Arabidopsis thaliana* F2 populations. Heredity

108:447–455.

Schwander, T., and B. J. Crespi. 2009. Multiple direct transitions from sexual reproduction to

apomictic parthenogenesis in Timema stick insects. Evolution 63:84–103.

Sharp, N. P., and A. F. Agrawal. 2016. Low Genetic Quality Alters Key Dimensions of the

806   Mutational Spectrum. PLoS Biol. 14:e1002419.

807   Simmons, M. J., and J. F. Crow. 1977. Mutations Affecting Fitness in *Drosophila*

808   Populations. Annu. Rev. Genet. 11:49–78.

809   Speijer, D., J. Lukeš, and M. Eliáš. 2015. Sex is a ubiquitous, ancient, and inherent attribute

810   of eukaryotic life. Proc. Natl. Acad. Sci. USA 112:8827–8834.

811   Spigler, R. B., K. Theodorou, and S. Chang. 2017. Inbreeding depression and drift load in

812   small populations at demographic disequilibrium. Evolution 71:81–94.

813   Stoffel, M. A., M. Esser, M. Kardos, E. Humble, H. Nichols, P. David, and J. I. Hoffman.

814   2016. inbreedR: an R package for the analysis of inbreeding based on genetic

815   markers. Methods Ecol Evol 7:1331–1339.

816   Takahashi, K. K., and H. Innan. 2022. Frequent somatic gene conversion as a mechanism

817   for loss of heterozygosity in tumor suppressor genes. Genome Res. 32:1017–1025.

818   The Arabidopsis Genome Initiative. 2000. Analysis of the genome sequence of the flowering

819   plant Arabidopsis thaliana. Nature 408:796–815.

820   Theologidis, I., I. M. Chelo, C. Goy, and H. Teotónio. 2014. Reproductive assurance drives

821   transitions to self-fertilization in experimental Caenorhabditis elegans. BMC Biol

822   12:93.

823   Tran Van, P., Y. Anselmetti, J. Bast, Z. Dumas, N. Galtier, K. S. Jaron, K. Martens, D. J.

824   Parker, M. Robinson-Rechavi, T. Schwander, P. Simion, and I. Schön. 2021. First

825   annotated draft genomes of nonmarine ostracods (Ostracoda, Crustacea) with

826   different reproductive modes. G3 Genes|Genomes|Genetics 11:jkab043.

827   Tucker, A. E., M. S. Ackerman, B. D. Eads, S. Xu, and M. Lynch. 2013. Population-genomic

828   insights into the evolutionary origin and fate of obligately asexual *Daphnia pulex*.

829   Proc. Natl. Acad. Sci. USA 110:15740–15745.

830   Vakhrusheva, O. A., E. A. Mnatsakanova, Y. R. Galimov, T. V. Neretina, E. S. Gerasimov, S.

831   A. Naumenko, S. G. Ozerova, A. O. Zalevsky, I. A. Yushenova, F. Rodriguez, I. R.

832   Arkhipova, A. A. Penin, M. D. Logacheva, G. A. Bazykin, and A. S. Kondrashov.

2020. Genomic signatures of recombination in a natural population of the bdelloid

rotifer Adineta vaga. Nat Commun 11:6421.

Weir, W., P. Capewell, B. Foth, C. Clucas, A. Pountain, P. Steketee, N. Veitch, M. Koffi, T.

De Meeûs, J. Kaboré, M. Camara, A. Cooper, A. Tait, V. Jamonneau, B. Bucheton,

M. Berriman, A. MacLeod, and D. Soldati-Favre. 2016. Population genomics reveals

the origin and asexual evolution of human infective trypanosomes. eLife 5:e11473.

Wright, S. 1951. The genetical structure of populations. Ann. Eugen. 15:323–354.

Wright, S. I., S. Kalisz, and T. Slotte. 2013. Evolutionary consequences of self-fertilization in

plants. Proc. R. Soc. B 280:20130133.

Xu, S., A. R. Omilian, and M. E. Cristescu. 2011. High Rate of Large-Scale Hemizygous

Deletions in Asexually Propagating *Daphnia*: Implications for the Evolution of Sex.

Mol. Biol. Evol. 28:335–342.

Yim, E., K. E. O'Connell, J. St. Charles, and T. D. Petes. 2014. High-Resolution Mapping of

Two Types of Spontaneous Mitotic Gene Conversion Events in *Saccharomyces*

cerevisiae. Genetics 198:181–192.