How do self-fertilising and facultative sexual populations differ in mutation accumulation?

Dominik Kopčak[1,2,3], Matthew Hartfield[1,*]

1) Institute of Ecology and Evolution, The University of Edinburgh, Edinburgh EH9 3FL, UK.
2) Institute of Science and Technology Austria, 3400 Klosterneuburg, Austria.
3) Okinawa Institute of Science and Technology Graduate University, Onna-son, Okinawa, Japan.
* Corresponding author: m.hartfield@ed.ac.uk

Key words: Self-fertilisation, facultative sex, gene conversion, deleterious mutation, dominance, mutation load.

Data availability statement: Simulation code and Conda environments with all used packages, together with all scripts have been archived on GitHub (https://github.com/dominik-kopcak/SelfClonGC).

Author contributions: MH designed the project with input from DK. DK wrote and ran the simulations, and analysed the resulting outputs with feedback from MH. MH developed and analysed the single-locus model. MH and DK wrote and edited the manuscript.

Funding: Matthew Hartfield is supported by a NERC Independent Research Fellowship (NE/R015686/1) and a UKRI Frontier Research Guarantee Grant (EP/X027570/1).

Conflict of interest statement: Nothing to declare.

Acknowledgements. We would like to thank Anders Poulsen Charmouh for advising on how to implement the non-constant s, h model, and members of the Hartfield lab for providing feedback on the work.

ORCID links:
Dominik Kopčak: https://orcid.org/0009-0005-3758-4390
Matthew Hartfield: https://orcid.org/0000-0001-6941-0647

Abstract

Self-fertilisation and asexual reproduction are both hypothesised to lead to long-term

extinction due to inefficient selection acting against deleterious mutations. Self-fertilisation can

counter these effects through creating homozygous genotypes and purging deleterious

mutations. Although complete asexuality lacks meiotic gene exchange, mitotic gene

conversion is a mechanism for creating homozygous regions that could limit deleterious

mutation accumulation in a similar manner to self-fertilisation. Here, we compare mutation

accumulation in self-fertilising and facultative sexual populations subject to mitotic gene

conversion, to determine if the latter can purge deleterious mutations in an analogous manner

to self-fertilisation. We first show analytically that this purging is most effective with high levels

of asexuality and gene conversion, and when deleterious mutations are recessive. We further

show using simulations that, when mitotic gene conversion becomes sufficiently high in

obligate asexuals, there is a sudden reduction in the mutation count and a jump in

homozygosity, reflecting mutation purging. However, this mechanism is not necessarily as

efficient at purging compared to high levels of self-fertilisation, and elevated rates of mitotic

gene conversion seem to be needed compared to empirical estimates. Hence, while mitotic

gene conversion can remove recessive deleterious mutations, self-fertilisation seems to be a

more effective mechanism for doing so.

Teaser Text

Asexuality has been argued to be an evolutionary 'dead end', due to a lack of gene

exchange causing inefficient selection acting against deleterious mutations. It has been

proposed that asexuals can counter these negative effects through mitotic gene conversion,

which exposes mutations to selection within individual lineages. Here, we theoretically

investigate how effective this mechanism is. We compare results to those obtained when

individuals reproduce by self-fertilisation, which has similar effects on exposing deleterious

variants. While mitotic gene conversion can be effective in removing recessive deleterious

mutations, high rates are required and it is not always as effective as selfing.

Introduction

The generation of genetic variation is essential for biological evolution. Mutations are

the ultimate source of genetic novelty in all organisms (Loewe and Hill 2010). Other processes

responsible for the generation and/or maintenance of genetic variability (genotype

combinations and their frequency) within eukaryotic populations are gene flow, selection,

genetic drift (fluctuations of the frequency of variants due to 'randomness', i.e., independent

of their fitness; Rice 2004, Charlesworth 2009), non-random mating (Sætre and Ravinet 2019),

recombination (primarily as meiotic crossing-over) and independent chromosome assortment

(Veller et al. 2019). However, the generation of variation is a double-edged sword; although

favourable mutations may emerge that are crucial for long-term survival, it may also result in

a burden of deleterious mutations, which are much more common than favourable ones (Eyre-

Walker and Keightley 2007).

The reproductive system, which determines how offspring are generated in a

population, has a profound impact on the prevalence of deleterious variation. Reproduction

can be either with or without sex. Sex usually involves two phenomena (Hartfield and Keightley

2012): recombination (the exchange of genetic information between homologous DNA

sequences, most commonly as meiotic crossing-over), and fusion of gametes (specialised

cells with halved ploidy and DNA content). In the simplest case when asexuality is equivalent

to clonal reproduction, the genotype of an offspring is identical to that of a parent (except for

de novo mutations). Clonal reproduction encompasses diverse mechanisms, including fission

or budding in unicellular eukaryotes, or budding (e.g., hydra, jellyfishes), fragmentation (e.g.,

corals) and whole-body fission (e.g., flatworms, starfish) in animals or many forms of

vegetative propagation in plants (De Meeûs et al. 2007). Nevertheless, most eukaryotic

organisms undergo some variant of sexual reproduction, at least occasionally (Speijer et al.

2015).

Gametes from which offspring originate may also be derived from the same individual,

leading to self-fertilisation or selfing. The opposite of selfing is panmictic outcrossing, with

anyone in the population having an equal chance of being a mate of anyone else, with self-

mating being rare or even prohibited (if, e.g., self-incompatibility mechanisms are in place to

prevent self-fertilisation; Nettancourt 2001). Cloning and selfing are two types of uniparental

reproduction, which forms a basis of their similarity (e.g., the lack of mate finding) and potential

ecological advantages (e.g., reproductive assurance; Baker 1955, 1967; Theologidis et al.

2014; Mráz and Mrázová 2021).

Despite these apparent similarities, cloning and selfing differ in the manner in which

they affect selection acting against deleterious mutations. Muller's ratchet is a model of a

stochastic mutation accumulation in asexual populations due to the inability to generate

offspring genomes that have fewer deleterious alleles than those of their ancestors (Muller

1964; Haigh 1978). This mechanism is a form of selection interference, which refers to the

effect of strong genetic linkage on the dynamics of variants under selection (Felsenstein 1974).

In completely clonal populations all sites are in absolute genetic linkage, therefore the

presence of multiple deleterious alleles selectively interferes with one another having a strong

impact on mutation accumulation (Hill and Robertson 1966; Roze and Barton 2006; Hartfield

and Keightley 2012). Recombination breaks this linkage and allows the dynamics of variants

to respond to selection independently, i.e., more efficient removal of deleterious mutations

and restoration of the most-fit genotype. Another important consequence due to the lack of

recombination and gamete fusion in diploid asexuals is the divergence of homologous

chromosomes found within one individual. The two genome copies are always transmitted

together across the lineage, gradually and independently accumulating mutations leading to

a high level of heterozygosity (Butlin 2002).

In contrast to cloning, the main genetic consequence of selfing is genome

homogenization. Under complete selfing, heterozygous genotypes *Aa* segregate into gamete

carrying alleles *A* and *a* in equal proportion. Hence, the probability of the offspring of a

homozygous parent losing heterozygosity is 50%, independently of the frequency of the alleles

in the population. The probability that heterozygosity is retained decreases exponentially every

generation, rapidly leading to overall genome homogenization. This homogenisation also

occurs, albeit at a lower rate, under partial selfing. As a consequence, the effective

recombination rate is reduced since recombination creates variability only when

heterozygosity is present (Nordborg et al. 1996; Nordborg 2000). An additional consequence

of increased homozygosity is the exposure of recessive deleterious alleles to selection, giving

rise to inbreeding depression (Charlesworth and Willis 2009). While this leads to an instant

reduction in the fitness of selfers, the subsequent purging and removal of deleterious

mutations from the population could eventually lead to elevated long-term fitness and

persistence of the selfing population (Abu Awad and Billiard 2017). An outcome of selfing

broadly shared with cloning is the removal of neutral variation through extensive linked

selection acting against deleterious mutations, manifesting itself as a reduction in effective

population size (Agrawal and Hartfield 2016).

From the phylogenetic distribution of cloning and selfing species, both reproductive

systems have been called 'dead-end' strategies, meaning that species with them have an

increased probability of extinction (Wright et al. 2013). In asexual populations, this

phenomenon may be explained by mutational meltdown, i.e., the accumulation of deleterious

mutations leading to extinction (Lynch et al. 1993; Olofsson et al. 2023). However, there are

also models in which asexuality does not increase the extinction rate but can still generate this

phylogenetic pattern (Janko et al. 2008; Schwander and Crespi 2009). For example, the rate

of the transition from sexuality to asexuality may be low due to mechanical constraints

associated with sexual reproduction, including the need for a haploid egg to become diploid

or to get activated by a sperm. Rare events might thus be required to overcome these

mechanical constraints, such as mutation to disable the requirement of activation by sperm

(Engelstädter 2008). Furthermore, there exist ancient asexuals that have seemingly existed

for long periods of time in the absence of sexual reproduction, including species of bdelloid

rotifers (Flot et al. 2013), oribatid mites (Brandt et al. 2021), ostracods (Tran Van et al. 2021),

and trypanosomes (Weir et al. 2016). The persistence of these ancient asexuals begs the

question of whether there are underexplored mechanisms at play that could prevent these

species from undergoing excessive accumulation of deleterious variants.

Although genome exchange is intrinsically part of sexual reproduction, gene

conversion (GC) is a mechanism of genome exchange that is also available to asexuals in the

form of mitotic recombination (Lee et al. 2009). GC results in homozygous tracts, akin to the

effect of selfing, and it has been shown to have an important impact on neutral diversity when

sex is rare (Hartfield et al. 2016; Hartfield 2021). Mitotic gene conversion has been argued to

affect deleterious mutation accumulation in asexual species, including the bdelloid rotifer

Adenata vaga (Flot et al. 2013; although see Vakhrusheva et al. (2020) who argue that these

species show evidence of inter-individual gene exchange), the crustacean *Daphnia pulex*

(Tucker et al. 2013) and the planarian worm *Schmidtea mediterranea* (Kershenbaum et al.

2023). Khakhlova and Bock (2006) also show experimental evidence of gene conversion

eliminating deleterious mutations in plastids of tobacco plants, which are uniparentally

inherited and are effectively asexual. However, there exist little theoretical predictions to

determine to what extent gene conversion can indeed purge deleterious mutations, and hence

maintain asexual populations, in an analogous manner to self-fertilisation. One exception is a

model in the supplement of Flot et al. (2013), showing how gene conversion slows down

Muller's Ratchet in asexual populations.

Here, we compare deleterious mutation prevalence and mutation purging in

populations that exhibit either self-fertilisation or facultative sex subject to mitotic gene

conversion. We use both single-locus analytical models and multi-locus simulations to

investigate what extent of genome homogenisation, either through selfing or mitotic gene

conversion, can reduce the frequency of deleterious mutations. Our principal goals are to

determine (i) if deleterious mutations can be removed from facultative sexual species under

realistic cases of gene conversion and fitness effects; (ii) what gene conversion and mutation

fitness parameters lead to purging; and (iii) if there is an equivalent level of selfing that leads

to a similar reduction in mutation prevalence, as compared to that observed under asexuality

with gene conversion.

Single-Locus Mathematical Model

To gain some insight into how mitotic gene conversion affects the purging of

deleterious mutations, we investigate a single-locus model of deleterious allele frequencies in

a diploid population subject to facultative sex, with asexuality occurring at frequency σ and

gene conversion with frequency γ. We consider the frequency of a deleterious allele (denoted

A) that arose from mutation from a wild-type allele *a* with frequency μ; heterozygous mutant

genotypes have fitness 1–*hs* and homozygotes 1–*s*, for *h* the dominance coefficient and *s* the

selection coefficient.

We can track the prevalence of deleterious mutations by creating numerical recursions

of how the frequency is affected by selection, reproduction, mutation and gene conversion

(following the methods of Agrawal and Hartfield (2016), Hartfield (2021); see Supplementary

Mathematica file for the full recursion equations). Although these recursions calculate the

change in genotype frequencies over time, we can convert these genotype frequencies to

allele frequencies; specifically, if the frequency of the mutant allele is p_a, then the frequency

of the mutant homozygote is $p_A^2 + F p_A (1 - p_A)$ (with similar equations for the wild-type

homozygote), and the heterozygote has frequency $2\,p_A\,(1 - p_A)\,(1 - F)$. Here, *F* denotes

the reduction in homozygosity compared to Hardy-Weinberg expectations (Wright 1951), and

shows to what extent gene conversion affects genotype composition by reducing heterozygote

frequencies. After making these transformations, we linearise the equations assuming $\mu, p_A \ll$

1 to obtain steady-state values of p_A and *F* (see Otto (2003) for a similar approach).

After these algebraic calculations, the following steady-state for the deleterious

mutation frequency is obtained:

$$\hat{p}_A = \frac{\mu}{s}\left(\frac{1 - \sigma(1 - s)(1 - \gamma)}{\gamma + h(1 - \gamma)(1 - \sigma(1 - s))}\right) \tag{1a}$$

$$\approx \frac{\mu}{s}\left(\frac{1 - \sigma}{\gamma + h(1 - \sigma)}\right) \tag{1b}$$

Equation 1b assumes that *s*, γ are small so $1 - s$, $1 - \gamma$ both approximate to one.

Both equations reduce to the classic mutation-selection balance result μ/hs if $\sigma = \gamma = 0$

(Wright 1931). We can see which factors have the greatest effect on reducing allele

frequency by dividing Equation 1b by μ/hs and denoting $\varsigma = 1 - \sigma$ as the frequency of sex, to

obtain:

$\hat{p}_A \approx \frac{1}{1+\alpha}$ (2)

where $\alpha = \gamma/h\varsigma$. Hence, the deleterious allele exhibits a lower relative reduction for higher

α, which can either be caused by higher mitotic gene conversion γ, or lower dominance *h*

and rates of sex ς. More specifically, both rare sex and gene conversion act together to not

only expose mutations to selection, but ensure they stay exposed due to a lack of genetic

segregation and are therefore subject to selection over several generations.

Mitotic gene conversion acts on deleterious mutations by increasing the frequency

of homozygotes and reduce that of heterozygotes. We can also quantify the relative

reduction of heterozygosity by calculating the steady-state value of *F*:

$\hat{F} = \frac{\gamma}{1 - \sigma (1 - s) (1 - \gamma)}$ (3)

Equation 3 demonstrates how a combination of gene conversion and rare sex both

lead to higher homozygosity.

Figure 1 illustrates these calculations for different values of the rate of asexuality and

mitotic gene conversion. If mutations are recessive (Figure 1a) then even small amounts of

gene conversion can reduce the prevalence of deleterious mutations compared to outcrossing

expectations, even for intermediate frequencies of sex. Conversely, the frequency of additive

alleles are close to outcrossing expectations (Figure 1b). These results illustrate that mitotic

gene conversion has the strongest effect on purging deleterious mutations if they are

recessive, with the purging effect magnified as sex becomes rarer. Plots of $\hat{F}$ show that while

this gene conversion effect leads to elevated homozygosity, the increase is mostly modest

unless sex is rare and deleterious mutations have a weaker selection effect (compare Figure

1c with 1d). Note that dominance has no effect on $\hat{F}$, presumably because it measures the

prevalence of homozygotes, whose fitness is not affected by the dominance coefficient.

To test the accuracy of these equations, we compared these values to exact values

obtained from numerical recursions (Supplementary Figure S1, with code in Supplementary

Mathematica file). Equation 1a for $\hat{p}_A$ and Equation 3 for $\hat{F}$ matches exact numerical recursions

over a wide range of parameters, showing them to be broadly accurate. The simplified version

of $\hat{p}_A$ (Equation 1b) is accurate as long as sex is frequent and selection is weak, but becomes

inaccurate for rare sex and/or stronger deleterious mutation.

Multi-Locus Simulation Methods

Simulation Outline

While the single-locus model provides insight into how rare sex and gene conversion

affect deleterious mutation frequencies, it does not show how gene conversion leads to

mutation purging across the genome. Hence, we use simulations to compare how gene

conversion leads to mutation removal in asexuals, compared to selfing species.

Simulations were written using SLiM v4.0.1 (Haller and Messer 2023), and were

implemented in either the Wright-Fisher framework (hereafter 'WF model') or in the non-

Wright-Fisher framework (hereafter 'non-WF model'). Simulated populations consist of 5,000

diploid individuals, each with a genome size of $L = 25 \ Mb$ that approximates the length of

Arabidopsis thaliana chromosome 1 (The Arabidopsis Genome Initiative 2000). Individuals do

not carry mutations at the start of the simulations. We let populations evolve for 60,000

generations, which is sufficient for the population variance to reach a steady-state

(Supplementary Figure S2). After this time, 50 individuals are sampled and the number of

mutations is calculated from SLiM's segregating mutations, as described in the 'Mutational

Load Measurement' section below. The mutation rate was set to 4×10^{-9}, which is on the same

order as that inferred from *Arabidopsis thaliana* (Ossowski et al. 2010). Per-base-position

recombination rate was set to 4×10^{-8}, reflecting estimates obtained from crosses of

Arabidopsis thaliana (Salome et al. 2012).

Simulated mutations were deleterious and mutation stacking is allowed by default. Two

types of fitness models were investigated. They differ in whether selection and dominance

coefficients were constant or drawn from a distribution (Table 1). In the first type, fixed values

of deleterious mutations and dominance coefficients were chosen. In the second type, the

selection and dominance coefficients were drawn from a distribution based on Spigler et al.

(2017) (see Table 1). Selection coefficient is modelled by a gamma distribution with shape

parameter α and mean $\underline{s}$. The dominance coefficient of a mutation with selection coefficient *s*

is then drawn from a uniform distribution bounded by 0 and e^{-ks} , where *k* is a constant (see

Table 1 for exact values). Free parameters in the model, *k*, $\underline{s}$, and α, were defined so $\underline{s}$ and $\underline{h}$

match with parameters in our main simulations (*s* = 0.01, *h* = 0.2) and the shape parameter

was based on Chen *et al.* (2017). The general rationale behind this model is to reflect empirical

observations that strongly deleterious mutations tend to be more recessive (Simmons and

Crow 1977; Agrawal and Whitlock 2011).

Implementing mitotic gene conversion in asexual reproduction

By default, recombination in SLiM (including gene conversion) only acts exclusively

when meiosis is involved. Therefore, the reproduction cycle was manually modified through a

reproduction call-back (a SLiM function for governing reproduction), so that mitotic gene

conversion in asexuals can be simulated. For simplicity, the model applies only to a completely

asexual population. The development of the code was facilitated by a post on the SLiM

discussion board (Dussert and Haller 2019) and was extended based on the gene conversion

model under SLiM Wright-Fisher simulations (Haller and Messer 2016).

When an offspring is generated, then the number of gene conversion events that acts

within it is drawn from a Poisson distribution with a mean $L\gamma$ where γ denotes the rate of GC

initiating at a site. For each GC event, the central position is first drawn uniformly through the

chromosome. Then, the tract is extended to the left and right independently by drawing each

length from a geometric distribution with parameter *p:*

$$p = \frac{-\lambda + \sqrt{\lambda}\sqrt{8+\lambda}}{2\lambda}$$ (2)

where λ is the total mean tract length we want to use in simulations. This parameterization is

used to account for the fact that GC tracts with length zero are discarded. This redefined value

of *p* means that the mean tract length is λ after accounting for discarding these tracts (see

Supplementary *Mathematica* notebook for further information). Gene conversion events that

overlap with the chromosome ends are truncated so they stop at the chromosome edges.

After the GC start- and end-points are drawn, they are checked and potentially

modified as follows: tracts of length 0 are removed; GC tracts may have the same start or end

positions, in which case, the start (or end) of one of them is shifted to the right (left), so that

GC tracts overlap; or two GC tracts overlap, in which case two interior GC positions are

removed so that the resulting GC tract is a union of the two smaller ones. Since multiple GC

tracts can overlap, this refinement process is repeated until all start and end positions are

unique and alternate after sorting.

GC can occur on either of the homologous chromosomes with equal probability and

results in the selected part of one chromosome being copied onto the same position at the

other chromosome. Note that for each offspring-production event, all GC tracts are in the same

direction, i.e., material is transferred solely from one strand to another during each

reproduction event. Since the 'initiation' strand is randomly chosen for each reproduction

event, then this effect should not lead to a major bias in results. The mean length of a GC

event is set to 4,000 base pairs; this value accounts for the fact that empirical studies of mitotic

gene conversion tends to find that GC tracts are bimodally distributed, with a combination of

short (on the order of 1kb) and long tracts (on the order of 10kb) (Yim et al. 2014; Chumki et

al. 2016; Jia et al. 2021). Using a mean length of 4,000bp will thus enable a wide distribution

of tract lengths to be drawn that cover both these possibilities.

Further changes to the non-WF model were made so that it behaves as a standard

Wright-Fisher model, based on section 16.15 in the SLiM Manual (Haller and Messer 2016).

Briefly, survival callback limits the age of an individual to a single generation, reproduction

callback samples constant number of parents with probability weighted by their fitness. By

comparing measurements of population fitness (Supplementary Figure S3) and mutation

count (Supplementary Figure S4) between the two simulations, we see that this new routine

yields similar behaviour under low GC rates to the original WF model with complete asexuality.

100 replicate simulations were run if s, h were fixed; 20 replicate simulations were run

for WF models where s, h are drawn from distributions; and 10 replicate simulations were run

for non-WF models where s, h are drawn from distributions. Fewer simulations replicates were

used in the latter two cases due to the much longer runtime of these simulations.

Mutational Load Measurement

Deleterious mutation load was assessed based on two measurements to reflect those

used in empirical genomic studies (e.g., Laenen et al. (2018) in an analysis of *Arabis alpina*).

Recessive load is defined as the number of homozygous mutant genotypes, while additive

load is defined as the total number of mutations present in the population. Both of these

population measures were estimated from 50 individuals' genomes sampled at the end of the

simulation.. We also calculated the relative homozygosity for each simulation, defined as

2*(number of homozygous mutations)/(total number of mutations). This measurement can be

interpreted as the proportion of deleterious mutations that are in homozygous state, across all

sites and individuals. Simulation outputs were analysed in R (R Core Team 2024).

Results

Comparing self-fertilisation and facultative sex

We first compared the mean fitness and mutation loads with different levels of self-fertilisation or asexual reproduction, to determine how mutation accumulation differs between the two mating systems. These simulations were run using the WF model.

Figure 2 shows how the mean fitness changes over time. Under self-fertilisation (Figure 2a), while the mean fitness is fairly high overall, it is lowest when there is complete outcrossing. It generally increases with the selfing rate, reaching a maximum for around 95% self-fertilisation, but then drops for higher rates than this. In contrast, the mean fitness drops dramatically for highly asexual populations (Figure 2b).

Figure 3 (and Supplementary Figure S5) shows mutation counts under these reproductive regimes. Selfing populations (Figure 3a) undergo mutation purging, as evidenced by a decreasing number of mutations in the population (approximately linearly with the selfing rate) and an increase in homozygous load reflecting higher inbreeding. In contrast, asexual populations (Figure 3b) have approximately the same mutation counts (both total and homozygote) as outcrossing populations, until the population becomes around 95% asexual. At this point, there is a spike in homozygosity that is then reduced with even higher levels of asexuality, while the total mutation count rapidly increases. The high mutation count reflects inefficient selection acting once asexuality becomes prevalent, as reflected in the drop in mean fitness (Figure 2). The spike in homozygosity likely reflects the fact that for high-but-incomplete asexuality, mutation accumulation starts acting but there is still sufficient segregation to create homozygous genotypes. In populations that become completely asexual (i.e., the level of clonal reproduction approaches 1), there is a lack of genetic segregation so genotypes only form as heterozygotes (also known as 'allelic sequence divergence' or the Meselson effect; Mark Welch and Meselson (2000); Butlin (2002); Hartfield (2016)). The relative homozygosity

355   (Supplementary Figure S6) also reflects these outcomes; it increases towards 1 with higher

356   selfing, reflecting greater homozygosity due to inbreeding. In contrast, it stays constant with

357   increased asexuality up to 95% asexual reproduction, but then decreases with higher rates.

358   No apparent spike at 95% asexuality is observed using this measure.

Impact of mitotic gene conversion

Given this increased mutation load, we next investigated to what extent mitotic GC can

reduce it. Here, we used the non-WF model with the new GC routine implemented. Figure 4

plots the mean fitness for different per-base-pair initiation rates of GC. For most values the

mean fitness remains low, similar to results in the absence of GC (Figure 2b). However, once

GC becomes sufficiently high (in this case, 10^{-7} and above) the mean fitness increases and

starts reflecting that of outcrossing populations. These results imply that purging is indeed

occurring, and this mechanism is validated when looking at mutation counts (Figure 5). At GC

rates of 10^{-7} and above, the mutation count drops while the homozygosity increases (as

reflected with heightened relative homozygosity; Supplementary Figure S7), in line with

purging reducing the total mutation prevalence and increasing the mean fitness.

Comparing mitotic gene conversion to self-fertilisation

We next compare the effects of mitotic GC in asexuals to self-fertilisation, to determine

to what extent the two effects are comparable. Figure 6 compares GC in asexuals against a

range of selfing rates. When looking at homozygosity, it seems that asexuality with the highest

gene conversion rates (10^{-5}) yield the same homozygous counts as complete selfing, implying

that the two are equivalent. However, the total mutation count is lower under any level of self-

fertilisation, compared to the highest gene conversion rate used. Hence, it seems that while

GC in asexuals can purge mutations, it does not produce the same benefits in terms of

reducing mutation load as observed under self-fertilisation.

379 Results where *s*, *h* are drawn from distributions

To test the generality of these results, we also investigate simulations where *s*, *h* are

drawn from probability distributions, using the same mutation rate. In these cases the fitness

variance also reaches a steady-state within 60,000 generations (Supplementary Figure S8).

Mean fitness results are generally similar to the fixed *s*, *h* case, although if looking just at

selfing populations, the highest fitness is realised for complete selfing (Supplementary Figure

S9). Mutation accumulation in the absence of gene conversion acts slightly differently; while

the highest mutation count occurs under high levels of sexual reproduction, the number of

homozygous genotypes is reduced under both high levels of selfing and asexual reproduction

(Supplementary Figures S10, S11). The relative homozygosity is higher than that for fixed *s*,

h cases under low rates of sex, but drastically drops once the rate of asexuality reaches 95%

and higher (Supplementary Figure S12). Gene conversion in asexuals restores population

fitness in a similar manner to the fixed *h* simulations (Supplementary Figure S13), with similar

levels of gene conversion needed to purge deleterious mutations (Supplementary Figure S14

– S16). Hence, it seems that the results found above do not strongly depend on the fitness

model used.

396 Discussion

397 Summary of results

In this study, we theoretically explored to what extent mitotic gene conversion in

facultative sexual species purge deleterious mutations. We additionally compared this process

to self-fertilisation, as the latter mating-system is known to purge deleterious mutations so we

can compare how effective the two are. Using a single-locus model (Figure 1), we first

demonstrated that mitotic gene conversion has the greatest effect in reducing mutation

frequency when asexuality is frequent, but also when mutations are recessive. We also show

that, while the process leads to elevated homozygosity (as measured by F_{IS}), it will not be

substantially elevated unless populations are fully asexual and gene conversion is high. We

subsequently used multi-locus simulations to explore the effect of mitotic gene conversion on

a genome-wide level. In the absence of gene conversion (Figures 2, 3), highly-asexual species

accumulate many mutations as heterozygotes, leading to a drastic drop in fitness. When gene

conversion is present in fully asexual populations (Figure 4), then we do see a purging effect:

once the gene conversion rate exceeds a critical value (generally an initialisation rate of 10^{-7}

per site) the total mutation count drops and more mutations are present as homozygotes.

These levels of gene conversion also dramatically increase the fitness of asexual populations

(Figure 5, Figure S13b). Despite these benefits of gene conversion, the total mutation count

is lowest still under any level of self-fertilisation (Figure 6), indicating that the benefits of gene

conversion do not fully counter the costs of inefficient selection under clonality.

Comparing mutation accumulation in self-fertilising and asexual populations

One notable feature of the asexual-selfing comparison is the threshold behaviour of

the mutation load. In the absence of gene conversion, neutral genetic diversity in asexual

populations is similar to outcrossers unless the rate of sex is of the order $1/N$ (Bengtsson 2003;

Ceplitis 2003; Hartfield et al. 2016; Hartfield 2021); background selection also incurs

noticeable reductions in diversity in species with infrequent sex (Agrawal and Hartfield 2016).

We see similar results with regards to the mutation count in Figure 3; it stays fairly stable under

facultative sex unless populations become highly asexual. Conversely, under high selfing we

generally see a drop in the mutation count and heightened homozygosity, in line with mutation

purging. With very high levels of selfing, we see an elevation in both the homozygous mutation

count and in the total mutation number. These increases likely arise due to an additional

increase in mutation number due to genetic drift, as N_e can be dramatically reduced in selfing

populations due to background selection effects (Kamran-Disfani and Agrawal 2014; Roze

2016). Note that this behaviour can change with the fitness model assumptions; under a model

where *s* and *h* are drawn from distributions, we instead see a dramatic reduction in the

mutation count under high selfing, in line with a model where genetic variation is purged

(Charlesworth and Charlesworth 1995; Lande and Porcher 2015; Abu Awad and Roze 2018).

Effects of mitotic gene conversion, and comparison with empirical estimates

Throughout this study, we compared purging via mitotic gene conversion in asexuals

to that via self-fertilisation. We showed that despite the benefits of mitotic gene conversion,

lower mutation counts were observed under any level of self-fertilisation (see, e.g., Figure 6).

These results indicate that self-fertilisation is a more efficient means of purging deleterious

mutations, likely because it affects the whole genome and homozygous regions can be formed

more rapidly. In contrast, mitotic gene conversion as employed here only affects a fraction of

the genome every generation, so purging acts in a piecemeal manner. This comparison seems

to hold over different mutation models, selfing and gene conversion rates; one caveat is that

there could be a specific, untested combination of mutation and recombination rates in selfers

that might lead to different results.

Our simulation model shows that a mitotic gene conversion *initiation rate* above 10^{-7}

per site can start facilitating the purging of deleterious mutations. This is in line with previous

theoretical work showing that, once gene conversion becomes sufficiently high, then it can

substantially reduce genetic variation in asexual populations (Hartfield et al. 2016; Hartfield

2021). However, note that this probability is not the same as the rate of a gene conversion

event per site; we have to multiply the initiation rate by 4,000 (the average length of each GC

event) to obtain this value, which gives a rate of around 10^{-4}. Although empirical estimates

are sparse, they are usually much lower than this value, ranging between 10^{-7} and 10^{-5}

depending on the study organism (Mandegar and Otto 2007; Xu et al. 2011; Flot et al. 2013;

Sharp and Agrawal 2016). Potentially higher rates were observed in rice; Jia et al. (2021)

inferred a mitotic recombination rate of 10^{-6} with mean tract lengths ranging between 30-50kb,

leading to a per-locus rate on the order of 10^{-2}. So it seems that while some larger empirical

estimates are sufficiently high to cause extensive deleterious mutation purging, on the whole

mitotic gene conversion is usually too weak a force to lead to extensive purging in nature.

However, note that the estimates of Jia et al. (2021) consider all mitotic recombination events,

both crossover and non-crossover events. Our simulation procedure only considers non-

crossover events, which converts more localised genomic regions. Mitotic crossover events,

in contrast, can convert much larger fractions of the genome (Jia et al. 2021). Even though

the net conversion rate per site seems the most important parameter, longer conversion

regions could have a stronger effect through affecting multiple sites at once, in a similar

manner to self-fertilisation. Other forms of asexual reproduction that also create extensive

heterozygous regions, such as terminal-fusion automixis (Mirzaghaderi and Hörandl 2016),

could also affect enough of the genome to cause large levels of mutation purging. Future

research can determine whether these different factors have a stronger effect on asexual

fitness, especially when mutations interact with one another (i.e., if epistasis is present).

Another factor not considered in our study are demographic feedbacks induced during the

purging period, where mutation purging also causes a reduction in population size (Abu Awad

and Billiard 2017). The presence of these feedbacks could increase the extinction probability

of populations during gene-conversion induced purging.

An interesting case where gene conversion may be important in asexual organisms is

tumour evolution. Here, cells reproduce asexually by mitosis and can potentially accumulate

deleterious mutations due to a lack of recombination (Ní Leathlobhair and Lenski 2022).

Hence, the gene conversion mechanisms investigated here may prevent transmissible

cancers from experiencing mutation meltdown (Ní Leathlobhair and Lenski 2022), and also

activate linked mutations that disable tumour-suppressing genes and enable carcinogenesis

(Takahashi and Innan 2022).

Table 1. The summary of model parameters and values used for simulations

Parameter	Values for simulations with f. e.*	
	constant	non-constant
Genome	Diploid, L = 25 Mbp	
Per-base mutation rate	4×10^{-9}	
Per-position recombination rate	4×10^{-8}	
Population size	5 000	
Sample size	50	
Selection	s = 0.01	s ~ Gamma($\underline{s}$, α) mean $\underline{s}$ = 0.01 shape α = 0.2
Dominance	h = 0.2	h ~ U(0, e^{-ks}) k = 91.63
Mean GC tract length	λ = 4 000 bp	

485   * f. e. - fitness effects; selection and dominance coefficients

Frequency of deleterious mutation $\hat{p}_A$, scaled by μ/hs

488
489

Reduction in homozygosity $\hat{F}$

490
491

Figure 1: Plots of single-locus analytical results. *Panels (a), (b) show the steady-state frequency of the deleterious mutation (Equation 1a), scaled to that expected under outcrossing (μ/hs), for different dominance values. Note that μ, s are not present in this scaled equation. Panels (c), (d) show $\hat{F}$ (Equation 3) for different selection coefficients.*

498 ***Figure 2: Mean fitness for different modes of uniparental reproduction.*** *Results are*

499 *shown as a function of time, for (a) different rates of self-fertilisation or (b) asexual*

500 *reproduction. Thick lines are mean values, while thinner shading around lines show 95%*

501 *confidence intervals. Notice the change of scale on the y-axis between (a) and (b).*

Figure 3: Deleterious mutation counts under different uniparental reproductive modes.

As a function of the rate of uniparental reproduction (either selfing or asexual reproduction),

these plots show (top) the total mutation count present in the sample of 50 individuals, and

(bottom) the number of homozygous genotypes present in the sample. Error bars represent

95% confidence intervals; if they are not shown then they lie within a point. See Supplementary

Figure S5 for the total count on a linear scale.

Figure 4: Mean fitness for obligately asexual populations when mitotic GC is present at

different rates. *Results are shown as a function of time. Thick lines are mean values, while*

thinner shading around lines show 95% confidence intervals.

Figure 5: Deleterious mutation counts for obligate asexual populations with mitotic GC

present. *Results are shown as a function of the rate of gene conversion initiation at a single*

site. Plots show (top) the total mutation count present in the sample of 50 individuals, and

(bottom) the number of homozygous genotypes present in the sample. Error bars represent

95% confidence intervals; if they are not shown then they lie within a point.

Figure 6: Comparing mutation counts under different levels of self-fertilisation to asexual populations subject to mitotic GC. *Plots show (top) the total mutation count present in the sample of 50 individuals, and (bottom) the number of homozygous genotypes present in the sample. Note that only GC rates of 10^{-5} and 10^{-6} are shown in the top panel. The points show results under different levels of self-fertilisation; error bars represent 95% confidence intervals, and if they are not shown then they lie within a point. Coloured lines are results for asexuality with different levels of mitotic GC, as indicated by the legend; shaded bands around lines indicate 95% confidence intervals.*

References

Abu Awad, D., and S. Billiard. 2017. The double edged sword: The demographic

consequences of the evolution of self-fertilization. Evolution 71:1178–1190.

Abu Awad, D., and D. Roze. 2018. Effects of partial selfing on the equilibrium genetic

variance, mutation load, and inbreeding depression under stabilizing selection.

Evolution 72:751–769.

Agrawal, A. F., and M. Hartfield. 2016. Coalescence with Background and Balancing

Selection in Systems with Bi- and Uniparental Reproduction: Contrasting Partial

Asexuality and Selfing. Genetics 202:313–326.

Agrawal, A. F., and M. C. Whitlock. 2011. Inferences About the Distribution of Dominance

Drawn From Yeast Gene Knockout Data. Genetics 187:553–566.

Baker, H. G. 1955. Self-Compatibility and Establishment After `Long-Distance' Dispersal.

Evolution 9:347–349.

Baker, H. G. 1967. Support for Baker's Law–As a Rule. Evolution 21:853–856.

Bengtsson, B. O. 2003. Genetic variation in organisms with sexual and asexual

reproduction. J. Evol. Biol. 16:189–199.

Brandt, A., P. Tran Van, C. Bluhm, Y. Anselmetti, Z. Dumas, E. Figuet, C. M. François, N.

Galtier, B. Heimburger, K. S. Jaron, M. Labédan, M. Maraun, D. J. Parker, M.

Robinson-Rechavi, I. Schaefer, P. Simion, S. Scheu, T. Schwander, and J. Bast.

2021. Haplotype divergence supports long-term asexuality in the oribatid mite

Oppiella nova. Proc. Natl. Acad. Sci. U.S.A. 118:e2101485118.

Butlin, R. 2002. The costs and benefits of sex: new insights from old asexual lineages. Nat.

Rev. Genet. 3:311–317.

Ceplitis, A. 2003. Coalescence times and the Meselson effect in asexual eukaryotes. Genet.

Res. 82:183–190.

Charlesworth, B. 2009. Effective population size and patterns of molecular evolution and

variation. Nat Rev Genet 10:195–205.

Charlesworth, D., and B. Charlesworth. 1995. Quantitative Genetics in Plants: The Effect of

the Breeding System on Genetic Variability. Evolution 49:911–920.

Charlesworth, D., and J. H. Willis. 2009. The genetics of inbreeding depression. Nat. Rev.

Genet. 10:783–796.

Chen, J., S. Glémin, and M. Lascoux. 2017. Genetic Diversity and the Efficacy of Purifying

Selection across Plant and Animal Species. Mol. Biol. Evol. 34:1417–1428.

Chumki, S. A., M. K. Dunn, T. F. Coates, J. D. Mishler, E. M. Younkin, and A. M. Casper.

2016. Remarkably Long-Tract Gene Conversion Induced by Fragile Site Instability in

Saccharomyces cerevisiae. Genetics 204:115–128.

De Meeûs, T., F. Prugnolle, and P. Agnew. 2007. Asexual reproduction: Genetics and

evolutionary aspects. Cell. Mol. Life Sci. 64:1355–1372.

Dussert, Y., and B. C. Haller. 2019. Mitotic recombination in clonal population. URL:

https://groups.google.com/g/slim-discuss/c/5bcUrRLA1cQ/m/kHg3nk4WAgAJ.

Accessed July 2024.

Engelstädter, J. 2008. Constraints on the evolution of asexual reproduction. BioEssays

30:1138–1150.

Eyre-Walker, A., and P. D. Keightley. 2007. The distribution of fitness effects of new

mutations. Nat. Rev. Genet. 8:610–618.

Felsenstein, J. 1974. The Evolutionary Advantage of Recombination. Genetics 78:737–756.

Flot, J.-F., B. Hespeels, X. Li, B. Noel, I. Arkhipova, E. G. J. Danchin, A. Hejnol, B.

Henrissat, R. Koszul, J.-M. Aury, V. Barbe, R.-M. Barthelemy, J. Bast, G. A. Bazykin,

O. Chabrol, A. Couloux, M. Da Rocha, C. Da Silva, E. Gladyshev, P. Gouret, O.

Hallatschek, B. Hecox-Lea, K. Labadie, B. Lejeune, O. Piskurek, J. Poulain, F.

Rodriguez, J. F. Ryan, O. A. Vakhrusheva, E. Wajnberg, B. Wirth, I. Yushenova, M.

Kellis, A. S. Kondrashov, D. B. Mark Welch, P. Pontarotti, J. Weissenbach, P.

Wincker, O. Jaillon, and K. Van Doninck. 2013. Genomic evidence for ameiotic

evolution in the bdelloid rotifer *Adineta vaga*. Nature 500:453–457.

Haigh, J. 1978. The accumulation of deleterious genes in a population–Muller's Ratchet.

Theor. Popul. Biol. 14:251–267.

Haller, B. C., and P. W. Messer. 2023. SLiM 4: Multispecies Eco-Evolutionary Modeling. The

American Naturalist 201:E127–E139.

Haller, B. C., and P. W. Messer. 2016. SLiM: An Evolutionary Simulation Framework. URL:

http://benhaller.com/slim/SLiM_Manual.pdf. Accessed July 2024.

Hartfield, M. 2021. Approximating the Coalescent Under Facultative Sex. Journal of Heredity

esaa036.

Hartfield, M. 2016. Evolutionary genetic consequences of facultative sex and outcrossing. J.

Evol. Biol. 29:5–22.

Hartfield, M., and P. D. Keightley. 2012. Current hypotheses for the evolution of sex and

recombination. Integr. Zool. 7:192–209.

Hartfield, M., S. I. Wright, and A. F. Agrawal. 2016. Coalescent Times and Patterns of

Genetic Diversity in Species with Facultative Sex: Effects of Gene Conversion,

Population Structure, and Heterogeneity. Genetics 202:297–312.

Hill, W. G., and A. Robertson. 1966. The effect of linkage on limits to artificial selection.

Genet. Res. 8:269–294.

Janko, K., P. Drozd, J. Flegr, and J. R. Pannell. 2008. Clonal Turnover Versus Clonal Decay:

A Null Model for Observed Patterns of Asexual Longevity, Diversity and Distribution.

Evolution 62:1264–1270.

Jia, X., Q. Zhang, M. Jiang, J. Huang, L. Yu, M. B. Traw, D. Tian, L. D. Hurst, and S. Yang.

2021. Mitotic gene conversion can be as important as meiotic conversion in driving

genetic variability in plants and other species without early germline segregation.

PLoS Biol 19:e3001164.

Kamran-Disfani, A., and A. F. Agrawal. 2014. Selfing, adaptation and background selection

in finite populations. J. Evol. Biol. 27:1360–1371.

Kershenbaum, S., M. J. Espejo, A. Griffin, and A. Aboobaker. 2023. Gene conversion limits

the cost of asexuality in somatically immortal worms. *bioRxiv*, doi:

10.1101/2023.03.20.533553.

Khakhlova, O., and R. Bock. 2006. Elimination of deleterious mutations in plastid genomes
by gene conversion. Plant J. 46:85–94.

Laenen, B., A. Tedder, M. D. Nowak, P. Toräng, J. Wunder, S. Wötzel, K. A. Steige, Y.
Kourmpetis, T. Odong, A. D. Drouzas, M. C. A. M. Bink, J. \AAgren, G. Coupland,
and T. Slotte. 2018. Demography and mating system shape the genome-wide impact
of purifying selection in *Arabis alpina*. Proc. Natl. Acad. Sci. USA 115:816–821.

Lande, R., and E. Porcher. 2015. Maintenance of Quantitative Genetic Variance Under
Partial Self-Fertilization, with Implications for Evolution of Selfing. Genetics 200:891–
906.

Lee, P. S., P. W. Greenwell, M. Dominska, M. Gawel, M. Hamilton, and T. D. Petes. 2009. A
Fine-Structure Map of Spontaneous Mitotic Crossovers in the Yeast Saccharomyces
cerevisiae. PLoS Genet 5:e1000410.

Loewe, L., and W. G. Hill. 2010. The population genetics of mutations: good, bad and
indifferent. Philos. Trans. R. Soc. Lond. B Biol. Sci. 365:1153–1167.

Lynch, M., R. Burger, D. Butcher, and W. Gabriel. 1993. The Mutational Meltdown in
Asexual Populations. J. Hered. 84:339–344.

Mandegar, M. A., and S. P. Otto. 2007. Mitotic recombination counteracts the benefits of
genetic segregation. Proc. R. Soc. B 274:1301–1307.

Mark Welch, D. B., and M. Meselson. 2000. Evidence for the Evolution of Bdelloid Rotifers
Without Sexual Reproduction or Genetic Exchange. Science 288:1211–1215.

Mirzaghaderi, G., and E. Hörandl. 2016. The evolution of meiotic sex and its alternatives.
Proc. R. Soc. B 283.

Mráz, P., and V. Mrázová. 2021. Greater reproductive assurance of asexual plant compared
with sexual relative in a low-density sympatric population: Experimental evidence for
pollen limitation. J of Evolutionary Biology 34:1503–1509.

Muller, H. J. 1964. The relation of recombination to mutational advance. Mutat. Res. 1:2–9.

Nettancourt, D. de. 2001. Incompatibility and incongruity in wild and cultivated plants. 2.,
totally rev. and enl. ed. Springer, Berlin Heidelberg New York Barcelona Hong Kong

London Milan Paris Singapore Tokyo.

Ní Leathlobhair, M., and R. E. Lenski. 2022. Population genetics of clonally transmissible

cancers. Nat Ecol Evol 6:1077–1089.

Nordborg, M. 2000. Linkage Disequilibrium, Gene Trees and Selfing: An Ancestral

Recombination Graph With Partial Self-Fertilization. Genetics 154:923–929.

Nordborg, M., B. Charlesworth, and D. Charlesworth. 1996. Increased Levels of

Polymorphism Surrounding Selectively Maintained Sites in Highly Selfing Species.

Proc. R. Soc. B 263:1033–1039.

Olofsson, P., L. Chipkin, R. C. Daileda, and R. B. R. Azevedo. 2023. Mutational meltdown in

asexual populations doomed to extinction. J. Math. Biol. 87:88.

Ossowski, S., K. Schneeberger, J. I. Lucas-Lledo, N. Warthmann, R. M. Clark, R. G. Shaw,

D. Weigel, and M. Lynch. 2010. The Rate and Molecular Spectrum of Spontaneous

Mutations in *Arabidopsis thaliana*. Science 327:92–94.

Otto, S. P. 2003. The Advantages of Segregation and the Evolution of Sex. Genetics

164:1099–1118.

R Core Team. 2024HarRoze LandeFlor. R: A Language and Environment for Statistical

Computing. R Foundation for Statistical Computing, Vienna, Austria.

Rice, S. H. 2004. Evolutionary theory: mathematical and conceptual foundations. Sinauer

Associates, Sunderland, Mass., USA.

Roze, D. 2016. Background Selection in Partially Selfing Populations. Genetics 203:937–

957.

Roze, D., and N. H. Barton. 2006. The Hill-Robertson Effect and the Evolution of

Recombination. Genetics 173:1793–1811.

Sætre, G.-P., and M. Ravinet. 2019. Changes in allele and genotype frequency. Pp. 49–68

in Evolutionary Genetics. Oxford University Press.

Salome, P. A., K. Bomblies, J. Fitz, R. A. E. Laitinen, N. Warthmann, L. Yant, and D. Weigel.

2012. The recombination landscape in *Arabidopsis thaliana* F2 populations. Heredity

108:447–455.

Schwander, T., and B. J. Crespi. 2009. Twigs on the tree of life? Neutral and selective

models for integrating macroevolutionary patterns with microevolutionary processes

in the analysis of asexuality. Molecular Ecology 18:28–42.

Sharp, N. P., and A. F. Agrawal. 2016. Low Genetic Quality Alters Key Dimensions of the

Mutational Spectrum. PLoS Biol. 14:e1002419.

Simmons, M. J., and J. F. Crow. 1977. Mutations Affecting Fitness in *Drosophila*

Populations. Annu. Rev. Genet. 11:49–78.

Speijer, D., J. Lukeš, and M. Eliáš. 2015. Sex is a ubiquitous, ancient, and inherent attribute

of eukaryotic life. Proc. Natl. Acad. Sci. USA 112:8827–8834.

Spigler, R. B., K. Theodorou, and S. Chang. 2017. Inbreeding depression and drift load in

small populations at demographic disequilibrium. Evolution 71:81–94.

Takahashi, K. K., and H. Innan. 2022. Frequent somatic gene conversion as a mechanism

for loss of heterozygosity in tumor suppressor genes. Genome Res. 32:1017–1025.

The Arabidopsis Genome Initiative. 2000. Analysis of the genome sequence of the flowering

plant Arabidopsis thaliana. Nature 408:796–815.

Theologidis, I., I. M. Chelo, C. Goy, and H. Teotónio. 2014. Reproductive assurance drives

transitions to self-fertilization in experimental Caenorhabditis elegans. BMC Biol

12:93.

Tran Van, P., Y. Anselmetti, J. Bast, Z. Dumas, N. Galtier, K. S. Jaron, K. Martens, D. J.

Parker, M. Robinson-Rechavi, T. Schwander, P. Simion, and I. Schön. 2021. First

annotated draft genomes of nonmarine ostracods (Ostracoda, Crustacea) with

different reproductive modes. G3 Genes|Genomes|Genetics 11:jkab043.

Tucker, A. E., M. S. Ackerman, B. D. Eads, S. Xu, and M. Lynch. 2013. Population-genomic

insights into the evolutionary origin and fate of obligately asexual *Daphnia pulex*.

Proc. Natl. Acad. Sci. USA 110:15740–15745.

Vakhrusheva, O. A., E. A. Mnatsakanova, Y. R. Galimov, T. V. Neretina, E. S. Gerasimov, S.

A. Naumenko, S. G. Ozerova, A. O. Zalevsky, I. A. Yushenova, F. Rodriguez, I. R.

Arkhipova, A. A. Penin, M. D. Logacheva, G. A. Bazykin, and A. S. Kondrashov.

2020. Genomic signatures of recombination in a natural population of the bdelloid

rotifer Adineta vaga. Nat Commun 11:6421.

Veller, C., N. Kleckner, and M. A. Nowak. 2019. A rigorous measure of genome-wide genetic

shuffling that takes into account crossover positions and Mendel's second law. Proc.

Natl. Acad. Sci. USA 116:1659–1668.

Weir, W., P. Capewell, B. Foth, C. Clucas, A. Pountain, P. Steketee, N. Veitch, M. Koffi, T.

De Meeûs, J. Kaboré, M. Camara, A. Cooper, A. Tait, V. Jamonneau, B. Bucheton,

M. Berriman, A. MacLeod, and D. Soldati-Favre. 2016. Population genomics reveals

the origin and asexual evolution of human infective trypanosomes. eLife 5:e11473.

Wright, S. 1931. Evolution In Mendelian Populations. Genetics 16:97–159.

Wright, S. 1951. The genetical structure of populations. Ann. Eugen. 15:323–354.

Wright, S. I., S. Kalisz, and T. Slotte. 2013. Evolutionary consequences of self-fertilization in

plants. Proc. R. Soc. B 280:20130133.

Xu, S., A. R. Omilian, and M. E. Cristescu. 2011. High Rate of Large-Scale Hemizygous

Deletions in Asexually Propagating *Daphnia*: Implications for the Evolution of Sex.

Mol. Biol. Evol. 28:335–342.

Yim, E., K. E. O'Connell, J. St. Charles, and T. D. Petes. 2014. High-Resolution Mapping of

Two Types of Spontaneous Mitotic Gene Conversion Events in *Saccharomyces*

cerevisiae. Genetics 198:181–192.